Small shifts, big ripples: multi-omics studies reveal how ambient temperatures

govern Chlamydomonas' cellular responses

Short title: Temperature acclimation in Chlamydomonas

Prateek Shetty,[a,b] Volker Wagner,[a] Dinara Myrzakhmetova,[a] Chia-Chi Peng,[b,c] Chulin

Li,[a] Wenshuang Li ,[a,†] Trang Vuong,[a] Joel Ching,[d] Ariane Zander,[d] Rosalind J Allen,[b,d]

Markus Lakemeyer,[b,c] Maria Mittag,[a,b,*]

[a] Matthias Schleiden Institute of Genetics, Bioinformatics and Molecular Botany,

Friedrich Schiller University, 07743 Jena, Germany

[b] Cluster of Excellence Balance of the Microverse, Friedrich Schiller University, 07743

Jena, Germany

[c] Institute for Organic Chemistry and Macromolecular Chemistry (IOMC), Humboldtstr.

10, Friedrich Schiller University, 07743 Jena

[d] Theoretical Microbial Ecology, Institute of Microbiology, Faculty of Biological Sciences,

Friedrich Schiller University, Jena, Germany

[†] Present address: Wuhan Institute of Biomedical Sciences, School of Medicine,

Jianghan University, 430056 Wuhan, China

[*] Address correspondence to m.mittag@uni-jena.de

ABSTRACT

Photosynthetic protists known as microalgae face increasing temperatures due to global climate change. The green biflagellate alga *Chlamydomonas reinhardtii* is commonly found in moist soil and serves as a model for investigating these effects. While responses to extreme thermal stress are well characterized, much less is known about the impact of modest, non-extreme temperature shifts. Understanding microalgal responses to subtle temperature changes is critical, as these primary producers drive ecosystem productivity and food web dynamics. Even small shifts may alter algal fitness and reshape interactions with other microbes. Here*, C. reinhardtii* grew at non-stress, ambient temperatures ranging from 18 °C to 33 °C. Transcriptomic profiling revealed extensive reorganization, with over 5,000 transcripts significantly affected, including those involved in algal-bacterial interactions, photoreception, RNA and lipid metabolism, cilia formation, and the secretome. Proteomic analysis of isolated cilia and secreted proteins corroborated these findings, uncovering major changes in abundance within these sub-proteomes, particularly intraflagellar transport (IFT) complexes in cilia and mating-related proteins in the secretome that are both upregulated at higher temperature. Together, these molecular alterations resulted in pronounced changes in growth, cell size, cilia length, swimming behavior, mating ability and bacterial antagonism. These data emphasize major cellular responses upon small temperature shifts.

INTRODUCTION

Chlamydomonas reinhardtii is a unicellular, biflagellate green microalga that lives in moist soil environments (Sasso et al., 2018). The microalga was originally isolated from a potato field in Massachusetts, USA (Harris, 1989). Subsequent studies have repeatedly found members of the genus *Chlamydomonas*, including *C. reinhardtii* in rice paddy soil samples (Barrett and Koch, 1982; Ghasemi et al., 2008; Lin et al., 2013), but also in rice wash water (Szydlowski et al., 2019; Carrasco et al., 2024).

C. reinhardtii, hereafter called Chlamydomonas, is characterized by a single cup-shaped chloroplast, including a pyrenoid along with RUBISCO for carbon dioxide fixation, and a primitive visual system, the eyespot, situated at the edge of the chloroplast near the cell equator (Harris, 1989; Schmidt et al., 2006). Chlamydomonas is a motile alga as it bears two anterior flagella, also called cilia. Each flagellum is anchored by the basal body to the cell. The flagella are structurally and functionally similar to the 9+2-type axonemes of most motile mammalian cilia (Pazour et al., 2005). The flagella contain an intraflagellar transport system (IFT) that is situated between the axoneme and the ciliary membrane and connected with flagellar length control (Marshall and Rosenbaum, 2001). The IFT complex itself is structurally homologous to the mammalian IFT machinery in both composition and functional organization (McCafferty et al., 2022). The ciliary axoneme consists of substructures such as dynein arms and radial spokes that control bending and are essential for motility, mating, and environmental sensing (Marshall, 2024).

The nuclear and organelle genomes of Chlamydomonas are fully sequenced and annotated. In addition, its defined sexual breeding cycle involving plus and minus gametes, along with extensive mutant collections of haploid vegetative cells, allows for genetic crosses and offers precise genetic manipulation using existing toolkits (Merchant et al., 2007; Crozet et al., 2018; Li et al., 2019; Craig et al., 2023). These features have firmly established Chlamydomonas as a model organism for studying a variety of biological processes such as photosynthesis, light perception, cilia biogenesis or synthetic biology (Scaife and Smith 2016; Dupuis and Merchant, 2023).

In recent years, algal-bacterial interactions have come into focus. Algae and bacteria coevolved for hundreds of millions of years, fostering highly relevant ecological associations (Burgunter-Delamare et al., 2024). Chlamydomonas emerged also as a model to study bacterial interactions. Examples include the vitamin B_{12}-based mutualism

(Helliwell et al., 2015; Bunbury et al., 2022) or the antagonism of the bacterium

Pseudomonas protegens (Aiyar et al., 2017). *P. protegens* secretes toxins including the

cyclic lipopeptide orfamide A that deflagellates the cell within a minute by increasing the

algal cytosolic Ca^{2+} levels. At least four different algal transient receptor potential channels

(TRPs) are involved in mediating the orfamide A signals (Hou et al., 2023).

Temperature is a critical factor in the life of many species. The effects of changes

in ambient temperature on microalgal life are, however, poorly understood. Climate

warming shifts the species distribution of animals, plants and phytoplankton (Mäkinen et

al., 2025) and thus can influence ecosystems. Ambient temperature changes may

influence the growth and metabolism of algae, while extreme temperatures may induce

stress responses in the short term and lead to cell death in the long term. With climate

change driving temperature fluctuations, understanding cellular responses to ambient

temperature changes in model algae like Chlamydomonas is essential. Chlamydomonas

is conventionally cultivated in the range of 20 °C to 25 °C in the laboratory (Harris, 1989).

The alga begins to experience heat stress above 37 °C, marked by the activation of *heat*

shock protein (*HSP*) genes, and experiences strong heat stress when the cultivation

temperature approaches 43.5 °C (Schroda et al., 2015). A lipidome analysis revealed that

Chlamydomonas cells exposed to heat stress of 42 °C exhibited changes in lipid

metabolism and increased lipid saturation as an early heat response (Légeret et al., 2016).

A recent system-wide analysis made use of transcriptomics and proteomics to study the

response of Chlamydomonas to moderate (35 °C) temperature stress that transiently

arrests the cell cycle and acute temperature stress (40 °C) disrupting cell division and

growth (Zhang et al., 2022). The results showed that some of the responses were shared

between the two treatments, although the effects of the 40 °C stress were typically more

extensive. For example, both the 35 °C and 40 °C treatments induced starch

accumulation, although through distinct mechanisms (Zhang et al., 2022). However, the

35 °C treatment also induced a unique set of responses that were absent at 40 °C. It is

also known that low temperatures below 13 °C can induce a cold stress response in

Chlamydomonas. Cold stress causes the arrest of cell division, increases the cell size and

induces palmelloid formation (Ermilova, 2020). Low temperature adaptation in microalgae

also involves production of betaine lipids that are nonphosphorous polar glycerolipids

(Murakami et al., 2018) and are specifically found in some microalgae including
Chlamydomonas (Riekhof et al., 2005).

Most previous studies investigating the effects of temperature on Chlamydomonas
have focused on the above-mentioned stress-inducing temperatures. Typically, these
studies examine responses to acute heat or cold shock treatments lasting only a few hours
to 24 hours, thereby primarily addressing short term acclimation mechanisms. Little is
known about how moderate changes in ambient temperature affect the algae when they
are grown over several diurnal cycles under these conditions; although some proteins
linked with RNA metabolism have been implicated in thermal acclimation (Li et al., 2018).
In our present study, Chlamydomonas was maintained under non-stress temperatures
ranging from 18 °C to 33 °C for four days, allowing system-wide profiling of its adaptive
responses during active growth. This temperature range is particularly relevant because
Chlamydomonas is commonly found in moist soils of rice fields, where the temperature
typically ranges between 15 °C and 30 °C (Chin et al., 1999). However, with ongoing
climate change, soil temperatures are projected to rise, and recent evidence indicates that
heat extremes in soil are increasing even faster than air temperature extremes in many
regions (García-García et al., 2023). By focusing on non-stress temperatures from 18 °C
to 33 °C over several day-night cycles, we sought to investigate how Chlamydomonas
might acclimate in the midterm and perform predictions in the longer term where soil
temperatures are elevated.

Our comprehensive analysis integrates results from multiple assays across four
ambient temperatures (18 °C, 23 °C, 28 °C and 33 °C) in Chlamydomonas. Transcriptomic
analysis reveals how temperature modulates the expression of transient receptor potential
(TRP) channels that are important for some algal-bacterial interactions, components of
RNA and betaine lipid metabolism, major photoreceptors, and numerous ciliary and
secretome-related transcripts. Based on growth data, we found that effects of increasing
temperatures reach a temperature tolerance range at 28 °C. Our transcriptomic results
are thus corroborated by immunoblots of selected photoreceptors, analyses of the ciliary
and secretory proteomes as well as by behavioral assays performed at 18 °C and 28 °C.
Elevated temperatures upregulate many secreted proteins, including those related to
mating, resulting in a significant increase in gamete mating efficiency. Furthermore,
proteins of the IFT complexes associated with flagellar length control, as well as other key

ciliary structural proteins, become more abundant at 28 °C, whereas acetylated tubulin abundance decreases. Importantly, morphological examinations reveal that cell size decreases and cilia length is shortened in cells grown at higher ambient temperatures. Adjustments in ciliary protein composition also affect swimming behavior. At 28 °C, the swimming path of algal cells is characterized by many changes in direction, leading to a reduced speed, while at 18 °C, the cells swim in a linear path without many turns and thus speed up. Together, these results demonstrate that thermal acclimation at different ambient temperatures impacts not only the transcriptome and proteome in Chlamydomonas, but also profoundly influences cell morphology, interspecies interactions, motility, and the sexual cycle.

RESULTS

Ambient temperatures influence the growth and cell size of Chlamydomonas

To address how Chlamydomonas may modulate its physiology and gene expression under non-stressful, ecologically relevant temperature settings, we systematically profiled growth, cell size, and global transcript abundances across a spectrum of ambient temperatures (18 °C, 23 °C, 28 °C, and 33 °C). Cells were grown in a light-dark cycle of 12 h light and 12 h dark (LD 12:12) and investigated during the day (see Methods).

Our results show that cultivation at 18 °C results in a significantly lower growth rate and reduced final cell density compared to the usual growth temperature of 23 °C, while elevated cultivation temperatures of 28 °C and 33 °C accelerated early growth and achieved greater final biomass (Figure 1A, Supplementary Table 1). The growth limitation at 18 °C becomes evident early in cultivation, with significant differences (compared to growth at the reference temperature of 23 °C) being observed consistently from day 3 onwards, ultimately resulting in a significantly reduced final cell density of 1.20×10^7 cells mL^{-1}. In contrast, cultures grown at 28 °C achieved a final cell density of 1.84×10^7 cells mL^{-1}, representing an approximately 20% increase over the 23 °C reference (1.54×10^7 cells mL^{-1}). Although the 33 °C cultivation yielded a slightly higher final cell density than the 28 °C cultivation (1.88×10^7 cells mL^{-1} at day 15), cell densities from days 3 to 14 were in most cases higher at 28 °C, indicating that the cells reach their temperature tolerance range around 28 °C (Figure 1A, Supplementary Table 1).

We also found that the vertical cell size of the algal cells, measured in the middle
of the day at LD6 (Figure 1B), was significantly reduced upon elevation of the cultivation
temperature. On day 4, cells grown at 18 °C averaged 8.80 µm in size. Cells cultivated at
23 °C exhibited a modest yet significant decrease in mean cell size (to 8.25 µm). Further
increasing the temperature to 28 °C and 33 °C intensified this effect, causing the cells to
shrink to 6.65 µm at 28 °C and 7.15 µm at 33 °C (Figure 1B, Supplementary Table 2). The
observation that temperature-dependent effects are even a bit increased at 28 °C versus
33 °C indicates the end of the temperature tolerance range. Similar data were obtained
when cell sizes were determined on day 7 for cultures grown at the higher temperatures
compared to 18 °C (Supplementary Figure 1).

In the next step, we performed comprehensive RNA-seq profiling of cells grown at
temperatures of 18 °C, 23 °C, 28 °C, and 33 °C, with three biological replicates at each
temperature. Our results revealed marked and temperature-dependent shifts in the global
Chlamydomonas transcriptome (Supplementary Table 3). All twelve libraries exceeded 58
million raw reads, with ≥62% of trimmed reads aligning to the Chlamydomonas reference
genome v6.1. The average mapping rate ranged from ~63% to ~73%. Principal-
component analysis (PCA) separated all four temperature conditions along PC1 (50%
variance) and PC2 (40% variance), indicating broad transcriptional reprogramming with
minimal between-replicate variation (Figure 1C). Together, PC1 and PC2 captured 90%
of the total variance. Differential expression analysis highlighted that thousands of
transcript abundances were either increased or decreased at elevated temperatures,
relative to 18 °C (Figure 1D and 1E). Cultivation at both 28 °C and 33 °C elicited large-
scale transcriptional remodeling, with a total of 5,357 and 6,381 genes, respectively, being
differentially abundant at a |$\log_2$fold change (FC)| ≥ 1 and adjusted *P*-value < 0.01.

Growth measurements indicated that Chlamydomonas grows better as
temperature increases respective to 18 °C, with the optimal growth being at 28 °C and no
further gains being observed above this temperature (Figure 1A). Thus, we focused our
comparison further on 18 °C vs. 28 °C. The top 1,000 upregulated and the bottom 1,000
downregulated genes between the 18 °C and 28 °C samples were identified, and the
expression values of these genes were extracted for all four temperature conditions (18
°C, 23 °C, 28 °C, 33 °C). This allowed us to construct a comprehensive heatmap revealing
distinct expression patterns across the temperature gradient (Figure 1F). These data

show that relatively small changes in ambient temperature efficiently remodel the algal
transcriptome.

Gene ontology enrichment of the transcripts that were differentially abundant
between the 18 °C and 28 °C samples revealed several functionally important categories
(Figure 1G, Supplementary Table 4). Extracellular matrix (ECM) remodeling emerges as
an important theme, with upregulated GO terms including "extracellular region
(GO:0005576)" and "polysaccharide catabolic process (GO:0000272)". The
Chlamydomonas ECM is composed of pherophorins and other hydroxyproline-rich
glycoproteins (HRGPs) and is closely connected to the secretome (Luxmi et al. 2018).
Based on these results, it seems that the ECM may undergo modification under different
ambient cultivation temperatures. Additionally, flagellar-related GO terms show
upregulation at 28 °C, including "dynein complex (GO:0030286)", "ATP hydrolysis activity
(GO:0016887)", and "protein tyrosine kinase activity (GO:0004713)". The increased
expression of dynein motor proteins and kinase activities suggests enhanced flagellar
motor activity. Besides, "ATP hydrolysis activity" is an enriched term, perhaps related to
the fact that ciliary motility is driven by dynein motors that hydrolyze ATP (Mitchell et al.,
2005). The upregulation of "protein import into peroxisome matrix (GO:0016558)" could
indicate altered protein trafficking pathways that may affect flagellar protein assembly.
Moreover, protein folding processes including "protein folding (GO:0006457)", "protein
refolding (GO:0042026)", and "unfolded protein binding (GO:0051082)" are
downregulated at 28 °C.

Transcripts encoding photoreceptors and TRP channels show temperature-
dependent regulation and bacterial antagonism against Chlamydomonas is altered
TRP channels such as the human TRPA1 (Moparthi et al., 2022) or TRP1 of
Chlamydomonas (McGoldrick et al., 2019) are known for sensing temperature. Other algal
TRPs participate in controlling algal-bacterial interactions, as mentioned before.
Transcriptomic profiling identified robust temperature-dependent regulation of transcripts
encoding TRP channels (Figure 2A, Supplementary Figure 2). 14 of the 29 known *TRP*s
(Hou et al., 2023) are significantly temperature-dependent altered with a |$\log_2$ FC| ≥ 1 in
their abundances (Supplementary Figure 2). *TRPs 34* and *5* are the most strongly

downregulated at 33 °C compared to 18 °C ($\log_2$ fold change close to −11 and −8,

respectively). Interestingly, three known TRP transcripts implicated in the orfamide A

driven signaling process (*TRP*s 5, 11 and 22; Hou et al., 2023) are altered in their

abundance. Besides the *TRP5* channel transcript, *TRP11*, also exhibits significant

repression at elevated temperatures, while *TRP22* is slightly enhanced. These changes

in the expression of *TRP*s that are involved in orfamide A perception suggest that higher

temperatures may alter the vulnerability of Chlamydomonas to orfamide-mediated

bacterial attacks. To test if there was any temperature-dependent change of bacterial

antagonism, we cocultured Chlamydomonas with *P. protegens* at a ratio of 1:250 in TAP

medium at 18 °C, 23 °C, 28 °C and 33 °C. In these cultures, algae are antagonized by the

bacteria, as evidenced by visual inspection of the green color of the chlorophyll (Figure

2B). We observed that the algal cells in algal-bacterial cocultures cultivated at higher

ambient temperatures (either 28 °C or 33 °C) apparently recovered by day 8 of growth in

coculture, while they had not recovered by this time at lower ambient temperatures (Figure

2B). To track algal cell density more quantitatively, we counted algal cell numbers in

monocultures and in cocultures together with *P. protegens*. At 18 °C, Chlamydomonas in

coculture with *P. protegens* exhibits reduced growth, with no or little recovery until 11 days

(Figure 2C). Similar results are obtained for a coculture grown at 23 °C (Supplementary

Figure 3A). In contrast, at 28 °C and 33 °C, algal cells exhibit a much faster recovery in

cocultures, resuming growth by day 7 and achieving cell densities closer to those of axenic

controls by day 15 (Figure 2D, Supplementary Figure 3B). These findings demonstrate

not only that higher ambient temperatures alter TRP-mediated bacterial antagonism but

that this also confers real-world protection to Chlamydomonas.

Our analysis also revealed temperature-dependent reprogramming of six of the 18

known/predicted photoreceptor transcripts (Figure 2E; Greiner et al., 2017; Sharma et al.,

2024a). Transcripts encoding channelrhodopsins (CHR1, CHR2), which are located in a

specialized part of the plasma membrane covering the eyespot area display pronounced

upregulation at higher temperatures. Conversely, two of the four known cryptochrome

transcripts, particularly CRY-DASH1, show a strong decrease as the temperature

increases. We examined whether the strong up- and down-regulation of the CHR1 and

CRY-DASH1 transcripts, respectively, is reflected at the protein level, using immunoblot

analyses. In both cases, the protein levels mirrored the tendency found in the transcript

profile. Our data reveals a more than fivefold increase in CHR1 protein abundance at

28 °C compared to 18 °C (Figure 2F) and a fivefold increase in CRY-DASH1 protein

abundance at 18 °C compared to 28 °C (Figure 2G).

Ambient temperature drives changes in transcripts encoding proteins of the ciliary

IFT, the secretory pathway, betaine lipid synthesis and RNA-binding proteins

To further elucidate the molecular adaptations underlying temperature-dependent

phenotypic changes in Chlamydomonas, we investigated the expression of additional key

gene families highlighted by the gene ontology enrichment, focusing on flagellar

assembly, ECM remodeling, RNA-binding proteins and enzymes involved in betaine lipid

synthesis.

Flagellar trafficking systems, specifically the IFT complexes, are essential for the

assembly, maintenance, and function of eukaryotic flagella (Pedersen et al., 2006; Katoh

et al., 2016; Ma et al., 2023). The IFT is composed of IFT-A and IFT-B complexes, which

coordinate the bidirectional transport of structural and signaling components along the

axoneme. Proper IFT activity is fundamental for Chlamydomonas motility, environmental

sensing, and cell division. In our transcriptome analysis (Figure 3A), we observed robust

and coordinated upregulation of many *IFT-A* and *IFT-B* subunit transcripts in cells grown

at both 28 °C and 33 °C compared to 18 °C. These findings, if reflected at the protein

level, imply that the upregulation of IFT machinery at higher temperatures reflects a state

of active flagellar remodeling.

Given the enrichment of ECM-related terms, we next examined changes in

transcripts encoding ECM proteins. The ECM in Chlamydomonad algae such as *Volvox*

carteri is primarily composed of hydroxyproline-rich glycoproteins, particularly

pherophorins and related structural proteins. Phylogenetic analyses comparing *V. carteri*

and Chlamydomonas have revealed that both species possess numerous pherophorin-

related proteins. In *V. carteri*, 74 of 99 of these proteins are species-specific, while in

Chlamydomonas, 10 of 30 are unique to the species (von der Heyde and Hallmann, 2022).

These ECM components play a pivotal role in maintaining cell wall integrity, protection,

signaling, and morphogenesis. They not only provide mechanical stability but also

participate in cell-cell communication and environmental adaptation (Hallmann, 2006).

Matrix metalloproteases (MMPs) are key regulators of ECM remodeling, cleaving
glycoproteins to facilitate matrix turnover and structural adaptation. Both are tightly
connected to the algal secretome (Luxmi et al., 2018). Our data show consistent and
dramatic downregulation of multiple transcripts encoding ECM proteins and MMPs at
higher temperatures (Figure 3B).

We also addressed betaine lipid synthesis that is known to be involved in abiotic
stresses, including cold temperatures (Salomon et al., 2025). We found that three
transcripts encoding enzymes involved in the betaine lipid (known as diacylglyceryl-N, N,
N-trimethylhomoserine) pathway are significantly downregulated at the lower ambient
temperature of 18 °C compared to 28 °C (Figure 3C). These include *S-*
adenosylmethionine synthetase1 (*SAS1*), the consequently acting *S-adenosyl-L-*
methionine decarboxylase (*SAMDC*) and the *betaine lipid synthase* (*BTA1*, Figure 3E).
These data highlight that betaine lipid synthesis is already affected at lower ambient
temperature compared to cold stress, at least at the transcript level.

Finally, we screened RNA-binding proteins (RBPs), which tune gene expression
post-transcriptionally or translationally. Nine transcripts encoding RBPs are responsive to
ambient temperatures (Figure 3D). Some RBP transcripts, encoding Pumilio or Musashi
proteins, are mostly significantly upregulated higher temperatures, while the C3 and C1
subunit transcripts of the RNA-binding protein CHLAMY1 (*CRB1* and *CRB3*; Zhao et al.,
2004) as well as NAB1 and DUS16 are downregulated at higher temperatures. Some of
these data are reflected at the protein level, as discussed later.

Temperature-dependent remodeling of the flagellar proteome results in shorter
flagella in cells grown at 28 °C
Transcriptome analysis revealed that many transcripts encoding ciliary IFT proteins are
altered in their abundances in cells grown at 18 °C compared to 28 °C (Figure 3A). To
examine whether these changes are reflected at the protein level, we performed label-
free quantitative proteomic analyses on the membrane-matrix-axoneme (MMA) fraction of
purified flagella from cells grown at 18 °C or 28 °C (see Methods) using single-pot phase-
enhanced sample preparation (Hughes et al., 2019) and liquid chromatography
electrospray ionization tandem mass spectrometry (LC-ESI-MS/MS, see Methods). We
identified a total of 3,271 proteins with at least two unique peptides (Supplementary Table

6). We generated a high-confidence ciliary protein list by matching all detected proteins

against the Chlamydomonas ciliary protein database established by Pazour et al., 2005

(https://chlamyfp.org/ChlamyFPv2/cr_read_sql.php). This comparison yielded 1,395

overlapping proteins, which we defined as the high-confidence ciliary MMA proteome

(Supplementary Table 7). All subsequent differential abundance analyses were restricted

to this set of 1,395 proteins. Using a differential abundance threshold of $|\log_2 FC| \geq 0.5$ and

adjusted P-value < 0.05, we identified 144 ciliary proteins as significantly upregulated and

47 as significantly downregulated at 28 °C compared to 18 °C (Figure 4A, Supplementary

Table 8). Most notably, there was a strong and coordinated upregulation of the IFT-

machinery: five of six IFT-A subunits and three of the 16 IFT-B subunits significantly

increased in abundance (Figure 4B). This proteomic signature mirrors some of the

transcript levels (*IFT*s 121 and 120 of complex A and *IFT*s 74 and 81 of complex B)

elevated at 28 °C (Figure 3A). It should be, however, noted that all other transcript levels

belonging to the upregulated IFT proteins (*IFT*s 43, 139, 144 of the IFT-A complex and

IFT-B 80) are also significantly upregulated in cells grown at 28 °C, but with a $|\log_2 FC|$ of

less than 1 (Supplementary Table 3).

It is known that Chlamydomonas can alter flagella length by activating axonemal

disassembly and stimulating IFT trafficking (Marshall and Rosenbaum 2001; Pan and

Snell, 2005). To determine whether the observed protein trends translate into a

morphological phenotype, we performed immunolocalization of cilia using anti-acetylated

tubulin antibodies (Figure 4C). On day 4 of growth, flagellar length was the longest in cells

cultivated at 18 °C, reaching a mean flagellar length of 8.50 µm under the growth

conditions that were used at LD6 (see Methods). As the cultivation temperature increased,

flagellar length progressively reduced, reaching an average of 6.14 µm at 28 °C and 6.03

µm at 33 °C (Figure 4D, Supplementary Table 9). Comparable results were observed on

day 7 of growth at LD6 (Supplementary Figure 4, Supplementary Table 9).

We also verified whether the reduction in cilia length is reflected in the amount of

the major ciliary tubulin, being α-tubulin1. We observed a small but non-significant

decrease in α-tubulin 1 abundance comparing its level in cells grown at 18 °C versus 28

°C, as measured by LC-ESI-MS/MS (Figure 4E, upper part). However, flagellar α-tubulin

differs from cell body α-tubulin, as it gets post-translationally modified by acetylation

(L'Hernault and Rosenbaum, 1985). To verify the amount of α-tubulin in the flagella, we
performed immunoblots with flagellar proteins along with anti-acetylated tubulin
antibodies. As loading control and for comparison, we used antibodies against casein
kinase 1 (CK1) that is slightly increased in cells grown at 28 °C. The immunoblot analyses
showed a clear reduction of acetylated tubulin while CK1 levels were enhanced in cells
grown at 28 °C (Figure 4E, lower parts; Supplementary Figure 5). These data corroborate
the length reduction of the cilia along with their axoneme.

Temperature-driven changes in flagellar proteins connected to motility alter algal
motility patterns

Further analysis of the flagellar proteome revealed substantial temperature-dependent
remodeling of Chlamydomonas' motility machinery (Figure 5A). A plethora of major motor
proteins, including axonemal and cytoplasmic dynein heavy chains, kinesin motor proteins,
and dynein light/intermediate chains, exhibited pronounced upregulation at 28 °C,
suggesting a heightened demand for active transport and mechanical work within the
flagellum at elevated temperatures. Similarly, proteins associated with ciliopathy
syndromes, especially the Bardet–Biedl syndrome (BBS) proteins, showed strong
upregulation at 28 °C. BBS proteins are connected to IFT trains and influence phototactic
behavior (Liu and Lechtreck, 2018). In contrast, ankyrin repeat proteins (FAP164 and
FAP208), which are critical for maintaining axonemal stability and regulating ciliary and
flagellar architecture, displayed reduced abundance. Moreover, downregulation of two
zinc chaperones (ZNG1, ZNG2) was observed in cells grown at 28 °C. A similar trend was
detected in the GO enrichment analyses, since protein folding and protein refolding were
enriched GO terms, with many transcripts showing downregulation at 28 °C (Figure 1G).
Such chaperones are important for maintaining protein homeostasis and might play a role
in flagellar assembly under changing environmental conditions, also at lower temperature.

We also sought to investigate the possible behavioral consequences of these
molecular changes in flagellar related proteins, by taking a close look at cell motility. For
this purpose, we examined the motility of Chlamydomonas cells using microscopy in a
temperature-controlled chamber, combined with the ImageJ plugin TrackMate (see
Methods), which quantifies various parameters associated with cell trajectories, including
average speed and the turning angle, indicating to what extent cells change their direction

between frames. For this purpose, we used the same cell densities for all samples (see Methods and Discussion). Representative trajectories for individual cells grown at each temperature illustrate severe shifts in motility patterns. Many cells grown at 18 °C followed relatively straight paths, with infrequent directional changes (Figure 5B, Video 1). In contrast, many cells grown at 28 °C exhibited more constrained paths characterized by frequent apparent abrupt directional changes (Figure 5C, Video 2). Figure 5B and C show the trajectories of the individual cells for 80 s, the videos show six exemplarily tracked cells for a total of 57 s. Quantitative analysis of over 150 tracked cells per biological replicate confirmed these observations. It should be noted that the basic swimming behavior of Chlamydomonas involves helical movement with breaststrokes when moving forward (Crockett, 2021). This ensures that the eyespot situated at the nearby *cis-*flagellum perceives light from all directions. The directional change between frames, averaged over a trajectory, was significantly higher in cells grown at 28 °C (Figure 5D). We also calculated the mean swimming speed along each trajectory. Averaging over trajectories, this was significantly reduced in cells cultivated at 28 °C (Figure 5E), consistent with the concept that frequent course corrections lead to slower cell speed.

Taken together, the proteome from cells grown at 28 °C reveals more dynamic flagella with heightened intraflagellar transport, increased abundance of axonemal motors and motility patterns that favor exploratory turning, compared to the more linear swimming that occurs at 18 °C.

Temperature-driven changes in the Chlamydomonas secretome and consequences for mating efficiency

Transcriptomic analyses revealed pronounced downregulation of transcripts encoding ECM components, specifically the pherophorins and MMPs, that are all linked to the secretome (Luxmi et al., 2018). Thus, in the next step, we analyzed whether the Chlamydomonas secretome also changed with cultivation temperature, using label-free quantitative proteomics (see Methods). Mass-spectrometric profiling of Chlamydomonas proteins in supernatants (see Methods) of cells cultured at 18 °C and 28 °C identified a total of 2,763 proteins with at least two identified peptides (Supplementary Table 10). We analyzed these 2,763 proteins using DeepLoc to predict extracellular proteins. Of the 2,763 proteins detected, 557 were predicted by the program as extracellular proteins and

constituted our final dataset of secreted proteins (Supplementary Table 11). Using

thresholds of $|\log_2 FC| \geq 0.5$ and adjusted P-value < 0.05, we found that 317 proteins were

significantly upregulated and 49 were significantly downregulated at 28 °C relative to

18 °C (Figure 6A, Supplementary Table 12).

While many proteins of the cilia coincide in their transcript and protein abundances,

the secretome responds in the opposite direction to the transcriptome. Many transcripts

encoding ECM proteins and other secretory proteins are downregulated (Figure 3C) or

unchanged, but a large number of secreted HRGPs, pherophorins and MMPs is strongly

upregulated in cells grown at 28 °C relative to 18 °C (Figure 6B). We also compared the

ambient temperature-altered secretomes against the published Chlamydomonas mating

secretome, measured at 22 °C (Luxmi et al., 2018), and identified interesting overlaps

between the two datasets. The ambient temperature-altered secretome contains

numerous proteins with established roles in mating, including multiple pherophorins,

gametolysins, and subtilisin-like proteases (these are all marked with a * in front of the

name in Figure 6B). The differential regulation of VLE1, a key subtilisin-like protease that

is also released during gametogenesis, was particularly relevant, as discussed later. The

temperature-dependent regulation of gametogenesis-related proteins could prime cells for

developmental programs that are advantageous under warmer conditions, potentially

accelerating the onset of mating competence.

To test this hypothesis functionally, we performed mating experiments using

Chlamydomonas strains CC-124 (mt⁺) and CC-125 (mt⁻), routinely used for mating assays

in our laboratory (Müller et al., 2017). Strains were cultivated under nitrogen-limited

conditions at different ambient temperatures (see Methods). Mating efficiency assays

confirmed that higher cultivation temperatures significantly increase gamete fusion

success, with cultures pre-conditioned at 28 °C showing enhanced mating competency

compared to those grown at 18 °C (Figure 6C).

DISCUSSION

We used transcriptome analysis to uncover gene expression changes in the biflagellate

model microalga Chlamydomonas acclimated to different ambient temperatures. We

complemented these data with proteomic and behavioral analyses that highlighted

structural changes in cilia impacting the swimming trajectories and speed. We observed shifts in algal-bacterial interactions, photoreceptor regulation, and marked changes in the secretome, connected to enhanced mating behavior, at higher temperatures. The temperature ranges investigated included 18 °C, 23 °C, 28 °C, and 33 °C for transcriptomic analyses, while our proteomic analyses focused on 18 °C and 28 °C. These conditions differed from those in previous system-level studies, which primarily addressed moderate or acute heat stress at 35 °C and 40 °C, compared to normal (25 °C) and heat-stressed growth (Zhang et al., 2022; Zhang et al., 2024), or examined cold stress below 15 °C (Valledor et al., 2013). Anticipating that climate change will both shift the typical range of temperatures and increase extreme stress situations, it is central to understand cellular responses over both stress-inducing and ambient temperature ranges. In their natural habitat of rice fields, Chlamydomonas has historically been exposed to temperatures ranging from 15 °C to 30 °C (Chin et al., 1999). In this study, a range of temperatures extending 3 °C beyond the natural limits experienced by Chlamydomonas were tested, allowing us to investigate the effects of moderate warming without exposing the algae to heat stress. Our results show that growth rates at 28 °C and 33 °C were similar, as seen in Figure 1A, indicating that maximum growth is achieved at 28 °C. Current temperature measurements in a flooded sandy rice soil field in Italy (dated June 20, 2025) revealed an external temperature of 35 °C, with soil temperature of 30 °C at 0 cm and 25 °C at 10 cm depth during the day (data put in order with Innovatech Agro, Italy). Thus, our studies capture actual relevant changes.

The transcriptome studies performed with cells grown in the range between 18 °C and 33 °C revealed changes in the abundances of more than 5,000 and 6,000 transcripts, respectively, between 18 °C and 28 °C (or 33 °C), covering roughly one third of all transcripts based on the genome vs 6.1 (Craig et al., 2023). These data highlight that ambient temperature changes remodel the algal transcriptome drastically.

While missing in land plants, animals and some algae have a high number of TRP channels (Verret et al., 2010). 14 of the 29 annotated TRP channel transcripts based on vs 6.1 of the Chlamydomonas genome (Hou et al., 2023) are significantly altered by temperature with a $|log_2FC| \geq 1$. Especially interesting findings include the changes in transcript abundances encoding the TRP channels 5, 11 and 22 that are involved in the antagonistic interaction of the bacterium *P. protegens* against Chlamydomonas (Hou et

al., 2023). These three channels group close to the vanilloid family (TRPV) tree that includes thermosensitive and calcium-permeable members in mammals, playing also a role in autoimmune diseases (Xiao et al., 2025). Among them, the *TRP5* transcript is most strongly downregulated in cells grown at 33 °C, but also shows a significant decrease at 28 °C. *TRP5* transcript abundance also decreases, in conjunction with a thickening of the algal cell wall, when algal cells are transferred to 3-dimensional (3-D) conditions in the presence of acetate to simulate nature-like conditions in rice fields (Vuong et al., 2025). Since algal fitness against antagonistic bacteria is enhanced under 3-D conditions, we analyzed whether ambient temperature changes would similarly influence algal-bacterial antagonism. We show that higher ambient temperatures indeed enhance algal fitness (Figure 2B - D, Supplementary Figure 3). This may be due to a combination of events happening at higher temperatures, including the decrease of two involved TRP channel transcripts, *TRP5* and *TRP11*. Based on our prior studies (Hou et al., 2023), we assume that the strong decrease of these two channels versus the slight increase of *TRP22* at the higher temperatures will result in a reduced deflagellation rate upon orfamide A exposure and thus strengthen algal fitness. But enhanced algal fitness may be also related to the increased ECM particles found at the higher temperature as shown in the secretome study (Figure 6B). The strongest altered *TRP* transcript (Supplementary Figure 2) was *TRP34* encoding also a member that groups closely to the TRPV family (Hou et al., 2023). In contrast, *TRP1* of Chlamydomonas whose heterologous expression and purification revealed the opening of the pore channel in response to a temperature of 28 °C or 29 °C (McGoldrick et al., 2019), groups closest to the TRP family of yeast and fungi (TRPY/F, Hou et al., 2023). *TRP1* was found to be significantly upregulated at 28 °C in our transcriptome study, however, with a $\log_2$ FC of less than 1 (Supplementary Table 3).

18 known/predicted photoreceptors are encoded in the genome of Chlamydomonas (Greiner et al., 2017; Sharma et al., 2024a). Transcriptome analyses revealed that transcripts encoding six of these photoreceptors are regulated by ambient temperature changes. For two photoreceptors, CHR1 and CRY-DASH1, changes in transcript abundance were also studied at the protein level; while CHR1 is being upregulated at 28 °C, CRY-DASH1 is downregulated at 28 °C (Figure 2F and 2G). In land plants, it was shown that photoreceptors sense stress based as well as ambient temperature changes (reviewed in Sharma et al., 2024b). Our data emphasizes that algal

photoreceptors can also sense ambient temperature variations. Moreover, CHR1, a microbial channelrhodopsin emerged as a major tool in modern biological research and was one of the first algal photoreceptors used for optogenetic approaches (Deisseroth and Hegemann, 2017; Christie and Zurbriggen, 2021). In the algal cell, CHR1 is involved in phototactic behavior and is situated in a specialized part of the plasma membrane surrounding the eyespot (Schmidt et al., 2006). Variations in CHR1 abundance could contribute to the altered motility pattern observed at higher temperatures, where algal cells displayed more frequent turns (Figure 5 D and E). The other photoreceptor, CRY-DASH1, helps balance the photosynthetic machinery by acting as a negative regulator that prevents hyper-stacking of thylakoid membranes (Rredhi et al., 2021). In addition, it is involved as a positive regulator in anoxic metabolism (Rredhi et al., 2024). Our data indicates that these processes may be changed in a temperature-dependent way, although this relationship requires further investigation.

As many transcripts involved in the ciliary structure and in the cellular secretory processes were modified at different temperatures, we extended our studies to include the ciliary and secretory sub-proteomes. Proteomic analysis of the ciliary MMA fraction isolated from cells grown at 18 °C and 28 °C, respectively, revealed a strong increase of most IFT-A and some IFT-B complex proteins, which are crucial for intraflagellar transport from the base of the cilia up to the tip (anterograde IFT train) and from the tip down to the basal body (retrograde IFT train) (Lacey et al., 2024). The increase in IFTs observed at the transcriptome and protein levels (Figures 3A and 4B, Supplementary Table 3) coincides with the upregulation of many motor proteins such as kinesin II motor proteins FLA8, FLA10 and KAP1 (FLA3) and several dyneins (Figure 5A) at higher temperature while the post-translationally modified acetylated α-tubulin is downregulated under these conditions (Figure 4E). The remodeling of many ciliary protein abundances, including the reduced α-tubulin levels in cells grown at 28 °C correlates with shorter cilia observed at this temperature in the middle of the day (LD6, Figure 4C and D). The control of flagellar length in Chlamydomonas is related to the diurnal cycle (Tuxhorn et al., 1998) and is linked with IFT particles as well as axonemal disassembly. This process has been discussed in various models (Marshall & Rosenbaum, 2001; Pan and Snell, 2005; Wemmer et al., 2020; Patel et al., 2024). Moreover, five proteins of the BBS complex (BBS-2, -4, -7-, -8 and -9) related to ciliopathies are increased in abundance in cells grown at 28 °C (Figure 5A).

Previous studies found that a purified BBS-like complex containing BBS1, -4, -5, -7, -8
associate with a subset of IFT proteins (Lechtreck et al., 2009). Three of these IFT-
associated BBS proteins are also temperature-regulated. Interestingly, some
Chlamydomonas BBSome mutants (*bbs1*, *bbs4*, *bbs7*), can assemble full-length, motile
flagella but become completely non-phototactic, due to an unregulated accumulation of
critical signaling proteins (Lechtreck et al., 2009). These results may thus also connect
with the observed modified swimming behavior at 18 °C and 28 °C (Figure 5 B-E).
Behavioral changes in swimming trajectories have been found before under cold and heat
stress conditions; they depend further on the acclimation duration (Meier et al., 2022). In
addition to temperature and acclimation, population density plays an important role in the
determination of swimming speed. The speed increases with higher cell densities due to
the secretion of a low-molecular weight compound (Folcik et al., 2020). Here we used the
same cell density for all measurements and kept it relatively low (5×10^5 cells mL^{-1}) for
comparative studies. This density was selected due to the limited growth of cultures at
18 °C by day 4 after inoculation (Figure 1A, Methods). The calculated speeds are in the
range of priorly observed speeds of Chlamydomonas cells based on similar cell numbers
(Folcik et al., 2020). Using a specialized temperature-controlled microscopy setup (see
Methods), we revealed that even ambient changes in temperature result in severe
alterations in swimming speed as well as in the frequency of turns performed by the algal
cells.

The analysis of the secreted proteome revealed a striking temperature-dependent
reorganization in Chlamydomonas characterized by the substantial upregulation of MMPs,
the vegetative lytic enzyme1, VLE1, and a diverse array of pherophorins at 28 °C, creating
a molecular environment primed for extensive ECM modification. MMPs belong to a family
of zinc-dependent proteinases and can cleave diverse substrates including components
of the ECM (Hey and Linder, 2024). One of the first MMPs of Chlamydomonas that was
characterized represents the gamete lytic enzyme (Kinoshita et al., 1992), but several
MMPs and other proteolytic enzymes such as VLE1 have been also found in the mating
secretome performed at 22 °C (Luxmi et al., 2018). Pherophorins are a major group of
extracellular hydroxyproline-rich proteins in Chlamydomonas (Barolo et al., 2022) and are
also important in *V. carteri* in ECM biosynthesis (von der Heyde and Hallmann, 2022). 18
pherophorins are upregulated at higher temperature as well as eight further

hydroxyproline rich proteins, including CWP2, FAPs 40, 113 and 137, ISGs 2 and 6,

MAW4 and VSP3. Several of these proteins (marked by an asterisk in Figure 6B) overlap

with proteins identified in the Chlamydomonas mating secretome study (Luxmi et al.,

2018). Sexual algal reproduction was shown earlier to depend on the secreted protein

profile (Snell et al., 1989). Our results highlight that cultivation at elevated ambient

temperatures significantly enhances the secretion of proteins required for mating. Thus,

we examined the mating efficiency of plus and minus gametes at different temperatures

and found that gametogenesis increases as temperature rises (Figure 6C). These data

highlight a direct influence of small ambient temperature changes on the sexual life cycle

of Chlamydomonas.

The ciliary and secretome proteomes identified more proteins (Supplementary

Tables 6 and 10) than are present in the ciliary database (Supplementary Table 7) and

predicted to be localized extracellularly by DeepLoc (Supplementary Table 11),

respectively. We are aware that these proteins may be contaminants. For example, at

higher temperature, some cells may have been subject to cell lysis and thus may have

influenced the results. It may, however, also be the case that some candidates have been

neglected based on the selection criteria used. For example, VLE1 is well known as

secreted protein (Luxmi et al, 2018), but it is not predicted to be localized to the

extracellular region with DeepLoc. Due to the prior study, it has been included in our list.

In some cases, the transcriptome changes agree with changes found at the protein

level, but in other cases, especially with the secretome, correlations are scarce or even

contradictory. Thus, post-transcriptional and translational control mechanisms along with

RBPs are most likely involved. The transcriptome studies have shown that transcripts

encoding a variety of RBPs are altered (Figure 3D). Former studies of RBPs selected at

different ambient temperatures had revealed that the C3 subunit of RBP CHLAMY1

(CRB3) is downregulated at the protein level in cells grown at 28 °C compared to 18 °C

(Li et al., 2018), which agrees with its transcript level (Figure 3D). Notably, the C3 subunit

confers thermal acclimation to Chlamydomonas as shown by mutant lines (Li et al., 2018).

Our present studies highlight that temperature acclimation to different ambient

temperatures represents an important step in the regulation of a variety of biological

processes in Chlamydomonas. Some of the depicted components such as TRP channels

and photoreceptors are likely to be members of the thermal acclimation pathway involved

in sensing ambient temperatures. Components like the C3 subunit seem to mediate sensed information. Finally, there are several members like IFT or ECM proteins involved in executing cellular responses like the shortening of the cilia or the enhancement of protein secretion. These events then induce behavioral changes such as swimming speed and turns, mating ability or bacterial antagonism. Altogether, our findings show that even modest changes in temperature, that may seem negligible at first glance, influence key properties of the organism and demand further focused study in the future. Thus, small shifts can indeed cause big ripples.

METHODS

Strains of *Chlamydomonas reinhardtii* (thereafter Chlamydomonas) used and quantification of growth

Chlamydomonas wild-type strain SAG 73.72 (mt$^+$) was obtained from the Culture Collection of Algae at Göttingen University and used for all experiments except for the mating assay. Strain SAG73.72 is an isolate of Tsubo's strain C8 that is close to 21 gr from Sager (Harris, 1989). Strain 21 gr served as a crossing partner for generating strain CC-4532, whose genome has been sequenced (Craig et al., 2023).

Cells were cultivated in Tris-Acetate-Phosphate (TAP) medium (Harris, 1989) under a 12h : 12h light:dark (LD) cycle at a light intensity of 50 µmol m^{-2} s^{-1} and stirring (100 rpm). Growth experiments were performed under four cultivation temperatures, 18 °C, 23 °C, 28 °C, and 33 °C. To initiate precultures, cells were transferred from TAP agar plates (Harris 1989) into 15 mL of liquid TAP medium in sterile NUNC flasks. These were incubated at the designated temperatures with orbital shaking at 100 rpm under the LD cycle for 4 days. On day 4, main cultures were prepared by measuring the cell density of each preculture and adjusting the starting concentration to 1 × 10^5 cells mL^{-1}. This step ensured uniform inoculum density across all temperature conditions. Culture densities were quantified daily by sampling 180 µL of sample and counting cells on an improved Neubauer chamber (no. T729.1; Carl Roth, Karlsruhe, Germany). Visual documentation of chlorophyll content (as a proxy for pigmentation and viability) was also captured every second day by photographing culture flasks. Cell samples for the different experiments were always taken in the middle of the day, at LD6, unless otherwise indicated.

Determination of cell size

Cell sizes were quantified on day 4 and day 7 of growth (Figure 1A). For this purpose, 2 mL aliquots from the main Chlamydomonas cultures were harvested and centrifuged at 500 g for 10 min. The algal cells were fixed with Lugol's solution and further treated as described previously (Vuong et al., 2025). Brightfield microscopy images along with a Zeiss Axiophot were taken to measure multiple cells with ImageJ (version 1.53t, Schneider et al., 2012) taking the vertical length of the cells into consideration.

Transcriptome sequencing and data analysis

Total RNA was isolated from cells grown in TAP media under four different ambient temperatures, with three biological replicates per condition. Total RNA was extracted using the Qiagen Plant RNeasy Mini kit (Qiagen, Germany) according to the manufacturer's instructions. RNA concentration and purity were measured using a NanoDrop ND-1000 UV/Vis spectrophotometer (Thermo FisherScientific, USA). Library preparation and sequencing were performed by Eurofins Genomics (Ebersberg, Germany). For each sample, stranded TruSeq cDNA libraries were prepared using poly(A) enrichment. Libraries were sequenced on an Illumina NovaSeq 5000 platform to generate 150 bp paired-end (PE) reads. The raw reads were processed using Fastp (v.0.23.4) (Chen et al., 2018) with default trimming options and base correction enabled to correct for low-quality bases. Kallisto (v0.50.1) (Bray et al., 2016) was used to quantify the abundance of each transcript against the reference *C. reinhardtii* v.6.1 transcriptome downloaded from Phytozome. All further analysis was carried out in Rstudio (v.2024.04.1, Posit team, 2023). R tximport package (v.1.32.0) (Soneson et al., 2015) was used to import the gene-level expression and normalize the data. Genes with read counts fewer than 10 reads in at least 6 samples were marked as lowly expressed and removed from further analysis. Differential expression analysis was carried out using DESeq2 package (v.1.44.0) (Love et al., 2014). A gene was considered statistically significant if the |$\log_2$ (fold change)| was ≥ 1 and the adjusted P-value < 0.01. Cultures grown at 18 °C served as the reference condition for all pairwise comparisons. Genes with positive $\log_2$ fold changes were designated as upregulated, and those with negative values were designated as

downregulated relative to the 18 °C condition. Gene ontology enrichment was carried out
using topGO (v.2.56.0).

Isolation of flagella and ciliary proteins from the membrane-matrix-axoneme (MMA)
fraction

Cells were grown under the above-mentioned conditions for four days at 18 °C and 28 °C,
respectively. Cells were harvested and washed according to Bösger et al. (2014). The
temperature of rotors and washing buffers was adjusted to 18 °C and 28 °C, respectively.
After deflagellation of the cells with dibucain and purification of the flagella using sucrose
step-gradient centrifugation, the MMA fraction was prepared by adding NP-40 to the
isolated flagella to achieve a final concentration of 1%, as described in Bösger et al. (2014).
Four independent biological replicates were prepared for each temperature condition,
from which three were used for further analysis.

Proteome analysis of the MMA fraction

For the analysis of the proteins in the MMA fraction, 0.5 mg proteins of each MMA fraction
were reduced and carbamidomethylated according to Zhang et al., 2005. Samples were
then subjected to the single-pot, solid phase-enhanced sample preparation (SP3)
according to Hughes et al., (2019). Briefly, the proteins were bound to magnetic SP3
beads, washed three times to remove contaminants, and digested overnight at 37 °C
directly on the beads by adding 20 µg of a Trypsin and LysC protease mix (MS grade,
Promega V507A) dissolved in 40 µL of the provided buffer per 0.5 mg sample in Eppendorf
Protein LoBind tubes that were shaken at 1,000 rpm in a Thermomixer. Prior to the mass
spectrometric analyses, the resulting peptides were further purified by reversed phase
chromatography using ZipTip microtips supplemented with additional Poros R2 10 µm
reversed phase material (Bösger et al., 2009), dried using a Speed-Vac and stored at –
80 °C before mass spectrometry analysis. Immediately before LC-ESI-MS/MS, peptides
were reconstituted in 25 µL of 5% MeCN/0.1% TFA by vortexing thrice and a 5 min cycle
of ultrasonic bath sonication. The samples were then centrifuged for 5 min at 13,000 rpm
and the supernatant was transferred to MS vials for LC-MS/MS measurements.

Preparation of samples for the secretome proteome

Secretome analysis was performed using cultures grown at 18 °C and 28 °C. Cells were cultivated for four days in TAP medium, as described above. Three independent biological replicates were prepared for each temperature condition. On day 4, cultures were harvested to collect cell-free supernatants. For each sample, the culture was centrifuged at 1,000 g for 10 min to remove intact cells and large particulates. The resulting supernatant was then filtered through a 0.22 µm pore-size membrane filter (Carl Roth, Germany) to ensure complete removal of any residual cells or debris.

10 mL of the supernatant from each sample were first concentrated to 250 µL using a 3 kDa MWCO Amicon centrifugal filter. Each concentrate was transferred into an Eppendorf Protein LoBind tube and adjusted to 15 µg total protein, as determined by BCA assay. The samples were finally brought to 250 µL with 1 M triethyl ammonium bicarbonate (TEAB) buffer, yielding a final concentration of 50 mM TEAB (pH ~8.5), and incubated at 60 °C for 10 min with shaking at 1,000 rpm.

The subsequent steps were performed at 22 °C with shaking at 1,500 rpm. Proteins were sequentially reduced by adding 2.5 µL TCEP to a final concentration of 5 mM and incubated for 10 min, then alkylated with 5 µL iodoacetamide to a final concentration of 10 mM for 30 min and finally quenched by adding 5 µL DTT to a final concentration of 10 mM for 15 min. Next, a 1:1 mixture of hydrophobic and hydrophilic Sera-Mag carboxylate coated magnetic beads (50 mg mL^{-1}; 2 mg total; particles E3/E7; Cytiva) were prewashed three times with 100 µL water. After washing, 2.5 µL of the bead suspension was added to each sample, followed by ethanol to a final concentration of 60%. Samples were incubated for 5 min to bind proteins. Beads were bound using a magnetic rack. The supernatant was removed, and the beads were washed three times with 400 µL of 80% ethanol and once with 400 µL of MeCN. After an air-drying step, the beads were resuspended in 50 µL of 50 mM TEAB (pH ~8.5).

Proteins on the beads were digested by adding 0.5 µL of recombinant LysC (0.2 µg µL^{-1}, MS grade, Promega V1671) and incubation at 37 °C for 1 h with shaking at 1,200 rpm. Subsequently, 0.4 µL of trypsin (0.5 µg µL^{-1}, MS grade, Promega VA9000) was added, and digestion continued overnight at 37 °C with shaking at 1,200 rpm. After digestion, the supernatant was collected into a new Protein LoBind tube. The beads were washed twice with 20 µL of 1% TFA, whereby each wash was incubated for 5 min at 22 °C

while shaking at 1,000 rpm. The combined supernatant and wash fractions were acidified

with 1 µL of 80% TFA (to pH < 3) and desalted using a 100 µL Pierce C18 Tips (Thermo

Fisher 87784). Tips were equilibrated with 100 µL of 50% MeCN/0.1% TFA twice, followed

by equilibration with 100 µL of 0.1% TFA twice. Samples were loaded by slowly aspirating

and dispensing the solution ten times. Tips were washed twice with 100 µL of 0.1% TFA,

and bound peptides were eluted three times with 75 µL of 80% MeCN/0.1% TFA. Eluates

were combined, dried in a vacuum concentrator at 35 °C, and stored at –80 °C until

analysis. Immediately before LC-ESI-MS/MS, peptides were reconstituted in 25 µL of 5%

MeCN/0.1% TFA by vortexing thrice and a 5 min cycle of ultrasonic bath sonication. The

samples were then centrifuged for 5 min at 13,000 rpm and the supernatant was

transferred to MS vials for LC-MS/MS measurements.

LC-ESI-MS/MS measurements

LC-MS/MS analysis was performed with an Ultimate3000 Nano-HPLC system (Thermo

Fisher Scientific) coupled to an Orbitrap Fusion Tribrid mass spectrometer (Thermo Fisher

Scientific). Samples were analyzed using an Acclaim PepMap 100 C18 trap column

(0.075 × 200 mm; Thermo Fisher Scientific) and an Aurora Ultimate XT C18 UHPLC

analytical emitter column (0.075 × 250 mm; Ionopticks, Collingwood, Australia) in an

EASY Spray source setting with HeatSync Column Heater (Ionopticks, Collingwood,

Australia). Each sample was loaded onto the trap column, washed with 0.1% TFA in water,

and then transferred to the analytical column. Peptides were separated in a 152-minute

run (buffer A: water with 0.1% FA, buffer B: MeCN with 0.1% FA, flowrate: 400 nL/min)

with the following gradient: from 5% B to 22% B in 105 min, from 22% B to 32% B in 10

min, from 32% B to 90% B in 10 min, hold at 90% B for 10 min, then to 5% B over 0.1

min, and hold at 5% B for 10 min. The Orbitrap Fusion Tribrid mass spectrometer was

operated in a data-dependent acquisition (DDA) mode with a cycle time of 3 s. Full scan

acquisition was performed in the Orbitrap at a resolution of 120,000 in the scan range of

300 – 1,500 m/z (AGC target of 50%, maximal injection time of 50 ms). Monoisotopic

precursor selection as well as dynamic exclusion (exclusion duration: 60 s) was enabled.

Precursors with charge states of 2 – 7 and intensities greater than 5e3 were selected for

fragmentation. Isolation was performed in the quadrupole using a window of 1.6 m/z.

Fragments were generated using higher-energy collisional dissociation (HCD, normalized

collision energy: 30%) and detected in the Ion Trap (scan rate: rapid, mass range: normal,

maximal injection time: 35 ms, AGC target: standard).

Proteomics data analysis along with label-free quantification

All raw LC–MS/MS data were processed with FragPipe version (v22.0), incorporating

MSFragger (v4.1) for peptide identification. We applied the built in LFQ MBR workflow to

search against the publicly available Chlamydomonas v6.1 reference proteome database

supplemented with common contaminants and decoy sequences

(CreinhardtiiCC_4532_707_v6.1.protein_primaryTranscriptOnly.fa). MSFragger

performed a closed search using initial precursor and fragment mass tolerances of 0.5 Da,

followed by automatic mass recalibration and selection of narrower tolerances. The

enzyme was set to strict trypsin, allowing up to two missed cleavages, with peptide lengths

constrained to 7–50 residues and masses between 500 and 5,000 Da. Methionine

oxidation (+15.9949 Da) and protein N terminal acetylation (+42.0106 Da) were set as

variable modifications, while cysteine carbamidomethylation (+57.0215 Da) was fixed.

Subsequently, MSBooster, Percolator, and Philosopher were employed to generate deep

learning augmented scores, perform PSM rescoring, and estimate false discovery rates.

Label-free quantification was then carried out with IonQuant (v1.10.27) under default

settings, using a mass tolerance of 10 ppm and a retention time tolerance of 0.4 min. Both,

"match between runs" and "MaxLFQ" algorithms were enabled. The "min scans"

parameter was set to 3, "min isotopes" was set to 2, and "MaxLFQ min ions" was set to 2.

Proteins were considered differentially regulated when they exhibited a $|\log_2$ fold change$|$

≥ 0.5 and an adjusted P-value < 0.05.

Predictions of extracellular proteins

DeepLoc 2.1 (Odum et al., 2024) was used to predict the subcellular localization of

identified secreted proteins, with the organism type set to "Plant" to optimize predictions

for the Chlorophyte alga Chlamydomonas. DeepLoc 2.1 predicts localization up to ten

subcellular compartments: nucleus, cytoplasm, extracellular/secreted, mitochondrion, cell

membrane, endoplasmic reticulum, chloroplast, Golgi apparatus, lysosome/vacuole, and

peroxisome. Proteins were classified as secreted if they received an "extracellular"
prediction with a confidence score > 0.5.

Crude extracts and immunoblots

Crude extracts of soluble proteins (Zhao et al., 2004) were used for CRY-DASH1 detection
and total protein extracts for CHR1 detection (Vuong et al., 2025). Cells were harvested
at LD4. For tubulin and CK1 detection, proteins from the MMA fraction were used.
Antibodies against CRY-DASH1 (Rredhi et al., 2021), CHR1 (Vuong et al., 2025) and CK1
(Schulze et al., 2013) were used. Antibodies against acetylated tubulin were obtained from
Sigma-Aldrich (T7451). Immunoblots were performed as described previously (Vuong et
al., 2025). Briefly, proteins were separated by SDS-PAGE and transferred to
polyvinylidene difluoride (PVDF) membranes that were stained with Coomassie Brilliant
Blue R 250. The following dilutions of primary antibodies were taken: 1:5,000 for anti-
CRY-DASH1, 1:1,000 for anti-CHR1, 1:5,000 for CK1 and 1:10,000 for anti-acetylated
tubulin (T7451, Sigma-Aldrich). The horseradish peroxidase-conjugated anti-rabbit IgGs
(Sigma-Aldrich) were used as the secondary antibody or horseradish peroxidase-
conjugated anti-mouse IgGs (Sigma-Aldrich) for acetylated tubulin. Dilutions of the
secondary antibody were 1:5,000 (CHR1, CK1), 1:6,666 (CRY-DASH1) and 1:10,000
(acetylated tubulin).

Determination of flagellar length

Flagellar length was quantified on day 4 and day 7 of cell growth. 2 mL aliquots from the
main Chlamydomonas cultures were harvested and centrifuged at 500 g for 10 min. Ciliary
immunolocalization studies were done according to Zou et al. (2017). Cells were
incubated with a primary antibody against acetylated tubulin (Sigma, clone 6-11B-1,
mouse monoclonal), followed by a goat anti-mouse secondary antibody conjugated to
AlexaFluor 555 (Life Technologies, A32727) at a 1:2,000 dilution. Cilia were visualized
using Zeiss Axiophot and Leica Stellaris 8 Fluorescent Microscopes, respectively. Flagella
length was measured with ImageJ (version 1.53t, Schneider et al., 2012). Statistical
analyses and graphing were conducted in RStudio (v.2024.04.1, Posit team, 2023).

Motility analysis

Motility analysis of algal cells was performed using day 4 cultures grown in TAP medium at the different temperatures, whose cell density had been standardized to 5×10^5 cells mL^{-1}. A 120 µL aliquot was transferred onto a sterile glass microscope slide fitted with a Gene Frame adhesive chamber (Thermo Fisher Scientific, catalog no. AB0578). Prior to placing the Gene Frame adhesive chamber, slides were cleaned with ethanol and sterilized under UV light for 15 min. After the sample was added, the chamber was sealed with a coverslip that had also been pre-cleaned with ethanol and UV-sterilized. Live-cell imaging was conducted using a Nikon Ti2 widefield fluorescence microscope under a 4x objective. Temperature control during imaging was achieved using a cooling/heating Incubation Insert P-Set 2000 coupled to a DYNEO DD-310F refrigerated and heating circulator (JULABO GmbH, Germany). The imaging temperature was maintained at either 18 °C or 28 °C to match the cultivation conditions of each sample. Time-lapse recordings were acquired at intervals of 200 ms per frame. Image sequences were subsequently imported into ImageJ to perform brightness and contrast correction, background subtraction and image thresholding. This results in a series of binary images in which Chlamydomonas cells can be clearly distinguished from the background. Individual cell trajectories were tracked using the TrackMate plugin, which quantified key motility parameters including turning frequency and swimming speed. The extracted tracking data were then exported for statistical analysis and visualization. Trajectory plots were visualized in RStudio.

Gametogenesis and mating ability assay (Jiang and Stern, 2009)

Chlamydomonas vegetative cells of strains CC-125 (mt$^+$) and CC-124 (mt$^-$) were first transferred from liquid TAP cultures onto solid TAP agar plates and incubated for 4 days at either 18 °C or 28 °C under a 12 h:12 h LD cycle. Following this, cells were transferred onto N10 agar plates and incubated in the dark at their respective temperatures for three additional days to induce nitrogen starvation and promote pregamete formation. Following this, cells were harvested from the N10 plates and resuspended in 10 mL of sterile distilled water that had been acclimatized to the corresponding temperature. Cell suspensions were transferred to sterile NUNC flasks and placed under continuous intense illumination at 70 µmol m^{-2} s^{-1} for 2–3 h without agitation to induce gamete formation. Following

induction, mt^+ and mt^- gametes were adjusted to a final concentration of 2×10^6 cells mL^{-1}
using a hemocytometer. 1 mL from each mating type were combined in fresh NUNC
flasks and kept under continuous light without shaking to allow mating. Mating efficiency
was evaluated 1 h after mixing the gametes. Samples from each flask were collected and
counted using a light microscope. Large, fused cells were counted as single units. Mating
efficiency was calculated using the formula:

Mating efficiency (%) = $2 \times (1 - N_1 / N_0) \times 100$

In this equation, N_0 represents the number of cells per mL immediately after mixing, and
N_1 is the number of cells per mL measured 1 h after incubation (Chiang et al., 1970).

Accession numbers
Transcriptome data from this article can be found under the BioProject PRJNA1306621.
Motility videos are deposited to Zenodo under DOI: 10.5281/zenodo.16885208. Proteome
data was uploaded to PRIDE PXD067459.

Supplementary Data

Data Set 1:

Supplementary Figure 1. Boxplot analysis of cell sizes measured on day 7

Supplementary Figure 2. Temperature-dependent regulation of transcripts encoding TRP channels

Supplementary Figure 3. Growth curves of Chlamydomonas mono- and cocultures with *P. protegens* at 23 °C (A) and 33 °C (B)

Supplementary Figure 4. Quantification of flagellar length in cells grown at different temperatures on day 7

Supplementary Figure 5. Quantification of α-tubulin and CK1 protein abundance within isolated flagella (second independent replicate)

Supplementary Figure 6. Cell movements with mean directional change (A) and median swimming speed (second independent replicate)

Supplementary Figure 7. Mating efficiency at different temperatures (second independent dataset)

Data Set 2:

Supplementary Table 1. Values for growth kinetics of Chlamydomonas at different ambient temperatures compared to 23 °C

Supplementary Table 2. Values for cell sizes at different ambient temperatures compared to 18 °C

Supplementary Table 3. Transcript abundances altered in cells grown at different ambient temperatures

Supplementary Table 4. Full list of gene ontology (GO) terms

Supplementary Table 5. List of selected candidates along with the genome model numbers from Figures 2 to 6 and Supplementary Figure 2

Supplementary Table 6. List of identified proteins identified in the ciliary MMA fractions from cells grown at 18 °C or 28 °C

Supplementary Table 7. List of identified ciliary MMA proteins present in the ciliary database

Supplementary Table 8. List of temperature-dependent significantly up- and downregulated ciliary proteins

Supplementary Table 9. List of ciliary lengths of cells grown at different ambient temperatures compared to 18 °C

Supplementary Table 10. List of identified secretory proteins from cells grown at 18 °C or 28 °C

Supplementary Table 11. List of identified secretory proteins determined by DeepLoc

Supplementary Table 12. List of temperature-dependent significantly up- and downregulated secretory proteins

Video 1. Motility patterns of Chlamydomonas cells grown at 18 °C
Video 2. Motility patterns of Chlamydomonas cells grown at 28 °C

ACKNOWLEDGMENTS

We thank Eduardo Saluzzo and Valentina Mandini from Innova-tech-agro (Italy) for helpful discussions, and Aurelie Jost from the Microverse Imaging Center for help with the Leica Stellaris 8 Fluorescence Microscope. PS, RJA, CCP, ML and MM received funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) within the Germany's Excellence Strategy – EXC 2051 – Project ID 390713860. JC was funded by DFG under CRC 1127, Project ID 239748522. CL was supported by a PhD grant of the China Scholarship Council (CSC). ML is grateful for support by the research profile line LIFE of FSU Jena and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) via the Emmy-Noether-Program (Project ID 528114058). We thank the Microverse Imaging Center also financed by EXC 2051 for providing microscope facility support for data acquisition.

AUTHOR CONTRIBUTIONS

PS, RJA, ML and MM designed the project, PS, VW, DM, CCP, WL, CL, TV, JC and AZ performed experiments, and all authors contributed to analysis and interpretation of results. PS and MM wrote the paper with input from all coauthors.

REFERENCES

Aiyar, P, Schaeme, D, García-Altares, M, Carrasco Flores, D, Dathe, H, Hertweck, C, Sasso, S and Mittag, M (2017) Antagonistic bacteria disrupt calcium homeostasis and immobilize algal cells. Nature Communications, **8**: 1756

Barolo, L, Commault, AS, Abbriano, RM, Padula, MP, Kim, M, Kuzhiumparambil, U, Ralph, PJ and Pernice, M (2022) Unassembled cell wall proteins form aggregates in the extracellular space of *Chlamydomonas reinhardtii* strain UVM4. Applied Microbiology and Biotechnology, **106**: 4145-4156

Barrett, MR and Koch, AR (1982) Effects of ammonium sulfate and urea on the growth of Chlorophycean algae from rice fields. Journal of Environmental Quality, **11**: 187-191

Bösger, J, Wagner, V, Weisheit, W and Mittag, M (2009) Analysis of flagellar phosphoproteins from *Chlamydomonas reinhardtii*. Eukaryotic Cell, **8**: 922-932

Bösger, J, Wagner, V, Weisheit, W and Mittag, M (2014) Comparative phosphoproteomics to identify targets of the clock-relevant casein kinase 1 in *C. reinhardtii* flagella. Methods in Molecular Biology, **1158**: 187-202

Bray, NL, Pimentel, H, Melsted, P and Pachter, L (2016) Near-optimal probabilistic RNA-seq quantification. Nature Biotechnology, **34**: 525-527

Bunbury, F, Deery, E, Sayer, AP, Bhardwaj, V, Harrison, EL, Warren, MJ and Smith, AG (2022) Exploring the onset of B_{12}-based mutualisms using a recently evolved Chlamydomonas auxotroph and B_{12}-producing bacteria. Environmental Microbiology **24:** 3134-3147

Burgunter-Delamare, B, Shetty, P, Vuong, T and Mittag, M (2024) Exchange or eliminate: The secrets of algal-bacterial relationships. Plants, **13**: 829

Carrasco Flores, D, Hotter V, Vuong T, Hou Y, Bando Y, Scherlach K, Burgunter-Delamare B, Hermenau R, Komor AJ, Aiyar P *et al*. (2024) A mutualistic bacterium rescues a green alga from an antagonist. Proceedings of the National Academy of Sciences USA, **121**: e2401632121

Chen, S, Zhou, Y, Chen, Y and Gu, J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics, **34**: i884-i890

Chiang, KS, Kates, JR, Jones, RF and Sueoka, N (1970) On the formation of a homogeneous zygotic population in *Chlamydomonas reinhardtii*. Developmental Biology, **22**: 655-669

Chin, KJ, Lukow, T and Conrad, R (1999) Effect of temperature on structure and function of the methanogenic archaeal community in an anoxic rice field soil. Applied and Environmental Microbiology, **65**: 2341-2349

Christie, JM and Zurbriggen, MD (2021) Optogenetics in plants. New Phytologist, **229**: 3108-3115

Craig, RJ, Gallaher, SD., Shu, S, Salomé, PA, Jenkins, JW, Blaby-Haas, CE, Purvine, SO, O'Donnell, S, Barry, K, Grimwood, J, et al. (2023) The Chlamydomonas

Genome Project, version 6: Reference assemblies for mating-type plus and minus strains reveal extensive structural mutation in the laboratory. The Plant Cell, **35**: 644-672

Crockett, C (2021) How a microorganism corkscrews while it breaststrokes. Physics **14:** s21

Crozet, P, Navarro, FJ, Willmund, F, Mehrshahi, P, Bakowski, K, Lauersen, KJ, Pérez-Pérez, ME, Auroy, P, Gorchs Rovira, A, Sauret-Gueto, S and Niemeyer, J, (2018) Birth of a photosynthetic chassis: a MoClo toolkit enabling synthetic biology in the microalga Chlamydomonas reinhardtii. ACS Synthetic Biology, **7**: 2074-2086

Deisseroth, K and Hegemann, P (2017) The form and function of channelrhodopsin. Science, **357**: eaan5544

Dupuis, S and Merchant, SS (2023) *Chlamydomonas reinhardtii*: A model for photosynthesis and so much more. Nature Methods, **20**: 1441-1442

Ermilova, E (2020) Cold stress response: an overview in Chlamydomonas. Frontiers in Plant Science, **11**: 569437

Folcik, AM, Cutshaw, K, Haire, T, Goode, J, Shah, P, Zaidi, F, Richardson, B and Palmer, A (2020) Quorum sensing behavior in the model unicellular eukaryote *Chlamydomonas reinhardtii*. iScience **23:** e101835

García-García, A, Cuesta-Valero, FJ, Miralles, DG, Mahecha, MD, Quaas, J, Reichstein, M, Zscheischler, J and Peng, J (2023) Soil heat extremes can outpace air temperature extremes. Nature Climate Change, **13**: 1237-1241

Ghasemi Y, Rasoul-Amini S, Morowvat MH, Raee MJ, Ghoshoon MB, Nouri F, Negintaji N, Parvizi R, Mosavi-Azam SB. (2008) Characterization of hydrocortisone biometabolites and 18S rRNA gene in *Chlamydomonas reinhardtii* cultures. Molecules, **13**: 2416-2425

Greiner, A, Kelterborn, S, Evers, H, Kreimer, G, Sizova, I and Hegemann, P (2017) Targeting of photoreceptor genes in *Chlamydomonas reinhardtii* via zinc-finger nucleases and CRISPR/Cas9. The Plant Cell, **29**: 2498-2518

Hallmann, A (2006) The pherophorins: common, versatile building blocks in the evolution of extracellular matrix architecture in Volvocales. Plant Journal, **45**: 292-307

Harris, EH (1989) The Chlamydomonas sourcebook: a comprehensive guide to biology and laboratory use. Academic Press, Cambridge

Helliwell, K.E., Collins, S., Kazamia, E., Purton, S., Wheeler, G.L., and Smith, A.G. (2015) Fundamental shift in vitamin B_{12} eco-physiology of a model alga demonstrated by experimental evolution. ISME J. **9:** 1446-1455.

Hey, S and Linder, S (2024) Matrix metalloproteinases at a glance. Journal of Cell Science, **137**: jcs261898

Hou, Y, Bando, Y., Carrasco Flores, D., Hotter, V., Das, R., Schiweck, B., Melzer, T., Arndt, H.D. and Mittag, M (2023) A cyclic lipopeptide produced by an antagonistic bacterium relies on its tail and transient receptor potential-type Ca^{2+} channels to immobilize a green alga. New Phytologist, **237**: 1620-1635

Hughes, CS, Moggridge, S, Müller, T, Sorensen, PH, Morin, GB and Krijgsveld, J (2019) Single-pot, solid-phase-enhanced sample preparation for proteomics experiments. Nature Protocols, **14**: 68-85

Jiang, X and Stern, D (2009) Mating and tetrad separation of *Chlamydomonas reinhardtii* for genetic analysis. Journal of Visualized Experiments, **30**: 1274

Katoh, Y, Terada, M, Nishijima, Y, Takei, R, Nozaki, S, Hamada, H and Nakayama, K (2016) Overall architecture of the intraflagellar transport (IFT)-B complex containing Cluap1/IFT38 as an essential component of the IFT-B peripheral subcomplex. Journal of Biological Chemistry, **291**: 10962-10975

Kinoshita, T, Fukuzawa, H, Shimada, T, Saito, T and Matsuda, Y (1992) Primary structure and expression of a gamete lytic enzyme in *Chlamydomonas reinhardtii*: similarity of functional domains to matrix metalloproteases. Proceedings of the National Academy of Sciences, **89**: 4693-4697

Lacey, SE, Graziadei, A and Pigino, G (2024) Extensive structural rearrangement of intraflagellar transport trains underpins bidirectional cargo transport. Cell, **187**: 4621-4636

Lechtreck, KF, Johnson, EC, Sakai, T, Cochran, D, Ballif, BA, Rush, J, Pazour, GJ, Ikebe, M and Witman, GB (2009) The *Chlamydomonas reinhardtii* BBSome is an IFT cargo required for export of specific signaling proteins from flagella. Journal of Cell Biology, **187**: 1117-1132

Légeret, B, Schulz-Raffelt, M, Nguyen, HM, Auroy, P, Beisson, F, Peltier, G, Blanc, G, Li-Beisson, Y (2016) Lipidomic and transcriptomic analyses of *Chlamydomonas reinhardtii* under heat stress unveil a direct route for the conversion of membrane lipids into storage lipids. Plant, Cell & Environment, **39**: 834-847

L'Hernault, SW and Rosenbaum, JL (1985) Chlamydomonas α-tubulin is posttranslationally modified by acetylation on the ε-amino group of a lysine. Biochemistry, **24**: 473-478

Li, W, Flores, DC, Füßel, J, Euteneuer, J, Dathe, H, Zou, Y, Weisheit, W, Wagner, V, Petersen, J and Mittag, M (2018) A musashi splice variant and its interaction partners influence temperature acclimation in Chlamydomonas. Plant Physiology, **178**: 1489-1506

Li, X, Patena, W., Fauser, F., Jinkerson, R. E., Saroussi, S., Meyer, M. T., Ivanova, N., Robertson, J. M., Yue, R., Zhang, R., et al. (2019) A genome-wide algal mutant library and functional screen identifies genes required for eukaryotic photosynthesis. Nature Genetics, **51**: 627-635

Lin, CS, Chou, TL and Wu, JT (2013) Biodiversity of soil algae in the farmlands of mid-Taiwan. Botanical Studies, **54**: 1-12

Liu, P and Lechtreck, KF (2018) The Bardet–Biedl syndrome protein complex is an adapter expanding the cargo range of intraflagellar transport trains for ciliary export. Proceedings of the National Academy of Sciences, **115**: E934-E943

Love, MI, Huber, W and Anders, S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology, **15**: 1-21

Luxmi, R, Blaby-Haas, C, Kumar, D, Rauniyar, N, King, SM, Mains, RE and Eipper, BA (2018) Proteases shape the Chlamydomonas secretome: comparison to classical neuropeptide processing machinery. Proteomes, **6**: 36

Ma, Y, He, J, Li, S, Yao, D, Huang, C, Wu, J and Lei, M (2023) Structural insight into the intraflagellar transport complex IFT-A and its assembly in the anterograde IFT train. Nature Communications, **14**: 1506

Mäkinen, J, Ellis, EE, Antão, LH, Davrinche, A, Laine, AL, Saastamoinen, M, Conenna, I, Hällfors, M, Santangeli, A, Kaarlejärvi, E, Heliölä, J et al. (2025) Thermal homogenization of boreal communities in response to climate warming. Proc Natl Acad Sci USA, **122:** e2415260122**.**

Marshall, WF (2024) Chlamydomonas as a model system to study cilia and flagella using genetics, biochemistry, and microscopy. Frontiers in Cell and Developmental Biology, **12**: 1412641

Marshall, WF and Rosenbaum, JL (2001) Intraflagellar transport balances continuous turnover of outer doublet microtubules: implications for flagellar length control. Journal of Cell Biology, **155**: 405-414

McCafferty, CL, Papoulas, O, Jordan, MA, Hoogerbrugge, G, Nichols, C, Pigino, G, Taylor, DW, Wallingford, JB and Marcotte, EM (2022) Integrative modeling reveals the molecular architecture of the intraflagellar transport A (IFT-A) complex. Elife, **11**: e81977

McGoldrick, LL, Singh, AK, Demirkhanyan, L, Lin, TY, Casner, RG, Zakharian, E and Sobolevsky, AI (2019) Structure of the thermo-sensitive TRP channel TRP1 from the alga *Chlamydomonas reinhardtii*. Nature Communications **10:** 4180

Meier, HS, Schuman, IJ, Layden, TJ, Ritz, A, Kremer, CT and Fey, SB (2022) Temperature-mediated transgenerational plasticity influences movement behaviour in the green algae *Chlamydomonas reinhardtii*. Functional Ecology, **36**: 2969-2982

Merchant, S. S., Prochnik, S. E., Vallon, O., Harris, E. H., Karpowicz, S. J., Witman, G. B., Terry, A., Salamov, A., Fritz-Laylin, L. K., Maréchal-Drouard, et al. (2007) The Chlamydomonas genome reveals the evolution of key animal and plant functions. Science, **318**: 245-250

Mitchell, BF, Pedersen, LB, Feely, M, Rosenbaum, JL and Mitchell, DR (2005) ATP production in *Chlamydomonas reinhardtii* flagella by glycolytic enzymes. Molecular Biology of the Cell, **16**: 4509-4518

Moparthi, L, Sinica, V, Moparthi, VK, Kreir, M, Vignane, T, Filipovic, MR, Vlachova, V and Zygmunt, PM (2022) The human TRPA1 intrinsic cold and heat sensitivity involves separate channel structures beyond the N-ARD domain. Nature Communications **13:** 6113

Müller, N, Wenzel, S, Zou, Y, Künzel, S, Sasso, S, Weiß, D, Prager, K, Grossman, A, Kottke, T and Mittag, M (2017) A plant cryptochrome controls key features of the Chlamydomonas circadian clock and its life cycle. Plant Physiology, **174**: 185-201

Murakami, H, Nobusawa, T, Hori, K, Shimojima, M and Ohta, H (2018) Betaine lipid is crucial for adapting to low temperature and phosphate deficiency in *Nannochloropsis*. Plant Physiology **177:** 181-193

Odum, M.T., Teufel, F., Thumuluri, V., Almagro Armenteros, J.J., Johansen, A.R., Winther, O., and Nielsen, H. (2024). DeepLoc 2.1: multi-label membrane protein type prediction using protein language models. Nucleic Acids Res. **52:** W215-W220.

Pan, J and Snell, WJ (2005) Chlamydomonas shortens its flagella by activating axonemal disassembly, stimulating IFT particle trafficking, and blocking anterograde cargo loading. Developmental Cell, **9**: 431-438

Patel, MB, Griffin, PJ, Olson, SF, Dai, J, Hou, Y, Malik, T, Das, P, Zhang, G, Zhao, W, Witman, GB and Lechtreck, KF, (2024) Distribution and bulk flow analyses of the intraflagellar transport (IFT) motor kinesin-2 support an "on-demand" model for Chlamydomonas ciliary length control. Cytoskeleton, **81**: 586-604

Pazour, GJ, Agrin, N, Leszyk, J and Witman, GB (2005) Proteomic analysis of a eukaryotic cilium. Journal of Cell Biology, **170**: 103-113

Pedersen LB, Geimer S, Rosenbaum JL (2006) Dissecting the molecular mechanisms of intraflagellar transport in Chlamydomonas. Current Biology **16:** 450-459

Riekhof, WR, Sears, BB and Benning, C (2005) Annotation of genes involved in glycerolipid biosynthesis in *Chlamydomonas reinhardtii*: discovery of the betaine lipid synthase BTA1Cr. Eukaryotic Cell **4:** 242-252

Rredhi A, Petersen J, Wagner V, Vuong T, Li W, Li W, Schrader L, Mittag M (2024) The UV-A receptor CRY-DASH1 up- and downregulates proteins involved in different plastidial pathways. Journal of Molecular Biology, **436**: 168271

Rredhi, A, Petersen, J, Schubert, M, Li, W, Oldemeyer, S, Li, W, Westermann, M, Wagner, V, Kottke, T and Mittag, M (2021) DASH cryptochrome 1, a UV-A receptor, balances the photosynthetic machinery of *Chlamydomonas reinhardtii*. New Phytologist, **232**: 610-624

Salomon, S, Schilling, M, Albrieux, C, Si Larbi, G, Jouneau, PH, Roy, S, Falconet, D, Michaud, M and Jouhet, J (2025) Betaine lipids overproduced in seed plants are not imported into plastid membranes and promote endomembrane expansion. Journal of Experimental Botany **76:** 980-996

Sasso, S, Stibor, H, Mittag, M and Grossman, AR (2018) The natural history of model organisms from molecular manipulation of domesticated *Chlamydomonas reinhardtii* to survival in nature. eLife, **7**: e39233

Scaife, MA and Smith, AG (2016) Towards developing algal synthetic biology. Biochemical Society Transactions, **44**: 716-722

Schmidt, M, Geßner, G, Luff, M, Heiland, I, Wagner, V, Kaminski, M, Geimer, S, Eitzinger, N, Reißenweber, T, Voytsekh, O and Fiedler, M (2006) Proteomic analysis

of the eyespot of *Chlamydomonas reinhardtii* provides novel insights into its components and tactic movements. The Plant Cell, **18**: 1908-1930

Schneider, CA, Rasband, WS and Eliceiri, KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nature Methods, **9**: 671-675

Schroda, M, Hemme, D and Mühlhaus, T (2015) The Chlamydomonas heat stress response. The Plant Journal, **82**: 466-480

Schulze, T, Schreiber, S, Iliev, D, Boesger, J, Trippens, J, Kreimer, G and Mittag, M (2013) The heme-binding protein SOUL3 of *Chlamydomonas reinhardtii* influences size and position of the eyespot. Molecular Plant, **6**: 931-944

Sharma, A, Samtani, H and Laxmi, A (2024b) Molecular dialogue between light and temperature signaling in plants: From perception to thermotolerance. Journal of Experimental Botany, **75**: erae356

Sharma, S, Sushmita, K, Singh, R, Sanyal, SK and Kateriya, S (2024a) Phototropin localization and interactions regulates photophysiological processes in Chlamydomonas reinhardtii. bioRxiv, doi:10.1101/2024.12.27.630506

Snell, WJ, Eskue, WA and Buchanan, MJ (1989) Regulated secretion of a serine protease that activates an extracellular matrix-degrading metalloprotease during fertilization in Chlamydomonas. Journal of Cell Biology, **109**: 1689-1694

Soneson, C, Love, MI and Robinson, MD (2015) Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. F1000Research, **4**: 1521

Szydlowski L, Sorokin A, Vasieva O, Boerner S, Fedorovich V, Goryanin I (2019). Evolutionary dynamics of microbial communities in bioelectrochemical systems. Journal of Computer Science and Systems Biology **13**: 1-9.

Tuxhorn, J, Daise, T and Dentler, WL (1998) Regulation of flagellar length in Chlamydomonas. Cell Motility and the Cytoskeleton, **40**: 133-146

Valledor, L, Furuhashi, T, Hanak, AM and Weckwerth, W (2013) Systemic cold stress adaptation of *Chlamydomonas reinhardtii*. Molecular & Cellular Proteomics, **12**: 2032-2047

Verret, F., Wheeler, G., Taylor, A.R., Farnham, G., and Brownlee, C. (2010) Calcium channels in photosynthetic eukaryotes: implications for evolution of calcium-based signalling. New Phytol. **187:** 23-43.

von der Heyde, EL and Hallmann, A (2022) Molecular and cellular dynamics of early embryonic cell divisions in Volvox carteri. The Plant Cell, **34**:1326-1353

Vuong, T, Shetty, P, Kurtoglu, E, Schultz, C, Schrader, L, Then, P, Petersen, J, Westermann, M, Rredhi, A., Chowdhury, S., Mukherji, R., et al. (2025) Metamorphosis of a unicellular green alga in the presence of acetate and a spatially structured three-dimensional environment. New Phytologist, **245**: 1180-1196

Wemmer, K, Ludington, W and Marshall, WF (2020) Testing the role of intraflagellar transport in flagellar length control using length-altering mutants of Chlamydomonas. Philosophical Transactions of the Royal Society B, **375**: 20190159

Xiao, Y., Xin, Y., Shen, K., Yang, M., and Wu, H. (2025). The roles of transient receptor potential vanilloid in autoimmune diseases. Autoimmun. Rev. **24:** 103872.

Zhang, Y, Wolf-Yadlin, A, Ross, PL, Pappin, DJ, Rush, J, Lauffenburger, DA and White, FM (2005) Time-resolved mass spectrometry of tyrosine phosphorylation sites in the epidermal growth factor receptor signaling network reveals dynamic modules. Molecular & Cellular Proteomics, **4**: 1240-1250

Zhang, N, Mattoon, EM, McHargue, W, Venn, B, Zimmer, D, Pecani, K, Jeong, J, Anderson, CM, Chen, C, Berry, JC, Xia, M (2022) Systems-wide analysis revealed shared and unique responses to moderate and acute high temperatures in the green alga *Chlamydomonas reinhardtii*. Communications Biology, **5**: 460

Zhang, N, Venn, B, Bailey, CE, Xia, M, Mattoon, EM, Mühlhaus, T and Zhang, R (2024) Moderate high temperature is beneficial or detrimental depending on carbon availability in the green alga *Chlamydomonas reinhardtii*. Journal of Experimental Botany, **75**: 979-1003

Zhao, B, Schneid, C, Iliev, D, Schmidt, EM, Wagner, V, Wollnik, F and Mittag, M (2004) The circadian RNA-binding protein CHLAMY 1 represents a novel type heteromer of RNA recognition motif and lysine homology domain-containing subunits. Eukaryotic Cell, **3**: 815-825

Zou, Y, Wenzel, S, Müller, N, Prager, K, Jung, EM, Kothe, E, Kottke, T and Mittag, M (2017) An animal-like cryptochrome controls the Chlamydomonas sexual cycle. Plant Physiology, **174**: 1334-1347

1
2
3 **Figure 1: Influence of different ambient cultivation temperatures on the growth and**
4 **transcript abundances of** Chlamydomonas*.* **A)** Growth curve of Chlamydomonas
5 cultured at 18 °C (blue), 23 °C (black), 28 °C (green), and 33 °C (red) over a 15-day period.

Data points represent mean cell densities (cells mL^{-1}) from two independent replicates, each comprising three biological replicates. Error bars indicate standard deviation. Asterisks indicate the statistically significant differences for 18 °C (black), 28 °C (green), and 33 °C (red) compared to 23 °C as calculated by Student's *t*-test (*: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$). **B)** Boxplot of vertical cell size measurements (µm) on day 4 of cultivation across the four temperature treatments. The cell sizes of day 7 are shown in Supplementary Figure 1. Each box represents measurements from three biological replicates (100 cells per replicate). Statistical significance between groups is indicated by asterisks. Asterisks represent significant differences as calculated by Student's *t*-test (****: $P < 0.0001$). **C)** Principal component analysis (PCA) of transcriptomic profiles from Chlamydomonas after 4 days of cultivation at each temperature. Each point corresponds to a biological replicate, colored by temperature (18 °C: blue, 23 °C: black, 28 °C: green, 33 °C: red). PC1 and PC2 account for 50% and 40% of the total variance, respectively, illustrating clear separation by temperature, especially at the extremes. **D–E)** Volcano plots depicting differentially abundant transcripts between 18 °C and 28 °C (D), and 18 °C and 33 °C (E), respectively. Transcripts with significant upregulation (orange) and downregulation (blue) are defined by adjusted *P*-value < 0.01 and |Log$_2$ fold change| ≥ 1. Non-significantly changed transcripts are shown in gray. Total transcript counts for each category are indicated in parentheses. Vertical dashed lines mark log$_2$ fold change thresholds of ±1; the horizontal dashed line marks the significance threshold. **F)** Heatmap showing the expression patterns of the 1,000 most differentially expressed transcripts between 18 °C and 28 °C, clustered across all temperature conditions. Each row represents a gene, and each column a temperature. Transcripts are grouped into clusters based on shared expression patterns, highlighting subsets with temperature-specific regulation. **G)** Exemplary results from a Gene Ontology (GO) enrichment analysis of differentially expressed genes, comparing 18 °C and 28 °C. The full list is shown in Supplementary Table 4. Bar plots display the percentage of enriched GO terms in Biological Process, Cellular Component, and Molecular Function categories. Orange bars indicate GO terms enriched at 28 °C relative to 18 °C, while blue bars represent downregulated terms.

38
39
40 **Figure 2: Temperature-dependent regulation of transcripts encoding selected TRP**
41 **channels and photoreceptors along with mitigation of bacterial antagonism. A)** $\log_2$

fold changes in the expression of selected *TRP* channel transcripts. A list of all significantly changed TRP channel transcripts with a $|\log_2 FC| \geq 1$ in Chlamydomonas in response to cultivation at varying temperatures (18 °C, 28 °C, 33 °C) is presented in Supplementary Figure 2 and Supplementary Table 5. TRPs 5, 11 and 22 are known to be involved in the orfamide A triggered signaling response (Hou et al., 2023). **B)** Visual evidence indicating the attenuation of bacterial antagonism during cocultivation of Chlamydomonas (abbreviated as *Cre*) with *P. protegens* (*Ppr)* at higher temperatures. Green chlorophyll coloration at 28 °C and 33 °C denotes improved algal viability compared to 18 °C, indicating that increased temperature diminishes bacterial antagonistic effects. **C, D)** Growth curves of axenic Chlamydomonas (monocultures) and *Cre–Ppr* cocultures at 18 °C and 28 °C, respectively. Cell densities (cells mL^{-1}) were measured over 15 days, highlighting temperature-associated changes in antagonism. Data points represent mean ± standard deviation cell densities from two independent replicates, each comprising three biological replicates. Statistical significance is indicated by asterisks. **E)** $\log_2$ fold changes ($|\log_2 FC| \geq 1$) in the expression of photoreceptor transcripts (Supplementary Table 5) under varying temperatures. Known domains of the photoreceptors analyzed by Phytozome 14 (https://phytozome-next.jgi.doe.gov/) are highlighted (Greiner et al., 2017). For simplicity, the size of the proteins has been standardized to a common bar length; the relative position of the domains is given. The bar size for pCRY with more than 1,000 amino acids has been enlarged. CHR, channelrhodopsin; aCRY, animal-like cryptochrome; pCRY, plant cryptochrome; CRY-DASH, *Drosophila Arabidopsis Synechocystis* human cryptochrome. **F)** Immunoblot analysis quantifying CHR1 protein abundances in cells grown at 18 °C and 28 °C. 75 µg of total protein per lane were separated by SDS-PAGE and immunoblotted with the respective antibodies as described in the Methods section. n = 3 biological replicates. Arrows indicate the positions of the target protein. Selected bands from a polyvinylidene difluoride membrane stained with Coomassie Brilliant Blue R250 were used as a loading control (LC). **G)** Immunoblot analysis quantifying CRY-DASH1 under similar conditions as in panel F. In this case, 30 µg of soluble proteins per lane were separated on SDS-PAGE. (**C, D, F, G**) Asterisks represent significant differences as calculated by Student's *t*-test. $P < 0.05$ (*), $P < 0.01$ (**), $P < 0.001$ (***), $P < 0.0001$ (****). (**A, E**) Adjusted *P*-values < 0.00001 (****), as calculated by DESeq2, are shown.

Figure 3: Temperature-dependent regulation of transcripts encoding IFT and extracellular proteins, enzymes of betaine lipid synthesis and RNA-binding proteins. A) Expression of transcripts encoding IFT proteins. Transcripts encoding IFT-A and IFT-B complex with a |log$_2$ FC| ≥ 1 show significantly higher abundance at temperatures of 28 °C and 33 °C. **B)** Expression of transcripts with a |log$_2$ FC| ≥ 1 encoding predicted extracellular proteins as identified by DeepLoc (see Methods) in cells grown at different temperatures. The first seven transcripts encode hydroxyproline-rich glycoproteins. CWP2, cell wall protein2, also known as FAP3; FAP, flagellar associated protein; ISG2/6, inversion-specific glycoprotein, found in cilia; MAW4, membrane associated wall protein; VSP3, serine-proline repeat hydroxyproline-rich cell wall protein found in cilia. **C)** Expression of transcripts with a |log$_2$ FC| ≥ 1 encoding enzymes for betaine lipid biosynthesis. SAS1 (S-adenosylmethionine synthetase1), SAMDC (S-adenosyl-L-methionine decarboxylase), BTA1 (betaine lipid synthase). **D)** Expression profiles of

transcripts encoding RNA-binding proteins, including Pumilio/PUF domain proteins, Musashi, CRB1/3 proteins being the C1 and C3 subunit of CHLAMY1, NAB1 (nucleic acid binding factor) and DUS16 (dull slicer-16). Known domains of the RBPs analyzed by Phytozome 14 (https://phytozome-next.jgi.doe.gov/) are highlighted. For simplicity, the size of the proteins has been standardized to a common bar length; the relative position of the domains is given. (**A-E**) Proteins are listed in Supplementary Table 5 along with their Cre accession numbers. Adjusted *P*-values < 0.01 (*), < 0.001 (**) or < 0.0001 (***) or < 0.00001 (****), as calculated by DESeq2, are shown.

100
101

Figure 4: Temperature-dependent remodeling of flagellar proteins alters the length of flagella. A) Volcano plot depicting differentially abundant proteins in the flagellar proteome of Chlamydomonas cultivated at 18 °C and 28 °C. Points represent individual proteins. The orange color indicates proteins upregulated at 28 °C, blue color indicates proteins downregulated at 28 °C, and black represents non-significant changes (adjusted P-value < 0.05 and $\log_2$ fold change threshold). Numbers in parentheses denote protein counts per category. Dashed vertical lines indicate $\pm \log_2$ fold change of 0.5; the horizontal dashed line marks the significance threshold. **B)** Barplot showing $\log_2$ fold change in IFT-A and IFT-B complex proteins that were significantly differentially abundant in cells grown at 28 °C compared to 18 °C (adjusted P-value < 0.05 and $|\log_2$ fold change| ≥ 0.5). Orange bars indicate upregulation in IFT component abundance, suggesting temperature-dependent modulations in the IFT machinery. **C)** Representative immunolocalization microscopy images of Chlamydomonas cells cultivated at either 18 °C or 28 °C. Left panels show brightfield microscopy images of cells and cilia, right panels depict immunofluorescence images of cilia stained with anti-acetylated tubulin antibodies (see the Methods section). **D)** Quantification of flagellar length in cells grown at 18 °C, 23 °C, 28 °C, and 33 °C on day 4, measured using immunolocalization images. Each boxplot

includes measurements from three biological replicates (50 flagella per replicate; n = 150
per temperature). Significant differences between temperature treatments were
determined by Student's *t*-test and are marked by asterisks (n.s.: not significant; ****:
P < 0.0001). **E)** Upper panel: Quantification of α-tubulin and CK1 protein abundance within
isolated flagella, shown as a barplot from proteome data. Lower panels: As indicated, 0.25
µg and 20 µg of proteins per lane from the MMA flagellar fraction isolated from cells grown
at 18 °C and 28 °C, respectively, were separated by SDS-PAGE and immunoblotted with
(i) anti-acetylated tubulin or (ii) anti-CK1 antibodies as described in the Methods section.
An independent biological replicate is shown in Supplementary Figure 5.
128

A

Motor proteins

Cilia formation and ciliopathies

Zinc chaperones

■ Upregulated at 28 °C compared to 18 °C   ■ Downregulated at 28 °C compared to 18 °C

B 18 °C
Few turns between movements

C 28 °C
Many turns between movements

D

E

● Starting position of both cells

Figure 5: Remodelling of ciliary proteins and increases in the frequency of algal cell rotations at higher ambient temperature. A) Log$_2$ fold change of selected flagellar proteins from multiple functional groups in Chlamydomonas compared between cultivation temperatures of 18 °C and 28 °C. Motor proteins (e.g., dyneins, kinesins), ciliopathy-related proteins (BBS proteins, ankyrins), and zinc chaperones display significant upregulation or downregulation, as indicated by colored bars. Orange bars denote proteins upregulated at 28 °C relative to 18 °C, while blue bars indicate downregulation. Statistically significant proteins are marked by asterisks. Adjusted *P*-values < 0.05 (*) and < 0.01 (**) are shown, as calculated by FragPipe. **B-C)** Motility paths tracked for a representative Chlamydomonas cell cultivated either at 18 °C (B) or at 28 °C (C). The first 80 s of the movement are depicted. Please note that the x-axes in B and C show different scales. Cells at 18 °C typically exhibit fewer turns between movements, resulting in relatively straight trajectories. In contrast, cells at 28 °C display frequent directional changes, producing more curved or meandering movement paths compared to 18 °C. **D)** Quantification of mean directional change (degrees) between frames (i.e. at 200 ms intervals) for cells grown at the different temperatures. Data reflects a minimum of 150 individual tracks per temperature per biological replicate, with two biological replicates in total. Statistical significance determined by Student's *t*-test is indicated by asterisks (****: *P* < 0.0001). A second independent biological replicate is shown in Supplementary Figure 6A. **E)** Mean swimming speed (µm/sec) for cells at each temperature, calculated from tracked cell movements. This is calculated by computing the mean swimming speed along each trajectory (in the x-y plane) then averaging this value over trajectories. Data reflects a minimum of 150 individual tracks per temperature per biological replicate, with two biological replicates in total. Statistical significance determined by Student's *t*-test is indicated by asterisks (****: *P* < 0.0001). A second independent biological replicate is shown in Supplementary Figure 6B.

Figure 6: Temperature effects on the secretome, extracellular proteins, and mating efficiency in Chlamydomonas A) Volcano plot depicting differentially abundant proteins in the Chlamydomonas secretome cultivated under 18 °C and 28 °C. Each point represents a unique protein; orange and blue indicate significant up- or downregulation. Axes indicate $\log_2$ fold change versus $-\log_{10}$ (adjusted P-value); numbers in parentheses denote protein counts per category. **B)** Barplot showing the $\log_2$ fold changes in abundance of key extracellular proteins as predicted by DeepLoc in the secretome. Notably, the protein VLE1 was detected among soluble mating secretory proteins in previous studies (Luxmi et al., 2018) but was not predicted as extracellular by DeepLoc. Other proteins detected in the soluble mating secretome are highlighted with an asterix and in bold. Adjusted P-values < 0.05 (*), < 0.01 (**), < 0.001 (***) and < 0.0001 (****) are shown, as calculated by FragPipe. **C)** Quantification of mating efficiency across temperature or experimental conditions. Data represents three biological replicates. The blots show the proportion of successfully fused mating pairs, with mean values marked. Significant differences in mating efficiency between groups were determined using Student's t-test; P-values are indicated (**: P < 0.01). A second independent dataset with three biological replicates is shown in Supplementary Figure 7.