Quenching of Chaos in externally driven metacommunities

Anusree Vinod Kunnath[a], Dweepabiswa Bagchi[a]

[a]*Trecent Systems,*
[b]*School of Physics, Indian Institute of Science Education and Research,*
Thiruvananthapuram-695 551, Kerala, India,

Abstract

The chaotic population dynamics of ecosystems across diverse spatial scales have been extensively investigated over the past three decades. Chaotic fluctuations in species populations, except under specific conditions, have generally been shown to increase extinction risk through cascading effects, environmental noise, and frequent periods of low species density. The persistence of chaotic ecosystems has been attributed to internal and external mechanisms of chaos control, which may be mediated by the ecosystems themselves or by environmental factors such as habitat heterogeneity. However, the control and particularly the quenching of chaos in metacommunities resulting from internal habitat heterogeneity remain unexplored. This study addresses this gap by analyzing the dynamics of a metacommunity composed of five chaotic ecosystems, each located in a distinct patch. The first patch functions as the drive, exerting external influence on the dynamics of the remaining four response patches. Each patch contains an inherently chaotic tritrophic food web, with habitat heterogeneity present among patches. Results indicate that in a drive-response metacommunity, strong influence from the drive quenches dynamical chaos in both drive and response patches, often leading to steady states. This phenomenon is observed in two distinct metacommunity network structures. The heterogeneity of the response systems and the dissimilarity between drive and response systems are found to play significant roles in suppressing chaotic population dynamics. These findings strongly imply that the persistence of inherently chaotic metacommunities

---

*Corresponding author
Email address:* `dweepabiswabagchi20@iisertvm.ac.in` (Dweepabiswa Bagchi)

may result from the quenching of chaos through the interplay of chaos, habitat heterogeneity, and network structure shaped by dispersal. Furthermore, metacommunities composed of inherently chaotic patches are also susceptible to extinction due to dispersal-induced synchronization. Accordingly, this study also investigates dispersal-induced complete synchronization among the constituent patches of the drive-response metacommunities.

Keywords:

1. Introduction

Chaotic dynamics have been reported in several population models describing the evolution of a plethora of natural ecosystems (Hastings and Powell (1991); Blasius et al. (1999); Upadhyay et al. (1998)). Most importantly, the extinction risk and the persistence of such inherently chaotic ecosystems have also been investigated in some detail. Several studies have associated chaotic fluctuation in population density with a low degree of persistence. Some studies revealed that ecosystems in dynamically chaotic states have a high probability of extinction, either through cascading effects or a synchronized response to external environmental noise (Williams and Martineza (2004); Boccaletti et al. (2000)). In particular, it has been shown that low-density populations have an elevated chance of extinction in the chaotic domain of phase space (Berryman and Millstein (1989)). This often leads to population extinctions under stochasticity (Thomas et al. (1980); Hastings (1982); Croteau (2010)). Additionally, chaotic dynamics might mask early warning signals and indirectly increase extinction risk (Krkosˇek and Drake (2014)). Further, Dietze (Dietze (2018)) reported that truly chaotic populations would fluctuate wildly, resulting in frequent population crashes, genetic bottlenecks, and a high risk of stochastic extinction.

Several studies (Jørgensen and Jørgensen (2002); Berryman and Millstein (1989); Doebeli (1993)) have asserted that natural selection constrains ecosystems from adopting parameter values that would induce chaotic population dynamics. Consequently, such ecosystems are thought to avoid chaotic behavior. Additionally, ecological mechanisms such as toxin release (Mandal et al. (2006)), omnivory (McCann and Hastings (1997)), and environmental factors like habitat heterogeneity (Krkosˇek and Drake (2014); Hilborn et al. (2003)) have been reported to regulate or suppress chaos in ecosystem

population dynamics (Doebeli (1993); Gamarra et al. (2001)). External interventions may also influence these dynamics. However, Solé and Gamarra (Solé and Gamarra (1998)) demonstrated that the chaotic domain is extensive for a metacommunity of chaotic ecosystems linked by dispersal. Within this domain, some regions exhibit population fluctuations that increase extinction risk, while others display high-amplitude species density fluctuations associated with lower extinction risk. Furthermore, strong dispersal can increase the extinction risk of the metacommunity by shifting it into the chaotic domain characterized by low fluctuations (Solé and Gamarra (1998)). In contrast, Allen et al. (Allen et al. (1993)) found that chaos in a weakly coupled metapopulation can have a decorrelating effect, thereby enhancing persistence. These findings indicate that chaotic dynamics may reduce persistence in a metacommunity, depending on the strength of inter-patch coupling (dispersal). The contrasting evidence suggests that the influence of dispersal on the persistence of metacommunities with chaotic population dynamics warrants further investigation.

Motivated by the above, this study examines the quenching of chaos in a metacommunity influenced by external invasionary dispersal. The metacommunity model consists of patches, each characterized by inherently chaotic food webs that describe the species dynamics within each patch. The structure we consider for our metacommunity is that of a drive-response network, for e.g., a river-distributary ecosystem. In such a network, the river controls the water level, nutrient level, and soil level of the distributaries (Larsen et al. (2021)), thus effectively controlling their habitat quality (Sarremejane et al. (2021); J. D. Yeakel and deAguiar (2014)) and 'driving' their dynamics. To ensure that our model adheres to real metacommunities, we consider an appreciable degree of heterogeneity between drive and response patches. We demonstrate how habitat heterogeneity in such metacommunities quenches the chaos in each patch. Further, we establish that the complete synchronization between homogeneous drive and response systems is absolute, but decreases considerably with increasing dispersal in the case of heterogeneous drive and response. We also propose that our model could also be used as an approach to understand the effect of invasion in impeding chaos in the metacommunity.

2. Models

In our study, both the drive and response patches contain of a tri-trophic, inherently chaotic foodweb described by the Blasius model (Blasius et al. (1999)). The basic equations of the Blasius model is given as

$$\dot{X} = aX - \alpha_1 \frac{XY}{1 + kX}, \tag{1a}$$

$$\dot{Y} = -bY + \alpha_1 \frac{XY}{1 + kX} - \alpha_2 YZ \tag{1b}$$

$$\dot{Z} = -c(Z - Z_s) + \alpha_2 YZ, \tag{1c}$$

, where the parameter a represents the growth rate of the resource in absence of inter-species interactions, while the parameters b and c represnts the mortality rate of the predator and consumer. The parameter α_1 and α_2 represents the predation rate of the predator and that of the consumer respectively, while the predator mortality rate is given by b.

The food web of the drive patch is described by the Blasius model (Appendix A). The predator mortality rate of the drive patch is given by b_d. The amplitude of drive dispersal is given by e. The predator mortality rate of the drive patch is given by b_d. The amplitude of drive dispersal is given by e. Equivalently, the food web of the response patch is governed by the same Blasius Model, with a different dispersal term as explained in Appendix B. For the i^{th} response patch, the parameter b_i describes the mortality of the predator of the i^{th} patch. The other parameters are same as above.

3. Results

Figure 1 shows the bifurcation diagram of the Blasius model food web with respect to the mortality of predator b. We have chosen both b_d and all b_r values such that our investigated food webs all exhibit chaotic dynamics, when unperturbed ($b \epsilon (0.8 - 1)$). The structure of our metacommunity is represented in 1(b), where the circle filled with red represents the drive patch, the arrows from it represent the drive-response diseprsal. The circles filled with green represent the response patches. The dotted lines represent inter-response dispersals, whose magnitude is 0 when we consider unconnected responses. As stated above, the habitat heterogeneity across various patches in the metacommnunity comes from the variation in predator mortality rate b from one patch to another.

Figure 1: (a) Bifurcation diagram of one single food web model described by the Blasius model, plotted as a function of predator mortality rate b.

3.1. Locally connected response patches

3.1.1. Homogeneous drive and response patches

First, we consider a group of locally connected, completely identical responses that are linked to an identical drive through dispersal, where we have fixed the predator mortality rate for both drive and response patches as $b_{drive} = b_{responses} = 0.9$. The temporal dynamics of the consumers from the drive patch Z_d and a representative response patch Z_r are depicted in Fig. 2(a) and 2(b) for the dispersal strengths $e = 0.2$ and 3.5, respectively. The chaotic attractor of the both drive patch (red line) and a representative response patch (green line) are also shown in the inset of both Figs. 2(a) and 2(b). It is evident from these figures that the time series of Z_d and Z_r evolve identically corroborating the complete synchronization among the drive and uncoupled response systems for both low and high values of the dispersal strength. In order to elucidate the nature of the dynamics exhibited by both drive and response systems, we have plotted their bifurcation diagrams, obtained by collecting the maxima Z_{dmax} and Z_{rmax} of the temporal dynamics of the drive and response patches, in Figs. 2(c) and 2(d), respectively, in the range of the dispersal strength $e \in (0.0, 3.5)$. It is evident from these bifurcation diagrams that the evolution of the consumer, and consequently resource X and predator Y, of both drive and response patches exhibit chaotic oscillation for their population density in the entire explored range of the dispersal strength. Thus, the homogeneous drive is not capable of quenching the chaotic evolution of the population density in response patches.

Figure 2: Locally connected response patches: Completely identical drive and homogeneous response patches. (a) Time series of consumer dynamics of both drive patch and a response patch for the low dispersal strength $e = 0.2$, (b) same as (a) for the high dispersal strength $e = 3.5$, (c) bifurcation diagram of the drive patch as a function of e, and (d) bifurcation diagram of a response patch as a function of e. Attractors of the drive (red line) and a response (green line) patches are depicted in the inset of (a) and (b). Predator mortality rate for drive $b_d = 0.9$, and for responses $b_r = 0.8$.

Figure 3: Locally connected response patches: Nonidentical drive and homogeneous response patches. (a) Time series of consumer dynamics of both drive patch and a response patch for the low dispersal strength $e = 0.2$, (b) same as (a) for the high dispersal strength $\varepsilon_d = 3.5$, (c) bifurcation diagram of the drive patch as a function of e, and (d) bifurcation diagram of a response patch as a function of e. Attractors of the drive (red line) and a response (green line) patches are depicted in the inset of (a) and (b). Predator mortality rate for drive $b_d = 0.9$, and for responses $b_r = 0.8$.

3.1.2. Heterogeneous drive and homogeneous response patches

Next, we consider a group of locally connected, completely identical responses that are linked to a non-identical drive through dispersal, where we have fixed the predator mortality rate for both drive and response patches as $b_{drive} = 0.9$ and $b_{responses} = 0.8$. The temporal dynamics of the consumers from the drive patch Z_d and a representative response patch Z_r are depicted in Fig. 3(a) and 3(b) for the dispersal strengths $e = 0.2$ and 3.5, respectively. The chaotic attractor of the both drive patch (red line) and a representative response patch (green line) are also shown in the inset of both Figs. 3(a) and 3(b). In order to elucidate the nature of the dynamics exhibited by both drive and response systems, we have plotted their bifurcation diagrams, obtained by collecting the maxima Z_{dmax} and Z_{rmax} of the temporal dynamics of the

drive and response patches, in Figs. 3(c) and 3(d), respectively, in the range of the dispersal strength $e \in (0.0, 3.5)$. It is evident from these bifurcation diagrams that the evolution of the consumer, and consequently resource X and predator Y, of both drive and response patches exhibit chaotic oscillation for their population density in the range of low dispersal strength. However, with increasing dispersal ($e\epsilon(0.8, 1.0)$), the dynamics of the drive patche undergo Hopf bifurcation from chaotic to a bistable state, consisting of 2 limit cycles. Upon further increase of dispersal ($e\epsilon(1.25, 3.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a bistable steady state. However, in case of the response patch, with increasing dispersal ($e\epsilon(0.8, 1.0)$) the dynamics undergoes Hopf Bifurcation from chaotic go mutli-stable limit cycles. Upon further increase of dispersal ($e\epsilon(1.25, 3.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a multi-stable steady state. Thus, the on-identical but similar drive quenches the chaotic evolution of the population density in completely identical locally connected response patches.

3.1.3. Heterogeneous drive and response patches

Next, we consider a group of locally connected, heterogeneous responses that are linked to a non-identical drive through dispersal, where we have fixed the predator mortality rate for both drive and response patches as $b_{drive} = 0.9$ and $b_{responses} = 0.77$, 0.76,0.82 and 0.85. The temporal dynamics of the consumers from the drive patch Z_d and a representative response patch Z_r are depicted in Fig. 4(a) and 4(b) for the dispersal strengths $e = 0.2$ and 3.5, respectively. The dynamics evidently changes from chaotic to steady state on increasing drive-response dispersal. The chaotic attractor of the both drive patch (red line) and a representative response patch (green line) are also shown in the inset of both Figs. 4(a) and 4(b). In order to elucidate the nature of the dynamics exhibited by both drive and response systems, we have plotted their bifurcation diagrams, obtained by collecting the maxima Z_{dmax} and Z_{rmax} of the temporal dynamics of the drive and response patches, in Figs. 4(c) and 4(d), respectively, in the range of the dispersal strength $\varepsilon_d \in (0.0, 3.0)$. It is evident from these bifurcation diagrams that the evolution of the consumer, and consequently resource X and predator Y, of both drive and response patches exhibit chaotic oscillation for their population density in the range of low dispersal strength ($e\epsilon(0.1, 0.25)$). However, with increasing dispersal ($e\epsilon(0.3, 0.8)$), the dynamics of the drive patch undergo Hopf bifurcation from chaotic to a bistable state, consisting of 2 limit cycles.

Figure 4: Same as Fig. 3 but predator mortality rate for drive and response patches are $b_d = 0.9$ and $b_r = 0.77$, 0.76,0.82 and 0.85.

Figure 5: N

Upon further increase of dispersal ($e\epsilon(0.8, 2.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a monostable steady state, with a decreasing trend in the average population density. On further increase in dispersal, ($e > 2.5$), the mono stable state is maintained but the average population density saturates. In case of the response patch, with increasing dispersal ($e\epsilon(0.1, 0.25)$) the dynamics stays chaotic. Upon further increase of dispersal ($e\epsilon(0.25, 0.8)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a multi-stable limit cycle. Upon further increase of dispersal ($e\epsilon(0.8, 2.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a bistable steady state, with a decreasing trend in the average population density. On further increase in dispersal, ($e > 2.5$), the bistable state is maintained but the average population density saturates. Thus, the non-identical but similar drive quenches the chaotic evolution of the population density in non-identical locally connected response patches.

3.1.4. Synchronization of the metacommunity

We investigate the emergence of synchronization in a group of oscillators as a function of coupling strength, considering identical, dissimilar and heterogenous systems. Our goal is to determine what degree of heterogeneity promotoes synchrony and what level of heterogeneity is required in the network to induce asynchronous steady states. We define synchronization error

for the group of oscillators. averaged over time as,

$$\eta = \frac{1}{T} \sum_t \sqrt{(\bar{Y})_t^2 - (\bar{Y^2})_t}. \tag{2a}$$

Here, $(\bar{Y})_t = \frac{1}{N} \sum_{i=1}^N Y_i$ and $(\bar{Y^2})_t = \frac{1}{N} \sum_{i=1}^N Y_i^2$ average values of the variables Y and Y^2 of oscillators $i = 1, ...N$, at an instant of time t. Further, we average η over different initial states to obtain an ensemble averaged synchronization error η.

Figure 5 demonstrates the synchronization between all the patches in the metacommunity, when the responses are nearest neighbor coupled. When the drive and responses are completely identical, all patches are synchronized. However, with slight habitat heterogeneity in the system, the synchronization error has a low but non-zero magnitude. However, with increasing invasionary dispersal e, the synchronization between the patches decrease. Synchronization is detrimental to the persistence of the metacommunity. Therefore, it is clearly demonstrated that in this case, invasion increases the persistence of the metacommunity as a whole. With further increase in the heterogeneity in the network, for the range of dispersal $e\epsilon(0.1 - 1.5$, there is an increasing but fluctuating trend in the synchronization error $< \eta >$. However, with further increase in dispersal e, the synchronization error keeps steadily increasing without fluctuation.

3.2. Globally connected response patches

3.2.1. Homogeneous drive and response patches

First, we consider a group of globally connected, completely identical responses that are linked to an identical drive through dispersal, where we have fixed the predator mortality rate for both drive and response patches as $b_{drive} = b_{responses} = 0.9$. The temporal dynamics of the consumers from the drive patch Z_d and a representative response patch Z_r are depicted in Fig. 6(a) and 6(b) for the dispersal strengths $e = 0.2$ and 3.5, respectively. The chaotic attractor of the both drive patch (red line) and a representative response patch (green line) are also shown in the inset of both Figs. 6(a) and 6(b). It is evident from these figures that the time series of Z_d and Z_r evolve identically corroborating the complete synchronization among the drive and uncoupled response systems for both low and high values of the

Figure 6: Globally connected response patches: Completely identical drive and homogeneous response patches. (a) Time series of consumer dynamics of both drive patch and a response patch for the low dispersal strength $e = 0.2$, (b) same as (a) for the high dispersal strength $\varepsilon_d = 3.5$, (c) bifurcation diagram of the drive patch as a function of e, and (d) bifurcation diagram of a response patch as a function of e. Attractors of the drive (red line) and a response (green line) patches are depicted in the inset of (a) and (b). Predator mortality rate for drive $b_d = 0.9$, and for responses $b_r = 0.8$.

Figure 7: Same as Fig. 6 but predator mortality rate for drive and response patches are $b_d = 0.9$ and $b_r = 0.8$.

dispersal strength. In order to elucidate the nature of the dynamics exhibited by both drive and response systems, we have plotted their bifurcation diagrams, obtained by collecting the maxima Z_{dmax} and Z_{rmax} of the temporal dynamics of the drive and response patches, in Figs. 6(c) and 6(d), respectively, in the range of the dispersal strength $e \in (0.0, 3.5)$. It is evident from these bifurcation diagrams that the evolution of the consumer, and consequently resource X and predator Y, of both drive and response patches exhibit chaotic oscillation for their population density in the entire explored range of the dispersal strength. Thus, the homogeneous drive is not capable of quenching the chaotic evolution of the population density in response patches.

3.2.2. Heterogeneous drive and homogeneous response patches

Next, we consider a group of globally connected, completely identical responses that are linked to a non-identical drive through dispersal, where we have fixed the predator mortality rate for both drive and response patches as $b_{drive} = 0.9$ and $b_{responses} = 0.8$. The temporal dynamics of the consumers

from the drive patch Z_d and a representative response patch Z_r are depicted in Fig. 7(a) and 7(b) for the dispersal strengths $e = 0.2$ and 3.5, respectively. The chaotic attractor of the both drive patch (red line) and a representative response patch (green line) are also shown in the inset of both Figs. 7(a) and 7(b). In order to elucidate the nature of the dynamics exhibited by both drive and response systems, we have plotted their bifurcation diagrams, obtained by collecting the maxima Z_{dmax} and Z_{rmax} of the temporal dynamics of the drive and response patches, in Figs. 7(c) and 7(d), respectively, in the range of the dispersal strength $\varepsilon_d \in (0.0, 3.5)$. It is evident from these bifurcation diagrams that the evolution of the consumer, and consequently resource X and predator Y, of both drive and response patches exhibit chaotic oscillation for their population density in the range of low dispersal strength. However, with increasing dispersal ($e\epsilon(0.8, 1.0)$), the dynamics of the drive patche undergo Hopf bifurcation from chaotic to a bistable state, consisting of 2 limit cycles. Upon further increase of dispersal ($e\epsilon(1.25, 3.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a bistable steady state. However, in case of the response patch, with increasing dispersal ($e\epsilon(0.8, 1.0)$) the dynamics undergoes Hopf Bifurcation from chaotic go mutli-stable limit cycles. Upon further increase of dispersal ($e\epsilon(1.25, 3.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a multi-stable steady state. Thus, the on-identical but similar drive quenches the chaotic evolution of the population density in completely identical locally connected response patches.

3.2.3. Heterogeneous drive and homogeneous response patches

Next, we consider a group of locally connected, heterogeneous responses that are linked to a non-identical drive through dispersal, where we have fixed the predator mortality rate for both drive and response patches as $b_{drive} = 0.9$ and $b_{responses} = 0.77$, 0.76,0.82 and 0.85. The temporal dynamics of the consumers from the drive patch Z_d and a representative response patch Z_r are depicted in Fig. 8(a) and 8(b) for the dispersal strengths $e = 0.2$ and 3.5, respectively. The dynamics evidently changes from chaotic to steady state on increasing drive-response dispersal. The chaotic attractor of the both drive patch (red line) and a representative response patch (green line) are also shown in the inset of both Figs. 8(a) and 8(b). In order to elucidate the nature of the dynamics exhibited by both drive and response systems, we have plotted their bifurcation diagrams, obtained by collecting the maxima Z_{dmax} and Z_{rmax} of the temporal dynamics of the drive and response patches,

Figure 8: Same as Fig. 7 but predator mortality rate for drive and response patches are $b_d = 0.9$ and $b_r = 0.77$, 0.76,0.82 and 0.85.

in Figs. 8(c) and 8(d), respectively, in the range of the dispersal strength $\varepsilon_d \in (0.0, 3.0)$. It is evident from these bifurcation diagrams that the evolution of the consumer, and consequently resource X and predator Y, of both drive and response patches exhibit chaotic oscillation for their population density in the range of low dispersal strength ($e\epsilon(0.1, 0.25)$). However, with increasing dispersal ($e\epsilon(0.4, 0.9)$), the dynamics of the drive patch undergo Hopf bifurcation from chaotic to a multistable state, consisting of 4 limit cycles to ultimately bistable states consisting of 2 limit cycles. Upon further increase of dispersal ($e\epsilon(0.8, 2.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a monostable steady state, with a decreasing trend in the average population density. On further increase in dispersal, ($e > 2.5$), the mono stable state is maintained but the average population density saturates. In case of the response patch, with increasing dispersal ($e\epsilon(0.1, 0.25)$) the dynamics stays chaotic. Upon further increase of dispersal ($e\epsilon(0.25, 0.8)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a multi-stable limit cycle. Upon further increase of dispersal ($e\epsilon(0.8, 2.5)$) the dynamics of the drive patch further undergoes reverse bifurcation to obtain a bistable steady state, with a decreasing trend in the average population density. On further increase in dispersal, ($e > 2.5$), the bistable state is maintained but the average population density saturates. Thus, the non-identical but similar drive quenches the chaotic evolution of the population density in non-identical locally connected response patches.

3.2.4. Synchronization of the network

Figure 9 demonstrates the synchronization between the different patches of the metacommunity. The synchronizations follow the same nature as that of Fig. 5.

4. Discussion and Conclusion

Chaotic population dynamics in ecosystems have been a staple subject of studies in recent years. Chaotic fluctuations in populations have generally been associated with low persistence as noise and rapid, large fluctuations subject populations to genetic bottlenecks and extinctions induced by stochasticity. However, inherently chaotic ecosystems continue to exist and thrive. This can be explained by control or quenching of chaos engineered by inherent ecological mechanisms or external forcings on ecosystems. The

Figure 9: N

question of chaos control or quenching of chaos in ecological systems coupled through dispersal has not been investigated in detail so far, as per the authors' knowledge goes. In this study, we consider an ecological network of five patches, with each patch containing a tri-trophic, chaotic foodweb. We looked at how quenching of chaos occurs in both a locally connected and a globally connected network of ecosystems when driven by invasionary dispersal from an external ecosystem.

For both types of network structures, we first considered the case of completely identical drive and responses. We demonstrated that for completely identical drives and responses, chaotic fluctuations are too strong to quench, irrespective of drive-response dispersal amplitude. Next, we introduced heterogeneity in predator mortality of the drive patch and response patches. This led to quenching of chaos and emergence of multistable steady states in the population dynamics of both drives and responses for high degrees of drive-response dispersal. Under a high degree of dispersal, the dynamics of the drive was that of a bistable state, and that of response was of a multi-stable steady state. Next, we increased the degree of heterogeneity even further and included heterogeneity in predator mortality across the response patches as well. As a result, the bistable steady state of the drive changed to a monostable steady state, and that of the responses changed to a bistable steady state. Our results strongly signify that heterogeneous metacommunities attain increased robustness and persistence through quenching of chaos.

Furthermore, for both kinds of network structure, we investigated various kinds of synchronizations induced by dispersal between the drive and response patches. Even with quenching of chaos, generalized synchronization is established in some, but not all, of the cases. The complete synchronization (quantified by synchronization error) and phase synchronization are nonexistent in most cases. These two types of synchronization are determined by network structure. However, the same investigation into synchronization between two response patches shows that in most cases, both phase and generalized synchronization exist. However, complete synchronization exists only when two responses are completely identical.

From the above results, we conclude that, barring a few idealized cases of completely identical drive and responses, dispersal primarily quenches chaos in real metacommunities consisting of inherently chaotic ecosystems and heterogeneous patches. Indeed, this study highlights the relevance of habitat heterogeneity in increasing persistence (or decreasing extinction risk) of metacommunities through control and quenching of chaos. This study also comments on the relevance of dispersal as one of the most fundamental mechanisms stabilizing the dynamics of metacommunities. Further, our work quantifies the level of drive dispersal required to quench chaos and whether such dispersal stabilizes the metacommunity or pushes it to extinction. Moreover, our investigation proves that whether the post-quenching steady-state dynamics has high species-density or low species-density depends on the degree of dispersal, habitat heterogeneity, and also on the network structure of the metacommunity.

Appendix A. Appendix: Blasius Model (Drive System)

The food web of the drive patch is described by the Blasius model Blasius et al. (1999) and governed by the equations

$$\dot{X}_d = aX_d - \alpha_1 \frac{X_d Y_d}{1 + kX_d} + \frac{e}{N} \sum_{i=1}^{N} (X_i - 4X_d), \qquad \text{(A.1a)}$$

$$\dot{Y}_d = -b_d Y_d + \alpha_1 \frac{X_d Y_d}{1 + kX_d} - \alpha_2 Y_d Z_d \qquad \text{(A.1b)}$$

$$\dot{Z}_d = -c(Z_d - Z_s) + \alpha_2 Y_d Z_d, \qquad \text{(A.1c)}$$

,

The parameter a represents the growth rate of the resource in absence of inter-species interactions, while the parameters b and c represnts the mortality rate of the predator and consumer. The parameter α_1 represents the predation rate of the predator and α_2 represents the predation rate of the consumer. The predator mortality rate of the drive patch is given by b_d. The amplitude of drive dispersal is given by e. The parameter N represents the number of response patches. For this entire study, $N = 4$.

Appendix B. Appendix: Blasius Model (Response network)

The food web of the response patches are governed by the following equations (Blasius Model):

$$\dot{X}_i = aX_i - \alpha_1 \frac{X_i Y_i}{1 + kX_i} +$$

$$e(X_d - X_i) + e_i \sum_{j=1, j \neq i}^{N} \gamma_{ij}(X_j - X_i), \tag{B.1a}$$

$$\dot{Y}_i = -b_i Y_i + \alpha_1 \frac{X_i Y_i}{1 + kX_i} - \alpha_2 f_2(Y_i, Z_i), \tag{B.1b}$$

$$\dot{Z}_i = -c(Z_i - Z_s) + \alpha_2 f_2(Y_i, Z_i), \tag{B.1c}$$

Γ_{ij} is the adjacency matrix detailing the network structure created through dispersal between patches. When the response patches are unconnected, all elements of the matrix $\gamma_{ij} \,\epsilon\, \Gamma_{ij} = 0$. For nearest neighbor coupling between the responses, elements of the matrix $\gamma_{ij} \,\epsilon\, \Gamma_{ij} \neq 0$, only if $i = j$. When the response patches are randomly coupled, elements of the matrix $\gamma_{ij} \,\epsilon\, \Gamma_{ij} \neq 0$ randomly. For the response patch i, the parameter b_i describes the mortality of the predator of the drive patch. The other parameters are given by $\alpha_1 = 0.2$, $\alpha_2 = 1, k = 0.05$, $Z_s = 0.006$, $c = 10.0$, $a = 1$ and $f = 0.25$. The magnitude of simple diffusive dispersal between the i^{th} resource patch and its nearest neighbors is given by the parameter e_i. Note that when we consider the scenario for **unconnected** response patches $\varepsilon = 0$. For the i^{th} response patch, the parameter b_i describes the mortality of the predator of the i^{th} patch. The other parameters are same as above.

Acknowledgements

The work of D.B. is supported by IISER Thiruvananthapuram.

References

Allen, J.C., Schaffer, W.M., Rosko, D., 1993. Chaos reduces species extinction by amplifying local population noise. Nature 364(6434), 229–232. doi:https://doi.org/10.1038/364229a0.

Berryman, A.A., Millstein, J.A., 1989. Are ecological systems chaotic — and if not, why not? Trends in Ecology & Evolution 4(1), 26–28. doi:https://doi.org/10.1016/0169-5347(89)90014-1.

Blasius, B., Huppert, A., Stone, L., 1999. Complex dynamics and phase synchronization in spatially extended ecological systems. Nature 399, 354–359. doi:https://doi.org/10.1038/20676.

Boccaletti, S., Grebogi, C., Lai, Y.C., Mancini, H., Maza, D., 2000. The control of chaos: theory and applications. Physics reports 329, 103–197.

Croteau, E., 2010. Causes and consequences of dispersal in plants and animals. Nature Education Knowledge 3, 12.

Dietze, M.C., 2018. Prediction in ecology: a first-principles framework. Ecological Applications 27(7), 2048–2060. doi:https://doi.org/10.1002/eap.1589.

Doebeli, M., 1993. The evolutionary advantage of controlled chaos. Proceedings: Biological Sciences 254, 281–285. doi:http://www.jstor.org/stable/49907.

Gamarra, J.G., Solé, R.V., Alonso, D., 2001. Control, synchrony and the persistence of chaotic populations. Chaos, Solitons & Fractals 12(2)), 235–249. doi:https://doi.org/10.1016/S0960-0779(00)00110-7.

Hastings, A., 1982. Dynamics of a single species in a spatially varying environment: the stabilizing role of high dispersal rates. Journal of mathematical biology 16, 49–55.

Hastings, A., Powell, T., 1991. Chaos in a three-species food chain. Ecology 72(3), 896–903. doi:https://doi.org/10.2307/1940591.

Hilborn, R., Quinn, Schindler, T.P., Rogers, D.E., 2003. Biocomplexity and fisheries sustainability. Proceedings of the national academy of sciences 100(11), 6564–6568. doi:https://doi.org/10.1073/pnas.1037274100.

J. D. Yeakel, J. W. Moore, P.R.J., deAguiar, M.A.M., 2014. Synchronisation and stability in river metapopulation networks. Ecology Letters doi:`10.1111/ele.12228`.

Jørgensen, S.E., Jørgensen, S.E., 2002. Chaos, fractals and ecology. Integration of Ecosystem Theories: A Pattern 1, 247–264. doi:`https://doi.org/10.1007/978-94-011-2682-3_10`.

Krkos˘ek, M., Drake, J., 2014. On signals of phase transitions in salmon population dynamics. Proc. R. Soc. B 281, 20133221. doi:`http://dx.doi.org/10.1098/rspb.2013.3221`.

Larsen, S., et al., 2021. A dispersal-induced paradox: synchrony and stability in stochastic metapopulations. Ecology letters .

Mandal, S., Ray, S., Roy, S., Jørgensen, S.E., 2006. Order to chaos and vice versa in an aquatic ecosystem. Ecological Modelling 197, 498–504. doi:`https://doi.org/10.1016/j.ecolmodel.2006.03.020`.

McCann, K., Hastings, A., 1997. Re–evaluating the omnivory–stability relationship in food webs. Proc. R. Soc. Lond. B 264, 1249–1254. doi:`http://doi.org/10.1098/rspb.1997.0172`.

Sarremejane, R., et al., 2021. Drought effects on invertebrate metapopulation dynamics and quasi-extinction risk in an intermittent river network. Global Change Biology doi:`https://doi.org/10.1111/gcb.15720`.

Solé, R.V., Gamarra, J.G., 1998. Chaos, dispersal and extinction in coupled ecosystems. Journal of theoretical Biology 193(3), 539–541. doi:`https://ui.adsabs.harvard.edu/link_gateway/1998JThBi.193..539S/doi:10.1006/jtbi.1998.0716`.

Thomas, W.R., Pomerantz, M., Gilpin, M.E., 1980. Chaos, asymmetric growth and group selection for dynamical stability. Ecology 61, 1312–1320. doi:`https://doi.org/10.2307/1939039`.

Upadhyay, R.K., , Iyengar, S.R.K., Rai, V., 1998. Chaos: An ecological reality? International Journal of Bifurcation and Chaos 8(6), 1325–1333. doi:`https://doi.org/10.1142/S0218127498001029`.

Williams, R., Martineza, N., 2004. Stabilization of chaotic and non-permanent food-web dynamics. Eur. Phys. J. B 38, 297–303. doi:`https://doi.org/10.1140/epjb/e2004-00122-1`.