1 Examining the fitness benefits of social plasticity to prey
2 availability in bottlenose dolphins

Running title: Social buffering in bottlenose dolphins

David N. Fisher*

School of Biological Sciences, University of Aberdeen, King's College, Aberdeen,
AB243FX

david.fisher@abdn.ac.uk

* corresponding author

Barbara J. Cheney

School of Biological Sciences, University of Aberdeen, Lighthouse Field Station,
George Street, Cromarty, IV11 8YL

b.cheney@abdn.ac.uk

17 Abstract

A key explanation for phenotypic plasticity is that changing behaviour to match
conditions increases fitness. Affiliative social associations with conspecifics can be
important for coping with challenging conditions, and therefore plasticity in social
behaviour in response to environment variation may bring fitness benefits. Here we
test the adaptive benefit of social plasticity by quantifying female bottlenose dolphins'
(*Tursiops truncatus*) plastic responses to salmon (*Salmo salar*) abundance and
determine if either this plasticity or mean social behaviour is associated with calving
success. We found that more gregarious females and those better connected to all
parts of the entire social network had higher calving success, but plasticity in social
behaviour in response to salmon abundance was not associated with this component

of fitness. This lack of relationship may have arisen as salmon abundance in either

current or previous years was not associated with the probability of producing a calf.

We therefore demonstrate that being more social is positively associated with

females' calving success in our study population, but social plasticity itself is not

associated with fitness benefits, questioning whether this is likely to be a mechanism

for this population of dolphins to cope with challenging environmental conditions.

Key words: dolphin, food abundance, environmental change, network, plasticity,

social buffering, Tursiops

Introduction

Animals are frequently exposed to adverse environmental conditions, including
extreme weather events or fluctuations in prey availability and predation risk, which
can impact their fitness. To mitigate these factors, animals may change their traits
(show phenotypic plasticity) to buffer any fitness costs that may otherwise be
inevitable in adverse conditions (Blumstein et al., 2023; Candolin & Wong, 2011;
López-Sepulcre & Kokko, 2012). For instance, brown trout *Salmo trutta* may
decrease their metabolism at times of low food availability (Auer et al., 2015), while
goldcrest *Regulus regulus* form larger groups to huddle and conserve heat in cold
conditions (Hogstad, 1984). Behaviours are the type of trait that are often thought to
be animals' first line of defence against adverse conditions, as they can change
relatively quickly compared to morphological or even physiological traits (Sih, 2013;
Sih et al., 2011). Consequently, changing behaviour to ameliorate the costs of
environmental stressors is thought to be widespread (Tuomainen & Candolin, 2010;
Wong & Candolin, 2015).

Many animals live in groups or near others, and so engage in social behaviours;
interactions with conspecifics that can be cooperative or competitive in nature
(Rubenstein & Abbot, 2017; Székely et al., 2010). Social behaviours are therefore a
key mechanism through which animals may be able to buffer against environmental
change (Komdeur & Ma, 2021). Global-scale comparative analyses have been used
to test the general idea that social interactions with conspecifics helps animals cope
with harsh environments. These relate aspects of a species' social structure, such as
the presence or absence of cooperative breeding, to environmental parameters of
the species' native range (e.g. variability in rainfall). For example, higher incidences
of communal living are found in social spiders in areas with higher rain intensity
which challenge the survival of lone females (Guevara & Avilés, 2015). These
analyses have demonstrated a link between cooperative breeding systems and
survival in harsh conditions, supporting the idea that social living helps animals cope
with environmental stressors (Firman et al., 2020; Guevara & Avilés, 2015; Jetz &
Rubenstein, 2011; Sheehan et al., 2015).

Comparative analyses, however, do not shed light on whether *plasticity* in behaviour
helps animals survive and reproduce in adverse conditions. Such studies require

multiple measures of individual social behaviour across a range of environments and

to associate variation in the social behaviours with a fitness component. Recent

studies have demonstrated a clear interaction between social behaviour or social

conditions (such as group size) and environmental factors in a regression on fitness

components like survival or reproductive success (Borger et al., 2023; Guindre-

Parker & Rubenstein, 2020). This suggests that animals can adjust their social

behaviours to limit the harm caused by stressful environments, and the authors

argue this provides evidence for adaptive social plasticity.

However, while the analyses conducted by Borger, Guindre-Parker, and colleagues

shows whether particular behaviours of individuals bring benefits in certain

conditions and not others, they do not explicitly test whether plasticity itself is

beneficial. Distinct subsets of individuals could show different behaviours when

environmental conditions are compared, which would still result in a clear statistical

interaction. Instead, as proposed by Warrington et al. (2024), a more appropriate test

for the adaptive value of social plasticity is to estimate the covariance between an

individual's plasticity in social behaviour in response to some environment factor and

a fitness component of that individual. This approach uses multivariate mixed-effect

modelling techniques (Arnold et al., 2019; Houslay & Wilson, 2017), which

specifically account for repeated measures of the behaviour but allow a single

measure of the fitness component *and* estimate the covariance at the correct

hierarchical level (the among-individual level). Such models estimate the

covariances between fitness and both individual trait means and trait plasticities,

testing the adaptive benefit of social behaviours overall (i.e., if being more gregarious

brings fitness benefits) and the adaptive benefit of social plasticity itself (i.e., if being

more flexible in connectedness brings fitness benefits). This approach has been

used to show the benefits of plasticity in traits such as life-history and phenology, for

instance female Ural owls *Strix uralensis* that show steeper declines in clutch size

with earlier lay date have higher fitness (Brommer et al., 2012). However, this

approach has not yet been used for social behaviours. We therefore have no current

evidence testing the suggestion that animals have evolved to plastically adjust their

social behaviour to buffer against environmental stressors.

Here, we test the associations between a fitness component (calving success) and

both mean and plasticity of individual social behaviours, in wild female bottlenose

dolphins (*Tursiops truncatus*). Bottlenose dolphins (*Tursiops* spp.) exhibit a fission-
fusion grouping pattern where social interactions and relationships are critical
components in their lives (Connor et al., 2000, 2019; Lusseau et al., 2006; Shane et
al., 1986). Further, we have shown that dolphin gregariousness, quantified as the
social network measure "strength", increases in months of high salmon abundance
(Fisher & Cheney, 2023, see also the correction: 2025), a key food source in this
study population (Santos et al., 2001). This raises the question of whether this
plasticity in response to salmon abundance is adaptive, i.e., does changing social
behaviour enhance fitness. This would be expected if social associations influence
the ability to acquire food (e.g., by cooperation or competition). Previous work has
shown mean levels of social behaviour are associated with fitness components in
toothed whales (Connor et al., 2022; Ellis et al., 2017; Gerber et al., 2022; Holmes et
al., 2024) but the fitness benefits of plasticity itself are not quantified.

We directly tested for a possible adaptive function of social plasticity by fitting models
that simultaneously estimate individual plasticity in social behaviours in response to
salmon abundance and individual variance in a fitness component (calving success).
Crucially, we estimate the covariance between calving success and both mean
individual social behaviour (testing for benefits of mean behaviour) and plasticity in
individual social behaviour in response to salmon abundance (testing for benefits of
plasticity in behaviour). Since we previously looked at plasticity at both the monthly
and yearly temporal scale (Fisher & Cheney, 2023), we repeat that two-scale
approach to test whether responding to changes in monthly or yearly salmon
abundance brings fitness benefits. Our aims are:

1. Determine if there are consequences for reproduction of changes in yearly
salmon abundance.
2. Estimate plasticity in social behaviour in response to salmon abundance (as
achieved by Fisher & Cheney, 2023 but with an updated dataset and with a
slightly different modelling approach).
3. Estimate the association with fitness of individual variation in mean social
behaviours.
4. Estimate the association with fitness of individual variation in plasticity in
social behaviours.

Methods

Data collection

We used data from a long-term individual-based study of a population of over 220 bottlenose dolphins in the North Sea (Cheney et al., 2013, 2014, 2024). Boat-based photo-identification surveys have been carried out each year from May to September 1990 to 2022 within the Moray Firth Special Area of Conservation (SAC; 92/43/EEC). Upon encountering dolphins (an "encounter"), each group is carefully followed and photographs taken of both sides of as many of the dolphins' dorsal fins as possible. Due to unique patterns of markings, scratches, and notches on each dolphin's back and dorsal fin, individual dolphins are identified in these photographs and tracked through time. For this study, we used data from 14,614 sightings of individual dolphins across 2,368 encounters (average 70 groups per year, range 24 – 145, average group size 6.3, range 1 - 56).

Identifying individuals allows us to determine which dolphins are still alive, who is in the same group and so associating, and to identify new calves (Cheney et al., 2019). The year of birth and so age can be estimated for calves ≤ 2 years old, but ages are unknown for individuals first sighted as older calves, immature or adults. Mothers are assigned to calves based on repeat observations of the calf surfacing in echelon position next to the same adult female (Cheney et al., 2019), but paternity cannot be assigned as genetic data are not available. For more information on survey methods, see Cheney et al. (2014), while Fisher and Cheney (2023) provide information on using these survey records to construct social networks. Photographic survey work was conducted under NatureScot Animal Scientific Licence (number 93751) and all work was approved by the University of Aberdeen School of Biological Sciences ethics board (project ID SBSEC/01/2023/March).

Social trait and fitness component calculation

We defined reproductive output per year in a binary manner as whether the female was known to give birth or not (Cheney et al., 2019). Sightings records are then used

to determine whether the calf survived beyond one or two years. Calves are regularly

observed with their mother in at least their first two years (Grellier et al., 2003) so

repeat sightings the mother without the calf at that age is a reliable indicator of calf

death rather than dispersal. We estimated a fitness component "calving success" as

the number of calves produced by each female that survived more than two years

divided by the number of years she lived as a reproductive individual (year last

observed – year of first reproduction). For females of known ages, the start of their

reproductive period was six years old (as this is the earliest known age a female has

been observed with a calf in our population, with a median age of nine; Cheney et

al., 2019), while for individuals of unknown age, it is from the year they were first

observed with a calf.

We can measure social traits of individuals by using the repeated observations of

individuals in groups to build social networks. In a social network, individuals who are

observed in the same group (photographed during an encounter, within 100m,

engaged in similar activities and if travelling moving in a single direction; Wells et al.,

1987) are connected, while those who are never observed in the same group are not

connected (the "gambit of the group"; Whitehead & Dufault, 1999). The relative

frequency two individuals are seen together defines the strength of the association –

precisely, the number of times two individuals were seen together in a given time

period divided by the total number of times both individuals were seen, either

together or alone (the "simple ratio index"), which ranges from 0 to 1 (Cairns &

Schwager, 1987). We constructed a social network per month and per year, using

observations of individuals of either unknown age or above three years old, i.e.,

excluding known calves. We used the whole network to calculate two measures of

each individual's social network position, representing two dimensions of social

behaviour. However, we then excluded individuals observed fewer than three times

in that month or year from any analysis, as their social network position would be

based on very few observations, and we excluded entire months (i.e., all rows in that

month) if there were fewer than five groups observed in that month.

For each individual in each month or year, we calculated "strength", the sum of all an

individual's weighted connections, giving an estimate of overall gregariousness. We

also calculated "closeness", which is based on the path lengths from an individual to

all others in the network, weighted by the inverse of association strengths,

representing connectedness to the entire population and has high values if an

individual has strong connections linking them to all parts of the network (Opsahl et

al., 2010). Both measures are therefore undirected and weighted. We use these

measures as we have previously found they show some variation in individual

change in response to measures of salmon abundance, which is necessary to then

be linked to among-individual differences in fitness (Fisher & Cheney, 2023). Fisher

& Cheney (2023) also found that dolphin social behaviour varies with salmon

abundance only at the monthly scale, and therefore we expect any association

between calving success and plasticity to be at this scale. We plot social networks in

Fig. 1 for 2011 and 2021 as examples.

Figure 1. Plots of yearly social networks for 2011 (a.) and 2021 (b.). Points represent

individual female dolphins and lines show associations between them. We have only

plotted associations with a simple ratio index above 0.2 to reduce visual clutter. Line

thickness is increased with stronger associations, and point size is increased if that

individual had higher calving success in that year.

Salmon abundance estimate

We followed Fisher and Cheney (2023) and Lusseau *et al.* (2004) in using data on

monthly catches by fishing rods (excluding nets) of Atlantic salmon (*Salmo salar*)

from five rivers feeding into the Moray Firth (the Alness, Beauly, Canon, Ness, and

Nairn) to quantify relative salmon abundance. These catch data are positively

correlated among months and among rivers and with other data sources, suggesting

they accurately reflect abundance (Thorley et al., 2005; Youngson et al., 2002).

Further, salmon abundance in the current year is positively associated with group

size (Lusseau et al., 2004) and in the current month with individual dolphin

gregariousness (Fisher & Cheney, 2023). We downloaded the data from

https://marine.gov.scot/data/marine-scotland-salmon-and-sea-trout-catches-salmon-

district-shinyapp, using "rod data", both one sea winter and multiple sea winter adults

and summing retained and released fish for a monthly index of salmon abundance.

We summed monthly catches within a calendar year for an annual index of salmon

abundance.

Data analysis

All models were fitted in MCMCglmm (v. 2.36) (Hadfield, 2010) in R (v. 4.4.1) (R

Development Core Team & Team, 2016). To test the effect of salmon abundance on

female reproduction (aim 1), we regressed the binary outcome of whether a female

had a calf or not in that year on two variables: salmon abundance in the current year

and salmon abundance in the previous year, as gestation in bottlenose dolphins is

around 12 months and so could be affected by food availability in either calendar

year. Both salmon abundances were z-transformed to a mean of zero and a

standard deviation of one. In this analysis, we only included females who had been

sighted a minimum of three times in that year, as we would otherwise be uncertain of

their reproductive success. Note that for this and all other analyses increasing the

minimum number of sightings per year decreases the number of observations and

the number of unique females included: for the analysis of salmon abundance and

reproductive success if we require a minimum of three encounters per individual per

year we have 712 observations of 100 unique females, with four per year we have

612 observations of 94 unique females, and with five sightings per year we have 530

observations of 82 unique females. There is therefore an unavoidable trade-off

between different aspects of statistical power. For this analysis the results are

qualitatively the same with a threshold of three, four, or five sightings (Tables S1-3).

In our statistical models we included random effects of maternal ID and year to

account for repeated observations of the same dolphins across years and for years

that might have had higher or lower reproductive success on average. We also

included a binary fixed effect of whether the female had given birth in the last two

years or not, as this would greatly reduce the chances of her reproducing in the

current year and needs to be accounted for (Cheney et al., 2019). We used a

categorical error structure, and uninformative priors with V = 1 and nu = 0.02 for both

random effects and the residual variance fixed at 1 (as in a model with a binary

response the residual variance is defined by the mean). We ran the model for

110,000 iterations, discarding the first 10,000 as the model would initially be far from

convergence, and using 1 out of every 100 remaining iterations for the posterior

distribution to minimise autocorrelation. We checked model trace plots for

satisfactory convergence and tested three runs of this model with the Gelman and

Rubin convergence diagnostic. All terms achieved satisfactory convergence

(potential scale reduction factor [PSRF] very close to 1). We report the posterior

distribution medians in the main text as they are likely the least biased (Pick et al.,

2023), but include the modes and means in tables reporting full model results in the

supplementary materials.

We then fitted four models to estimate mean and plasticity in social behaviour and

testing for associations with calving success; aims 2-4, for all combinations of

strength and closeness at monthly and yearly scales. All four models have an

identical structure except that the monthly models also include a random effect of

month, to account for seasonal variation in factors, aside from salmon abundance,

that might influence social behaviour. For each model, we include a social network

trait (either strength or closeness) and calving success as the two response

variables. As before, we only included measures of social network traits where the

female had been observed at least three times in a given month or year. Increasing

this threshold to four or five approximately halves or quarters the dataset

respectively for the monthly analyses, resulting in a dataset too small to fit our

chosen models. The effect is less dramatic in the yearly analysis as relatively more

females have at least four or five sightings per year, but we keep the same threshold

for both temporal scales to allow comparisons of the results. We also only included

females in these analyses that had been seen for at least five years as reproductive,

as some individuals below that threshold had artificially high calving success due to

having a surviving calf in one year and then not being seen again.

We used salmon abundance in that year or month (z-transformed) as a fixed effect

for the social network trait only. We include the random effects of individual ID for

both traits and year (and month for the monthly models) for the social network trait

only. We include random intercepts for the social network trait and for calving

success, and random slopes for the social network trait with salmon abundance. We

estimate all pairwise covariances between social network trait intercepts (mean

individual social behaviour), social network trait slopes (individual plasticity in social

behaviour in response to salmon abundance), and individual calving success. These

are estimated at the among-individual level, and so the covariance between the

individual slopes and calving success specifically tests whether individual plasticity is

associated with fitness (Houslay & Wilson, 2017). We used Gaussian error

structures for both traits, uninformative priors with V = 1 and nu = 0.02 for the

residual variance of the social network trait and the random effect of year. The

residual variance for calving success was fixed at 0.0001 to aid estimation as the

trait does not vary within individuals. Finally, for the variance-covariance matrix at the

among-individual level, we used a parameter-expanded prior with V = 1, nu = 3,

alpha.mu = 0, and alpha.V = 625 following Houslay and Wilson (2017). We checked

the trace plots for all four models for satisfactory convergence and tested three runs

of each model with the Gelman and Rubin convergence diagnostic. Terms in all

models achieved satisfactory convergence (PSRF < 1.005), apart from the random

effect of month in the model for monthly closeness (PSRF = 1.204) and to a lesser

extent the same random effect in the model for monthly strength (PSRF = 1.055).

Since these are control terms and not of specific interest we do not believe this is

problematic.

Results

To test the effect of salmon abundance on yearly calf production and survival, we

had 712 observations of known reproductive status across 100 different females.

Salmon abundance in both current (posterior distribution median [PDM] = -0.218,

95% credible intervals [CIs] = -0.647 to 0.206) and previous years (PDM = -0.137,

95% CIs = -0.579 to 0.291) did not influence reproductive success. Having a calf in

the previous two years reduced the chance of a female producing a calf (PDM = -

1.977, 95% CIs = -2.597 to -1.369; full model results in Table S1).

To test for the fitness associations between mean and plasticity in social behaviour at

the monthly scale, we had 736 measures of social behaviour across 63 different

females. For the yearly scale analysis, we had 602 measures of social behaviour

across 66 different females.

At the monthly level, there were no covariances between calving success and

plasticity in strength or closeness (Table 1). There were also no covariances

between intercepts of strength or closeness and calving success (Figs. 2a & b &

Table 1). There were no intercept-slope covariances for either trait (Table 1). Full

model results are given in Tables S4 & 5.

Table 1. Estimates of the covariance parameters for the models with strength and

closeness. In each cell are the posterior distribution median (and the 95% credible

intervals in parentheses) for the covariance between the two terms shown. Above

the diagonal (filled with Xs) are values for the monthly analyses, while below the

diagonal are values for the yearly analyses. Covariances that do not overlap with

zero are shown in bold.

	Strength		
	Individual intercepts	Individual slopes	Calving success
Individual intercepts	X	-0.250 (-0.964 to 0.123)	0.053 (-0.000 to 0.116)
Individual slopes	-0.034 (-0.289 to 0.138)	X	-0.015 (-0.048 to 0.008)
Calving success	**0.046 (0.005 to 0.091)**	0.000 (-0.015 to 0.017)	X
	Closeness		
Individual intercepts	X	-0.000 (-0.001 to 0.000)	0.001 (-0.001 to 0.003)
Individual slopes	-0.000 (-0.002 to 0.000)	X	-0.000 (-0.002 to 0.001)
Calving success	**0.003 (0.001 to 0.006)**	-0.000 (-0.002 to 0.001)	X

Figure 2. Plots of the relationship between monthly (a. & b.) and yearly (c. & d.) means of social network traits (a. & c show strength, b. & d. show closeness) and calving success. Each point represents an individual females' mean social network trait for a given month or year against that individual's calving success (one measure per individual). For both traits at the yearly level (c. and d.), there was a clear positive covariance, and so we show a trend line as a solid coloured line with the standard errors shown in grey (both calculated using the "stat_smooth" function in ggplot2; Wickham, 2009).

At the yearly level, there were no covariances between plasticity in strength and closeness (Table 2) and calving success. There were weak positive covariances between calving success and intercepts of both strength (Fig. 2c & Table 2, correlation median = 0.333, 95% CRIs = 0.106 to 0.548) and closeness (Fig. 2d & Table 2, correlation median = 0.345, 95% CRIs = 0.117 to 0.538). There were no intercept-slope covariances for either trait (Table 2). Full model results are given in Tables S6 & 7.

Discussion

We found that salmon abundance was not associated with female reproductive

success (aim 1). We estimated both individual means and plasticities of social

behaviour (aim 2) and demonstrated that there are positive covariances between

individual means in strength and closeness and calving success, but only at the

yearly timescale (aim 3). We found no evidence that social plasticity might be

adaptive as there were no covariances between plasticity in strength or closeness in

response to salmon abundance and calving success (aim 4).

Female dolphins who had high strength and closeness over an entire year had

higher calving success. Positive associations between mean social behaviour and

fitness components, such as longevity, has been reported multiple times in terrestrial

mammals, such as rhesus macaques *Macaca mulatta* (Ellis et al., 2019), big horn

sheep *Ovis canadensis* (Wal et al., 2015) and spotted hyenas *Crocuta crocuta*

(Yoshida et al., 2016). Further, in a closely studied population of Indo-Pacific

bottlenose dolphins *T. aduncus* (Allen et al., 2016), Stanton and Mann (2012) found

that young males with higher eigenvector centrality (a social network measure of

connectedness accounting for the connections possessed by one's social

associates) had better survival from weaning to 10 years of age. More recent

research on the same population showed that adult males with strong bonds and

who are well integrated in their cooperative alliance sire more offspring (Gerber et

al., 2022). However, females in Stanton and Mann's (2012) study, which is the sex

we studied here, showed a non-significant negative relationship between eigenvector

centrality and survival (although not directly tested for a difference from zero, it

seems likely the relationship in females is at least not positive). Given that we

studied different fitness components and a different species with a different ecology

to Stanton and Mann, whether our results are truly in contrast is questionable.

Clearly, further comparative work on associations between fitness components and

social behaviour in both sexes and multiple populations of *Tursiops* spp. is

warranted. Such work should take temporal scale into account, as we did not find a

clear relationship between social network traits and calving success at the monthly

scale. While we might expect the same animals to have high trait values in both the

yearly and monthly analysis, the additional month-to-month variation in the monthly

analysis may have made detecting the covariance more difficult. Additionally, the two

analyses had marginally different sets of females, which could have led to the slightly

different results

In general, fitness benefits of sociality are thought to arise due to detection of and

protection from predators, opportunities to share information, and the ability to

complete tasks more efficiently compared to when alone (Rubenstein & Abbot,

2017). In this bottlenose dolphin population, social grouping is unlikely to function as

protection from predators, as their natural predators are absent from these waters

(Cheney et al., 2019). Additionally, while the dolphins in the Moray Firth SAC tend to

be observed in groups (median size approximately 6-10 individuals; Cheney et al.,

2024), they are generally observed hunting individually (B. Cheney, pers obs.),

suggesting increased hunting efficiency may not be the key benefit. Instead, a

possibility is that high gregariousness and being connected to all subgroups in the

population may help adult female dolphins find food patches or maintain a larger

number of female associates with whom to share calf care responsibilities, reducing

the risk of infanticide by males and/or allowing mothers to spend more time foraging

to provide energy for lactation (Methion et al., 2023). In the Indo-Pacific bottlenose

dolphin, females tend to associate with conspecifics sharing similar reproductive

state (Möller & Harcourt, 2008), and a female's reproductive success tends to be

higher if she has successful female relatives or associates with other successful

females (Frère et al., 2010). However, it is possible that the causality of the

relationship between gregariousness and calving success could be in the opposite

direction; regular presence of calves may make a female dolphin socially attractive

(such an association in fact drove a spurious association between maternal social

behaviour and offspring survival in Amboseli baboons Papio spp.; Creighton et al.,

2025). Further, this relationship could be observed if female dolphins in poorer

condition or those with adverse early life effects are less likely to have calves and

spend more time alone hunting (Thompson, 2019). Since we analysed an

association between repeated measures of social network traits and a single

measure of reproductive success, confounding factors must act on a similar temporal

scale. Fine scale data, such as those analysed by Creighton *et al.* (2025), and

appropriate randomisations or experimental approaches, may be necessary to

confirm causal relationships.

We did not detect any fitness benefits to plasticity. Female dolphins do not appear to

adjust their social behaviour to buffer low salmon abundance or take advantage of

high salmon abundance. This observational dataset has inherent limitations, for

example, not all females are observed every year, not all calves are identified, and

some calves may die before they are observed. These limitations add noise to the

dataset and so prevent us detecting genuine relationships between fitness

components and other traits. Also, as we did not manipulate social behaviour, we

cannot be certain of the causal link to calving success. Nevertheless, given we show

that salmon abundance is not related to calf production, it is perhaps not surprising

that changes in social behaviour are not required to "buffer" low salmon numbers.

This could indicate that the benefits of social behaviour identified here are not related

to more effective salmon location and predation. Dolphins may switch to other prey

when salmon are scarce, such as whiting *Merlangius merlangus*, saithe *Pollachius*

virens, and cod *Gadus morhua* (Santos et al., 2001), instead of changing their social

behaviour. Estimating the resource requirements of the population should be a

priority in the face of a changing climate. Meanwhile, social behaviour, which shows

considerable within-individual variation, may instead change in response to

reproductive opportunities, to shifts in local currents and fronts, or other

environmental factors such as anthropogenic disturbance, and the adaptive value of

plasticity might show itself in any of these contexts.

Our study is the first to estimate the covariance between plasticity in social behaviour

and a fitness component, but two previous studies on birds have tested whether

different social behaviours yield fitness benefits in different conditions. In superb

starlings *Lamprotornis superbus*, males have higher survival in larger groups only in

wet years but the reverse is true in dry years (Guindre-Parker & Rubenstein, 2020).

Interestingly, in females, larger groups are always associated with better survival,

matching our result (although we looked at reproduction rather than survival).

Meanwhile, in Seychelles warblers *Acrocephalus sechellensis*, low rainfall is

associated with lower rates of reproduction but, in years of little rain, alloparenting

was *less* common, in opposition to predictions that social behaviours help buffer

harsh conditions (Borger et al., 2023). Therefore, while there is a range of evidence

to show that social behaviour plastically varies with environmental conditions (Fisher

et al., 2021), there is currently no evidence that this plasticity buffers harsh
environmental conditions.

In general, there is variable evidence for selection on plasticity. For example,
Pelletier *et al*. (2007) found selection on seasonal plasticity in female bighorn sheep
ewes, with more plastic individuals having higher reproduction. Meanwhile, female
red deer *Cervus elaphus* that give birth early after dry autumns and late after wet
autumns may have higher fitness, but this trend was marginally non-significant (p =
0.06) and only applied if those females were born in a high density period (Nussey et
al., 2005). Meanwhile, female Ural owls show selection on quadratic reaction norms
between laying date and clutch size, but no selection on the linear reaction norms
(Brommer et al., 2012), which is what we evaluated here. These studies give an idea
of the diversity of findings but we are not aware of any meta-analyses collecting
selection estimates on plasticity in animals. Arnold et al. (2019) reviewed evidence
for selection on thermal plasticity in plants and found it to be lacking, while Davidson
et al. (2011) found that invasive plants were more plastic than non-invasive plants,
but that this plasticity did not confer especial fitness benefits. Therefore, while we
have spent much of the 21st century wondering what the costs of plasticity are
(Pigliucci, 2005), the benefits might not be as empirically well-supported as
expected.

Future studies on the benefits of social plasticity in bottlenose dolphins could focus
on "developmental plasticity" – the ability of a genotype to produce different
phenotypes over the course of development, which cannot be reversed (Taborsky,
2017). This is in contrast to "reversible plasticity", where an adult individual changes
its phenotype and can reverse the changes back (measured here). For example,
stress exposure in juvenile rats *Rattus norvegicus* reduces sociality in later life (Toth
et al., 2008). Evidence that developmental plasticity in social behaviour is adaptive,
however, is inconclusive, with seemingly maladaptive changes often observed
(Spencer, 2017). In Indo-Pacific bottlenose dolphins, juveniles can show
considerable variation in social behaviour (Gibson & Mann, 2008), which persists
into adulthood (Evans et al., 2021), and has fitness consequences (Gerber et al.,
2022; Holmes et al., 2024; Stanton & Mann, 2012). Further work should therefore
identify what factors drive early variation in social behaviour and the adaptive
consequences of early-life effects.

Acknowledgements

We are forever grateful to Paul Thompson for conception and development of this long-term individual-based research study, and for commenting on this manuscript. We are especially thankful for our colleagues at the Lighthouse Field Station, past and present, who have contributed to and advanced this long-term study. Simon Allen and three anonymous reviewers made numerous useful comments and suggestions on an earlier draft. The contribution of numerous funders has been key for this long-term study, including University of Aberdeen, NatureScot, Beatrice Offshore Windfarm Ltd., Moray Offshore Renewables Ltd., Marine Scotland, the Crown Estate, Highlands and Islands Enterprise, the BES, ASAB, Greenpeace Environmental Trust, Whale and Dolphin Conservation, Talisman Energy (UK) Ltd., Department of Energy and Climate Change, Chevron and Natural Environment Research. Survey work was conducted under NatureScot Animal Scientific Licences.

Authors' contributions

BJC collects data on this long-term project and curates the database. DNF conceived the idea of this particular manuscript and conducted preliminary analyses. DNF and BJC discussed the ideas and the analysis before DNF finalised the analysis and wrote a draft. BJC edited the draft and together the authors revised the final version.

Conflict of interest declaration

We declare we have no competing interests.

References

Allen, S. J., Bryant, K. A., Kraus, R. H. S., Loneragan, N. R., Kopps, A. M., Brown, A. M., Gerber, L., & Krützen, M. (2016). Genetic isolation between coastal and

508   fishery-impacted, offshore bottlenose dolphin (*Tursiops* spp.) populations.

509   *Molecular Ecology*, *25*(12), 2735–2753. https://doi.org/10.1111/mec.13622

510   Arnold, P. A., Nicotra, A. B., & Kruuk, L. E. B. (2019). Sparse evidence for selection

511   on phenotypic plasticity in response to temperature. *Philosophical*

512   *Transactions of the Royal Society B: Biological Sciences*, *374*(1768),

513   20180185. https://doi.org/10.1098/rstb.2018.0185

514   Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J., & Metcalfe, N. B. (2015).

515   Flexibility in metabolic rate confers a growth advantage under changing food

516   availability. *Journal of Animal Ecology*, *84*(5), 1405–1411.

517   https://doi.org/10.1111/1365-2656.12384

518   Blumstein, D. T., Hayes, L. D., & Pinter-Wollman, N. (2023). Social consequences of

519   rapid environmental change. *Trends in Ecology & Evolution*, *38*(4), 337–345.

520   https://doi.org/10.1016/j.tree.2022.11.005

521   Borger, M. J., Richardson, D. S., Dugdale, H., Burke, T., & Komdeur, J. (2023).

522   Testing the environmental buffering hypothesis of cooperative breeding in the

523   Seychelles warbler. *Acta Ethologica*, *26*(3), 211–224.

524   https://doi.org/10.1007/s10211-022-00408-y

525   Brommer, J. E., Kontiainen, P., & Pietiäinen, H. (2012). Selection on plasticity of

526   seasonal life-history traits using random regression mixed model analysis.

527   *Ecology and Evolution*, *2*(4), 695–704. https://doi.org/10.1002/ece3.60

528   Cairns, S. J., & Schwager, S. J. (1987). A comparison of association indices. *Animal*

529   *Behaviour*, *35*(5), 1454–1469. https://doi.org/10.1016/S0003-3472(87)80018-

530   0

531   Candolin, U., & Wong, B. B. M. (2011). *Behavioural Responses to a Changing World:*

532   *Mechanisms & Consequences* (U. Candolin & B. B. M. Wong, Eds.). Oxford

533   University Press. https://global.oup.com/academic/product/behavioural-

534   responses-to-a-changing-world-9780199602575?cc=gb&lang=en&

535   Cheney, B. J., Arso Civil, M., Hammond, P. S., & Thompson, P. M. (2024). *Site*

536   *Condition Monitoring of bottlenose dolphins within the Moray Firth Special*

537   *Area of Conservation 2017-2022* (1360; NatureScot Research Report).

538   NatureScot. https://www.nature.scot/doc/naturescot-research-report-1360-

539   site-condition-monitoring-bottlenose-dolphins-within-moray-firth

540   Cheney, B. J., Corkrey, R., Durban, J. W., Grellier, K., Hammond, P. S., Islas-

541   Villanueva, V., Janik, V. M., Lusseau, S. M., Parsons, K. M., Quick, N. J.,

542   Wilson, B., & Thompson, P. M. (2014). Long-term trends in the use of a

543   protected area by small cetaceans in relation to changes in population status.

544   *Global Ecology and Conservation*, *2*, 118–128.

545   https://doi.org/10.1016/j.gecco.2014.08.010

546 Cheney, B. J., Thompson, P. M., & Cordes, L. S. (2019). Increasing trends in

547   fecundity and calf survival of bottlenose dolphins in a marine protected area.

548   *Scientific Reports*, *9*(1), 1767. https://doi.org/10.1038/s41598-018-38278-9

549 Cheney, B. J., Thompson, P. M., Ingram, S. N., Hammond, P. S., Stevick, P. T.,

550   Durban, J. W., Culloch, R. M., Elwen, S. H., Mandleberg, L., Janik, V. M.,

551   Quick, N. J., ISLAS-Villanueva, V., Robinson, K. P., Costa, M., Eisfeld, S. M.,

552   Walters, A., Phillips, C., Weir, C. R., Evans, P. G. H., … Wilson, B. (2013).

553   Integrating multiple data sources to assess the distribution and abundance of

554   bottlenose dolphins Tursiops truncatus in Scottish waters. *Mammal Review*,

555   *43*(1), 71–88. https://doi.org/10.1111/j.1365-2907.2011.00208.x

556 Connor, R. C., Krützen, M., Allen, S. J., Sherwin, W. B., & King, S. L. (2022).

557   Strategic intergroup alliances increase access to a contested resource in male

558   bottlenose dolphins. *Proceedings of the National Academy of Sciences*,

559   *119*(36), e2121723119. https://doi.org/10.1073/pnas.2121723119

560 Connor, R. C., Sakai, M., Morisaka, T., & Allen, S. J. (2019). The Indo-Pacific

561   Bottlenose Dolphin (Tursiops aduncus). In B. Würsig (Ed.), *Ethology and*

562   *Behavioral Ecology of Odontocetes* (pp. 345–368). Springer International

563   Publishing. https://doi.org/10.1007/978-3-030-16663-2_16

564 Connor, R. C., Wells, R. S., Mann, J., & Read, A. J. (2000). The bottlenose dolphin.

565   In *Cetacean Societies: Field studies of whales and dolphins* (pp. 91–125).

566   University of Chicago Press Chicago, Illinois.

567   https://books.google.com/books?hl=en&lr=&id=W-

568   UQNoxMONwC&oi=fnd&pg=PA91&dq=The+bottlenose+dolphin.+In%C2%A0

569   Cetacean+Societies:+Field+studies+of+whales+and+dolphins&ots=WVoTXS

570   wbyk&sig=ElaUH2Qk9T9v3N_aTx_-csPPcpc

571 Creighton, M. J. A., Lerch, B. A., Lange, E. C., Silk, J. B., Tung, J., Archie, E. A., &

572   Alberts, S. C. (2025). Reevaluating the relationship between female sociality

573   and infant survival in wild baboons. *Proceedings of the National Academy of*

574   *Sciences*, *122*(20), e2417378122. https://doi.org/10.1073/pnas.2417378122

Davidson, A. M., Jennions, M., & Nicotra, A. B. (2011). Do invasive species show
higher phenotypic plasticity than native species and, if so, is it adaptive? A
meta-analysis. *Ecology Letters*, *14*(4), 419–431. https://doi.org/10.1111/j.1461-
0248.2011.01596.x

Ellis, S., Franks, D. W., Nattrass, S., Cant, M. A., Weiss, M. N., Giles, D., Balcomb,
K. C., & Croft, D. P. (2017). Mortality risk and social network position in
resident killer whales: Sex differences and the importance of resource
abundance. *Proceedings. Biological Sciences*, *284*(1865), 20171313.
https://doi.org/10.1098/rspb.2017.1313

Ellis, S., Snyder-Mackler, N., Ruiz-Lambides, A., Platt, M. L., & Brent, L. J. N. (2019).
Deconstructing sociality: The types of social connections that predict longevity
in a group-living primate. *Proceedings of the Royal Society B: Biological*
Sciences, *286*(1917), 20191991. https://doi.org/10.1098/rspb.2019.1991

Evans, T., Krzyszczyk, E., Frère, C., & Mann, J. (2021). Lifetime stability of social
traits in bottlenose dolphins. *Communications Biology*, *4*(1), 1–8.
https://doi.org/10.1038/s42003-021-02292-x

Firman, R. C., Rubenstein, D. R., Moran, J. M., Rowe, K. C., & Buzatto, B. A. (2020).
Extreme and Variable Climatic Conditions Drive the Evolution of Sociality in
Australian Rodents. *Current Biology*, *30*(4), 691-697.e3.
https://doi.org/10.1016/j.cub.2019.12.012

Fisher, D. N., & Cheney, B. J. (2023). Dolphin social phenotypes vary in response to
food availability but not the North Atlantic Oscillation index. *Proceedings of the*
Royal Society B: Biological Sciences, *290*(2008), 20231187.
https://doi.org/10.1098/rspb.2023.1187

Fisher, D. N., & Cheney, B. J. (2025). Correction to: 'Dolphin social phenotypes vary
in response to food availability but not the North Atlantic Oscillation index'
(2023), by Fisher and Cheney. *Proceedings of the Royal Society B: Biological*
Sciences, *292*(2046), 20242078. https://doi.org/10.1098/rspb.2024.2078

Fisher, D. N., Kilgour, R. J., Siracusa, E. R., Foote, J. R., Hobson, E. A., Montiglio, P.
O., Saltz, J. B., Wey, T. W., & Wice, E. W. (2021). Anticipated effects of abiotic
environmental change on intraspecific social interactions. *Biological Reviews*,
96(6), 2661–2693. https://doi.org/10.1111/brv.12772

Frère, C. H., Krützen, M., Mann, J., Connor, R. C., Bejder, L., & Sherwin, W. B.
(2010). Social and genetic interactions drive fitness variation in a free-living

dolphin population. *Proceedings of the National Academy of Sciences*,

107(46), 19949–19954. https://doi.org/10.1073/pnas.1007997107

Gerber, L., Connor, R. C., Allen, S. J., Horlacher, K., King, S. L., Sherwin, W. B.,

Willems, E. P., Wittwer, S., & Krützen, M. (2022). Social integration influences

fitness in allied male dolphins. *Current Biology*, *32*(7), 1664–1669.

Gibson, Q. A., & Mann, J. (2008). Early social development in wild bottlenose

dolphins: Sex differences, individual variation and maternal influence. *Animal*

Behaviour, *76*(2), 375–387. https://doi.org/10.1016/j.anbehav.2008.01.021

Grellier, K., Hammond, P. S., Wilson, B., Sanders-Reed, C. A., & Thompson, P. M.

(2003). Use of photo-identification data to quantify mother◻calf association

patterns in bottlenose dolphins. *Canadian Journal of Zoology*, *81*(8), 1421–

1427. https://doi.org/10.1139/z03-132

Guevara, J., & Avilés, L. (2015). Ecological predictors of spider sociality in the

Americas. *Global Ecology and Biogeography*, *24*(10), 1181–1191.

https://doi.org/10.1111/geb.12342

Guindre-Parker, S., & Rubenstein, D. R. (2020). Survival Benefits of Group Living in

a Fluctuating Environment. *The American Naturalist*, *195*(6), 1027–1036.

https://doi.org/10.1086/708496

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed

models: The MCMCglmm R package. *Journal of Statistical Software*, *33*(2),

1–22.

Hogstad, O. (1984). Variation in numbers, territoriality and flock size of a Goldcrest

Regulus regulus population in winter. *Ibis*, *126*(3), 296–306.

https://doi.org/10.1111/j.1474-919X.1984.tb00252.x

Holmes, K. G., Krützen, M., Ridley, A. R., Allen, S. J., Connor, R. C., Gerber, L.,

Flaherty Stamm, C., & King, S. L. (2024). Juvenile social play predicts adult

reproductive success in male bottlenose dolphins. *Proceedings of the*

National Academy of Sciences, *121*(25), e2305948121.

https://doi.org/10.1073/pnas.2305948121

Houslay, T. M., & Wilson, A. J. (2017). Avoiding the misuse of BLUP in behavioural

ecology. *Behavioral Ecology*, *12*(Pt A), S9.

https://doi.org/10.1093/beheco/arx023

Jetz, W., & Rubenstein, D. R. (2011). Environmental Uncertainty and the Global
Biogeography of Cooperative Breeding in Birds. *Current Biology*, *21*(1), 72–
78. https://doi.org/10.1016/J.CUB.2010.11.075

Komdeur, J., & Ma, L. (2021). Keeping up with environmental change: The
importance of sociality. *Ethology*, *127*(10), 790–807.
https://doi.org/10.1111/ETH.13200

López-Sepulcre, A., & Kokko, H. (2012). Understanding behavioural responses and
their consequences. In U. Candolin & B. B. M. Wong (Eds.), *Behavioural*
Responses to a Changing World: Mechanisms and Consequences. Oxford
University Press.
https://www.oxfordscholarship.com/view/10.1093/acprof:osobl/978019960256
8.001.0001/acprof-9780199602568-chapter-1

Lusseau, D., Williams, R., Wilson, B., Grellier, K., Barton, T. R., Hammond, P. S., &
Thompson, P. M. (2004). Parallel influence of climate on the behaviour of
Pacific killer whales and Atlantic bottlenose dolphins. *Ecology Letters*, *7*(11),
1068–1076. https://doi.org/10.1111/j.1461-0248.2004.00669.x

Lusseau, D., Wilson, B., Hammond, P. S., Grellier, K., Durban, J. W., PARSONS, K.
M., Barton, T. R., & Thompson, P. M. (2006). Quantifying the influence of
sociality on population structure in bottlenose dolphins. *Journal of Animal*
Ecology, *75*(1), 14–24. https://doi.org/10.1111/j.1365-2656.2005.01013.x

Methion, S., Giralt Paradell, O., Padín, X. A., Corrège, T., & Díaz López, B. (2023).
Group size varies with climate and oceanographic conditions in bottlenose
dolphins. *Marine Biology*, *170*(1), 7. https://doi.org/10.1007/s00227-022-
04154-4

Möller, L. M., & Harcourt, R. G. (2008). Shared Reproductive State Enhances
Female Associations in Dolphins. *International Journal of Ecology*, *2008*(1),
498390. https://doi.org/10.1155/2008/498390

Nussey, D. H., Clutton-Brock, T. H., Elston, D. A., Albon, S. D., & Kruuk, L. E. B.
(2005). Phenotypic Plasticity in a Maternal Trait in Red Deer. *Journal of*
Animal Ecology, *74*(2), 387–396.

Opsahl, T., Agneessens, F., & Skvoretz, J. (2010). Node centrality in weighted
networks: Generalizing degree and shortest paths. *Social Networks*, *32*(3),
245–251. https://doi.org/10.1016/j.socnet.2010.03.006

Pelletier, F., Réale, D., Garant, D., Coltman, D. W., & Festa-Bianchet, M. (2007).
Selection on heritable seasonal phenotypic plasticity of body mass. *Evolution*,
61(8), 1969–1979. https://doi.org/10.1111/j.1558-5646.2007.00160.x
Pick, J. L., Kasper, C., Allegue, H., Dingemanse, N. J., Dochtermann, N. A.,
Laskowski, K. L., Lima, M. R., Schielzeth, H., Westneat, D. F., Wright, J., &
Araya-Ajoy, Y. G. (2023). Describing posterior distributions of variance
components: Problems and the use of null distributions to aid interpretation.
Methods in Ecology and Evolution, *14*(10), 2557–2574.
https://doi.org/10.1111/2041-210X.14200
Pigliucci, M. (2005). Evolution of phenotypic plasticity: Where are we going now?
Trends in Ecology & Evolution, *20*(9), 481–486.
R Development Core Team, & Team, R. D. C. (2016). R: A language and
environment for statistical computing. In *R Foundation for Statistical*
Computing (Vol. 1, p. 409) [Computer software]. R Foundation for Statistical
Computing. https://doi.org/10.1007/978-3-540-74686-7
Rubenstein, D. R., & Abbot, P. (2017). *Comparative Social Evolution* (D. R.
Rubenstein & P. Abbot, Eds.). Cambridge University Press.
https://doi.org/10.1017/9781107338319
Santos, M. B., Pierce, G. J., Reid, R. J., Patterson, I. a. P., Ross, H. M., & Mente, E.
(2001). Stomach contents of bottlenose dolphins (Tursiops truncatus) in
Scottish waters. *Journal of the Marine Biological Association of the United*
Kingdom, *81*(5), 873–878. https://doi.org/10.1017/S0025315401004714
Shane, S. H., Wells, R. S., & Würsig, B. (1986). Ecology, Behavior and Social
Organization of the Bottlenose Dolphin: A Review. *Marine Mammal Science*,
2(1), 34–63. https://doi.org/10.1111/j.1748-7692.1986.tb00026.x
Sheehan, M. J., Botero, C. A., Hendry, T. A., Sedio, B. E., Jandt, J. M., Weiner, S.,
Toth, A. L., & Tibbetts, E. A. (2015). Different axes of environmental variation
explain the presence vs. Extent of cooperative nest founding associations in
Polistes paper wasps. *Ecology Letters*, *18*(10), 1057–1067.
https://doi.org/10.1111/ele.12488
Sih, A. (2013). Understanding variation in behavioural responses to human-induced
rapid environmental change: A conceptual overview. *Animal Behaviour*.
https://www.sciencedirect.com/science/article/pii/S0003347213000936?casa_
token=2-

UuMOtyassAAAAA:eHXDj44BrAgcu7HfDrip6_IzwDas_b4SSR0q6EJbwryT_c

as6MqS3IB_cBN33xRlfTVSKNOxmg

Sih, A., Ferrari, M. C. O., & Harris, D. J. (2011). Evolution and behavioural responses

to human-induced rapid environmental change. *Evolutionary Applications*,

4(2), 367–387. https://doi.org/10.1111/j.1752-4571.2010.00166.x

Spencer, K. A. (2017). Developmental stress and social phenotypes: Integrating

neuroendocrine, behavioural and evolutionary perspectives. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *372*(1727),

20160242. https://doi.org/10.1098/rstb.2016.0242

Stanton, M. A., & Mann, J. (2012). Early social networks predict survival in wild

bottlenose dolphins. *PloS One*, *7*(10), e47508.

https://doi.org/10.1371/journal.pone.0047508

Székely, T. (Tamás), Moore, A. J. (Allen J., & Komdeur, J. (2010). *Social behaviour:*

Genes, ecology and evolution. Cambridge University Press.

https://www.cambridge.org/gb/academic/subjects/life-sciences/animal-

behaviour/social-behaviour-genes-ecology-and-

evolution?format=PB&isbn=9780521709620

Taborsky, B. (2017). Chapter Three - Developmental Plasticity: Preparing for Life in a

Complex World. In M. Naguib, J. Podos, L. W. Simmons, L. Barrett, S. D.

Healy, & M. Zuk (Eds.), *Advances in the Study of Behavior* (Vol. 49, pp. 49–

99). Academic Press. https://doi.org/10.1016/bs.asb.2016.12.002

Thompson, N. A. (2019). Understanding the links between social ties and fitness

over the life cycle in primates. *Behaviour*, *156*(9), 859–908.

https://doi.org/10.1163/1568539X-00003552

Thorley, J. L., Eatherley, D. M. R., Stephen, A. B., Simpson, I., MacLean, J. C., &

Youngson, A. F. (2005). Congruence between automatic fish counter data and

rod catches of Atlantic salmon (Salmo salar) in Scottish rivers. *ICES Journal*

of Marine Science, *62*(4), 808–817.

https://doi.org/10.1016/j.icesjms.2005.01.016

Toth, E., Avital, A., Leshem, M., Richter-Levin, G., & Braun, K. (2008). Neonatal and

juvenile stress induces changes in adult social behavior without affecting

cognitive function. *Behavioural Brain Research*, *190*(1), 135–139.

https://doi.org/10.1016/j.bbr.2008.02.012

Tuomainen, U., & Candolin, U. (2010). Behavioural responses to human-induced

environmental change. *Biological Reviews*, *86*(3), 640–657.

https://doi.org/10.1111/j.1469-185X.2010.00164.x

Wal, E. V., Festa-Bianchet, M., Vander Wal, E., Festa-Bianchet, M., Réale, D.,

Coltman, D. W., & Pelletier, F. (2015). Sex-based differences in the adaptive

value of social behavior contrasted against morphology and environment.

Ecology, *96*(3), 631–641. https://doi.org/10.1890/14-1320.1

Warrington, M., Fisher, D., Komdeur, J., Pilakouta, N., & Griesser, M. (2024).

Stronger together? A framework for studying population resilience to climate

change impacts via social shielding. Life Sciences.

https://doi.org/10.32942/X2QG9C

Wells, R. S., Scott, M. D., & Irvine, A. B. (1987). The Social Structure of Free-

Ranging Bottlenose Dolphins. In H. H. Genoways (Ed.), *Current Mammalogy*

(pp. 247–305). Springer US. https://doi.org/10.1007/978-1-4757-9909-5_7

Whitehead, H., & Dufault, S. (1999). Techniques for Analyzing Vertebrate Social

Structure Using Identified Individuals: Review and Recommendations.

Advances in the Study of Behavior, *28*.

http://www.sciencedirect.com/science/article/pii/S0065345408602156

Wickham, H. (2009). *ggplot2: Elegant graphics for data analysis* [Computer

software]. Springer New York. http://had.co.nz/ggplot2/book

Wong, B. B. M., & Candolin, U. (2015). Behavioral responses to changing

environments. *Behavioral Ecology*. https://doi.org/10.1093/beheco/aru183

Yoshida, K. C. S., Van Meter, P. E., & Holekamp, K. E. (2016). Variation among free-

living spotted hyenas in three personality traits. *Behaviour*, *153*(13–14), 1665–

1722. https://doi.org/10.1163/1568539X-00003367

Youngson, A. F., MacLean, J. C., & Fryer, R. J. (2002). Rod catch trends for early-

running MSW salmon in Scottish rivers (1952–1997): Divergence among

stock components. *ICES Journal of Marine Science*, *59*(4), 836–849.

https://doi.org/10.1006/jmsc.2002.1195