Diverging effects of global change on future invasion risks of *Agama picticauda* between

invaded regions: same problem, different solutions

Nicolas Dubos[1], Steven Calesse[2, 3], Kathleen C. Webster[4, 5], Chloé Bernet[3], Gregory Deso[6], Thomas

W. Fieldsend[7], Saoudati Maoulida[8], Soule Mounir[8, 9], Xavier Porcel[10], Jean-Michel Probst[11],

Hindatou Saidou[8], Jérémie Souchet[3], Mohamed Thani Ibouroi[8, 12]*, Markus A. Roesch[3, 10]*

* equal contribution

[1] Université de Rennes, CNRS, ECOBIO (Ecosystèmes, Biodiversité, Evolution), 35000, Rennes,

France

[2] Earth and Life Instistute, Université Catholique de Louvain, 1348 Louvain-La-Neuve, Belgique

[3] Association Nature Océan Indien, 46 rue des Mascarins 97429 Petite-Île, La Réunion, France

[4] Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057, Zurich, Switzerland

[5] Museum of Southwestern Biology and Department of Biology, University of New Mexico,

MSC03-2020, 1 University of New Mexico, Albuquerque, NM, USA

[6] AHPAM-Association Herpétologique de Provence Alpes Méditerranée, Maison des Associations,

384 route de Caderousse, 84100 Orange, France

[7] Department of Biological Sciences, Florida International University, Miami 33199, Florida, USA

[8] Groupe de Recherches et de Protection de la Faune et de la Flore des Iles de l'océan Indien

(GRPFOI), 6 rue Mnarajou, 97660, Dembeni, Mayotte, France

[9] Faculté des Sciences, Université de Mahajanga, Campus Universitaire d'Ambondrona, BP 652

401, Mahajanga, Madagascar

[10] CIBIO Research Centre in Biodiversity and Genetic Resources, InBIO Associate Laboratory,

University of Porto, Rua Padre Armando Quintas 7, 4485-661 Vairão, Portugal

[11] Association Nature and Patrimoine, Sainte Clotilde, La Réunion, France

[12] Réserve Naturelle Nationale de l'îlot Mbouzi, Naturalistes, Environnement et Patrimoine de

Mayotte, 5 Rue Georges Pompidou, 97660, Dembeni, Mayotte, France

Corresponding author: Nicolas Dubos, dubos.research@gmail.com

Abstract

Biological invasions can be predicted with modelling approaches to help understand the spread of invasive alien species and mitigate their impact. Many factors can be at play, potentially acting in contrasting directions and with heterogeneous effects through space. With an Ecological Niche Modelling approach, we predicted the invasion risks of Peters's Rock Agama *Agama picticauda*, a species that is spreading globally in non-forested areas through freight transport and un-/intentional releases from the pet trade. The potential establishment of the species in new areas is of concern for multiple endemic species throughout the world. We quantified the effects of climate, anthropogenic activity and forest cover on invasion risk. We used verified records from the native and non-native range and accounted for the latest methodological recommendations. We predicted how invasion risk will vary in the future (2070) using projections from two scenarios (SSP2 and SSP5). We predict that invasion risks will vary in diverging directions, depending on the region. The risk will increase in human-populated regions and on small islands but will decrease in Florida. We recommend increasing surveillance in vehicular transportation of material especially within the Comoros and the Mascarenes archipelagos. Since many introductions are related to the pet trade in Florida, we recommend stronger legal regulations and the promotion of public awareness. Promoting tree cover may be locally beneficial to prevent establishment of *A. picticauda*. The effect of climate change, land use change and human activities may differ between and within both, the native and the invaded regions.

Keywords

Climate change, Deforestation, Ecological Niche Modelling, Human activity, Pet trade, freight transport.

1. Introduction

Global transportation of goods as well as the international pet trade have led to many incidental or deliberate introductions of non-native species throughout the world (Gippet and Bertelsmeier, 2021). Once introduced to new environments, some species can establish and colonize further grounds, causing severe ecological and economic impact (Bellard et al., 2016; Diagne et al., 2021). New introductions are being reported every year, with no sign of saturation whatsoever (Clements et al., 2025; Seebens et al., 2017). Invasive alien species (IAS) are one of the main causes of native species decline, and the risk of establishment and spread of IAS is likely to vary in the future with global change (Bellard et al., 2013). Climate change can affect invasion risks in either direction, i.e., increasing or reducing the risk (Bellard et al., 2013; Dubos et al., 2023a), and the effect of climate and anthropogenic activities may sometimes also be contrasting (e.g., Dubos et al., 2025). Land use change can also alter invasion risk either positively or negatively, depending on the ability of the species to exploit anthropogenic structures and resources (e.g. Mitchell et al., 2021). Environmental change can affect populations of the native and the non-native region in opposite directions, due to different socio-economic contexts (Dubos et al., 2025). Similarly, for IAS that have spread globally, the mechanisms of invasion may differ between the invaded regions. However, few studies report quantifications of invasion risk while distinguishing between invaded regions, which can make management recommendations unsuitable to the local context in some cases.

The best strategy to mitigate the ecological and economic impacts incurred by IAS is the prevention of new introductions (Pearson, 2024). This can be achieved with the promotion of proactive surveillance and early detection at the main entry points such as airports and harbours in target regions (Cuthbert et al., 2022). The regions that are the most at risk in the present and future can be identified with predictive modelling approaches such as Ecological Niche Models (ENMs; also known as Species Distribution Models, SDMs). These models are widely used to predict species response to future environmental change (Dubos et al., 2023b), prioritizing conservation areas (Dubos et al., 2022b; Leroy et al., 2014), testing ecological hypotheses (Raxworthy et al., 2007) or

assessing invasion risk (Gallien et al., 2012). However, increasing accessibility with user-friendly

interfaces of highly complex models designs has led to a widespread misuse or misinterpretation of

SDMs (e.g. ignorance of biological or statistical assumptions, subjective reliance on performance

metrics, systematic use of default parameters; Leroy, 2023; Yackulic et al., 2013; Zurell et al.,

2020). As a result, projections of future invasion risk can largely differ and become potentially

misleading. However, provided the model design is in line with ecological theory and results are

interpreted specific to the natural history of the study organism, SDMs can produce maps that help

target priority entry points of potential IAS for stakeholders (Hui, 2022).

Modelling the environmental niche of non-native species that are still spreading is challenging

because of the non-equilibrium hypothesis (Gallien et al., 2012; Hui, 2022). After being introduced

outside of their native range, IAS have the potential to extend their realized niche as a result of

ecological release (i.e. fewer biotic constraints) and in situ adaptation (Fieldsend et al., 2021;

Refsnider et al., 2015). At an early stage of invasion, the species is not at equilibrium and is

susceptible to extend further its range. Hence the available occurrence data does not fully represent

the true extent of suitable environments that are likely to be colonized. Therefore, models might

downplay the importance of some suitable regions that have not yet been invaded and underestimate

the potential spread. To mitigate this effect, it is recommended to include both data from the native

and non-native range to train the models with all available information on the ecological niche of

the species (Barbet-Massin et al., 2018). It is further possible to exclude zones that have not yet

been reached by the species from the model training to reduce model omission error and the effect

of non-equilibrium (Chapman et al., 2019). To do so, the model training area (i.e. background)

should be limited to a restricted area near the known occurrences, and more weight should be given

to the most visited places (i.e. the equivalent of correcting for sample bias). Eventually, since the

non-native species could be introduced into places that are spatially separated, it is of critical

importance to evaluate models using approaches that enable the assessment of spatial transferability

(e.g. block-cross validation). To maximize the reliability of predictions of invasion risks, models

must be designed in close interaction between modellers and experts.

Peters's rock agama *Agama picticauda*, also known as the African red-headed agama or African

rainbow lizard*,* is native to central-western Africa (Leaché et al., 2014). Its current global

distribution is the result of both incidental escapes from the pet trade, deliberate introductions from

pet owners, hitchhiking with maritime transportation of goods and natural dispersion through

natural or anthropogenic structures (Gray, 2020; Guillermet et al., 1998; Mitchell et al., 2021). The

species was confounded with *A. agama* species complex until its systematic and taxonomic

clarification by Leaché et al. (2014) The identity of introduced populations of this agama has also

been confirmed as *A. picticauda* (Nuñez et al., 2016; Roesch et al., 2025). The species is now

present and well-established outside of its native range in Florida, Réunion Island and Grand

Comore, where it may represent a threat to local species. Further introductions have already begun

to be documented in the Bahamas, the British Virgin Islands, Madagascar, Cape Verde Islands and

the USA (where establishment is yet to be confirmed; Enge, 2024; Wagner et al., 2009 and

references therein). Additional introductions and establishments are expected to occur in presently

unreached regions (e.g. in the Lesser Antilles; van den Burg et al., 2024).

Agama picticauda can be found in a variety of habitat types in its native region, but inhabits mostly

open habitats such as savannah or forest edges (Krishnan et al., 2019). The species particularly

benefits from anthropogenic structures, where it can find shelter, thermoregulation sites, a high prey

availability and few competitors (Mitchell et al., 2021). The species is an opportunistic insectivore,

and introduced populations have recently been found to show dietary niche conservatism preserving

Hymenoptera as their main prey (Harris et al., 2025). However, it has also been observed eating

fruits or discarded human food items (Gray, 2020; Ofori et al., 2018). The preference for

hymenopteran diet and high flexibility of the species to exploit anthropogenic food sources grants it

a concerning potential for further spread in the tropics.

The presence of *A. picticauda* could represent a major threat to native reptiles. There is evidence of a risk of predation on smaller individuals (including cannibalism and predation on reptiles; Henigan et al., 2019; Ofori et al., 2023; van den Burg et al., 2024). The species may also impact native reptiles through interspecific competition (van den Burg et al., 2024). *Agama picticauda* displays an aggressive territoriality (Yeboah 1982) to which endemic reptiles can be highly sensitive (e.g. *Phelsuma inexpectata* in Réunion Island; Deso et al., 2023). Multiple regions are of concern, as for example in Grand Comore, where the endemic *Oplurus cuvieri comorensis* is already the focus of conservation efforts (Hawlitschek et al., 2011). On Réunion Island, *A. picticauda* has rapidly spread across all coastal areas since its introduction (first record in 1995; Guillermet et al., 1998) and is progressing up to higher elevations (Roesch et al., 2025). The positive effect of anthropogenic structures is all the more concerning for species that dwell within urban areas such as the critically endangered *P. inexpectata*, or those living in the vicinity of villages such as *P. borbonica* (Augros et al., 2017; Dubos, 2013; Dubos et al., 2014). Its introduction is expected to affect many reptile species in the Lesser Antilles (> 45 species; van den Burg et al., 2024). This calls for the urgent need to predict invasion risks at the global scale.

Agama picticauda is absent from dense forests, suggesting a lower impact on natural forested areas (van den Burg et al., 2024). However, the high transition rate of forests to urban areas (e.g. in the Caribbean; van den Burg et al., 2024) and the alarming rates of deforestation at the global scale suggest a potential increase in the area of habitat suitable for this species in the future. We present this study as a response to an urgent need to identify areas that are predicted to be suitable for *A. picticauda* by 2070, accounting for future projections of climate, land use and human activity.

We modelled the invasion risks of *A. picticauda*, accounting for its environmental niche (climate and habitat) and factors of introduction and spread. With a robust approach which accounts for the latest methodological recommendations, we identify regions that are the most at risk, quantify the effects of future climate change and deforestation for the native and the three invaded regions separately, and provide guidelines for targeted future surveillance efforts.

2. Methods

2.1 Overview – we used an ecological niche modelling approach with a combination of the two best performing algorithms (RandomForest down-sampled and MaxEnt; details below), correcting for sample bias (coping with non-equilibrium assumptions), with a spatial partitioning for model evaluation (to assess spatial transferability), a robust predictor selection process, accounting for anthropogenic factors, habitat and high-resolution global climate data that are appropriate for small islands.

2.2 Occurrence data – We used verified occurrence data from multiple sources from both the native and non-native regions, as recommended to best inform the models of the environmental conditions already filled by the species (Hui, 2022). We selected verified data points from multiple sources: i.e. the Global Assessment of Reptile Distributions (GARD; Roll et al., 2017), GBIF (gbif.org), iNaturalist research grade, the association Nature Océan Indien (NOI), Système d'information de l'inventaire du patrimoine naturel (SNIP La Réunion; borbonica.re), and from the Early Detection & Distribution Mapping System (EDDMaps; eddmaps.org) and from published and unpublished literature (Augros et al., 2019; Leaché et al., 2017). For the native range, we only used data from Leaché et al. (2014, 2017) to avoid the inclusion of misidentified agamas. We complemented those data with further recent opportunistic field observations from Réunion Island and a dedicated survey in the Comoros. The latter was conducted across the entire island of Grande Comore between June and November 2024, where administrative regions were surveyed (Soule et al., in press). We performed our sampling across a range of habitats to discriminate suitable and non-suitable habitats. In each location, presence data were gathered (when available) along trails, roads, or along line transects established both in intact and degraded natural forests. Whenever possible, we selected routes that crossed different habitat types (e.g., urban areas, agricultural land, or forest)

to maximise the likelihood of detecting agama across a variety of environments within each

geographical area.

This results in an initial sample size of unique point coordinates of 4140 points (native + non-native

ranges). To avoid pseudo-replication, we resampled one point per pixel of environmental data at

1 km resolution (i.e. a process of spatial data thinning; (Steen et al., 2021), resulting in a final

sample of 1289 points (90 from the native range and 1199 from the non-native range).

2.3 Climate data – Predictions of climate suitability, as well as invasion risks, can be highly

sensitive to the choice of climate data source (Dubos et al., 2023a, 2022a). We used 19 bioclimatic

variables (description available at https://www.worldclim.org/data/bioclim.html; see also Booth et

al., 2014) at 30 arc seconds (approximately 1 × 1 km) resolution for the current and future (2070)

climate from CHELSA version 1.2 (Karger et al., 2017). There is a significant uncertainty related to

the source of climate data in ENM, and it is recommended to use multiple climate data sources

(Baker et al., 2016; Dubos et al., 2022a). Worldclim is the only alternative available at the global

scale in high resolution (1 × 1 km) and with projections of future conditions for multiple scenarios

and Global Circulation Models, GCMs (Fick and Hijmans, 2017). However, Worldclim is known to

be less reliable in small oceanic islands (IPCC, 2021), which is why we preferred to use CHELSA

only, which better takes into account orographic factors.

For future projections, we used three commonly used GCMs (GCMs; BCC-CSM1-1, MIROC5, and

HadGEM2-AO) to account for uncertainty in the methodology. We used two greenhouse gas

emission scenarios (one optimistic RCP45-SSP2 and the most pessimistic one RCP85-SSP5) to

consider a wide panel of possible environmental change by 2070 (SSP1 being now considered

unrealistic and therefore, no longer recommended; (IUCN Standards and Petitions Committee,

2024).

2.4 Land use data – We used current data and future projections of land use at the global scale

(resolution 0.05°, approximately 5 × 5 km; (Chen et al., 2020). To represent the overall absence of

the species in forests and increase the realism of our projections, we performed a post-hoc filtering

to remove all forested areas from the predictions of invasion risks. Provided the species is strongly

associated (either negatively or positively) with a given habitat type, using a filter (or a mask)

enables to account for habitat requirements without adding any habitat predictor to the models,

therefore preserving statistical parsimony. We used current (2015) and future (2070) ensemble

model projections (i.e. the mean projection across five GCMs) for the same scenarios as our climate

data (RCP45-SSP2 and RCP85-SSP5; Fig. S1). We selected all predictors of temperate and tropical

forest cover (expressed as percentages), i.e. temperate evergreen trees, tropical evergreen trees,

temperate evergreen trees, tropical deciduous trees and temperate deciduous trees. We summed up

all types of forest cover to produce a single map of percentage of tree cover for each projection

(current, future SSP2 and future SSP5). To match the resolution of our projections (1 km), we

resampled the tree cover maps with a bilinear interpolation. We set the predictions of invasion risks

to zero where tree cover was over 95% (we only removed pixels which were nearly filled with

forest because the species can be found in forest edges and glades, and because the vast majority of

presence points were located in pixels with < 95% tree cover; Fig S2). To visualize the effect of

future tree cover change (i.e. the difference between future and current tree cover), we used a mask

of current and future tree cover, respectively, on our projections of future invasion risk, for each

corresponding scenario.

2.5 Factors of spread – To account for the risk of introduction and spread of the species, we used

anthropogenic predictors, which we assumed to be related to propagule pressure and spread. The

species has been introduced accidentally through transportation of goods, notably horticultural

exports and shipments of consumable goods (Gray, 2020; Moore, 2019; van den Burg et al., 2024),

and spread within islands through transport of construction material (M.T. Ibouroi pers obs. 2024).

We used three types of drivers of propagule pressure and spread as proxies for factors of

introduction and spread. We used distance to port and airports as a proxy for introduction risk. We

obtained port data from the World Port Index (https://msi.nga.mil/Publications/WPI, accessed

December 2024) and airport data from the OpenFlights Airport database

(https://openflights.org/data.html, accessed December 2024). These data were computed as the

Euclidean distance between all pixels and the closest port / airport. We used primary road density as

an indicator of potential spread facilitation (Lanner et al., 2022), because the species is known to

spread through vehicular transportation of goods. We used data from the Global Roads Inventory

Project (GRIP4) dataset (Meijer et al., 2018). We included human population density as a proxy for

both introduction and expansion risks, since the species has been intentionally released in the wild

or escapes from pet owners (e.g. in Florida; Gray 2020). We obtained human population density

data for the current period and all four future scenarios from Gao (2020) at a resolution of 1×1

km². We were unable to include future scenarios for distance to ports, airports and roads because

such data are not available; thus, our projections assumed no change in these predictors.

2.6 Background selection – It is recommended to use a restricted background to mitigate the effect

of sample bias, follow the hypotheses of past dispersal and account for the Equilibrium hypothesis

(Acevedo et al., 2012; Dubos et al., 2022c; Vollering et al., 2019). In the case of invasive species

that are still spreading, it is preferable to exclude areas that are potentially suitable to the species but

have not yet been reached to avoid generating of pseudo-absences in these zones and prevent false

negatives (i.e. suitable environments predicted as unsuitable). We defined the background as a

rectangle of minimal surface that includes all presence points, which we extend by 10 % in each

cardinal direction. This results in a map which includes Florida, central Africa, the Comoros

Archipelago, Réunion Island and Mauritius (Fig. S3). After model training on this background

extent, we projected the predictions of invasion risk on a global map, excluding cold temperate

climates from both hemispheres (latitude > |40°|).

2.7 Pseudo-absence selection and sample bias correction – We generated 40,000 pseudo-absences with the spatSample function of the terra package (Hijmans, 2025). This number represents a fair trade-off between model reliability and computation time (Valavi et al., 2021). We generated three sets of pseudo-absence. We applied a sample bias correction technique consisting of a selection of pseudo-absence points that imitate the spatial bias of presence points (Phillips et al., 2009). To do so, we used a null geographic model, which is the most appropriate approach for species that are still spreading and have not reached equilibrium, because it downplays the areas that have not been invaded yet and therefore mitigates the effect of non-equilibrium. We used the null geographic model as a probability weight for pseudo-absence selection (Fig. S3).

2.8 Predictor selection – The selection of environmental predictors is an unresolved issue in SDM (Leroy, 2023). To follow the latest methodological recommendations (Araújo et al., 2019; Gábor et al., 2019; Sillero et al., 2021; Sillero and Barbosa, 2021), we performed a predictor selection process in five steps: (1) removing predictors with low variability in the study region, therefore unlikely to explain the presence of the species (Araújo et al., 2019), (2) removing collinear predictors (Dormann et al., 2013), (3) keeping predictors with the clearest causal relationship with species presence and spread (Dubos et al., 2022d; Fourcade et al., 2018; Hui, 2022), (4) selecting predictors with the highest relative importance (Bellard et al., 2016; Thuiller et al., 2009), and (5) considering potential interactive effects (Gábor et al., 2019). (1) We discarded bio3 (isothermality) because the relationship with species presence was not straightforward (bio3 being an interaction between daily and yearly variation in temperature), and displayed little variability on small islands (hence, yielding little capacity to explain species presence). We kept the remaining 18 variables because they all may explain species distribution (mostly through direct effects of temperature and water balance on physiology, or indirect effects on ecosystem productivity and food availability (Dubos et al., 2018). (2) For each climate data source and for each species-specific background

(details below), we selected one predictor variable per group of inter-correlated variables to avoid

collinearity (Pearson's r > 0.7; Fig. S4) using the removeCollinearity function of the virtualspecies

R package (Leroy et al., 2016). (3) When mean values were collinear with extremes, we selected the

variables representing extreme conditions (e.g., warmest/driest condition of a given period) because

these are more likely to drive mortality and local extirpation and be causally related to the presence

of the species (Maxwell et al., 2019; Mazzotti et al., 2016). (4) We excluded the predictors with the

lowest relative importance (assessed by the GINI index of the RandomForest R package). Relative

importance was assessed with centred and scaled predictors (Fig. S5). (5) The effect of temperature

can differ according to the level of drought (Dubos et al., 2019). To consider potential interactive

effects between temperature and precipitation while accounting for propagule pressure, we kept a

final set of two predictors related to temperature and two related to precipitation (in addition to the

selected factors of spread). The latter step implies that we may re-qualify predictors which were

formerly excluded.

2.9 Modelling technique – We modelled and projected climatic niches using two algorithms

evaluated as the best performing ones (Valavi et al., 2023, 2021), i.e. Random Forest down-sampled

(hereafter RF down-sampled), i.e. RF parametrised to deal with a large number of background

samples and few presence records (Prasad et al., 2006) and MaxEnt (Phillips et al., 2009). We set

RF down-sampled to run for 1000 bootstrap samples/trees, each iteration randomly samples the

same number of pseudo-absences as the number of presences, with replacement permitted. For

MaxEnt, we allowed all available types of relationships except threshold ones, because these are

more likely to be driven by overfit. After discarding poorly performing models (see *Model*

evaluation section), we averaged all projections across replicates.

2.10 Model evaluation – We used block-cross validation approach which better assesses model

transferability compared to random partitioning (Valavi et al., 2023, 2019). We spatially partitioned

our data into four folds using blocks of 500 × 500 km (Fig. S6). We assessed model spatial transferability with the Boyce Index (CBI; Hirzel et al., 2006), because other discrimination metrics (e.g. AUC and TSS) require absence data and may be misleading with biased presence-only data (Dubos et al., 2022d; Leroy et al., 2018), especially in the case of IAS at non-equilibrium. An index of 1 suggests that suitability models predicted perfectly the presence points, a zero value suggests that models are no better than random and a negative value implies a counter prediction. We discarded all models with a Boyce index < 0.5. We present projections of the mean suitability obtained across all highly transferable models (total number of models: 4 block-cross validation runs × 3 pseudo-absence runs × 2 modelling techniques = 24; for future projections: 24 × 2 scenarios × 3 GCMs = 144 projections).

2.11 Quantifying the effect of environmental change – We quantified the projected change in environmental suitability at the presence points (a) of the native range and (b) of the three invaded regions separately. We extracted the predictions of current and future invasion risk to represent the effect of environmental change on current populations. To highlight the effect of projected deforestation, we extracted suitability scores from the forecast maps (future projections) which were filtered with current (2015) and future (2070) tree cover (see *Land use data* sub-section).

2.12 Quantifying climate niche overlap – We assessed the level of similarity between the climate conditions of the native and the non-native regions. We computed a Principal Component Analysis between the selected climate predictors (see *Predictor selection* subsection) of the native and non-native ranges following Broennimann et al. (2012). For each of the three invaded regions (Florida-Caribbean, Comoros and Mascarenes), we quantified niche overlap with Schoener's D overlap index and visualized both climate niches (as represented by two principal components) using the *ecospat.plot.niche.dyn* function of the ecospat R package (Di Cola et al., 2017).

3. Results

We selected nine predictors (five climate variables and four factors of introduction and spread; Table 1). All models were highly transferable, with a mean Boyce index of 0.92 ± 0.09 SD. No model was discarded.

The species was absent from regions with a high temperature seasonality (Fig. S7). It established in warm regions (between 25 and 40°C of maximum summer temperature, Bio5), although the predicted values showed multiple modalities, reflecting differences between occupied regions. *Agama picticauda* is only present in regions with warm winters (> 13°C) and is absent from extremely arid regions (rainfall < 10 mm for the driest month, and < 100 mm for the warmest quarter; Fig. S7). Invasion risks are the highest near ports and airports (decreasing rapidly beyond 50 km), and are lowest in regions with low primary road densities and human populations. We identify several regions of high invasion risk that have not been reached yet, i.e. the Caribbean, East and Southeast Asia, the Southeast South American urban corridor (corresponding to the MERCOSUR core economic region), Madagascar, Australia and Pacific Islands (Fig. 1).

352 **Table 1**. Selected environmental predictors of invasion risk for *Agama picticauda*, with expected

353 responses and hypotheses of causal relationship with the species presence.

Selected predictor	Expected species response	Causality with species presence	Reference
Bio4 (temperature seasonality)	Higher presence in low seasonality	Adaptation to humid tropical regions, effects on survival, reproduction and phenotype-related fitness	Steele & Warner, 2020
Bio5 (maximum temperature)	High presence in warm regions	Adaptation to Savannah climates	Leaché et al. 2014
Bio6 (minimum temperature)	Avoids the coldest regions	Effect on survival and sex-ratio	Steele & Warner, 2020
Bio14 (precipitation of driest month)	Avoids the most arid regions	Higher hydric stress, lower food availability on arid conditions	Dubos et al., 2018
Boi18 (precipitation of warmest quarter)	Interactive effects with maximum temperature	Effects on water balance	Dupoué et al., 2017
Distance to closest port	Higher presence near ports	The species is introduced through maritime transportation of goods	Guillermet et al., 1998
Distance to closest airport	Higher presence near airports	The species is introduced with the pet trade through aerial shipments and escapes may occur; high traffic of material around airports	Krysko et al., 2016; Wagner et al., 2009
Primary road density	Higher presence in road-dense areas	The species is spread through terrestrial transportation of goods	Gray 2020
Human population	Higher presence in populated areas	Human infrastructures facilitate species establishment; intentional releases and unintentional escapes from pet owners and breeders	Gray 2020

Fig. 1 Global invasion risk of *Agama picticauda* under current conditions, projected from an ensemble model which accounts for climate, habitat and anthropogenic factors of introduction and spread. Red: high invasion risk; Green: low invasion risk.

The combined effects of climate, land use and human population changes was variable through space (Fig. 2, 3, 4). Invasion risk may increase in human-populated areas, especially in specific locations of Central America, Mayotte, Réunion Island, Japan and South America. Future environmental change will decrease invasion risks in Florida and China overall (Fig. 2). At the location of current populations, invasion risk will vary differently between regions (Fig. 3). The risk will increase in Comoros, decrease in Florida, and become more variable in the Mascarenes (Fig. 3). Tree cover will vary in either direction in the future and depend on the scenario. It will increase in China and Florida (especially in the SSP5 scenario), which will contribute to the mitigation of *A. picticauda* propagation (Fig. 4). Deforestation will create some new areas at risk, but mostly in areas that are not suitable. In regions with a high environmental suitability, deforestation will represent a small surface and are hard to perceive compared to climate and anthropogenic change (Fig. 4).

The niche overlap analysis revealed a high overlap between the native and the non-native regions, as shown by the blue areas of figure 5. In non-native regions, the species is present in climates that are not represented in the native region (solid red areas), which provides evidence for niche expansion. In the Mascarenes and Comoros, some regions share the same conditions as found in the

native range, but have not been invaded yet (indicated by the solid green areas; Fig. 5). This

suggests that there is room for further spread in both regions. In Florida on the other hand, most of

the conditions that are represented in the native regions are occupied, suggesting that the known

realized niche is filled. However, the species has expanded its niche beyond the conditions available

in the native range (light red) and might keep spreading. A significant part of the niche overlap

between the native and invaded regions is related to anthropogenic factors of spread (i.e. human

population and primary road density; Fig. S8).

Fig. 2 Effect of future environmental change (climate, tree cover and human population) on

projections of global invasion risk by *Agama picticauda*. We show two scenarios (A: SSP2, B:

SSP5) for the 2061-2080 period.

Fig. 3 Predictions of current and future invasion risk (2070, two scenarios) by *Agama picticauda* at

the current populations of the native (West-Central Africa) and the three invaded regions (the

Comoros archipelago, Florida-Caribbean and the Mascarenes). Invasion risk was predicted from a

combination of climate and habitat suitability, accounting for anthropogenic factors of spread.

Boxes are composed of the first decile, the first quartile, the median, the third quartile and the ninth

decile. Each point is an occupied 1 km²-pixel.

Fig. 4 Predictions of invasion risk by *Agama picticauda* for four regions of interest (left panels: current, middle panels: future, SSP2; right panels: future, SSP5). We show the density of predicted invasion risk for each region and scenario, accounting for tree cover change (red) and assuming no change in tree cover (yellow).

Fig. 5 Realized climate niche of *Agama picticauda* obtained from Principal Component Analysis (five climate predictors and four anthropogenic factors of spread: bio4, bio5, bio6, bio14, bio18, distance to ports, distance to airports, primary road density and human population; see predictor contribution in figure S8). Each panel compares an invaded region with the native region (Florida-Caribbean, Mascarenes and Comoros). Green: climate niche of the native range (contour: environmental conditions of the native range; light green: occupied regions with conditions represented in the native region only; solid green: unoccupied regions of the non-native area with conditions that are represented in the native range); red: conditions of the non-native range (contour: environmental conditions of the non-native range, solid red: niche expansion under common conditions between the native and non-native regions, light red: niche expansion in novel conditions; blue: niche stability (i.e. overlapping conditions). The shades of black represent the density of occurrence points at a given set of principal component (PC) score.

4. Discussion

We predicted the current and future global invasion risks of *Agama picticauda*, accounting for climate, habitat and anthropogenic factors of introduction and spread. Global change will affect invasion risk, but in different direction and magnitude depending on the region. Our niche analysis indicated little room for further invasion in Florida and the risk should decrease in the future. This contrasts with the Comoros, where the species may keep spreading, and where we predict an aggravation of the risk. The risk will become more variable in the Mascarenes. We identify priority areas for proactive surveillance and highlight the need for region-specific management guidelines.

4.1 Drivers of invasion

4.1.1 Anthropogenic factors – Factors of human-driven introduction and spread were the most influential predictors of the current distribution of *A. picticauda*, which is a common finding in studies of invasion risk (Bellard et al., 2016b; Lanner et al., 2022). The large amount of observation recorded from Florida, along with the important pet trade in this region may have largely contributed to the strong explanatory power of distance to ports and airports. The high density of presence points around airports suggests that many cases of introduction might result from escapes from the pet trade, as formerly observed for other marketed reptiles (e.g. *Phelsuma* spp.; Dubos et al., 2014). This can also be related to incidental introductions through the ornamental plant trade, as is the case for the green iguana (Villanueva et al., 2025). However, the first reports of *A. picticauda* in the three invaded regions were not in airports directly. In this specific case, the importance of anthropogenic factors was possibly due to intense vehicular movement of goods and vehicle traffic between highly populated urban areas and airports. The effect of human population (i.e. absence in unpopulated areas) may be related to both the habitat use of the species and intentional and unintentional releases. *A. picticauda* is commonly found in human-modified habitats, where it can find shelter and food items (Mitchell et al., 2021). The increasing availability of human structures may benefit the species in the future and increase invasion risk in the modified regions (Gray 2020).

Invasion risks will increase at locations where human population is projected to increase, suggesting that anthropogenic factors may increase the chances of introduction and spread. This contrasts with projections of climate change effects, which echoes another predictive study of invasion risks in the tropics (Dubos et al., 2025).

In the Comoros, the spread and colonization of new habitats by *A. picticauda* appear to be closely linked to urbanization and the expansion of built infrastructure. One of the main vectors of dispersal is the inter-regional transport of construction materials, particularly sand, where the species lays its eggs. Additionally, porous concrete blocks used in construction often harbour eggs, juveniles, or even adult individuals, thus facilitating unintentional translocation. Although the species was initially introduced in the capital city of Moroni, it is now distributed across nearly all regions of the island. These findings suggest that human-mediated transport, especially through the construction sector, plays a key role in the dissemination of the species across the island landscape.

4.1.2 Precipitation – The species was strongly influenced by summer precipitation, presumably due to water balance requirements during the warmest period. It is also possible that the avoidance of the most arid environments is related to indirect effects, through lower primary and secondary production, thus limiting food availability (Dubos et al., 2018). A decrease in summer precipitation might affect evaporative water loss in the hottest regions and impact negatively the persistence of the species.

4.1.3 Temperature – *Agama picticauda* is currently distributed in regions with low temperature seasonality. This is consistent with the native region of the species, which is characterized by a tropical savannah climate (Peel et al., 2007). Fluctuating temperatures can affect sex-dependent survival and sex ratio (Steele and Warner, 2020). An increase in temperature seasonality may disrupt sex-dependent determination (towards more females; Steele and Warner, 2018) and affect population dynamics in the future. The establishment of the species in Florida may therefore seem surprising given the low temperatures occasionally occurring during winter. This suggests a high potential for behavioural or physiological plasticity or adaptation to new environments (Card et al.,

2018; Fieldsend et al., 2021; Lapwong et al., 2021). Ecological release may also offer opportunities to colonize colder regions. This is the case for *Hemidactylus frenatus, Anolis sagrei, Leiocephalus carinatus* and *Phelsuma grandis*, which were able to spread in areas colder than their native range (Angetter et al., 2011; Claunch et al., 2021; Dubos et al., 2023a; Lapwong et al., 2021). Although our niche analysis indicated that the species already occupies all the environmental conditions that are represented in its native range, the width of its fundamental niche remains unknown, and *A. picticauda* may keep spreading towards novel conditions. This is supported by recent findings in new, colder locations (although establishment needs to be confirmed; Enge, 2024), and by the ability of reptiles to adapt or acclimate to colder conditions after experiencing cold snaps (Stroud et al., 2020).

4.1.4 Tree cover – We found that change in tree cover will not significantly increase invasion risk from a macroecological point of view. Most of the regions with projections of deforestation are located in regions of low environmental suitability. Tree cover is projected to increase in some regions (e.g. north Florida and eastern China), which might mitigate the spread. However, deforestation might create new areas available for establishment at the local scale in specific regions (e.g. In the Lesser Antilles; van den Burg et al., 2024). Accounting for tree cover may have increased the realism of our projections of suitable macroecological conditions. However, the resolution of our study (1 km²) may be limiting at the local scale, because *A. picticauda* is observed at the edge of large forests and in the surroundings of small tree patches, which cannot be perceived with our data. In addition, future projections may be uncertain, especially in regards to changes in local policies and stochastic events (e.g. in Comoros, where tree cover has been decimated across the archipelago in 2024 after a cyclone).

4.2 Niche analysis

We found that *A. picticauda* has colonized all the environmental conditions that are represented in the native region in Florida, including regions where the species is absent. This supports that the species was constrained by biotic factors in the native region and suggests that niche expansion may

be the result of ecological release (Fieldsend et al., 2021). Nonetheless, the species has extended its niche even beyond the conditions that are represented in Africa, suggesting either that the fundamental niche of the species is largely underestimated, or that the niche was enlarged through in situ adaptation (Fieldsend et al., 2021). In our specific case, a large proportion of the overlap is driven by road density and human population (Fig. S8), which emphasizes the importance of anthropogenic factors in explaining the current global distribution, especially in Florida. The situation contrasts in Comoros where we observe no niche expansion, and in the Mascarenes where niche expansion was low. The absence of niche expansion is presumably due to similar climates between western Africa and the archipelagos, along with the smaller size of the islands that limits potential variation in climate and habitats compared to continental regions. Nonetheless, we found a high potential for further spread in both regions. We showed that the Comoros may become completely invaded, since the available environmental conditions are all suitable to the species.

Although *A. picticauda* displays traits commonly associated with invasive species, such as ecological flexibility, territorial aggression and strong affinity for human-modified environments, there is hitherto little empirical evidence of direct negative ecological impacts. While *A. picticauda* has been identified as a potential threat to native herpetofauna, especially on islands with high endemism, there is currently no documented evidence linking its presence to confirmed population declines (van den Burg et al., 2024). This differs from arboreal IAS such as *Phelsuma laticauda* in Réunion Island (Deso et al., 2023), where there is greater niche overlap with endemic reptiles. However, the presence of *A. picticauda* on forest edges could affect arboreal species on small patches, which constitutes the majority of the habitat of some endemic species such as *P. inexpectata* (Dubos et al., 2022c). Native species would be better preserved at the core of larger forest patches, which supports the creation of large tree patches as a potential management solution.

4.3 Management recommendations

4.3.1 Non-invaded regions – We identified priority for proactive surveillance at the global scale in

regions that are still untouched by the species. Our models support that incidental escapes from the

pet trade might be a major driver of invasion risk. We recommend to (1) increase biosecurity

measures to promote early detection in ports and airports and (2) implement strong trade regulations

to prevent further introductions, such as the banishment of *A. picticauda* from the pet trade in

regions that are the most at risk (e.g. in the South American urban corridor, East and Southeast

Asia). We recommend informing all stakeholders linked to the pet trade (including port managers,

maritime transit, pet traders and pet owners) of the risk of escape and the subsequent ecological and

economic hazards.

4.3.2 Invaded regions – In Florida in particular, introductions are largely caused by deliberate

releases from pet owners. This suggests that stronger regulations and public awareness may be a

worthy investment to mitigating the risk in this region. Urban development is high in this region,

which tends to create further opportunities for establishment (Clements et al., 2025). The species

largely benefits from human structures and resources, either through the availability of shelters and

food items, or through the low abundance of competitors (Mitchell et al., 2021). A key to slowing

down the spread of the species is to promote the restoration of natural habitats. Such intervention

may benefit both the native flora and fauna while reducing the amount of available space for *A.*

picticauda.

In Comoros, the species is largely spread through transportation of building material. This calls for

the need to integrate biosecurity into urban planning and increase surveillance between construction

sites, logistic platforms and factories. Although representing a small surface at the global scale,

deforestation may facilitate the establishment and spread of the species. We recommend the

prevention of deforestation in the most sensitive areas (e.g. in the Caribbean, Mauritius and

Mayotte). An increase in tree cover may be another potential way to mitigate the risk, provided

native tree species are promoted. To mitigate the exposition of native arboreal species, the creation

of large patches should be preferred over smaller patches. In regions that are exposed to cyclone

activity such as the Comoros, extreme winds can deforest large areas and create colonization

opportunities for *A. picticauda*. It is urgent to facilitate tree plantation in order to restore native

habitats and prevent new establishments. Global change will increase the risk in the Comoros and

Mascarene archipelagos, which stresses the need to account for future environmental suitability in

habitat restoration plans. We recommend the development of monitoring programmes for *A.*

picticauda populations, and the study of the response of the native vegetation to environmental

change. Implementing these recommendations will require coordinated efforts among stakeholders,

targeted awareness campaigns, and integration into local land-use planning and trade regulation

frameworks.

Acknowledgements

We are grateful to all the staff of Nature Océan Indien for their help in data collection.

CRediT authorship contribution statement

Nicolas Dubos: Conceptualization, Data Curation, Formal analysis, Methodology, Investigation,

Supervision, Writing – original draft. **Steven Calesse**: Data Curation, Formal analysis,

Investigation. **Kathleen C. Webster**: Data Curation, Investigation, Writing – review and editing.

Chloé Bernet: Supervision, Project administration, Funding acquisition, Writing – review and

editing. **Gregory Deso**: Data Curation, Investigation, Writing – review and editing. **Thomas W.**

Fieldsend: Investigation, Writing – review and editing. **Saoudati Maoulida**: Data curation,

Writing – review and editing. **Soule Mounir**: Data curation, Writing – review and editing. **Xavier**

Porcel: Data Curation, Investigation, Writing – review and editing. **Jean-Michel Probst:** Data

curation, Investigation, Writing – review and editing. **Hindatou Saidou**: Data curation, Writing –

review and editing. **Jérémie Souchet**: Supervision, Project administration, Investigation, Writing

– review and editing. **Mohamed Thani Ibouroi**: Conceptualization, Data Curation, Investigation,

Supervision, Writing – review and editing. **Markus A. Roesch**: Conceptualization, Data Curation,

Methodology, Investigation, Supervision, Funding acquisition, Writing – review and editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships

that could have appeared to influence the work reported in this paper.

References

Acevedo, P., Jiménez-Valverde, A., Lobo, J.M., Real, R., 2012. Delimiting the geographical
background in species distribution modelling. J. Biogeogr. 39, 1383–1390.
https://doi.org/10.1111/j.1365-2699.2012.02713.x

Angetter, L.S., Lötters, S., Rödder, D., 2011. Climate niche shift in invasive species: The case of the
brown anole. Biol. J. Linn. Soc. 104, 943–954. https://doi.org/10.1111/j.1095-
8312.2011.01780.x

Araújo, M.B., Anderson, R.P., Barbosa, A.M., Beale, C.M., Dormann, C.F., Early, R., Garcia, R.A.,
Guisan, A., Maiorano, L., Naimi, B., O'Hara, R.B., Zimmermann, N.E., Rahbek, C., 2019.
Standards for distribution models in biodiversity assessments. Sci. Adv. 5, 1–12.
https://doi.org/10.1126/sciadv.aat4858

Augros, S., Faipoux, L., Bodin, M., Goff, A. Le, Sanchez, M., 2017. Evidence for colonisation of
anthropogenic habitats by the Réunion day gecko Phelsuma borbonica (Mertens , 1966)
(Réunion Island , France): conservation implications 10, 563–571.

Augros, S., Hawlirschek, O., Fabulet, P.-Y., Eudeline, R., 2019. Atlas des reptiles et amphibiens
terrestres de l'archipel des Comores. Biotope - Muséum National d'Histoire Naturelle.

Baker, D.J., Hartley, A.J., Butchart, S.H.M., Willis, S.G., 2016. Choice of baseline climate data
impacts projected species' responses to climate change. Glob. Chang. Biol. 22, 2392–2404.
https://doi.org/10.1111/gcb.13273

Barbet-Massin, M., Rome, Q., Villemant, C., Courchamp, F., 2018. Can species distribution models
really predict the expansion of invasive species? PLoS One 13, 1–14.
https://doi.org/10.1371/journal.pone.0193085

Bellard, C., Leroy, B., Thuiller, W., Rysman, J.F., Courchamp, F., 2016. Major drivers of invasion
risks throughout the world. Ecosphere 7, 1–14. https://doi.org/10.1002/ecs2.1241

Bellard, C., Thuiller, W., Leroy, B., Genovesi, P., Bakkenes, M., Courchamp, F., 2013. Will climate change promote future invasions? Glob. Chang. Biol. 19, 3740–3748. https://doi.org/10.1111/gcb.12344

Booth, T.H., Nix, H.A., Busby, J.R., Hutchinson, M.F., 2014. Bioclim: The first species distribution modelling package, its early applications and relevance to most current MaxEnt studies. Divers. Distrib. 20, 1–9. https://doi.org/10.1111/ddi.12144

Broennimann, O., Fitzpatrick, M., Pearman, P., Petitpierre, B., Pellissier, L., Yoccoz, N.G., Thuiller, W., Fortin, M.-J., Randin, C., Zimmermann, N.E., Graham, C.H., Guisan, A., 2012. Measuring Ecological Niche Overlap From Occurrence. Development 1–34.

Card, D.C., Perry, B.W., Adams, R.H., Schield, D.R., Young, A.S., Andrew, A.L., Jezkova, T., Pasquesi, G.I.M., Hales, N.R., Walsh, M.R., Rochford, M.R., Mazzotti, F.J., Hart, K.M., Hunter, M.E., Castoe, T.A., 2018. Novel ecological and climatic conditions drive rapid adaptation in invasive Florida Burmese pythons. Mol. Ecol. 27, 4744–4757. https://doi.org/10.1111/mec.14885

Chapman, D., Prescott, O.L., Roy, Helen, E., Tanner, R., 2019. Improving species distribution models for invasive non-native species with biologically informed pseudo-absence selection. J. Biogeogr. 46, 1029–1040.

Chen, M., Vernon, C.R., Graham, N.T., Hejazi, M., Huang, M., Cheng, Y., Calvin, K., 2020. Global land use for 2015–2100 at 0.05° resolution under diverse socioeconomic and climate scenarios. Sci. Data 7, 1–11. https://doi.org/10.1038/s41597-020-00669-x

Claunch, N.M., Goodman, C., Reed, R.N., Guralnick, R., Romagosa, C.M., Taylor, E.N., 2021. Invaders from islands: Thermal matching, potential or flexibility? Biol. J. Linn. Soc. 134, 587–603. https://doi.org/10.1093/biolinnean/blab103

Clements, S., Ackerman, F.M., Olensky, I.M., White, E.C., Rogers, M.E., Searcy, C.A., 2025. Non-native herpetofauna continue to proliferate in the world's most invaded herpetofauna community. Ecol. Evol. 15, e71556.

Cuthbert, R.N., Diagne, C., Hudgins, E.J., Turbelin, A., Ahmed, D.A., Albert, C., Bodey, T.W., Briski, E., Essl, F., Haubrock, P.J., Gozlan, R.E., Kirichenko, N., Kourantidou, M., Kramer, A.M., Courchamp, F., 2022. Biological invasion costs reveal insufficient proactive management worldwide. Sci. Total Environ. 819, 153404. https://doi.org/10.1016/j.scitotenv.2022.153404

Deso, G., Roesch, M.A., Porcel, X., Claudin, J., Probst, J., Luspot, J., Dubos, N., 2023. Interactions between the endemic gecko Phelsuma inexpectata and the introduced Phelsuma laticauda: understanding the drivers of invasion on Reunion island. Herpetol. Bull. 165, 21–23.

Di Cola, V., Broennimann, O., Petitpierre, B., Breiner, F.T., D'Amen, M., Randin, C., Engler, R., Pottier, J., Pio, D., Dubuis, A., Pellissier, L., Mateo, R.G., Hordijk, W., Salamin, N., Guisan, A., 2017. ecospat: an R package to support spatial analyses and modeling of species niches and distributions. Ecography (Cop.). 40, 774–787. https://doi.org/10.1111/ecog.02671

Diagne, C., Leroy, B., Vaissière, A.C., Gozlan, R.E., Roiz, D., Jarić, I., Salles, J.M., Bradshaw, C.J.A., Courchamp, F., 2021. High and rising economic costs of biological invasions worldwide. Nature 592, 571–576. https://doi.org/10.1038/s41586-021-03405-6

Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B., Lafourcade, B., Leitão, P.J., Münkemüller, T., Mcclean, C., Osborne, P.E., Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013. Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. Ecography (Cop.). 36, 27–46. https://doi.org/10.1111/j.1600-0587.2012.07348.x

Dubos, N., 2013. New locality record for Phelsuma grandis (Sauria: Gekkonidae) in Reunion, in sympatry with the Critically Endangered Phelsuma inexpectata. Herpetol. Notes 6, 309–311.

Dubos, N., Augros, S., Deso, G., Probst, J., Notter, J., Roesch, M., 2022a. Here be dragons: important spatial uncertainty driven by climate data in forecasted distribution of an endangered insular reptile. Anim. Conserv. 25, 704–717. https://doi.org/10.1111/acv.12775

Dubos, N., Dehorter, O., Yves, P., Viol, I. Le, 2019. Thermal constraints on body size depend on the population ' s position within the species ' thermal range in temperate songbirds. Glob. Ecol. Biogeogr. 28, 96–106. https://doi.org/10.1111/geb.12805

Dubos, N., Fieldsend, T.W., Roesch, M.A., Stephane, A., Aurélien, B., Arthur, C., Ineich, I., Kenneth, K., Boris, L., Malone, S.L., Probst, J., Raxworthy, C.J., Crottini, A., 2023a. Choice of climate data influences predictions of current and future global invasion risks for two Phelsuma geckos. Biol. Invasions. https://doi.org/10.1101/2022.08.04.502765

Dubos, N., Geslin, B., Jourdan, H., Renault, D., Zakardjian, M., 2025. Contrasting effects of climate and anthropogenic change on future invasion risk of a solitary bee Amegilla pulchra. J. Environ. Manage. 385. https://doi.org/10.1016/j.jenvman.2025.125711

Dubos, N., Havard, A., Crottini, A., Seglie, D., Andreone, F., 2023b. Predicting future conservation areas while avoiding sympatry in two alpine urodele amphibians severely threatened by climate change. J. Nat. Conserv. 76, 126490. https://doi.org/10.1016/j.jnc.2023.126490

Dubos, N., Le Viol, I., Robert, A., Teplitsky, C., Ghislain, M., Dehorter, O., Julliard, R., Henry, P.Y., 2018. Disentangling the effects of spring anomalies in climate and net primary production on body size of temperate songbirds. Ecography (Cop.). 41, 1319–1330. https://doi.org/10.1111/ecog.03413

Dubos, N., Lenormand, M., Castello, L., Oberdorff, T., Guisan, A., Luque, S., 2022b. Protection gaps in Amazon floodplains will increase with climate change: Insight from the world's largest scaled freshwater fish. Aquat. Conserv. Mar. Freshw. Ecosyst. 32, 1830–1841. https://doi.org/10.1002/aqc.3877

Dubos, N., Montfort, F., Grinand, C., Nourtier, M., Deso, G., Probst, J.-M., Razafimanahaka, J.H., Andriantsimanarilafy, R.R., Rakotondrasoa, E.F., Razafindraibe, P., Jenkins, R., Crottini, A., 2022c. Are narrow-ranging species doomed to extinction? Projected dramatic decline in future climate suitability of two highly threatened species. Perspect. Ecol. Conserv. 20, 18–28. https://doi.org/10.1016/j.pecon.2021.10.002

Dubos, N., Piludu, N., Andriantsimanarilafy, R.R., Randrianantoandro, J.C., Andreone, F., 2014. New findings of Phelsuma grandis and P. laticauda (Sauria : Gekkonidae) at the southern edge of the range of the endangered Phelsuma serraticauda in eastern Madagascar. Herpetol. Notes 7, 21–23.

Dubos, N., Préau, C., Lenormand, M., Papuga, G., Monsarrat, S., Denelle, P., Louarn, M. Le, Heremans, S., Roel, M., Roche, P., Luque, S., 2022d. Assessing the effect of sample bias correction in species distribution models. Ecol. Indic. 145, 109487. https://doi.org/10.1016/j.ecolind.2022.109487

Dupoué, A., Rutschmann, A., Le Galliard, J.F., Miles, D.B., Clobert, J., DeNardo, D.F., Brusch, G.A., Meylan, S., 2017. Water availability and environmental temperature correlate with geographic variation in water balance in common lizards. Oecologia 185, 561–571. https://doi.org/10.1007/s00442-017-3973-6

Enge, K., 2024. New county records for Peters's rock agama (Agama picticauda) in Florida, Georgia, South Carolina, Texas, Virginia, and Wisconsin, USA. Herpetol. Rev. 55, 53.

Fick, S.E., Hijmans, R.J., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. Int. J. Climatol. 37, 4302–4315. https://doi.org/10.1002/joc.5086

Fieldsend, T.W., Dubos, N., Krysko, K.L., Raxworthy, C.J., Malone, S.L., 2021. In situ adaptation and ecological release facilitate the occupied niche expansion of a non-native reptile. Ecol. Evol. 11, 9410–9422. https://doi.org/10.1002/ece3.7749

Fourcade, Y., Besnard, A.G., Secondi, J., 2018. Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. Glob. Ecol. Biogeogr. 27, 245–256. https://doi.org/10.1111/geb.12684

Gábor, L., Moudrý, V., Barták, V., Lecours, V., 2019. How do species and data characteristics affect species distribution models and when to use environmental filtering? Int. J. Geogr. Inf. Sci. 00, 1–18. https://doi.org/10.1080/13658816.2019.1615070

Gallien, L., Douzet, R., Pratte, S., Zimmermann, N.E., Thuiller, W., 2012. Invasive species distribution models - how violating the equilibrium assumption can create new insights. Glob. Ecol. Biogeogr. 21, 1126–1136. https://doi.org/10.1111/j.1466-8238.2012.00768.x

Gao, J., 2020. Global 1-km Downscaled Population Base Year and Projection Grids Based on the Shared Socioeconomic Pathways, Revision 01 [WWW Document]. Palisades, New York NASA Socioecon. Data Appl. Cent. https://doi.org/https://doi.org/10.7927/q7z9-9r69.

Gippet, J.M.W., Bertelsmeier, C., 2021. Invasiveness is linked to greater commercial success in the global pet trade. Proc. Natl. Acad. Sci. U. S. A. 118. https://doi.org/10.1073/PNAS.2016337118

Gray, R.J., 2020. Exotic Hobos: Release, escape, and potential secondary dispersal of African red-headed agamas (*Agama picticauda* PETERS, 1877) through the Florida railway systems. https://doi.org/10.1101/2020.05.11.089649

Guillermet, C., Couteyen, S., Probst, J., 1998. Une nouvelle espèce de reptile naturalisée à La Réunion, l'Agame des colons Agama agama (Linnaeus). Bull. Phaethon 8, 67–69.

Harris, D.J., Roesch, M.A., Visconcelos, D.S., Bernet, C., Rato, C., 2025. Metabarcoding assessment of the diet of an introduced continental lizard to an oceanic island reveals dietary niche conservatism. https://doi.org/doi.org/10.21203/rs.3.rs-6830034/v1

Hawlitschek, O., Brückmann, B., Berger, J., Green, K., Glaw, F., 2011. Integrating field surveys and remote sensing data to study distribution, habitat use and conservation status of the herpetofauna of the Comoro Islands. Zookeys 144, 21–79. https://doi.org/10.3897/zookeys.144.1648

Henigan, L.E., Blais, C.B., Dzikowski, N.T., Romagosa, C., Claunch, N.M., 2019. Agama picticauda. Herpetol. Rev. 50.

Hijmans, 2025. terra: Spatial Data Analysis.

Hirzel, A.H., Le Lay, G., Helfer, V., Randin, C., Guisan, A., 2006. Evaluating the ability of habitat suitability models to predict species presences. Ecol. Modell. 199, 142–152. https://doi.org/10.1016/j.ecolmodel.2006.05.017

Hui, C., 2022. The dos and don'ts for predicting invasion dynamics with species distribution models. Biol. Invasions. https://doi.org/10.1007/s10530-022-02976-3

IPCC, 2021. Climate Change 2021 Synthesis Report. Contrib. Work. Groups I, II III to Fifth Assess. Rep. Intergov. Panel Clim. Chang. 1–112. https://doi.org/10.1017/CBO9781107415324

IUCN Standards and Petitions Committee, 2024. Guidelines for Using the IUCN Red List Categories and Criteria. Version 16, Prepared by the Standards and Petitions Committee.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann, N.E., Linder, H.P., Kessler, M., 2017. Climatologies at high resolution for the earth's land surface areas. Sci. Data 4, 1–20. https://doi.org/10.1038/sdata.2017.122

Krishnan, S., Ofori-Boateng, C., Fujita, M.K., Leaché, A.D., 2019. Geographic variation in West African Agama picticauda: insights from genetics, morphology and ecology. African J. Herpetol. 68, 33–49. https://doi.org/10.1080/21564574.2018.1509139

Krysko, K.L., Somma, L.A., Smith, D.C., Gillette, C.R., Cueva, D., Wasilewski, J.A., Enge, K.M., Johnson, S.A., Campbell, T.S., Edwards, J.R., Rochford, M.R., Tompkins, R., Fobb, J.L., Mullin, S., Lechowicz, C.J., Hazelton, D., Warren, A., 2016. New Verified Nonindigenous Amphibians and Reptiles in Florida through 2015, with a Summary of More Than 152 Years of Introductions. IRCF Reptil. Amphib. 23, 110–143.

Lanner, J., Dubos, N., Geslin, B., Leroy, B., Hernández-Castellano, C., Dubaić, J.B., Bortolotti, L., Calafat, J.D., Ćetković, A., Flaminio, S., Le Féon, V., Margalef-Marrase, J., Orr, M., Pachinger, B., Ruzzier, E., Smagghe, G., Tuerlings, T., Vereecken, N., Meimberg, H., 2022. On the road: Anthropogenic factors drive the invasion risk of a wild solitary bee species. Sci. Total Environ. 827, 154246. https://doi.org/10.1016/j.scitotenv.2022.154246

Lapwong, Y., Dejtatadol, A., Webb, J., 2021. Shifts in thermal tolerance of the invasive Asian house gecko (Hemidactylus frenatus) across native and introduced ranges. Biol. Invasions 23, 1–8.

Leaché, A.D., Grummer, J.A., Miller, M., Krishnan, S., Fujita, M.K., Böhme, W., Schmitz, A., Lebreton, M., Ineich, I., Chirio, L., Ofori-boateng, C., Eniang, E.A., Greenbaum, E., Rödel, M.O., Wagner, P., 2017. Bayesian inference of species diffusion in the West African Agama agama species group (Reptilia, Agamidae). Syst. Biodivers. 15, 192–203. https://doi.org/10.1080/14772000.2016.1238018

Leaché, A.D., Wagner, P., Linkem, C.W., Böhme, W., Papenfuss, T.J., Chong, R.A., Lavin, B.R., Bauer, A.M., Nielsen, S. V., Greenbaum, E., Rödel, M.O., Schmitz, A., LeBreton, M., Ineich, I., Chirio, L., Ofori-Boateng, C., Eniang, E.A., Baha El Din, S., Lemmon, A.R., Burbrink, F.T., 2014. A hybrid phylogenetic-phylogenomic approach for species tree estimation in african agama lizards with applications to biogeography, character evolution, and diversification. Mol. Phylogenet. Evol. 79, 215–230. https://doi.org/10.1016/j.ympev.2014.06.013

Leroy, B., 2023. Choosing presence-only species distribution models. J. Biogeogr. 50, 247–250. https://doi.org/10.1111/jbi.14505

Leroy, B., Bellard, C., Dubos, N., Colliot, A., Vasseur, M., Courtial, C., Bakkenes, M., Canard, A., Ysnel, F., 2014. Forecasted climate and land use changes, and protected areas: the contrasting case of spiders. Divers. Distrib. 20, 686–697. https://doi.org/10.1111/ddi.12191

Leroy, B., Delsol, R., Hugueny, B., Meynard, C.N., Barhoumi, C., Barbet-Massin, M., Bellard, C., 2018. Without quality presence–absence data, discrimination metrics such as TSS can be misleading measures of model performance. J. Biogeogr. 45, 1994–2002. https://doi.org/10.1111/jbi.13402

Leroy, B., Meynard, C.N., Bellard, C., Courchamp, F., 2016. virtualspecies, an R package to generate virtual species distributions. Ecography (Cop.). 39, 599–607. https://doi.org/10.1111/ecog.01388

Maxwell, S.L., Butt, N., Maron, M., McAlpine, C.A., Chapman, S., Ullmann, A., Segan, D.B., Watson, J.E.M., 2019. Conservation implications of ecological responses to extreme weather and climate events. Divers. Distrib. 25, 613–625. https://doi.org/10.1111/ddi.12878

Mazzotti, F.J., Cherkiss, M.S., Parry, M., Beauchamp, J., Rochford, M., Smith, B., Hart, K., Brandt, L.A., 2016. Large reptiles and cold temperatures: Do extreme cold spells set distributional limits for tropical reptiles in Florida? Ecosphere 7, 1–9. https://doi.org/10.1002/ecs2.1439

Meijer, J.R., Huijbregts, M.A.J., Schotten, K.C.G.J., Schipper, A.M., 2018. Global patterns of current and future road infrastructure. Environ. Res. Lett. 13. https://doi.org/10.1088/1748-9326/aabd42

Mitchell, T.S., Folt, B., Hall, J.M., 2021. Dumpsters and other anthropogenic structures as habitat for invasive African rock agama lizards in Florida. Biol. Invasions 23, 2689–2693. https://doi.org/10.1007/s10530-021-02537-0

Moore, J.A., 2019. Agama picticauda (Peters's Rock Agama) Dispersal. Herpetol. Rev. 50, 360.

Nuñez, L.P., Krysko, K.L., Avery, M.L., 2016. Confirmation of introduced Agama picticauda in Florida based on molecular analyses. Bull. Florida Museum Nat. Hist. 54, 138–146.

Ofori, B.Y., Ewool, J.M., Asamoah, A., Adarkwah, P.A., Oduro, D., 2023. Diversity and
Composition of Gastrointestinal Helminths and Gram-Negative Bacteria of the West African
Rainbow Lizard (Agama picticauda) in a Human-Modified Landscape: Implications for
Conservation and Zoonosis.

Ofori, B.Y., Martey, P., Attuquayefio, D., 2018. Observations of the west african rainbow lizard,
Agama picticauda peters, 1877, from Ghana feeding on bread. Herpetol. Notes 11, 955–957.

Pearson, D.E., 2024. Biological Invasions: An Overview, Encyclopedia of Biodiversity, Third
Edition: Volume 1-7. Elsevier Ltd. https://doi.org/10.1016/B978-0-12-822562-2.00037-2

Peel, M.C., Finlayson, B.L., McMahon, T.A., 2007. Updated world map of the Köppen-Geiger
climate classification. Hydrol. Earth Syst. Sci. Discuss. 4, 439–473.
https://doi.org/10.1127/0941-2948/2006/0130

Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J., Ferrier, S., 2009.
Sample selection bias and presence-only distribution models: implications for background and
pseudo-absence data. Ecol. Appl. 19, 181–197. https://doi.org/10.1890/07-2153.1

Prasad, A.M., Iverson, L.R., Liaw, A., 2006. Newer classification and regression tree techniques:
Bagging and random forests for ecological prediction. Ecosystems 9, 181–199.
https://doi.org/10.1007/s10021-005-0054-1

Raxworthy, C.J., Ingram, C.M., Rabibisoa, N., Pearson, R.G., 2007. Applications of ecological
niche modeling for species delimitation: a review and empirical evaluation using day geckos
(Phelsuma) from Madagascar. Syst. Biol. 56, 907–23.
https://doi.org/10.1080/10635150701775111

Refsnider, J.M., Des Roches, S., Rosenblum, E.B., 2015. Evidence for ecological release over a fine
spatial scale in a lizard from the White Sands formation. Oikos 124, 1624–1631.
https://doi.org/10.1111/oik.02406

Roesch, M.A., Webster, K.C., Huet, N., Hawlitschek, O., Mohamed, Y., Sauroy-Toucouère, S.,
Wagner, P., Schmitz, A., 2025. Molecular analysis confirms Agama picticauda invasion on
Western Indian Ocean islands. Biol. Invasions 27. https://doi.org/10.1007/s10530-025-03552-1

Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A.M., Böhm, M., Castro-Herrera, F.,
Chirio, L., Collen, B., Doan, T.M., Grismer, L.L., Itescu, Y., Kraus, F., LeBreton, M., Lewin,
A., Martins, M., Maza, E., Meirte, D., Nagy, Z.T., de Nogueira, C.C., Pincheira-Donoso, D.,
Powney, G.D., Sindaco, R., Tallowin, O., Torres-Carvajal, O., Trape, J.-F., Vidan, E., Uetz, P.,
Wagner, P., Wang, Y., David Orme, C.L., Grenyer, R., Meiri, S., 2017. The global distribution
of tetrapods reveals a need for targeted reptile conservation. Nat. Ecol. Evol.

Seebens, H., Blackburn, T.M., Dyer, E.E., Genovesi, P., Hulme, P.E., Jeschke, J.M., Pagad, S.,
Pyšek, P., Winter, M., Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G., Capinha, C.,
Celesti-Grapow, L., Dawson, W., Dullinger, S., Fuentes, N., Jäger, H., Kartesz, J., Kenis, M.,
Kreft, H., Kühn, I., Lenzner, B., Liebhold, A., Mosena, A., Moser, D., Nishino, M., Pearman,
D., Pergl, J., Rabitsch, W., Rojas-Sandoval, J., Roques, A., Rorke, S., Rossinelli, S., Roy, H.E.,
Scalera, R., Schindler, S., Štajerová, K., Tokarska-Guzik, B., Van Kleunen, M., Walker, K.,

Weigelt, P., Yamanaka, T., Essl, F., 2017. No saturation in the accumulation of alien species
worldwide. Nat. Commun. 8, 1–9. https://doi.org/10.1038/ncomms14435

Sillero, N., Arenas-Castro, S., Enriquez-Urzelai, U., Vale, C.G., Sousa-Guedes, D., Martínez-
Freiría, F., Real, R., Barbosa, A.M., 2021. Want to model a species niche? A step-by-step
guideline on correlative ecological niche modelling. Ecol. Modell. 456.
https://doi.org/10.1016/j.ecolmodel.2021.109671

Sillero, N., Barbosa, A.M., 2021. Common mistakes in ecological niche models. Int. J. Geogr. Inf.
Sci. 35, 213–226. https://doi.org/10.1080/13658816.2020.1798968

Steele, A.L., Warner, T.W., 2020. Sex-specific effects of developmental temperature on morphology,
growth and survival of offspring in a lizard with temperature-dependent sex determination.
Biol. J. Linn. Soc. 130, 320–335.

Steele, A.L., Warner, T.W., 2018. Revisiting the first report of temperature-dependent sex
determination in a vertebrate, the African redhead agama. J. Zool. 306, 16–22.

Steen, V.A., Tingley, M.W., Paton, P.W.C., Elphick, C.S., 2021. Spatial thinning and class
balancing: Key choices lead to variation in the performance of species distribution models with
citizen science data. Methods Ecol. Evol. 12, 216–226. https://doi.org/10.1111/2041-
210X.13525

Stroud, J.T., Mothes, C.C., Beckles, W., Heathcote, R.J.P., Donihue, C.M., Losos, J.B., 2020. An
extreme cold event leads to community-wide convergence in lower temperature tolerance in a
lizard community: Convergent shifts in extreme events. Biol. Lett. 16.
https://doi.org/10.1098/rsbl.2020.0625

Thuiller, W., Lafourcade, B., Engler, R., Araújo, M.B., 2009. BIOMOD - A platform for ensemble
forecasting of species distributions. Ecography (Cop.). 32, 369–373.
https://doi.org/10.1111/j.1600-0587.2008.05742.x

Valavi, R., Elith, J., Lahoz-Monfort, J.J., Guillera-Arroita, G., 2023. Flexible species distribution
modelling methods perform well on spatially separated testing data. Glob. Ecol. Biogeogr. 32,
369–383. https://doi.org/10.1111/geb.13639

Valavi, R., Elith, J., Lahoz-Monfort, J.J., Guillera-Arroita, G., 2019. blockCV: An r package for
generating spatially or environmentally separated folds for k-fold cross-validation of species
distribution models. Methods Ecol. Evol. 10, 225–232. https://doi.org/10.1111/2041-
210X.13107

Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J.J., Elith, J., 2021. Predictive performance of
presence-only species distribution models: a benchmark study with reproducible code. Ecol.
Monogr. e01486, 1–27. https://doi.org/10.1002/ecm.1486

van den Burg, M.P., Wasilewski, J.A., Thorpe, R.S., Debrot, A.O., Daltry, J.C., Angin, B., Boman,
E.M., Brannon, L., Corbett, E.A., Horrocks, J.A., Mukhida, F., Paranthoën, N., Pascoe, N.K.,
Petrovic, C., Verhoeven, A., Yokoyama, M., 2024. The threat of Peters's Rock Agama (Agama
picticauda) to reptile diversity across the Lesser Antilles. Anim. Conserv.
https://doi.org/10.1111/acv.12889

Villanueva, C.N.D.J., Prado, G.P.M., Gould, W., García-Quijano, C., Kolbe, J.J., 2025. Interviews
with farmers suggest negative direct and indirect effects of the invasive green iguana (Iguana
iguana) on agriculture in Puerto Rico. Manag. Biol. Invasions 13, 781–797.
https://doi.org/10.3391/mbi.2022.13.4.13

Vollering, J., Halvorsen, R., Auestad, I., Rydgren, K., 2019. Bunching up the background betters
bias in species distribution models. Ecography (Cop.). 42, 1717–1727.
https://doi.org/10.1111/ecog.04503

Wagner, P., Glaw, F., Glaw, K., Böhme, W., 2009. Studies on African Agama IV: First record of
Agama agama (Sauria: Agamidae) from Madagascar and identity of the alien population on
Grande Comore Island. Herpetol. Notes 2, 73–77.

Yackulic, C.B., Chandler, R., Zipkin, E.F., Royle, J.A., Nichols, J.D., Grant, E.H.C., Veran, S.,
2013. Presence-only modelling using MAXENT: when can we trust the inferences? Methods
Ecol. Evol. 4, 236–243. https://doi.org/10.1111/2041-210x.12004

Zurell, D., Franklin, J., König, C., Bouchet, P.J., Dormann, C.F., Elith, J., Fandos, G., Feng, X.,
Guillera-Arroita, G., Guisan, A., Lahoz-Monfort, J.J., Leitão, P.J., Park, D.S., Peterson, A.T.,
Rapacciuolo, G., Schmatz, D.R., Schröder, B., Serra-Diaz, J.M., Thuiller, W., Yates, K.L.,
Zimmermann, N.E., Merow, C., 2020. A standard protocol for reporting species distribution
models. Ecography (Cop.). 43, 1261–1277. https://doi.org/10.1111/ecog.04960