An integrated multi-tissue atlas of epigenomic landscapes and regulatory elements in the bovine genome

Dailu Guan[1,†], Jennifer Jessica Bruscadin[1,2†], Wenjing Yang[1,†], Claire Prowse-Wilkins[1,†], Junjian Wang[3], Bruna Petry[4], Houcheng Li[5], Huicong Zhang[5], Xiaoqin (Shally) Xu[1], Ying Wang[1], Zhangyuan Pan[1], Yuri Utsunomiya[6], Gordon Murdoch[7], Kimberley Davenport[7], Guosong Wang[8], Yue Xing[9], Christian Maltecca[3], Li Ma[10], Gonzalo Rincon[11], Sarah Corum[11], Pengcheng Lyu[12] Joseph (Nader) Deeb[6], Zhiping Weng[13], Jicai Jiang[3], Lingzhao Fang[5], Honglin Jiang[12], Brenda M. Murdoch[14], Stephanie D. McKay[15], Monique Rijnkels[8], Claire A. Gill[8], Timothy P. L. Smith[16], Zhihua Jiang[17], Wansheng Liu[18], Jim Reecy[4], Juan F. Mendrano[1], James E. Koltes[4], Pablo Ross[1], Huaijun Zhou[1]

[1]Department of Animal Science, University of California, Davis, CA, 95616, USA
[2]Center of Biological and Health Sciences, Federal University of Sao Carlos, Sao Carlos, Brazil
[3]Department of Animal Science, North Carolina State University, Raleigh, NC 27695, United States
[4]Department of Animal Science, Iowa State University, Ames, Iowa 50011, USA
[5]Center for Quantitative Genetics and Genomics (QGG), Aarhus University, Aarhus, Denmark
[6]STgenetics, 22515 Highway 6 South, Navasota, Texas 77868
[7]Department of Animal, Veterinary & Food Sciences, University of Idaho, Moscow, ID, USA
[8]Department of Veterinary Integrative Biosciences, School of Veterinary Medicine, Texas A&M University, College Station, TX, 77843, USA
[9]Department of Animal Science, Texas A&M University, College Station, TX 77843, USA
[10]Department of Animal and Avian Sciences, University of Maryland, College Park, MD 20742, USA
[11]Zoetis Inc.
[12]School of Animal Sciences, Virginia Tech, Blacksburg, VA 24060, USA
[13]Program in Bioinformatics and Integrative Biology, University of Massachusetts Chan Medical School, Worcester, MA, USA
[14]Department of Animal, Veterinary & Food Sciences, University of Idaho, Moscow, ID, USA
[15]Division of Animal Sciences, University of Missouri, Columbia, MO, 65211, USA
[16]USDA, ARS, US Meat Animal Research Center, Clay Center, Nebraska, USA
[17]Department of Animal Sciences and Center for Reproductive Biology, Washington State University, Pullman WA 99164-7620, USA.
[18]Department of Animal Science, Center for Reproductive Biology and Health (CRBH), College of Agricultural Sciences, The Pennsylvania State University, University Park, PA, USA

†These authors contributed equally.

Corresponding authors*:

Prof. Huaijun Zhou: Department of Animal Science, University of California-Davis, Davis, CA, 95616, USA

E-mail: hzhou@ucdavis.edu

Emails for all authors:

dguan@ucdavis.edu;

jennifer.j.bruscadin@gmail.com;

wenyang@ucdavis.edu;

c.prowsewilkins@gmail.com;

bpetry@iastate.edu;

jwang259@ncsu.edu;

houcheng.li@qgg.au.dk;

huicongz@qgg.au.dk;

xqxu@ucdavis.edu;

bpetry@iastate.edu;

yuri.utsunomiya@stgen.com;

ucywang@ucdavis.edu;

zhypan01@163.com;

yuri.utsunomiya@stgen.com

gordon.murdoch@wsu.edu

Sarah.Corum@zoetis.com;

Pengcheng@vt.edu;

joseph.deeb@stgen.com

zhipingweng@gmail.com

jicai_jiang@ncsu.edu;

lingzhao.fang@qgg.au.dk;

hojiang@vt.edu;

bmurdoch@uidaho.edu;

stephanie.mckay@uvm.edu;

mrijnkels@cvm.tamu.edu;

clare.gill@ag.tamu.edu;

tim.smith2@usda.gov;

jiangz@wsu.edu;

wul12@psu.edu;

jreecy@iastate.edu;

jekoltes@iastate.edu;

pross@ucdavis.edu;

hzhou@ucdavis.edu

Abstract

Deciphering the regulatory grammar of the genome is essential to understand the genetic and
molecular architecture of complex traits, as most trait-associated variants lie in non-coding
regions. Yet, functional annotation of the bovine genome remains limited, hindering our ability to
unravel the mechanisms underpinning complex traits of economic and ecological importance in
cattle. Here, we present a comprehensive epigenetic atlas comprising 1,147 genome-wide
epigenetic profiles, including chromatin accessibility, six histone modifications, CTCF
transcription factor binding sites, DNA methylation, chromatin conformation, and transcriptomes
across 53 adult tissues, 6 fetal tissues, and 7 primary cell types. This atlas covers 99.8% of the
bovine genome with at least one epigenomic feature, and annotates 45% of the genome as
putative regulatory elements exhibiting tissue- or cell-specific regulatory activity. Leveraging
sequence-to-function deep learning models, we discovered 301 sequence motifs and predicted
the functional impact of genetic variants through *in silico* mutagenesis, thereby facilitating the
decoding of the regulatory syntax of the cattle genome and fine-mapping of GWAS loci for 22
complex traits. Cross-species analysis further revealed evolutionarily conserved features of
regulatory architecture and provided evolutionary insights into complex traits and diseases in
humans. Together, this atlas offers a foundational resource for advancing cattle functional
genomics, sustainable breeding, and studies of regulatory evolution.
96

Introduction

Cattle are a key agricultural species, providing meat, milk, and leather, while also serving as a valuable model for studying embryo development, epigenetics and imprinting, immunology, and the genetics of complex traits[1,2]. Advances in high-throughput sequencing and genotyping technologies have substantially enhanced our understanding of the genetic architecture of economically important traits in cattle and accelerated the pace of genetic improvement[3]. For instance, incorporation of single nucleotide polymorphisms (SNPs) into dairy breeding programs has doubled the rate of genetic gain for key production traits[4]. As of the latest update, the CattleQTLdb[5] catalogs approximately 193,453 quantitative trait loci (QTL), the majority of which reside in non-coding genomic regions[6]. This strongly suggests that many QTL act through regulatory mechanisms, modulating transcriptional activity of nearby or distal gene regulatory elements[7]. Comprehensive annotation of the non-coding bovine genome is therefore essential to uncover mechanistic insights into GWAS signals and pinpoint causal variants. Such insights may ultimately enable precision breeding strategies and the development of "ultimate genotypes"[8] through genome editing[9] and synthetic biology[10], while further enhancing its utility as a biomedical model.

Initial efforts to characterize bovine regulatory landscape have provided important foundations. For example, Kern *et al.*[11] performed genome-wide annotation of regulatory elements by analyzing chromatin immunoprecipitation followed by sequencing (ChIP-seq) for four histone modifications (H3K4me3, H3K27ac, H3K4me1, H3K27me3) and the DNA-binding protein CTCF, as well as Assay for Transposase-Accessible Chromatin using sequencing (ATAC-seq), across eight core tissues (cortex, cerebellum, hypothalamus, liver, lung, spleen, muscle, adipose). Subsequent studies extended these annotations to rumen tissues [12,13] and rumen epithelial primary cells[12], and incorporated additional assays such as Cap Analysis Gene Expression (CAGE) sequencing[14], as well as broader genomic feature mapping[15]. Although these studies parallel large-scale efforts like ENCODE[16] and Roadmap Epigenomics[17] in humans towards annotating the non-coding genome, the bovine non-coding genome remains incompletely annotated. A systematic, large-scale effort spanning diverse tissues, cell types, and developmental stages (hereafter referred to as "tissue contexts") is essential to fully characterize context-specific regulatory mechanisms.

Complementary transcriptomic resources, including the Cattle Genotype-Tissue Expression (CattleGTEx)[18,19] and large-scale allele-specific (AS) analysis[20], have identified millions of regulatory variants influencing gene expression and splicing. Yet, these resources have not yet advanced to systematic identification of causal regulatory variants underlying GWAS associations of complex traits in cattle. Meanwhile, the emergence of sequence-to-function (S2F) deep learning models offers powerful approaches to decode the regulatory syntax of the genome and to prioritize functional variants. Tools like SpliceAI[21] and Pangolin[22] have demonstrated utility in predicting splicing effects in cattle[23], but broader application of S2F models for interpreting regulatory elements and context-specific variants remains largely unexplored.

In this study, we constructed a comprehensive bovine epigenomic atlas by integrating 540 newly generated datasets from 29 previously unprofiled tissues and assays with public available data[12,13,15,24,25]. In total, our resource comprises 1,147 genome-wide epigenomic profiles including 158 RNA-seq, 204 ATAC-seq, 91 whole genome bisulfite sequencing (WGBS), 682 ChIP-seq (six histone marks and CTCF), and 12 chromatin conformation (Hi-C) datasets. We used this data atlas to annotate chromatin states, link enhancers to their target genes, decode

regulatory syntax via deep learning approaches, fine-map GWAS loci, and perform multi-tissue
cross-species comparisons (**Fig. 1a**). The resulting resource provides a powerful foundation for
mechanistic dissection of complex traits, prioritization of functional variants, and comparative
regulatory biology.

Results

Near-complete bovine epigenomic landscape

We have compiled a comprehensive atlas of 1,147 genome-wide epigenomic maps through
uniform processing and quality control (**Fig. 1a-b**, **Supplementary Figures S1-S3**, **Table S1-S4**,
see **Methods**). These include 158 RNA-seq maps to quantify gene expression (**Supplementary**
Figure S4, Table S1), 204 ATAC-seq datasets to profile open chromatin regions (OCRs)
(**Supplementary Figure S5-S6**, **Table S2**), 91 WGBS datasets to assess DNA methylation
(**Supplementary Figure S7-S8**, **Table S3**), and 682 ChIP-seq datasets capturing histone
modifications and transcription factor (TF) binding (**Table S4**). The ChIP-seq datasets
encompass H3K4me1 (n = 131), marking enhancer regions; H3K4me3 (n = 132), associated with
promoter activity; H3K27ac (n = 125), indicative of active enhancer and promoter regions;
H3K27me3 (n = 123), marking Polycomb-mediated repression; H3K9me3 (n = 20), linked to
heterochromatin; H3K36me3 (n = 27), associated with actively transcribed regions; and CTCF (n
= 124), marking insulator elements[26] (**Supplementary Figure S9**). Among these, 540 ChIP-seq
datasets (47.5%) from 29 tissues were newly generated in this study, and 75 datasets were
computationally imputed using the ChromImpute pipeline[27], including H3K4me1 (n = 11),
H3K4me3 (n = 10), H3K27ac (n = 12), H3K27me3 (n = 22), and CTCF (n = 20), with genome-
wide Pearson correlation ranging from 0.56 (H3K27me3) to 0.91 (H3K4me3), comparable to
those observed in human datasets[27] (**Extended data Fig. 1, Supplementary Figure S10-S11**).
Peak calling from experimental ChIP-seq datasets yielded 895,108 H3K4me1 (52.7% of the
autosomal genome), 737,720 H3K4me3 (18.2%), 802,966 H3K27ac (26.9%), 1,038,083
H3K27me3 (47.8%), 442,773 H3K9me3 (7.2%), 567,480 H3K36me3 (18.4%), 831,902 CTCF
(19.8%) non-overlapping peaks. ATAC-seq analysis identified 953,179 non-overlapping open
chromatin regions (OCRs) covering 21% of the genome, while WGBS profiling identified
approximately 1.8 million methylation blocks (i.e. regions with ≥4 continuous CpGs), covering
73.4% of the genome. Additionally, we uniformly processed 12 public Hi-C datasets from 3
tissue contexts, yielding on average 4,979 topologically associating domains (TADs) per sample
at 10 kb resolution (**Table S5**). Altogether, these datasets cover 99.8% of the bovine genome with
at least one epigenomic feature (71% when excluding DNA methylation and TADs), providing a
comprehensive resource for exploring the regulatory landscape of the cattle genome.

Nearly half of the bovine genome annotated as regulatory elements

Leveraging the atlas-level maps above, we defined a 10-state "core" chromatin model across 72
complete bovine epigenomes representing 33 tissue contexts (**Fig. 1c**) (**Fig. 2a**, **Supplementary**
Figure S12). Here, a "complete epigenome" is defined as a biosample with all four core histone
marks (i.e., H3K4me1, H3K4me3, H3K27ac, and H3K27me3). These states included three
promoter-like states (PLS): TssA (active transcription start site [TSS], covering 2.2% of the
genome across tissue context), TssFlnk (flanking TSS, 3%), and TssWk (weak flanking TSS,

3.5%); three enhancer-like states (ELS): EnhA (active enhancer, 7.6%), EnhWk (weak enhancer,
11.7%), and EnhPos (poised enhancer, 12.5%); two bivalent states: BivTss (bivalent TSS, 0.98%)
and BivEnh (bivalent enhancer, 0.68%); as well as ReprPC (repressed Polycomb, 4.6%) and
Quies (quiescent/low signal) (**Fig. 2a**). These chromatin states, as expected[26,28,29], exhibited
distinct patterns of gene expression (**Extended data Fig. 2a**), chromatin accessibility (**Extended**
data Fig. 1a), DNA methylation (**Extended data Fig. 2b**), evolutionary conservation (**Fig. 2a,**
Extended data Fig. 2d, Supplementary Figure S13), and transposable elements enrichment
(**Supplementary Figure S14-15**). Of note, we found ELS from fetal epigenomes were
significantly enriched for CR1 and L2 subfamilies (**Supplementary Figure S15**), and a high
number of ELS overlapped with LINE (e.g. L1 and L2) and SINE subfamilies (**Supplementary**
Figure S16).

Compared to the previous annotation of eight core tissues by Kern et al[11], our atlas incorporates
25 additional tissues (16 adult, 5 fetal and 4 primary cell types), resulting in an additional 106
Mb of annotated sequence (19% increase). For example, we annotated 146,445 non-overlapping
EnhA, of which 66% were novel[11] (**Fig. 2b**). Across 65 complete bovine epigenomes with two
additional marks (OCRs, CTCF), we further constructed a 15-state "expanded" model. This
model introduced additional states such as TssFlnD (downstream flanking TSS), EnhAHet
(active enhancer lacking ATAC signal), EnhCtcf (enhancer with CTCF), CtcfIsl (CTCF island),
AtacIsl (ATAC island), and BivTssCtcf (bivalent TSS with CTCF) (**Extended data Fig. 3**).

To link enhancers to their target genes, moreover, we employed the Activity-by-Contact (ABC)
approach[30], and successfully linked 25,567 unique enhancers to 26,100 genes, resulting in a total
of 182,596 enhancer-gene pairs (**Fig. 2c-e, Supplementary Figure S17**). On average, each
enhancer was associated with 7.1 genes (**Fig. 2c**), and each gene with 7 enhancers (**Fig. 2d**), the
mean enhancer–gene distance was 168 kb (**Fig. 2e**). Furthermore, using ROSS software[31], we
predicted an average of 1,242 super-enhancers across 32 tissues (covering 18.6% of the genome),
ranging from 6 in liver to 3,128 in cerebral cortex (**Supplementary Figure S18-19**). Altogether,
45% of the bovine genome (excluding Quies) was annotated as regulatory elements, representing
the most comprehensive regulatory annotation to date in cattle.

Dynamic epigenomic signatures across tissues and cell types

To investigate context-specific regulation of gene expression, we systematically analyzed
chromatin state dynamics across tissues, developmental stages, and primary cell types. Tissue-
specific states were first identified (**Supplementary Figure S20** and **Table S6**). Focusing on
active enhancer (EnhA), we observed the highest number of tissue-specific EnhA elements in
muscle tissues, including muscle, heart, fetal heart, fetal muscle, and myoblasts (**Table S6**). In
contrast, digestive tissues such as omasum, reticulum, and duodenum (excluding rumen) showed
relatively few unique EnhA elements (**Fig. 2f**). These tissue-specific EnhA elements were
strongly enriched for biologically relevant functional terms (**Supplementary Figure S21, Table**
S7) and TF motifs[32] (**Fig. 2g, Table S8**). For instance, MYOG, MYF5, and MYOD motifs were
enriched in muscle tissues, NEUROG2, NEUROD1, and NF1 in central nervous system (CNS)
tissues, JUN, FOS and MYB73 in myoblasts (**Fig. 2g**), consistent with their established roles in
cell proliferation and lineage-specific differentiation[33,34]. Comparisons between adult, fetal, and
primary cells states, revealed additional layers of regulatory dynamics (**Supplementary Figure**
S22-24, Table S9-S10). As illustrated in **Fig. 2h**, we identified 27,255 adult-specific (30%),

13,425 fetal-specific (15%), and 29,946 myoblast-specific (34%) EnhA elements. A
representative example involves the myogenic regulators *MYF5* and *MYF6*. While *MYF6* was
highly expressed in adult and fetal muscle, but not in myoblasts, *MYF5* shows high expressions
in fetal muscle and myoblasts, but not in adult muscle (**Fig. 2i, Supplementary Figure S24**).
Chromatin accessibility data mirrored these expression patterns, with myoblasts- and fetal
muscle-specific enhancers flanking the *MYF5* locus (**Fig. 2i**). These findings align with the
established role of *MYF5* in early myogenic differentiation and *MYF6* in terminal myogenic
differentiation[35], mediated by a combinatorial network of enhancer elements during
development[36].

We next explored DNA methylation dynamics using 91 newly generated WGBS datasets
(**Supplementary Figure S7-S8**), identifying 207,436 differentially methylated blocks (DMBs)
(see **Methods**, **Table S11-S12**) across approximately 1.8 million tested regions (**Table S11**).
Fetal liver exhibited the highest number of DMBs (196,231). To construct a bovine tissue-
specific hypomethylated atlas, we selected the top 50 differentially unmethylated regions per
tissue (for tissues with >50 DMBs), resulting in a total of 1,120 DMBs (**Extended Data Fig. 4**).
These DMBs exhibited low methylation level (average 18%) and strong co-enrichment with
chromatin accessibility and histone modifications, exemplified by the cerebellum (**Extended
Data Fig. 5a**). Motif enrichment analysis using HOMER[32] further revealed tissue-relevant TFs
(**Table S13**, **Extended Data Fig. 5b**), such as *NEUROD1* in cerebellum, *MYB81* in muscle, and
MYF5 in myoblast (**Extended Data Fig. 5b**). Notably, *MYF5* encodes a critical regulator of
skeletal myogenesis[37]. The tissue-specific patterns for chromatin accessibility are shown in
Supplementary Fig. 25, and for gene expression in **Supplementary Fig. 26.** Finally, integration
of tissue-specific OCRs with GWAS data revealed significant enrichment of complex trait loci in
context-matched regulatory landscapes (**Extended Data Fig. 6**), underscoring the functional
relevance of these dynamic epigenomic signatures.

Deep learning reveals regulatory sequence syntax across tissue contexts

Gene expression is initiated and regulated through the binding of TF to specific DNA sequence,
known as TF binding site (TFBS)[38]. Recent advances in S2F deep learning models have
demonstrated strong performance in decoding regulatory syntax of DNA sequences and
predicting TFBS[7,39]. To explore the sequence-level grammar of regulatory elements in cattle, we
trained ChromBPNet[40], a convolutional neural network (CNN), to model the relationship
between genomic sequence and chromatin accessibility across 55 tissue contexts. For each tissue,
model performance, evaluated as the Pearson correlation between predicted and observed log-
transformed ATAC-seq read counts on held-out peaks (**Methods**), achieved a median of 0.76
(**Extended Data Fig. 7a, Table S14**), comparable to reports in human datasets[40,41]. Importantly,
Pearson correlations were stable across distances to the nearest transcription start site (TSS)
(**Extended Data Fig. 7b**). To assess generalizability, we applied the models to independent
ATAC-seq data from Yuan et al.[24] across 19 overlapping tissues. For example, in abomasum, our
model achieved a Pearson correlation of 0.84 on our datasets (**Extended Data Fig. 7c**) and 0.75
on Yuan et al.'s dataset[24] (**Extended Data Fig. 7d**), with minor differences potentially reflecting
breed or age effects[24]. A representative locus at *DGAT1* (chromosome 14: 603,067-604,765)
across tissues in both datasets, achieved a Pearson correlation of 0.69, with bone marrow
showing the highest chromatin accessibility (**Extended Data Fig. 7e**). Additionally, the model
could also predict tissue-specific accessibility peaks, as demonstrated in **Extended Data Fig. 7f**.

We next used high-performing tissue models (Pearson's $r \geq 0.72$, corresponding to the lower
quantile of model performances across tissues) for *de novo* motif discovery (44 tissue models
retained, **Extended Data Fig. 7a**). Using DeepLIFT [42] to score nucleotide contribution and TF-
MoDISco[43] to cluster predictive subsequences, we identified a non-redundant set of 301 *de novo*
motifs across 44 selected tissues (average: 16 per tissue, range: 9–23; **Fig. 3a**). The canonical
CTCF motif was consistently recovered in all tissue contexts, with contribution weight matrices
(CWM) strongly resembling the JASPAR positional weight matrix (PWM) (**Fig. 3b**), and
closely matching motifs identified in humans[41]. We also recovered some well-known TF motifs
(e.g., NEUROD1 in cerebellum and MEF2B in heart) (**Fig. 3c**) as well as motifs with limited
similarity to entries in the JASPAR database (*q*-value < 0.05), suggesting under-characterized or
potentially novel TFBSs (**Fig. 3d**). Scanning across all OCRs revealed the motif for KLF15, a
zinc-finger transcription factor with key roles in metabolic processes[44], as the most frequently
occurring one across tissues (**Fig. 3e**). Collectively, our findings demonstrate that ChromBPNet
generalizes well to cattle, enabling accurate prediction of chromatin accessibility and discovery
of conserved as well as novel sequence motifs. This supports its utility in decoding *cis*-regulatory
logic and prioritizing functional variants in livestock.

An atlas of allele-specific regulatory variations

To assess the impact of genetic variants on regulatory activity, we performed allele-specific (AS)
analyses across about 10 million SNPs identified from two individuals (see **Methods**). Out of
3,568,140 heterozygous SNPs overlapping epigenomic peaks, 91,125 (2.55%) exhibited
significant allelic imbalance (adjusted $P < 0.05$, **Fig. 4a, Table S15**). AS SNPs were enriched in
gene-dense and regulatory regions, showing positive correlation with gene density (Pearson's $r =$
0.23, $P = 3.92 \times 10^{-32}$), ChIP-seq peak density (Pearson's $r = 0.15$, $P = 8.7 \times 10^{-15}$) and OCRs
(Pearson's $r = 0.18$, $P = 3.04 \times 10^{-20}$) (**Fig. 4a**). The majority (70%) of AS SNPs were located in
intronic or intergenic regions (**Fig. 4b**), with significant enrichment within 5 kb upstream of TSS
(**Fig. 4c**). AS effects on gene expression correlated strongly with those affecting active
epigenomic marks (**Fig. 4d**), highlighting coordinated regulatory roles. Although effect sizes of
RNA-seq AS SNPs showed limited concordance with eQTLs from the multi-breed cattle GTEx
dataset[18,45] (**Supplementary Figure S27a**), they were highly and significantly correlated with
AS SNPs previously reported by Prowse-Wilkins et al.[46] (**Fig. 4e**). Furthermore, effect sizes of
epigenomic AS SNPs identified in our analysis showed high concordance with the Prowse-
Wilkins dataset[46] (**Fig. 4f**).

To evaluate their functional relevance to complex traits, we intersected AS SNPs with trait-
associated SNPs curated in the CattleQTLdb[5], identifying 439 putative variants (**Supplementary**
Figure S27b, Table S16). These AS SNPs exhibited strong assay- and tissue-specific
epigenomic effects (**Fig. 4g**). For example, SNP 3:15345227 in the *PKLR* promoter, a gene
associated with milk traits[47] and highly expressed in brain-related tissues, exhibits a strong
H3K27ac AS signal in jejunum ($\log_2 FC = 3.42$), coinciding with a tissue-specific BivTssCtcf
state (bivalent transcription start site with CTCF binding) (**Fig. 4h-i**). Another variant, SNP
29:42051788 in the *RTN3* promoter, showed the largest AS fold change in muscle, affecting
H3K4me3 signal (**Supplementary Figure S27c-d**). *RTN3* has been associated with tenderness
score[48] and lipid metabolism[49], suggesting potential causal functionality.

Together, these findings establish an atlas of allele-specific regulatory variants in cattle, linking genetic variation to tissue-specific regulatory effects and providing candidate functional variants underlying complex traits.

In silico prioritization of regulatory variants

S2F modeling enables quantitative prediction of the regulatory impact of genetic variants through *in silico* mutagenesis[39]. To systematically prioritize functional variants across tissue contexts, we employed deltaSVM[50], which balances context coverage (e.g. tissue and assay types) with computational efficiency (**Extended Data Fig. 8a**). Models were trained using epigenomic peaks as positives and GC-matched controls as negatives across 206 assay–tissue combinations (55 OCRs, 29 CTCF, 32 H3K27ac, 27 H3K27me3, 30 H3K4me1, 33 H3K4me3). Model performance was high, with average the receiver operating characteristic curves (AUCs) of 0.82–0.95 across assays (**Extended Data Fig. 8b**). Robustness was confirmed using OCR datasets from dairy cattle[24] with a median AUC of 0.95.

We then scored 28 million SNPs derived from 3,530 whole-genome sequences across diverse cattle breeds. We observed that deltaSVM scores were positively correlated with effect sizes of AS SNPs across epigenomic assays (**Fig. 5a, Supplementary Figure 28**), validating their use as proxies for regulatory impact of allelic changes. Following a strategy used in humans[51], we then defined a "regulatory magnitude" for each SNP as its maximum deltaSVM score across all contexts. This score was weakly but significantly correlated with both context specificity (i.e., the number of contexts in which the SNP overlapped with regulatory peaks) and genomic proximity to TSS (i.e., the distance of a SNP to the nearest gene TSS) (**Fig. 5b-c**), as well as with conservation metrics (phyloP and GERP), but not with the FAETH score[52]. We further categorized SNPs by regulatory context overlap: 1) Null: no overlap; 2) Specific: overlapping peaks in a single context; 3) Multiple: overlapping peaks in more than one but fewer than 80% of contexts; and 4) Shared: overlapping peaks in more than 80% of contexts. Stratification of SNPs by regulatory overlap revealed that tissue-shared variants were enriched near TSS and associated with promoter-like states, whereas tissue-specific variants overlapped primarily with enhancers (**Fig. 5d-e**). Moreover, both eQTL and trait-associated QTLs were significantly enriched for variants with high regulatory magnitude scores (**Supplementary Figure 28c**).

To evaluate their utility for fine mapping, we incorporated regulatory magnitude–based priors into Bayesian analyses of 22 complex traits (**Table S17**). This approach significantly reduced credible set sizes compared to unweighted analyses (**Fig. 5f**). For protein percentage, for example, regulatory weighting identified an additional putative causal SNP (6:84709163) (**Fig. 5g**). This variant is located in an intronic of *MGC152010*, which is highly expression in liver (**Supplementary Figure 28d**, **Fig. 5g, Table S17**), overlaps a liver-specific enhancer, and is predicted to disrupt high-mobility group factor binding (**Fig. 5g**). These results demonstrate that incorporating predicted regulatory effects into GWAS fine-mapping enhance causal-variant resolution, providing a framework for functional prioritization in livestock.

Comparative analysis of bovine and human epigenomes

To explore the evolutionary conservation of *cis*-regulatory elements (CREs), we compared chromatin states between cattle and human across 22 matched tissues. For humans, we analyzed

1,089 Roadmap Epigenomics datasets[17] spanning six regulatory marks (OCR, H3K27ac,
H3K27me3, H3K4me1, H3K4me3, and CTCF). Human chromatin states were re-annotated
using our pipeline to ensure methodological consistency, producing comparable emission
profiles and state numbers in both species (**Fig. 6a**). We classified human CREs into four
conservation categories: (1) sfCREs: conserved in both sequence and function; (2) sdCREs:
conserved in sequence but divergent in function; (3) soCREs: conserved in sequence only; and (4)
ssCREs: human-specific elements (**Fig. 6a**). Approximately 30% of human CREs were sdCREs,
indicating dynamic functional divergence despite sequence conservation. PLS were more often
functionally conserved (sfCREs) than ELS (**Fig. 6b**). Among active enhancers (EnhA),
sequence-conserved types (sfCREs, sdCREs, and soCREs) exhibited significantly higher phyloP
scores than ssCREs (q-value $< 10^{-300}$), with sfCREs being the most evolutionarily constrained
(**Fig. 6c**). Notably, ssCREs were more proximal to TSS, whereas sequence-conserved elements,
particularly sfCREs, were enriched at distal regions (**Fig. 6d**).

To assess trait relevance, we performed heritability partitioning using Linkage Disequilibrium
Score Regression (LDSC)[53] across 25 human GWAS traits (**Table S18**). The analysis identified
46 significant tissue-trait associations (FDR < 0.05), including enrichment of heart failure
heritability in myoblasts and fetal heart (**Fig. 6e**). Remarkably, sfCREs and ssCREs contributed
disproportionately to these enrichments, accounting for 41.3% and 37.0% of all significant
tissue-trait pairs, respectively (**Fig. 6f**). These findings highlight that both evolutionarily
conserved and lineage-specific enhancers play prominent roles in shaping the genetic
architecture of complex traits.

Discussion

Here we present a comprehensive bovine epigenomic atlas that integrates 1,147 datasets across a
broad range of tissues, developmental stages, and primary cell types. By combining RNA-seq,
ATAC-seq, WGBS, ChIP-seq, and Hi-C data, we mapped chromatin states and regulatory
elements that collectively cover approximately 45% of the bovine genome, representing a ~20%
increase over previous annotations[11]. This expanded regulatory landscape displayed extensive
context specificity, supported by enriched tissue-relevant TFBS motifs such as *NEUROD1* in
neural tissues and *MYF5* in myoblasts. Leveraging this atlas, we applied S2F models,
including ChromBPNet[40] and deltaSVM[50], to decode regulatory grammar and predict variant
effects. These models captured base-resolution accessibility patterns, recovered canonical (e.g.
CTCF) and novel sequence motifs, and assigned quantitative regulatory scores to millions of
variants. Integration with AS analyses and GWAS fine-mapping, highlighted putative causal
variants. Cross-species comparisons further revealed that both evolutionarily conserved and
lineage-specific enhancers contribute significantly to genetic architecture of complex traits:
conserved elements (sfCREs) were distal and under strong sequence constraint, whereas species-
specific elements (ssCREs), though often proximal to TSS, explained a large proportion of
heritability for agricultural and biomedical traits. Together, these findings establish cattle as a
powerful model for dissecting regulatory biology and complex traits, with relevance extending
beyond agriculture to comparative and evolutionary genomics.

Despite these advances, several limitations remain. First, reliance on bulk tissue datasets limits
resolution, masking cell-type–specific activity. Incorporating single-cell epigenomic and
transcriptomic profiles will be critical to capture cellular heterogeneity and dynamic transitions

during development or disease progression[54]. Second, the small number of individuals profiled restricts detection of population-level regulatory variation. Population-scale efforts will enable discovery of molecular QTLs, construction of gene regulatory networks, and more precise genotype–phenotype links[10,18,45,55,56]. Third, deeper sequencing and enhanced data quality will improve sensitivity for detecting weak or transient regulatory elements, annotation and enable assessment of complex variants, such as structural variants[57], which are currently underexplored in functional contexts. Fourth, while our S2F models[58] perform robustly, further refinements—especially for those leveraging multimodal data and single-cell resolution—are needed to improve prediction of subtle or combinatorial variant effects. Fifth, comparative epigenomics across additional species will provide deeper insights into regulatory evolution and shared principles of trait heritability. Finally, functional validation remains essential. Massively parallel reporter assays[59], STARR-seq[60], and CRISPR-based perturbation strategies will be key for functionally confirming enhancer activity and directly testing variant effects in their native chromatin context.

In summary, this study provides the first atlas-scale annotation of regulatory elements in cattle, offering a foundational resource for mechanistic interpretation of non-coding variation. By integrating functional genomics with predictive modeling and comparative analysis, our work supports applications in precision breeding, evolutionary biology, and functional genomics. As future efforts expand to single-cell resolution and population scale, this resource lays essential groundwork for a comprehensive understanding of gene regulation and complex trait variation in mammals.

Acknowledgments:

We thank all of the people that contributed to the breeding and collection of tissues used in this study. We also thank all the researchers who have contributed to the publicly available data used in this research.

Funding:

This study was supported by Agriculture and Food Research Initiative Competitive Grant no. 2018-67015-27500 (H.Z., P.R. etc.) and sample collection was supported by no. 2015-67015-22940 (H.Z. and P.R.) from the USDA National Institute of Food and Agriculture. D.G. was supported by Agriculture and Food Research Initiative Competitive grants nos. 2018-67015-27500 (H.Z.) from the USDA National Institute of Food and Agriculture. J.B. received scholarships financed by Foundation to Support the Research of São Paulo State (FAPESP) (Grants Numbers: 2020/01369-1 and 2022/12631-4). H.Z. acknowledges funding from the USDA National Institute of Food and Agriculture, Multistate Research Project NRSP8 and NC1170 and the California Agricultural Experimental Station.

Author contribution statement:

H.Z., L.F. and D.G. conceived the manuscript. D.G. led all the data analysis. J.B. and D.G. analyzed RNA-seq and linked enhancer to target genes, J.B., D.G., C.A.G, G.W. and Y.X. processed WGBS data, D.G. and W.Y. annotated super-enhancers. W.Y., J.B., D.G., B.P., and

C.P.W. performed allele-specific analysis, C.P.W., D.G. analyzed chromatin accessibility data,
D.G., J.W. and J.J. conducted complex trait integration analysis, S.C., X.X., Y.Wang, Y. Z. and
D.G. led new data generation and quality control, H.L., H.Z. and L.F. provided eQTL resources
used in this study, Y.U. and J.D. provide GWAS summary statistics from beef cattle, L.M., C.M.,
J.W. and J.J. provided GWAS summary statistics from dairy cattle. G.W. and Y.X. analyzed
ATAC-Seq. P.L. and H.J. generated ChIP-seq data and primary cells. G.R., C.G., T.P.L.S., H.J.,
Z.J., W.L., J.R., J.F.M., J.D., P.R., J.E.K., H.Z. contributed to funding acquisition. D.G. prepared
the initial manuscript draft with input from L.F. and H.Z. D.G., L.F. and H.Z., and revised and
finalized the manuscript. All authors read, edited and approved the final manuscript.

Competing interests:

457 Y.T.U., V.M.P.R. and J.D. are employed by STgenetics. P.R.S acquired the funding for this
458 research while affiliated to the University of California, Davis, and is currently employed by
459 STgenetics. S.C. and G.R. is from Zoetis Inc.

Data and materials availability:

We have deposited newly generated RNA-Seq, ChIP-Seq, ATAC-seq WGBS in FAANG portal
(https://data.faang.org/), NCBI SRA database (https://www.ncbi.nlm.nih.gov/sra/), and European
Nucleotide Archive. Accession IDs will be available upon the acceptance of the manuscript.
Accession IDs for public data includes: PRJEB41939, PRJNA672996, PRJNA531208, E-
MTAB-11825 and E-MTAB-11826 (ENA). Metadata details are available in Supplementary
Tables   S1-S4.   All   processed   data   are   available   at   Zenodo   database
(https://zenodo.org/10.5281/zenodo.12216791, available upon the acceptance). Cattle traits
QTLs retrieved from CattleQTLdb (https://www.animalgenome.org/cgi-bin/QTLdb/BT/index),
and eQTL data retrieved from https://cattlecell.kiz.ac.cn/. Other data sets used in this study,
published previously, are described in the Methods. All the computational scripts and codes are
available at the GitHub website, including https://github.com/guandailu/BovineFAANG,
https://github.com/guandailu/ChIPQC, and https://github.com/guandailu/SnakeHiC.

Online Methods

Sample collection

Samples used in this study were collected by the Functional Annotation of Animal Genomes
(FAANG) community. The sampling procedure has been previously described in detail by Tixier-
Boichard et al[61]. Briefly, tissue samples for beef cattle were collected from two male and two
female Hereford cattle within 1–2 hours post-euthanasia, flash-frozen in liquid nitrogen, and
stored at -80°C until further processing. The same protocol was applied to collect mammary
gland tissues from Holstein cows at different physiological stages. All animal experiments were
conducted in accordance with Protocol for Animal Care and Use No. 18464, approved by the
Institutional Animal Care and Use Committee at the University of California, Davis.

Library preparation and sequencing

For RNA-seq experiments, frozen tissue samples were pulverized in liquid nitrogen using a
mortar and pestle. Total RNA was then extracted with TRIzol reagent (Thermo Fisher Scientific,
#15596026) and treated with DNase I (Thermo Fisher Scientific, #EN0521). RNA integrity was
assessed using an Agilent Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA), and
samples with RNA integrity number (RIN) > 8 were used for library preparation, followed by
sequencing on an Illumina Hiseq 4000 platform (Illumina, San Diego, CA, USA). For ChIP-seq
(targeting H3K4me3, H3K4me1, H3K27ac, H3K27me3, and CTCF) experiments, samples were
processed as previously described[28,29]. ATAC-seq libraries were prepared using a modified
protocol through incorporating cryopreserved nuclei workflows, as described in [62,63]. Both
ATAC-seq and ChIP-seq libraries were sequenced on an Illumina HiSeq 4000 platform using 50-
bp paired-end and single-end reads, respectively. For whole-genome bisulfite sequencing
(WGBS), genomic DNA of tissue samples was extracted, fragmented, and sequencing libraries
were prepared using methylation sequencing adaptors. Libraries were then treated with sodium
bisulfite, which converted unmethylated cytosines to uracils while preserving methylated
cytosines. The bisulfite-converted libraries were then PCR-amplified and subjected to Illumina
paired-end 150 bp sequencing to assess genome-wide DNA methylation at single-base resolution.

Raw data processing and quality control

For ChIP-seq sequencing data, we processed them using a uniform pipeline available at
https://github.com/guandailu/ChIPQC, after gathering newly generated sequencing data in this
study and previously published datasets (details provided in **Table S1**). Briefly, sequencing
adapters were trimmed using Trim Galore (v0.6.7). Cleaned reads were aligned to the
ARS_UCD1.2 reference genome (Ensembl release v105) using BWA-MEM[64]. PCR duplicated
reads were removed prior to peak calling. Deduplicated BAM files were used to identify peaks
with MACS2[65] with the options of "--q 0.01" for narrow marks (i.e. H3K27ac, CTCF), and of "--
q 0.05 --broad" for broad marks (i.e. H3K4me1, H3K27me3, H3K36me3, and H3K9me3). Then,
we filtered out samples with the clean reads < 10 million, the number of peaks < 10,000 and
Jensen-Shannon Divergence (JSD) between two replicates < 0.05 for broad marks and < 0.1 for
narrow marks. This filtering step finally left 607 datasets that remained for subsequent analyses.
For ATAC-seq, we applied the same processing steps, including adaptor trimming, read
alignments, the removal of PCR duplicated reads, and peak calling, while the peaking calling
was applied with the specific options of "-f BEDPE --nomodel". The filtering criteria for ATAC-
seq included a FRiP score >0.3 and checked TSS enrichment manually. For RNA-seq, a similar
pipeline was applied although the read aligner employed STAR (v2.7.0)[66]. Gene expression
levels were quantified as Transcripts Per Million (TPM) using StringTie2[67], and raw read counts
were obtained using featureCounts (release 2.0.6)[68] based on the ARS_UCD1.2 gene annotation
(Ensembl release v105). For 12 public Hi-C datasets, we uniformly processed them using the
pipeline SnakeHiC (https://github.com/guandailu/SnakeHiC). For DNA methylation data
generated by WGBS, adaptors were trimmed using Trim Galore (v0.6.6) and we preprocessed
using the Bismark pipeline (Version 0.24.2). The resulting alignment files in BAM format were
subsequently converted into compressed formats, including "beta and "pat", using wgbs_tools
(version 0.2.2)[69,70]. Homogeneously methylated blocks (defined as contiguous CpG sites with
similar methylation levels) were therefore identified using the "*segment*" command, and blocks
with fewer than 4 CpG sites were excluded from downstream analyses. For each segmented

block, we computed the average methylation level across CpG sites based on the corresponding
beta formatted files.

Imputation of missing and low-quality epigenomic signals

To obtain complete epigenomes for chromatin state annotations, we employed the ChromImpute
pipeline (v1.0.5) developed by the Roadmap Epigenomics Consortium[17,26,27]. Briefly, we
generated signal tracks for experimental samples using the *bdgcmp* function of the MACS2
tool[65] , computing $-\log_{10}P$ as recommended by[26,27]. For brevity, we only considered signals for
autosomes 1-29. ChIP-seq signal tracks were converted to 25-bp resolution by averaging signal
intensity across each 25-bp bin. Imputation of missing or low-quality signals followed four main
steps: 1) computing global distance between datasets with the *ComputeGlobalDist* function; 2)
generating training features with the *GenerateTrainData* command; 3) training mark-specific and
sample-specific predictors with *Train* command; and 4) applying these predictors to generate
imputed signal tracks using *Apply*. To assess imputation accuracy, we implemented a cross-
validation-like procedure. Two experimental samples were randomly masked, and imputation
was performed on these held-out samples. The correlation between imputed and observed signals
was then assessed using the *Eval* function. This process was repeated 15 times to ensure that
each epigenetic mark was tested at least twice. In addition, imputed signals were manually
inspected to ensure quality. Additionally, we validated the robustness of the pipeline using ChIP-
seq datasets published by Prowse-Wilkins et al.[25], masking 15 samples using the same procedure.
Comparable high imputation accuracy was observed across these held-out datasets.

Predicting chromatin states and super-enhancers

We employed ChromHMM (v1.26)[71,72] to predict chromatin states for 72 bovine epigenomes,
each of which included the core set of four histone modifications: H3K4me1, H3K4me3,
H3K27ac, and H3K27me3. Briefly, we used the *LearnModel* function in ChromHMM to train
models with state numbers ranging from 2 to 15—the maximum number supported by the
software. To identify the optimal model, we compared emission probability across models
following the approach described by Gorkin et al.[73] The optimal number of states was
determined as the point at which the median correlation of emission probability across two or
more replicates plateaued. For the 72 bovine epigenomes, a 10-state model was ultimately
selected, and state labels were manually assigned based on an extensive literature review,
particularly as described in ref[74]. For 65 bovine complete epigenomes with two additional marks
(OCRs and CTCF), we ran a similar pipeline and learned 15 chromatin states. In order to
compare cattle data with that of humans, we also conducted the chromatin state annotation for
134 human epigenomes with the same pipeline.

We also annotated super-enhancers, a class of regulatory elements that differ from typical
enhancers in terms of their size, transcription factor density and content, transcriptional
activation potential, and sensitivity to perturbation[31,75]. Super-enhancer identification was
performed using the ROSE algorithm (version 0.1)[31] across 33 tissue contexts. For each tissue,
H3K27ac ChIP-seq profiles from two biological replicates were merged using SAMtools
(v1.19.2)[76]. As input, we used the genomic regions annotated as typical enhancers (i.e. chromatin

states E4, E5, and E6) to define candidate super-enhancer regions. These were then ranked based
on H3K27ac signal intensity to identify super-enhancers following the standard ROSE pipeline[31].

Linking enhancers to genes

To predict target genes of enhancer elements, we employed the Activity-by-Contact (ABC)
model[30]. We applied the ABC pipeline across 62 samples from 37 tissues that had matched
ATAC-seq and H3K27ac ChIP-seq data. The ABC score for a given enhancer-gene pair was
calculated as:

$$ABC\ score\ _{(E,G)} = \frac{A_E \times C_{E,G}}{\sum A_e \times C_{e,G}}$$

where E and G represent the enhancer element and the gene, respectively. A(E) represents the
activity of element E, and C(E, G) denotes the Hi-C contact frequency between element E and
the promoter of gene G. The denominator sums this product over all candidate elements within
±5 Mb of the gene. Activity was quantified using read counts from ATAC-seq and H3K27ac
ChIP-seq data. Candidate enhancer elements were defined as ~500 bp regions centered on
ATAC-seq peaks.

Context specificity analyses
For gene expression: We employed the Tspex[77] as described previously[55]. For a given tissue, we
computed the average gene expression across biological replicates. Tissue-specific expression
was then defined using a Z-score approach across all tissues, with genes considered tissue-
specific if Z-score >= 0.8. Adult and fetal tissues were separate in this analysis. Subsequently, we
performed enrichment analysis on the identified tissue-specific gene sets using the Enrichr
package (v3.2)[78] and the GO_Biological_Process_2023 database. The significant threshold was
set as adjusted *P*-value threshold of <= 0.05 with the False Discovery Rate (FDR) approach[79].
For chromatin states: We applied an intuitive comparison between the target tissue and the test to
identify tissue-specific chromatin states with BEDtools (version 2.31.0)[80]. That is, a state
observed in one tissue but not in others. Motif enrichment of tissue-specific chromatin states was
conducted using the Hypergeometric Optimization of Motif EnRichment (HOMER) tool (v5.1,
7-16-2024)[32].
For open chromatin regions: To identify tissue-specific OCRs, we first gathered all peaks across
valid samples to create the consensus set of peaks. Then we quantified raw reads counts of
consensus peaks across samples. Tissue-specific OCRs were identified by using DESeq2 tool (v
1.49.1)[81] when comparing a given tissue with the rest. The significant threshold was set as FDR-
adjusted $P < 0.05$ and fold change at $\log_2$ scale > 1.
For DNA methylation: Tissue-specific differentially methylated regions (DMRs) were identified
by comparing each focal tissue against all others using the *find_markers* function in the
wgbs_tools[69] package (v0.2.2). Statistical significance was assessed via two-sided t-tests, and
DMRs with $P > 0.05$ were excluded using the --pval 0.05 option.
Comparing adult and fetal samples: We identified differential chromatin states between adult and
fetal stages using an BEDtools (version 2.31.0). That is, a state observed in one stage but not in
another one.

Allele-specific identification

We identified genome-wide single nucleotide polymorphisms (SNPs) from two biological
replicates (M08 and M22) using the Genome Analysis Toolkit (GATK, v4.5.0.0) following the
"best practices" pipeline[82]. To minimize reference mapping bias and enable accurate detection of
allele-specific (AS) events, we applied the WASP (v0.16.3) pipeline[83] to all epigenomic datasets.
Read counts for each SNP were quantified using the *ASEReadCounter* module in GATK. SNPs
were retained for AS analysis if they (i) were heterozygous, (ii) overlapped epigenetic peaks
previously identified by MACS2[65], (iii) were located on autosomes, and (iv) had >6 total reads,
with each allele supported by at least 3 reads. For each retained SNP within a sample, a two-
sided binomial test was performed to assess allelic imbalance. *P*-values were then adjusted for
multiple testing using the Benjamini-Hochberg (BH) procedure[79], and SNPs with a false
discovery rate (FDR) < 5% were considered significant AS events.

Training ChromBPNet to learn sequence motifs

We employed ChromBPNet (v1.0)[40], a CNN-based deep learning framework, to learn sequence
regulatory syntax of chromatin accessibility in the bovine genome. Briefly, for each of the 55
tissue contexts, we merged BAM files from biological replicates to generate input data. The
analysis was restricted to ATAC-seq data, as ChromBPNet is specifically designed to model
chromatin accessibility and also for simplicity. OCR peaks, previously called by MACS2[65], were
merged by tissue using the Irreproducible Discovery Rate (IDR) framework (version 2.0.3)[84] and
filtered to exclude regions overlapping with highly repetitive sequences. Genome mappability for
the ARS-UCD1.2 reference genome was computed using GenMap (v1.3.0) [85]. ChromBPNet
models were trained using a chromosome-based data split: training set (chr2, chr4-5, chr7, chr9-
19, chr21-29), validation set (chr8, chr20), and testing set (chr1, chr3, chr6). As negative sets, we
generated GC-matched background regions using the "prep nonpeaks" function of the
ChromBPNet pipeline. Model training proceeded in two steps: (1) training a bias model to
account for Tn5 insertion bias, and (2) training a bias-factorized ChromBPNet model. Chromatin
accessibility in OCRs from the test set was then predicted using the *pred_bw* function, which
outputs both read counts and signal shape of profiles. Prediction accuracy was evaluated using
Pearson's correlation, Spearman's correlation, and JSD. In accordance with the original
publication[40], Pearson's correlation between observed and predicted read counts demonstrated
desirable evaluations and was therefore selected for subsequent analyses. To interpret model
predictions and identify sequence motifs underlying OCRs, nucleotide-level contribution scores
were computed and stored as BigWig files. These scores were used for *de novo* motif discovery
with TF-MoDISCo (v1.0.0)[43]. To construct a non-redundant motif catalog across tissue contexts,
we clustered predicted motif instances and merged clusters with pairwise similarity scores
greater than 0.1, yielding a final set of 301 non-redundant sequence motifs.

Training gkm-SVM classifier

To assess the regulatory impact of DNA sequence variants in the cattle genome, we adopted the
deltaSVM approach[50], which offers both high predictive accuracy and computational efficiency,
making it suitable for genome-wide variant scoring across diverse epigenetic marks and tissue
contexts. Specifically, we trained gapped k-mer Support Vector Machine (gkm-SVM)
classifiers[86,87] for each tissue-epigenetic mark combination. For each epigenetic mark, we first
merged MACS2[65] identified peaks from two biological replicates using the IDR framework

(v2.0.3) [84] and filtered out regions overlapping with highly repetitive sequences. The remaining
peaks were then aggregated across all tissues to define a positive set. To generate the negative set,
we sampled GC-matched genomic regions that did not overlap any peak regions. We used LS-
GKM (v0.1.1) [88], an optimized implementation of gkm-SVM with improved performance[87,88], to
train the classifiers. Model performance was evaluated using 10-fold cross-validation, with
parallelization enabled via the "-i" option.

To compute deltaSVM scores, we first generated all non-redundant 11-mers using the *nrkmers.py*
script. Then, for ~28 million SNPs identified from 3,530 whole-genome sequences (generated by
collaborators, unpublished), we synthesized 40-bp allelic oligonucleotides for both reference and
alternative alleles (centered on the SNP with ±19 bp flanking sequence). Variant impact scores
were computed using the deltasvm.pl script with the following command: perl deltasvm.pl
chr${chr}.ref.fa chr${chr}.alt.fa weights.txt deltaSVM.tsv. For each SNP, we defined the
maximum deltaSVM score across all tissue-mark combinatory contexts as its regulatory
magnitude, representing the predicted change in regulatory activity between alleles. To assess
whether CattleGTEx eQTLs[18,45] and traits QTLs[5] are enriched for high regulatory magnitude
scores, we conducted an enrichment analysis by fitting a linear model. For eQTLs, we compared
fine-mapped causal variants (posterior inclusion probability [PIP] > 0.8) to non-causal eQTL
SNPs (PIP < 0.01). For trait QTLs, we contrasted them with randomly selected SNPs matched
for minor allele frequency (MAF). In both analyses, we regressed out genomic confounders,
including distance to the nearest TSS, context specificity, and GERP conservation scores, as
these features were significantly correlated with regulatory magnitude.

GWAs fine-mapping analyses
Genome-wide significant association peaks for 22 dairy bull traits were identified from GWAS
summary statistics (Wang et al., in press) using a significance threshold of $P < 5 \times 10^{-8}$. For fine-
mapping analysis, candidate regions were defined as 2-Mb windows centered on the lead SNP of
each associated peak. Fine-mapping was performed using SuSiE-adj[89], an adaptation of SuSiE[90]
that improves fine-mapping accuracy in samples with related individuals by incorporating linear
mixed model (LMM)-derived summary statistics. We applied SuSiE-adj both with and without
scaled regulatory magnitude as prior weights across 22 traits. Conditional analysis was
conducted by including the lead SNP as a covariate and performing association analysis using
the SLEMM-GWA method[91].

Linkage disequilibrium score regression analysis

To investigate whether the conservation of regulatory elements defined in this study could
inform the genetic architecture of human complex traits, we collected GWAS summary statistics
for 25 traits (details in **Table S18**). We performed tissue-specific (equivalent to cell-type-
specific[92]) heritability enrichment analyses using the LDSC package (v1.0.0)[93]. GWAS summary
statistics were standardized using the *munge_sumstats.py* script to remove SNPs with MAF $\leq$
0.01, INFO score $\leq$ 0.9, genotype call rate $\leq$ 0.75, duplicate rsIDs, out-of-bounds *P*-values,
extreme chi-squared statistics, strand ambiguity, or discordance with SNPs used in LD score
calculations. For each category of regulatory elements, we identified tissue-specific segments by

contrasting the target tissue with others using BEDtools (v2.31.0)[80]. Annotations were generated
for LDSC analysis using the *make_annot.py* script, and stratified LD scores were then calculated
within a 500 kb window using SNPs from the 1000 Genomes Project Phase 3. We used default
SNP weights and reference panels obtained from https://zenodo.org/records/7768714. Tissue-
specific heritability enrichment analyses were then conducted using the "--h2-cts" option in
LDSC[92].
707

Figure 1. Overview of the study design and datasets. (a) Schematic overview of the analytical framework used in this study, including data processing, imputation of missing or low-quality histone marks, chromatin state annotation, enhancer-gene linking, tissue-specificity analysis, sequence grammar modelling using deep learning, fine-mapping GWAS loci of complex traits, and cross-species epigenomic comparison. (b) Summary of multi-omic datasets integrated in this study. (c) Illustration of tissue samples used for chromatin state annotation in this study. Icons and the cattle image were adapted from BioRender.com.

Figure 2. Chromatin state annotation and context-specific regulatory dynamics. (a) Definition and characterization of chromatin states, including emission probabilities, functional descriptions (annotations), abbreviations, enrichment near transcription start sites (TSS), overlap with genomic features, associated gene expression levels, and evolutionary constraint based on GERP scores[94] (Ensembl release v105). CDS: coding sequence. UTR: untranslated regions. (b) Genome coverage of strong enhancers (EnhA) compared to annotations from that of Kern et al.[11] (c)-(e) Enhancer-gene linking based on the Activity-by-Contact (ABC) model[30]: number of genes linked per enhancer (c), number of enhancers linked per gene (d), and genomic distance between enhancers and their predicted target genes (e). (f) Number of tissue-specific strong enhancers (EnhA) identified across tissues. (g) Motif enrichment analysis of tissue-specific EnhA elements. Representative enriched motifs and associated transcription factors (TFs) with known tissue relevance are shown at right. (h) Venn diagram depicting the overlap of strong enhancers (EnhA) identified in myoblasts (cell line), adult muscle and fetal muscle (bulk tissue). (i) Genome browser tracks displaying gene expression and epigenomic signals around the *MYH5* and *MYH6* loci (chr5:10262831-10302829 bp).

Figure 3. Sequence motif discovery across tissue contexts. (a) Number of seqlets (i.e., short informative subsequences) supporting each of the 301 unique motifs (y-axis) across 42 tissue contexts (x-axis). (b) Comparison between the CTCF motif identified by ChromBPNet[40] in this study (top) and the canonical CTCF motif from the JASPAR database (bottom). (c) Representative motifs identified in this study showing high similarity to known TF motifs curated in the JASPAR database. (d) Five motifs discovered in the rumen, a ruminant-specific organ, which show low similarity to any known motifs in JASPAR motifs (q-value > 0.05). (e) Proportion of motif instances identified across tissue contexts. The KLF15 motif was the most frequently detected. The unit of y-axis for motifs patterns is "bits", while the ChromBPNet derived motifs are represented using contribution weight matrix (CWM) score. The known non-redundant motifs from the JASPAR database were last accessed on February 11, 2022. TF: transcription factor.

Figure 4. Identification and characterization of allele-specific (AS) epigenomic variation. (a) Genome-wide distribution of AS SNPs across bovine autosomes. (b) Proportion of SNPs (x-axis) by variant categories (y-axis). The variant category was obtained using SnpEff[95]. (c) Genomic distribution of significant AS SNPs relative to gene transcription start sites (TSS). (d) Correlation of AS effect sizes ($\log_2$ scaled) in RNA-seq (x-axis) and matched epigenomic assays (y-axis). (e) Comparison of AS effect sizes ($\log_2$ scale) in RNA-seq between this study (x-axis) and Prowse-Wilkins et al.[46] (y-axis). (f) Comparison of AS effect sizes ($\log_2$ scale) in epigenomic assays between this study (x-axis) and Prowse-Wilkins et al.[46] (y-axis). *P*-values in (d–f) were computed using the asymptotic *t*-distribution approximation. FC: fold change. (g) Heatmap showing fold changes ($\log_2$ scale) for 439 AS SNPs overlapping known trait-associated QTLs curated in the CattleQTLdb[5]. (h) Gene expression profile of *PKLR* across tissues measured in

transcripts per million (TPM). (i) Genomic tracks showing an AS SNP in the promoter region of
PKLR, detected in jejunum H3K27ac data. This SNP overlaps a predicted ETV5:FIGLA binding
motif (identified using FIMO[96]).

Figure 5. Predicting regulatory effects of sequence variants. (a) Correlation between
deltaSVM scores (y-axis) and allelic fold change (log□-scaled, x-axis) across four assays:
ATAC-seq, CTCF, H3K27ac, and H3K4me3 (left to right, respectively). (b) Relationship
between regulatory magnitude (left y-axis: defined as the maximum deltaSVM score across all
assay-tissue contexts) and distance to the nearest transcription start site (TSS; right y-axis) as a
function of context specificity (i.e., number of assay–tissue contexts affected). (c) Mean distance
to TSS for genetic variants stratified by deciles of regulatory magnitude. (d) Density plot
showing the distribution of variant distances to TSS categorized by context specificity, including:
"null" (impacting 0 assay-tissue contexts), "specific" (impacting only 1 assay-tissue context),
"multiple" (impacting > 1 but <80% of assay-tissue contexts); and "shared" (impacting >80% of
assay-tissue contexts). (e) Proportion of SNPs residing within enhancer-like chromatin states
(EnhPos, EnhWk, EnhA) and promoter-like chromatin states (TssWk, TssFlnk, TssA), stratified
by "Specific" and "Shared" variant categories. (f) Regulatory magnitude weighted fine mapping
produces smaller median credible set sizes across 22 complex traits (P=4.43 × 10^{-8}, Wilcoxon
test). (g) Fine-mapping a cattle GWAS locus for protein percentage in milk using regulatory
magnitude as prior weights in SuSiE-adj[89,90]. The panels from top to down are: GWAS
significance plot (-log$_{10}P$); posterior inclusion probability (PIP) for fine-mapped variants without
prior weights; posterior inclusion probability (PIP) for fine-mapped variants with regulatory
magnitude prior weights; chromatin state annotations across 65 bovine epigenomes. The vertical
red line highlights the fine-mapped variant overlapping both chromatin states and transcription
factor binding sites (TFBS).

Figure 6. Cross-species comparison and conservation of regulatory elements. (a) Chromatin
state annotations comprising 15 states were independently predicted in cattle and human. The top
panel summarizes chromatin state labels, functional descriptions, abbreviations, and
corresponding emission probabilities for both species. The bottom panel categorizes cis-
regulatory elements (CREs) based on conservation: 1) sfCREs, sequence- and functionally
conserved elements; 2) sdCREs, sequence conserved but functionally divergent; 3) soCREs,
sequence-only conserved; and 4) ssCREs, human-specific elements. (b) Proportions of regulatory
elements stratified by conservation category. (c) phyloP conservation scores for strong enhancers
(EnhA), grouped by conservation category. Significance was assessed using a two-sided
Student's t-test with FDR adjustment. phyloP scores were obtained from Cactus 241-way
mammalian alignments provided by the Zoonomia Project[97]. (d) Genomic distance between
strong enhancers (EnhA) and the nearest transcription start site (TSS), stratified by conservation
category. Color legend is same as panel (b). (e) Tissue-specific heritability enrichment of EnhA
elements stratified by conservation category, represented as -log$_{10}P$ values. (f) Number of
significantly enriched tissue–trait pairs (x-axis) contributed by EnhA elements from each
conservation category (y-axis). Trait name abbreviations are provided at **Table S28.**

793
794

Reference:

1. Hubert, J.-N., Perret, M., Riquet, J. & Demars, J. Livestock species as emerging models for genomic imprinting. *Front. Cell Dev. Biol.* **12**, (2024).
2. Kemper, K. E. & Goddard, M. E. Understanding and predicting complex traits: knowledge from cattle. *Human Molecular Genetics* **21**, R45–R51 (2012).
3. Georges, M., Charlier, C. & Hayes, B. Harnessing genomic information for livestock improvement. *Nat Rev Genet* **20**, 135–156 (2019).
4. Wiggans, G. R., Cole, J. B., Hubbard, S. M. & Sonstegard, T. S. Genomic Selection in Dairy Cattle: The USDA Experience. *Annual Review of Animal Biosciences* **5**, 309–327 (2017).
5. Hu, Z.-L., Park, C. A. & Reecy, J. M. Bringing the Animal QTLdb and CorrDB into the future: meeting new challenges and providing updated services. *Nucleic Acids Research* **50**, D956–D961 (2022).
6. Koufariotis, L., Chen, Y.-P. P., Bolormaa, S. & Hayes, B. J. Regulatory and coding genome regions are enriched for trait associated variants in dairy and beef cattle. *BMC Genomics* **15**, 436 (2014).
7. Zhou, H. *et al.* Comparative Genomics and Epigenomics of Transcriptional Regulation. *Annual Review of Animal Biosciences* **13**, 73–98 (2025).
8. Hayes, B. J. *et al.* Potential approaches to create ultimate genotypes in crops and livestock. *Nat Genet* **56**, 2310–2317 (2024).
9. Tait-Burkard, C. *et al.* Livestock 2.0 – genome editing for fitter, healthier, and more productive farmed animals. *Genome Biology* **19**, 204 (2018).
10. Fang, L. *et al.* The Farm Animal Genotype–Tissue Expression (FarmGTEx) Project. *Nat Genet* 1–11 (2025) doi:10.1038/s41588-025-02121-5.
11. Kern, C. *et al.* Functional annotations of three domestic animal genomes provide vital resources for comparative and agricultural research. *Nat Commun* **12**, 1821 (2021).
12. Fang, L. *et al.* Functional annotation of the cattle genome through systematic discovery and characterization of chromatin states and butyrate-induced variations. *BMC Biology* **17**, 68 (2019).
13. Gao, Y. *et al.* Functional annotation of regulatory elements in cattle genome reveals the roles of extracellular interaction and dynamic change of chromatin states in rumen development during weaning. *Genomics* **114**, 110296 (2022).
14. Salavati, M. *et al.* Improving the annotation of the cattle genome by annotating transcription start sites in a diverse set of tissues and populations using CAGE sequencing. *G3 Genes|Genomes|Genetics* jkad108 (2023) doi:10.1093/g3journal/jkad108.
15. Beiki, H. *et al.* Enhanced bovine genome annotation through integration of transcriptomics and epi-transcriptomics datasets facilitates genomic biology. *GigaScience* **13**, giae019 (2024).
16. Moore, J. E. *et al.* Expanded encyclopaedias of DNA elements in the human and mouse genomes. *Nature* **583**, 699–710 (2020).
17. Boix, C. A., James, B. T., Park, Y. P., Meuleman, W. & Kellis, M. Regulatory genomic circuitry of human disease loci by integrative epigenomics. *Nature* **590**, 300–307 (2021).
18. Liu, S. *et al.* A multi-tissue atlas of regulatory variants in cattle. *Nat Genet* **54**, 1438–1447 (2022).
19. Liu, S. & Fang, L. The CattleGTEx atlas reveals regulatory mechanisms underlying complex traits. *Nat Genet* **54**, 1273–1274 (2022).
20. Li, J. *et al.* A multi-tissue atlas of allelic-specific expression reveals the characteristics, mechanisms, and relationship with dominant effects in cattle. *BMC Biology* **23**, 148 (2025).

21. Jaganathan, K. *et al.* Predicting Splicing from Primary Sequence with Deep Learning. *Cell*
176, 535-548.e24 (2019).
22. Zeng, T. & Li, Y. I. Predicting RNA splicing from DNA sequence using Pangolin. *Genome*
Biology **23**, 103 (2022).
23. Charles, M. *et al.* Functional impact of splicing variants in the elaboration of complex traits
in cattle. *Nat Commun* **16**, 3893 (2025).
24. Yuan, C. *et al.* An organism-wide ATAC-seq peak catalog for the bovine and its use to
identify regulatory variants. *Genome Res.* (2023) doi:10.1101/gr.277947.123.
25. Prowse-Wilkins, C. P. *et al.* Putative Causal Variants Are Enriched in Annotated Functional
Regions From Six Bovine Tissues. *Front. Genet.* **12**, (2021).
26. Kundaje, A. *et al.* Integrative analysis of 111 reference human epigenomes. *Nature* **518**, 317–
330 (2015).
27. Ernst, J. & Kellis, M. Large-scale imputation of epigenomic datasets for systematic
annotation of diverse human tissues. *Nat Biotechnol* **33**, 364–376 (2015).
28. Pan, Z. *et al.* Pig genome functional annotation enhances the biological interpretation of
complex traits and human disease. *Nat Commun* **12**, 5848 (2021).
29. Pan, Z. *et al.* An atlas of regulatory elements in chicken: A resource for chicken genetics and
genomics. *Science Advances* **9**, eade1204 (2023).
30. Fulco, C. P. *et al.* Activity-by-contact model of enhancer–promoter regulation from
thousands of CRISPR perturbations. *Nat Genet* **51**, 1664–1669 (2019).
31. Whyte, W. A. *et al.* Master Transcription Factors and Mediator Establish Super-Enhancers at
Key Cell Identity Genes. *Cell* **153**, 307–319 (2013).
32. Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription Factors Prime
cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell* **38**,
576–589 (2010).
33. Li, L., Chambard, J. C., Karin, M. & Olson, E. N. Fos and Jun repress transcriptional
activation by myogenin and MyoD: the amino terminus of Jun can mediate repression. *Genes*
Dev. **6**, 676–689 (1992).
34. Pataskar, A. *et al.* NeuroD1 reprograms chromatin and transcription factor landscapes to
induce the neuronal program. *The EMBO Journal* **35**, 24–45 (2016).
35. Ott, M.-O., Bober, E., Lyons, G., Arnold, H. & Buckingham, M. Early expression of the
myogenic regulatory gene, myf-5, in precursor cells of skeletal muscle in the mouse embryo.
Development **111**, 1097–1107 (1991).
36. Hadchouel, J. *et al.* Analysis of a key regulatory region upstream of the Myf5 gene reveals
multiple phases of myogenesis, orchestrated at each site by a combination of elements
dispersed throughout the locus. *Development* **130**, 3415–3426 (2003).
37. Francetic, T. & and Li, Q. Skeletal myogenesis and Myf5 activation. *Transcription* **2**, 109–
114 (2011).
38. Spitz, F. & Furlong, E. E. M. Transcription factors: from enhancer binding to developmental
control. *Nat Rev Genet* **13**, 613–626 (2012).
39. Sokolova, K., Chen, K. M., Hao, Y., Zhou, J. & Troyanskaya, O. G. Deep Learning Sequence
Models for Transcriptional Regulation. (2024) doi:10.1146/annurev-genom-021623-024727.
40. Pampari, A. *et al.* ChromBPNet: bias factorized, base-resolution deep learning models of
chromatin accessibility reveal cis-regulatory sequence syntax, transcription factor footprints
and regulatory variants. 2024.12.25.630221 Preprint at
https://doi.org/10.1101/2024.12.25.630221 (2024).

41. Liu, B. B. *et al.* Dissecting regulatory syntax in human development with scalable
multiomics and deep learning. 2025.04.30.651381 Preprint at
https://doi.org/10.1101/2025.04.30.651381 (2025).
42. Shrikumar, A., Greenside, P. & Kundaje, A. Learning Important Features Through
Propagating Activation Differences. Preprint at https://doi.org/10.48550/arXiv.1704.02685
(2019).
43. Shrikumar, A. *et al.* Technical Note on Transcription Factor Motif Discovery from
Importance Scores (TF-MoDISco) version 0.5.6.5. Preprint at
https://doi.org/10.48550/arXiv.1811.00416 (2020).
44. Fan, L. *et al.* Muscle Krüppel-like factor 15 regulates lipid flux and systemic metabolic
homeostasis. *J Clin Invest* **131**, (2021).
45. Li, H. *et al.* An atlas of cell type specific regulatory effects in cattle. 2025.06.23.661035
Preprint at https://doi.org/10.1101/2025.06.23.661035 (2025).
46. Prowse-Wilkins, C. P., Wang, J., Garner, J. B., Goddard, M. E. & Chamberlain, A. J. Allele
specific binding of histone modifications and a transcription factor does not predict allele
specific expression in correlated ChIP-seq peak-exon pairs. *Sci Rep* **13**, 15596 (2023).
47. Du, A. *et al.* Genetic polymorphisms of PKLR gene and their associations with milk
production traits in Chinese Holstein cows. *Front. Genet.* **13**, (2022).
48. Fonseca, P. A. de S., Caldwell, T., Mandell, I., Wood, K. & Cánovas, A. Genome-wide
association study for meat tenderness in beef cattle identifies patterns of the genetic
contribution in different post-mortem stages. *Meat Science* **186**, 108733 (2022).
49. Guo, D. *et al.* Lipid overload-induced RTN3 activation leads to cardiac dysfunction by
promoting lipid droplet biogenesis. *Cell Death Differ* **31**, 292–308 (2024).
50. Lee, D. *et al.* A method to predict the impact of regulatory variants from DNA sequence. *Nat*
Genet **47**, 955–961 (2015).
51. Marderstein, A. R. *et al.* Mapping the regulatory effects of common and rare non-coding
variants across cellular and developmental contexts in the brain and heart.
2025.02.18.638922 Preprint at https://doi.org/10.1101/2025.02.18.638922 (2025).
52. Xiang, R. *et al.* Quantifying the contribution of sequence variants with regulatory and
evolutionary significance to 34 bovine complex traits. *Proceedings of the National Academy*
of Sciences **116**, 19398–19408 (2019).
53. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).
54. Preissl, S., Gaulton, K. J. & Ren, B. Characterizing cis-regulatory elements using single-cell
epigenomics. *Nat Rev Genet* 1–23 (2022) doi:10.1038/s41576-022-00509-1.
55. Guan, D. *et al.* Genetic regulation of gene expression across multiple tissues in chickens. *Nat*
Genet 1–11 (2025) doi:10.1038/s41588-025-02155-9.
56. Teng, J. *et al.* A compendium of genetic regulatory effects across pig tissues. *Nat Genet* **56**,
112–123 (2024).
57. Scott, A. J., Chiang, C. & Hall, I. M. Structural variants are a major source of gene
expression differences in humans and often affect multiple nearby genes. *Genome Res.*
gr.275488.121 (2021) doi:10.1101/gr.275488.121.
58. Sasse, A., Chikina, M. & Mostafavi, S. Unlocking gene regulation with sequence-to-function
models. *Nat Methods* **21**, 1374–1377 (2024).

59. Agarwal, V. *et al.* Massively parallel characterization of transcriptional regulatory elements
in three diverse human cell types. 2023.03.05.531189 Preprint at
https://doi.org/10.1101/2023.03.05.531189 (2023).
60. Arnold, C. D. *et al.* Genome-Wide Quantitative Enhancer Activity Maps Identified by
STARR-seq. *Science* **339**, 1074–1077 (2013).
61. Tixier-Boichard, M. *et al.* Tissue Resources for the Functional Annotation of Animal
Genomes. *Frontiers in Genetics* **12**, 847 (2021).
62. Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables
interrogation of frozen tissues. *Nat Methods* **14**, 959–962 (2017).
63. Halstead, M. M. *et al.* Systematic alteration of ATAC-seq for profiling open chromatin in
cryopreserved nuclei preparations from livestock tissues. *Sci Rep* **10**, 5230 (2020).
64. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
Preprint at https://doi.org/10.48550/arXiv.1303.3997 (2013).
65. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biology* **9**, R137
(2008).
66. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).
67. Kovaka, S. *et al.* Transcriptome assembly from long-read RNA-seq alignments with
StringTie2. *Genome Biology* **20**, 278 (2019).
68. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).
69. Loyfer, N., Rosenski, J. & Kaplan, T. wgbstools: A computational suite for DNA methylation
sequencing data representation, visualization, and analysis. 2024.05.08.593132 Preprint at
https://doi.org/10.1101/2024.05.08.593132 (2024).
70. Loyfer, N. *et al.* A DNA methylation atlas of normal human cell types. *Nature* **613**, 355–364
(2023).
71. Ernst, J. & Kellis, M. Chromatin-state discovery and genome annotation with ChromHMM.
Nat Protoc **12**, 2478–2492 (2017).
72. Ernst, J. & Kellis, M. ChromHMM: automating chromatin-state discovery and
characterization. *Nat Methods* **9**, 215–216 (2012).
73. Gorkin, D. U. *et al.* An atlas of dynamic chromatin landscapes in mouse fetal development.
Nature **583**, 744–751 (2020).
74. Ernst, J. & Kellis, M. Discovery and characterization of chromatin states for systematic
annotation of the human genome. *Nat Biotechnol* **28**, 817–825 (2010).
75. Lovén, J. *et al.* Selective Inhibition of Tumor Oncogenes by Disruption of Super-Enhancers.
Cell **153**, 320–334 (2013).
76. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).
77. Camargo, A. P., Vasconcelos, A. A., Fiamenghi, M. B., Pereira, G. A. G. & Carazzolle, M. F.
tspex: a tissue-specificity calculator for gene expression data. *Research Square* Preprint at
https://doi.org/10.21203/rs.3.rs-51998/v1 (2020).
78. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server
2016 update. *Nucleic Acids Research* **44**, W90–W97 (2016).
79. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
(Methodological) **57**, 289–300 (1995).

80. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).
81. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).
82. McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
83. van de Geijn, B., McVicker, G., Gilad, Y. & Pritchard, J. K. WASP: allele-specific software
for robust molecular quantitative trait locus discovery. *Nat Methods* **12**, 1061–1063 (2015).
84. Li, Q., Brown, J. B., Huang, H. & Bickel, P. J. Measuring reproducibility of high-throughput
experiments. *Ann. Appl. Stat.* **5**, (2011).
85. Pockrandt, C., Alzamel, M., Iliopoulos, C. S. & Reinert, K. GenMap: ultra-fast computation
of genome mappability. *Bioinformatics* **36**, 3687–3692 (2020).
86. Ghandi, M., Lee, D., Mohammad-Noori, M. & Beer, M. A. Enhanced Regulatory Sequence
Prediction Using Gapped k-mer Features. *PLOS Computational Biology* **10**, e1003711
(2014).
87. Ghandi, M. *et al.* gkmSVM: an R package for gapped-kmer SVM. *Bioinformatics* **32**, 2205–
2207 (2016).
88. Lee, D. LS-GKM: a new gkm-SVM for large-scale datasets.
89. Wang, J. *et al.* Fine-mapping methods for complex traits: essential adaptations for samples of
related individuals. 2021.10.10.463846 Preprint at
https://doi.org/10.1101/2021.10.10.463846 (2025).
90. Zou, Y., Carbonetto, P., Wang, G. & Stephens, M. Fine-mapping from summary data with the
"Sum of Single Effects" model. *PLOS Genetics* **18**, e1010299 (2022).
91. Cheng, J. *et al.* SLEMM: million-scale genomic predictions with window-based SNP
weighting. *Bioinformatics* **39**, btad127 (2023).
92. Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies
disease-relevant tissues and cell types. *Nat Genet* **50**, 621–629 (2018).
93. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide
association summary statistics. *Nat Genet* **47**, 1228–1235 (2015).
94. Davydov, E. V. *et al.* Identifying a High Fraction of the Human Genome to be under
Selective Constraint Using GERP++. *PLOS Computational Biology* **6**, e1001025 (2010).
95. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff. *Fly* **6**, 80–92 (2012).
96. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**, 1017–1018 (2011).
97. Christmas, M. J. *et al.* Evolutionary constraint and innovation across hundreds of placental
mammals. 2023.03.09.531574 Preprint at https://doi.org/10.1101/2023.03.09.531574 (2023).
1013

a

301 unique motifs

42 unique tissue contexts

motifs

tissues detected

seqlets
- 1000
- 2000
- 3000
- 4000

Tissue
- Abomasum
- Adipose
- Bladder
- BoneMarrow
- Cecum Cerebellum
- C o l o n
- Cortex Epididymis
- Esophagus
- FetalBrain
- FetalGonad
- FetalHeart FetalKidney
- FetalLiver
- FetalSpleen Follicle
- Heart
- Hypothalamus
- Jejunum
- Kidney
- Liver Lung
- LymphNode
- MammaryLatePregnancy
- MammaryMiddlePregnancy
- Myometrium
- Myotubes Omasum
- Ovary
- Pancreas Pituitary
- PlacentalCaruncle
- PlacentalCotyledon
- Preadipocytes
- Reticulum
- Rumen
- Skin
- Spleen
- Testis
- Uterus

b CTCF

This study

JASPAR

c

| Cerebellum | Abomasum | Heart | Fetal heart | Preadipocytes |

CWM

JASPAR

TF - $\log_{10}(q)$

| NEUROD1 | FOSB::JUNB | MEF2B | THAP11 | Creb5 |
| 1.93 | 4.69 | 4.32 | 4.68 | 4.77 |

d

Rumen CWM

e

instances (×10⁵)

KLF15

%instances

301 unique motifs