Texas 2-Step: A new Model for YcgR::c-di-GMP Action at the Flagellar Motor

Nabin Bhattarai[1], Wangbiao Guo[2], Jonathan D Partridge[1], and Rasika M. Harshey*[1]

[1]Department of Molecular Biosciences and

LaMontagne Center for Infectious Diseases,

The University of Texas at Austin,

Austin, Texas, USA

[2]Microbial Sciences Institute & Department of Microbial Pathogenesis, Yale School of

Medicine, New Haven, Connecticut, USA

* Corresponding author: rasika@austin.utexas.edu

Abstract

YcgR is a c-di-GMP effector that inhibits chemotaxis and swimming speed in *Escherichia coli*

and *Salmonella.* Genetic, biochemical, and structural studies suggest that YcgR interacts

with both the bidirectional flagellar rotor and the stator to bias rotation toward

counterclockwise (CCW) and reduce motor speed, but the underlying mechanism remains

unresolved. Recent cryo-EM structures revealing conformational changes in the rotor–stator

complex during directional switching suggested to us a mechanism by which YcgR acts. We

call this the **Texas 2-Step** model, after the country dance in which partners move smoothly

in a CCW arc with quick steps followed by slow ones. In this model, YcgR first binds a MotA

subunit when the rotor adopts the CCW conformation, in which stators are largely displaced

from the C-ring. In the next step, the rotating MotA pentamer delivers YcgR to the rotor

protein FliG, thereby slowing motor speed. We provide evidence for the first step of this

model, offering testable predictions for future work.

Importance

The mechanism of YcgR action has been investigated by multiple laboratories using diverse

approaches, yet no consensus has emerged. Some studies implicate the rotor, others the

stator. A key complication is the involvement of four interacting proteins—MotA, FliG, FliM,

and YcgR—with multiple contact sites in several of them. Recent rotor–stator cryo-EM

structures revealing conformational changes during directional switching suggested a

mechanism that we set out to test. Our experiments show that rotor conformation is crucial

for YcgR function.

Introduction

Among the many outputs of c-di-GMP signaling, a ubiquitous one is inhibition of bacterial motility (1). The inhibitory effect on flagellar motility is well-studied in *E. coli* and *Salmonella* and is orchestrated via interaction of the c-di-GMP effector YcgR with both the flagellar rotor and stator (2-8). In these bacteria, YcgR expression is under the control of the flagellar regulon, where a three-tiered regulatory cascade integrates various environmental cues and ensures an ordered assembly of the flagellum (9). Curiously, both YcgR and PdeH (also called YhjH), a strong phosphodiesterase that degrades c-di-GMP, are expressed in the last tier of the cascade (10-13). PdeH is expected to keep c-di-GMP levels low when cells are all set to move, while YcgR is likely on guard to arrest motility if environmental conditions elevate c-di-GMP levels via the multiple diguanylate cyclases and phosphodiesterases that these bacteria encode (14). Accordingly, mutation of *pdeH*, expected to elevate c-di-GMP levels, has an inhibitory effect on swimming motility in both bacteria (10, 12, 15). A study that first demonstrated c-di-GMP-binding to the C-terminal PilZ domain of YcgR, also showed that the inhibitory effect of the *pdeH* mutant was abrogated by deleting *ycgR*, establishing later that YcgR::c-di-GMP is the only effector for motility inhibition (16, 17).

The flagellar motors of *E. coli* and *Salmonella* consist of a bi-directional rotor and dynamic stators that engage and disengage with the rotor in a load-dependent manner (18-22). The rotor includes the cytoplasmic C-ring made of three proteins (FliG, FliM, FliN). FliG is at the top of the ring, interlaced with inner membrane MS ring protein FliF, which in turn links to the periplasmic rod, external hook and the long helical flagellar filament. The stators are composed of pentameric MotA and dimeric MotB proteins (23, 24). They are independent units that drift in the inner membrane in a "closed" ion-nonconducting form until they encounter the flagellar basal body, upon which they dock on top of FliG in the C-ring and adopt an "open" ion-conducting conformation. Protons transiting through the MotAB complex cause conformational changes in MotA that power CW rotation of the pentameric MotA ring

around a stationary MotB, thought to attach to peptidoglycan or some stationary moiety in

the periplasm (25). Rotary MotA pushes on FliG, wherein a conserved patch of charged

residues in MotA interact with compatible residues in the 'torque helix' of FliG, generating

torque that results in the rotation of the C-ring and all parts linked to it (26). The C-ring can

accommodate ~11 stator units, but their numbers vary depending on external load. When

attached to the flagellar filament (high load), *E. coli* motors spin at ~125 Hz; when the

filament is absent (low load), motors rotate at 300 Hz or faster (27, 28).

The default state of the bi-directional C-ring is CCW. The chemotaxis signal CheY~P,

interacts with FliM/FliN located below FliG to elicit a large change in the C-terminal domain

of FliG (FliG$_C$) where the torque helix is located (29), as well as producing changes at FliG$_N$,

concomitantly switching the orientation of charges in the torque helix of FliG and the

rotational direction of the C-ring from CCW to CW (30-32). Between FliG$_C$ and FliG$_N$ is

located a 'cleft' whose width changes from 30Å in CCW state to 40Å in the CW state. As first

observed in *Borrelia burgdorferi* where the entire structure of the stator-associated rotor is

visible, changes in FliG$_C$ not only change the size of the FliG ring, but also change the

disposition of the stators, which move inwards in the CW state and outward in the CCW

state (33). Similar events at the C-ring were reported in *Salmonella* (30-32), where the

diameter of the upper FliG ring toggles between ~446 Å in the CW state to ~466 Å in the

CCW state (32). In the CW state, the rotating MotA pentamer would interact not only with

FliG$_C$ but also with FliG$_N$, where charged residues similar to the torque helix in FliG$_C$ are

located. A genetic fusion between stators of a closely related bacterium and the torque helix

in FliG$_c$ (30), as well as docking simulations (31, 32), show that like seen in *B. burgdorferi*,

stators must relocate in and out of the FliG ring.

YcgR action at the motor has been examined for over a decade, yet its mechanism remains

unresolved. Some studies implicate the stators, others the rotor, and still others both. A

persistent challenge is that YcgR interacts with multiple components—rotor proteins FliG

and FliM and the stator protein MotA—with contacts mapped to the N- and C-termini of both

FliG and YcgR. Recent rotor–stator cryo-EM structures revealing conformational changes

during directional switching suggested a mechanism that we now explore experimentally.

Results

A new model for YcgR action.

Multiple studies have demonstrated that in a *pdeH* mutant, where c-di-GMP levels are

elevated, chromosomal YcgR produces two distinct motor phenotypes – increased CCW

rotor bias and a reduction in motor speed (2-5, 7, 8, 13). In a *pdeH ycgR* double mutant

carrying an inducible plasmid expressing YcgR, the rotor bias of some motors shifted to

CCW before motor speed decreased (17), demonstrating that changes in bias and speed

can occur sequentially. This temporal separation suggests that two mechanically distinct

steps may be involved.

To explain these observations in the context of recent insights into the two C-ring

conformations, we propose a model we call the **Texas 2-step**, delineated in Figure 1, which

outlines the key components involved. The crystal structure of YcgR bound to c-di-GMP (6)

is shown in Figure 1A. In this structure, the PilZ domain of YcgR ($YcgR_C$) binds c-di-GMP

between $YcgR_N$ and $YcgR_C$ (16), triggering a rearrangement of $YcgR_C$ that exposes an α-

helix at the extreme C-terminus (C-tail) (Fig. 1A, green helix) (6). The exposed C-tail of

$YcgR_C$ interacts with MotA residues previously identified as contacting the torque helix of

FliG (26). This interaction is supported by mutations in both $YcgR_C$ and MotA that abolish

MotA-YcgR binding *in vitro* and YcgR-mediated motility inhibition *in vivo* (6). Consistent with

this, extragenic suppressors of YcgR also map to the same MotA residues (2). Moreover,

multiple studies have also demonstrated YcgR binding to FliG in the rotor (3, 4, 8).

The model. In the first step, YcgRc can access MotA only in stators engaged with the CCW

rotor conformation, where they project outward from the C-ring. This interaction is likely

prevented in the CW conformation, where the stators are expected to shift inward, shielding

the YcgR-binding region of MotA (compare Fig. 1B vs 1C).

In a second step, as the $YcgR_C$-bound MotA pentamer begins to rotate, $YcgR_N$ is positioned

to contact FliG. We envision two possible scenarios for this interaction. In the first, YcgR

bridges $FliG_C$ and MotA via its two domains, either arresting rotation of the bound MotA

pentamer (Fig. 1C, Scenario 1; Movie 1), or more plausibly, remaining attached to $FliG_C$

while MotA dissociates (not shown). In the second scenario, YcgR disengages from rotating

MotA and moves across the top of the C-ring to bind $FliG_N$ (Fig. 1C, Scenario 2; Movie 2)

from where it may weaken or prevent MotA binding. In both cases, YcgR occupation of FliG

stabilizes the CCW state, reducing rotation speed as additional YcgR molecules engage FliG.

Involvement of FliG in our proposed second step is supported by several lines of evidence.

Genetic analyses show that mutation in the $FliG_C$ torque helix residues relieve YcgR-

mediated motility inhibition (4). Two-hybrid and pull-down assays demonstrate that YcgR

interacts with both $FliG_N$ and $FliG_C$ (3). Moreover, TIRF experiments measuring eGFP-YcgR

fluorescence at the motor before and after addition of NaN_3 – an uncoupler that dissipates

the proton motive force and causes stators to dissociate – showed no change in

fluorescence intensity, indicating that YcgR remains stably associated with the rotor (5). The

closest rotor protein to MotA-bound YcgR is FliG, consistent with these observations.

Involvement of $YcgR_N$ in motility inhibition is also supported by genetic evidence: even when

the MotA-interacting α-helix in YcgR☐ remains intact and functional, mutations in $YcgR_N$

residues disrupt motility (4, 8). Thus, interaction of YcgR with MotA alone cannot fully

account for all YcgR-dependent effects.

In this study, we focus exclusively on **Step 1** of the proposed model, which makes a clear

prediction: this step should be blocked in CW-locked rotors.

<u>Testing the first step of the model</u>. We used two *E. coli* parent strains - HCB5 and MG1655 -

to ensure that observed motor behaviors were not strain-dependent. HCB5 carried only a

pdeH deletion (WT*), while MG1655 was deleted for both *pdeH* and *ycgR* (WT**). To

generate motors that rotated predominantly in one direction, we introduced into these strains

deletions in genes encoding the chemotaxis signaling components or the rotor protein FliG:

Δ*cheY* - CCW, Δ*cheZ* - CW, *fliG*$_{\Delta PAA}$ (deletion of PAA residues 169-171) - extreme or 'locked'

CW (34). In both strain backgrounds, YcgR was supplied from an inducible plasmid, as

described previously (17).

Motor behavior was monitored using a bead assay, where a 0.75 μm polystyrene bead was

attached to a stub of a sheared flagellar filament on bacteria fixed to a glass slide, and bead

rotation was captured by high-speed videos (see Methods). For most strains, at least 15

motors were recorded continuously for 10 min following inducer addition.

The behavior of the *pdeH ycgR* (WT**) strain upon YcgR induction was reported previously

(17) and is reproduced here in Fig. S1A (top trace). The *pdeH* strain (WT*) showed a similar

response (Fig. 2A and S1A bottom trace). This strain expresses chromosomal levels of

YcgR, and the motors display the expected variation in CCW bias and speed. Upon plasmid-

driven YcgR induction, we observed three distinct motor behaviors: a strong CCW bias with

no reduction in initial speed (Fig. 2A, top trace); a CCW bias accompanied by a gradual

speed reduction (Fig 2A, bottom trace); or no change in bias prior to sudden motor arrest

(Fig. S1A, bottom trace). These observations indicate that the motor response to YcgR

induction is independent of the strain background.

The Δ*cheY* strain, which has an initial CCW bias, also displayed two behaviors: a graded

speed reduction over time (Fig. 2B, top trace), or an abrupt stop without prior speed

reduction (Fig. 2B, bottom trace).

The two CW-biased strains - Δ*cheZ* and FliGΔPAA – were largely impervious to YcgR (Fig.

2C-D). Δ*cheZ* motors rotated predominantly CW, with an 'unstable' bias characterized by

occasional excursions into the CCW zone, yet they remained mostly unresponsive to YcgR

(Fig. 2C). When a larger bead (0.99 μm) was used, the CW bias of Δ*cheZ* motors became

even more unstable, switching rapidly between the two conformations (Fig. S1B, traces

before YcgR induction). Upon YcgR induction, two Δ*cheZ* motors exhibited a shift to CCW

accompanied by speed reduction (Fig. S1B, top and bottom traces at 9 and 6 min,

respectively), followed in one case by a pronounced CCW bias and speed reduction (Fig.

S1B, bottom), resembling the motor traces shown in Figure 3A. These observations

indicated that even Δ*cheZ* motors that respond to YcgR do so primarily in the CCW regime.

FliGΔPAA motors displayed the expected stable or 'locked' CW bias, with most motors

resistant to YcgR (Fig. 2D, top and bottom). However, in both WT* and WT**, approximately

10% of the CW-locked motors showed either a gradual decrease in speed (Fig. S1CD, top),

a gradual decrease followed by a stop (Fig. S1C, bottom), or a sudden stop (Fig. S1D,

bottom). These observations indicate that YcgR can access a minority of CW-locked motors.

We note that the cleft between $FliG_N$ and $FliG_C$ is larger in the CW rotor, and that a density

consistent with YcgR has been observed in one data set from FliGΔPAA motors (31).

A summary of the motors tracked in this study is shown in Figure 3. While majority of the

data were generated with 0.75μm beads, we also monitored WT** motors with smaller

(0.45μm) and larger (0.99μm) beads, which are expected to impose lower and higher loads

on the motor, respectively (Fig. 3C). A clear trend emerged: upon YcgR induction, more

motors stopped when smaller beads were used compared with larger ones. This trend is

consistent with the expected relationship between bead size and motor torque, as smaller

beads engage fewer stators at the rotor, while larger beads engage more, explaining the

load-dependent differences in motor response.

Discussion

Using strains whose motors rotate predominantly in the CCW or CW directions, we show

that YcgR primarily affects motors in the CCW conformation, supporting the proposed first

step our model (Fig. 1C). Dependence on the CCW conformation would facilitate YcgR

access to the stators, where YcgR binds MotA—an interaction supported by genetic and

biochemical evidence (2, 6). Binding of YcgR to MotA is expected to disrupt MotA's normal

torque-generating interaction with FliG, thereby reducing motor speed. This first step,

however, cannot fully account for YcgR function, as YcgR also alters rotor bias.

We therefore propose a second step in which $YcgR_N$ binds $FliG_C$, the rotor subunit most

proximal to MotA-bound YcgR (Fig.1, Scenario 1) (see 'A new model' above for supporting

evidence). Such an interaction would drive the rotor towards a CCW conformation, coupling

bias stabilization with progressive speed reduction, and eventually leading to motor arrest as

more YcgR molecules engage the rotor (Movie1).

Scenario 1 in the proposed second step could, in principle, explain both the CCW bias shift

and the accompanying speed reduction observed in WT motors. To account for WT motors

that shift to a CCW bias **without** an initial decrease in speed (Fig. 2A & S1A, top traces),

one would have to assume that YcgR can remain bound to the rotating MotA pentamer

without affecting torque, which seems unlikely. Instead, Scenario 2 - where $YcgR_N$ binds

$FliG_N$ - better accommodates this behavior. This scenario is supported by findings that YcgR

interacts with both $FliG_N$ and $FliG_C$ in two-hybrid and pull-down assays (3). A trivial

explanation for these dual interactions could be that sequence homologies between the two

FliG domains reflect their shared MotA-contacting surfaces in the distinct rotor conformations.

Figure 4 presents a final model, shown as an aerial view of the C-ring, summarizing support

for Step 1 from the data obtained in this study.

<u>Additional Mechanistic Insights</u>

1. Given the variety of experimental protocols demonstrating binding of YcgR to the rotor (3-

5, 8), why did extragenic suppressors of YcgR only map to MotA residues that interact with

FliG (2), and not to FliG itself? To address this question, we re-isolated YcgR suppressors

and recovered the same MotA mutations - G93E, G93R, G93V and S96L – along with a new

mutation, Q91P. None mapped to *fliG*. We also obtained these suppressors from the Jenal

lab (Basel, Switzerland) and characterized their motor behavior (Fig. S2). The suppressors

were identified by their ability to swim in soft agar plates (Fig. S2A), an assay that requires

functional chemotaxis. Motor assays revealed that the suppressors exhibited normal motor

bias but reduced rotation speeds (50–70 Hz) compared to wild type (80 Hz) (Fig.S2 B,C).

These reduced speeds were intrinsic to the suppressor mutations, as they persisted even in

the absence of YcgR (Fig. S2B, $\Delta ycgR$, S96L shown).

We consider two explanations for why YcgR suppressors did not map to *fliG*. First, because

YcgR can interact with both $FliG_N$ and $FliG_C$, the probability of mutationally disrupting both

binding sites is low. Alternatively, mutations in these regions of FliG may compromise motor

bias and thus fail to be recovered in the chemotaxis-dependent motility screens used for

suppressor isolation. Indeed, mutations targeting the torque helix of FliG severely impair

motility in soft agar (4).

2. In a FRAP study using TIRF microscopy, 36% of mEGFP-YcgR molecules expressed in a

pdeH mutant were found to be weakly bound to the motor, with the remaining fraction tightly

bound (7). Evidence that the tightly bound YcgR population localizes to the rotor rather than

the stator comes from a similar TIRF-based analysis showing that depletion of the proton

motive force did not alter YcgR fluorescence (5). Moreover, two independent studies that

directly interrogated interactions with both MotA and FliG identified MotA as the primary

target of YcgR (6, 8). Taken together, these findings suggest that YcgR initially associates

with MotA, which induces a conformational change that enables its subsequent tight binding

to FliG.

3. Motor behavior observed at chromosomal levels of YcgR (Fig. S2; *pdeH*) as well as at

plasmid-induced levels (Fig. S1B; *pdeH ycgR cheZ*) shows that, despite reduced speed and

a CCW bias, these motors are still capable of switching. We envision two possibilities: (i)

YcgR-bound FliG subunits do not impede rotor switching, or (ii) YcgR-occupied subunits

become functionally or spatially isolated from the remainder of the rotor, which continues to

switch normally.

4. Significant binding of YcgR to FliM has been reported in only one study, which employed

three independent approaches—pulldown assays, visualization of YcgR–GFP at the rotor,

and motility analyses (4). Interestingly, two YcgR variants, R118D and S147A, which differ in

their ability to bind c-di-GMP and modulate motor function, were both found to interact with

FliM, albeit more weakly than with FliG (3, 4, 16). These findings collectively make a credible

case for YcgR–FliM interaction. Given that the R118D mutant, reported to be defective in c-

di-GMP binding, was nevertheless observed at the motor (4), we consider the possibility that

FliM serves as a reservoir for YcgR, positioning it to engage the motor rapidly upon an

increase in intracellular c-di-GMP levels.

5. In contrast to *E. coli* YcgR, the *Bacillus subtilis* homolog MotI (also known as DgrA) does

not influence motor bias but instead functions as a molecular clutch, binding to MotA and

disengaging it from the C-ring to halt rotation (35). The evolutionary advantages of

employing distinct mechanisms to control motor activity through these related proteins

remain speculative. Nevertheless, their behaviors converge under certain conditions: at

subinhibitory concentrations, MotI mimics YcgR by reducing swimming speed, whereas at

high concentrations, YcgR mimics MotI by stopping the motor.

Acknowledgments

243 This model was inspired by the first published images of the *Borrelia* motor from Jun Liu's

244 lab showing differences in stator disposition at the rotor; we thank him for discussions. We

245 also thank Urs Jenal and Birgit Scharf for strains. This work was supported by NIH grant

246 GM118085 to RMH.

Methods

Strains and growth conditions

Strains and plasmids used in this study are listed in Table S1. Bacterial cultures were grown in Lennox broth (LB) base (20 g/liter) (Fisher BioReagents). For motility assays in soft agar, 8 μl of an exponential-phase culture (optical density at 600 nm [OD_{600}], ~0.6) was inoculated onto plates made with 0.3% Bacto agar (Difco) and incubated at 30°C. All plate images shown are representative of three biological replicates, each in triplicate. Where required, the following antibiotics were used: ampicillin (100 μg/ml), chloramphenicol (20 μg/ml), and kanamycin (50 μg/ml). For inducible plasmids, 50uM IPTG (isopropyl-β-d-thiogalactopyranoside) and 0.2% L-arabinose were used.

Genetic manipulation

The WT parent strains for *E. coli* were MG1655 and HCB5. Mutant strains were constructed by inserting a kanamycin resistance cassette (KAN) into the designated gene as previously described (36) or sourced from the Keio collection (37). Mutations were transferred to fresh backgrounds by phage P1 (P1 Cm) transduction. Excision of the inserted KAN cassettes was achieved by expression of the FLP recombinase encoded by pCP20 (36).The resulting strains were confirmed by DNA sequencing. A 3 amino acid deletion of residues 169-171 (PAA) in FliG was introduced using the CRMAGE protocol (38), which leverages CRISPR-Cas9 technology as follows: A cocktail of 3 primer pairs for making guide RNAs (gRNA) targeting *fliG* were cloned into pMAZ-SK plasmid via Golden Gate assembly; gRNAs are expressed with anhydrotetracycline (aTetracycline). These constructs were introduced along with repair oligonucleotides into NBN28 containing pMA7CR_2.0, which expresses the λ Red β-protein and the CRISPR/Cas9 protein upon induction with L-arabinose. The insertion was confirmed by PCR amplification followed sequencing. All the primers and

oligonucleotides used are listed in Table S2.

Measurement of motor dynamics using bead assay

The bead assay was conducted as described in the previous work (17, 39). *E. coli fliC*

deletion strains (containing additional chromosomal mutations) were complemented with

pFD313 expressing the 'sticky' FliC variant (40), and pSEVA224 expressing YcgR. First,

cells were passed through two syringes connected by a 7" polyethylene capillary to shear

the flagella. A 40 µL cell suspension was then added to a poly-l-lysine-coated cover slip,

incubated for 10 minutes, and rinsed with motility buffer to remove unattached cells (41). A

40 µL solution of polystyrene beads (1:50 dilution) was added and left for 10 minutes before

excess beads were washed away, prior to introducing IPTG for inducing YgcR expression.

Bead rotations were captured at 1,000 frames/s using a CCD camera (ICL-B0620M-KC0;

Imperx, Boca Raton, FL). Videos were processed using custom analytical programs within

LabVIEW 2012 as described before (17).

Animation

Animation illustrating YcgR-mediated regulation of the stator–C-ring complex. was created

using ChimeraX 1.9 and assembled in Keynote. Structural models used include YcgR (PDB:

5Y6F), C-ring in CW and CCW conformations (PDB: 8UMX and 8UMD, respectively), and

MotA (PDB: 6YKM).

Figure Legends

Fig 1. Two-Step Model of YcgR Action at the Motor. **A.** Crystal structure of YcgR bound to c-di-GMP (PDB: 5Y6F). The α-helix involved in MotA interaction is shown in green, C-terminal domain (YcgR$_C$) in orange, and N-terminal domain (YcgR$_N$) in magenta. **B.** CW (Clockwise) C-ring conformation (PDB: 8UMX; FliNMG subunits are in aqua) with a docked CW-rotating MotA pentamer (PDB: 6YKM; yellow). **C.** CCW (Counterclockwise) C-ring conformation (PDB: 8UMD). Unlike in B, the α-helix of YcgR has clear access to the bottom of MotA. This is Step 1. There are two possibilities for Step 2. In Scenario 1, as the MotA pentamer rotates, YcgR moves with it until YcgR$_N$ contacts the C-terminal domain of a FliG subunit (FliG$_C$). Straddling the MotA–FliG$_C$ interface, YcgR obstructs MotA rotation. In an alternate Scenario 2, YcgR traverses the top of the C-ring to bind FliG$_N$, settling into the cleft between FliG$_N$ and FliG$_C$.

Fig 2. Response of WT, CCW and CW motors to YcgR. **A.** Two representative motors from WT* HCB5 strain are shown. YcgR was supplied from an IPTG-inducible low copy plasmid pSEVA224. Motor behavior was monitored by recording the motion of 0.75 μm polystyrene beads attached to flagella filament stubs (see Methods). After IPTG addition (arrow), 60-second representative segments from each consecutive 3-minute recording interval are shown. CCW (-) and CW (+) speeds in Hz are shown on the Y-axis. The profiles are representative of 15 individual motors tracked (see Fig. 3). **B.** As in A, except the strain is WT** MG1655 and has *cheY* deleted. **C.** As in B, except *cheZ* is deleted. **D.** As in B, except *fliG* is deleted for nucleotides encoding PAA at residues 169-171. Relevant genotypes of the strains are indicated on the top right-hand corner.

Fig 3. Summary of all motor traces. **A,B.** The relevant mutation expected to affect motor behavior is indicated directly under each bar graph. Genotype of WT* and WT** strains, common to all motors, is under the line. Bead size is 0.75 μm for all graphs, except for Δ*cheZ**, where the size was 0.99 μm. Color key indicates the motor behaviors observed. **C.** Behavior of WT** motors under varying bead sizes. p-values were calculated only for motors showing 'no effect' compared to WT. Fisher's exact test was used to assess statistical significance; *** indicates $p < 0.001$.

Fig 4. Final model with an aerial view of the FliG ring. **A.** YcgR only affects motors in the CCW conformation, supporting the proposed first step of our model. The MotA pentamer is accessible for binding in this conformation. **B.** To affect both speed and bias, YcgR must engage with both the stator and the rotor. This second step could be achieved by delivery of YcgR to either the outer (FliG$_C$) or inner (FliG$_N$) FliG rings by the rotating MotA pentamer.

References

1. Jenal U, Reinders A, Lori C. 2017. Cyclic di-GMP: second messenger extraordinaire. Nat Rev Microbiol 15:271-284.

2. Boehm A, Kaiser M, Li H, Spangler C, Kasper CA, Ackermann M, Kaever V, Sourjik V, Roth V, Jenal U. 2010. Second messenger-mediated adjustment of bacterial swimming velocity. Cell 141:107-16.

3. Fang X, Gomelsky M. 2010. A post-translational, c-di-GMP-dependent mechanism regulating flagellar motility. Mol Microbiol 76:1295-305.

4. Paul K, Nieto V, Carlquist WC, Blair DF, Harshey RM. 2010. The c-di-GMP binding protein YcgR controls flagellar motor direction and speed to affect chemotaxis by a "backstop brake" mechanism. Molecular cell 38:128-39.

5. Wang R, Wang F, He R, Zhang R, Yuan J. 2018. The Second Messenger c-di-GMP Adjusts Motility and Promotes Surface Aggregation of Bacteria. Biophys J 115:2242-2249.

6. Hou YJ, Yang WS, Hong Y, Zhang Y, Wang DC, Li DF. 2020. Structural insights into the mechanism of c-di-GMP?bound YcgR regulating flagellar motility in Escherichia coli. Journal of Biological Chemistry 295:808-821.

7. Liu X, Zhang C, Zhang R, Yuan J. 2022. The Effect of the Second Messenger c-di-GMP on Bacterial Chemotaxis in Escherichia coli. Appl Environ Microbiol 88:e0037322.

8. Han Q, Wang SF, Qian XX, Guo L, Shi YF, He R, Yuan JH, Hou YJ, Li DF. 2023. Flagellar brake protein YcgR interacts with motor proteins MotA and FliG to regulate the flagellar rotation speed and direction. Front Microbiol 14:1159974.

9. Chevance FF, Hughes KT. 2008. Coordinating assembly of a bacterial macromolecular machine. Nature Rev Microbiol 6:455-65.

10. Ko M, Park C. 2000. Two novel flagellar components and H-NS are involved in the motor function of *Escherichia coli*. J Mol Biol 303:371-82.

11. Wang Q, Mariconda S, Suzuki A, McClelland M, Harshey RM. 2006. Uncovering a large set of genes that affect surface motility in *Salmonella enterica* serovar Typhimurium. J Bacteriol 188:7981-4.

12. Frye J, Karlinsey JE, Felise HR, Marzolf B, Dowidar N, McClelland M, Hughes KT. 2006. Identification of new flagellar genes of *Salmonella enterica* serovar Typhimurium. J Bacteriol 188:2233-43.

13. Girgis HS, Liu Y, Ryu WS, Tavazoie S. 2007. A comprehensive genetic characterization of bacterial motility. PLoS Genet 3:1644-60.

14. Hengge R. 2009. Principles of c-di-GMP signalling in bacteria. Nat Rev Microbiol 7:263-73.

15. Simm R, Morr M, Kader A, Nimtz M, Romling U. 2004. GGDEF and EAL domains inversely

328   regulate cyclic di-GMP levels and transition from sessility to motility. Mol Microbiol 53:1123-34.

329 16. Ryjenkov DA, Simm R, Romling U, Gomelsky M. 2006. The PilZ domain is a receptor for the
330   second messenger c-di-GMP: the PilZ domain protein YcgR controls motility in enterobacteria.
331   J Biol Chem 281:30310-4.

332 17. Nieto V, Partridge JD, Severin GB, Lai RZ, Waters CM, Parkinson JS, Harshey RM. 2019.
333   Under Elevated c-di-GMP in Escherichia coli, YcgR Alters Flagellar Motor Bias and Speed
334   Sequentially, with Additional Negative Control of the Flagellar Regulon via the Adaptor Protein
335   RssB. J Bacteriol 202.

336 18. Berg HC. 2008. Bacterial flagellar motor. Current biology : CB 18:R689-91.

337 19. Lele PP, Hosu BG, Berg HC. 2013. Dynamics of mechanosensing in the bacterial flagellar
338   motor. Proc Natl Acad Sci U S A 110:11839-44.

339 20. Tipping MJ, Delalez NJ, Lim R, Berry RM, Armitage JP. 2013. Load-dependent assembly of
340   the bacterial flagellar motor. MBio 4.

341 21. Baker AE, O'Toole GA. 2017. Bacteria, Rev Your Engines: Stator Dynamics Regulate
342   Flagellar Motility. J Bacteriol 199.

343 22. Wadhwa N, Tu Y, Berg HC. 2021. Mechanosensitive remodeling of the bacterial flagellar
344   motor is independent of direction of rotation. Proc Natl Acad Sci U S A 118.

345 23. Deme JC, Johnson S, Vickery O, Aron A, Monkhouse H, Griffiths T, James RH, Berks BC,
346   Coulton JW, Stansfeld PJ, Lea SM. 2020. Structures of the stator complex that drives rotation
347   of the bacterial flagellum. Nat Microbiol 5:1553-1564.

348 24. Santiveri M, Roa-Eguiara A, Kuhne C, Wadhwa N, Hu H, Berg HC, Erhardt M, Taylor NMI.
349   2020. Structure and Function of Stator Units of the Bacterial Flagellar Motor. Cell 183:244-
350   257 e16.

351 25. Guo S, Liu J. 2022. The Bacterial Flagellar Motor: Insights Into Torque Generation, Rotational
352   Switching, and Mechanosensing. Front Microbiol 13:911114.

353 26. Kojima S, Blair DF. 2001. Conformational change in the stator of the bacterial flagellar motor.
354   Biochemistry 40:13041-50.

355 27. Yuan J, Berg HC. 2008. Resurrection of the flagellar rotary motor near zero load. Proc Natl
356   Acad Sci U S A 105:1182-5.

357 28. Berg HC. 2003. The rotary motor of bacterial flagella. Annu Rev Biochem 72:19-54.

358 29. Zhou J, Lloyd SA, Blair DF. 1998. Electrostatic interactions between rotor and stator in the
359   bacterial flagellar motor. Proc Natl Acad Sci U S A 95:6436-41.

360 30. Johnson S, Deme JC, Furlong EJ, Caesar JJE, Chevance FFV, Hughes KT, Lea SM. 2024.
361   Structural basis of directional switching by the bacterial flagellum. Nat Microbiol 9:1282-1292.

362 31. Singh PK, Sharma P, Afanzar O, Goldfarb MH, Maklashina E, Eisenbach M, Cecchini G,
363   Iverson TM. 2024. CryoEM structures reveal how the bacterial flagellum rotates and switches
364   direction. Nat Microbiol 9:1271-1281.

365 32. Tan J, Zhang L, Zhou X, Han S, Zhou Y, Zhu Y. 2024. Structural basis of the bacterial
366   flagellar motor rotational switching. Cell Res 34:788-801.

367   33.   Chang Y, Zhang K, Carroll BL, Zhao X, Charon NW, Norris SJ, Motaleb MA, Li C, Liu J. 2020.
368   Molecular mechanism for rotational switching of the bacterial flagellar motor. Nat Struct Mol
369   Biol 27:1041-1047.

370   34.   Togashi F, Yamaguchi S, Kihara M, Aizawa SI, Macnab RM. 1997. An extreme clockwise
371   switch bias mutation in fliG of Salmonella typhimurium and its suppression by slow-motile
372   mutations in motA and motB. J Bacteriol 179:2994-3003.

373   35.   Subramanian S, Gao X, Dann CE, 3rd, Kearns DB. 2017. MotI (DgrA) acts as a molecular
374   clutch on the flagellar stator protein MotA in Bacillus subtilis. Proc Natl Acad Sci U S A
375   114:13537-13542.

376   36.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia*
377   *coli* K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640-5.

378   37.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner
379   BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout
380   mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

381   38.   Ronda C, Pedersen LE, Sommer MO, Nielsen AT. 2016. CRMAGE: CRISPR Optimized
382   MAGE Recombineering. Sci Rep 6:19452.

383   39.   Partridge JD, Nieto V, Harshey RM. 2015. A new player at the flagellar motor: FliL controls
384   both motor output and bias. MBio 6:e02367.

385   40.   Ryu WS, Berry RM, Berg HC. 2000. Torque-generating units of the flagellar motor of
386   *Escherichia coli* have a high duty ratio. Nature 403:444-7.

387   41.   Partridge JD, Harshey RM. 2020. Investigating Flagella-Driven Motility in Escherichia coli by
388   Applying Three Established Techniques in a Series. J Vis Exp doi:10.3791/61364.