Somatic Programmed DNA Elimination is widespread in free-living Rhabditidae nematodes

Caroline Launay*, Eva Wenger*, Brice Letcher[#], Marie Delattre[#]

[#]**Corresponding author**: marie.delattre@ens-lyon.fr, brice.letcher@ens-lyon.fr

*Contributed equally

Affiliations

Laboratory of Biology and Modelling of the Cell, Ecole Normale Supérieure de Lyon, CNRS, Inserm, UCBL,

69007 Lyon, France

Abstract

All cells of a multicellular organism usually share an identical genome, faithfully transmitted through

successive divisions. Yet, a number of animal species deviate from this dogma, as parts of their DNA are

systematically eliminated in all their somatic nuclei, in a process called Programmed DNA Elimination

(PDE). PDE leads to the unexpected reorganisation of the genome at every generation in all somatic cells but

its molecular mechanism, evolutionary origins, and functional significance remain unknown. This lack of

understanding partially stems from limitations in genetically tractable model species.

PDE can target an entire chromosome, or involve chromosome fragmentation followed by selective

fragment retention and elimination, raising further questions on genome stability, genome integrity and

mechanisms of DNA repair. PDE by chromosome fragmentation has been described in parasitic nematodes

in the family Ascarididae, copepods in the genus Cyclops and unicellular ciliates. More recently, PDE has

been discovered in three non-parasitic, lab-tractable nematode species from the Rhabditidae family, opening

new perspectives.

In this study, we used cytological approaches to screen 25 new Rhabditidae species for PDE. We

found evidence of PDE in 17 species. Our work reveals that PDE is present in 12 out of 17 tested genera,

demonstrating its widespread presence in Rhabditidae nematodes, with the notable exception of *C. elegans*.

Genetic tools have already been established for some species. This work provides a collection of lab-

tractable species that can be used to test many aspects of somatic Programmed DNA Elimination by

chromosome fragmentation in animals.

Introduction

Some species systematically eliminate portions of their genome in their soma, in a process called Programmed DNA Elimination (PDE) [1,2]. PDE can target an entire chromosome, as in songbirds or lamprey for instance [3,4]. PDE can also happen by the fragmentation and partial destruction of chromosomes, as spotted for the first time in 1887 in the parasitic nematode *Parascaris univalens* [5]. PDE by chromosome fragmentation was later identified across the phylum of ciliate protists, Ciliophora [6]. Ciliates are unicellular organisms with two nuclei: a transcriptionally inactive micronucleus (MIC) transmitted to descendants after a sexual phase, and a transcriptionally active macronucleus (MAC), copied from the MIC and amplified. During MAC development, ciliates undergo extensive PDE, often involving both excision of internal sequences, healed by non-homologous end-joining, as well as chromosome fragmentation, as found in ascarid parasitic nematodes. PDE is pervasive in ciliates. Its study in a few species has led to seminal discoveries on the role of small RNAs in genome rearrangement [6]. It also revealed that internal eliminated sequences are remnants of transposable elements, excised by a domesticated transposase [7–9]. This, together with many internal eliminated sequences being also transposon-derived, has led a widely accepted scenario that excision of internal sequences emerged as a selfish, TE-driven process in ciliates [10]. In striking contrast, PDE through chromosome fragmentation remains poorly understood in ciliates, and its molecular dissection in animals has been largely precluded because ascarids are obligate parasites and not experimentally tractable. Moreover, the functional importance of somatic PDE is also unknown because it has not been possible to abolish PDE in any animal species so far.

Recently, PDE was identified in two nematode genera belonging to the Rhabditidae family, in a different sub-order than the one including ascarids. In *Oscheius tipulae*, telomere-to-telomere genome reconstruction revealed that 0.6 % of the genome is eliminated at chromosome ends [11,12]. Almost concomitantly, and by serendipity, we discovered substantial PDE in two other Rhabditidae species - although distant from *O. tipulae* - using cytology initially: *Mesorhabditis belari* and *Mesorhabditis spiculigera*. These two species eliminate approximately 30% and 20% of their genome, respectively, at all chromosome ends but also in large internal chromosomal blocks, leading to an increased somatic karyotype, as shown in Ascarids [13].

We could easily follow chromosome fragmentation in *Mesorhabditis* embryos, across the successive

embryonic divisions. Indeed, in a large range of nematode species, including ascarids and Rhabditidae, the

embryonic cell lineage is highly reproducible and conserved. Moreover, the germline precursor cell is easily

identified: it is the smallest and most posterior cell; it does not divide until late in embryogenesis, or even

after hatching in some species; and it harbours highly compacted chromatin [14]. Somatic cells also divide in

a relatively conserved order, allowing for easy identification of cell types from one species to the other, by

analogy with what has been well described in *C. elegans* [14]. In *Mesorhabditis* embryos, as previously

described in ascarids, we found that PDE starts with chromosome fragmentation in all five somatic precursor

cells, successively and independently, as they enter mitosis. This is followed by fragment exclusion during

mitosis and DNA destruction later in development. We also found in *Mesorhabditis*, as shown for ascarids or

O. tipulae, that telomeres of the germline inherited chromosomes are systematically eliminated in the soma,

and that new telomeres are assembled on the new extremities of all somatic chromosomes [12,13,15].

The finding that free-living nematode species undergo PDE following the same mechanism as in

ascarids demonstrates that PDE is not a parasitic oddity in nematodes and raises the possibility that PDE,

rather than having evolved independently in two distantly related clades, could be more widespread than

anticipated in nematodes.

In this study, we asked whether PDE is widespread in family Rhabditidae, which originated

approximately 200 million years ago [16]. Using simple DNA staining, we reveal that 17 out of 25 newly

tested species show evidence of PDE. This corresponds to 12 out of 17 genera in which at least one species

undergoes PDE. This work reveals that PDE is widespread in Rhabditidae, with the notable exception of *C.*

elegans, and has gone unnoticed for more than a century of nematode research.

Results

Identification of Programmed DNA Elimination by DNA staining

In ascarids and *Mesorhabditis* species, DNA elimination is immediately visible in early embryos because chromosomes are fragmented into large blocks [5,13,17]. Prior to elimination, no fragments are visible and somatic nuclei harbour the same number of chromosomes as germ cells. During elimination, chromosome fragments can be seen excluded from the spindle during mitosis or are visible in the cytoplasm in interphasic cells. After elimination, DNA fragments are eventually destroyed after a few cell cycles and old embryos do not show cytoplasmic DNA fragments [13,15]. When fragments internal to chromosomes are also eliminated, as in ascarids or *Mesorhabditis*, the chromosome number is increased in somatic nuclei compared to germline nuclei. Overall, it is possible to detect PDE using simple DNA staining on embryos at different time points during their development.

In order to detect PDE, we performed DNA staining on embryos from 25 new species. These species belong to 17 different genera in the Rhabditidae family, which include previously studied *Oscheius* and *Mesorhabditis* genera. We unambiguously identified cytoplasmic DNA fragments in 17 Rhabditidae new species spanning 12 genera (Fig. 1, Supp. Fig. 1).

We reconstructed the dynamics of fragment production in all species (Fig.1 and Supp. Fig. 1). We found that fragments were initially absent in 1, 2 and 4-cell stage embryos. Fragments then appeared rapidly (possibly during a single mitotic division) and, for any given strain, reproducibly in time. After a few cell divisions, fragments were no longer detectable suggesting they had been physically destroyed. Importantly, fragments were not detected in the germ cell primordium (Fig.1 and Supp. Fig. 1). For some species we could confirm that in cells that have generated cytoplasmic fragments, the number of chromosomes in their nuclei was higher than the number of chromosomes found in early embryos or in germ cells (Supp. Fig. 1). For instance, we counted 2n=12 chromosomes in the germline of *Pelodera teres* JU4407, and found more than 26 small fragments in somatic cells from old embryos (Supp Fig. 1). This is compatible with a scenario where for each chromosome, at least one internal chromosomal fragment has been eliminated, as shown in ascarids or *Mesorhabditis* [13,15]. We ruled out that our observed fragments corresponded to intracellular

bacteria, as cytoplasmic DNA fragments appeared and disappeared specifically in somatic cells, and not in

the germline primordium.

We found large variation in the amount of eliminated DNA (Fig. 1 and Supp. Fig. 1). In some cases,

we found very large and numerous fragments, as in *Mesorhabditis simplex*, or more dramatically even in

Cruznema sp. JU4406 (Fig. 1). For other species, the amount of eliminated DNA seems more limited as

fragments are small. This is notably the case in *Rhabditella axei* (Fig.1), *Oscheius dolichura* and

Caenorhabditis monodelphis (Supp. Fig. 1). With our DNA staining method, we did not detect any

fragments in *O. tipulae*, most likely because it eliminates 0.6% of its genome only (Supp. Fig. 1) [11,12].

This suggested that we may have underestimated the number of eliminating species using this method.

We next asked if as found in ascarids, *Mesorhabditis belari* and *Oscheius tipulae*, germline

telomeres are also eliminated from somatic cells in our newly identified species. To this end, we performed

DNA-FISH against the conserved telomeric repeat TTAGGC in several species (Fig. 1). We confirmed

germline telomere elimination in a few species only, as the FISH signal was often too noisy to

unambiguously conclude (see Material and Method). However, germline telomere elimination could be

confirmed in *Pelodera teres*, *Rhabditis rainai*, *Oscheius tipulae*, *Auanema freiburgensis* and *Caenorhabditis*

monodelphis. As a control, we showed that germline telomeres are maintained in *C. elegans* somatic nuclei,

as expected from a species that does not undergo PDE (Fig. 1). We also found maintenance of germline

telomeres in *Pelodera strongyloides*, a species for which we did not detect cytoplasmic DNA fragments,

strongly suggesting that this species also does not undergo PDE (Supp. Fig. 1). In some species, following

DNA elimination the telomere signal did not reappear in somatic nuclei, including in *Oscheius tipulae* (Fig.

1), and in *Mesorhabditis belari*, confirming our previous finding [13]. This is consistent with the detection,

from sequencing reads, of shorter neo-telomeres at the new somatic chromosome ends than in the germline

in *Oscheius tipulae* [11]. In contrast, the telomeric signal progressively reappears in the somatic nuclei of

late embryos in *Pelodera teres* and *Rhabditis rainai*, suggesting that somatic telomeres are longer in these

species than in *Mesorhabditis* species. This will need to be confirmed using genomics.

Altogether, our results reveal that at least 17 out of these 25 newly studied Rhabditidae species

undergo PDE, that they most likely all eliminate germline telomeres, and in some cases additionally

eliminate internal parts of chromosomes as well.

Evolution of PDE within Rhabditidae

To identify phylogenetic patterns of PDE presence/absence, we generated a new phylogeny containing the 28

species we studied here (*M. belari*, *M. spiculigera*, *O. tipulae* and 25 new species), using 18S and 28S

ribosomal DNA (rDNA) sequencing data (Fig. 2). Our phylogeny is consistent with previous publications

[18], except on one occasion that we discuss below.

We found that PDE occurs in two large monophyletic groups, both with strong bootstrap support

(Fig. 2 and Supp. Fig. 2). The first clade encompasses the *Mesorhabditis* genus plus three other genera:

Pelodera, Teratorhabditis and *Rhomborhabditis*. All five analysed *Mesorhabditis* species undergo PDE.

However, we found that *Pelodera strongyloides* does not eliminate whereas *Pelodera teres* does. We note

that the phylogenetic placement of *Pelodera strongyloides* is ambiguous in our tree, due to highly divergent

18S and 28S sequences (see Material and Methods). Previous publications also reported weak branch support

for this specific species [18,19]. It thus remains to be determined whether *P. teres* and *P. strongyloides* truly

belong to the same genus, and where exactly *P. strongyloides* branches. At this stage, we do not know if

PDE was present in the common ancestor of this clade and subsequently lost in the lineage leading to *P.*

strongyloides (Fig. 2), or whether it emerged after the divergence from *P. strongyloides*.

The second large monophyletic clade with PDE encompasses the *Oscheius* genus plus seven other

genera (Fig. 2). All ten studied species from this clade undergo PDE, with striking variability in the amount

of eliminated DNA, ranging from very little in *O. tipulae* to massively in *Cruznema sp.* JU4406 (Fig. 1).

During our survey, we also found that DF5015, another putative strain of *Cruznema tripartitum* displayed

much less eliminated DNA than JU4406 (Supp. Fig. 1). While this suggested strong intra-species variability,

crosses between JU4406 and DF5015 produced dead embryos only, likely suggesting they are in fact

different species (see Materials and Methods). Overall, our results suggest that the extent of PDE can vary

over a relatively short evolutionary time scale, i.e. at least within the same genus (Fig. 2).

Outside of these two clades with many tested eliminating species, we also detected PDE in a third

clade with one eliminating species, *Caenorhabditis monodelphis*. In this clade, we found PDE is absent in *C.*

elegans, C. briggsae and *C. angaria*, demonstrating again variability at the genus level. In a recent study,

deeper investigation within this genus confirmed using genomics that PDE occurs in *C. monodelphis*, as

found here, and also identified PDE in two other basal *Caenorhabditis* species, *C. auriculariae* and *C.*

parvicauda, with cytological confirmation in *C. auriculariae* [20]. That study found an absence of PDE in all

other *Caenorhabditis* species - including the three we tested here - leading to the conclusion that PDE was

subsequently lost in the genus *Caenorhabditis* [20].

Finally, we did not detect PDE in four other species, each belonging to a different genus:

Diploscapter sp., Protorhabditis sp., *Choriorhabditis cristata* and *Pristionchus pacificus*. The position of

Pristionchus pacificus is consistent with previous publications placing it basal to all other species studied

here [16,18]. However, the position of *Choriorhabditis cristata* is inconsistent with one previously published

phylogeny [18],that also placed it basally, while here *C. cristata* falls between the *Caenorhabditis* genus and

the clade containing *Oscheius*. Future whole genome-based phylogenies will enable definitively resolving

this inconsistency. .

Irrespective of the placement of *C. cristata*, our results suggest that PDE has been lost and/or gained

at least three times independently within the family Rhabditidae (Fig. 2).

Variation in the timing of PDE

In ascarids, chromosomes start to be fragmented in ABa and ABp cells at the 4-cell stage [5,15,17]. We

found the same early fragmentation starting in ABa and ABp (the two largest and most anterior cells) in

Teratorhabditis palmarum and also in all *Mesorhabditis* species, although embryos already had five

blastomeres when ABa and ABp started dividing in *Mesorhabditis* [13] (Supp. Fig. 1). However, DNA

fragments are visible only when embryos reach ~9 cells for *Pelodera teres* and *Rhomborhabditis regina*.

This delay of approximately one cell cycle for the onset of DNA fragmentation was also observed in all

species of the second large clade and in *C. monodelphis*. *Mesorhabditis*, *Pelodera* and *Rhomborhabditis* are

closer phylogenetically than either is to *C. monodelphis* (Fig. 2). Thus, the timing of PDE initiation varies

between species, but with no obvious ties to the topology of the phylogeny.

PDE occurs regardless of reproductive systems or chromosome number

Last, we analysed PDE in the context of the reproductive system or the chromosome number of the species,

two other parameters that were known [14,21] or that we collected during our analysis (see Materials and

Methods and Supp. Fig. 1). Various reproductive systems are found in Rhabditidae nematodes: regular

sexual reproduction, hermaphroditism, a combination of the two with the presence of three sexes [22], strict

parthenogenesis (full asexuality) [23] or autopseudogamy (pseudo-sexuality) [24]. We found no obvious link

between PDE presence/absence and reproductive systems (Fig. 2). For instance, some sexual species

undergo PDE(e.g. *Pelodera teres)*, while others do not (e.g. *Pelodera strongyloides* and *Caenorhabditis*

angaria). Similarly, PDE is also found in species that reproduce through all reproductive modes:

hermaphroditism (e. g. *Oscheius* species), species with three sexes such as *Auanema freiburgensis*,

or through autopseudogamy such as *Mesorhabditis belari* and *Mesorhabditis simplex,* but is not found in

other hermaphrodites (e.g. *C. elegans*) or asexuals (e.g. *Diploscapter*).

Finally, we found no correlation between the occurrence of PDE and the number of germline

chromosomes (Supp. Fig. 1 and Fig. 2). PDE can occur in species with a large number of germline

chromosomes, such as *Mesorhabditis longespiculosa* (n=17) or a small number, such as *Teratorhabditis*

palmarum (n=4). Conversely, PDE is found neither in *Diploscapter coronatus*, a species with a single pair of

germline chromosomes, nor *Pelodera strongyloides* that has 11. Many species in Rhabditidae have six

chromosomes, some of which undergo PDE, such as *Caenorhabditis monodelphis* or *Oscheius dolichura* and

others that do not, such as *Caenorhabditis elegans* (Fig. 2).

Discussion

Programmed DNA Elimination (PDE) was first described in 1887 in a parasitic nematode [5] and we had

recently uncovered the same process in free-living *Mesorhabditis* species that eliminate up to 30% of their

genome in the soma [13]. In this study, we revealed PDE in 17 other free-living species (out of 25 newly

tested species). This corresponds to 12 out of 17 genera in which at least one species undergoes PDE. We

have discovered that PDE is widespread in Rhabditidae. To confirm this finding, more species in this family

can be tested, beyond our collection. Our work also reveals that this process was missed by early pioneers of

nematode research, even before *C. elegans*- an outsider when considering PDE- was chosen as a model

system [25].

Within Rhabditidae, we found three distinct clades in which PDE is found in at least one species.

Given our current sampling, it is not possible to conclude whether PDE has emerged independently three

times within this family, was present in the common ancestor with independent losses, or a combination of

both. In particular, we have found that the amount of eliminated DNA can vary substantially within a genus,

calling for extending observations to more species per genus. Nevertheless, if we consider PDE in ascarids, it

is tempting to speculate that PDE is an ancient process in nematodes and has been lost several times,

including in the well-studied species *C. elegans* [20]. An interesting parallel is the unicellular ciliates, an

extremely broad phylum in which most studied representatives so far have been found to undergo PDE,

except in genus *Loxodes* [26]; this includes species in genera *Blepharisma*, *Euplotes*, *Oxytricha,*

Tetrahymena and *Paramecium* [6,27]. Here too, the timing and amount of eliminated DNA vary widely. To

what extent PDE is ancestral in ciliates, and/or evolved repeatedly and convergently, remains to be fully

elucidated.

In the case of nematodes, exploration of more species - in particular, outside the Rhabditidae family -

will be required to further explore the history of PDE. Identification of the mechanisms of the molecular

effectors of PDE, none of which are known so far in nematodes, in all these branches will also shed light on

its evolutionary trajectory.

Interestingly, elimination of the germline telomeres is a feature shared by ascarids, *O. tipulae*,

Mesorhabditis. Indeed *O. tipulae* which eliminates very little DNA and does not fragment its chromosomes

internally, eliminates all its germline chromosome telomeres. We confirmed this feature in four other species

and it has also been confirmed for *Caenorhabditis parvicauda* and *Caenorhabditis auriculariae* [20] To date,

no species has been found to eliminate exclusively internal portions while retaining chromosome ends,

indicating that telomere elimination may represent a conserved, ancestral feature.

We found variation in the developmental timing at which chromosome fragments appear in the

cytoplasm. One possibility is that fragmentation begins at the same stage in all species, but that fragments are

subsequently processed or sorted at different times, preventing their cytoplasmic detection. Alternatively,

this variation may reflect species-specific differences in the timing at which the PDE machinery is

transcribed or becomes active.

Within our collection, *C. elegans* appears as one of a minority of species that does not undergo PDE.

Perhaps our understanding of PDE would have been different if the scientific community had picked another

Rhabditidae species as a model species, in the mid-20th century [28]. Nevertheless, our work now reveals

that PDE is found in many free-living species, all of which are potentially lab-tractable. Genetic tools are

already available in some of them, including *Oscheius tipulae* [12,29] and *Auanema* [30,31] , and we have

also established CRISPR and RNAi in some *Mesorhabditis and Pelodera* species (unpublished). This at last

opens avenues for the mechanistic exploration of the PDE in animals.

We originally discovered PDE in *Mesorhabditis* by serendipity when analysing chromosome

inheritance in young embryos [13]. Our study calls for a systematic search for signs of PDE by chromosome

fragmentation, using cytological approaches as described here, when embryos are easy to stain, or using

bioinformatic approaches in any animal species (for example using the tool delfies [32]), because PDE could

be more widespread than anticipated even outside nematodes.

Acknowledgements

Funding: This work has been supported by a grant from the ANR-22-CE12-0027 to MD, and by a grant from the

Ecole Normale Supérieure de Lyon.

Authors contributions: MD designed the experiments. BL established the phylogeny. CL and EW performed the

experimental work and built the figures. BL and MD wrote the manuscript. We acknowledge the contribution of

the imaging plateform PLATIM from SFR Biosciences Lyon. We also thank Garance Fayon, Clara Bourgeay,

Heloise Brun and Basile Bergeron for technical help.

Competing interests: All the authors declare having no competing interests.

Material and methods

Species collection

Strains are maintained at 20°C on NGM plates seeded with *E. coli* OP50, following *C. elegans* protocols, as

described in [33]. Strains are all listed in Supplementary Figure 1. Strains were obtained from the CGC stock

center, or were kind gifts from Marie-Anne Félix or from David Fitch.

Genetic Crosses

DF5015 and JU4406 were both matching *Cruznema tripartitum* on public databases, based on the 18S

sequencing. Because they revealed contrasting amounts of eliminated DNA, we wondered whether they

belong to the same species. To test this, we crossed 3 virgin females from one strain with one male of the

other strain. In both cross directions, females produced dead embryos only. From this result, we concluded

that DF5015 and JU4406 belong to two distinct species, though they are probably closely related as males

and females can still mate.

DNA staining and DNA FISH and immunno-staining

Samples were fixed with methanol and the freeze-cracking method, as described in [13]. We fixed

embryos to analyse DNA fragments or count chromosomes. We also fixed dissected female gonads

in order to count the number of chromosomes in meiotic cells. At this stage, chromosomes are

compact and paired as homologs, facilitating counting. After washing in PBS, the slides were

counter-stained with Hoechst (0.5 µg/ml). For DNA-FISH, we used the following fluorescent oligo

probes at 500 nM (ordered at Sigma) to detect telomeres: 5'

[TAM]TTAGGCTTAGGCTTAGGCTTAGGC 3'. In many species, the telomeric signal

was found inside nuclei, as expected, but also outside nuclei, which we interpreted as noise. This prevented

us from concluding on the presence/absence of telomeres. Images were taken on a Zeiss LSM-800 confocal.

Phylogeny based on 18S and 28S sequences

Sequencing and sequence extraction

To build a phylogeny of the 28 species we studied in family Rhabditidae, we amplified the DNA sequences

of 18S and 28S ribosomal RNA by PCR using universal primers, as described in [18].

In some cases, PCR amplification or sequencing did not work or gave truncated sequences. In these

cases, we turned to public databases to obtain sequences from the same species that we studied by cytology.

In these cases, we used the SILVA database of ribosomal gene sequences, release 138.2, [34], downloading

all SSU (18S) and LSU (28S) sequences for taxon Chromadorea (taxon ID: 119089), a superset of

Rhabditidae. For any given species, when multiple hits were available in SILVA, we used the hit referenced

in [18] in which the authors built a phylogeny of many of the same species of Rhabditidae as studied here,

also using 18S and 28S sequences (plus the large subunit of RNA Polymerase II where available). These hits

can be found in Table S1 of Kiontke *et al.* (2007)[18].

We summarise the origin of each sequence for each species we studied in table 1 below. All sequences are

available on Zenodo: https://doi.org/10.5281/zenodo.18890442

Species	18S origin	18S strain	28S origin	28S strain
Mesorhabditis belari	This study	JU2817	This study	JU2817
Mesorhabditis simplex	This study	JU2864	-	-
Mesorhabditis monhystera	This study	JU2855	-	-
Mesorhabditis longespiculosa	This study	DF5017	This study	DF5017
Mesorhabditis spiculigera	This study	AF72		
Teratorhabditis palmarum	U13937.1	DF5019	This study	DF5019
Pelodera strongyloides	U13932.1	DF5022	-	-

Pelodera teres	AF083002.1	EM437	This study	JU4407
Rhomborhabditis regina	This study	DF5012	This study	DF5012
Oscheius tipulae	KP756939.1	CEW1	EU195969.1	CEW1
Oscheius dolichura	EU196010	JU460	EU195971.1	JU460
Oscheius myriophilus	U81588.1	DF5020	AY602176.1	DF5020
Rhabditis rainai	AF083008.1	PS1191	EU195966.1	DF5091
Auanema freiburgensis	KY680647.1	SB372	KY680646.1	SB372
Rhabditella axei	U13934.1	DF5006	AY602177.1	DF5006
Rhabditis terricola	This study	JU4408	EF417152.1	NK3
Pellioditis typica	U13933.1	DF5026	EU195962.1	DF5025
Phasmarhabditis neopapillosa	This study	JU1104	-	-
Cruznema tripartitum	U73449.1	DF5015	EU195974.1	SB202
Cruznema sp.	This study	JU4406		
Choriorhabditis cristata	EU196013.1	SB300	EU195976.1	SB300
Caenorhabditis elegans	This study	N2	This study	N2
Caenorhabditis briggsae	U13929.1	PB102	AY604481.1	PB800
Caenorhabditis angaria	JN636068.1	PS1010	AY604482.1	PS1010
Caenorhabditis monodelphis	JN636070.1	SB341	AY602170.1	SB341
Diploscapter coronatus	This study	JU4418	BBTA01000078.4838	PDL0010
Protorhabditis sp.	This study	JB122	EU195958.1	JB122
Pristionchus pacificus	AF083010.1	PS312	EU195982.1	PS312
Bunonema reticulatum	EU196017.1	PDL1002	This study	PDL1002

Table 1. **Origin and strain of the 18S and 28S rDNA sequences used to build our Rhabditidae**

phylogeny. For each of our 28 species (rows), the origin columns show whether we used our sequencing,

where successful ('This study'), or if publicly-available data was used, in which case we provide the NCBI

Nucleotide accession. The sequenced strains, of our study, or of the publicly-available data, are also

provided. '-' indicates we did not sequence and no public data was available - this was only the case for 28S

sequences.

We note from Table 1 that, in a number of cases, the publicly-available sequences we used did not come

from the same strain that we studied by cytology (see Supp. Figure 1), and/or the public sequences we used

were not from the same strains between 18S and 28S. However, we checked our resulting phylogeny against

previously-built phylogenies of Rhabditidae, and found our final phylogeny to be consistent with previous

knowledge, except in one instance that is discussed above; validating the use of different strains when the

strain we imaged was not available. In addition, when we used our own sequencing, we consistently found

that our sequencing matched almost identically with publicly-available sequences of the same species (where

available).

Phylogeny building

For each set of gene sequences (18S and 28S), we next produced multiple sequence alignments using mafft

v7.490 [35] (`--adjustdirection --reorder`), and trimmed the alignment with clipkit v2.3.0 [36] to remove

aligned positions with more than 80% gaps (-m gappy -g 0.8). We then built a phylogeny using IQ-TREE

v2.3.6 [37] with auto-detection of the best-fitting substitution model (here, GTR+F+I+R3 for both the 18S

tree and the 28S tree) and 1000 bootstrap replicates (-B 1000). We then used weighted ASTRAL v1.23 [38](-

-treeweights) to combine the two gene trees into a species tree, using a hybrid weighting scheme (--mode 1)

that accounts for both bootstrap support and branch lengths to resolve gene tree discordance, and using 16

rounds of placement and subsampling (-R). We also gave twice as much weight to the 28S tree compared to

the 18S tree, as the 18S tree placed the *Diploscapter*/*Protorhabditis* group incorrectly relative to the

Caenorhabditis group (Supp. Fig. 2).

Phylogenetic trees were visualised and rendered with FigTree (https://github.com/rambaut/figtree/).

The raw Newick-formatted tree files, and code to generate the trees from the input sequences, are available

on Zenodo: https://doi.org/10.5281/zenodo.18890442

 Figure legend

 Figure 1: Somatic chromosome fragmentation and telomere elimination is found in several

 Rhabditidae species.

 A. Embryos from different species and at various stages of development (from the youngest on the left to the

 oldest on the right) were fixed and stained. DNA is shown in blue. The contours of the cells are shown in

 dotted white lines. The DNA of the germline precursor cell is shown in red. Polar bodies, when visible, are

 circled with a solid white line. For the pseudogamous species *Mesorhabditis simplex*, the paternal DNA stay

 highly condensed after fertilization [33] and is shown circled with a solid yellow line. White arrows point

 towards cytoplasmic fragments that become visible from the 5-cell stage in *Mesorhabditis simplex* and later

 in *Cruznema sp.* and *Rhabditella axei*. The name of the strain for each species is shown in parentheses. A

 zoomed-in view of the image of each embryo in the third column is shown in the inset on the right. In the

 inset, the nuclei are encircled in white and the DNA fragments excluded from the nuclei are indicated by an

 arrow. **B.** Embryos from different species are shown and at various stages of development, after DNA FISH

 treatment against telomeric repeats. On the left, embryos are undergoing chromosome fragmentation in a

 least one of their somatic cells. Enlargement of one cell is shown in the inset. DNA is in blue and telomeres

 are in red. In the inset, the nuclei are encircled in white and the telomeres excluded from the nuclei are

 indicated by an arrow. Older embryos, after chromosome fragmentation are shown on the right. In

 Caenorhabditis elegans, all cells in young and old embryos maintain their telomeres, whereas in species

 undergoing PDE, somatic telomeres do not reappear (*Oscheius tipulae, Caenorhabditis monodelphis*) or

 reappear progressively (*Rhabditis rainai* and *Pelodera teres*).

 Figure 2: Phylogenetic distribution of species undergoing Programmed DNA Elimination in family

 Rhabditidae

 Cladogram of species analysed in this study, based on the 18S and 28S rDNA sequencing. Species in which

 we detected PDE are shown in blue and others in orange. PDE has not been scored in *Bunonema reticulatum*,

 which serves here as an outgroup for the phylogeny only. Species undergoing PDE are grouped in two main

 monophyletic groups. We also found that *Caenorhabditis monodelphis* undergoes PDE whereas other

Caenorhabditis species do not. The reproductive modes are shown on the right (asexual, male/female,

hermaphrodite and three sexes), as well as chromosome numbers (also visible on Supp. Fig. 1). *Pelodera*

strongyloides is not monophyletic with *Pelodera teres*, (see discussion in the main text).

Supplementary Figure 1: Cytological results for all 28 analysed species

Embryos from all species analysed in this study are shown. DNA is in blue. The contours of the cells are

shown in dotted white lines. Polar bodies, when visible, are circled with a solid white line. For the

pseudogamous species *M. belari, M. simplex, M. monhystera* [33] *and Protorhabditis sp. (*unpublished*)*, the

paternal DNA stay highly condensed after fertilization [33] and is shown circled with a solid yellow line.

White arrows point towards cytoplasmic fragments. On the left, embryos at the 4 or 5-cell stage are shown.

On the right, are shown embryos at the time of chromosome fragmentation for species undergoing PDE

(shown with a star) or at equivalent stage for species in which we never detected cytoplasmic fragments.

Images used to quantify chromosome number are shown as well. In most cases, we analysed meiotic cells

during diakinesis of prometaphase. At this stage, chromosomes are paired and we therefore count a haploid

number of chromosomes. For some species, it was possible to count chromosome number in 1 or 2 cell

embryos during prometaphase. In this case, we counted a diploid number. Of note, for *Diploscapter*

coronatus the 2 DNA bodies observed in diakinesis correspond to the two unpaired chromosome. This

species as a single chromosome which does not pair during meiosis. For *Oscheius tipulae*, *Auanema sp.*

VSL2220 and *Caenorhabditis monodelphis*, the images come from DNAFISH stainings, revealing the

telomeres in red. This allows validating elimination in these species which eliminate very little DNA. In

Teratorhabditis palmarum, the number of chromosomes, some being very small, is more than 2n=8 in

mitotic cells, revealing that chromosome fragmentation starts at the 4-cell stage (embryos on the left).

Supplementary Figure 2: Phylogeny of studied species using 18S and 28S sequences

Phylogenies of species analysed in this study, based on the 18S rDNA sequencing or 28S rDNA sequencing

References

1. Wang J, Davis RE. Programmed DNA elimination in multicellular organisms. Curr Opin Genet Dev. 2014;27: 26–34. doi:10.1016/j.gde.2014.03.012

2. Dedukh D, Krasikova A. Delete and survive: strategies of programmed genetic material elimination in eukaryotes. Biol Rev Camb Philos Soc. 2022;97: 195–216. doi:10.1111/brv.12796

3. Smith JJ, Timoshevskaya N, Ye C, Holt C, Keinath MC, Parker HJ, et al. The sea lamprey germline genome provides insights into programmed genome rearrangement and vertebrate evolution. Nat Genet. 2018;50: 270–277. doi:10.1038/s41588-017-0036-1

4. Kinsella CM, Ruiz-Ruano FJ, Dion-Côté A-M, Charles AJ, Gossmann TI, Cabrero J, et al. Programmed DNA elimination of germline development genes in songbirds. Nat Commun. 2019;10: 5468. doi:10.1038/s41467-019-13427-4

5. Boveri T. Ueber Differenzierung der Zellkerne wahrend der Furchung des Eies von Ascaris megalocephala. Anat Anz. 1887;2: 688–693.

6. Bétermier M, Klobutcher LA, Orias E. Programmed chromosome fragmentation in ciliated protozoa: multiple means to chromosome ends. Microbiol Mol Biol Rev. 2023;87: e0018422. doi:10.1128/mmbr.00184-22

7. Baudry C, Malinsky S, Restituito M, Kapusta A, Rosa S, Meyer E, et al. PiggyMac, a domesticated piggyBac transposase involved in programmed genome rearrangements in the ciliate Paramecium tetraurelia. Genes Dev. 2009;23: 2478–2483. doi:10.1101/gad.547309

8. Bischerour J, Bhullar S, Denby Wilkes C, Régnier V, Mathy N, Dubois E, et al. Six domesticated PiggyBac transposases together carry out programmed DNA elimination in Paramecium. Elife. 2018;7: e37927. doi:10.7554/eLife.37927

9. Nowacki M, Higgins BP, Maquilan GM, Swart EC, Doak TG, Landweber LF. A functional role for transposases in a large eukaryotic genome. Science. 2009;324: 935–938. doi:10.1126/science.1170023

10. Arnaiz O, Mathy N, Baudry C, Malinsky S, Aury J-M, Denby Wilkes C, et al. The Paramecium germline genome provides a niche for intragenic parasitic DNA: evolutionary dynamics of internal eliminated sequences. PLoS Genet. 2012;8: e1002984. doi:10.1371/journal.pgen.1002984

11. Gonzalez de la Rosa PM, Thomson M, Trivedi U, Tracey A, Tandonnet S, Blaxter M. A telomere-to-telomere assembly of Oscheius tipulae and the evolution of rhabditid nematode chromosomes. G3 (Bethesda). 2021;11: jkaa020. doi:10.1093/g3journal/jkaa020

12. Dockendorff TC, Estrem B, Reed J, Simmons JR, Zadegan SB, Zagoskin MV, et al. The nematode Oscheius tipulae as a genetic model for programmed DNA elimination. Curr Biol. 2022;32: 5083-5098.e6. doi:10.1016/j.cub.2022.10.043

13. Rey C, Launay C, Wenger E, Delattre M. Programmed DNA elimination in Mesorhabditis nematodes. Curr Biol. 2023;33: 3711-3721.e5. doi:10.1016/j.cub.2023.07.058

14. Haag ES, Fitch DHA, Delattre M. From "the Worm" to "the Worms" and Back Again: The Evolutionary Developmental Biology of Nematodes. Genetics. 2018;210: 397–433. doi:10.1534/genetics.118.300243

15. Wang J, Veronezi GMB, Kang Y, Zagoskin M, O'Toole ET, Davis RE. Comprehensive Chromosome End Remodeling during Programmed DNA Elimination. Curr Biol. 2020;30: 3397-3413.e4. doi:10.1016/j.cub.2020.06.058

16.   Picao-Osorio J, Bouleau C, Gonzalez De La Rosa PM, Stevens L, Fekonja N, Blaxter M, et al.
Evolution of developmental bias explains divergent patterns of phenotypic evolution in two nematode
clades. Proc Natl Acad Sci USA. 2025;122: e2507529122. doi:10.1073/pnas.2507529122

17.   Estrem B, Wang J. Programmed DNA elimination in the parasitic nematode Ascaris. PLoS Pathog.
2023;19: e1011087. doi:10.1371/journal.ppat.1011087

18.   Kiontke K, Barrière A, Kolotuev I, Podbilewicz B, Sommer R, Fitch DHA, et al. Trends, Stasis, and
Drift in the Evolution of Nematode Vulva Development. Current Biology. 2007;17: 1925–1937.
doi:10.1016/j.cub.2007.10.061

19.   Sudhaus W. Phylogenetic systematisation and catalogue of paraphyletic "Rhabditidae" (Secernentea,
Nematoda). Journal of Nematode Morphology and Systematics. 2011;14: 113–178.

20.   Stevens L, Sun S, Haruta N, Xiao L, Uwatoko N, Kieninger M, et al. Programmed DNA elimination
was present in the last common ancestor of Caenorhabditis nematodes. bioRxiv; 2025. p.
2025.10.23.681605. doi:10.1101/2025.10.23.681605

21.   Kiontke K. The phylogenetic relationships of Caenorhabditis and other rhabditids. WormBook. 2005
[cited 30 Jan 2016]. doi:10.1895/wormbook.1.11.1

22.   Kanzaki N, Kiontke K, Tanaka R, Hirooka Y, Schwarz A, Müller-Reichert T, et al. Description of two
three-gendered nematode species in the new genus Auanema (Rhabditina) that are models for
reproductive mode evolution. Scientific Reports. 2017;7. doi:10.1038/s41598-017-09871-1

23.   Fradin H, Kiontke K, Zegar C, Gutwein M, Lucas J, Kovtun M, et al. Genome Architecture and
Evolution of a Unichromosomal Asexual Nematode. Curr Biol. 2017;27: 2928-2939.e6.
doi:10.1016/j.cub.2017.08.038

24.   Grosmaire M, Launay C, Siegwald M, Brugière T, Estrada-Virrueta L, Berger D, et al. Males as
somatic investment in a parthenogenetic nematode. Science. 2019;363: 1210–1213.
doi:10.1126/science.aau0099

25.   Nigon V. Modalités de la reproduction et déterminisme du sexe chez quelques nématodes libres.
1949;11: 1–105.

26.   Seah BKB, Singh A, Vetter DE, Emmerich C, Peters M, Soltys V, et al. Nuclear dualism without
extensive DNA elimination in the ciliate Loxodes magnus. Proc Natl Acad Sci U S A. 2024;121:
e2400503121. doi:10.1073/pnas.2400503121

27.   Singh M, Seah BKB, Emmerich C, Singh A, Woehle C, Huettel B, et al. Origins of genome-editing
excisases as illuminated by the somatic genome of the ciliate Blepharisma. Proc Natl Acad Sci U S A.
2023;120: e2213887120. doi:10.1073/pnas.2213887120

28.   Félix M-A. RNA interference in nematodes and the chance that favored Sydney Brenner. J Biol.
2008;7: 34. doi:10.1186/jbiol97

29.   Felix MA. Oscheius tipulae. WormBook. 2006; 1–8. doi:10.1895/wormbook.1.119.1

30.   Yamashita T, Pires-daSilva A, Oomura S, Kusano T, Haruta N, Hasumi M, et al. Microparticle
Bombardment as a Method for Transgenesis in Auanema and Tokorhabditis. MicroPubl Biol.
2025;2025. doi:10.17912/micropub.biology.001585

31.   Adams S, Pathak P, Shao H, Lok JB, Pires-daSilva A. Liposome-based transfection enhances RNAi
and CRISPR-mediated mutagenesis in non-model nematode systems. Sci Rep. 2019;9: 483.
doi:10.1038/s41598-018-37036-1

32.   Letcher B. delfies: a Python package for the detection of DNA breakpoints with neo-telomere addition.
Journal of Open Source Software. 2025;10: 7385. doi:10.21105/joss.07385

33.   Launay C, Félix M-A, Dieng J, Delattre M. Diversification and hybrid incompatibility in auto-
pseudogamous species of Mesorhabditis nematodes. BMC Evol Biol. 2020;20: 105.
doi:10.1186/s12862-020-01665-w

34.   Silva. [cited 21 Aug 2025]. Available: https://www.arb-silva.de/

35.   Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30: 3059–3066.
doi:10.1093/nar/gkf436

36.   Steenwyk JL, Iii TJB, Li Y, Shen X-X, Rokas A. ClipKIT: A multiple sequence alignment trimming
software for accurate phylogenomic inference. PLOS Biology. 2020;18: e3001007.
doi:10.1371/journal.pbio.3001007

37.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective Stochastic
Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol. 2015;32: 268–274.
doi:10.1093/molbev/msu300

38.   Zhang C, Mirarab S. Weighting by Gene Tree Uncertainty Improves Accuracy of Quartet-based
Species Trees. Mol Biol Evol. 2022;39: msac215. doi:10.1093/molbev/msac215

Fig.1

A. DNA staining
Low elimination
Rhabditella axei DF5006

Moderate elimination
Mesorhabditis simplex JU2864

Massive elimination
Cruznema sp. JU4406

B. Telomere DNA FISH

Caenorhabditis elegans N2

Oschieus tipulae CEW1

Caenorhabditis monodelphis JU1667

Pelodera teres JU4407

Rhabditis rainai PS1191

Fig.2