TITLE

Multiple scales of coordination along the body axis during *Drosophila* larval locomotion

AUTHORS / AFFILIATIONS

Marie R Greaney

• Committee on Neurobiology, The University of Chicago, Chicago, IL, United States.

Ellie S Heckscher *

• Department of Molecular Genetics and Cell Biology, The University of Chicago, Chicago, IL, United States.

• Neuroscience Institute, The University of Chicago, Chicago, IL, United States.

Matthew T Kaufman *

• Department of Organismal Biology and Anatomy, The University of Chicago, Chicago, IL, United States.

• Neuroscience Institute, The University of Chicago, Chicago, IL, United States.

• NSF-Simons National Institute for Theory and Mathematics in Biology

 * co-corresponding authors, equal contributors

   heckscher@uchicago.edu

   mattkaufman@uchicago.edu

AUTHOR CONTRIBUTIONS

All authors designed the study; MRG performed the experiments and analyses; MRG wrote the paper with assistance from ESH and MTK; all authors edited the paper; MTK designed the block modeling; ESH provided funding.

CONFLICT OF INTEREST STATEMENT

The authors declare no competing financial interests.

ACKNOWLEDGMENTS

Research reported in this publication was supported by the Pritzker Fellowship in the Neurosciences to MRG; and by the National Institute of Neurological Disorders and Stroke of the National Institutes of Health under award number F31NS118835 to MRG, and R01NS105748 and R56NS134862 to ESH. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

We gratefully acknowledge the assistance of H. Grier in setting up motion tracking and C. Wreden in animal care as well as the University of Chicago's Statistics Consulting Group for helpful discussion of block modeling.

ABSTRACT

Coordinated movement along the body axis is critical to locomotion. In segmented, limbless animals, anterior (head) and posterior (tail) segments play different roles in locomotion, leading to a need for flexible coordination across body regions. Larval *Drosophila melanogaster* present a tractable experimental model for limbless, segmented crawling given the extensive genetic tools available and the optical clarity of the body. Prior work has suggested that, during crawling, all larval body segments contract similarly, despite the fact that each crawl cycle comprises two overlapping phases: an piston involving the most posterior segments, and a peristaltic wave involving all body segments. To test whether coordination varies regionally during locomotion, we expressed GCaMP in all body wall muscles, and recorded segmental contraction kinematics and muscle recruitment during many cycles of locomotion in linear channels. Facilitated by machine vision techniques, we discovered new features of larval crawling at multiple scales. First, the propagation of both contraction and recruitment waves slowed approaching mid-body segments, then sped up towards the head. Second, the timing relationship between contraction and recruitment waves could be highly variable in anterior segments. Third, contraction durations showed particularly strong intersegmental correlations among posterior segments. These data suggest posterior segments coordinately power the piston phase while anterior segments tolerate greater flexibility to enable reorienting behaviors. Our results depict an unanticipated degree of axial heterogeneity in the coordination of limbless crawling, opening new avenues to study the origins of whole body coordination and the consequences of segmental diversity for locomotion.

SIGNIFICANCE STATEMENT

Most animals move through the world by coordinating muscle contractions across the body's many segments. Studying how segments are coordinated in limbless animals can uncover principles that are fundamental for movement across the animal kingdom. However, in one of the best genetic model organisms to address this question, the larval fruit fly, it was unknown how the nearly identical body segments are coordinated across the body. We used machine vision techniques to discover an unanticipated degree of heterogeneity: the timings of muscle activity and movements are more coordinated among tail-end than head-end segments. This provides new insights into animal movements, and opens opportunities to study how interactions of the body and nervous system coordinate segmental movements across an entire animal.

INTRODUCTION

The coordination of various movement features across body regions—timings, amplitudes, rates, and durations of contraction—is critical for effective locomotion. Coordination is well-characterized for certain types of locomotion—anguilliform swimming, vermiform crawling, and inchworm crawling—in certain limbless animals, including leech, earthworm, tadpole, and lamprey, whose bodies are composed of near-identical segments arrayed along the body axis. A general picture emerges that coordination of segmental movements can be highly consistent or can vary cycle-to-cycle, depending on environmental substrate (Gray et al., 1938; Kristan et al., 1974; Kahn et al., 1982; Wallén & Williams, 1984; Stern-Tomlinson et al., 1986).

Larval insects are becoming an important model for studies of behavior, neurophysics, motor control, development, and evolution. Insect larvae possess limbless, segmented bodies. They navigate through their various environments using a behavioral repertoire composed of alternating lateral head bends or turns and linear forward "visceral pistoning" strides (Simon et al., 2010; Green et al., 1983). However, the coordination of segmental movements during visceral pistoning remains poorly characterized.

Visceral pistoning locomotion comprises overlapping piston and wave phases that together constitute a stride cycle (Simon et al., 2010; Heckscher et al., 2012). During the piston phase, the animal's tail end, head end and internal viscera move with near synchrony, translocating the animal's center of mass, while the rest of the external body remains largely stationary. During the wave phase, each segment of the body wall

contracts sequentially in a posterior-to-anterior peristaltic wave, returning the animal's exterior into register with the interior.

Visceral pistoning has been modeled as uniform propagation of a contraction wave through a series of identical and identically coupled units (Gjorgjieva et al., 2013; Pehlevan et al., 2016; Loveless et al., 2019). Kinematic studies that sample a small number of cycles have suggested that body segments' contractions are similar and have consistent timing relationships (Gjorgjieva et al., 2013; Pulver et al., 2015; Sun et al., 2022). However, evidence from dissected preparations reveals substantial differences in the intrinsic properties of segmental circuits along the body axis, particularly between posterior and anterior segments (Jonaitis et al., 2022). These intrinsic differences raise questions about the presumed identical nature of segments' motor outputs, and the strength and spatial scale of segmental coordination.

The field has made major progress in understanding the neural control of visceral pistoning, but despite this, the behavior and how it relates to neuromuscular activity remains incompletely understood. Researchers have delineated neural components of the circuits that subserve visceral pistoning (e.g., Kohsaka et al., 2014; Berni, 2015; Fushiki et al., 2016; Kohsaka et al., 2019) and recorded nervous system activity in dissected preparations (Pulver et al., 2015; Tastekin et al., 2018; Jonaitis et al., 2022). However, the relationships between CNS-driven muscle recruitment and the kinematics of movement are not understood.

This study aimed to determine how segmental contraction and muscle recruitment are coordinated across the Drosophila larval body axis during visceral pistoning locomotion. To do so, aided by machine vision techniques, we recorded >3000

cycles of forward visceral pistoning from larvae expressing calcium indicators in all body wall muscles as they crawled inside linear channels. In visceral pistoning locomotion, unlike other forms of limbless locomotion, segments' contraction rates, durations, and timing relationships changed in nonlinear trends from posterior to anterior. Movement features operating at various scales— segments, adjacent segments, and groups of segments—all varied along the body axis, each with a transition point around the mid-body. Moreover, distinct regions of the larval body axis displayed different degrees of flexibility in converting muscle recruitment to movement. These data support the model that posterior segments coordinately power the visceral piston phase while anterior segments tolerate greater flexibility to enable bending and turning. In short, our findings reveal an unanticipated degree of axial heterogeneity in the coordination of limbless crawling, opening new avenues to study the origins of whole-body coordination and the nature and consequences of segmental diversity.

MATERIALS AND METHODS

Animal care and genotypes

Drosophila melanogaster strains used were the following:

w; +/+; Mef2-GAL4/TM6B-Dfd-YFP (derived from RRID:BDSC_27390)

w; +/+; UAS-jGCaMP7f (RRID:BDSC_79032)

w; +/+; Mef2-GAL4, UAS-Gerry/TM6, Tb, TubP-GAL80 (derived from

RRID:BDSC_27390 and RRID:BDSC_80141)

Drosophila stocks were maintained at 25º C on molasses-based fly food. Fly crosses and embryo/larval collections were maintained on yeasted apple juice agar plates at 25º C, humidified to at least 75%.

<u>Obtaining larvae for imaging</u>

Larvae of genotype *w; +/+; Mef2-GAL4,UAS-Gerry/Mef2-GAL4,UAS-Gerry* ("Gerry") were obtained from in-crosses of same-genotype adults. Larvae were screened for presence of GCaMP in body wall muscles prior to imaging sessions.

Larvae of genotype *w; +/+; Mef2-GAL4/UAS-jGCaMP7f* ("jGC7f") were obtained from adult *w; +/+; UAS-jGCaMP7f* crossed to *w; +/+; Mef2-GAL4/TM6B-Dfd-YFP*. Larvae were screened for presence of GCaMP in body wall muscles prior to imaging sessions.

Embryos were obtained from crosses during a 24-hour collection window, aged for an additional 24 to 48 hours, and larvae to be imaged were subsequently screened and selected by size (average 1mm after 24 hours for first-instar jGC7f larvae; 2.5mm after 48 hours for second-instar Gerry larvae). Larval staging was confirmed by examining larval mouth hooks under magnification to check for characteristic second-instar size and shape. Larvae were not sexed, as sexual characteristics have not yet emerged at these instars. Larvae were handled using soft-bristled paintbrushes.

Imaging larval crawling inside agarose channels

For in-depth explanation of steps, including recipes used, see Greaney and Heckscher, 2024.

<u>Channel preparation</u>

Linear agarose channels were cast with 2% (w/v, in distilled water) LE agarose

(APExBIO, Houston, TX) against raised channel molds (Heckscher et al., 2012;

Greaney and Heckscher, 2024). Channels used for first-instar jGC7f larvae were either

200 μm (n=14 larvae) or 250 μm (n=4 larvae) wide by 200 μm deep; channels used for

second-instar Gerry larvae were approximately 450 μm wide by 450 μm deep. After

agarose fully set, channels were unmolded, trimmed to fit on slides, and kept at 4° C

immersed in water until 15 minutes prior to use. Prior to use the channels were

transferred to room temperature. Larvae were matched to channels of appropriate

widths relative to body size: larvae were snugly enclosed by channel walls and

coverslip, which reduced the speed of crawling (facilitating tracking) and prevented

larvae from turning around inside channels.

Imaging larval locomotion

After screening larvae for expression of the appropriate fluorophore, larvae were

kept on unyeasted apple juice agar plates for the duration of an imaging session

(average 2 hours). We observed no effect of time within session on animals' crawling

speeds, periods, or stride distances, and therefore grouped all animals together for

analysis. Prior to imaging, larvae were washed in distilled water, then placed into an

agarose channel, immersed in distilled water, and covered with a glass coverslip (Figure

1A). Channels were saturated with water throughout the duration of video capture.

Larvae were imaged one at a time for 5-20 minutes. jGC7f larvae were imaged

using a Marianas spinning disk confocal microscope (3i, Denver, CO) with 488nm laser

illumination, at 20 Hz, using laser intensities that prevented range saturation. Gerry

larvae were imaged using a Zeiss LSM 800 point-scanning confocal microscope (Zeiss,

Oberkochen, Germany) with 488nm and 561nm laser illumination, at 13.3 Hz, using laser intensities that prevented saturation in either channel. All larvae were imaged using a 5X, 0.15 NA objective (Zeiss). To keep larvae in the field of view, motorized stage movements were commanded manually throughout the imaging session.

Larvae were allowed to crawl until the end of video capture (12k frames, for jGC7f larvae), until they paused crawling for a period of at least 20 seconds, or until they reached the end of the linear channel, whichever happened first. To acquire additional cycles, some larva were allowed to recover for 1 minute outside of the channels, then were replaced and recorded again. All videos recorded of a single larva were combined for analysis.

Developing DeepLabCut (DLC) segment tracking pipeline

To track larval segment boundaries, the DeepLabCut (DLC) Python package was used (Mathis et al., 2018; RRID:SCR_021391). The following steps resulted in boundary positions over time for segments T3 through A7. Subsequent analysis was performed in Matlab version 2021b (RRID:SCR_001622).

<u>Initial training</u>

Videos were oriented so that the larva's head pointed left. If larvae were rolled relative to the imaging plane, videos were flipped so that the more visible side was the larva's left side.

Separate DLC networks were trained, each using different tracking keypoints, for videos of jGCaMP7f-expressing larvae ("GCaMP network") and videos of Gerry-expressing larvae ("Gerry network").

The GCaMP network tracked two keypoints per segment boundary (Figure 1B), as well as two keypoints each to estimate locations of segments T2 and T1 and two points to estimate the position of a pair of longitudinal tail muscles (dorsal external oblique 1; Campos-Ortega and Hartenstein, 1997). Keypoints were placed at the exterior-most visible point of the left and right sides of a segment boundary. In frames where an animal was rolled substantially to one side, the ventral-anterior corner of 'muscle 4'/'lateral longitudinal 1' was labeled instead.

The Gerry network tracked five keypoints per segment anterior boundary (T3 through A7). Keypoints were placed at the exterior-most visible anterior edges of the left and right sides of each boundary; the central anterior corner of dorsal longitudinal muscles (dorsal oblique 1; Landgraf et al., 1997); and an additional point at the ventral-anterior corner of the left side's dorsal acute 3 (Figure 1C). This network also tracked four keypoints for the anterior boundary of segment A8, two points on the tail segment's dorsal external obliques, and two keypoints to estimate the anterior boundary of segment T2.

Training data included 33 of 35 larvae from the analyzed dataset, as well as data from 61 additional larvae that were not analyzed further due to few overall locomotor cycles, unstable roll angle with respect to the imaging plane, poor manual tracking, or too few segments in view at any given time. 3550 total training frames were used. In both networks, the boundary between segments A8 and A9 was not tracked, and the anterior boundaries of T2 and T1 could not be tracked reliably. Therefore, only segments T3 through A7 were included for subsequent analyses.

Refining DLC network performance

An iterative refinement process was used to add training frames and arrive at the final GCaMP and Gerry DLC networks. First, a DLC network was trained using an initial set of training frames, then used to predict keypoint locations for all videos (either GCaMP or Gerry) in the dataset. Next, prediction quality was evaluated using a multi-step process. To do so, keypoints were converted into segment lengths (see below), then both algorithmic and manual inspection of segment lengths were used to generate a list of potential problem frames. These frames were validated manually. Keypoint locations were corrected for a representative selection of these frames, then the network was retrained with the addition of the corrected frames. This process was repeated until all videos were predicted with few problem frames. Any remaining problem frames were corrected manually.

Converting keypoints to kinematic parameters

<u>Filtering, finding segment boundary positions and lengths</u>

Keypoint positions were smoothed using a 2nd-order Savitzky-Golay filter with a window of 15 frames, then converted into segment boundary positions by taking the mean of all keypoint positions belonging to the same segment boundary as the 'midpoint' for that boundary. Segment lengths were defined as the Euclidean distance between the midpoints of a segment's anterior and posterior boundaries (Figure 1E). A "baseline" length for each segment was defined as the 95th percentile of its length distribution over all frames.

To estimate how far a larva traveled on each wave, segment positions were converted from frame (pixel) coordinates into real spatial coordinates.

For the Gerry videos, frame coordinate positions were converted to real spatial coordinates using recorded stage movement data. For the jGC7f videos, which were captured using software/hardware that did not allow for motorized stage logging, stage movement was estimated from a consensus of pixel velocities in segments that were not actively contracting. Results agreed well with visual inspection.

Finally, segments' boundary positions were extrapolated on frames where they temporarily left the field of view, as follows. First, the expected distance between that boundary and its nearest visible neighbor was defined as the median distance between them in the original dataset (prior to any velocity and position estimation). Then, each time a boundary (typically at the head or tail end of the body) went out of frame, its positions following the out-of-frame period were compared to the position of the adjacent, more central boundary. A single value was added or subtracted to all of the former boundary's positions after the out-of-frame period, such that the median distance between the two boundaries after the out-of-frame period was corrected to match their expected distance. This process was repeated following any block of missing frames. N.B.: extrapolated boundary positions were only used to determine body segment movement distances and cycle stride lengths; they did not affect the calculation of tail movement times, segment lengths, or segment contraction timings.

Segment and tail forward movement times

To define periods of forward movement for each boundary, first, its positions over time (in real spatial coordinates) were converted to velocities. Boundary forward movement periods were defined as times when a segment's frame-to-frame displacement exceeded 1% of its baseline length. Since no baseline length could be

reliably found for the tail, an empirically-determined threshold of 2 μm/frame was used as the tail forward movement threshold. This threshold produced good agreement between manual definition of tail movement times (based on visual inspection of videos) and algorithmically-defined tail movement times.

<u>Segment movement distances</u>

Having established each segment's moving periods, a segment's movement distance on a given cycle was calculated using its anterior boundary. First, the forward movement period was found that overlapped with the segment's contraction during that cycle's locomotor wave (see below). Then, the boundary's median positions before and after the movement period were determined, using all 'not moving' frames since the preceding 'moving' period to determine before-moving position, and all 'not moving' frames until the following 'moving' period to determine the after-moving position. The difference between before-moving and after-moving positions was taken as the movement distance for that cycle.

<u>Segment contraction times and lengths</u>

Segments' lengths traces were processed to determine contraction onset, peak, and offset times as follows.

First, contraction peaks (length minima) were isolated by inverting the lengths traces and applying Matlab's findpeaks function. To avoid finding small length changes that did not correspond to the primary contraction during a wave, a minimum peak prominence was enforced of 20% of the difference between segment's 90th and 10th percentile lengths.

To find contraction onset and offset times, each contraction peak was first isolated from the rest of the lengths trace by finding the length peaks on either side. Then, during this window, the contraction onset was defined as the (interpolated) time when the length decreased to 75% of the difference between maximum length before contraction and the minimum length during contraction. Similarly, the contraction offset was similarly defined as the time when the length increased to 75% of the difference between the minimum length during contraction and the maximum length after contraction (Figure 1F). This method was found to be robust to variation over time in between-cycle segment length.

<u>Forward locomotor waves</u>

Forward locomotor waves were defined as tail-to-head progressions of segmental contractions, as follows. The start of a forward wave was defined as the time at which the tail began moving forward. After this time, a "completed" forward wave occurred if each segment from A7 to T3 contracted sequentially. To allow for the scenario in which neighboring segments began shortening at around the same time, up to five 'out-of-order' frames were permitted in which an anterior segment apparently contracted before its posterior neighbor. No rule was applied that segments had to de-contract in order.

To ensure that only complete, reasonably comparable forward waves were analyzed, waves were excluded if they met any of the following criteria:

- Not all 8 segments (T3-A7) contracted.

- One or more segments contracted more than once.

- The wave preceded or followed a long pause, or a long pause happened partway through the wave. A long pause was defined as 1 second elapsing without any segment boundary moving forward.

- A cycle period could not be defined. (This could happen if the tail was not visible when moving forward, so the cycle start time was missing.)

- The head end (including at least T3 and A1) was retracted, including at the beginning or end of the wave. Head retractions were defined as periods in which one or more contiguous anterior segments (between T3-A3) contracted and de-contracted without simultaneous co-contraction of the next posterior segment.

- After applying earlier exclusions, any remaining wave whose period (z-scored) exceeded 4 standard deviations away from the mean cycle period in either direction was excluded.

<u>Cycle stride distance, period, and speed</u>

Typically, to estimate forward stride distance for each locomotor wave, the visual centroid of a larva is tracked over time (Risse et al., 2017; Thane et al., 2023). To obtain an estimate that depended less on body perimeter estimation, stride distance for each cycle was estimated as the mean of distances moved by segments A2 through A7 on that cycle. Segments A1 and T3 were excluded from the estimation because they occasionally moved as part of a head movement between contractions, resulting in less reliable movement distance calculation for these two segments.

The start of each locomotor wave was defined (above) as the time at which the tail began moving forward in advance of segmental contractions. Cycle periods were defined as the duration from this tail movement until the following cycle's tail movement.

Finally, each cycle was assigned an average speed by dividing the cycle stride distance by the cycle period.

<u>Segment contraction phases</u>

For each cycle, each segment's contraction onset was converted into a cycle phase value. This was done by calculating the delay between the start of the wave and the segment's contraction onset, then normalizing this delay by the cycle period.

Extracting segment fluorescence traces

The calcium sensor GCaMP was used to obtain information about segments' muscle recruitment on each cycle. Larvae expressed the Mef2-GAL4 driver, which drove expression of either jGCaMP7f (Dana et al., 2019) or Gerry (tethered GCaMP6m-mCherry, as in Ugur et al., 2017) in all body wall muscles. However, because of the high likelihood of movement artifacts during crawling, recruitment was only evaluated in Gerry-expressing larvae, in which dynamic calcium sensor (GCaMP) fluorescence could be normalized by stable fluorophore (mCherry) fluorescence.

<u>Region of interest definition</u>

To obtain calcium sensor activity over time (a "fluorescence trace") for each segment, the DLC-tracked keypoints belonging to a segment's anterior and posterior boundaries were connected to define the boundary around a region of interest. Then, all pixels fully contained within the region of interest were averaged. This produced a single fluorescence intensity value per frame, per segment. GCaMP and mCherry fluorescence were processed separately. If a segment was missing any keypoint from either boundary on a given frame, that frame was skipped.

<u>Estimating green-to-red fluorescence ratios</u>

To account for increases in GCaMP fluorescence due to movement or contraction rather than muscle recruitment, segment fluorescence traces were expressed as a change in green-to-red ratio over time ($dGR - GR_0$). To obtain a baseline green-to-red ratio (GR_0), the green-to-red fluorescence ratio (GR) was averaged across all frames during which a segment's length exceeded its baseline length (i.e., the segment was unlikely to be recruited). A segment's muscle recruitment was defined as the difference between GR and GR_0 on each frame.

Fluorescence traces were Savitzky-Golay filtered, using the same kernel applied to segment lengths traces, prior to analysis.

Controlling fluorescence data for viscera movement

For each segment, the middle 50% of pixels were defined by drawing lines passing through the 25% and 75% points of line segments connecting the anterior and posterior external keypoints. Any pixels falling fully or partially between these lines were excluded when calculating green and red fluorescence traces for the segment. Visual inspection confirmed that this removed most or all of visible gut fluorescence.

Matching fluorescence and kinematic data (cycles)

Segments were processed one at a time to obtain the following recruitment metrics and when matching recruitment events into locomotor cycles:

<u>Recruitment event timings and rates</u>

To determine the peak, onset and offset timings and rates of segment recruitment events, fluorescence traces were treated identically to lengths traces, with the exception of a minimum peak prominence of 30% instead of 20%.

<u>Matching recruitment events to cycles</u>

Processing the trace as above led to one onset/offset time, one duration, and one rate value per detected recruitment event. Recruitment events were occasionally missing for a given segment's contraction (due to missing keypoints), and some segments had more than one detected recruitment event on a given locomotor cycle. To find the recruitment events most likely to correspond to a segment's contraction on each cycle, a "best match" algorithm was used that paired a single recruitment event to each cycle. First, recruitment peak timings were compared to their segment's length minimum timings, and each peak was provisionally "assigned" to its closest matching minimum in time. Then, if there were any duplicate assignments (multiple recruitment peaks assigned to the same length minimum), only the nearest match was retained. After this process, the paired recruitment events (peaks, plus their associated onset/offset timings and rate values) were assigned to the same locomotor cycle as the matching contraction.

After matching, quality control steps were applied to discard any probable bad matches. First, if any recruitment onsets or offsets assigned to a given cycle occurred prior to an earlier cycle's length minimum, or later than a later cycle's length minimum, they were discarded. Applying a stronger assumption, if any recruitment onset, specifically, occurred later than the same cycle's segment length minimum, all of that cycle's matched recruitment events were discarded. Finally, if any delay between a segment's recruitment onset and its contraction onset exceeded the median period of a full locomotor cycle, that cycle's recruitment events were discarded. Thus, the remaining matched recruitment events occurred within a reasonable time frame relative to their assigned contraction events.

<u>Recruitment phases</u>

Having assigned recruitment events into cycles, phases were determined for these events using the same procedure as for contraction events: the delay between a cycle's start time (tail movement) and a segment's recruitment onset was normalized to the total cycle period.

Accounting for cycle speed information

To investigate the extent to which intersegmental correlations arose from the relationships between segments' contractions and locomotor speed, speed information was regressed away from segments' kinematic data. To do so, separately for each segment, linear regression was performed to obtain a relationship between cycle speed and each of the segment's contraction kinematic features (contraction amplitude, contraction rate, contraction duration). The predicted value of each feature on each cycle was then subtracted, leaving residual values. Pearson's correlation coefficients were calculated between segments using these residual values in place of the original values.

Modeling block correlation structure

To evaluate the spatial structure of contraction correlations among segments, a method was designed to find "blocks" of contiguous segments that share a correlation strength. This method generates and compares block-based models of segments' correlation strengths for a single feature of contraction kinematics (e.g., contraction amplitude).

<u>Model structure and assumptions</u>

The central assumption of the method is that, within a block, adjacent segment pairs share the same correlation strength for the kinematic feature in question. Then, all further pairwise correlations within the block are modeled as the expected correlation under "local-only" correlations, which decrease exponentially as a function of distance. The correlation strength is determined for each block separately. Outside of blocks, segment pairs share a single baseline correlation strength (see example in Figure 9A).

In this method, blocks potentially range in size from including just one segment to all eight segments. Two blocks may overlap but blocks may not nest fully inside other blocks and a segment may not be contained in more than two blocks. The 8-block model corresponds to a null model in which each segment is uncoupled to any others, except for a constant average correlation strength for all segment pairs. Conversely, the 1-block model (Figure 9B) corresponds to a null model in which correlation between all eight segments is purely dependent on distance (local-only), with no other variation. For all other models, in which subsets of segments group together inside blocks, expected correlation between segments is a function of whether or not they belong to the same block and, if applicable, the correlation strength for that block.

<u>Defining and evaluating models</u>

To fit a model given a set of block edges, the expected pairwise correlations were fit as follows (see Figure 9A).

- Within a block, we averaged the d'th root of all correlations, where d is the distance between the segments in the pair (e.g., for adjacent segments, d=1). To permit this operation, correlations less than zero were set to zero.

- When two blocks overlapped, we performed the same operation but took the d'th root of the correlation as being the average of the values in the two blocks. All within-block values could be estimated simultaneously in this way by linear regression.

- Out-of-block elements, which correspond to segment pairs that were not in the same block, were averaged.

- To prevent fitting for lower-than-expected long-range correlations, any within-block values lower than the out-of-block average value were raised to this value.

To find the most likely block structure for each larva's empirical correlations, first, every possible model was fit (n=1094 possible block combinations). Then, the log likelihood of the data under each model was calculated assuming a Gaussian distribution with a covariance matrix equal to the fitted model. Finally, standard model comparison was performed using the Bayesian Information Criterion (Schwarz, 1978), which penalizes parameters to account for models with more parameters being likely to fit better. Results were similar when the Akaike Information Criterion (Akaike, 1974) was used for model selection instead.

The model as structured has one degeneracy: nearly-uncorrelated segments can be included in the same block with a very low fitted correlation, which model comparison considers more parsimonious than using several 1-segment blocks. To analyze results of block modeling, we therefore only evaluated "strongly-coupled" blocks (R > 0.5 for segments with distance=1). However, the same analyses were performed when instead considering all blocks (Figures S6, S7), producing qualitatively similar results.

Alternative model assumptions

The method as described above assumes that correlations between segments within a block decrease with distance, until they reach a minimum (out-of-block average) value. A variant of the method was tested that assumed correlations did not decrease with distance (Figures S6, S7). In this version of the method, all segment pairs in a block were given the same correlation (the mean correlation strength of all in-block pairs from the empirical data). Out-of-block pairs were averaged as in the model above. Results from this model variant were similar. A second variant was tested in which correlation values within blocks were allowed to decrease with distance to values below the out-of-block average; results were again similar (results not shown).

Experimental Design and Statistical Analysis

The number of cycles required per larva was estimated early in data collection, using the first five jGC7f larvae successfully predicted by a DLC network. For five individual larvae, a threshold of at least 50 "good" forward locomotor cycles was determined by eye to produce reasonable error bounds around pairwise intersegmental correlations. Imaging sessions were designed to maximize the number of cycles that could be well-tracked and predicted per animal, and the minimum threshold of 50 cycles was maintained for the remaining imaged animals (Figure 1D).

To compare distributions of kinematic features across eight body segments in Figure 2, data were combined across larvae. Normality of distributions was rejected using the Shapiro-Wilk test ($p << 0.05$) for all distributions analyzed in Figure 2. Therefore, distributions were compared using the nonparametric Friedman's test and Dunn-Šidak post-hoc multiple comparison correction.

For significance testing of correlation strengths at multiple intersegmental distances (Figure 8A), one-tailed Student's T-tests were used to compare correlation strengths to a null value of zero (Figure 8A, right-tailed test) or to the original, pre-regression correlation strength (Figure 8B, left-tailed test) with a Bonferroni-corrected alpha of 0.002.

To distinguish observed correlations from the null model that correlation strength would depend on intersegmental distance (Figure 9C), a bootstrapping approach was used to generate an expected distribution of correlation strengths under the null model. A separate distribution was generated for each larva. To do so, first, the larva's cycles were sampled with replacement (up to the original number of cycles). Next, pairwise intersegmental correlations were calculated using the resampled data. Then, the expected correlation at each intersegmental distance was calculated by averaging over all pairs of segments at a given distance (7 pairs of segments were averaged with distance=1, 6 pairs with distance=2, etc.). Lastly, the process was repeated (n=10,000 iterations). After generating bootstrapped distributions of expected correlation strength, empirical correlations were compared to the bootstrapped distributions and a p-value calculated for each.

Data and Code Accessibility

Experiments and analyses were not preregistered. Code will be made available at the following GitHub repository: https://github.com/kaufmanlab/larvariability-public. Raw video data and DLC annotations will be made available at the following FigShare repository: *[url to be generated]*

RESULTS

Our goal was to determine how segments are coordinated across the body axis during Drosophila larval visceral pistoning. We asked to what extent segmental coordination matches a model of uniform wave propagation through identical segments. We focused on three kinematic features of segmental movement during crawling — contraction amplitude, contraction rate, and contraction duration – as most directly impacting locomotion speed and therefore potentially needing to be coordinated across segments.

To study visceral pistoning, specifically, we restricted larval behavior: In free crawling on flat surfaces, Drosophila perform a complex behavioral repertoire (Green et al., 1983; Kohsaka et al., 2017; Clark et al., 2018). To limit this repertoire to multiple consecutive cycles of visceral pistoning, we placed larvae inside linear channels (Heckscher et al., 2012; Greaney and Heckscher, 2024) (Figure 1A). To gain a broad understanding, we used various sizes (stages) and genotypes: 18 first-instar larvae expressing jGCaMP7f in all body wall muscles (Figure 1B; Movie 1) and 17 second-instar larvae expressing "Gerry," GCaMP6m and mCherry, in all body wall muscle cells (Figure 1C; Movie 2). Distributions of larval sizes and cycle statistics (speeds, periods, and stride distances) are shown in Figure S1. We obtained an average of 100 forward cycles per animal (see Methods; Figure 1D). From all larvae, we identified segment boundary positions over time (see Methods; Videos S1 and S2), calculated the lengths of each segment from thoracic segment 3 (T3) to abdominal segment 7 (A7) over time (Figure 1E), and assessed the kinematic features of interest for each segment on each cycle (Figure 1F). From Gerry-expressing larvae, we also extracted each segments'

calcium fluorescence over time, a proxy for neuromuscular activity that we compare to kinematics (Figure 1G).

Segment contraction kinematics vary non-monotonically across the body axis

As a first step in testing a model of uniform segmental wave propagation during visceral pistoning, we analyzed the contraction kinematics of individual segments across the body axis. All three kinematic features — contraction amplitude, rate, and duration – varied non-monotonically. For amplitudes, the thoracic segment T3 contracted significantly less than the abdominal segments A1-A7 (Figure 2A). For rates, shortening was slowest in the mid-body (A4 and A5; Figure 2B). For durations, posterior segments contracted for longer than anterior segments in the following order: A5, A4, A6, and A7 (Figure 2C). Thus, segmental contraction waves do not uniformly propagate along the body. Instead, different segmental kinematic features vary in different non-monotonic patterns along the body axis.

Propagation of the segmental contraction wave slows around mid-body

Coordinating the relative timings of specific movements is critical for behavior. In the case of visceral pistoning, the peristaltic wave depends on an ordered sequence of segmental contractions. The relative time at which each segment contracts can be expressed as a fraction of the total cycle period (i.e., its contraction phase, Figure 2D). The phase relationships between segments' contractions may be linear, as in the uniformly propagating contraction wave during some types of swimming (Hill et al., 2003; Di Santo et al., 2021), or more variable, as in some types of terrestrial crawling

(Stern-Tomlinson et al., 1986; Cacciatore et al. 2000, Trimmer and Issberner, 2007).

Therefore, we asked whether *timing coordination* was uniform or variable along the body axis during visceral pistoning. We quantified the intersegmental phase delays between adjacent segments' contractions (Figure 2D). Phase delays varied along the body axis (Figure 2E), and were slowest around mid-body (A4-A3; Figure 2E). Accordingly, segments' contraction phases were distributed nonlinearly (Figure 2F); overall, a sigmoidal model was a better fit than a linear model for segmental contraction phase distributions, both for the combined data and for a vast majority of individual larvae. Thus, the segmental timing relationships for visceral positioning in channels feature a slowing of the contraction wave around mid-body.

Mid-body contraction wave slowing was not explained by tail movement or channel interactions

The axial variation in contraction wave propagation (Figure 2) could arise from several factors, including differences across segments in neural circuit activity and/or in conversion of muscle activation into forces via body-environment interactions (see, e.g., Peron et al., 2009; Srinivasan et al., 2012; Jonaitis et al., 2022; Kohsaka, 2023). Particularly because we constrained larvae inside channels as part of our approach, we wanted to evaluate the likelihood that physical forces such as body-channel interactions could explain the differences in wave propagation.Therefore, we performed two analyses to test for the influence of physical forces.

First, we tested whether differences in contraction kinematics could be explained by the forward movement of the tail. In forward visceral pistoning, the tail segments

contract, pushing the viscera and center of mass forwards (Simon et al., 2010; Heckscher et al., 2012). We considered the possibility that waves with farther tail movement might slow at a more anterior point than waves with less tail movement. To test this, we split each larva's cycles at the median into cycles with longer vs. shorter tail movement distances. We did not observe different axial trends for either contraction rates or durations (Figure 3), arguing that the location of mid-body slowing was not a consequence of tail displacement.

A second possibility was that the axial differences in segmental coordination might directly result from physical interactions between the sides of the larva and walls of the channel (Figure 4A). Because the sizes of the larvae in our dataset varied, correlated with larval stage (Figure S1), we analyzed the first-instar and second-instar larvae separately. Our second-instar but not first-instar larvae appeared increasingly impeded by the channels at larger body sizes, leading to longer cycle periods (Figure 4B-C). Regardless of size and stage, we observed similar axial trends in the distributions of contraction amplitudes, rates, durations, and intersegmental phase delays (Figure 4D). We conclude the structure of these distributions was unlikely to be a consequence of segments differing in their physical interactions with the channels.

Trends in the recruitment wave led similar trends in the contraction wave by one segment to the anterior

Since we could not explain the mid-body slowing of the contraction wave using proxies for physical forces, we next asked whether the non-uniform wave could be neural in origin. If so, similar non-monotonic axial trends would also be present in

segments' muscle activations. To assess segmental trends in neuromuscular activity, we quantified a proxy of muscle recruitment: segment calcium activity in the Gerry-expressing larvae. We measured each segment's recruitment amplitude, rate, and duration, which were closely analogous to the kinematic features.

First, we compared axial trends for recruitment and contraction features. Overall, trends for recruitment features were similar to analogous contraction features (compare Figure 2A-C to Figure 5A-C). Further, the recruitment wave, like the contraction wave, slowed around mid-body (compare Figure 2E-F to Figure 5D-E). A sigmoidal model was a better fit than a linear model for recruitment phase distributions, both for the combined data and most individual larvae.

The trends in the recruitment wave led similar trends in the contraction wave by one segment to the anterior. A spatial offset thus emerged between recruitment and contraction features: the slowest recruitment rates, longest durations, and longest phase delays each occurred one segment anterior relative to the slowest or longest analogous contraction feature.

Unlike the contraction wave, for the recruitment wave there was a striking increase in variance in wave propagation. This manifested particularly in the phase delays between A3 and A2 (Figure 5D) and as large cycle-to-cycle phase delay variability within individual larvae (example in Figure 5F). We conclude that, in addition to slowing around mid-body, recruitment wave propagation can progress differently on each crawl cycle, and the point of greatest variation in its progression appears to be in the anterior mid-body.

The relationship between recruitment and contraction waves becomes variable at mid-body

Because, generally, we found high variance in the recruitment wave at mid-body (Figure 5D-F), we next investigated the timing relationships between the recruitment and contraction waves in single animals on a cycle-by-cycle basis. The relationship between a segment's recruitment and its contraction can inform the role of muscle activations in locomotion. For instance, in anguilliform swimming, the phase delay between muscle recruitment and segment contraction changes systematically along the body axis, supporting different roles of muscle activity at the tail end and head end (Williams et al., 1989; Wardle et al., 1995; McMillen et al., 2008). Therefore, we directly asked whether any such systematic differences existed in larval crawling.

For each segment and each cycle, we calculated the phase delays between recruitment and contraction (Figure 6A-C). In most individual larvae, in at least one segment, the difference between recruitment and contraction phases was much larger and more variable than for other segments (example: Figure 6A-B, segments A2, A3). Overall, the propagation of the recruitment wave was furthest in advance of the contraction wave at segment A3 (Figure 6C). Moreover, the phase delays between recruitment and contraction, while consistent among posterior segments, became substantially more variable toward the head end, both within- and across larvae (Figure 6C-D), an effect that did not appear to depend on animal size (Figure S2: larvae sorted by size).

Interestingly, we observed relatively frequent negative phase delays between T3's recruitment and contraction, indicating that shortening began before muscles had

been substantially activated, possibly due to pushing from behind (Figure 6C, Figure S2).

One possible confound was the variable movement of viscera during visceral pistoning; because the viscera were visible in the fluorescence channels, this movement might have affected apparent segmental recruitment timings. To control for this possibility, we repeated this analysis while excluding the central 50% of each segment's pixels. Results were unchanged (Figure 6E-F), arguing that visceral movement was not the source of either the phase delay or the variability.

In sum, delays in propagating recruitment and contraction between segments did not reliably align, leading to variable phase relationships between the waves beginning in the mid-body.

Segments' contractions are correlated over long ranges, only partially due to locomotor speed

Thus far, we have considered kinematic features on a segment-by-segment basis (Figures 2A-C, 5A-C) and their short-range adjacent pair relationships (Figures 2D-F, 5D-F). Short range coordination might explain contraction kinematics during certain types of locomotion. However, a unique feature of visceral pistoning compared to other forms of locomotion is that head and tail movements are highly correlated as they are physically coupled to each other via the viscera and the fluid-filled body interior (Heckscher et al., 2012). Thus, for visceral pistoning locomotion, we must consider longer-range coordination. Below, we evaluated correlations between segments' kinematic features over distance.

We determined which contraction features were significantly correlated and estimated the ranges, in intersegmental distance, over which correlations were observed. We found an average correlation strength for each feature at each intersegmental distance. For instance, at an intersegmental distance of three, the correlation was calculated by averaging values for the following segmental pairs: T3-A2, A1-A3, A2-A4, A3-A5, A4-A6, and A5-A7; whereas an intersegmental distance of eight was calculated using only the T3-A7 segmental pair (Figure 7A). At almost all intersegmental distances, all contraction features were significantly correlated (Pearson's R > 0; one-tailed T-test; p < (0.05/21)). Contraction durations were the most strongly correlated (Figure 7A), with some correlations being quite strong (R > 0.5). We conclude that during visceral pistoning, segmental contractions are correlated over long ranges.

A somewhat trivial explanation for long-range correlations could be that they result from changes related to whole animal locomotor speed. To account for this possibility, we generated a best-fit linear model for the relationship between speed and segments' cycle-by-cycle contraction kinematics (Methods), subtracted this prediction from the data, and recalculated correlations using the residuals. Contraction amplitude correlations remained almost independent of speed (Figure 7B). For contraction rates and durations, correlation strengths were significantly diminished but did not disappear at intersegmental distances beyond adjacent segments. This suggests that speed-related factors play a role in correlating certain features across the body, but cannot fully explain longer-distance correlations.

***Contraction and recruitment durations are strongly correlated among posterior
segments***

The above analysis revealed the existence of long-range coordination of
segmental contractions (Figure 7), but did not localize correlations to a particular region
of the body. Given that both segmental contraction and muscle recruitment varied
across the body with a prominent mid-body transition, we asked whether long-range
coordination also varied across the body, and if so, how (Figure 8A). In visceral
pistoning locomotion, all segmental contractions move the exterior body wall; however,
contraction of the posterior segments additionally pushes the interior viscera forward.
The strength of correlations across groups of segments may reflect how tightly different
body regions must be coordinated for behavior. Therefore, we wondered if coordination
might differ among anterior or posterior groups of segments with different roles in
behavior.

We performed the following analysis both with and without speed-corrected
correlations, with similar outcomes (Figures S3, S4). First, for each kinematic feature—
contraction amplitude, rate, and duration—we generated a correlation matrix with
segments on each axis (Figure 8B; Figure S3A-C). Of note, segment T3 tended to be
less correlated with all other segments, pointing to potential thoracic-abdominal
differences in coordination. Most strikingly, for contraction durations, correlations were
much stronger in the posterior compared to the anterior (Figure 8B). We compared to a
bootstrapped null model (Methods), finding correlations in segmental contraction
duration were consistently higher than expected toward the tail end of the animal, at
ranges spanning multiple segments (Figure 8C, right). Next, we repeated these

analyses for muscle recruitment features. Generally, the correlations for muscle recruitment features were stronger than for analogous kinematic features (Figure S5), suggesting stronger coordination of segmental neuromuscular activity compared to segmental contractions.

Thus, regardless of speed, contraction and muscle recruitment durations are coordinated more strongly and at longer ranges among posterior segments.

A new statistical model of segment interactions

Above, we observed that non-adjacent pairs of segments exhibited correlated contraction durations, and further that these correlations were more commonly elevated in the posterior segments (Figure 8). We hypothesized that there might be a strongly-coordinated posterior block of segments, whose contraction durations are adjusted as a unit from cycle to cycle, but weaker coordination among anterior segments, whose contraction durations could be more independently modulated.

To explicitly test the idea that groups of segments behaved as blocks during visceral pistoning, we developed a new statistical method. This method finds groups of segments with similar correlation strengths and identifies where along the body axis correlation strengths change. It compares actual data to all possible models in a model family, wherein segments are grouped into different numbers and locations of blocks of shared correlation strengths (Fig. 9A; Methods). The simplest model groups all segments in a single large block, which means correlation strengths do not change along the body axis (Figure 9B). More complex models group segments into two or more blocks, indicating that correlations change somewhere along the body axis. The

best-fitting model was identified for each larva (Figure 9C); often, the best fit model had two strongly-correlated blocks of segments (Figure 9D). We thresholded the best-fitting model for each larva to retain blocks of segments with strong correlations (R > 0.5), and identified the segmental location of the blocks' edges.

Block edges were infrequent between posterior segments A4-A7 (Figure 9E), suggesting that correlation strengths rarely changed between these segments. To check this more directly, we asked for how many larvae did the best-fitting models combine each pair of segments into the same block (Figure 9F). Posterior segments A4 through A7 were included in the same block with the other posterior segments for almost every larva (Figure 9F). Anterior segments, with the exception of A1 and A2, less often appeared together in the same block. The same observations held with and without accounting for speed or using variants of the model assumptions (Figure S6). In contrast to contraction durations, modeling blocks for contraction amplitude or rate correlations did not reveal consistent groups of segments (Figure S7). We repeated this analysis for muscle recruitment durations. The block results for recruitment duration correlations were strongly similar to those for contraction duration correlations (Figure 9G-I). Thus, generally, posterior segments comprised a coordinated block, and less often anterior segments also comprised a coordinated block.

DISCUSSION

We determined how segments contract and how muscles are recruited during larval Drosophila visceral pistoning inside linear channels, finding a previously unknown diversity in coordination across the body. First, segment-by-segment, muscle recruitment and contraction kinematics had their longest durations around the mid-body. Second, for adjacent pairs of segments, the propagation of contraction and recruitment waves slowed around the mid-body. Also around mid-body, the recruitment wave's propagation became quite variable, and the phase relationships between contraction and recruitment waves became inconsistent. Lastly, for a group of posterior segments, contraction and recruitment durations were strongly correlated. In summary, we discovered that features of visceral pistoning operating at various scales—segments, adjacent segments, and groups of segments—all vary along the body axis, each with a transition point around the mid-body.

For visceral pistoning locomotion, mid-body variation was unexpected. Compared to our study, previous studies of larval locomotion on flat surfaces sampled fewer cycles of movement (Gjorgjieva et al., 2013; Pulver et al., 2015; Sun et al., 2022; Liu et al., 2023), and did not find differences in segments' contraction durations or amplitudes. Thus, the prevailing model has been uniform propagation of contraction waves along the body at all speeds (Gjorgjieva et al., 2013; Pehlevan et al., 2016; Liu et al., 2023). However, differences between the intrinsic properties, inputs, and input responses of anterior and posterior body segments have been described (Srinivasan et al., 2012; Takagi et al., 2017; Tastekin et al., 2018; Murawski et al., 2020; Jonaitis et al.,

2022; Jonaitis et al., 2024), motivating us to reevaluate the prevailing model for visceral pistoning locomotion.

Our study used linear channels to enable the collection of large numbers of similar visceral pistoning cycles. Consequently, we did not study other features of free behavior such as head sweeps or turns (Green et al., 1983; Lahiri et al., 2011; Berni et al., 2012; Gershow et al., 2012; Gomez-Marin and Louis, 2014; Wystrach et al., 2016). Notably, channels versus flat surfaces provide different sensory environments, which might encourage differences in segmental coordination (Vaadia et al., 2019, Hughes and Thomas, 2007). However, we did not find evidence that physical forces due to the channel environment—assessed using forward travel distance or size of the larva relative to channels— related to axial differences (Figures 3, 4). Thus, it seems unlikely that the channel environment per se explains the observed structure of coordination. Imaging technologies are emerging that enable simultaneous recording of neural/ muscular recruitment and segmental kinematics while larvae move in unconstrained environments (Karagyozov et al., 2018; Bonnard et al., 2022; McNulty et al., 2024). These advances will permit testing how the segmental coordination in channels changes when larvae navigate and perform more complex behaviors in different environments and will contribute insight into the role of sensory feedback in diverse contexts.

Because our setup allowed simultaneous imaging of muscle contraction and muscle calcium activity across large numbers of locomotor cycles, we could compare coordination of segmental recruitment to coordination of contraction kinematics. Generally, trends in the recruitment wave led similar trends in the contraction wave by

one segment to the anterior (Figure 5). In *Manduca sexta* caterpillars, a limbed model of

visceral pistoning locomotion (Simon et al., 2010), anterior segments' muscle activations

interact with the substrate to produce tension that pulls posterior segments forward (Lin

and Trimmer, 2010). Our data suggest that similarly, in limbless visceral pistoning

locomotion, posterior body segments are pulled forward by muscles in anterior body

segments. A notable exception is that T3 segments can shorten before muscle

recruitment (Figure 6C, Figure S2), possibly due to pushing from behind.

The timing relationship between the muscle recruitment and contraction waves

varied along the body, with a tight relationship near the tail end that became quite

variable starting in mid-body segments (Figure 6). The variability in muscle recruitment

could signal that anterior segments are sensitive to sensory feedback about mechanical

or internal state and adjust the tension generated in advance of the contraction wave.

Alternatively, the variability in muscle recruitment may be intrinsic to central circuits:

recent work in fictive nerve cord preparations showed variable propagation of neural

activity through anterior segments, even in the absence of sensory feedback (Francis et

al., 2024). The variable delay between muscle recruitment and segment contraction in

anterior segments demonstrates that converting neuromuscular activity to segment

shortening is not a segment-autonomous process, and other factors contribute to the

conversion. Such factors could include features of the channel environment (e.g.,

interaction with the channel walls), or dynamic (e.g., neuromuscular) or static (e.g.,

tissue mechanics) properties of the larval body, or a combination. To test these

possibilities in future, it will be essential to determine the mechanical forces that relate

larval muscle activity to movement in a specific environment (Tytell et al., 2011; Ormerod et al., 2022; Sun et al., 2022; Kohsaka, 2023; Booth et al., 2024).

Additionally, we found differences in the strength of correlations for both contraction and recruitment durations among posterior and anterior segments. During visceral pistoning, we observed stronger correlations among a block of posterior segments, even when accounting for cycle speed (Figures 8, 9). In general, stronger correlations among groups of segments point to areas of the body that might need to be coordinated more tightly for behavior. An attractive possibility is that the stronger posterior correlations may be due to differing behavioral roles for posterior and anterior segments. In the most posterior segments, contraction has a dual function—to move the exterior body wall and to push the internal viscera. The dual function of the posterior may require a tightly coordinated contraction; indeed, coordinated posterior activity, or inhibition of activity, has been noted even in the absence of locomotor waves (Pulver et al., 2015; Tastekin et al., 2018; Jonaitis et al., 2022). During visceral pistoning, anterior segments primarily move the exterior body wall, and can be recruited between waves for lateral head movements (Lahiri et al., 2011; Berni et al., 2012; Gomez-Marin and Louis, 2014; Pulver et al., 2015; Wystrach et al., 2016). Therefore, the larva's anterior segments may benefit from greater flexibility to respond to sensory inputs, enabling reorienting maneuvers.

What is the contribution of development to axial differences in locomotor coordination? Some axial trends could be explained by known developmental factors. For example, thoracic and abdominal regions differ in their Hox gene expression, which leads to anatomical, neural, and behavioral differences (Roberts, 1971; Campos-Ortega

and Hartenstein, 1997; Dixit et al., 2007; Tsuji et al., 2008; Heckscher et al., 2014; Rotheray, 2019; Jonaitis et al., 2022; Diao et al., 2024; Vasudevan et al., 2024). Indeed, compared to abdominal segments, thoracic segment T3 contracted less tightly (Figure 2) and was less correlated with abdominal segments (Figure 8, Figure S5), and could shorten before muscles' recruitment (Figure 6C, Figure S2), any of which could be due to differences in neural circuits. Therefore, an attractive hypothesis posits segments with differing Hox codes contain diverse neuromuscular circuits and thus differ in their coordination. However, our data suggest that a differential Hox code is only part of the explanation for differing axial coordination, because axial trends within the abdomen are not well-correlated with known Hox expression borders. Specifically, we did not find evidence that segments A7-A5, which share a Hox code that differs from the Hox code of other segments (Singh et al., 2014), were more self-similar than other segments in terms of muscle recruitment or kinematics (Figures 2, 5). Second, although the transitions in contraction and recruitment features occurs in the mid-body, the specific segment varies, further suggesting a role for factors beyond those that are hardwired early in development.

Irrespective of developmental boundaries, within the abdomen, a variety of segmental boundaries have been proposed as decision or reversal points with respect to locomotor waves and behavioral decision-making (e.g., Berni, 2015; Takagi et al., 2017; Tastekin et al., 2018; Murawski et al., 2020; Jonaitis et al., 2022). Our results suggest that, in addition to producing different behavioral responses to stimuli, intersegmental differences affect the coordination and propagation of locomotor waves.

Thus, it becomes of greater interest to understand the extent to which locomotor circuits differ along the body axis and how these circuit differences support axial coordination.

The need for an accurate model of visceral pistoning is at least three-fold: It contributes to the field's understanding of embodied behavior; it is a required precursor for dissecting underlying neural mechanisms; and it motivates further investigation into the development of axially diverse motor circuits. In summary, our findings of complex variation in how body segments are recruited, contracted, and correlated across the body axis, along with varying relationships between muscle recruitment and kinematics, make visceral pistoning locomotion in Drosophila larvae a rich system for interrogating locomotor control. Combined with the power of Drosophila genetics, this model unlocks new possibilities for probing the scales, mechanisms, and behavioral outcomes of whole-body coordination in segmented nervous systems.

REFERENCES

Akaike H (1974) A new look at the statistical model identification. IEEE Trans Automat
    Contr 19:716–723 Available at: http://dx.doi.org/10.1109/TAC.1974.1100705.

Berni J (2015) Genetic Dissection of a Regionally Differentiated Network for Exploratory
    Behavior in Drosophila Larvae. Current Biology 25:1319–1326 Available at: http://
    dx.doi.org/10.1016/j.cub.2015.03.023.

Berni J, Pulver SR, Griffith LC, Bate M (2012) Autonomous Circuitry for Substrate
    Exploration in Freely Moving Drosophila Larvae. Current Biology 22:1861–1870
    Available at: http://dx.doi.org/10.1016/j.cub.2012.07.048.

Booth JH, Meek AT, Kronenberg NM, Pulver SR, Gather MC (2024) Optical mapping of
    ground reaction force dynamics in freely behaving Drosophila melanogaster
    larvae. eLife 12 Available at: http://dx.doi.org/10.7554/elife.87746.3.

Cacciatore TW, Rozenshteyn R, Kristan WB Jr (2000) Kinematics and Modeling of
    Leech Crawling: Evidence for an Oscillatory Behavior Produced by Propagating
    Waves of Excitation. J Neurosci 20:1643–1655 Available at: http://dx.doi.org/
    10.1523/jneurosci.20-04-01643.2000.

Campos-Ortega JA, Hartenstein V (1997) The Embryonic Development of Drosophila
    melanogaster. Springer Berlin Heidelberg. Available at: http://dx.doi.org/
    10.1007/978-3-662-22489-2.

Clark MQ, Zarin AA, Carreira-Rosario A, Doe CQ (2018) Neural circuits driving larval
    locomotion in Drosophila. Neural Dev 13 Available at: http://dx.doi.org/10.1186/
    s13064-018-0103-z.

Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R, Macklin JJ, Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS (2019) High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. Nat Methods 16:649–657 Available at: http://dx.doi.org/10.1038/s41592-019-0435-6.

Di Santo V, Goerig E, Wainwright DK, Akanyeti O, Liao JC, Castro-Santos T, Lauder GV (2021) Convergence of undulatory swimming kinematics across a diversity of fishes. Proc Natl Acad Sci USA 118 Available at: http://dx.doi.org/10.1073/pnas.2113206118.

Diao F, Vasudevan D, Heckscher ES, White BH (2024) Hox gene–specific cellular targeting using split intein Trojan exons. Proc Natl Acad Sci USA 121 Available at: http://dx.doi.org/10.1073/pnas.2317083121.

Dixit R, VijayRaghavan K, Bate M (2007) Hox genes and the regulation of movement in Drosophila. Developmental Neurobiology 68:309–316 Available at: http://dx.doi.org/10.1002/dneu.20589.

Dorgan KM (2010) Environmental Constraints on the Mechanics of Crawling and Burrowing Using Hydrostatic Skeletons. Exp Mech 50:1373–1381 Available at: http://dx.doi.org/10.1007/s11340-010-9399-2.

Francis J, Gibeily CR, Smith WV, Petropoulos IS, Anderson M, Heitler WJ, Prinz AA, Pulver SR (2024) Generation of motor program diversity and variability through inhibitory circuit motifs in the Drosophila larval locomotor system. bioRxiv 610491. Available at: http://dx.doi.org/10.1101/2024.09.02.610491.

Fushiki A, Zwart MF, Kohsaka H, Fetter RD, Cardona A, Nose A (2016) A circuit

mechanism for the propagation of waves of muscle contraction in Drosophila.

eLife 5 Available at: http://dx.doi.org/10.7554/eLife.13253.

Gjorgjieva J, Berni J, Evers JF, Eglen SJ (2013) Neural circuits for peristaltic wave

propagation in crawling Drosophila larvae: analysis and modeling. Front Comput

Neurosci 7 Available at: http://dx.doi.org/10.3389/fncom.2013.00024.

Gomez-Marin A, Louis M (2014) Multilevel control of run orientation in Drosophila larval

chemotaxis. Front Behav Neurosci 8 Available at: http://dx.doi.org/10.3389/

fnbeh.2014.00038.

Gray J, Lissmann HW, Pumphrey RJ (1938) The Mechanism of Locomotion in the

Leech (Hirudo Medicinalis Ray). Journal of Experimental Biology 15:408–430

Available at: http://dx.doi.org/10.1242/jeb.15.3.408.

Greaney MR, Heckscher ES (2024) Studying Drosophila Larval Behavior in Agarose

Channels. Cold Spring Harb Protoc Available at: http://dx.doi.org/10.1101/

pdb.prot108420.

Green CH, Burnet B, Connolly KJ (1983) Organization and patterns of inter- and

intraspecific variation in the behaviour of Drosophila larvae. Animal Behaviour

31:282–291 Available at: http://dx.doi.org/10.1016/s0003-3472(83)80198-5.

Heckscher ES, Lockery SR, Doe CQ (2012) Characterization ofDrosophilaLarval

Crawling at the Level of Organism, Segment, and Somatic Body Wall

Musculature. J Neurosci 32:12460–12471 Available at: http://dx.doi.org/10.1523/

jneurosci.0222-12.2012.

Heckscher ES, Long F, Layden MJ, Chuang CH, Manning L, Richart J, Pearson JC, Crews ST, Peng H, Myers E, Doe CQ (2014) Atlas-builder software and the eNeuro atlas: resources for developmental biology and neuroscience. Development 141(12):2524-32 Available at: http://dx.doi.org/10.1242/dev.108720.

Hill AAV, Masino MA, Calabrese RL (2003) Intersegmental Coordination of Rhythmic Motor Patterns. Journal of Neurophysiology 90:531–538 Available at: http://dx.doi.org/10.1152/jn.00338.2003.

Hughes CL, Thomas JB (2007) A sensory feedback circuit coordinates muscle activity in Drosophila. Molecular and Cellular Neuroscience 35:383–396 Available at: http://dx.doi.org/10.1016/j.mcn.2007.04.001.

Hunter I, Coulson B, Zarin AA, Baines RA (2021) The Drosophila Larval Locomotor Circuit Provides a Model to Understand Neural Circuit Development and Function. Front Neural Circuits 15 Available at: http://dx.doi.org/10.3389/fncir.2021.684969.

Jonaitis J, MacLeod J, Pulver SR (2022) Localization of muscarinic acetylcholine receptor-dependent rhythm-generating modules in the Drosophila larval locomotor network. Journal of Neurophysiology 127:1098–1116 Available at: http://dx.doi.org/10.1152/jn.00106.2021.

Jonaitis J, Hibbard KL, Layte KM, Hiramoto A, Cardona A, Truman JW, Nose A, Zwart MF, Pulver SR (2024) Steering From the Rear: Coordination of Central Pattern Generators Underlying Navigation by Ascending Interneurons. bioRxiv 598162. Available at: http://dx.doi.org/10.1101/2024.06.17.598162.

Kahn JA, Roberts A, Kashin SM (1982) The neuromuscular basis of swimming movements in embryos of the amphibian Xenopus Laevis. Journal of Experimental Biology 99:175–184 Available at: http://dx.doi.org/10.1242/jeb.99.1.175.

Karagyozov D, Mihovilovic Skanata M, Lesar A, Gershow M (2018) Recording Neural Activity in Unrestrained Animals with Three-Dimensional Tracking Two-Photon Microscopy. Cell Reports 25:1371-1383.e10 Available at: http://dx.doi.org/10.1016/j.celrep.2018.10.013.

Kohsaka H (2023) Linking neural circuits to the mechanics of animal behavior in Drosophila larval locomotion. Front Neural Circuits 17 Available at: http://dx.doi.org/10.3389/fncir.2023.1175899.

Kohsaka H, Takasu E, Morimoto T, Nose A (2014) A Group of Segmental Premotor Interneurons Regulates the Speed of Axial Locomotion in Drosophila Larvae. Current Biology 24:2632–2642 Available at: http://dx.doi.org/10.1016/j.cub.2014.09.026.

Kohsaka H, Guertin PA, Nose A (2017) Neural Circuits Underlying Fly Larval Locomotion. CPD 23:1722–1733 Available at: http://dx.doi.org/10.2174/1381612822666161208120835.

Kohsaka H, Zwart MF, Fushiki A, Fetter RD, Truman JW, Cardona A, Nose A (2019) Regulation of forward and backward locomotion through intersegmental feedback circuits in Drosophila larvae. Nat Commun 10 Available at: http://dx.doi.org/10.1038/s41467-019-10695-y.

Kristan WB Jr, Stent GS, Ort CA (1974) Neuronal control of swimming in the medicinal

   leech. J Comp Physiol 94:97–119 Available at: http://dx.doi.org/10.1007/

   BF00617837.

Lahiri S, Shen K, Klein M, Tang A, Kane E, Gershow M, Garrity P, Samuel ADT (2011)

   Two Alternating Motor Programs Drive Navigation in Drosophila Larva Skoulakis

   EMC, ed. PLoS ONE 6:e23180 Available at: http://dx.doi.org/10.1371/

   journal.pone.0023180.

Landgraf M, Bossing T, Technau GM, Bate M (1997) The Origin, Location, and

   Projections of the Embryonic Abdominal Motorneurons ofDrosophila. J Neurosci

   17:9642–9655 Available at: http://dx.doi.org/10.1523/

   jneurosci.17-24-09642.1997.

Lin H-T, Trimmer B (2010) Caterpillars use the substrate as their external skeleton.

   Communicative & Integrative Biology 3:471–474 Available at: http://dx.doi.org/

   10.4161/cib.3.5.12560.

Liu Y, Hasegawa E, Nose A, Zwart MF, Kohsaka H (2023) Synchronous multi-segmental

   activity between metachronal waves controls locomotion speed in Drosophila

   larvae. eLife 12 Available at: http://dx.doi.org/10.7554/eLife.83328.

Loveless J, Lagogiannis K, Webb B (2019) Modelling the mechanics of exploration in

   larval Drosophila Ayers J, ed. PLoS Comput Biol 15:e1006635 Available at: http://

   dx.doi.org/10.1371/journal.pcbi.1006635.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018)

   DeepLabCut: markerless pose estimation of user-defined body parts with deep

learning. Nat Neurosci 21:1281–1289 Available at: http://dx.doi.org/10.1038/s41593-018-0209-y.

McMillen T, Williams T, Holmes P (2008) Nonlinear Muscles, Passive Viscoelasticity and Body Taper Conspire To Create Neuromechanical Phase Lags in Anguilliform Swimmers Daniel T, ed. PLoS Comput Biol 4:e1000157 Available at: http://dx.doi.org/10.1371/journal.pcbi.1000157.

McNulty P, Wu R, Yamaguchi A, Heckscher EH, Haas A, Nwankpa A, Skanata MM, Gershow MH (2024) CRASH2p: Closed-loop Two Photon Imaging in a Freely Moving Animal. bioRxiv 595209. Available at: http://dx.doi.org/10.1101/2024.05.22.595209.

Murawski C, Pulver SR, Gather MC (2020) Segment-specific optogenetic stimulation in Drosophila melanogaster with linear arrays of organic light-emitting diodes. Nat Commun 11 Available at: http://dx.doi.org/10.1038/s41467-020-20013-6.

Ormerod KG, Scibelli AE, Littleton JT (2021) Regulation of excitation-contraction coupling at the Drosophila neuromuscular junction. The Journal of Physiology 600:349–372 Available at: http://dx.doi.org/10.1113/jp282092.

Pehlevan C, Paoletti P, Mahadevan L (2016) Integrative neuromechanics of crawling in D. melanogaster larvae. eLife 5 Available at: http://dx.doi.org/10.7554/elife.11031.

Peron S, Zordan MA, Magnabosco A, Reggiani C, Megighian A (2009) From action potential to contraction: Neural control and excitation–contraction coupling in larval muscles of Drosophila. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 154:173–183 Available at: http://dx.doi.org/10.1016/j.cbpa.2009.04.626.

Pulver SR, Bayley TG, Taylor AL, Berni J, Bate M, Hedwig B (2015) Imaging fictive

locomotor patterns in larval Drosophila. Journal of Neurophysiology 114:2564–

2577 Available at: http://dx.doi.org/10.1152/jn.00731.2015.

Risse B, Berh D, Otto N, Klämbt C, Jiang X (2017) FIMTrack: An open source tracking

and locomotion analysis software for small animals. Poisot T, ed. PLoS Comput

Biol 13:e1005530 Available at: http://dx.doi.org/10.1371/journal.pcbi.1005530.

Roberts MJ (1971) On the locomotion of cyclorrhaphan maggots (Diptera). Journal of

Natural History 5:583–590 Available at: http://dx.doi.org/

10.1080/00222937100770421.

Rotheray GE (2019) Forms, Functions and Names. Zoological Monographs:53–86

Available at: http://dx.doi.org/10.1007/978-3-319-92546-2_3.

Schwarz G (1978) Estimating the Dimension of a Model. Ann Statist 6 Available at:

http://dx.doi.org/10.1214/aos/1176344136.

Simon MA, Woods WA Jr, Serebrenik YV, Simon SM, van Griethuijsen LI, Socha JJ, Lee

W-K, Trimmer BA (2010) Visceral-Locomotory Pistoning in Crawling Caterpillars.

Current Biology 20:1458–1463 Available at: http://dx.doi.org/10.1016/

j.cub.2010.06.059.

Singh NP, Mishra RK (2014) Role of abd-A and Abd-B in Development of Abdominal

Epithelia Breaks Posterior Prevalence Rule. Desplan C, ed. PLoS Genet

10:e1004717 Available at: http://dx.doi.org/10.1371/journal.pgen.1004717.

Srinivasan S, Lance K, Levine RB (2012) Segmental differences in firing properties and

potassium currents in Drosophila larval motoneurons. Journal of

Neurophysiology 107:1356–1365 Available at: http://dx.doi.org/10.1152/
jn.00200.2011.

Stern-Tomlinson W, Nusbaum MP, Perez LE, Kristan WB Jr (1986) A kinematic study of
crawling behavior in the leech, Hirudo medicinalis. J Comp Physiol 158:593–603
Available at: http://dx.doi.org/10.1007/bf00603803.

Sun X, Liu Y, Liu C, Mayumi K, Ito K, Nose A, Kohsaka H (2022) A neuromechanical
model for Drosophila larval crawling based on physical measurements. BMC Biol
20 Available at: http://dx.doi.org/10.1186/s12915-022-01336-w.

Takagi S, Cocanougher BT, Niki S, Miyamoto D, Kohsaka H, Kazama H, Fetter RD,
Truman JW, Zlatic M, Cardona A, Nose A (2017) Divergent Connectivity of
Homologous Command-like Neurons Mediates Segment-Specific Touch
Responses in Drosophila. Neuron 96:1373-1387.e6 Available at: http://dx.doi.org/
10.1016/j.neuron.2017.10.030.

Tastekin I, Khandelwal A, Tadres D, Fessner ND, Truman JW, Zlatic M, Cardona A,
Louis M (2018) Sensorimotor pathway controlling stopping behavior during
chemotaxis in the Drosophila melanogaster larva. eLife 7 Available at: http://
dx.doi.org/10.7554/elife.38740.

Thane M, Paisios E, Stöter T, Krüger A-R, Gläß S, Dahse A-K, Scholz N, Gerber B,
Lehmann DJ, Schleyer M (2023) High-resolution analysis of individual Drosophila
melanogaster larvae uncovers individual variability in locomotion and its
neurogenetic modulation. Open Biol 13 Available at: http://dx.doi.org/10.1098/
rsob.220308.

Trimmer B, Issberner J (2007) Kinematics of Soft-bodied, Legged Locomotion in

Manduca sexta Larvae. The Biological Bulletin 212:130–142 Available at: http://

dx.doi.org/10.2307/25066590.

Tsuji T, Hasegawa E, Isshiki T (2008) Neuroblast entry into quiescence is regulated

intrinsically by the combined action of spatial Hox proteins and temporal identity

factors. Development 135:3859–3869 Available at: http://dx.doi.org/10.1242/

dev.025189.

Tytell E, Holmes P, Cohen A (2011) Spikes alone do not behavior make: why

neuroscience needs biomechanics. Current Opinion in Neurobiology 21:816–822

Available at: http://dx.doi.org/10.1016/j.conb.2011.05.017.

Ugur B, Bao H, Stawarski M, Duraine LR, Zuo Z, Lin YQ, Neely GG, Macleod GT,

Chapman ER, Bellen HJ (2017) The Krebs Cycle Enzyme Isocitrate

Dehydrogenase 3A Couples Mitochondrial Metabolism to Synaptic Transmission.

Cell Reports 21:3794–3806 Available at: http://dx.doi.org/10.1016/

j.celrep.2017.12.005.

Vaadia RD, Li W, Voleti V, Singhania A, Hillman EMC, Grueber WB (2019)

Characterization of Proprioceptive System Dynamics in Behaving Drosophila

Larvae Using High-Speed Volumetric Microscopy. Current Biology 29:935-944.e4

Available at: http://dx.doi.org/10.1016/j.cub.2019.01.060.

Vasudevan D, Wang Y, Carr H, Paniel E, Corcoran S, Wreden CC, Kushkowski E, Lee-

Smith C, Heckscher ES (2024) How serially homologous neuroblasts produce

different temporal cohorts along the Drosophila larval body axis. bioRxiv 622783.

Available at: http://dx.doi.org/10.1101/2024.11.09.622783.

Wallén P, Williams TL (1984) Fictive locomotion in the lamprey spinal cord in vitro compared with swimming in the intact and spinal animal. The Journal of Physiology 347:225–239 Available at: http://dx.doi.org/10.1113/jphysiol.1984.sp015063.

Wardle CS, Videler JJ, Altringham JD (1995) Tuning In To Fish Swimming Waves: Body Form, Swimming Mode And Muscle Function. Journal of Experimental Biology 198:1629–1636 Available at: http://dx.doi.org/10.1242/jeb.198.8.1629.

Williams TL, Grillner S, Smoljaninov VV, Wallén P, Kashin S, Rossignol S (1989) Locomotion in Lamprey and Trout: the Relative Timing of Activation and Movement. Journal of Experimental Biology 143:559–557 Available at: http://dx.doi.org/10.1242/jeb.143.1.559.

Wystrach A, Lagogiannis K, Webb B (2016) Continuous lateral oscillations as a core mechanism for taxis in Drosophila larvae. eLife 5 Available at: http://dx.doi.org/10.7554/elife.15504.

FIGURES

Figure 1: Segment-level kinematic and muscle recruitment data were obtained for many cycles of larval locomotion.

A: Schematic of confocal imaging setup.

B: Example frame for a crawling jGCaMP7f larva. Grayscale values, jGCaMP7f fluorescence intensity (inverted). Colored dots, DeepLabCut keypoints at the edges of each segment boundary.

C: Example frame for a crawling Gerry larva. Red (top) and green (bottom) emitted fluorescence were collected in separate channels.

D: Locomotor cycles obtained per larva imaged, n=35 larvae.

E: Schematic representation of larval position. Black dots, keypoints; white dots, segment boundary centroids; double-headed arrow, example segment length.

F: Schematic of contraction features.

G: Schematic of fluorescence features.

Movie 1: DeepLabCut network tracking of two minutes from example jGC7f larva video (ID: "larva 1"). Green channel gamma-adjusted for visualization purposes and displayed in inverted greyscale. Colored dots, tracked keypoints. Video display rate: 20 fps (~real time).

Movie 2: DeepLabCut network tracking of two minutes from example Gerry larva (ID: "larva 22"). Red channel only, gamma-adjusted for visualization purposes and displayed in greyscale. Colored dots, tracked keypoints. Video display rate: 13 fps (~real time).

Figure 2: Contraction kinematics and phase relationships of individual body segments varied along the anterior-posterior body axis.

A: Distributions of per-cycle normalized contraction amplitudes for segments T3 through A7. Box plots show inter-quartile range (IQR) with line for median and asterisk for mean; whiskers show 1.5 times IQR. Outliers were few and are not shown. All pairs of distributions were significantly different (p<0.05) except for A2 vs. A3 and A4 vs. A5.

B: Same as A for maximum shortening rates. All pairs of distributions were significantly different (p<0.05) except for A1 vs. A2 and A3 vs. A5.

C: Same as A for total contraction durations. All pairs of distributions were significantly different (p<0.05) except for A3 vs. A7.

D: Schematic of calculating intersegmental contraction phase delays. Vertical offset for clarity. A5-A4 phase delay for the simulated cycle shown (black double-headed arrow) would be 0.1.

E: Distributions of intersegmental contraction phase delays. Each box plot represents per-cycle delay distribution between a pair of segments. All pairs of delay distributions were significantly different (p<0.05).

F: Distributions of per-cycle contraction phases for segments T3 through A7. Histogram and box plot (without outliers) given for each segment.

All panels: n=3497 cycles, 35 larvae. P-values from nonparametric Friedman's test with post-hoc Dunn-Šidak corrected multiple comparisons.

Figure 3: Axial trends in contraction kinematics did not depend on tail forward movement distance.

A: (Left) Schematic of segment width calculation, which used the segment's anterior boundary. (Right) Segmental distributions of mean boundary widths, n=35 larvae.

B: Segmental distributions for the indicated feature of segmental contraction, for the 50% of cycles where larvae moved their tails forward by a greater distance (left column) or a lesser distance (right column). Median split of cycles was performed separately for each larva. n=1723 cycles for each column.

Figure 4: Axial trends in contraction kinematics are conserved across populations that appear differently constrained inside channels.

A: Scatter plot of individual larvae's mean segment lengths against mean cycle periods. n=35 larvae; points colored per larva as in Figure S1.

B: Scatter plots of same data as in (A), separated into jGC7f (left) and Gerry (right) subpopulations.

C: Segmental distributions for the indicated feature of segmental contraction, given separately for jGC7f (left column) and Gerry (right column) subpopulations.

Figure 5: Recruitment and recruitment phase relationships of individual body segments varied along the anterior-posterior body axis.

A: Distributions of per-cycle total recruitment amplitudes for segments T3 through A7. Box plots show inter-quartile range (IQR) with line for median and asterisk for mean; whiskers show 1.5 times IQR. Outliers were few and are not shown. All pairs of distributions were significantly different (p<0.05) except for A5 vs. A7 and A6 vs. A7.

B: Same as A for recruitment rates (maximum rate of increase in $\Delta G{:}R\text{-}G{:}R_0$). All pairs of distributions were significantly different (p<0.05) except for T3 vs. A5, A5 vs. A7, and A6 vs. A7.

C: Same as A for recruitment durations. All pairs of distributions were significantly different (p<0.05) except for A1 vs. A2.

D: Distributions of intersegmental recruitment phase delays; each box plot represents per-cycle delay distribution between a pair of adjacent segments. All pairs of distributions were significantly different (p<0.05) except for T3 vs. A1, T3 vs. A3, and A1 vs. A3.

E: Distributions of per-cycle recruitment phases for segments T3 through A7. Histogram and box plot (without outliers) are given for each segment. Note that recruitment phases can be negative if (for example) segment A7 began increasing in fluorescence before the tail movement marking the start of a cycle began.

F: Distributions of per-cycle intersegmental recruitment phase delays (colored dots) for a single example larva. Mean phase delay values given as black dots.

A-E: n=1633 cycles, 17 larvae. P-values from nonparametric Friedman's test with post-hoc Dunn-Šidak corrected multiple comparisons.

Figure 6: Phase delays between recruitment and contraction became longer and more variable in the mid-body.

A: Segmental distributions of recruitment phases (grey dots) and contraction phases (colored dots) for all cycles from an example larva ("l21" in Figures S1, S2, S3). Black dots, means.

B: Per-cycle phase delays between a segment's recruitment and its contraction (colored dots) for an example larva. Black dots, means.

C: Segmental distributions of per-cycle phase delays between a segment's recruitment and contraction, for all larvae. Histogram and box plot (without outliers) shown for each segment. Note that recruitment-to-contraction phase delays can be negative if a segment began shortening before substantially increasing in fluorescence. All pairs of distributions were significantly different (p<0.05) except for A6 vs. A7.

D: Segmental distributions of variance in the phase delay between recruitment onset and contraction onset. Box plot (without outliers) is given for each segment. n=17 larvae.

E: Same as B but after excluding central 50% of pixels.

F: Same as C but after excluding central 50% of pixels. All pairs of distributions were significantly different (p<0.05) except for T3 vs. A5.

C and F: n=1633 cycles, 17 larvae. P-values from nonparametric Friedman's test with post-hoc Dunn-Šidak corrected multiple comparisons.

Figure 7: Contraction kinematics were correlated over long intersegmental ranges, with partial dependence on crawling speed.

A: Pairwise intersegmental correlation strength of the indicated kinematic feature, averaged over all segment pairs at each intersegmental distance. Distance=1 indicates neighboring segments, etc. Black asterisks: correlations significantly greater than zero, one-tailed T-test, Bonferroni-corrected alpha (p<0.002). Error bars: S.E.M, n=35 larvae.

B: Pairwise intersegmental correlation strengths of speed-corrected residuals for each kinematic feature, plotted over the original correlation strengths (grey bars, same as in A). To correct for speed, values for each segment were regressed against cycle speed and the result subtracted; correlations were calculated using residual values. Blue asterisks: correlations significantly less than original values, one-tailed T-test, Bonferroni-corrected alpha (p<0.002). Error bars: S.E.M, n=35 larvae.

Figure 8: Range and strength of contraction duration correlations were higher among posterior segments.

A: Schematic of calculating pairwise intersegmental correlation matrices. Example contraction duration relationships shown as scatter plots for two pairs of segments (left). Example plots correspond to two pairwise comparisons in the full correlation matrix (right), marked with boxes.

B: Mean pairwise intersegmental correlations for the indicated feature of segmental

contraction after speed correction (averaged over n=35 larvae). All values along the

diagonal = +1.

C: For each pair of segments, fraction of larvae with significantly (p<0.05) higher-than-

expected pairwise correlations after speed correction.

Figure 9: Contraction duration correlations were frequently higher among a group of posterior segments.

A: Illustration of example model tested by the model-fitting procedure. Indicators "A" and "B" are adjacent-segment-pair correlation strength for different blocks; "C" is the out-of-block correlation strength.

B: 1,094 total models were fit to each larva's empirical correlation matrix, including the null model shown: all segments are included in the same block and correlation strength between segments is entirely a function of intersegmental distance.

C: Two example best-performing block models (right) fit to empirical contraction duration correlation matrices for two individual larvae (left). After all possible block models are fit to a larva's empirical correlation matrix, they are compared using Bayesian Information Criterion (BIC) and the best-scoring model is selected for further analysis. Color display scales and segment order as in B.

D: Number of high-correlation blocks (adjacent-segment R>0.5) found by the best-performing block model for speed-corrected contraction durations for each of 35 larvae.

E: Location of the edges of high-correlation blocks found by the best-performing models (n=35), indicating how often blocks ended between a given pair of segments. Edges between overlapping, similar-strength blocks were excluded (overlapping blocks with <0.2 difference in adjacent-segment R).

F: Frequency with which best-performing models (n=35) included each pair of segments inside the same high-correlation block. The diagonal shows the frequency with which each individual segment was included in a high-correlation block, while the 'exterior corner' (T3 and A7 included in the same block) is equivalent to how many larvae's correlations were best modeled as a single large block: the null model (as in B).

G-I: analogous to D-F for recruitment metrics.