1 TITLE:
2 How sex shapes transcriptome evolution in the songbird brain
3
4 RUNNING TITLE: (39/45 characters)
5 How sex shapes transcriptome evolution
6

7 AUTHORS
8 Isaac Miller-Crews[1], Sara E. Lipshutz[1,2], Ben Fulton[1,3], Jason Bertram[4], Matthew W. Hahn[1,5],
9 and Kimberly A. Rosvall[1]
10
11 *correspondence to: imillerc@iu.edu or krosvall@iu.edu
12
AFFILIATIONS:
1: Department of Biology, Indiana University, Bloomington, IN, USA
2: Department of Biology, Duke University, Durham, NC, USA
3: University Information Technology Services, Indiana University, Bloomington, IN, USA
4: Department of Mathematics, Western University, London, ON, Canada
5: Department of Computer Science, Indiana University, Bloomington, IN, USA
19

20 **ABSTRACT (226 of 250)**
21
Sex differences have captivated scientists for a long time, yet the evolutionary rate of change in
sex-biased gene expression has not been directly quantified. To address this issue, we leverage
brain gene expression data, and we introduce new options in the software package CAGEE
(Computational Analysis of Gene Expression Evolution), specifically unbounded Brownian
motion and variable evolutionary rates among genes. We applied these new features to 10 focal
songbird species, half of which have convergently evolved obligate cavity-nesting, an element of
reproductive ecology linked to sex-specific changes in competition. We find that the degree of
sex bias – measured as the male:female ratio in expression for each gene – evolves twice as fast
on the Z chromosome vs. autosomes, but otherwise, Z gene expression does not evolve at
different rates in males vs. females. Most Z-linked genes are male-biased in their expression,
though some exhibit roughly equal patterns of expression. These sex-balanced genes are not
skewed in their rate of evolution, contrary to the hypothesis that some genes experience selection
for balance and therefore may evolve more slowly. Finally, the degree of sex bias in gene
expression evolves more quickly along obligate-cavity nesting lineages, suggesting that changes
in sex-specific ecological selection shape the evolution of brain sex differences, or lack thereof.
Together, these tools and results provide new insights on the interplay between sex and gene
expression evolution.
39
KEYWORDS: sex, evolution, brain, transcriptome, songbird, behavior

INTRODUCTION

Biologists have long been captivated by why species vary in the degree of sexually differentiated
traits (Darwin, 1871). This variation may come through various mechanisms at different levels of
biological organization, including evolutionary changes to sex chromosomes, gene expression,
and protein sequences (Rowe et al., 2018). For example, within many different species, we know
that sex differences in brain gene expression correspond to sex differences in other components
of the phenotype (Harris & Hofmann, 2014; Renn et al., 2008) (but see (De Vries, 2004).
Comparative studies of small numbers of lineages further re-iterate connections between sex
differences in gene expression and trait evolution (Houle & Cheng, 2021; Ko et al., 2021;
Schumer et al., 2011). There is also evidence that male- vs. female-biased genes differ in how
they evolve (Mank, 2017; Rowe et al., 2018), yet the evolutionary rate of those sex differences in
gene expression has not been directly quantified, underscoring the need for empirical testing
(Tosto et al., 2023).

One challenge to filling this knowledge gap is methodological. Most phylogenetic analyses of
transcriptomic data estimate the rate of evolution in each gene individually, using post-hoc
analyses to infer larger patterns across the transcriptome (e.g. (Pease et al., 2022; Yang et al.,
2019)). Recently, the software CAGEE (Computational Analysis of Gene Expression Evolution)
was developed to assess evolutionary rates while accounting for the fact that genes evolving
along a lineage do not evolve independently from one another (Bertram et al., 2023). CAGEE
v1.0 estimates rates of expression evolution across an entire transcriptome or a subset of the
transcriptome (e.g. a particular chromosome). Variation in these rates can be modeled across one
or more branches of a phylogeny (Fig. 1A), or across different tissues, treatments, or sampling
timepoints within multiple species. However, CAGEE v1.0 and later v1.1 (which fixed several
bugs) otherwise assume all genes share the same average rate of evolution. This assumption
could obscure interesting patterns, considering that different genes could evolve at different
rates. The existing CAGEE software also uses a bounded Brownian motion model, which only
considers continuous positive traits (e.g. absolute gene expression levels). While this approach
works for some comparisons, for instance analyses of each sex independently, the rate of
evolution of sex bias is better captured as a male-to-female ratio of gene expression (hereafter,
'M:F ratio'). Once log-transformed for analyses, this metric includes negative values, thereby
necessitating an unbounded Brownian motion model. By expanding CAGEE to include an
unbounded Brownian motion model (Fig. 1B), alongside the option for different evolutionary
rates among subsets of genes (Fig. 1C), we can directly test long-standing predictions related to
the evolution of sex-biased genes, and in doing so, we also broaden the types of research
questions that can be addressed in comparative transcriptomics.

Songbirds are well-suited to these questions for several reasons. For one, avian chromosomes are
relatively stable, with a high degree of synteny (Ellegren, 2010; Shetty et al., 1999), including on
the Z sex chromosome (Nanda et al., 2008; Zhou et al., 2014). Z genes tend to evolve more
rapidly than those of autosomes, in both coding sequence (Wright, Harrison, et al., 2015) and
expression (Dean et al., 2015), and sex differences in expression may be an important driver of
behavioral variation, even controlling for hormones or gonadal tissue (Wade et al., 1999). Gene
expression on the Z is also 30-40% higher in males compared to females (Itoh et al., 2007; Julien
et al., 2012). This is a notable feature because male birds have two Z chromosomes and females
have one Z and one W. Most animals handle the problem of chromosome number variation

between the sexes via dosage compensation, in which expression outputs are equalized between
the shared sex chromosome in the homogametic and heterogametic sexes (Chen et al., 2020; Gu
& Walters, 2017). However, birds do not exhibit global dosage compensation (Graves, 2016);
instead, individual Z genes vary in their degree of sex difference, and some genes exhibit relative
balance in expression between the sexes (Itoh et al., 2010; Ko et al., 2021; Naurin et al., 2011;
Wolf & Bryk, 2011), prompting questions as to whether these genes are under selection to be
balanced. If so, we may expect that sex-balanced genes would maintain balance across the
phylogeny, and therefore, these more balanced genes would exhibit both a balanced ancestral
state and a slower evolutionary rate in their degree of sex bias. Whether some Z genes
consistently exhibit balance due to evolutionary constraint is unclear and requires direct testing
through quantifying evolution of the degree of sex bias.
98
Beyond these chromosomal considerations, there also may be ecological drivers of gene
expression evolution, for example with foraging guilds (Nesta & Ledon-Rettig, 2025),
subterranean living (Partha et al., 2017), predation pressure (Ghalambor et al., 2015), or sexual
selection (Harrison et al., 2015). For birds, cavity-nesting is another relevant ecological factor,
which has arisen independently over 30 times (Fang et al., 2018). Obligate secondary cavity-
nesters require a hole in a tree or other substrate in order to reproduce, and they cannot excavate
one for themselves. Cavity-nesting is also associated with distinct selection pressures and
phenotypic effects, including slower developmental trajectories, altered predation and thermal
regimes, and more competition for limited breeding sites (Bosque & Bosque, 1995; Drury et al.,
2020; Martin et al., 2017; Martin & Li, 1992; Zenil-Ferguson et al., 2023). Critically for our
consideration of sex and its role in evolution, obligate cavity-nesting is associated with female-
specific increases in conspecific aggressiveness (Lipshutz et al., 2025; Lipshutz & Rosvall,
2021). This increase in aggression is thought to stem from heightened pressures on cavity-
nesting females to find and secure a limited nesting site – pressures that are softened among
females in more flexibly nesting species whose reproductive success does not require such a
restricted ecological niche. How these ecological changes relate to the pace of sex-associated
gene expression evolution is unknown.
116
Here, we use brain gene expression data from 10 songbird species to explore how sex differences
in gene expression evolve, using a new version of CAGEE (v1.2). This updated CAGEE
introduces two new features: (i) an unbounded Brownian motion model and (ii) the option to
allow genes to have different evolutionary rates via gamma-distributed rate categories (see Fig.
1). We apply these new features to test three hypotheses: First, we hypothesize that the degree of
sex differences in gene expression on the Z will evolve more quickly than that of the autosomes,
driven by faster evolution of the female Z. Second, we hypothesize that some Z genes may be
selected for more balanced expression between the sexes; if so, this will be represented by both
more balanced expression and a slower evolutionary rate for the M:F ratio. Third, we
hypothesize that changes in sex-specific selection will shape how sex differences in the brain
evolve, with sex-biased gene expression evolving more rapidly alongside the transition to an
obligate cavity-nesting phenotype. Together, these hypotheses, and the new CAGEE features we
test alongside them, open new doors into our understanding of sex and its interaction with
transcriptomic evolution.

MATERIALS AND METHODS

Sample collection
Samples were collected from mid-March to mid-May 2018-2021 in Indiana, Kentucky, or
Illinois, USA, as detailed in Lipshutz et al. (2025). This earlier paper controlled for sex in an
analysis of convergent evolution of gene expression and behavior; here, we develop and apply
novel analyses that dive deeper into the degree of sex differences and their pace of evolutionary
change. We focused on 10 songbird species, initially selected for a pairwise comparison of
obligate secondary cavity nesting (listed first) vs. their more flexibly nesting relatives (listed
second): Eurasian tree sparrow (*Passer montanus*) vs. house sparrow (*Passer domesticus*),
prothonotary warbler (*Protonotaria citrea*) vs. yellow warbler (*Setophaga petechia*), Eastern
bluebird (*Sialia sialis*) vs. American robin (*Turdus migratorius*), house wren (*Troglodytes
aedon*) vs. Carolina wren (*Thryothorus ludovicianus*), tree swallow (*Tachycineta bicolor*) vs.
barn swallow (*Hirundo rustica*) (Fig. 2A). All birds were in the same pre-breeding stage of
territory establishment and pair bonding, confirmed via behavioral observations and post-mortem
gonadal inspection.
146
Each subject was captured on its territory and euthanized with an overdose of isoflurane
followed by immediate decapitation, brain dissection, and flash freezing on dry ice (n = 6/sex per
species, with n=7 for female tree swallow, for total n=121). Brains were later microdissected
following (Soma et al., 2003), and we focused on the ventromedial telencephalon (VmT), which
includes the avian medial amygdala (or TnA (Reiner et al., 2004) or AMV (Mello et al., 2019)),
bed nucleus of the stria terminalis (BNST), and lateral septum (LS). We isolated total RNA using
Trizol (Invitrogen, USA).
154
Sequencing, alignment, and mapping
Sequencing and pre-processing is detailed in Lipshutz et al. (2025); briefly, we performed
RNAseq (Illumina Nextseq500, 75 cycle), which yielded ~28 million reads per sample.
Transcriptomes for each species were assembled with Trinity version 2.13.2 (Grabherr et al.,
2011) and aligned to the zebra finch (Taeniopygia guttata) proteome
(GCF_003957565.2_bTaeGut1.4.pri_protein.faa) (Rhie et al., 2021), yielding 10,672 one-to-one
orthologs present across all 10 species.
162
We assigned chromosomal location to genes using Ensembl (Harrison et al., 2024) and the zebra
finch reference genome. Because this reference is from a male that does not have a W
chromosome, we did not assign W-linked genes. W-linked genes tend to experience purifying
selection (Bellott et al., 2017; Moghadam et al., 2012), and they may be involved in restoring
dosage balance with their Z-linked homologs (Smeds et al., 2015; Xu & Zhou, 2020).
Nonetheless, they are relatively small in number (~30-50 genes (Ko et al., 2021)), such that their
absence in our dataset should have a minimal impact on our broader findings.
170
171 Most of the genes that differ in their expression between female and males are found on the Z
172 (see Supplement, §1, Fig. S1). Although Z genes have a mainly male-biased distribution (i.e.
173 higher in males than in females), a small fraction of Z genes show approximately equal
174 (balanced) or female-biased expression (see Fig. 2B), providing a strong foundation for testing
175 hypotheses on the degree of sex bias in gene expression.
176

Modelling the evolution of gene expression in CAGEE

Using normalized gene expression counts from DESeq2 (Supplemental Dataset S1), we calculate three main dependent variables: (1) median gene expression for each species; (2) median gene expression for males and females separately, within each species; and (3) ratio of male median and female median gene expression within a species (M:F ratio), which provides insight into the degree of species-level sex differences for each gene (Supplemental Dataset S2). This lattermost metric is the first implementation of an unbounded Brownian motion model, now available in CAGEE v1.2 (https://github.com/hahnlab/CAGEE; Fig. 1B). In practice, both the original bounded Brownian and the new unbounded Brownian motion models require an artificial upper bound to allow numerical analyses to finish in a reasonable time; the new unbounded Brownian motion model also includes an artificial lower bound that is simply the negative reflection of the upper bound.

First, we focus on the role of the Z chromosome in the rate of gene expression evolution, using both pre-existing and new features of CAGEE. We apply a series of nested models to calculate the average rate evolution per million years (σ^2) and a log-likelihood (-lnL) that is used to compare fit among iterations of competing models (Bertram et al., 2023). Specifically, iterations of Model 1 ignore sex, asking only whether the species median rate of gene expression evolution is faster on the Z vs. autosomes (Table 1). Iterations of Model 2 isolate both sex and sex chromosome, asking whether median male and median female gene expression levels evolve at different rates, and whether any differences are driven by the Z chromosome in one or both sexes (Table 2). Iterations of Model 3 focus on the M:F ratio of gene expression, asking directly in a single phylogenetic model how the degree of sex bias evolves on the Z vs. autosome (Table 3).

We also use the second new feature in CAGEE v1.2—gamma-distributed rate categories—to explore how the rate of evolution differs among genes on the same branch of the phylogeny (Fig. 1C). This feature allows multiple, discrete equiprobable rate categories drawn from a gamma distribution (for details see: (Mendes et al., 2020; Yang, 1996). The gamma distribution is calculated from the data, with the shape parameter (α) equal to the scale parameter (β). It is possible for genes to be partial members of multiple rate categories, in that the rate would better be reflected by the inclusion of an additional intermediate category, but a gene can also be assigned to its most likely rate category using maximum likelihood; the σ^2 for that gene is then presumed to be the σ^2 for that category. After validating our approach in a simulated dataset (see Supplement §2, Fig. S3), we modeled our ~10k gene dataset with k=10 rate categories, for both species (median) gene expression and M:F ratio. We chose k=10 rate categories to increase granularity while maintaining a robust number of genes in each category, targeting on average 1,000 genes per category. Although it is not possible to identify an optimal value of k for a dataset (Yang 1996; Mendes et al. 2020), our validations across various k values demonstrate that this choice does not affect our conclusions (see Supplement §2, Fig. S4).

Outputs from these variable rate models were used in several downstream analyses, assigning individual genes to a single rate-category. A) We used a one-tailed Fishers exact test to ask whether genes whose expression evolves quickly are also genes whose sex-ratio of expression evolves quickly (and vice versa, for slowly evolving genes). B) We used a one sample two-tailed t-test to ask directly whether, in each rate category, there was a different percentage of Z chromosome genes compared to other chromosomes, for both gene expression and M:F ratio. C)

We used gene ontology (GO) analyses to ask whether genes with the fastest (or slowest) rates of
evolution are enriched for specific functional pathways, for both species (median) gene
expression and M:F ratio (elaborated in Supplement §3). D) Focused on the Z, we used an
ordinal logistic regression to ask whether ancestral state of sex bias (i.e. M:F ratio at the root
node of the tree (Bertram et al., 2023)), predicts the evolutionary rate category, as is expected if
there is selection for some genes to be more balanced in their expression between the two sexes.
229
Finally, we focus on cavity nesting as a potential ecological driver of sex-specific gene
expression evolution. We modeled species (median) gene expression with bounded Brownian
motion (Table 4), and M:F ratio with unbounded Brownian motion (Table 5). In each case, we
compared three model iterations: i) all branches together (i.e. one σ^2 across the phylogeny); ii)
internal and external branches having separate values of σ^2; iii) and internal, obligate cavity-
nesting, and more flexibly nesting branches each having their own σ^2 parameter. To avoid over-
parameterization, these models were agnostic to chromosome and used a single (average) σ^2 for
all genes.

RESULTS
Modelling the role of sex and sex chromosomes
Focusing first on median gene expression and without considering sex, we found that rates of
evolution on the Z chromosome are faster than on the autosomes (σ^2 = 0.01420, vs. 0.01128);
this model allowing for separate rates among chromosomes is a significantly better fit than the
model that does not allow for different rates (Table 1). In models that directly include both sex
and sex chromosome, the best fit model allows for three rates of gene expression evolution: one
for autosomes in males (σ^2 = 0.01219), another (slightly lower) value for autosomes in females
(σ^2 = 0.01187), and a third (higher) value for the Z, regardless of sex (σ^2 = 0.01513; Table 2). To
test the robustness of these results, we used a random sub-sampling approach to explore the
impact of intra-specific variation on σ^2 (see Supplement §4). We found the same pattern of a
higher σ^2 in male autosomal genes compared to female autosomal genes, and a higher σ^2 of Z
genes with no difference between the sexes (Fig. S6), suggesting that the median value does not
obscure broad patterns. Together, these results show that gene expression on the Z chromosome
evolves ~25% faster than all autosomes, and among autosomes, male gene expression evolves
~5% more quickly than female gene expression values, yet there is no sex difference in
evolutionary rate of gene expression on the Z.

Looking next at the M:F ratio of gene expression, we found that the degree of sex bias on the Z
evolves almost twice as fast as that of autosomes (σ^2 = 0.00070, vs. 0.00037); this model is a
significantly better fit than the model that does not allow for different rates among chromosomes
(Table 3). Note that σ^2-values for this trait are more than an order-of-magnitude lower than for
the median gene expression above. This likely reflects the fact that male and female expression
are generally highly correlated, so that changes in their relative magnitude (reflected in the M:F
ratio) occur more rarely.

Modelling with distinct rate categories
To allow genes to vary in their evolutionary rate regardless of their chromosomal position or pre-
specified hypotheses, we fit a discrete-gamma distribution with 10 rate categories (k = 10) to the
data from both gene expression ($\alpha = \beta$ = 1.12887, σ^2 = 0. 01138) and M:F ratio ($\alpha = \beta$ =

0.792558, $\sigma^2 = 0.00056$), and we found significant overlap among rates categories for these two
metrics (Fig. 3A; FDR p < 0.05). This indicates that, in general, genes evolve quickly in both
their overall expression and their sex ratio (upper right corner Fig. 3A), and genes evolve slowly
in both overall expression and sex ratio (lower left corner). However, the median rate of gene
expression evolution is not necessarily predictive of pace of sex ratio evolution, as there are
many genes in the fastest rate category for M:F ratio that have moderate to low rates for overall
expression evolution, and vice-versa (Fig. 3A, top row and right column).
275
The best-fit gamma distribution shows that most genes evolve slowly, with a long tail of rapidly
evolving genes, for both median expression (Fig. 3B) and M:F ratio (Fig. 3C). Among the fastest
evolving genes, we found no significantly enriched GO categories for either median gene
expression or M:F ratio. For genes with the slowest rate of gene expression evolution, however,
there is significant enrichment for GO pathways involved with cell signaling, protein acetylation,
and mRNA metabolism (Fig. S5A). For the slowest evolving genes in M:F ratio, there is
significant enrichment for GO pathways involved DNA regulation, cell cycle regulation, protein
and mRNA processing (Fig. S5B).
284
Since this multi-rate CAGEE model allows genes to vary in their rate of evolution, without
respect to pre-specified groups of genes, we used this model to test again whether genes on the Z
are evolving faster than those on autosomes. Consistent with our earlier analyses (Tables 2, 3),
the Z chromosome has a higher proportion of genes in the fastest rate categories, and lower
proportion in the slowest rate category, compared to the average autosome for both gene
expression (Fig. 3B) and M:F ratio (Fig. 3C) (see Supplemental Dataset S3 for full results).
Specifically, for gene expression, 8.67% of Z genes were in the fastest rate category (vs.
autosome mean = 6.17%, SE = 0.37) and 4.21% of Z genes were in the slowest rate category (vs.
autosome mean = 9.15%, SE = 0.38). Similarly, for M:F ratio, 14.60% of Z genes were in the
fastest rate category (vs. autosome mean = 5.84%, SE = 0.58) and 0.99% of Z genes were in the
slowest rate category (vs. autosome mean = 8.93%, SE = 0.55).
296
Modelling ancestral states of male:female ratio
Though most genes on the Z are expressed more highly in males, some are more balanced or
even female-biased in their expression (Fig. 2B). Focused on this degree of sex bias (i.e. M:F
ratio) among Z genes, we found no significant relationship between a gene's ancestral state (as
estimated by CAGEE) and its evolutionary rate (ordinal logistic regression: β = -0.137, SE = 0.
4497, p = 0.761; Fig. 4).
303
Modelling the role of nesting strategy
Regarding nesting ecology and focusing first on median gene expression without considering
sex, the best fit model allows for three rates: one for all internal branches along the phylogeny
($\sigma^2 = 0.00625$), another (higher) value for obligate cavity-nesters ($\sigma^2 = 0.01328$), and a third (even
higher) value for species that are not obligate cavity-nesters ($\sigma^2 = 0.01400$). Based on likelihood
ratios, this three-parameter model is a significantly better fit than alternatives (Table 4). These
results indicate that brain gene expression levels evolve ~5% *slower* along obligate cavity-
nesting vs. more flexible-nesting lineages. Analyzing the sexes separately reveals the same
pattern (Tables S1, S2), suggesting that cavity-nesting does not differentially affect rates of brain
gene expression evolution in one vs. the other sex.

In contrast, we find a different outcome when using the M:F ratio. Specifically, obligate cavity-
nesting species have ~18% *faster* evolution of M:F ratio compared to flexible-nesting species (σ^2
$= 0.00059$ vs. 0.00050; Table 5). Thus, even though median gene expression and sex ratio of
expression are not completely independent, these different measures provide different insights on
the interplay between sex and transcriptomic evolution.

DISCUSSION

We developed new features in CAGEE (v1.2, Fig. 1), applied here to test three hypotheses on the
evolution of sex-biased brain gene expression in songbirds. In support of our first hypothesis,
genes on the Z chromosome exhibit a faster evolutionary rate of neural median gene expression
and M:F ratio, compared to autosomal genes. However, there is no strong evidence that among
our focal species gene expression on the Z evolves faster in males than females. Additionally, Z
genes with slower evolutionary rates in sex ratio are not ancestrally more balanced between the
sexes, a pattern that is inconsistent with our second hypothesis, that the relatively balanced
expression seen in some Z genes may be a result of historical selection for balance (Mank &
Ellegren, 2009). Finally, we find that sex ratios in gene expression evolve more rapidly along
songbird lineages that have convergently evolved obligate cavity-nesting, a trait associated with
sex-specific changes in selection related to territorial aggression (Lipshutz et al., 2025). Across
multiple results, our direct analyses of sex ratio evolution provide different effect sizes from
indirect analyses that measure the sexes separately (Table 2 vs. 3; Table 4 vs. 5; Fig. 3B vs. 3C).
Together, these findings demonstrate (i) that the sexes need to be studied concurrently to
understand how evolution shapes gene expression and (ii) how new features in CAGEE v1.2 can
advance the field of comparative transcriptomics.

Z chromosomes and the evolution of gene expression
All ten focal species displayed similar levels of male-biased expression on the Z (Fig. 2B &
Supplemental §1), with Z gene expression averaging 59% higher for males than for females. This
unequal gene expression has been documented previously (Itoh et al., 2007; Julien et al., 2012)
and is important for the study of sex differences because there is evidence in birds that secondary
sexual traits are driven by sex differences in expression of Z genes, even when controlling for
differences in gonadal hormones or global methylation (Diddens et al., 2021; Ioannidis et al.,
2021; Wade et al., 1999). We extend these findings here, documenting faster gene expression
evolution on the Z, coupled with a faster rate of evolution of the sex ratio of expression as well.
These higher rates on the Z could be the result of reduced effective population size of the Z
compared to autosomes, differences in selection on the two sets of chromosomes, or some
combination of these (Dean et al., 2015; Hayes et al., 2020; Ottenburghs, 2022). Furthermore,
the higher evolutionary rates we observe do not distinguish among the causes of these
differences, which would require future study. However, we do not find evidence for faster gene
expression evolution on the *female* Z relative to the male Z (Table 2), as has previously been
shown in peripheral tissue of other non-songbirds (Dean et al., 2015). It is therefore possible that,
unlike peripheral tissue, neural tissue does not show a pattern of a faster evolutionary rate on the
female Z, suggesting the potential for tissue-specific rates of sex ratio gene expression evolution.

With the new unbounded Brownian motion feature introduced here, we now provide a non-
mutually exclusive alternative interpretation– that outcomes from direct analyses on the degree

of sex bias give different answers than analyses of each sex separately. Modelling gene
expression evolution for the sexes separately, though, we found only small differences in the
pace of change on autosomes, with male gene expression evolving about 5% faster than that of
females (Table 2). This pattern is consistent with a previous analysis in peripheral tissue from
Galliform birds, which reported a greater number of evolutionary transitions of male-biased
compared to female-biased autosomal genes (Harrison et al., 2015). While our sex-specific
analysis of gene expression on the Z vs. autosomes revealed only modest differences, our direct
analysis of the degree of sex difference (i.e. M:F ratio) revealed that sex-biased gene expression
on the Z evolves almost twice as fast as that of autosomes. Importantly, there is evidence in birds
that sex-biased gene expression on the Z may drive tissue-specific sexual differentiation, which
is only weakly controlled via sex hormones after gonadal differentiation (Deviatiiarov et al.,
2023; Ioannidis et al., 2021; Zhao et al., 2010). Moving forward, it will be critical to apply these
direct approaches beyond our focal songbird species and in other tissues, to assess generality.
More broadly and beyond the context of sex, the unbounded Brownian motion feature can also
apply to model the degree of widely used treatment effects (e.g. response to social challenge
sensu (Rittschof et al., 2014)), as well as other non-positive continuous data, like those extracted
from PCA, WGCNA, or other multi-omics datasets.
376
Variable rates of expression evolution among genes
The second new CAGEE feature we developed– variable rates of evolution for different genes on
the same branch of the phylogeny – further advances understanding of the evolution of sex-
biased gene expression. For one, by assigning genes into ten rate categories without *a priori*
hypotheses, we provide a separate inference on the rate of Z vs. the autosomes, mirroring the
single rate analyses (Tables 1-3) in that genes with the fastest evolutionary rates are concentrated
on the Z chromosome (Fig. 3B,C).
384
The variable-rate feature also opens doors to asking questions about why some genes evolve
more quickly in their degree of sex bias vs. balance, a particularly salient issue in birds due to
their lack of global dosage compensation (Fallahshahroudi et al., 2025; Itoh et al., 2007; Julien et
al., 2012). Nonetheless, in each of our 10 focal species, we find that some Z-linked genes are
expressed relatively equally between females and males (Fig. 2B), consistent with prior work
(Mank & Ellegren, 2009; Zimmer et al., 2016). Past comparisons of two or three bird species
also provide some evidence that there may be consistency in which Z genes are most balanced
between the sexes over the course of evolutionary time (Itoh et al., 2010; Ko et al., 2021; Naurin
et al., 2011; Wolf & Bryk, 2011), as would be expected if there is some form of gene-by-gene
purifying selection for balance (Mank & Ellegren, 2009). If so, those genes on the Z under
selection for a more balanced sex ratio would be conserved, as reflected in both a relatively
balanced ancestral state and a slower rate of gene expression evolution. Contrary to this
hypothesis, though, we find no relationship between the rate of gene expression evolution and
the degree of sex bias at the ancestral root node of these 10 songbirds (Fig. 4). There is a
possibility that this lack of purifying selection manifests in other tissues, life-history stages, or
even within other lineages. For example, other bird lineages, but not songbirds, have
hypermethylated regions of the Z that may represent the first step towards the evolution of
complete dosage compensation (Melamed & Arnold, 2007; Sun et al., 2019; Wright, Zimmer, et
al., 2015). We also find that most Z genes are not significantly male-biased across our 10 focal
songbird species (Fig. S1), further indicating that there is standing natural variation in the degree

of sex bias. Other processes, like transcriptional bursting (Papanicolaou et al., 2025), post-
transcriptional compensation (Fallahshahroudi et al., 2025; Lister et al., 2024; Uebbing et al.,
2015), gene-specific hormonal effects (van Nas et al., 2009), or variation across developmental
time (Mank & Ellegren, 2009) may still operate, and future analyses should further explore how
these factors interact with rates of gene expression evolution.
410
Another non-mutually exclusive hypothesis is that genes with certain functions evolve at
different rates. This issue has been explored in depth in the past, demonstrating the slower
sequence evolution of essential genes related to the synthesis of DNA, RNA, and proteins across
animals (Rancati et al., 2018; Zhan & Boutros, 2016). We approached this question with GO
analyses of the genes in the most extreme rate categories, finding no functional enrichment for
genes that evolve most quickly– either in terms of their median levels of expression or their
degree of sex bias. Focused on genes with the slowest rates of gene expression evolution,
however, we do find enrichment of several basic cell functions, including cell cycle, chromatin
organization, RNA and protein processing. These findings are consistent with the likely
constraints that shape these mission critical pathways, with little wiggle room for gene
expression, or its degree of sex bias, to change among species.
422
Our analyses underscore open questions as to why and how it is that some Z genes exhibit
relative balance in their expression despite the fact that males have two copies of the Z and
females have just one. Notably in our analyses, genes that evolve most quickly in their degree of
sex bias also tend to evolve most quickly in their median expression level. Exceptions occur,
with hundreds of genes that evolve quickly in their sex ratio but not their overall expression, and
vice versa (Fig. 3A). Thus, even though these two metrics are related, they can vary semi-
independently, including cases in which overall expression evolves slowly but the sex ratio of
expression still evolves quickly. We urge future research to explore these relationships across
tissues and across different taxa, alongside further optimization of the best number of rate
categories for these and other research questions.
433
Sex associated gene expression evolution and reproductive ecology
Past research provides key insights on the ecological drivers of sex differences (Gonzalez-Voyer
et al., 2022), and we extend this here in relation to obligate cavity-nesting in songbirds, which
involves many changes in the phenotype and has evolved convergently across half of the species
in our study (Fig. 1A). A previous analysis, using these same transcriptomic data from the
ventromedial telencephalon, identified a small set of genes whose expression convergently
evolves alongside higher levels of aggression and obligate cavity-nesting (Lipshutz et al., 2025).
Here, we add new findings linking this element of reproductive ecology with the degree of sex
bias in gene expression and the pace of gene expression evolution. Specifically, along obligate
cavity-nesting lineages, we find a lower overall rate of gene expression evolution (Table 4),
suggesting an ecological constraint on transcriptome evolution. Alternative models that treat the
sexes separately are not a significantly better fit (Tables S1, S2), implying that the sex-related
gene expression does not evolve differently alongside the convergent evolution of obligate
cavity-nesting. Critically, though, with the new ratio-feature in CAGEE v1.2, we see that the
M:F expression evolves significantly *faster* along obligate-cavity nesting lineages (Table 5), an
insight we would have missed from studying the sexes independently or without regard to the
degree of sex bias. These results suggest that, in association with a key component of

reproductive ecology, brain gene expression is evolving more slowly in absolute terms, but the
expression relationship between males and females is more volatile, with shifts that are more
marked in one sex versus the other. While we are unable to determine causality, we speculate
that this ecologically-associated acceleration of sex bias evolution relates to the need for
different mechanisms to drive convergent behavior across the sexes (sensu: (De Vries, 2004; Ko
et al., 2021). More broadly, these results speak to an emerging view that research on sex should
consider the degree of sex differences (McCarthy et al., 2012; Smiley et al., 2024) as we seek to
understand the interplay between sex and evolutionary processes.
459

TABLES

Table 1. Species gene expression evolutionary rate

Model	-lnl	rate	σ^2
All	337733	All	0.01138
Autosomal vs Z	**337690**	**A**	**0.01128**
		Z	**0.01420**

Table 2. Gene expression evolutionary rate per sex

Model	-lnl	rate	σ^2
All	680226	All	0.01203
Male vs Female	680220	M	0.01219
		F	0.01187
Autosomal vs Z	680149	A	0.01194
		Z	0.01508
Autosomal vs Male Z vs Female Z	680149	MA+FA	0.01194
		MZ	0.01563
		FZ	0.01451
Male Autosomal vs Female Autosomal vs Z	**680143**	**MA**	**0.01209**
		FA	**0.01180**
		MZ+FZ	**0.01513**
Male Autosomal vs Female Autosomal vs Male Z vs Female Z	680143	MA	0.01210
		FA	0.01176
		MZ	0.01529
		FZ	0.01444

Table 3. M:F ratio gene expression evolutionary rate

Model	-lnl	rate	σ^2
All	214132	All	0.00038
Autosomal vs Z	**213901**	**A**	**0.00037**
		Z	**0.00070**

Table 4. Cavity nesting gene expression evolutionary rate

Model	-lnl	rate	σ^2
All	337733	All	0.01138
Internal vs External	336583	Internal	0.00629
		External	0.01360
Nesting Phenotype	**336577**	**Internal**	**0.00625**
		Obligate	**0.01328**
		Flexible	**0.01401**

Table 5. M:F ratio cavity nesting gene expression evolutionary rate			
Model	-lnl	rate	σ^2
All	214132	All	0.00038
Internal vs External	208787	Internal	0.00006
		External	0.00055
Nesting Phenotype	**208723**	**Internal**	**0.00006**
		Obligate	**0.00059**
		Flexible	**0.00050**

FIGURES

473
Figure 1:

CAGEE v1.2 now integrates two new features to extend the original CAGEE. **A)** With CAGEE
v1.0, a transcriptome-wide evolutionary rate (σ^2) can be calculated across the phylogeny, or
different σ^2-values can be calculated for specific branches (green vs brown lineages vs black
internal branches). σ^2 represents the evolutionary variance over time, with larger values indicating
a faster amount of change. CAGEE also uses trait values at the tips of the tree to conduct ancestral
state reconstruction at internal nodes. **B)** CAGEE v1.0 models incremental changes over time with
a bounded Brownian motion model (top), for use with continuous positive traits (e.g. gene
expression). CAGEE v1.2 also includes an unbounded Brownian motion (bottom), for use with
any continuous trait (e.g. ratio of gene expression between sexes, converted to a log scale). **C)**
CAGEE v1.0 calculates a single σ^2 for the average σ^2 (black triangle) across the distribution of all
genes (top). CAGEE v1.2 allows for variation in σ^2 among genes (bottom) by fitting data to a
discrete-gamma distribution, calculating k equiprobable rate categories (e.g. k = 10), and assigning
the average rate (triangle) to all genes in that category.

488
Figure 2:

A) Phylogeny of 10 focal species from TimeTree (Kumar et al., 2022). External branches are
colored by cavity-nesting phenotype (brown = obligate cavity-nesting, green = not obligate
cavity-nesting), with internal branches colored black (*illustrations by Tessa Patton*). **B)** Density
plots show M:F gene expression ratio for each species. Each species is a row, with autosomal
genes in grey and Z chromosome genes in red. Positive values indicate higher expression in
males compared to females, and vice versa.
496

497
Figure 3:
A) Comparison of the number of genes in each rate category (σ^2), based on M:F ratio of
expression (y-axis) and median expression (x-axis). Significance is calculated for enrichment
between every rate-category separately; *p < 0.05, **p < 0.01, ***p < 0.001. **B)** Percentage of
genes assigned to different rate categories (σ^2) on the Z chromosome (red) and autosomes (black;
mean and SE) for median gene expression. **C)** Percentage of genes assigned to different rate
categories (σ^2) on the Z chromosome (red) and autosomes (black; mean and SE) for M:F ratio.
Significance is calculated for Z vs. autosomes for every rate-category separately; *p < 0.05, **p
< 0.01, ***p < 0.001.

Figure 4:

509 On the Z chromosome, the sex ratio of the ancestral state at the root of the tree (x-axis) is
510 unrelated to the rate of gene expression evolution for M:F ratio (y-axis). Box and whisker plots
511 show distribution of sex ratio ancestral states for Z genes in each M:F σ^2 category, with outliers
512 shown as points. Dashed vertical line is the average ancestral state for all Z genes.

ACKNOWLEDGMENTS
For discussion and advice, we thank members of the Rosvall and Hahn labs, as well as three anonymous reviewers. For support in data collection and sequencing, we thank AB Bentz, AM Buechlein, TA Empson, EM George, ME Hauber, DB Rusch, WM Schelsky, AM Turner, SE Wolf, MJ Woodruff, and the IU Center for Genomics and Bioinformatics.

FUNDING
Common Themes in Reproductive Diversity NIH 2T32HD049336 (to IMC), National Science Foundation (NSF) grants DBI-1907134 (to SEL), DBI-2146866 to MWH, IOS-1942192 (to KAR).

DATA AVAILABILITY STATEMENT
All data are available in the main text and/or supplementary materials, with scripts available on Zenodo (Miller-Crews, 2026). The CAGEE software is available at https://github.com/hahnlab/CAGEE.

BENEFITS GENERATED
Benefits of this research come from the public sharing of our data and results, discussed above.

AUTHOR CONTRIBUTIONS: SEL, IMC, MWH, and KAR conceived of the study; SEL led the sample collection and sequencing. IMC analyzed and interpreted the data, supported by BF and JB. BF, JB, and MWH developed computational tools. IMC, MWH, and KAR wrote the paper, and all authors contributed to revisions. The manuscript reflects the contributions and ideas of all authors.

Reference:
Bellott, D. W., Skaletsky, H., Cho, T. J., Brown, L., Locke, D., Chen, N., Galkina, S., Pyntikova,
T., Koutseva, N., Graves, T., Kremitzki, C., Warren, W. C., Clark, A. G., Gaginskaya, E.,
Wilson, R. K., & Page, D. C. (2017). Avian W and mammalian Y chromosomes
convergently retained dosage-sensitive regulators. *Nat Genet*, *49*(3), 387-394.
https://doi.org/10.1038/ng.3778

Bertram, J., Fulton, B., Tourigny, J. P., Pena-Garcia, Y., Moyle, L. C., & Hahn, M. W. (2023).
CAGEE: Computational Analysis of Gene Expression Evolution. *Molecular Biology and*
Evolution, *40*(5). https://doi.org/10.1093/molbev/msad106

Bosque, C., & Bosque, M. T. (1995). Nest Predation as a Selective Factor in the Evolution of
Developmental Rates in Altricial Birds. *American Naturalist*, *145*(2), 234-260.
https://doi.org/Doi 10.1086/285738

Chen, J., Wang, M., He, X., Yang, J.-R., & Chen, X. (2020). The evolution of sex chromosome
dosage compensation in animals. *Journal of Genetics and Genomics*, *47*(11), 681-693.
https://doi.org/10.1016/j.jgg.2020.10.005

Darwin, C. (1871). *Descent of man, and selection in relation to sex*.

De Vries, G. J. (2004). Minireview: Sex differences in adult and developing brains:
compensation, compensation, compensation. *Endocrinology*, *145*(3), 1063-1068.
https://doi.org/10.1210/en.2003-1504

Dean, R., Harrison, P. W., Wright, A. E., Zimmer, F., & Mank, J. E. (2015). Positive Selection
Underlies Faster-Z Evolution of Gene Expression in Birds. *Molecular Biology and*
Evolution, *32*(10), 2646-2656. https://doi.org/10.1093/molbev/msv138

Deviatiiarov, R., Nagai, H., Ismagulov, G., Stupina, A., Wada, K., Ide, S., Toji, N., Zhang, H.,
Sukparangsi, W., Intarapat, S., Gusev, O., & Sheng, G. (2023). Dosage compensation of
Z sex chromosome genes in avian fibroblast cells. *Genome Biology*, *24*(1), 213.
https://doi.org/10.1186/s13059-023-03055-z

Diddens, J., Coussement, L., Frankl-Vilches, C., Majumdar, G., Steyaert, S., Ter Haar, S. M.,
Galle, J., De Meester, E., De Keulenaer, S., Van Criekinge, W., Cornil, C. A., Balthazart,
J., Van Der Linden, A., De Meyer, T., & Vanden Berghe, W. (2021). DNA Methylation
Regulates Transcription Factor-Specific Neurodevelopmental but Not Sexually
Dimorphic Gene Expression Dynamics in Zebra Finch Telencephalon. *Frontiers in Cell*
and Developmental Biology, *9*, 583555. https://doi.org/10.3389/fcell.2021.583555

Drury, J. P., Cowen, M. C., & Grether, G. F. (2020). Competition and hybridization drive
interspecific territoriality in birds. *Proceedings of the National Academy of Sciences of*
the United States of America, *117*(23), 12923-12930.
https://doi.org/10.1073/pnas.1921380117

Ellegren, H. (2010). Evolutionary stasis: the stable chromosomes of birds. *Trends Ecol Evol*,
25(5), 283-291. https://doi.org/10.1016/j.tree.2009.12.004

Fallahshahroudi, A., Yousefi Taemeh, S., Rodriguez-Montes, L., Trost, N., Frank, D., Lafrenz,
P., Koubek, J., Tellez, G., Jr., Ballantyne, M., Idoko-Akoh, A., Taylor, L., Sherman, A.,
Davey, M., Ma, C., Sorato, E., Johnsson, M., Grozou, C., Xue, Y., Liu, L.,…Kaessmann,
H. (2025). A male-essential miRNA is key for avian sex chromosome dosage
compensation. *Nature*. https://doi.org/10.1038/s41586-025-09256-9

Fang, Y. T., Tuanmu, M. N., & Hung, C. M. (2018). Asynchronous evolution of interdependent
nest characters across the avian phylogeny. *Nature Communications*, *9*(1), 1863.
https://doi.org/10.1038/s41467-018-04265-x

Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., & Hughes, K. A.
(2015). Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in
nature. *Nature*, *525*(7569), 372-375. https://doi.org/10.1038/nature15256

Gonzalez-Voyer, A., Thomas, G. H., Liker, A., Kruger, O., Komdeur, J., & Szekely, T. (2022).
Sex roles in birds: Phylogenetic analyses of the influence of climate, life histories and
social environment. *Ecol Lett*, *25*(3), 647-660. https://doi.org/10.1111/ele.13938

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X.,
Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A.,
Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K.,…Regev, A.
(2011). Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nature Biotechnology*, *29*(7), 644-652. https://doi.org/10.1038/nbt.1883

Graves, J. A. (2016). Evolution of vertebrate sex chromosomes and dosage compensation. *Nat*
Rev Genet, *17*(1), 33-46. https://doi.org/10.1038/nrg.2015.2

Gu, L., & Walters, J. R. (2017). Evolution of Sex Chromosome Dosage Compensation in
Animals: A Beautiful Theory, Undermined by Facts and Bedeviled by Details. *Genome*
Biology and Evolution, *9*(9), 2461-2476. https://doi.org/10.1093/gbe/evx154

Harris, R. M., & Hofmann, H. A. (2014). Neurogenomics of behavioral plasticity. *Adv Exp Med*
Biol, *781*, 149-168. https://doi.org/10.1007/978-94-007-7347-9_8

Harrison, P. W., Amode, M. R., Austine-Orimoloye, O., Azov, A. G., Barba, M., Barnes, I.,
Becker, A., Bennett, R., Berry, A., Bhai, J., Bhurji, S. K., Boddu, S., Branco Lins, P. R.,
Brooks, L., Ramaraju, S. B., Campbell, L. I., Martinez, M. C., Charkhchi, M., Chougule,
K.,…Yates, A. D. (2024). Ensembl 2024. *Nucleic Acids Res*, *52*(D1), D891-D899.
https://doi.org/10.1093/nar/gkad1049

Harrison, P. W., Wright, A. E., Zimmer, F., Dean, R., Montgomery, S. H., Pointer, M. A., &
Mank, J. E. (2015). Sexual selection drives evolution and rapid turnover of male gene
expression. *Proceedings of the National Academy of Sciences of the United States of*
America, *112*(14), 4393-4398. https://doi.org/10.1073/pnas.1501339112

Hayes, K., Barton, H. J., & Zeng, K. (2020). A Study of Faster-Z Evolution in the Great Tit
(Parus major). *Genome Biology and Evolution*, *12*(3), 210-222.
https://doi.org/10.1093/gbe/evaa044

Houle, D., & Cheng, C. (2021). Predicting the Evolution of Sexual Dimorphism in Gene
Expression. *Molecular Biology and Evolution*, *38*(5), 1847-1859.
https://doi.org/10.1093/molbev/msaa329

Ioannidis, J., Taylor, G., Zhao, D., Liu, L., Idoko-Akoh, A., Gong, D., Lovell-Badge, R., Guioli,
S., McGrew, M. J., & Clinton, M. (2021). Primary sex determination in birds depends on
DMRT1 dosage, but gonadal sex does not determine adult secondary sex characteristics.
Proceedings of the National Academy of Sciences of the United States of America,
118(10), e2020909118. https://doi.org/10.1073/pnas.2020909118

Itoh, Y., Melamed, E., Yang, X., Kampf, K., Wang, S., Yehya, N., Van Nas, A., Replogle, K.,
Band, M. R., Clayton, D. F., Schadt, E. E., Lusis, A. J., & Arnold, A. P. (2007). Dosage
compensation is less effective in birds than in mammals. *Journal of Biology*, *6*(1), 2.
https://doi.org/10.1186/jbiol53

Itoh, Y., Replogle, K., Kim, Y. H., Wade, J., Clayton, D. F., & Arnold, A. P. (2010). Sex bias
and dosage compensation in the zebra finch versus chicken genomes: general and
specialized patterns among birds. *Genome Research*, *20*(4), 512-518.
https://doi.org/10.1101/gr.102343.109

Julien, P., Brawand, D., Soumillon, M., Necsulea, A., Liechti, A., Schutz, F., Daish, T.,
Grutzner, F., & Kaessmann, H. (2012). Mechanisms and evolutionary patterns of
mammalian and avian dosage compensation. *PLoS Biology*, *10*(5), e1001328.
https://doi.org/10.1371/journal.pbio.1001328

Ko, M. C., Frankl-Vilches, C., Bakker, A., & Gahr, M. (2021). The Gene Expression Profile of
the Song Control Nucleus HVC Shows Sex Specificity, Hormone Responsiveness, and
Species Specificity Among Songbirds. *Frontiers in Neuroscience*, *15*, 680530.
https://doi.org/10.3389/fnins.2021.680530

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., Stecher, G., &
Hedges, S. B. (2022). TimeTree 5: An Expanded Resource for Species Divergence
Times. *Molecular Biology and Evolution*, *39*(8). https://doi.org/10.1093/molbev/msac174

Lipshutz, S. E., Hibbins, M. S., Bentz, A. B., Buechlein, A. M., Empson, T. A., George, E. M.,
Hauber, M. E., Rusch, D. B., Schelsky, W. M., Thomas, Q. K., Torneo, S. J., Turner, A.
M., Wolf, S. E., Woodruff, M. J., Hahn, M. W., & Rosvall, K. A. (2025). Repeated
behavioural evolution is associated with convergence of gene expression in cavity-
nesting songbirds. *Nature Ecology & Evolution*, *9*(5), 845-856.
https://doi.org/10.1038/s41559-025-02675-x

Lipshutz, S. E., & Rosvall, K. A. (2021). Nesting strategy shapes territorial aggression but not
testosterone: A comparative approach in female and male birds. *Hormones and behavior*,
133, 104995. https://doi.org/10.1016/j.yhbeh.2021.104995

651   Lister, N. C., Milton, A. M., Patel, H. R., Waters, S. A., Hanrahan, B. J., McIntyre, K. L.,
652   Livernois, A. M., Horspool, W. B., Wee, L. K., Ringel, A. R., Mundlos, S., Robson, M.
653   I., Shearwin-Whyatt, L., Grutzner, F., Graves, J. A. M., Ruiz-Herrera, A., & Waters, P.
654   D. (2024). Incomplete transcriptional dosage compensation of chicken and platypus sex
655   chromosomes is balanced by post-transcriptional compensation. *Proceedings of the*
656   *National Academy of Sciences of the United States of America*, *121*(32), e2322360121.
657   https://doi.org/10.1073/pnas.2322360121

658   Mank, J. E. (2017). The transcriptional architecture of phenotypic dimorphism. *Nature Ecology*
659   *& Evolution*, *1*(1), 6. https://doi.org/10.1038/s41559-016-0006

660   Mank, J. E., & Ellegren, H. (2009). All dosage compensation is local: gene-by-gene regulation of
661   sex-biased expression on the chicken Z chromosome. *Heredity (Edinb)*, *102*(3), 312-320.
662   https://doi.org/10.1038/hdy.2008.116

663   Martin, T. E., Boyce, A. J., Fierro-Calderón, K., Mitchell, A. E., Armstad, C. E., Mouton, J. C.,
664   & Bin Soudi, E. E. (2017). Enclosed nests may provide greater thermal than nest
665   predation benefits compared with open nests across latitudes. *Functional Ecology*, *31*(6),
666   1231-1240. https://doi.org/10.1111/1365-2435.12819

667   Martin, T. E., & Li, P. (1992). Life History Traits of Open- vs. Cavity-Nesting Birds. *Ecology*,
668   *73*(2), 579-592. https://doi.org/10.2307/1940764

669   McCarthy, M. M., Arnold, A. P., Ball, G. F., Blaustein, J. D., & De Vries, G. J. (2012). Sex
670   differences in the brain: the not so inconvenient truth. *J Neurosci*, *32*(7), 2241-2247.
671   https://doi.org/10.1523/JNEUROSCI.5372-11.2012

672   Melamed, E., & Arnold, A. P. (2007). Regional differences in dosage compensation on the
673   chicken Z chromosome. *Genome Biology*, *8*(9), R202. https://doi.org/10.1186/gb-2007-8-
674   9-r202

675   Mello, C. V., Kaser, T., Buckner, A. A., Wirthlin, M., & Lovell, P. V. (2019). Molecular
676   architecture of the zebra finch arcopallium. *The Journal of Comparative Neurology*,
677   *527*(15), 2512-2556. https://doi.org/10.1002/cne.24688

678   Mendes, F. K., Vanderpool, D., Fulton, B., & Hahn, M. W. (2020). CAFE 5 models variation in
679   evolutionary rates among gene families. *Bioinformatics*, *36*(22-23), 5516-5518.
680   https://doi.org/10.1093/bioinformatics/btaa1022

681   Miller-Crews, I. (2026). imillercrews/10sp_CAGEE: 10sp_CAGEE_v1.1 (v1.1.0). *Zenodo*.
682   https://doi.org/https://doi.org/10.5281/zenodo.16883711

683   Moghadam, H. K., Pointer, M. A., Wright, A. E., Berlin, S., & Mank, J. E. (2012). W
684   chromosome expression responds to female-specific selection. *Proceedings of the*
685   *National Academy of Sciences*, *109*(21), 8207-8211.
686   https://doi.org/10.1073/pnas.1202721109

Nanda, I., Schlegelmilch, K., Haaf, T., Schartl, M., & Schmid, M. (2008). Synteny conservation
of the Z chromosome in 14 avian species (11 families) supports a role for Z dosage in
avian sex determination. *Cytogenet Genome Res*, *122*(2), 150-156.
https://doi.org/10.1159/000163092

Naurin, S., Hansson, B., Hasselquist, D., Kim, Y. H., & Bensch, S. (2011). The sex-biased brain:
sexual dimorphism in gene expression in two species of songbirds. *BMC Genomics*,
12(1), 37. https://doi.org/10.1186/1471-2164-12-37

Nesta, D. J., & Ledon-Rettig, C. C. (2025). Cryptic genetic variation in brain gene expression
precedes the evolution of cannibalism in spadefoot toad tadpoles. *Proceedings of the*
National Academy of Sciences of the United States of America, *122*(19), e2418431122.
https://doi.org/10.1073/pnas.2418431122

Ottenburghs, J. (2022). Avian Introgression Patterns are Consistent With Haldane's Rule. *J*
Hered, *113*(4), 363-370. https://doi.org/10.1093/jhered/esac005

Papanicolaou, N., Lentini, A., Wettersten, S., Hagemann-Jensen, M., Kruger, A., Zhang, J.,
Coucoravas, C., Petrosian, I., Xin, X., Ceyhan, I., Rorbach, J., Wright, D., & Reinius, B.
(2025). Multi-layered dosage compensation of the avian Z chromosome by increased
transcriptional burst frequency and elevated translational rates. *Nature Communications*,
16(1), 9088. https://doi.org/10.1038/s41467-025-64817-w

Partha, R., Chauhan, B. K., Ferreira, Z., Robinson, J. D., Lathrop, K., Nischal, K. K., Chikina,
M., & Clark, N. L. (2017). Subterranean mammals show convergent regression in ocular
genes and enhancers, along with adaptation to tunneling. *eLife*, *6*.
https://doi.org/10.7554/eLife.25884

Pease, J. B., Driver, R. J., de la Cerda, D. A., Day, L. B., Lindsay, W. R., Schlinger, B. A.,
Schuppe, E. R., Balakrishnan, C. N., & Fuxjager, M. J. (2022). Layered evolution of gene
expression in "superfast" muscles for courtship. *Proceedings of the National Academy of*
Sciences of the United States of America, *119*(14), e2119671119.
https://doi.org/10.1073/pnas.2119671119

Rancati, G., Moffat, J., Typas, A., & Pavelka, N. (2018). Emerging and evolving concepts in
gene essentiality. *Nat Rev Genet*, *19*(1), 34-49. https://doi.org/10.1038/nrg.2017.74

Reiner, A., Perkel, D. J., Bruce, L. L., Butler, A. B., Csillag, A., Kuenzel, W., Medina, L.,
Paxinos, G., Shimizu, T., Striedter, G., Wild, M., Ball, G. F., Durand, S., Gunturkun, O.,
Lee, D. W., Mello, C. V., Powers, A., White, S. A., Hough, G.,…Avian Brain
Nomenclature, F. (2004). Revised nomenclature for avian telencephalon and some related
brainstem nuclei. *The Journal of Comparative Neurology*, *473*(3), 377-414.
https://doi.org/10.1002/cne.20118

Renn, S. C., Aubin-Horth, N., & Hofmann, H. A. (2008). Fish and chips: functional genomics of
social plasticity in an African cichlid fish. *The Journal of Experimental Biology*, *211*(Pt
18), 3041-3056. https://doi.org/10.1242/jeb.018242

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M.,…Jarvis, E. D. (2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature*, *592*(7856), 737-746. https://doi.org/10.1038/s41586-021-03451-0

Rittschof, C. C., Bukhari, S. A., Sloofman, L. G., Troy, J. M., Caetano-Anolles, D., Cash-Ahmed, A., Kent, M., Lu, X., Sanogo, Y. O., Weisner, P. A., Zhang, H., Bell, A. M., Ma, J., Sinha, S., Robinson, G. E., & Stubbs, L. (2014). Neuromolecular responses to social challenge: common mechanisms across mouse, stickleback fish, and honey bee. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(50), 17929-17934. https://doi.org/10.1073/pnas.1420369111

Rowe, L., Chenoweth, S. F., & Agrawal, A. F. (2018). The Genomics of Sexual Conflict. *Am Nat*, *192*(2), 274-286. https://doi.org/10.1086/698198

Schumer, M., Krishnakant, K., & Renn, S. C. (2011). Comparative gene expression profiles for highly similar aggressive phenotypes in male and female cichlid fishes (Julidochromis). *The Journal of Experimental Biology*, *214*(Pt 19), 3269-3278. https://doi.org/10.1242/jeb.055467

Shetty, S., Griffin, D. K., & Graves, J. A. (1999). Comparative painting reveals strong chromosome homology over 80 million years of bird evolution. *Chromosome Research*, *7*(4), 289-295. https://doi.org/10.1023/a:1009278914829

Smeds, L., Warmuth, V., Bolivar, P., Uebbing, S., Burri, R., Suh, A., Nater, A., Bures, S., Garamszegi, L. Z., Hogner, S., Moreno, J., Qvarnstrom, A., Ruzic, M., Saether, S. A., Saetre, G. P., Torok, J., & Ellegren, H. (2015). Evolutionary analysis of the female-specific avian W chromosome. *Nature Communications*, *6*(1), 7330. https://doi.org/10.1038/ncomms8330

Smiley, K. O., Munley, K. M., Aghi, K., Lipshutz, S. E., Patton, T. M., Pradhan, D. S., Solomon-Lane, T. K., & Sun, S. D. (2024). Sex diversity in the 21st century: Concepts, frameworks, and approaches for the future of neuroendocrinology. *Hormones and Behavior*, *157*, 105445. https://doi.org/10.1016/j.yhbeh.2023.105445

Soma, K. K., Schlinger, B. A., Wingfield, J. C., & Saldanha, C. J. (2003). Brain aromatase, 5 alpha-reductase, and 5 beta-reductase change seasonally in wild male song sparrows: relationship to aggressive and sexual behavior. *Journal of Neurobiology*, *56*(3), 209-221. https://doi.org/10.1002/neu.10225

Sun, D., Maney, D. L., Layman, T. S., Chatterjee, P., & Yi, S. V. (2019). Regional epigenetic differentiation of the Z Chromosome between sexes in a female heterogametic system. *Genome Research*, *29*(10), 1673-1684. https://doi.org/10.1101/gr.248641.119

Tosto, N. M., Beasley, E. R., Wong, B. B. M., Mank, J. E., & Flanagan, S. P. (2023). The roles of sexual selection and sexual conflict in shaping patterns of genome and transcriptome

variation. *Nature Ecology & Evolution*, *7*(7), 981-993. https://doi.org/10.1038/s41559-023-02019-7

Uebbing, S., Konzer, A., Xu, L., Backstrom, N., Brunstrom, B., Bergquist, J., & Ellegren, H. (2015). Quantitative Mass Spectrometry Reveals Partial Translational Regulation for Dosage Compensation in Chicken. *Molecular Biology and Evolution*, *32*(10), 2716-2725. https://doi.org/10.1093/molbev/msv147

van Nas, A., Guhathakurta, D., Wang, S. S., Yehya, N., Horvath, S., Zhang, B., Ingram-Drake, L., Chaudhuri, G., Schadt, E. E., Drake, T. A., Arnold, A. P., & Lusis, A. J. (2009). Elucidating the role of gonadal hormones in sexually dimorphic gene coexpression networks. *Endocrinology*, *150*(3), 1235-1249. https://doi.org/10.1210/en.2008-0563

Wade, J., Swender, D. A., & McElhinny, T. L. (1999). Sexual differentiation of the zebra finch song system parallels genetic, not gonadal, sex. *Hormones and behavior*, *36*(2), 141-152. https://doi.org/10.1006/hbeh.1999.1537

Wolf, J. B., & Bryk, J. (2011). General lack of global dosage compensation in ZZ/ZW systems? Broadening the perspective with RNA-seq. *BMC Genomics*, *12*(1), 91. https://doi.org/10.1186/1471-2164-12-91

Wright, A. E., Harrison, P. W., Zimmer, F., Montgomery, S. H., Pointer, M. A., & Mank, J. E. (2015). Variation in promiscuity and sexual selection drives avian rate of Faster-Z evolution. *Molecular Ecology*, *24*(6), 1218-1235. https://doi.org/10.1111/mec.13113

Wright, A. E., Zimmer, F., Harrison, P. W., & Mank, J. E. (2015). Conservation of Regional Variation in Sex-Specific Sex Chromosome Regulation. *Genetics*, *201*(2), 587-598. https://doi.org/10.1534/genetics.115.179234

Xu, L., & Zhou, Q. (2020). The Female-Specific W Chromosomes of Birds Have Conserved Gene Contents but Are Not Feminized. *Genes*, *11*(10). https://doi.org/10.3390/genes11101126

Yang, J., Ruan, H., Xu, W., & Gu, X. (2019). TreeExp2: An Integrated Framework for Phylogenetic Transcriptome Analysis. *Genome Biology and Evolution*, *11*(11), 3276-3282. https://doi.org/10.1093/gbe/evz222

Yang, Z. (1996). Among-site rate variation and its impact on phylogenetic analyses. *Trends Ecol Evol*, *11*(9), 367-372. https://doi.org/10.1016/0169-5347(96)10041-0

Zenil-Ferguson, R., McEntee, J. P., Burleigh, J. G., & Duckworth, R. A. (2023). Linking Ecological Specialization to Its Macroevolutionary Consequences: An Example with Passerine Nest Type. *Syst Biol*, *72*(2), 294-306. https://doi.org/10.1093/sysbio/syac083

Zhan, T., & Boutros, M. (2016). Towards a compendium of essential genes - From model organisms to synthetic lethality in cancer cells. *Crit Rev Biochem Mol Biol*, *51*(2), 74-85. https://doi.org/10.3109/10409238.2015.1117053

Zhao, D., McBride, D., Nandi, S., McQueen, H. A., McGrew, M. J., Hocking, P. M., Lewis, P.
D., Sang, H. M., & Clinton, M. (2010). Somatic sex identity is cell autonomous in the
chicken. *Nature*, *464*(7286), 237-242. https://doi.org/10.1038/nature08852

Zhou, Q., Zhang, J., Bachtrog, D., An, N., Huang, Q., Jarvis, E. D., Gilbert, M. T., & Zhang, G.
(2014). Complex evolutionary trajectories of sex chromosomes across bird taxa. *Science*,
346(6215), 1246338. https://doi.org/10.1126/science.1246338

Zimmer, F., Harrison, P. W., Dessimoz, C., & Mank, J. E. (2016). Compensation of Dosage-
Sensitive Genes on the Chicken Z Chromosome. *Genome Biology and Evolution*, *8*(4),
1233-1242. https://doi.org/10.1093/gbe/evw075
808