Plasma membrane accessible cholesterol is regulated by ACC1 and lipid droplets

Kalani M. Wijesinghe[1], Chai-wan Kim[2], Emily O. Schad[1], Shuo Li[1], Summer Chen[1], Erin Takeshima[1], Chandni B. Khandwala[1], Desiree Tillo[3], Andres M. Lebensohn[4], James A. Olzmann[5], Rajat Rohatgi[1], Maia Kinnebrew[1,6]

[1] Departments of Biochemistry and Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA
[2] Center for Human Nutrition and Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046, USA
[3] Center for Cancer Research Genomics Core, National Cancer Institute, National Institutes of Health, NIH, Building 41, RM D701, Bethesda, MD 20892, USA.
[4] Laboratory of Cellular and Molecular Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, NIH, Building 37, RM 2056C, Bethesda, MD 20892, USA.
[5] Department of Molecular and Cell Biology, University of California at Berkeley, Berkeley, CA, USA and Department of Nutritional Sciences and Toxicology, University of California at Berkeley, Berkeley, CA 94720, USA.
[6] Lead contact

Abstract

Proper maintenance of plasma membrane (PM) cholesterol is essential for diverse processes ranging from animal development to pathogen evasion. Despite decades of study, the mechanisms governing cellular cholesterol regulation are incomplete. Using genome-wide screens we find that ACC1, the rate-limiting enzyme in fatty acid biosynthesis, regulates PM cholesterol transport. ACC1 loss causes a ~10-fold increase in PM accessible cholesterol in cells and mice. Mechanistically, we find that ACC1 regulates lipid droplet (LD) catabolism, and LDs are intimately tied to PM accessible cholesterol levels since reductions or elevations in their numbers block or promote cholesterol trafficking, respectively. Furthermore, LDs are required for cholesterol trafficking induced by 25-hydroxycholesterol, a modulator of inflammation and an interferon-stimulated second messenger that protects cells from pathogen invasion. This work identifies an unrecognized role for ACC1 and LDs in cholesterol regulation, which has implications for diseases where LD numbers are altered, from metabolic syndromes to neurodegeneration.

Keywords: Cholesterol, accessible cholesterol, lipid droplet, plasma membrane, membrane trafficking, ACC1, ATGL, SREBP2, ACAT1 and ACAT2, AMPK, 25-hydroxycholesterol, triacylglycerol, cholesterol esters

Introduction

Cholesterol is a vastly important molecule: it makes up 40% of all lipids in the plasma membrane (PM) through the coordinated effort of nearly 100 proteins that regulate its

synthesis, transport and metabolism[1,2]. While being vital for cell and organismal health, it is also
linked to numerous human diseases, including cardiovascular disease, metabolic syndrome,
cancer and neurodegeneration. Despite a rich history of investigation dating to the 1700s and
multiple Nobel prizes awarded to its study, we lack a complete understanding of how cholesterol
is regulated[3]. For example, only in the last 15 years has a pivotal concept come to light: while
the majority of cholesterol plays structural roles in the PM, ~2% of PM cholesterol has high
chemical activity[4–7]. This pool of cholesterol, often referred to as *active* or *accessible*
cholesterol, can dynamically bind to proteins and drive cell signaling events. The complete
importance of accessible cholesterol in vivo is unknown, but thus far it has been shown to
control de novo cholesterol synthesis, viral and bacterial pathogenesis, and the Hedgehog
signaling pathway[8–13]. We hypothesize that significant breakthroughs related to the study of
cholesterol in human diseases have been obscured because the majority of cholesterol
research (99.98% of published articles) did not measure the accessible cholesterol pool. This
study seeks to redefine our understanding of cholesterol by investigating this understudied yet
functionally essential accessible cholesterol pool.
Accessible cholesterol is continually transported from the PM to the Endoplasmic
Reticulum (ER) where its levels are sensed to control de novo cholesterol synthesis. When ER
cholesterol is high, SCAP and SREBP2 are retained in the ER and cholesterol synthesis is
blocked. When ER cholesterol is low, SCAP escorts SREBP2 to the Golgi where it is
proteolytically cleaved into a transcriptionally active fragment that drives the expression of
cholesterol biosynthesis genes[12,14–17]. A long-standing mystery is how changes in PM accessible
cholesterol are conveyed to the cholesterol-sensing machinery in the ER. Additionally, are there
proteins that sense and control the amount of PM accessible cholesterol directly?
The high accessible cholesterol content of vertebrate PMs has been exploited by both
bacterial and viral pathogens for their entry into cells[10,13,18,19]. To counteract this, the immune
system has evolved a mechanism to evade pathogens where PM accessible cholesterol levels
are rapidly depleted. Interferon gamma signaling induces the expression of the enzyme
cholesterol 25-hydroxylase (CH25H), which converts cholesterol to 25-hydroxycholesterol
(25HC). 25HC diffuses across the cell membrane to surrounding cells, triggering rapid (~1-6
hours) PM cholesterol depletion by allosterically activating the ER localized proteins ACAT1 and
ACAT2 (genes *SOAT1* and *SOAT2*). ACAT1/2 esterify cellular cholesterol, which is then stored
in lipid droplets[20–22]. 25HC also reduces PM cholesterol on longer timescales that require the
activation of transcriptional programs: 25HC elevates the expression of ABC transporters by
activating LXR and it also activates INSIG to block new cholesterol synthesis[23–27]. How
cholesterol is transported out of the PM in response to 25HC is unknown.
GRAMD1B is known to regulate accessible cholesterol-specific trafficking from the PM to
the ER; it localizes in the ER and moves cholesterol down its concentration gradient at PM-ER
contact sites[28–31]. Knockout of GRAMD1B (and triple knockout of GRAMD1A, GRAMD1B and
GRAMD1C) does not affect cell viability, demonstrating that additional pathways for cholesterol
transport out of the PM must exist. A subset of cytoplasmically localized START-domain
containing proteins (StarD4, StarD5, and StarD6) have also been shown to bind cholesterol and
control its intracellular trafficking; their relative contribution to PM cholesterol homeostasis has
not been systematically investigated[32,33]. Finally, cholesterol can be internalized from the PM
through vesicular transport, although there is no known endocytic pathway dedicated

specifically to maintaining cholesterol homeostasis[34–37]. In summary, we lack a complete picture
of how accessible cholesterol is regulated in cells, which obscures our understanding of its
function for cell and organismal physiology. We thus set out to solve this foundational mystery
using an unbiased genetic screen to identify new genes controlling PM accessible cholesterol
homeostasis. From here forward in this manuscript, unless specifically stated otherwise, any
reference to cholesterol refers to accessible cholesterol.
92

Results
Random-insertion mutagenesis screens identify new cholesterol regulatory genes.
To test the known pathway of PM to ER cholesterol transport, we knocked out GRAMD1A,
GRAMD1B and GRAMD1C (*GRAMD1A/B/C* TKO) in human haploid HAP1 cells (**Extended**
data Fig. 1a)[28–31]. *GRAMD1A/B/C* TKO cells showed 1.2-fold elevated PM cholesterol,
consistent with a defect in cholesterol trafficking out of the PM (**Fig. 1a**). However, when treated
with 25HC, cholesterol was depleted from the PM with similar kinetics to wild-type (WT) cells
(**Fig. 1b**). These results were indistinguishable to GRAMD1B single KO cells (**Extended data**
Fig. 1b-**d**). 25HC can also reduce PM cholesterol levels by activating LXR and increasing the
expression of ABC transporters that export cholesterol from the cell[26]. Overnight treatment with
an LXR inhibitor strongly reduced ABCA1 transcript levels and slightly elevated PM cholesterol
levels; however, it did not block PM cholesterol depletion induced by 25HC (**Fig. 1c**, **d and**
Extended data Fig. 1e-**g**). These data demonstrate that when cells are stimulated with 25HC,
cholesterol can exit the PM in a GRAMD1A/B/C- and LXR-independent manner. This raises the
question: how does cholesterol exit the PM in response to 25HC (**Fig. 1e**)?
To find genes that regulate PM cholesterol, we devised a new genome-wide screening
strategy. We reasoned that strong regulators of cellular cholesterol are likely to be required for
cell viability, and therefore avoided CRISPR knockout screening. Instead, we used random-
insertion mutagenesis to generate a library of ~100 million cells in the human haploid HAP1 cell
line. This mutational strategy can generate hypomorphic alleles by truncating proteins or by
reducing the transcriptional output of genes. Next, we stained live cells at the outer leaflet of the
PM with the accessible cholesterol binding protein ALOD4 fused to mNeonGreen (mNG-
ALOD4) to enrich cells whose cholesterol levels were altered (**Fig. 1f**) [38,39]. Using this strategy,
we performed two screens probing distinct biological processes. We first treated cells with the
NPC1 inhibitor U18666A for 20 hours, which reduces PM cholesterol by blocking low density
lipoprotein (LDL)-derived cholesterol export from lysosomes. We then sorted for cells with the
top 10% of mNG-ALOD4 fluorescence to identify mutants that rescue PM cholesterol in the
absence of NPC1 function. In a second screen, we treated cells with 25HC for 6 hours and then
sorted for cells with the top 10% of mNG-ALOD4 fluorescence to identify mutants in which
cholesterol transport in response to 25HC was blocked. Strikingly, both screens identified ACC1
(*ACACA* gene) as the top hit by a large margin (**Fig. 1g**, **h**). ACC1 has been the subject of
extensive research due to its role in human diseases, but it has not been linked to cholesterol
regulation in cells; this motivated us to decipher this mystery.
ACC1 catalyzes the rate limiting step in fatty acid biosynthesis, converting acetyl-CoA
into malonyl-CoA, which is then elongated by fatty acid synthase (FASN) to produce long-chain
saturated fatty acids[40]. Interestingly, the majority of ACC1 insertions enriched in the screens

Fig. 1: Random insertion mutagenesis screens identify new cholesterol regulatory genes.

a-d, Plasma membrane (PM) cholesterol was measured by mNeonGreen-ALOD4 (mNG-ALOD4) staining followed by flow cytometry. Wild type (WT) HAP1 cells are compared to GRAMD1A, GRAMD1B, and GRAMD1C triple knockout HAP1 cells (*GRAMD1* TKO) either left untreated (**a**) or treated with 4 µM 25-hydroxycholesterol (25HC) for the times indicated (**b**). WT cells were treated with or without LXR inhibitor (LXRi; 1 µM GSK2033) for 16 hrs in 5% FBS and either left untreated (**c**) or treated with 25HC for the times indicated (**d**). **e**, Schematic showing cholesterol trafficking from the PM to the ER membrane where it binds to proteins including SCAP and ACAT1 or ACAT2 (ACAT). ACAT esterifies cholesterol with a fatty acid (FA), enabling it to be stored in lipid droplets (LD). Cholesterol is transported to the PM from the Golgi[89]. **f**, Schematic of genetic screening pipelines. The mutant cell library was treated with either of NPC1i (1 µM U18666A) for 20 hrs or 4 µM 25HC for 6 hrs and then cells were stained with mNG-ALOD4. The top 10% of mNG-ALOD4 fluorescent cells were isolated for sequencing. g,h, Screen results showing enriched genes with an FDR-corrected *p*-value of less than 2E-4 for cells treated with NPC1i (**g**) or 25HC (**h**). IGTIOB is a measure of the inactivating potential of the mapped insertions, and circle size shows the number of insertions for each gene. **i,j**, mNG-ALOD4 flow cytometry analysis of PM cholesterol in WT or *ACC1* KO HAP1 cells treated with or without 1 µM of NPC1i for 20 hrs (**i**) or WT and *NPC1* KO HAP1 cells treated with or without ACC1i (30 µM Firsocostat) for 16 hrs (**j**). **k-m**, Cells were left untreated (**k**) or treated for 16 hrs with 30 µM ACC1i and then treated with 4 µM 25HC for the times indicated (**l** and **m**). The amount of mNG-ALOD4 bound to the PM was measured by flow cytometry. Data is normalized to cells not treated with 25HC for each condition, and each circle is the mean of three biological replicates; error bars are the standard deviation. Statistical tests compare two conditions at the same time point of 25HC treatment (**d** and **k-m**). Statistical significance was determined by Student's t-test with a Welch's correction (**a-d** and **j-m**) or an Ordinary one-way ANOVA (**i**). Exact *p*-values are reported in the Methods.

mapped to intron 52, which truncates ACC1 at its carboxyltransferase domain (**Extended data**
Fig. 1h, **i**). Using CRISPR to generate a series of ACC1 truncations, we found that an exon 52
truncation behaves similarly to an exon 11 truncation lacking both catalytic domains, and is thus
referred to as *ACC1* KO (**Extended data Fig. 1j-l**). Surprisingly, ACC1 truncated at exon 3 was
lethal to cells, suggesting that ACC1 early exons have functionality beyond malonyl-CoA
synthesis, perhaps through a scaffolding function (**Extended data Fig. 1m**)[41]. This finding
highlights the strength of random insertion mutagenesis: since CRISPR libraries have sgRNAs
targeting early exons, they would fail to identify ACC1 as a hit due to cell lethality.
The lysosomal cholesterol transporters NPC1 and NPC2 are mutated in Niemann-Pick
disease type C (NPC disease), an incurable fatal syndrome. We and others have shown that
NPC1 inhibition or knockout causes reduced PM cholesterol levels (**Extended data Fig. 1n-**
p)[42]. Since the cell biological changes that drive NPC disease symptoms remain an area of
active research, treatments that restore cells to homeostasis could be investigated as potential
therapies. Remarkably, *ACC1* KO cells treated with U18666A had PM cholesterol levels
indistinguishable from WT untreated cells (**Fig. 1i**). To more closely mirror a chronic disease
state, we treated *NPC1* KO cells with the ACC1 inhibitor Firsocostat, a drug tested in human
clinical trials for the treatment of metabolic dysfunction-associated steatotic liver disease
(MASLD). Remarkably, Firsocostat elevated PM cholesterol in both *NPC1* KO HAP1 cells and
mouse embryonic fibroblasts; the amount of Firsocostat could be reduced so that PM
cholesterol abundance more closely matches control levels. This demonstrates that ACC1
inhibition does not raise PM cholesterol levels through a mechanism involving NPC1 (**Fig. 1j**
and **Extended data Fig. 1p**). We thus suggest that ACC1 inhibition could be a testable
treatment for NPC disease.
Cells protect themselves from pathogen invasion by transiently synthesizing 25HC,
which drives a ~10-fold reduction in PM cholesterol (**Fig. 1b**, **d**). Under non-infection conditions,
elevated 25HC promotes inflammation linked to elevated stroke risk, osteoarthritis,
atherosclerosis, and fibrosis[43–46]. Therefore, understanding how 25HC drives PM cholesterol
changes, and reducing these changes for healthy individuals, could have broad clinical impacts.
Treatment of *ACC1* KO cells with 25HC revealed a significant defect in cholesterol depletion
compared to WT cells, with >60% of cholesterol remaining at the PM after 6 hours (**Fig. 1k**).
The ACC1 inhibitor Firsocostat also prevented PM cholesterol depletion induced by 25HC in
HAP1, 293T and 3T3 cells, demonstrating that this response is conserved across cell types
from different lineages (**Fig.1l**, **m**)[47]. Given the broad importance of 25HC in human disease, we
set out to understand how ACC1 loss can block 25HC-mediated cholesterol trafficking.
164
ACC1 functions independently of ACAT1/2 to maintain plasma membrane cholesterol
levels.
Using both microscopy and flow cytometry, ACC1 loss caused a striking increase in PM
cholesterol in untreated cells (**Fig. 2a**, **b** and **Extended data Fig. 1k**, **l**). This effect was
observed in multiple cell types and with structurally distinct ACC1 inhibitors, including the
dimerization inhibitor Firsocostat and the carboxyltransferase domain active site inhibitor CP-
640186 (**Fig. 2c**, **d**)[48,49]. ACC1 loss also increased cell death caused by the cholesterol-
dependent cytolysin PFO* (**Fig. 2e**). PFO* binds to accessible cholesterol and above 4°C it
oligomerizes to generate pores in the PM[10,50]. We hypothesized that ACC1 could raise PM

cholesterol accessibility by reducing sphingomyelin levels since sphingomyelin binds and
sequesters cholesterol in the PM. Using mass spectrometry, a sphingomyelin-binding probe,
and a sphingomyelinase assay, we showed that ACC1 loss does not change sphingomyelin
levels (**Extended data Fig. 2a-d**). Altogether, our data suggests that ACC1 is critical for
maintaining PM cholesterol in untreated cells.
25HC-driven PM cholesterol depletion was previously shown to require ACAT1 and
ACAT2; these proteins esterify cholesterol with a fatty acid, promoting its storage in lipid
droplets[20]. We hypothesized that ACC1 loss reduces the pool of available fatty acids that are
used by ACAT1/2 to make cholesterol esters (CEs) (**Fig. 2f**). If cholesterol cannot be stored, PM
cholesterol may rise as observed in *ACC1* KO cells. To test this, we treated cells with the
ACAT1/2 inhibitor SZ58-035, which blocked CE formation induced by 25HC (**Extended data**
Fig. 2e)[51]. ACAT1/2 inhibition reduced PM cholesterol depletion induced by 25HC, however the
magnitude of the effect was small compared to previous reports and also less than what was
observed for ACC1 inhibition (**Extended data Fig. 2f** and **Fig. 1k**)[20]. Therefore, we generated
ACAT1/2 double knockout (DKO) cells with a dual-sgRNA CRISPR strategy (**Fig. 2g**). As
expected, *ACAT1/2* DKO cells showed undetectable levels of CEs (**Fig. 2h**). Similarly to SZ58-
035 treated cells, *ACAT1/2* DKO cells showed reduced cholesterol depletion in response to
25HC compared to WT cells, however, in contrast to previous reports, a significant fraction of
PM cholesterol was still depleted (**Fig. 2i**)[20].
If ACC1 and ACAT1/2 are acting in the same pathway, then knocking out ACC1 and
ACAT1/2 together should not have a combinatorial effect. We treated *ACAT1/2* DKO cells with
ACC1 inhibitor, which caused a dramatic increase in PM cholesterol in the presence of 25HC
(**Fig. 2j**). Additionally, ACAT1/2 inhibition further blunted the effect seen in *ACC1* KO cells
(**Extended data Fig. 2g**). Finally, we made a triple knockout (TKO) cell line lacking ACC1,
ACAT1 and ACAT2, which had an additive effect compared to knocking out either ACC1 or
ACAT1/2 alone (**Fig. 2k**, **l**). Moreover, the PM cholesterol levels of *ACAT1/2* DKO cells were not
significantly different from WT cells in agreement with previous reports[20]. Inhibition of ACC1 in
ACAT1/2 DKO cells led to a dramatic increase in PM cholesterol (**Fig. 2m**). These epistasis
experiments cleanly demonstrate that ACC1 acts through a distinct pathway from the ACAT1/2
proteins.
25HC can deplete *total cellular cholesterol* through multiple known pathways: 1)
LXR/ABC transporter-driven cholesterol efflux, 2) INSIG activation leading to reduced
cholesterol synthesis, 3) ACAT1/2-mediated cholesterol esterification, and 4) by elevating the
expression of oxysterol-producing enzymes like CH25H[20,26,52]. With the exception of LXR-driven
cholesterol efflux, which we previously showed had minor effects (**Fig. 1d**), the remaining
mechanisms all require cholesterol to leave the PM in order to reveal a depletion. We observed
a decrease in total cellular cholesterol upon 25HC treatment in WT cells, but not in *ACC1* KO
cells, further suggesting that ACC1 loss traps cholesterol at the PM (**Fig. 2n**). Additionally,
blocking HMGCR activity with a statin had minimal effects on PM cholesterol after 6 hours,
suggesting that the reduced PM cholesterol we observed is not due to reduced synthesis
(**Extended data Fig. 2h**).

Fig. 2: ACC1 functions independently of ACAT1/2 to maintain plasma membrane cholesterol levels.

a, Fluorescence microscopy image showing wild-type (WT) 3T3 cells left untreated or treated with 0.3 mM MβCD:cholesterol or ACC1i (30 μM Firsocostat) and stained with mNG-ALOD4 and DAPI. Scale bar is 10 microns. **b-d**, Cell lines were left untreated (**b**) or treated with 30 μM Firsocostat (**c**) or 20 μM CP-640186 for 16 hrs (**d**) and then stained with mNG-ALOD4 and analyzed by flow cytometry. **e**, Cells were treated with 4 μM 25HC for 4 hrs and then media was replaced with PFO* for 30 minutes at 37°C before cell viability was measured with an XTT assay. **f**, Schematic of cholesterol ester synthesis. ACAT1 and ACAT2 (ACAT) enzymes transfer a fatty acid from acyl-CoA to cholesterol, forming cholesterol esters. **g**, Western blot (ACAT1) and PCR (*ACAT2*) analysis of *ACAT1/2* DKO clones. **h**, WT and *ACAT1/2* DKO cell lines were treated with 4 μM 25HC for 6 hrs and the esterified cholesterol was quantified using the Amplex Red assay. **i,j**, Cells were left untreated (**i**) or treated for 16 hrs with ACC1i (30 μM Firsocostat) (**j**); cells were then treated with 4 μM 25HC for the times indicated and then the amount of mNG-ALOD4 bound to the PM was measured by flow cytometry. **k**, Western blot (ACAT1 and ACC1) or PCR (*ACAT2*) analysis of *ACC1/ACAT1/2* TKO clones**. l**, Cells were treated with 4 μM 25HC for the indicated times and then mNG-ALOD4 staining was measured by flow cytometry. **m**, mNG-ALOD4 flow cytometry analysis of PM cholesterol in WT and *ACAT1/2* DKO HAP1 cells left untreated or treated with ACC1i (30 μM Firsocostat) for 16 hrs. **n,** Quantification of total cellular free (unesterified) cholesterol in WT HAP1 cells or *ACC1* KO HAP1 cells treated with or without 4 μM 25HC for 8 hrs. Cells (**b**, **d**, and **m**) were grown in lipoprotein depleted serum media for 16 hours prior to treatments. For each condition (**i,j** and **l**), data is normalized to cells not treated with 25HC, and each data point represents the average of three biological replicates, with error bars denoting the standard deviation. Statistical significance was determined by a Student's t-test with a Welch's correction (**b**,**c**,**h-j**, **l** and **n**) or an Ordinary one-way ANOVA (**d,** and **m**). Exact *p*-values are found in the Methods.

Lipid droplet catabolism increases plasma membrane cholesterol.

Interestingly, ACC1 inhibition caused rapid changes in PM cholesterol levels. Treatment with Firsocostat resulted in observable effects in 2 hours in HAP1 cells and in 15 minutes in 3T3 cells (**Fig. 3a** and **Extended data Fig. 3a**). These changes are not consistent with the induction of a transcriptional program like SREBP2 signaling, which should show a lag as new genes are synthesized. Consistent with SREBP2 independent effects, treatment of cells with the HMGCR inhibitor Lovastatin did not reverse the effect of ACC1 loss on PM cholesterol (**Extended data Fig. 3b**, **c**)[53]. How does ACC1 loss cause elevated PM cholesterol?

We used immunofluorescence imaging to observe ACC1 in cells and found the formation of ring-like puncta. Co-staining with Bodipy 493:503 revealed that ACC1 localizes on the surface of lipid droplets (LD), which was further confirmed through density gradient centrifugation (**Fig. 3b** and **Extended data Fig. 3d**). ACC1 was previously identified in a LD-proximity labeling study, though its localization at LDs was not further explored[54,55]. We found that ACC1 loss-of-function caused a striking reduction in LD numbers, suggesting that ACC1 activity controls LD abundance[56]. Overnight treatment with the ACC1 inhibitor Firsocostat decreased the median number of LDs per cell from ~six to zero and *ACC1* KO cells lacked LDs (**Fig. 3c** and **Extended data Fig. 3e**).

LDs have a hydrophobic core made of triacylglycerol (TAG) and CE; the TAG/CE ratio is cell-type dependent with TAG making up the major component in most cell types[57–59]. The abundance of LDs in a cell is determined by a balance of synthesis and degradation. We observed that blocking LD synthesis by inhibiting the TAG-synthesizing enzymes DGAT1 and DGAT2 did not eliminate LDs after 24 hours (**Fig. 3d** and **Extended data Fig. 3f**). These drug treatments were robust since they blocked new LD formation induced by exogenous oleic acid (OA) addition. This suggests that blocking synthesis does not cause rapid changes in LD numbers. In contrast, inhibition of ACC1 caused a 50% depletion in LD numbers after ~6 hours in lipoprotein-depleted serum (LDS) media (**Fig. 3e**). The same assay performed in lipoprotein-replete 5% fetal bovine serum (FBS) showed slower kinetics, with 50% depletion after ~10 hours (**Fig. 3f**). We therefore propose that ACC1 inhibition triggers LD catabolism, which is further explored in Figure 4. We hypothesize that catabolism occurs more robustly in LDS media because catabolic enzymes are more active under conditions of nutrient deprivation. For example, the rate limiting enzyme in TAG breakdown, adipose triacylglycerol lipase (ATGL), is activated by AMPK when cellular energy stores are low[60]. LD numbers showed a striking inverse correlation with PM cholesterol levels in both LDS and FBS, leading us to speculate that LD loss could cause an increase in PM cholesterol (**Fig. 3e**, **f**).

If LDs are indeed required for PM cholesterol trafficking, then we should be able to demonstrate this through orthogonal mechanisms. Knockout of FASN, the enzyme acting downstream of ACC1 to generate fatty acids, showed reduced LD numbers and a ~2.5-fold increase in PM accessible cholesterol (**Extended data Fig. 3g-i**). *FASN* KO also caused a defect in cholesterol trafficking in response to 25HC (**Extended data Fig. 3j**). The effects of *FASN* KO were overall smaller than those measured for *ACC1* KO, which we speculate was due to poor cell health; for example, these cells required growth in 20% FBS-containing media in order for survival. High serum likely provides significant exogenous fatty acids, which may mask the full impact of FASN loss. The FASN inhibitor C75 caused no change to PM cholesterol levels, and also did not induce LD catabolism after 48 hours (**Extended data Fig. 3k**). This

260 suggests that in contrast to ACC1 inhibition, FASN inhibition does not trigger LD catabolism;
261 interestingly, FASN did not colocalize with LDs (**Extended data Fig. 3l**).
262

Fig. 3: Lipid droplet catabolism increases plasma membrane cholesterol.

a, Plasma membrane (PM) cholesterol was measured by mNG-ALOD4 staining followed by flow cytometry in WT HAP1 cells treated with ACC1i (30 µM Firsocostat) for the indicated times. **b,c**, Microscopy images of HAP1 cells left untreated or treated with ACC1i (30 µM Firsocostat), and then stained with ACC1 monoclonal antibody (**b**) or 647-conjugated ACC1 polyclonal antibody (**c**), Bodipy 493/503 to measure lipid droplets (LDs) and DAPI (blue). Scale bar is 5 microns. **d**, Schematic showing that enzymes involved in the synthesis of fatty acyl-CoAs, which are then converted into triacylglycerol (TAG), cholesterol esters (CE), sphingolipids (SL) and phospholipids (PL). TAG and CE are the main components of the LD core (green oval). Inhibitors targeting enzymes in this process are shown in gray. **e-g** mNG-ALOD4 flow cytometry analysis of PM cholesterol (blue curves) and the corresponding fluorescence microscopy quantification of the number of LDs per cell (purple curves) in HAP1 cells grown in 5% LDS (**e**) or 5% FBS (**f,g**) and then treated with ACC1i (30 µM Firsocostat) (**e,f**) or ACSLi (5 µM Triacsin-C) (**g**). LDs were quantified by taking the mean number of LDs per cell in four fields of view with ~75 cells per field. **h,i**, mNG-ALOD4 flow cytometry analysis of PM cholesterol in WT HAP1 cells treated with or without ACSLi (5 µM Triacsin-C) for 16 hrs in 5% LDS, followed by treatment with 4 µM 25HC (**h**) or 30 µM ACC1i (**i**) for the indicated times. **j,k**, LD quantification (**j**) and mNG-ALOD4 flow cytometry analysis of PM cholesterol (**k**) of WT HAP1 cells treated with or without 30 µM ACC1i or 5 µM ACSLi for 16 hrs, followed by 200 µM oleic acid treatment for 6 hrs in 5% FBS. Each flow cytometry data point (**e-i**) represents the mean of three biological replicates, with error bars indicating the standard deviation. Values were normalized to the untreated control at the corresponding time point. Statistical significance was determined by Ordinary one-way ANOVA (**a**) or Student's t-test with a Welch's correction (**h-k**). Exact *p*-values are found in the Methods.

The most common method to acutely reduce LD numbers is treatment with the acyl-CoA
synthetase long-chain (ACSL1,3-6 are henceforward collectively called ACSL) inhibitor Triacsin-
C, which both prevents LD synthesis and stimulates LD catabolism by starving the cell of fatty
acyl-CoA (**Fig. 3d**)[61,62]. Remarkably, ACSL inhibition led to an ~8-fold increase in PM cholesterol
levels that inversely correlated with LD levels (**Fig. 3g**). ACSL inhibition showed a maximal
effect on PM cholesterol after ~18 hours; at 24 hours, PM cholesterol started to decrease and at
48 hours cell death occurred. Similarly to ACC1 loss, ACSL loss also robustly blocked PM
cholesterol depletion induced by 25HC (**Fig. 3h**). This provides additional evidence that LDs
mediate cholesterol trafficking out of the PM in response to 25HC. To test whether ACC1 and
ACSLs were acting in the same pathway, we treated cells overnight with Triacsin-C to eliminate
LDs, and then performed a Firsocostat treatment time course. In cells lacking LDs, ACC1
inhibition had no added effect on PM cholesterol levels (**Fig. 3i**). This result suggests that the
differences we observe upon ACC1 inhibition arise from changes in LD numbers; if LDs are
absent to begin with, ACC1 does not cause further changes.
If trafficking of cholesterol out of the PM requires LDs, then the addition of exogenous
OA should reverse the ACC1 phenotype. Treatment with 6 hours of OA elevated the number of
LDs in both control cells and ACC1 inhibited cells (**Fig. 3j**). Importantly, ACSL inhibition blocked
LD formation induced by OA treatment, since ACSL enzymes are required to convert fatty acids
into fatty acyl-CoAs, the precursors for both TAG and CE. OA reversed the high PM cholesterol
levels of ACC1 inhibited cells but not ACSL inhibited cells (**Fig. 3k**). The finding that OA addition
to cells can reverse the effects of ACC1, but not ACSL, provides strong evidence that LDs are
the crucial mediator of PM cholesterol trafficking.
286
ACC1 inhibition triggers lipid droplet catabolism through ATGL.
Since both ACC1 and ACSL loss raised PM cholesterol levels, we hypothesized that cholesterol
homeostasis is controlled by a common product of these enzymes, which could include fatty
acyl-CoAs or LDs made of TAG (**Fig. 4a**). ATGL is the rate limiting enzyme in LD catabolism,
hydrolyzing TAG to generate diacylglycerol (DAG) and a fatty acid[63]. To test if TAG breakdown is
required for the PM cholesterol changes seen upon ACC1 inhibition, we generated CRISPR
knockouts of ATGL (**Fig. 4b**). *ATGL* KO cells showed similar numbers of LDs compared to WT
cells, although they were significantly larger in size (**Fig. 4c**, **d**). When ACC1 was inhibited, WT
cells showed a ~50% depletion in LD numbers after six hours. In contrast, *ATGL* KO cells
showed no change in LD numbers with six hours of ACC1 inhibition (**Fig. 4e-g**). This suggests
that ACC1 inhibition triggers ATGL-dependent LD catabolism.
ACC1 inhibition leads to a rapid increase in PM cholesterol in WT cells. Importantly, the
effects of ACC1 inhibition were largely blocked in *ATGL* KO cells, which we propose is due to
the stable LD population (**Fig. 4h**). The newly characterized human-specific inhibitor of ATGL,
NG-497, also blocked PM cholesterol changes induced by ACC1 loss of function (**Fig. 4i**)[64].
ATGL inhibition also reversed the defect in 25HC-induced cholesterol transport observed in
ACC1 inhibited cells (**Extended data Fig. 4a**). These data demonstrate that ACC1 inhibition
causes PM cholesterol changes by depleting cells of LDs, and not through another mechanism.
When TAG is hydrolyzed by ATGL, a fatty acid is liberated, which can then be converted into a
fatty acyl-CoA by ACSL enzymes (**Fig. 4a**). The ability of *ATGL* KO to reverse the ACC1

phenotype argues against fatty acyl-CoA's playing a role in PM cholesterol homeostasis since
ATGL loss further reduces fatty acyl-CoA levels.
We hypothesized that PM cholesterol rises upon LD catabolism because CEs are
hydrolyzed, and subsequently the newly liberated (unesterified) cholesterol traffics to the PM. To
test this, we imaged LDs in *ACAT1/2* DKO cells, which had slightly reduced LD numbers
compared to WT cells, though LD size was unaltered (**Extended data Fig. 4b-d**). ACC1
inhibition induced LD loss in *ACAT1/2* DKO cells similarly to WT cells (**Extended data Fig. 4e**).
Finally, Firsocostat treatment raised PM cholesterol levels in *ACAT1/2* DKO cells (lacking CE),
although with slightly reduced kinetics compared to WT cells (**Extended data Fig. 4f**). These
results suggest that CE hydrolysis partially contributes to the *initial* increases in PM cholesterol
seen upon ACC1 inhibition. Our data showing that *ACC1* KO leads to a sustained increase
(over weeks of passage) in PM cholesterol suggests that LDs loss has long lasting effects on
cellular cholesterol (**Fig. 2b**, **Extended data Fig. 1k**). Cells lacking ACC1 are not able to return
to the WT homeostatic set point of PM cholesterol.

Fig. 4: ACC1 inhibition triggers lipid droplet catabolism through ATGL.

a, Schematic illustrating the formation of fatty acyl-CoAs from de novo synthesis of fatty acids (top) or from the breakdown of triacylglycerol (TAG) into diacylglycerol and a fatty acid by ATGL (bottom). ATGL localizes at the lipid droplet (LD) surface. **b**, Western blot analysis of *ATGL* KO HAP1 cells compared to a wild-type (WT) control. **c,d**, Quantification of LD numbers per cell (**c**) or LD area (**d**) in WT and *ATGL* KO cells. **e-g** Microscopy images (**e,f**) and quantification (**g**) of LDs in WT (**e**) and *ATGL* KO (**f**) HAP1 cells treated with or without ACC1i (30 µM Firsocostat) for 6 hrs. Cells were stained with Bodipy 493/503 and DAPI (blue). Scale bar is 5 microns. Each data point is the total number of LDs per cell in a single field of view, with ~75 cells per field (**c** and **g**). **h**, **i**, mNG-ALOD4 flow cytometry analysis of PM cholesterol in WT and *ATGL* KO cells (**h**) or WT cells treated with or without ATGLi (50 µM NG-497) (**i**), followed by treatment with ACC1i (30 µM Firsocostat) for the indicated times. All experiments were carried out in cells pre-cultured for 16 hours in lipoprotein depleted serum. Data points represent the average of three biological replicates, with error bars indicating the standard deviation. All values were normalized to the untreated control (**h,i**). Statistical significance was determined by a Student's t-test with a Welch's correction (**c,d** and **g-i**). Exact *p*-values are found in the Methods.

322
Increasing lipid droplet abundance reduces plasma membrane cholesterol.
What is the consequence of increasing LD numbers? We found that the ATGL inhibitor NG-497
increased LD numbers in cells (**Fig. 5a**, **b** and **Extended data Fig. 5a**). This contrasts the *ATGL*
KO cells which showed enlarged but not more numerous LDs; it is not uncommon for knockout
cell lines to show different effects compared to drug treatments since cells will adapt to the
chronic loss of a gene. Interestingly, ATGL inhibition caused a ~7-fold decrease in PM
cholesterol levels (**Fig. 5c** and **Extended data Fig. 5b**). This data suggests that elevating LD
numbers reduces PM cholesterol. Combined treatments of the ATGL inhibitor with an ACC1 or
ACSL inhibitor showed intermediate effects on LD numbers at 24-hour time points. In all cases,
PM cholesterol showed a clear inverse relationship with LD numbers. We next sought to
increase LD numbers through an orthogonal mechanism.

AMPK plays central roles in cellular energy metabolism by controlling the balance
between anabolic and catabolic processes. When cellular energy stores are low, AMPK inhibits
anabolic proteins such as ACC1, HMGCR and RAPTOR, a member of the mTORC1 complex.
Conversely, AMPK activates catabolic enzymes like ATGL, CPT1 and ULK1, which, together
with mTORC1 inhibition, promotes autophagic flux (**Fig. 5d**)[65,66]. We therefore asked whether
AMPK inhibition would change LD numbers by activating ACC1 and simultaneously inhibiting
ATGL. Treatment of cells with the AMPK inhibitor Dorsomorphin (also named Compound C)
reduced the levels of phosphorylated inactive ACC1, which is expected to increase the active
ACC1 pool (**Fig. 5e**)[67]. Consistent with ACC1 activation leading to elevated fatty acid synthesis,
AMPK inhibition led to a ~2-fold increase in the number of LDs per cell (**Fig. 5f**, **g**). This effect
appeared to be driven by ACC1 activity rather than ATGL inactivation, because treatment with
the ACC1 inhibitor Firsocostat reversed the changes in LD numbers.

Elevating LD numbers by inhibiting AMPK caused a ~5-fold decrease in PM accessible
cholesterol levels (**Fig. 5h**). This provides further evidence that elevating LD abundance can
reduce PM cholesterol, possibly by enhancing its transport out of the PM. Alternatively, ACC1
activation could reduce PM cholesterol by increasing the availability of fatty acids for cholesterol
esterification thereby promoting the conversion of free cholesterol into CEs. We show this is
unlikely because AMPK inhibition elevated LD numbers and reduced PM cholesterol in *ACAT1/2*
DKO cells similarly to WT cells (**Fig. 5i**, **j**). Surprisingly, OA treatment, which causes the rapid
formation of large LDs, did not reduce PM cholesterol below WT levels, despite reversing the
effects of ACC1 inhibition (**Fig. 3j, k**). Analysis of OA-induced LDs by microscopy showed that
they had reduced ACC1 localization at their surface, compared to untreated cells or cells treated
with ATGL or AMPK inhibitors (**Extended data Fig. 5c**). This suggests that LDs formed from OA
addition may partially lack proteins that are important for LD localization, function, or LD-
organelle contact sites. In conclusion, two structurally distinct inhibitors that raise LD numbers
both cause a decrease in PM cholesterol levels. These findings provide additional evidence that
LDs are important for PM cholesterol homeostasis.

Fig. 5: Increasing lipid droplet abundance reduces plasma membrane cholesterol.

a-c, Wild-type (WT) HAP1 cells were left untreated or treated with ACSLi (5 µM Triacsin-C), ATGLi (50 µM NG-497) or both for 16 hrs and then analyzed by microscopy to quantify lipid droplet (LD) numbers per cell (**a**, **b**) or by mNG-ALOD4 flow cytometry analysis to determine plasma membrane (PM) cholesterol levels (**c**). **d**, Schematic demonstrating that AMPK inactivates ACC1 and activates ATGL. Inhibitors for each enzyme are shown in gray. **e**, Western blot analysis of ACC1 phosphorylation in WT HAP1 cells left untreated or treated with AMPKi (2.5 µM Dorsomorphin), ACC1i (30 µM Firsocostat) or both. **f-h** WT HAP1 cells were left untreated or treated with 30 µM ACC1i, 2.5 µM AMPKi or both and then analyzed by microscopy to quantify LD numbers per cell (**f**, **g**) or by mNG-ALOD4 flow cytometry analysis to determine PM cholesterol levels (**h**). **i,j**, LD quantification (**i**) and flow cytometry analysis of PM cholesterol using mNG-ALOD4 staining (**j**) in *ACAT1/2* DKO cells left untreated or treated with 2.5 µM AMPKi, 30 µM ACC1i or both. All cells were grown in lipoprotein depleted serum media for 16 hours prior to drug treatments. Lipid droplets are visualized with Bodipy 493:503 and nuclei are visualized with DAPI. Scale bars are 5 microns. Statistical significance was determined by an Ordinary one-way ANOVA (**b-c** and **g-j**). Each data point is the total number of LD per cell in a single field of view (**b,g** and **i**). Exact *p*-values are found in the Methods.

ACC1 loss traps accessible cholesterol at the plasma membrane and triggers de novo
cholesterol synthesis in a mouse model.
To strengthen our findings, we extended our studies with an animal model. In hepatocytes,
ACC2 contributes to the synthesis of ~10% of cellular fatty acids[68]. Therefore, we induced
recombination-mediated removal of both ACC1 and ACC2 using Albumin-CRE *Acc1/Acc2*
flox/flox mice (with flox sites around exon 21/22 and 26/27 for ACC1 and ACC2, respectively) to
generate a double liver knockout of *Acc1/Acc2* [68]. Hepatocytes isolated from *Acc1/Acc2* dLKO
mice showed undetectable levels of ACC1 and ACC2 proteins, and fatty acid synthesis was
markedly reduced (**Fig. 6a**, **b**). These hepatocytes also showed a reduction in TAG and CE
levels compared to controls (**Fig. 6c**, **d**).
Strikingly, dLKO primary hepatocytes showed a 10-fold increase in PM cholesterol (**Fig.**
6e, **f**). These data highlight that ACC1 loss leads to a sustained increase in PM cholesterol that
cannot be restored to WT levels in vivo. Liver cells are specialized to export total cellular
cholesterol, both as VLDL and as bile acids[69,70]. The inability of cells lacking ACC1 to restore

their PM cholesterol levels suggests that ACC1 is a critical component of cholesterol
homeostasis machinery.
378   Trapping cholesterol at the PM is expected to reduce ER-localized cholesterol and
trigger de novo cholesterol biosynthesis. To test the cellular response to ACC1 loss in an
unbiased way, we performed RNA sequencing on cells treated with inhibitors against either
ACC1 (Firsocostat), ACSL (Triacsin-C), HMGCR (Lovastatin) or FASN (C75). Throughout this
work, we showed that ACC1 and ACSL inhibition reduced LD numbers and increased PM
cholesterol (**Fig. 3**); in contrast, HMGCR inhibition reduced PM cholesterol (**Extended data Fig.**
3b, **c**), and FASN inhibition did not change LD or PM cholesterol levels (**Extended data Fig.**
3k). FASN inhibition serves as a useful control for ACC1 inhibition, since both conditions block
fatty acid synthesis. Strikingly, we observed that ACC1 and ACSL inhibition potently elevated the
expression of cholesterol biosynthesis genes, and closely resembled HMGCR inhibition (**Fig.**
6g). To the best of our knowledge, prior to this work, there was no known mechanism to explain
why blocking fatty acid biosynthesis would trigger SREBP2 activation. FASN inhibition had small
to no effects on SREBP2 targets, consistent with its inability to trigger LD breakdown. Inspection
of SREBP2 target gene induction across treatments yielded unexpected findings for future
studies- for example, FDFT1, the first committed step in cholesterol biosynthesis was not
induced by HMGCR inhibition, though it was significantly induced by ACC1 inhibition.
394   Cholesterol and fatty acid biosynthetic pathways have evolved to be coregulated, which
is evidenced by significant work showing that SCAP, an ER-localized cholesterol sensor,
controls the activation of both SREBP1 and SREBP2 transcription factors[14,71,72]. This work
implies that when ER cholesterol levels are low, both SREBP1 and SREBP2 target genes are
induced. Investigating known SREBP1 targets revealed that multiple genes (FASN, FADS2,
SCD, ACSL3, ACACA, and SREBP1 itself) were elevated more strongly through the inhibition of
ACC1 or ACSL than by HMGCR (**Fig. 6h**). This was not due to the relative potency of different
inhibitors since HMGCR inhibition had the strongest effects for multiple cholesterol biosynthesis
genes (including HMGCR, MVK, SQLE and CYP51A1). Factors such as insulin and
polyunsaturated fatty acids are known to regulate SREBP1 independently of SREBP2; future
investigations are required to determine which of these are driving the discordance seen
between SREBP2 and SREBP1 targets in our study[73–76].
406   To determine which pathways were most significantly affected by LD loss, we performed
a Gene Set Enrichment Analysis (GSEA) using the top 200 most differentially regulated genes
from each treatment condition (ranked by FDR-corrected p-value). Among the pathways
enriched, cholesterol metabolism was number one for both ACC1- and ACSL-inhibited cells
(**Fig. 6i**). All four treatments showed similar effects on VEGFA-VEGFR2 signaling and mRNA
processing pathways. Finally, ACSL and FASN inhibition showed multiple overlapping enriched
pathways. Altogether, RNA sequencing provided strong evidence that ER cholesterol is depleted
by ACC1 and ACSL inhibitor treatments, which supports the model that LD loss reduces
cholesterol trafficking out of the PM.
415   Finally, we tested cholesterol synthesis rates in the *Acc1/Acc2* dLKO primary
hepatocytes. Consistent with our cellular findings, radiotracing of tritiated water showed dLKO
hepatocytes had markedly elevated rates of cholesterol synthesis compared to controls (**Fig.**
6j). This result, combined with the observed increase in PM cholesterol levels (**Fig. 6e**, **f**),
suggests that ACC1 loss reduces cholesterol trafficking from the PM to the ER. Despite having

elevated PM cholesterol and elevated cholesterol synthesis, *total* cellular cholesterol levels were
not significantly elevated in vivo or in cell culture (**Fig. 6k**, **Fig. 2n** and **Extended data Fig. 6a**,
b). What is the fate of the newly synthesized excess cholesterol? Analysis of fecal cholesterol
levels revealed that it was excreted in mice (**Fig. 6l**). These findings demonstrate that while
maintaining high PM accessible cholesterol, cells undergo homeostasis to maintain total cellular
cholesterol levels. This observation is at odds with the dogma which states that cells sense and
regulate the accessible pool of cholesterol specifically. In conclusion, our study finds that ACC1,
the master regulator of fatty acid biosynthesis, is critical for controlling LD catabolism through
ATGL. Most strikingly, we show that LDs are crucial for cellular cholesterol homeostasis– their
loss traps cholesterol at the PM and induces de novo cholesterol synthesis in the ER (**Fig. 6m**).

Fig. 6: ACC1 loss traps cholesterol at the plasma membrane and triggers de novo cholesterol synthesis in a mouse model.

a, Western blot analysis of ACC1 and ACC2 proteins in primary hepatocytes isolated from *Acc1/Acc2* flox/flox and Albumin-Cre-*Acc1/Acc2* dLKO mice. **b-d**, Hepatic fatty acid synthesis rates (**b**), hepatic triacylglycerol (TAG) levels (**c**), and hepatic cholesterol ester (CE) levels (**d**) in *Acc1/Acc2* flox/flox and Albumin-Cre-*Acc1/Acc2* dLKO mice. **e**, Fluorescence microscopy images of primary hepatocytes stained with mNG-ALOD4 and DAPI. Scale bar is 10 microns. Quantification is shown in (**f**). **g,h**, Bubble plot showing differential gene expression from RNA sequencing in WT HAP1 cells. Cells were pre-cultured in lipoprotein depleted serum (LDS) media for 16 hours and then left untreated or treated for 6 hrs with HMGCRi (10 μM Lovastatin), ACC1i (30 μM Firsocostat), ACSLi (5 μM Triacsin-C) or FASNi (10 μM C75). Differential gene expression was computed by comparing each treatment to the untreated (LDS only) control. Bubble size shows the FDR-corrected *p*-value and bubble color shows the fold change (see legends) for each SREBF2 (**g**) and SREBP1 (**h**) target gene. SREBP1 target genes are ranked based on their enrichment in a ChIP-Seq dataset. **i**, Gene Set Enrichment Analysis from RNA sequencing data using the top 200 genes sorted by the FDR-corrected *p*-value. "nf" refers to not found. **j-l**, Hepatic sterol synthesis rate (**j**), hepatic cholesterol content (**k**), or fecal cholesterol measurement (**l**) in chow-fed *Acc1/Acc2* flox/flox and Albumin-Cre-*Acc1/Acc2* dLKO mice. **m,** Schematic showing a proposed model: ACC1 inhibits ATGL, blocking the breakdown of TAG to diacylglycerol (DAG) and fatty acids (FA). When ACC1 is inhibited (right), ATGL is activated, leading to TAG hydrolysis. Cell diagrams show that lipid droplets (LD) control cholesterol trafficking from the PM to the ER. When LDs are depleted (right), cholesterol becomes trapped at the PM, and SREBP2 drives the expression of cholesterol biosynthesis genes. Statistical significance was determined by a Student's t-test with a Welch's correction (**b-d, f** and **j-l**). Exact *p*-values are found in the Methods.

Discussion

In this study we set out to find new regulators of PM cholesterol levels. We were surprised to discover that ACC1, a protein extensively studied for its role in fatty acid synthesis, also regulates cellular cholesterol homeostasis. ACC1 knockout raises PM cholesterol levels 5- to 10-fold, restores PM cholesterol changes observed in NPC1-deficient cells, and also blocks PM cholesterol depletion induced by 25HC through a mechanism that is independent of cholesterol esterification by ACAT1/2. We implicate LDs as playing a critical role in cholesterol movement out of the PM. Reducing LD numbers traps cholesterol at the PM and strongly induces de novo cholesterol biosynthesis both in vitro and in an animal model. Our screen did not identify positive regulators of LD formation because many of these proteins are redundant and their knockout is not sufficient to cause LD loss– this redundancy speaks to the importance of LDs for cell survival. Given that the majority of cells in the human body have LDs, the mechanism we have discovered could represent a broad principle of cholesterol trafficking[77,78].

PM accessible cholesterol levels are thought to be faithfully maintained by continuous transport to the ER, where sensors like SCAP, ACAT1/2, and GRAMD1B localize[10,11,79]. Our discovery that LD loss causes a sustained increase in PM accessible cholesterol is surprising, and suggests that LDs are essential for the trafficking of a large pool of PM cholesterol. Why would evolution link PM-to-ER cholesterol trafficking to LD levels? We draw inspiration from the seemingly peculiar and inelegant arrangement observed in animal cells: SCAP controls both SREBP1 (fatty acid) and SREBP2 (cholesterol) activation, and yet SCAP is solely a cholesterol sensor[71]. Why not independently couple fatty acid sensing to fatty acid synthesis and cholesterol sensing to cholesterol synthesis? We believe our work raises an intriguing hypothesis to reconcile this mystery– LD levels convey information about fatty acid storage. When fatty acid storage is low (LDs are low), cholesterol cannot effectively traffic to the ER, leading to the activation of both SREBP1 and SREBP2. Conversely, when lipid storage is high and LDs are abundant, cholesterol trafficking is intact, which blocks de novo lipid synthesis; in other words, it is disadvantageous to make new lipids when significant lipids are already stored.

This paper raises the question of how LDs control cholesterol trafficking. LDs within a single cell are heterogeneous, with unique protein binding and organelle contact sites, suggestive of distinct functions[57,59]. We propose two hypotheses for how LDs control PM cholesterol transport. First, when cholesterol molecules are trafficked out of the PM, they may move onto the LD phospholipid monolayer. The LD monolayer could either accumulate cholesterol, or this membrane may act as a bridge intermediate between the PM and another organelle such as the ER. Secondly, LDs may be required for endosome maturation[80]. In this model, a loss of LDs causes early endosomes to recycle more rapidly, which would resemble a defect in cholesterol trafficking out of the PM and cause PM cholesterol to rise if anterograde trafficking of cholesterol is intact. Future studies will distinguish between these models.

While our study provides valuable insights into the role of ACC1 inhibition in promoting LD catabolism by ATGL, further investigation is needed to fully elucidate the molecular mechanism underlying this process. ATGL is known to be regulated by AMPK, however, we observed ACC1 inhibition induced LD breakdown even in the presence of AMPK inhibition, suggesting that ATGL activation was AMPK independent. We propose that ACC1 could directly inhibit ATGL through an unknown mechanism– when ACC1 is inactivated or lost, ATGL

becomes active and LDs are broken down. Future studies will be needed to determine how
ACC1 triggers LD catabolism through ATGL.
Our study has implications for the myriad of disease pathologies being linked to the
enzyme CH25H and its product 25HC. Recent publications show that CH25H has roles beyond
pathogen evasion; its expression can promote diabetes, microglial activation, adipose tissue
inflammation and fibrosis[43–45,81]. We show that ACC1 is required for 25HC-induced cholesterol
trafficking out of the PM, and suggest that ACC1 inhibitors could reverse some of the disease
pathologies linked to elevated 25HC levels. Additionally, LDs play clear roles in metabolic
syndromes, and they have also been linked to changes in neuronal function and
neurodegeneration[82,83]. ACC1 inhibition could alleviate LD-driven disease symptoms by
triggering their catabolism. For example, we find that Firsocostat more robustly, and with less
toxic effects, eliminates LDs compared to Triacsin-C, a previous gold standard in the field. In
conclusion, we find a novel function of ACC1 in ATGL activation and LD homeostasis, and also
demonstrate that LDs control PM cholesterol transport.
488
489

Acknowledgements

491 We thank Dr. Yvonne Lange, Dr. Theodore Steck, Dr. Pehr Harbury and Dr. Giovanni Luchetti
492 for helpful discussions.
493
Funding: MK was supported by GM118082, the Ara Parseghian Medical Research Foundation,
the Howard Hughes Medical Institute Hanna H. Gray Fellows Program and 5DP5OD03615502.
RR was supported by GM118082 and the Ara Parseghian Medical Research Foundation; CBK
was also supported by GM118082. JAO was supported by R01DK128099. KMW, EOS and SL
were supported by 5DP5OD03615502. SC was supported by a Fellowship from the Arc
Institute. ET was supported by the Stanford Community College Outreach Program (CCOP),
with funding provided by the Biochemistry Department at Stanford University and an
anonymous donor gift to CCOP. CK was supported by P01HL160487 and P30DK127984. DT
was supported by the Intramural Research Program of the National Cancer Institute at the
National Institutes of Health. AL was supported by the Center for Cancer Research, National
Cancer Institute, National Institutes of Health Intramural Research Program project number ZIA
BC 011901.
506
507 Competing Interests: JAO is a member of the scientific advisory board for Vicinitas
508 Therapeutics.

509
510

Materials and Methods

Cell culture

HAP1 cells were grown in HyClone Isocove's Modified Dulbecco's Medium (IMDM)
(SH30228FS, Fisher/Cytiva), supplemented with 5% Fetal Bovine Serum (FBS) (F0926,
Millipore Sigma), 2 mM L-glutamine (MT25005CI, Fisher/corning), Penicillin (50 U/mL) and
Streptomycin (50 µg/mL) (15140122, Fisher/Gibco).

3T3 and 293T cells were cultured in HyClone™ Dulbecco's Modified Eagles Medium
(SH30081FS, Fisher/Cytiva), enriched with 10% FBS (F0926, Millipore Sigma), 2 mM L-
glutamine (MT25005CI, Fisher/Corning), Penicillin (50 U/mL) and Streptomycin (50 μg/mL)
(15140122, Fisher/Gibco), 1 mM sodium pyruvate (11360070, Fisher/Gibco), 1× MEM Non-
Essential Amino Acids Solution (11140050, Fisher/Gibco).
Lipoprotein Depleted Serum (LDS) was prepared by supplementing media with 5% LDS
instead of 10% FBS. To prepare LDS, FBS was incubated with Cab-o-sil (1 gram per 50 mL
FBS) overnight at 4°C with constant shaking. After 24 hours, the suspension was centrifuged at
10,000 rpm to pellet silica complexes. The supernatant was carefully decanted without
disturbing the pellet, followed by addition of fresh Cab-o-sil at the original 1 g/50 mL ratio. After a
second overnight incubation at 4°C with shaking, the suspension was centrifuged again and
then the supernatant was filtered through 0.2 μm filter, aliquoted and snap-frozen in liquid
nitrogen for future use.

Drug treatments
Small molecules used in this study include: 25-hydroxycholesterol (25HC) (11097, Cayman
Chemical), the LXR inhibitor GSK2033 (1 μM) (25443, Cayman Chemical), the NPC1 inhibitor
U18666A (1 μM) (10009085, Cayman Chemical) the ACC1 inhibitor Firsocostat (also named
GS-0976 and ND-630) (30 μM) (HY-16901, MedChemExpress), the ACC1 inhibitor CP-640186
(20 μM) (HY15259, MedChemExpress), the HMGCR inhibitor Lovastatin (10 μM) (10010338,
Cayman Chemical), the ACAT1/2 inhibitor SZ58-035 (50 μM) (S9318, Millipore Sigma), the
FASN inhibitor C75 (10 μM) (10005270, Cayman Chemical), the AMPK inhibitor Dorsomorphin
dihydrochloride (2.5 μM) (HY-13418, MedChemExpress), the long-chain fatty acyl CoA
synthetase (ACSL) inhibitor Triacsin-C (5 μM) (BMLEI218, Enzo), human adipose triglyceride
lipase (ATGL) inhibitor NG-497 (50 μM) (HY-148756, MedChemExpress), Oleic acid (200 μM)
(O1383,Millipore Sigma), the *DGAT1* inhibitor T863 (20 μM) (SML0539, Sigma-Aldrich) and the
DGAT2 inhibitor PF-06424439 (10 μM) (PZ0233, Sigma-Aldrich). Drug treatments were carried
out in various media compositions indicated in the Figure legends, for 16 hours at 37°C unless
otherwise noted.
Flow cytometry analysis of plasma membrane cholesterol
The cells of interest were seeded in a 24-well cell culture plate (3 replicates for each condition)
aiming for ~80% confluency on the day of flow cytometry analysis. The kinetics of 25-
hydroxycholesterol (25HC) were measured by introducing 4 μM 25HC for various times prior to
flow cytometry analysis both with and without 16 hour prior incubation with the noted inhibitors.
To prepare cells for flow cytometry they were first rinsed with 1x Phosphate Buffered
Saline (PBS) (P38135, Sigma), harvested with Trypsin (0.05%) - 1x EDTA (25300120,
Fisher/Gibco), and quenched with 4°C 5% FBS containing IMDM (or 10% FBS containing
DMEM) media while gently pipetting up and down to generate a single cell suspension. Cells
were then transferred to a 1.5 mL eppendorf tube on ice, and then spun for 5 minutes at 4°C
(HAP1 WT at 1000 x*g* and 3T3/293T at 750 x*g*). The supernatant was removed and cell pellets
were resuspended in ~25 μL of 2 μM mNeonGreen-ALOD4 (mNG-ALOD4) diluted in 1x Probe
Block Buffer (PBB- 1x PBS, 1% BSA, filter sterilized through a 0.2 μm filter). The cells were
incubated on ice for 12 minutes before being diluted to 150 μL in PBB, and then transferred to a
96-well plate. Flow cytometry was performed using the BD Accuri C6 Plus Flow Cytometer.

During flow cytometry single live cell populations were selected based on forward scatter (FSC)
versus side scatter (SSC) density plots. The fluorescence of mNG-ALOD4 was measured by
exciting using a 488 nm laser and detecting the emission through a 585/40 nm bandpass filter.
The FCS files generated from BD CSampler software were loaded to FCS Express software
and then exported directly to GraphPad Prism for analysis.
565

Measuring sphingomyelin levels with SMase
HAP1 WT cells and HAP1 *ACC1* KO cells were seeded in 5% FBS IMDM in a 24-well plate and
grown to ~50% confluency. Cells were then left untreated or treated for 16 hours with 30 µM
Firsocostat in 5% FBS IMDM. The day of the assay, cells were first washed with 1x PBS and
then incubated at 37 ºC with sphingomyelinase (SMase) (150 mU/mg protein,S8633, Millipore
Sigma) for 30 minutes to hydrolyze sphingomyelin. After incubation, the cells were washed and
trypsinized for flow cytometry analysis as described above.
573

Measurement of free cholesterol and cholesterol esters
Cells were plated in a 12-well tissue culture dish in 5% FBS IMDM media and grown to 80%
confluency. On the day of the assay, cells were washed with 1x PBS and lysed for 10 minutes at
4ºC with 150 µL of lysis buffer containing 10% glycerol, 2x Protease Inhibitor Cocktail (S8830,
Millipore sigma), 150 mM NaCl, 50 mM tris-HCl (pH 8), 1% NP-40 and 1 mM $MgCl_2$. The
Amplex Red Cholesterol Assay Kit (A12216, Invitrogen/Fischer) was used to measure the free
cholesterol and cholesterol esters in each sample. After lysis, lysates were diluted with an equal
volume of 1x reaction buffer from the kit. Total cholesterol and free cholesterol were measured
by preparing two working solutions (with and without cholesterol esterase) as described in the
assay protocol and mixing the lysates with the mastermix in a 1:1 ratio in a 96-well plate. The
plate was incubated at 37 ºC for 30 minutes and the fluorescence emission was measured at
590 nm using a microplate reader.
586

Microscopy analysis of lipid droplets
For lipid droplet analysis, the cells of interest were seeded on acid-washed coverslips in a 24-
well plate. Various treatments were performed as described in the figure legends. To stain for
lipid droplets, cells were washed with 1x PBS buffer and fixed in a solution containing 4%
paraformaldehyde (PFA) and 1x PBS for 10 minutes at room temperature. After fixing, cells
were washed 3 times with 1x PBS, and then cells were permeabilized with immunofluorescence
(IF) block buffer (1x PBS, 1% BSA and 0.1% Triton X-100, filter sterilized through a 0.2 µm filter)
for 15 minutes at room temperature. Cells were then incubated with 10 µM Bodipy 493/503
(Cayman Chemical; Cat# 25892) in IF block buffer for 1 hour followed by 3 washes with 1x PBS.
Coverslips were mounted on glass slides using ProLong Diamond Antifade Mountant (P36962,
Invitrogen/thermo fisher) and left to cure overnight at room temperature before imaging. Imaging
was conducted using a Keyence BZ-X800L fluorescence microscope.
To co-stain for proteins of interest (ACC1 or FASN) and lipid droplets, cells were treated
the same as above. After the 15 minute permeabilization step, cells were stained with ACC1
primary antibody (RRID: AB_2882621) diluted 1:1000 in IF block buffer for 1 hour at room
temperature. Coverslips were then washed 3x with IF block buffer before adding a mixture of 10
µM BODIPY 493/503 and secondary antibody (anti- mouse 647; RRID:AB_3073503) diluted

1:500 in IF block buffer for 1 hour. Alternatively (**Fig. 3c**), CoraLite Plus 647-conjugated ACC1
mouse polyclonal antibody (RRID: AB_11042445) was diluted 1:500 along with 10 µM BODIPY
493/503 in IF block buffer. After 1 hour of incubation, cells were washed 3x with IF block buffer,
and then mounted on glass slides and imaged as described above.
608
Quantification of lipid droplets
To quantify lipid droplets (LDs), images were processed using ImageJ software. First, images
were imported into ImageJ and then the images were converted to an 8-bit grayscale format.
Thresholding was performed to eliminate background by manually adjusting the threshold
intensity. Within a single experiment, all threshold values were kept constant for equal
comparisons of the images.
LDs were then quantified using the analyze particles function in ImageJ and the
parameters were configured to limit particle size to a minimum of 2 pixels² to eliminate very
small thresholded noise; circularity was set to 0.00-1.00. The results window provided raw data,
including the LD area (pixels² was converted into µm² using the conversion factor for our
microscope = 0.12581 µm/ pixel), which was copied into excel for analysis. LDs per cell were
calculated by dividing the total number of LDs for a field from the total number of nuclei in a
given field. In Extended data Fig. 3e, LDs were counted manually and assigned to a specific cell
to get a measurement of per-cell LD numbers. This latter method was not a preferred method
because when cells reach >70% confluency, it is often difficult to ascertain which LD belongs to
which cell.
625
Cellular fractionation to isolate lipid droplets
HAP1 WT cells were seeded on two 15 cm plates, with one plate treated with 30 µM Firsocostat
for 16 hours. Cells were harvested by gentle scraping in ice cold 1x PBS. The cell suspensions
were then centrifuged at 1000 xg for 5 minutes at 4°C to obtain cell pellets. After removing the
supernatant the pellets were resuspended in 8% sucrose buffer containing 0.25 M sucrose, 50
mM Tris-HCl (pH 7.4), 150 mM NaCl, 1x Protease Inhibitor Cocktail, and 1x Phosphatase
Inhibitor (1 PhosSTOP tablet per 10 ml cell extract or buffer, Millipore Sigma Cat #
4906845001). After incubating for 10 minutes on ice, the cells were lysed by passage through a
chilled ball-bearing cell homogenizer (isobiotec) with a 12 µm clearance 8 times. Successful
lysis was confirmed using Trypan blue staining (Gibco, Cat#15250061). A fraction (~50 µL) of
the homogenate was saved for Western blotting (whole-cell lysate). The remainder of the
homogenate was centrifuged twice at 1000 xg for 10 minutes each to pellet nuclei. The
supernatant (~600 µL from the top) was pipetted into a separate chilled microtube and the
pelleted nuclei was saved for Western blotting.
To isolate lipid droplets (LDs), 600 µL of 0% sucrose buffer (150 mM NaCl and 50 mM
Tris-HCl pH 7.4) was slowly added on top of the supernatant isolated above while trying to
maintain the separation of different sucrose densities. The tubes were centrifuged at 130,000 xg
for 1 hour at 4°C. LDs were collected in four, 50 µL fractions (total volume 200 µL) using a pre-
wetted pipette tip starting from the top floating layer, which appeared cloudy. The remaining
~400 µL of 0% sucrose buffer was removed. The bottom ~600 uL cytoplasmic fraction was
collected and saved for Western blotting. Finally, the pelleted plasma membrane fraction was
resuspended in Lysis Buffer (10% glycerol, 2x Protease Inhibitor Cocktail (S8830, Millipore

sigma), 150 mM NaCl, 50 mM tris-HCl (pH 8), 1% NP-40 and 1 mM $MgCl_2$) and saved for
Western blotting.
650   The four LD fractions were washed with 500 µL 0% sucrose buffer by spinning at 20,000
651   xg for 30 minutes at 4°C. Next using a syringe needle, ~600 µL from the bottom of the tube was
652   removed and the top portion was saved for Western blotting of the LD fraction.
653   For Western blot analysis, the whole cell fraction was lysed by adding NP-40 to 1%. The
nuclear fraction was also resuspended in Lysis Buffer. The whole cell, nuclear, and plasma
membrane fractions were mechanically shaken at 4°C for 45 minutes to aid in lysis. The lysates
were then clarified by spinning at 18,000 xg for 30 minutes to remove aggregated material and
DNA. Finally, protein concentrations for the whole cell, nucleus, cytoplasm and plasma
membrane fractions were measured with a BCA assay, and equal amounts of protein were
saved for Western blots. LD fractions were also lysed by adding 1% NP-40, but BCA assays
revealed very low protein concentrations. For Western blot, the ~40 µL (a maximum possible
volume) of lysed LDs were loaded.
662

Western blotting
Cells were plated on a 6 cm dish and treated with small molecules when indicated. In **Fig. 6a**,
primary hepatocytes were isolated from 8-week old mice and then plated on collagen-coated
coverslips prior to analysis. Cells were then washed and scraped with chilled 1x PBS, and then
pelleted by spinning at 1000 xg for 5 minutes at 4°C. After removing the supernatant, the cells
were lysed with Lysis Buffer (10% glycerol, 2x Protease Inhibitor Cocktail (S8830, Millipore
sigma), 150 mM NaCl, 50 mM tris-HCl (pH 8), 1% NP-40 and 1 mM $MgCl_2$) at 4°C using a
mechanical shaker for 1 hour. Lysates were spun for 30 minutes at 18,000 xg and 4°C, and then
supernatants were transferred to a new tube. A BCA assay was performed on the supernatant to
determine the total protein concentration. Samples containing equal amounts of protein (~40 µg
per well) in 1X mPAGE LDS Sample Buffer (MPSB, millipore sigma) and 100 mM Dithiothreitol
(BP17225, Fisher) were incubated at 37 °C for 30 minutes before SDS-PAGE.
675   After completion of the SDS-PAGE, samples were transferred from the gel to a PVDF
membrane (IPVH85R, millipore sigma). The bands were cut after staining the membrane with
Ponceau stain and blocked with 5% nonfat milk in TBST (1x TBS, 10% tween). For analysis of
knockout cell lines, the membranes were incubated with the appropriate dilutions of
commercially available primary antibodies (ACAT1, ACC1, NPC1, FASN, ATGL) in 5% nonfat
milk diluted in TBST for 2 hours at room temperature, washed 3 times with TBST for 5 minutes
each, and then incubated with HRP-conjugated secondary antibodies diluted to 1:10,000 for 1
hour at room temperature. After washing the membrane three times with TBST for 5 minutes
each the blot was developed using ECL solutions (PI32209, Fisher/ Pierce).
684

Generation of HAP1 mutant cell library for random insertion mutagenesis screen
To randomly mutate the genome we used a retrovirus bearing a Gene Trap (GT) cassette that
introduces a splice acceptor site and a poly A tail, which terminates transcription regardless of
whether it inserts into introns or exons [84]. In contrast to CRISPR knockout screens, GT
mutagenesis can identify regulatory mechanisms that originate in non-coding regions like
promoters, and it can identify hypomorphic and dominant negative alleles by truncating proteins.

691   The production of GT virus was previously reported[85]. Briefly, to produce GT retrovirus,
692 15 million HEK 293FT cells were seeded into six T-175 flasks in 10% FBS DMEM media lacking
693 antibiotics. When cells reached 80% confluency, they were transected with two retroviral
694 transfer plasmids containing the GT in two reading frames (3.3 µg pGT-mCherry and 3.3 µg pG-
695 +1-mCherry), a plasmid with viral packaging genes (4 µg of pCMV-Gag-Pol), a plasmid with viral
696 envelope genes (2.6 µg of pCMV-VSV-G), and a plasmid that enhances translation (1.7 µg of
697 pAdVAntage). First, 45 µL of X-tremeGENE HP was added to 450 µL of OptiMEM for 5 minutes
698 at room temperature, and then plasmids were added to the mixture, which was briefly vortexed
699 and then incubated for 30 minutes at room temperature. The transfection reaction was added
700 dropwise to the 293FT cells. After 16 hours, virus-containing media was harvested, filtered
701 through a 0.45 µm low-protein binding filter and then concentrated by ultracentrifugation at
702 65,000 xg and 4°C for 1.5 hours in a swinging bucket rotor. Eight hours later, the virus was
703 harvested and concentrated for a second time. After centrifugation, the supernatant was
704 removed, leaving ~ 1 mL of media to avoid disrupting the viral pellet. Tubes were then sealed
705 with parafilm and incubated overnight at 4°C.
706   To generate the HAP1 cell library WT HAP1 cells were pre-sorted for haploidy using a
707 Hoechst stain. Then, 20 million haploid HAP1 cells were seeded into each of three T-175 flasks
708 in antibiotic free 10% FBS IMDM. Immediately before infecting with GT-bearing retrovirus, the
709 viral pellets were combined and resuspended in 76 mL of antibiotic free 10% FBS IMDM
710 containing 4 µg/mL polybrene. One-third of this mixture was added to each of the three flasks
711 containing WT HAP1 cells. Cells were expanded and maintained in multiple flasks, such that 50-
712 100 million cells were kept in passage at any given time to ensure the genotypic richness of the
713 library was maintained. Flow cytometry analysis measuring the GT-based mCherry fluorescence
714 was used to assess transduction efficiency, which was determined to be >90%.
715
716 **Random insertion mutagenesis screens**
717 For each screen, 20 million HAP1-GT cells were seeded in 7x T-175 flasks in 10% FBS IMDM
718 media. To find genes that increase PM cholesterol after NPC1 loss of function, all 7 flasks were
719 treated 8 hours after seeding with 1 µM U18666A. 20 hours later, 1 flask of cells was harvested
720 by trypsinization, pelleted at 1000 xg, and then snap frozen as an unsorted control. The
721 remaining flasks were prepared for Fluorescence Activated Cell Sorting (FACS). Plates were
722 processed one at a time, one per hour. After washing with 1x PBS, cells were trypsinized and
723 then spun at 1000 xg to isolate a cell pellet. Cells were then resuspended in 4 µM mNG-ALOD4
724 diluted into ice cold Probe Block Buffer (1x PBS, 1% BSA, filter sterilized). After 20 minutes of
725 staining on ice, cells with the top 10% of mNG-ALOD4 fluorescence were isolated with FACS.
726 Approximately 1-2 million cells were harvested per T-175 flask. After FACS, cells were spun
727 down and then seeded into the same 15 cm plate. Cells were then expanded until 6x T-175
728 flasks could be reseeded. FACS was carried out 2 consecutive times to enrich the cells with
729 high mNG-ALOD4 fluorescence. After the second sort, 30 million cells were pelleted and snap
730 frozen in liquid nitrogen. Library preparation and the computational analysis that follows has
731 been previously described in detail [86]. The only difference is that for the calculation of FDR-
732 corrected *p*-values (**Fig. 1g**, **h**) we included all GT insertions, regardless of orientation, for both
733 the selected and unsorted cell populations.

To identify genes that regulate cholesterol trafficking in response to 25-
hydroxycholesterol, 20 million HAP1-GT cells were seeded in 7x T-175 flasks in 10% FBS
IMDM, as described above. On the day of FACS sorting, one plate was isolated as the unsorted
control. The remaining 6 plates were treated in a staggered manner, with 1 flask treated every
hour with 4 µM 25-hydroxycholesterol. Each total treatment time was 6 hours. After 6 hours of
treatment, cells were harvested by trypsinization, pelleted, and then isolated using the same
protocol described above. For this screen, cells were sorted a total of three times, which may
give rise to the increased FDR-corrected p-values observed for this screen compared to the
U18666A screen (**Fig. 1g**, **h**).
743
Mass spectrometry analysis of lipid species
WT HAP1 cells were grown on 4x 15 cm dishes per treatment condition until ~75% confluent in
5% FBS IMDM. Cells were then moved into 5% LDS media with or without inhibitor (30 µM
Firsocostat) for 24 hours. Media was removed and cells were washed gently 3x with room
temperature 1x PBS to remove dead cell debris. Cells were then scraped in ice cold 1x PBS on
an ice metal rack in the tissue culture hood. Each plate of cells was moved to a single 15 mL
conical tube and then spun at 1000 x*g* to pellet. PBS was then aspirated, cells were snap frozen
in liquid nitrogen and shipped on dry ice to Metabolon Inc. for analysis.
Raw values obtained from Metabolon Inc. are areas under the curve for each detected
lipid species, normalized to protein concentrations for each sample. Values are not exact
quantitative measurements since internal standards were not used. All comparisons made
should only be considered as relative changes compared to the control sample (5% LDS lacking
inhibitors).
For analysis of lipids from *Acc1/Acc2* flox/flox mice and Albumin-CRE *Acc1/Acc2*
flox/flox mice, mice were chow-fed, with n=4 per group. Lipids were extracted by liquid-liquid
extraction and analyzed by LC-MS/MS using internal standards. Each value was normalized to
tissue weight.
761
Measurements of the rate of fatty acid and cholesterol synthesis in mouse livers
For analysis of fatty acid synthesis rates, mice were maintained on a chow diet and subjected to
a time-restricted feeding regimen (feeding from 6 PM to 6 AM; fasting from 6 AM to 6 PM) for 3
consecutive days prior to the assay. [^{3}H]H$_2$O (50 mCi) was administered intraperitoneally, and
hepatic fatty acid synthesis rates were determined by quantifying ^{3}H incorporation into newly
synthesized fatty acids, as previously described [87]. Hepatic sterol synthesis rates were
assessed by measuring [^{3}H] incorporation into newly synthesized sterols, as previously
described [87].
770
Measurement of total liver cholesterol and fecal cholesterol content in mice
For hepatic liver cholesterol measurements, approximately 100 mg of liver tissue was used for
lipid extraction, and cholesterol quantification was measured using the Infinity™ Cholesterol
Liquid Stable Reagent.
For fecal cholesterol measurements, feces were collected from mice (n = 7 per group),
which were individually housed overnight. Around 100 mg of feces was homogenized in Folch
reagent (chloroform:methanol, 2:1), and the organic phase was extracted by phase separation

with saline and isolated by centrifugation. A portion of the organic phase was dried in the
presence of Triton X-100, and cholesterol content was measured using the Infinity™ Cholesterol
Liquid Stable Reagent.
781
RNA sequencing analysis
WT HAP1 cells were grown in 6-well plates, with 4 wells per condition, until ~75% confluent in
5% FBS IMDM. Cells were then moved into 5% LDS media for 16 hours. Cells were then
treated with or without 10 µM Lovastatin, 30 µM Firsocostat, 5 µM Triacsin-C or 10 µM C75 for 6
hours. Cells were then washed twice with 1x PBS, and then all PBS was aspirated.
RNA was isolated using Trizol-chloroform extraction. Briefly, 800 µL Trizol was added
directly to the wells, incubated for 5 minutes, and then transferred to DNase/RNase-free 1.5 mL
eppendorf tubes. After adding 160 µL chloroform, samples were shaken vigorously for
15 seconds and incubated at room temperature for 3 minutes. Phase separation was achieved
by centrifugation at 12,000 xg for 15 minutes at 4 °C, and then the aqueous phase was
transferred to a new tube. Next, 400 µL isopropanol was added and then samples were
incubated for 10 minutes at room temperature. RNA was then pelleted by centrifugation at
12,000 xg for 10 minutes at 4 °C. The RNA pellet was washed with 800 µL 75% ethanol,
vortexed briefly, and centrifuged at 7,500 xg for 5 minutes at 4 °C. The supernatant was
removed, and pellets were spun again for 5 minutes. Residual ethanol was removed and pellets
were air-dried at room temperature for 5–10 minutes until the pellet turned translucent. RNA was
resuspended in 40 µL DNase/RNase-free water, incubated at room temperature for 10 minutes,
and gently mixed. RNA was stored at -80 °C and shipped to Novogene USA Inc. for standard
RNA sequencing analysis. The returned FASTQ files were analyzed using the PartekFlow
software using previously described methods from Xiaowen Wang (Field Application Specialist
from Partek Incorporated). Data was processed using a differential gene set enrichment
analysis (DeSeq2).
The bubble plots (**Fig. 6g**, **h**) were generated using both GraphPad Prism 10 and Adobe
Illustrator. First, a list of genes of interest were curated for SREBP2 (**Fig. 6g**) and SREBP1 (**Fig.
6h**). SREBP1 targets were found using a publically available ChIP-Seq database (ChIP-Atlas).
Next, FDR-corrected p-values were -Log$_{10}$ transformed, and Log$_2$ fold changes values were
subjected to exponential transformation (2^(Log$_2$(fold-change values)). The transformed p-
values and fold change values for genes of interest were found for each treatment and given a
unique identifier (e.g. HMGCRi-ACAT2, ACC1i-ACAT2, ACSLi-ACAT2, and FASNi-ACAT). All
genes were then plotted into one column using a Multiple Variable analysis plot type in Graph
Pad Prism 10. Bubble size was set to be determined by the transformed p-value, and bubble
color was set to be the fold change; this software also assigned x-axis and y-axis positioning for
the bubbles, however these were not used in the final figure generation. GraphPad Prism 10
was simply used to generate bubbles of meaningful size and color for each gene in its
respective treatment. Finally, the GraphPad Prism 10 file was exported as a .emf file, opened in
Adobe Illustrator, and the bubbles were re-arranged into a column graph.
818
Gene Set Enrichment Analysis
To analyze the top pathways affected by ACC1, ACSL, FASN, and HMGCR inhibition, we
utilized the Gene Set Enrichment Analysis (GSEA) website. First, we navigated to the Explore

the Molecular Signatures Database (MSigDB), and then to the "Investigate" tab in the Human
Collections. In the "Input Gene Identifiers" box, we pasted the top 200 genes that were
differentially regulated for each treatment- rankings were made by sorting based on FDR-
corrected p-values. In the Compute Overlaps window, we selected the WikiPathways gene sets,
and then computed the overlaps. This returned the top 10 WikiPathways for each set of genes.
Enterocyte Cholesterol Metabolism (WP5333), Cholesterol Metabolism (WP5304), Cholesterol
Biosynthesis Pathway (WP197), and Cholesterol Biosynthesis Pathway in Hepatocytes
(WP5329) were excluded from the final plot due to their significant overlap with the genes found
in Cholesterol Metabolism with Bloch and Kandutsch-Russell Pathways (WP4718) and
Cholesterol Synthesis Disorders (WP5193), which were used in the final plot. If a pathway was
not found in the top 10 pathways identified, it was denoted in the plot as "nf" for not found.
833
Crispr Cas9 mediated knockout of genes
Two single guide RNA (sgRNA) sequences were designed using the Benchling software to
direct Cas9 binding and cleavage at two separate genomic sites[88]. Next the sgRNA guide
sequences were cloned into px458-GFP and px458-mCherry plasmids. The sequence verified
plasmids were next transfected into HAP1 cells and after 24 hours both mCherry- and GFP-
positive cells were single cell sorted into a 96 well plate using FACS. Subsequently, these
single-cell clones were expanded and then genotyped in order to identify knockouts.
841
Statistical analysis
Data analysis was done using GraphPad Prism 10.3.1. All statistical comparisons between two
datasets were conducted using a Student's t-test with Welch's correction. Ordinary one-way
ANOVA was applied when comparing one dataset to multiple others. P-values from GraphPad
Prism 10.4.1 are presented on the graphs using the following notation: not significant (ns) $p>$
0.1234, *$p≤0.0332$, **$p≤0.0021$, ***$p≤0.0002$, and ****$p<0.0001$. Unless stated otherwise, all
experiments were repeated three times.
849
Figure 1 statistics: p-values: **a**, WT vs *GRAMD1* TKO $p<0.0001$; **b**, WT vs *GRAMD1* TKO (2 hr
25HC) $p=0.8044$, WT vs *GRAMD1* TKO (4 hr 25HC) $p=0.5201$, and WT vs *GRAMD1* TKO (6 hr
25HC) $p=0.2595$; **c**, Control vs LXR inhibitor $p=0.0468$; **d**, Control vs LXR inhibitor (2 hr 25HC)
$p=0.2070$, Control vs LXR inhibitor (4 hr 25HC) $p=0.0667$ and Control vs LXR inhibitor (6 hr
25HC) $p=0.0233$; **i**, WT vs NPC1i $p=0.0160$ and WT vs *ACC1* KO (NPC1i) $p=0.0608$; **j**, WT vs
NPC1 KO $p=0.0124$ **k**, WT vs *ACC1* KO (3 hr 25HC) $p=0.0022$, WT vs *ACC1* KO (4.5 hr 25HC)
$p=0.0006$ and WT vs *ACC1* KO (6 hr 25HC) $p<0.0001$; **l** Control vs ACC1i (3 hr 25HC)
$p=0.0552$, Control vs ACC1i (4.5 hr 25HC) $p=0.0236$ and Control vs ACC1i (6 hr 25HC)
$p=0.0087$; **m**, 3T3 control vs ACC1i (3 hr 25HC) $p=0.0097$, 3T3 control vs ACC1i (4.5 hr 25HC)
$p=0.0069$ and 3T3 control vs ACC1i (6 hr 25HC) $p=0.0222$, 293T control vs ACC1i (3 hr 25HC)
$p=0.1192$, 293T control vs ACC1i (4.5 hr 25HC) $p=0.0452$ and 293T control vs ACC1i (6 hr
25HC) $p=0.0208$.
862
863   Figure 2 statistics: p-values are: **b**, WT vs *ACC1* KO $p<0.0001$; **c,** WT HAP1 untreated vs ACC1i
864    $p<0.0001$, 293T untreated vs ACC1i $p<0.0001$ and 3T3 untreated vs ACC1i $p<0.0001$; **d**,
865   untreated vs CP-640186 $p<0.0001$, untreated vs Firsocostat $p<0.0001$; **e**, WT vs KO (0.2

μg/mL) p=0.1871, WT vs KO (0.5 μg/mL) p=0.1171, WT vs KO (1.3 μg/mL) p=0.0024, WT vs KO (3.2 μg/mL) p=0.0056, WT vs KO (8 μg/mL) p=0.0344 and WT vs KO (20 μg/mL) p=0.7060; **h**, WT untreated vs 25HC p=0.0061 and *ACAT1/2* DKO untreated vs 25HC p=0.5563; **i**, WT vs *ACAT1/2* DKO (3 hr 25HC) p=0.1164, WT vs *ACAT1/2* DKO (4.5 hr 25HC) p=0.0176 and WT vs *ACAT1/2* DKO (6 hr 25HC) p=0.0001; **j**, *ACAT1/2* DKO control vs *ACAT1/2* DKO + ACC1i (2 hr 25HC) p=0.0012, *ACAT1/2* DKO control vs *ACAT1/2* DKO + ACC1i (4 hr 25HC) p=0.0003, and *ACAT1/2* DKO control vs *ACAT1/2* DKO + ACC1i (6 hr 25HC) p=0.0016; **l**, *ACAT1/2* DKO vs *ACC1/ACAT1/2* TKO (3 hr 25HC) p<0.0001, *ACAT1/2 DKO* vs *ACC1/ACAT1/2* TKO (4.5 hr 25HC) p<0.0001 and *ACAT1/2* DKO vs *ACC1/ACAT1/2* TKO (6 hr 25HC) p<0.0001; **m**, WT untreated vs *ACAT1/2* DKO untreated p=0.8348, *ACAT1/2* DKO vs *ACAT1/2* DKO + ACC1i p<0.0001; **n**, WT untreated vs 25HC p<0.0001, ACC1 KO untreated vs 25HC p=0.6349.

Figure 3 statistics: p-values are: **a**, WT 0 hr vs 3hr ACC1i p=0.0005, WT 0 hr vs 6hr ACC1i p<0.0001, WT 0 hr vs 24hr ACC1i p<0.0001, and WT HAP1 0 hr vs 48hr ACC1i p<0.0001; **h**, WT control vs ACSLi (2 hr 25HC) p=0.0116, WT control vs ACSLi (4 hr 25HC) p=0.0015 and WT control vs ACSLi (6 hr 25HC) p=0.0077; **i**, WT control vs ACSLi (2 hr ACC1i) p=0.0565, WT control vs ACSLi (4 hr ACC1i) p=0.0294 and WT control vs ACSLi (6 hr ACC1i) p=0.0273; **j,** control untreated vs oleic acid p=0.0050, ACC1i vs ACC1i + oleic acid p=0.0006, and ACSLi vs ACSLi + oleic acid p=0.0502 and **k**, control untreated vs oleic acid p=0.9312, ACC1i vs ACC1i + oleic acid p=0.0006, and ACSLi vs ACSLi + oleic acid p=0.0278.

Figure 4 statistics: p-values are: **c**, WT vs *ATGL* KO p= 0.1690; **d**, WT vs *ATGL* KO p<0.0001; **f**, WT untreated vs ACC1i p=0.0362 and *ATGL* KO untreated vs ACC1i p= 0.6774; **g**, WT vs *ATGL* KO (2 hr ACC1i) p= 0.1746, WT vs *ATGL* KO (4 hr ACC1i) p= 0.0511 and WT vs *ATGL* KO (6 hr ACC1i) p= 0.0121; and **h**, control vs ATGLi (2 hr ACC1i) p=0.0257, control vs ATGLi (4hr ACC1i) p=0.0079 and control vs ATGLi (6hr ACC1i) p=0.0138.

Figure 5 statistics: p-values are: **b**, untreated vs ACSLi p<0.0001, untreated vs ATGLi p<0.0001, ATGLi vs ATGLi + ACSLi p<0.0001; **c**, untreated vs ACSLi p<0.0001, untreated vs ATGLi p=0.0011, and ATGLi vs ATGLi + ACSLi p<0.0001; **g**, untreated vs ACC1i p<0.0001, untreated vs AMPKi p<0.0001, and AMPKi vs ACC1i + AMPKi p<0.0001; **h**, untreated vs ACC1i p<0.0001, untreated vs AMPKi p=0.0005, and AMPKi vs ACC1i + AMPKi p<0.0001; **i**, *ACAT1/2* DKO untreated vs ACC1i p<0.0001, *ACAT1/2* DKO untreated vs AMPKi p<0.0001, and *ACAT1/2* DKO AMPKi vs ACC1i+ AMPKi p<0.0001; **j**, *ACAT1/2* DKO untreated vs ACC1i p<0.0001, *ACAT1/2* DKO untreated vs AMPKi p<0.0001, and *ACAT1/2* DKO AMPKi vs ACC1i + AMPKi p<0.0001.

Figure 6 statistics: p-values are: **b**, WT vs *Acc* dLKO p<0.0001; **c**, WT vs *Acc* dLKO p=0.0022; **d**, WT vs *Acc* dLKO p=0.0005; **f**, WT vs *Acc* dLKO p=0.0004; **j**, WT vs *Acc* dLKO p<0.0001; **k**, WT vs *Acc* dLKO p=0.1915 and **l**, WT vs *Acc* dLKO p=0.0434.

References

1.   Kumar, G. A., Jafurulla, M. & Chattopadhyay, A. The membrane as the gatekeeper of

infection: Cholesterol in host-pathogen interaction. *Chem. Phys. Lipids* **199**, 179–185
(2016).
2. Goluszko, P. & Nowicki, B. Membrane cholesterol: a crucial molecule affecting interactions
of microbial pathogens with mammalian cells. *Infect. Immun.* **73**, 7791–7796 (2005).
3. Radhakrishnan, A., Rohatgi, R. & Siebold, C. Cholesterol access in cellular membranes
controls Hedgehog signaling. *Nat. Chem. Biol.* **16**, 1303–1313 (2020).
4. Hussain, G. *et al.* Role of cholesterol and sphingolipids in brain development and
neurological diseases. *Lipids Health Dis.* **18**, 26 (2019).
5. Lönnfors, M., Doux, J. P. F., Killian, J. A., Nyholm, T. K. M. & Slotte, J. P. Sterols have
higher affinity for sphingomyelin than for phosphatidylcholine bilayers even at equal acyl-
chain order. *Biophys. J.* **100**, 2633–2641 (2011).
6. Demel, R. A., Jansen, J. W., van Dijck, P. W. & van Deenen, L. L. The preferential
interaction of cholesterol with different classes of phospholipids. *Biochim. Biophys. Acta*
465, 1–10 (1977).
7. Slotte, J. P. & Bierman, E. L. Depletion of plasma-membrane sphingomyelin rapidly alters
the distribution of cholesterol between plasma membranes and intracellular cholesterol
pools in cultured fibroblasts. *Biochem. J* **250**, 653–658 (1988).
8. Endapally, S. *et al.* Molecular Discrimination between Two Conformations of Sphingomyelin
in Plasma Membranes. *Cell* **176**, 1040–1053.e17 (2019).
9. Kinnebrew, M. *et al.* Cholesterol accessibility at the ciliary membrane controls hedgehog
signaling. *Elife* **8**, (2019).
10. Das, A., Brown, M. S., Anderson, D. D., Goldstein, J. L. & Radhakrishnan, A. Three pools of
plasma membrane cholesterol and their relation to cholesterol homeostasis. *Elife* **3**, (2014).
11. Infante, R. E. & Radhakrishnan, A. Continuous transport of a small fraction of plasma
membrane cholesterol to endoplasmic reticulum regulates total cellular cholesterol. *Elife* **6**,
(2017).
12. Radhakrishnan, A., Goldstein, J. L., McDonald, J. G. & Brown, M. S. Switch-like control of
SREBP-2 transport triggered by small changes in ER cholesterol: A delicate balance. *Cell*
Metab. **8**, 512–521 (2008).
13. Sun, L.-P., Seemann, J., Goldstein, J. L. & Brown, M. S. Sterol-regulated transport of
SREBPs from endoplasmic reticulum to Golgi: Insig renders sorting signal in Scap
inaccessible to COPII proteins. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 6519–6526 (2007).
14. Brown, M. S. & Goldstein, J. L. Cholesterol feedback: from Schoenheimer's bottle to Scap's
MELADL. *J. Lipid Res.* **50 Suppl**, S15–27 (2009).
15. Brown, M. S. & Goldstein, J. L. The SREBP pathway: regulation of cholesterol metabolism
by proteolysis of a membrane-bound transcription factor. *Cell* **89**, 331–340 (1997).
16. Abrams, M. E. *et al.* Oxysterols provide innate immunity to bacterial infection by mobilizing
cell surface accessible cholesterol. *Nat. Microbiol.* **5**, 929–942 (2020).
17. Zhou, Q. D. *et al.* Interferon-mediated reprogramming of membrane cholesterol to evade
bacterial toxins. *Nat. Immunol.* **21**, 746–755 (2020).
18. Wang, S. *et al.* Cholesterol 25-Hydroxylase inhibits SARS-CoV-2 and other coronaviruses
by depleting membrane cholesterol. *EMBO J.* **39**, e106057 (2020).
19. Zhao, J., Chen, J., Li, M., Chen, M. & Sun, C. Multifaceted functions of CH25H and 25HC
to modulate the lipid metabolism, immune responses, and broadly antiviral activities.

Viruses **12**, (2020).

20. Heisler, D. B. *et al.* A concerted mechanism involving ACAT and SREBPs by which oxysterols deplete accessible cholesterol to restrict microbial infection. *Elife* **12**, (2023).

21. Lichtenstein, A. H. & Brecher, P. Esterification of cholesterol and 25-hydroxycholesterol by rat liver microsomes. *Biochim. Biophys. Acta* **751**, 340–348 (1983).

22. Lange, Y., Ye, J. & Strebel, F. Movement of 25-hydroxycholesterol from the plasma membrane to the rough endoplasmic reticulum in cultured hepatoma cells. *J. Lipid Res.* **36**, 1092–1097 (1995).

23. Liu, Y. *et al.* Activation of liver X receptor plays a central role in antiviral actions of 25-hydroxycholesterol. *J. Lipid Res.* **59**, 2287–2296 (2018).

24. Adams, C. M. *et al.* Cholesterol and 25-hydroxycholesterol inhibit activation of SREBPs by different mechanisms, both involving SCAP and Insigs. *J. Biol. Chem.* **279**, 52772–52780 (2004).

25. Goldstein, J. L., DeBose-Boyd, R. A. & Brown, M. S. Protein sensors for membrane sterols. *Cell* **124**, 35–46 (2006).

26. Venkateswaran, A. *et al.* Control of cellular cholesterol efflux by the nuclear oxysterol receptor LXR alpha. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 12097–12102 (2000).

27. Wong, J., Quinn, C. & Brown, A. J. SREBP-2 positively regulates transcription of the cholesterol efflux gene, ABCA1, by generating oxysterol ligands for LXR. *Biochem. J.* **400**, 485–491 (2006).

28. Ercan, B., Naito, T., Koh, D. H. Z., Dharmawan, D. & Saheki, Y. Molecular basis of accessible plasma membrane cholesterol recognition by the GRAM domain of GRAMD1b. *EMBO J.* **40**, e106524 (2021).

29. Naito, T. & Saheki, Y. GRAMD1-mediated accessible cholesterol sensing and transport. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* **1866**, 158957 (2021).

30. Naito, T. *et al.* Regulation of cellular cholesterol distribution via non-vesicular lipid transport at ER-Golgi contact sites. *Nat. Commun.* **14**, 5867 (2023).

31. Ferrari, A. *et al.* Aster-dependent nonvesicular transport facilitates dietary cholesterol uptake. *Science* **382**, eadf0966 (2023).

32. Soccio, R. E. *et al.* The cholesterol-regulated StarD4 gene encodes a StAR-related lipid transfer protein with two closely related homologues, StarD5 and StarD6. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 6943–6948 (2002).

33. Soccio, R. E. The STARD4 subfamily: STARD4 and STARD5 in cholesterol metabolism. in *Cholesterol Transporters of the START Domain Protein Family in Health and Disease* 139–171 (Springer New York, New York, NY, 2014).

34. Subtil, A. *et al.* Acute cholesterol depletion inhibits clathrin-coated pit budding. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 6775–6780 (1999).

35. Lippincott-Schwartz, J. & Phair, R. D. Lipids and cholesterol as regulators of traffic in the endomembrane system. *Annu. Rev. Biophys.* **39**, 559–578 (2010).

36. Möbius, W. *et al.* Recycling compartments and the internal vesicles of multivesicular bodies harbor most of the cholesterol found in the endocytic pathway: Cholesterol in the endocytic pathway. *Traffic* **4**, 222–231 (2003).

37. Linder, M. D. *et al.* Rab8-dependent recycling promotes endosomal cholesterol removal in normal and sphingolipidosis cells. *Mol. Biol. Cell* **18**, 47–56 (2007).

38. Johnson, K. A. & Radhakrishnan, A. The use of anthrolysin O and ostreolysin A to study
cholesterol in cell membranes. *Methods Enzymol.* **649**, 543–566 (2021).
39. Endapally, S., Infante, R. E. & Radhakrishnan, A. Monitoring and modulating intracellular
cholesterol trafficking using ALOD4, a cholesterol-binding protein. *Methods Mol. Biol.* **1949**,
153–163 (2019).
40. Wakil, S. J. & Abu-Elheiga, L. A. Fatty acid metabolism: target for metabolic syndrome. *J.*
Lipid Res. **50 Suppl**, S138–43 (2009).
41. Li, Q. *et al.* CircACC1 regulates assembly and activation of AMPK complex under metabolic
stress. *Cell Metab.* **30**, 157–173.e7 (2019).
42. Lu, A. *et al.* CRISPR screens for lipid regulators reveal a role for ER-bound SNX13 in
lysosomal cholesterol export. *J. Cell Biol.* **221**, (2022).
43. Russo, L. *et al.* Cholesterol 25-hydroxylase (CH25H) as a promoter of adipose tissue
inflammation in obesity and diabetes. *Mol. Metab.* **39**, 100983 (2020).
44. Raselli, T. *et al.* The oxysterol synthesising enzyme CH25H contributes to the development
of intestinal fibrosis. *J. Crohns. Colitis* **13**, 1186–1200 (2019).
45. Toral-Rios, D. *et al.* Cholesterol 25-hydroxylase mediates neuroinflammation and
neurodegeneration in a mouse model of tauopathy. *J. Exp. Med.* **221**, (2024).
46. Nguyen, C., Saint-Pol, J., Dib, S., Pot, C. & Gosselet, F. 25-Hydroxycholesterol in health
and diseases. *J. Lipid Res.* **65**, 100486 (2024).
47. Harriman, G. *et al.* Acetyl-CoA carboxylase inhibition by ND-630 reduces hepatic steatosis,
improves insulin sensitivity, and modulates dyslipidemia in rats. *Proc. Natl. Acad. Sci. U. S.*
A. **113**, E1796–805 (2016).
48. Luo, D.-X. *et al.* Targeting acetyl-CoA carboxylases: small molecular inhibitors and their
therapeutic potential. *Recent Pat. Anticancer Drug Discov.* **7**, 168–184 (2012).
49. Zhang, H., Tweel, B., Li, J. & Tong, L. Crystal structure of the carboxyltransferase domain of
acetyl-coenzyme A carboxylase in complex with CP-640186. *Structure* **12**, 1683–1691
(2004).
50. Das, A., Goldstein, J. L., Anderson, D. D., Brown, M. S. & Radhakrishnan, A. Use of mutant
125I-perfringolysin O to probe transport and organization of cholesterol in membranes of
animal cells. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 10580–10585 (2013).
51. Ross, A. C., Go, K. J., Heider, J. G. & Rothblat, G. H. Selective inhibition of acyl coenzyme
A:cholesterol acyltransferase by compound 58-035. *J. Biol. Chem.* **259**, 815–819 (1984).
52. Liu, Y. *et al.* 25-Hydroxycholesterol activates the expression of cholesterol 25-hydroxylase
in an LXR-dependent mechanism. *J. Lipid Res.* **59**, 439–451 (2018).
53. Alberts, A. Discovery, biochemistry and biology of lovastatin. *Am. J. Cardiol.* **62**, 10J–15J
(1988).
54. Peterson, C. W. H., Deol, K. K., To, M. & Olzmann, J. A. Optimized protocol for the
identification of lipid droplet proteomes using proximity labeling proteomics in cultured
human cells. *STAR Protoc.* **2**, 100579 (2021).
55. Bersuker, K. *et al.* A proximity labeling strategy provides insights into the composition and
dynamics of lipid droplet proteomes. *Dev. Cell* **44**, 97–112.e7 (2018).
56. Bacci, M. *et al.* Acetyl-CoA carboxylase 1 controls a lipid droplet-peroxisome axis and is a
vulnerability of endocrine-resistant ER+ breast cancer. *Sci. Transl. Med.* **16**, eadf9874
(2024).

57. Hsieh, K. *et al.* Perilipin family members preferentially sequester to either triacylglycerol-
specific or cholesteryl-ester-specific intracellular lipid storage droplets. *J. Cell Sci.* **125**,
4067–4076 (2012).

58. Dumesnil, C. *et al.* Cholesterol esters form supercooled lipid droplets whose nucleation is
facilitated by triacylglycerols. *Nat. Commun.* **14**, 915 (2023).

59. Majchrzak, M. *et al.* Perilipin membrane integration determines lipid droplet heterogeneity in
differentiating adipocytes. *Cell Rep.* **43**, 114093 (2024).

60. Ahmadian, M. *et al.* Desnutrin/ATGL is regulated by AMPK and is required for a brown
adipose phenotype. *Cell Metab.* **13**, 739–748 (2011).

61. Mathiowetz, A. J. *et al.* CLCC1 promotes hepatic neutral lipid flux and nuclear pore complex
assembly. *bioRxivorg* (2024) doi:10.1101/2024.06.07.597858.

62. Roberts, M. A. *et al.* Parallel CRISPR-Cas9 screens identify mechanisms of PLIN2 and lipid
droplet regulation. *Dev. Cell* **58**, 1782–1800.e10 (2023).

63. Zimmermann, R. *et al.* Fat mobilization in adipose tissue is promoted by adipose
triglyceride lipase. *Science* **306**, 1383–1386 (2004).

64. Grabner, G. F. *et al.* Small-molecule inhibitors targeting lipolysis in human adipocytes. *J.*
Am. Chem. Soc. **144**, 6237–6250 (2022).

65. Fang, C. *et al.* The AMPK pathway in fatty liver disease. *Front. Physiol.* **13**, 970292 (2022).

66. Hardie, D. G., Schaffer, B. E. & Brunet, A. AMPK: An energy-sensing pathway with multiple
inputs and outputs. *Trends Cell Biol.* **26**, 190–201 (2016).

67. Zhou, G. *et al.* Role of AMP-activated protein kinase in mechanism of metformin action. *J.*
Clin. Invest. **108**, 1167–1174 (2001).

68. Kim, C.-W. *et al.* Acetyl CoA carboxylase inhibition reduces hepatic steatosis but elevates
plasma triglycerides in mice and humans: A bedside to bench investigation. *Cell Metab.* **26**,
394–406.e6 (2017).

69. Schwarz, M., Russell, D. W., Dietschy, J. M. & Turley, S. D. Marked reduction in bile acid
synthesis in cholesterol 7α-hydroxylase-deficient mice does not lead to diminished tissue
cholesterol turnover or to hypercholesterolemia. *J. Lipid Res.* **39**, 1833–1843 (1998).

70. Gruffat, D., Durand, D., Graulet, B. & Bauchart, D. Regulation of VLDL synthesis and
secretion in the liver. *Reproduction Nutrition Development* **36**, 375–389 (1996).

71. Brown, M. S., Radhakrishnan, A. & Goldstein, J. L. Retrospective on cholesterol
homeostasis: The central role of Scap. *Annu. Rev. Biochem.* **87**, 783–807 (2018).

72. Edwards, P. A., Tabor, D., Kast, H. R. & Venkateswaran, A. Regulation of gene expression
by SREBP and SCAP. *Biochim. Biophys. Acta* **1529**, 103–113 (2000).

73. Kim, J. B. *et al.* Nutritional and insulin regulation of fatty acid synthetase and leptin gene
expression through ADD1/SREBP1. *J. Clin. Invest.* **101**, 1–9 (1998).

74. Zhang, Y., Yin, L. & Hillgartner, F. B. SREBP-1 integrates the actions of thyroid hormone,
insulin, cAMP, and medium-chain fatty acids on ACCalpha transcription in hepatocytes. *J.*
Lipid Res. **44**, 356–368 (2003).

75. Kim, H.-J., Miyazaki, M. & Ntambi, J. M. Dietary cholesterol opposes PUFA-mediated
repression of the stearoyl-CoA desaturase-1 gene by SREBP-1 independent mechanism. *J.*
Lipid Res. **43**, 1750–1757 (2002).

76. Sekiya, M. *et al.* Polyunsaturated fatty acids ameliorate hepatic steatosis in obese mice by
SREBP-1 suppression. *Hepatology* **38**, (2003).

77. Walther, T. C. & Farese, R. V., Jr. Lipid droplets and cellular lipid metabolism. *Annu. Rev.*
Biochem. **81**, 687–714 (2012).
78. Olzmann, J. A. & Carvalho, P. Dynamics and functions of lipid droplets. *Nat. Rev. Mol. Cell*
Biol. **20**, 137–155 (2019).
79. Naito, T. *et al.* Movement of accessible plasma membrane cholesterol by the GRAMD1 lipid
transfer protein complex. *Elife* **8**, (2019).
80. Peng, W. *et al.* Endosomal trafficking participates in lipid droplet catabolism to maintain lipid
homeostasis. *Nat. Commun.* **16**, 1917 (2025).
81. Forwell, A. L. *et al.* Analysis of CH25H in multiple sclerosis and neuromyelitis optica. *J.*
Neuroimmunol. **291**, 70–72 (2016).
82. Haney, M. S. *et al.* APOE4/4 is linked to damaging lipid droplets in Alzheimer's disease
microglia. *Nature* **628**, 154–161 (2024).
83. Zadoorian, A., Du, X. & Yang, H. Lipid droplet biogenesis and functions in health and
disease. *Nat. Rev. Endocrinol.* **19**, 443–459 (2023).
84. Carette, J. E. *et al.* Haploid genetic screens in human cells identify host factors used by
pathogens. *Science* **326**, 1231–1235 (2009).
85. Khandwala, C. B. *et al.* Direct ionic stress sensing and mitigation by the transcription factor
NFAT5. *bioRxivorg* (2023) doi:10.1101/2023.09.23.559074.
86. Lebensohn, A. M. *et al.* Comparative genetic screens in human cells reveal new regulatory
mechanisms in WNT signaling. *Elife* **5**, (2016).
87. Shimano, H. *et al.* Overproduction of cholesterol and fatty acids causes massive liver
enlargement in transgenic mice expressing truncated SREBP-1a. *J. Clin. Invest.* **98**, 1575–
1584 (1996).
88. Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**,
2281–2308 (2013).
89. Ikonen, E. & Zhou, X. Cholesterol transport between cellular membranes: A balancing act
between interconnected lipid fluxes. *Dev. Cell* **56**, 1430–1436 (2021).
90. Zhou, F. *et al.* Filament structures unveil the dynamic organization of human acetyl-CoA
carboxylase. *Sci. Adv.* **10**, eado4880 (2024).
91. Hunkeler, M. *et al.* Structural basis for regulation of human acetyl-CoA carboxylase. *Nature*
558, 470–474 (2018).

1115

1116

1117

1118

1119

1120

1121

1122

a,b, PCR analysis of *GRAMD1A/B/C* TKO clones (**a**) and *GRAMD1B* KO clones (**b**). **c,d**, Plasma membrane (PM) cholesterol analyzed by mNG-ALOD4 staining followed by flow cytometry in wild-type (WT) HAP1 and *GRAMD1B* KO clones either left untreated (**c**) or treated with 25HC for the indicated times (**d**). **e**, quantitative real-time PCR analysis of *Abca1* mRNA levels normalized to *Gapdh* levels from 3T3 cells treated for 20 hrs with LXRi (1 µM GSK2033). **f,g**, PM cholesterol analyzed by flow cytometry analysis of mNG-ALOD4 staining in 3T3 cells treated for 20 hrs with or without 1 µM of LXRi alone (**f**) or followed by treatment with 25-hydroxycholesterol (25HC) for the indicated times (**g**). **h**, Domain architecture of ACC1 (top) including the biotin carboxylase domain (BC), biotin transferase domain (BT), biotin carboxyl carrier protein domain (BCCP), central domain (CD), and the carboxyltransferase domain (CT) [90,91]. Residues implicated in catalysis are shown in dark orange. Exon map of ACC1 (bottom) with exon colors matching the domain they encode. **i**, The number of insertions mapped per intron are normalized to intron length in base pairs (bp). **j**, Western blot (top) and PCR analysis (bottom) of WT HAP1 and *ACC1* KO clonal cell lines. **k,l**, mNG-ALOD4 flow cytometry analysis of PM cholesterol in WT HAP1 and *ACC1* KO cells made with CRISPR sgRNAs targeting the indicated exons either left untreated (**k**) or treated with 30 µM ACC1i (Firsocostat) for 20 hrs (**l**). **m**, Table showing the number of clones isolated from two 96-well dishes and the percentage of those clones that were found to be knockouts with PCR analysis. **n**, Western blot analysis of WT and *NPC1* KO HAP1 cell lines. **o,p**, Western blot analysis (**o**) or mNG-ALOD4 flow cytometry analysis of PM cholesterol (**p**) of *Npc1*[-/-] mouse embryonic fibroblasts (MEFs) or *Npc1*[-/-] MEFs with NPC1 stably re-introduced using MSCV retrovirus, and then treated with or without ACC1i (30 µM Firsocostat) for 20 hrs. Each data point is the average of three biological replicates (**d** and **g**). Cells in **k** and **l** were grown in 5% lipoprotein depleted serum for 16 hours prior to analysis. Statistical significance was determined by a Student's t-test with a Welch's correction (**c-g** and **l**) or an Ordinary one-way ANOVA (**k** and **p**). Exact *p*-values: **c**, WT vs *GRAMD1B* KO *p*=0.0043; **d**, WT vs *GRAMD1B* KO (2 hr 25HC) *p*=0.9961, WT vs *GRAMD1B* KO (4 hr 25HC) *p*=0.0847, and WT vs *GRAMD1B* KO (6 hr 25HC) *p*<0.0001; **e**, untreated vs LXRi *p*<0.0001; **f**, control vs LXRi *p*<0.0001; **g**, control vs LXRi (2 hr 25HC) *p*=0.0064, control vs LXRi (4 hr 25HC) *p*=0.0132 and control vs LXRi (6 hr 25HC) *p*=0.9640; **k**, WT vs *ACC1* KO exon 11 *p*<0.0001, WT vs *ACC1* KO exon 21 *p*<0.0001, WT vs *ACC1* KO exon 31 *p*<0.0001, WT vs *ACC1* KO exon 41 *p*<0.0001 and WT vs *ACC1* KO exon 52 *p*<0.0001; **l**, WT untreated vs ACC1i *p*=0.0076, *ACC1* KO exon 11 untreated vs ACC1i *p*=0.1371 and *ACC1* KO exon 52 untreated vs ACC1i *p*=0.0563; and **p**, *Npc1*[-/-] untreated vs ACC1i *p*<0.0001 and *Npc1*[-/-] vs *Npc1*[-/-]:NPC1 *p*=0.0004.

Extended data Fig. 2

a, Schematic demonstrating that sphingomyelin (SM, purple) binds to cholesterol in membranes. ACC1 loss could elevate cholesterol by 1) increasing cholesterol or by 2) reducing SM levels. **b**, Mass spectrometry analysis of SM levels in wild-type (WT) HAP1 cells either left untreated or treated with ACC1i (30 μM Firsocostat) for 16 hrs. **c**, OlyA-EGFP flow cytometry analysis of WT HAP1 cells left untreated or treated with 0.3 mM MβCD: cholesterol or 150 mU/mg SMase for 30 minutes, or 30 μM ACC1i for 16 hrs. **d**, *ACC1* KO cells or WT HAP1 cells were left untreated or treated for 16 hrs with 30 μM ACC1i. Cells were then treated with or without 150 mU/mg SMase in 5% FBS for 30 minutes and stained with mNG-ALOD4 and analyzed by flow cytometry. **e**, Quantification of esterified cholesterol in WT HAP1 cells treated with ACATi (60 μM SZ58-035) for 16 hrs or left untreated, followed by a 6 hr treatment with or without 4 μM 25HC using the Amplex Red assay. **f**, Flow cytometry analysis of mNG-ALOD4 staining in WT HAP1 cells treated with or without 60 μM ACATi for 16 hrs, followed by 4 μM 25HC for the times indicated. **g**, **h**, mNG-ALOD4 flow cytometry analysis of *ACC1* KO cells left untreated or treated for 16 hours with ACATi (60 μM SZ58-035) followed by 4 μM 25HC treatment for the indicated times (**g**) and WT HAP1 cells grown in 5% FBS and treated with HMGCRi (10 μM Lovastatin) for the indicated times (**h**). Statistical significance was determined by a Student's t-test with a Welch's correction (**b** and **d-g**) or an Ordinary one-way ANOVA (**c** and **h**). Exact *p*-values: **b**, untreated vs ACC1i *p*=0.6634; **c**, untreated vs cholesterol *p*<0.0001, untreated vs SMase *p*=0.0038, untreated vs ACC1i *p*=0.0036; **d**, control with vs without SMase *p*=0.0002, ACC1i with vs without SMase *p*<0.0001, *ACC1* KO with vs without SMase *p*<0.0001; **e**, control untreated vs 25HC *p*=0.0013 and ACATi vs ACATi + 25HC *p*=0.6469; **f**, control vs ACATi (3 hr 25HC) *p*=0.0004, control vs ACATi (4.5 hr 25HC) *p*=0.0024 and control vs ACATi (6 hr 25HC) *p*=0.0146; **g**, *ACC1* KO vs *ACC1* KO + ACATi (3 hr 25HC) *p*=0.0134, *ACC1* KO vs *ACC1* KO + ACATi (4.5 hr 25HC) *p*=0.0654 and *ACC1* KO vs *ACC1* KO + ACATi (6 hr 25HC) *p*=0.0735; **h**, 0 vs 2 hrs HMGCRi *p*=0.2028, 0 vs 4 hrs HMGCRi *p*=0.0137 and 0 vs 6 hrs HMGCRi *p*=0.0495.

1125

1126

1127

1128

Extended data Fig. 3

a, 3T3 cells were treated with 30 µM ACC1i (Firsocostat) for the indicated times in 5% FBS, then analyzed by flow cytometry after staining with mNG-ALOD4. **b**, mNG-ALOD4 flow cytometry analysis of WT HAP1 cells cultured in 5% FBS and either left untreated or treated with HMGCRi (10 µM Lovastatin), ACC1i (30 µM Firsocostat) or both for 16 hrs. **c**, mNG-ALOD4 flow cytometry analysis of WT HAP1 cells and *ACC1* KO cells grown in 5% FBS and either left untreated or treated with HMGCRi (10 µM Lovastatin) for 16 hrs. **d**, Western blot analysis of fractionated LDAH-GFP expressing HAP1 cells with various cellular fractions including whole cell (WC), nucleus (Nuc.; detected by LaminA/C), cytoplasm (Cyt.; detected by GAPDH), plasma membrane (PM; detected by LDLR), and lipid droplets (LD; detected by LDAH-GFP). Cells were either left untreated or treated with ACC1i (30 µM Firsocostat) for 24 hrs. **e**, Quantification of LD numbers per cell in WT HAP1 cells treated with or without ACC1i (30 µM Firsocostat) for 16 hrs (left) or WT HAP1 cells compared to *ACC1* KO HAP1 cells (right). The solid middle line shows the median and dashed lines show the interquartile range. **f**, Microscopy images of LDs in WT HAP1 cells left untreated or treated with DGAT1 and DGAT2 inhibitors (DGATi) for 16 hrs followed by 150 µM oleic acid (OA) treatment for 6 hrs in 5% FBS stained with Bodipy 493/503. Scale bar is 10 microns. **g**, Western blot analysis of WT HAP1 cells compared to *FASN* KO HAP1 clones. **h**, Microscopy images of LDs in WT HAP1 and *FASN* KO cells stained with Bodipy 493/503 and DAPI (blue). Scale bar is 10 microns. **i,j**, Flow cytometry analysis of mNG-ALOD4 PM staining in WT HAP1 and *FASN* KO clones grown in 5% LDS media for 24 hrs left untreated (**i**) or then treated with 4 µM 25HC for the indicated times (**j**). **k**, mNG-ALOD4 flow cytometry analysis of PM accessible cholesterol (blue curves) and the corresponding fluorescence microscopy quantification of the number of LDs per cell (purple curves) in HAP1 cells grown in 5% FBS and then treated with FASNi (10 µM C75) for the indicated times in 5% LDS. LDs were quantified by taking the mean number of LDs per cell in four fields of view with ~75 cells per field. **l**, Microscopy images of FASN localization relative to LD staining with Bodipy 493:503. Scale bar is 5 microns. Statistical significance was determined by a Student's t-test with a Welch's correction (**a-c,e,i** and **j**). Exact *p*-values: **a,** 0 vs 15 min ACC1i *p*=0.0419; **b,** WT HMGCRi vs HMGCRi + ACC1i *p*=0.0008 ; **c,** WT HMGCRi vs *ACC1* KO HMGCRi *p*<0.0001; **e,** WT untreated vs ACC1i *p*<0.0001 and WT vs *ACC1* KO *p*<0.0001; **i,** WT vs *FASN* KO *p*<0.0001; **j,** WT vs *FASN* KO (2 hr 25HC) *p*=0.0001, WT vs *FASN* KO (4 hr 25HC) *p*<0.0001 and WT vs *FASN* KO (6 hr 25HC) *p*=0.0003.

Extended data Fig. 4

1130

a, Flow cytometry analysis of PM mNG-ALOD4 staining. WT HAP1 cells were left untreated, treated with ACC1i (30 µM Firsocostat), or treated with both ACC1i and ATGLi (50 µM NG-497), followed by treatments with 4 µM 25HC for the indicated times. **b**, Microscopy images of LDs in WT HAP1, untreated *ACAT1/2* DKO and *ACAT1/2* DKO treated with ACC1i (30 µM Firsocostat) and stained with Bodipy 493/503 and DAPI (blue). Scale bar is 10 microns. **c**, Quantification of LD numbers per cell in WT and *ACAT1/2* DKO HAP1 cells cultured in 5% LDS for 24 hrs. **d**, LD area for untreated WT and *ACAT1/2* DKO HAP1 cells. Each violin plot shows the area for LDs measured in three fields of view. **e**, LD quantification for WT and *ACAT* DKO cells, either untreated or treated with ACC1i (30 µM Firsocostat) for 6 hrs. Each data point is the total number of LDs per cell in one field of view. **f**, Flow cytometry analysis of mNG-ALOD4 staining in WT and *ACAT1/2* DKO HAP1 cell lines cultured in 5% LDS for 16 hrs, followed by treatment with ACC1i (30 µM Firsocostat) for the indicated times. Statistical significance was determined by Student's t-test with Welch's correction (**a** and **c-f**). Exact *p*-values: **a,** WT ACC1i vs ACC1i + ATGLi (2 hr 25HC) *p*=0.0196, WT ACC1i vs ACC1i + ATGLi (4 hr 25HC) *p*=0.0256, and WT ACC1i vs ACC1i + ATGLi (2 hr 25HC) *p*=0.0017; **c**, WT vs *ACAT 1/2* DKO *p*=0.0151; **d**, WT vs *ACAT 1/2* DKO *p*=0.0588; **e**, WT untreated vs ACC1i *p*=0.0186 and *ACAT 1/2* DKO untreated vs ACC1i *p*=0.0146; **f**, WT vs *ACAT1/2* DKO (2 hr ACC1i) *p*=0.1519, WT vs *ACAT1/2* DKO (4 hr ACC1i) *p*=0.0099 and WT vs *ACAT1/2* DKO (6 hr ACC1i) *p*=0.0160.

Extended data Fig. 5

% LDs with colocalization: 33%   3%   20%   13%

a,**b**, Quantification of lipid droplet (LD) numbers per cell (**a**) or flow cytometry analysis of PM mNG-ALOD4 staining (**b**) in WT HAP1 cells left untreated, treated with ATGLi (50 µM NG-497), ACC1i (30 µM Firsocostat), or both for 16 hrs. **c**, Microscopy images of ACC1 localization relative to LD staining (Bodipy 493:503) in cells treated with 100 µM oleic acid for 6 hrs, ATGLi (50 µM NG-497) for 16 hrs, or AMPKi (2.5 µM Dorsomorphin) for 16 hrs. The percentage of LDs with ACC1 co-localization was determined by dividing the number of LDs with adjacent ACC1 puncta by the total number of LDs. n>250 LDs for all conditions. Scale bar is 5 microns. Statistical significance was determined by an Ordinary one-way ANOVA (**a** and **b**). Exact *p*-values: **a**, untreated vs ATGLi *p*<0.0001, untreated vs ACC1i *p*<0.0001 and ACC1i vs ACC1i + ATGLi *p*=0.0058; and **b**, untreated vs ATGLi *p*=0.0002, untreated vs ACC1i *p*<0.0001 and ACC1i vs ACC1i + ATGLi *p*<0.0001.

1131

Extended data Fig. 6

1132

1133

1134

1135

1136

1137

1138

a, Quantification of free (unesterified) cholesterol normalized to total protein in WT HAP1 cells treated with ACC1i (30 µM Firsocostat) for the indicated times using the Amplex Red assay. **b**, Quantification of free (unesterified) cholesterol in WT HAP1 cells and *ACC1* KO cells using the Amplex Red assay. Statistical significance was determined by an Ordinary one-way ANOVA (**a**) or a Student's t-test with a Welch's correction (**b**). Exact *p*-values: **a**, untreated vs 2 hrs ACC1i *p*=0.8245, untreated vs 4 hrs ACC1i *p*>0.9999, untreated vs 6 hrs ACC1i *p*=0.5931, untreated vs 24 hrs ACC1i *p*=0.0004, untreated vs 48 hrs ACC1i *p*=0.2352 and **b**, WT vs *ACC1* KO *p*=0.6323.