# Sex-dependent interferon signaling drives female-biased vulnerability in Alzheimer's disease

Verónica López-López,[1] Gerard Iniesta,[1] Marcos Galán-Ganga,[2,3,4] Alejandro Expósito-Coca,[1] Aysha Bhojwani-Cabrera,[1] Marina Guillot-Fernández,[1] Albert Giralt,[2,3,4,5] José P. López-Atalaya[1]

[1]Instituto de Neurociencias, Consejo Superior de Investigaciones Científicas, Universidad Miguel Hernández, Alicante, Spain.

[2]Departament de Biomedicina, Facultat de Medicina, Institut de Neurociències, Universitat de Barcelona, Barcelona, Spain.

[3]Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain.

[4]Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Madrid, Spain.

[5]Production and Validation Centre of Advanced Therapies (Creatio), Faculty of Medicine and Health Science, University of Barcelona, Barcelona, Spain.

Correspondence: José P. López-Atalaya, Instituto de Neurociencias, Consejo Superior de Investigaciones Científicas, Universidad Miguel Hernández, Alicante, Spain. Phone: 34.965.23.3744. Email: jose.lopezatalaya@csic.es.

Conflict of interest: The authors have declared that no conflict of interest exists.

## Abstract

Alzheimer's disease (AD) disproportionately affects women, yet the neurobiological mechanisms underlying this sex bias remain poorly understood. Here, we identify a sex-dependent activation of type I interferon (IFN-I) signaling as a contributor to this disparity. Transcriptomic profiling of brain from AD patients revealed selective enrichment of IFN-I pathway in females. This immune signature was mirrored in the APP/PS1 mouse model of AD, where females exhibited more pronounced amyloid accumulation, neuroinflammation, and neurodegeneration. Acute IFN-I activation reproduced pathological features of AD, whereas chronic IFN-I elevation in APP/PS1 mice aggravated disease progression. In contrast, pharmacological targeting of IFN-I response by inhibiting the cGAS-STING pathway in APP/PS1 female mice reduced neuropathological burden, and preserved cognitive performance. Together, these findings identify interferon signaling as a modifiable and sex-linked driver of AD pathology. Our study uncovers a critical neuroimmune mechanism contributing to female-biased vulnerability and highlights interferon modulation as a promising therapeutic strategy in AD.

**Keywords:** Alzheimer's disease; Sex differences; Type I interferon; Interferon signaling; Neuroinflammation; cGAS–STING pathway; Learning and memory

## Introduction

Alzheimer's disease (AD) is a progressive and ultimately fatal neurodegenerative disorder and the most common cause of dementia worldwide (1–3). With an aging global population, the number of individuals affected by AD is projected to exceed 100 million by 2050, placing an enormous burden on healthcare systems and caregivers (1, 3–5). The neuropathological hallmarks of AD - extracellular amyloid-β plaques, intracellular tau neurofibrillary tangles, and chronic neuroinflammation - are accompanied by cognitive and behavioral impairments that progressively erode autonomy and quality of life (2, 3, 6). Despite decades of intense research, currently approved treatments provide only modest symptomatic relief, and no curative or disease-modifying therapies have yet been established (6–8).

A consistently reported yet underexplored feature of AD is its disproportionate impact on females. Females comprise nearly two-thirds of individuals diagnosed with AD (1, 3, 5), and lifetime risk estimates indicate that approximately 1 in 5 females, compared to 1 in 10 males, will develop the disease (1, 4). Although increased longevity contributes to this imbalance, it does not fully account for the observed differences (9–12). Clinical and neuroimaging studies have revealed that females with AD often exhibit faster cognitive decline, greater neuropathological burden, and more rapid progression from mild cognitive impairment to dementia than their male counterparts (9, 13–18). Biological sex thus acts as an independent modifier of AD risk and trajectory, shaping both susceptibility and disease course.

Despite its clinical significance, the mechanistic basis for sex-related differences in AD remains poorly defined. Historically, biological sex has been underrepresented or insufficiently analyzed in both clinical trials and experimental models (19–22). Many preclinical studies rely exclusively on male animals, or include both sexes without

stratified analysis, thereby missing potentially critical sex-dependent effects (20, 23–25). This limitation not only undermines the translational relevance of preclinical findings but also hinders the development of sex-informed therapeutic strategies.

Addressing this knowledge gap requires a targeted focus on molecular and cellular pathways that are both central to AD pathogenesis and modulated by sex. In this regard, the innate immune system - particularly neuroimmune signaling within glial networks - has emerged as a key candidate. Genome-wide association studies (GWAS) have identified numerous AD susceptibility loci involved in immune regulation, including variants in *TREM2*, *CD33*, and *HLA* genes, among others (26–29). These discoveries have expanded the conceptual framework of AD beyond classical amyloid and tau cascades, implicating immune dysregulation and glial biology as core features of disease pathogenesis (30–34).

In parallel, a growing body of research has shown that immune responses are strongly shaped by biological sex. Females generally exhibit stronger baseline and stimulus-induced immune responses, both peripherally and in the central nervous system (35–39). While this amplified responsiveness may enhance pathogen clearance, it is also linked to greater vulnerability to chronic inflammation and autoimmunity (35, 36, 38). In the context of neurodegeneration, sex-biased immune traits may alter the threshold, magnitude, or persistence of neuroinflammatory responses, potentially influencing susceptibility to and progression of AD (14, 40–42).

Whether such immunological differences translate into distinct molecular trajectories or pathological outcomes in AD remains an open question. Biological sex may shape diverse cellular programs - including glial activation states (43–46), neuronal vulnerability, and synaptic remodeling (47–50) - that collectively influence disease

dynamics. However, these hypotheses require systematic and mechanistic investigation.

Understanding how sex interacts with disease-relevant pathways is essential for

explaining observed clinical disparities and for identifying novel therapeutic targets that

reflect this dimension of disease biology.

In this study, we sought to investigate the molecular mechanisms underlying sex-

associated differences in AD vulnerability. We combined transcriptomic analysis of

*postmortem* human brain tissue with functional studies in established mouse models of

AD. By stratifying human samples by sex, age, and brain region, we identified a

pronounced activation of type I interferon (IFN-I) signaling in females. To explore the

causal role of sex-dependent IFN-I signaling in disease pathogenesis, we employed both

pharmacological and genetic manipulations *in vivo* and assessed the impact of modulating

IFN-I signaling on neuroinflammation, neuropathology, and cognitive performance. This

integrative approach establishes IFN-I signaling as a key contributor to sex-associated

differences in AD vulnerability and highlights its potential as a therapeutic target.

## Results

**Sex-Based Differences in Interferon Pathway Activation Are Associated with Increased Alzheimer's Risk and Prevalence in Women**

Epidemiological data from the Alzheimer's Facts and Figures 2024 report highlight a markedly increased lifetime risk of AD in women (51). At both 45 and 65 years of age, the estimated lifetime risk of developing AD is approximately 20% for women versus 10% for men (51–54). Among individuals aged 65 and older, women account for approximately 61% of all AD cases, while men account for 39% (51, 53, 55). These sex disparities in disease prevalence and risk are illustrated schematically (Fig. 1A).

While several clinical studies have reported faster disease progression and greater pathological burden in females (13, 14, 16, 17), the molecular mechanisms driving this increased vulnerability remain poorly understood. To address this, we leveraged RNA-sequencing data from the parahippocampal gyrus (PHG) of AD patients in the Mount Sinai Brain Bank (MSBB). To minimize potential confounding effects of age, we focused on individuals aged 60 to 90 years, yielding a final cohort of 34 females and 32 males (Fig. 1B and Supplementary Data Table 1).

Principal component analysis (PCA) confirmed robust sex-based separation and revealed one sample annotated as male that clustered instead with female samples (Extended Data Fig. 1A). Further analysis of sex chromosome gene expression - including *XIST* and *TSIX* from the X chromosome, and *UTY*, *RPS4Y1*, and *DDX3Y* from the Y chromosome - confirmed this misclassification (Extended Data Fig. 1B). This sample (marked in bold) was excluded from all subsequent analyses.

Differential gene expression analysis revealed a prominent female-biased transcriptional signature related to interferon signaling in AD patients, including genes

such as *GBP4*, *IFIT2*, *PSMB8*, *PSMB9*, *RTP4*, *B2M*, and *MX1* (Fig. 1C,D). Functional enrichment analysis revealed that "Interferon Alpha Response" was the most significantly enriched "hallmark" gene set (56) in females, followed by "Interferon Gamma Response" and "p53 Pathway" (Fig. 1E). Gene Set Enrichment Analysis (GSEA) (57) further confirmed strong and coordinated upregulation of the interferon alpha response pathway in female AD samples (Fig. 1F).

To place these sex-stratified findings in the broader context of AD pathology, we compared AD versus age-matched control samples without stratifying by sex (n = 108; 42 controls and 66 AD) (Supplementary Data Table 1). In this pooled analysis, functional profiling identified "Epithelial Mesenchymal Transition" as the top enriched pathway, followed by "Interferon Alpha Response" (Extended Data Fig. 1C), in addition to other classical AD-associated pathways such as 'Cholesterol Homeostasis' (Supplementary Data Table 1). However, stratifying by sex revealed a marked shift in the dominant transcriptional programs: in males (n = 55; 23 controls and 32 AD), "Epithelial Mesenchymal Transition" emerged as the principal upregulated pathway, whereas in females (n = 53; 19 controls and 34 AD), "Interferon Alpha Response" became the leading enriched pathway with, "Epithelial Mesenchymal Transition" ranking lower (Extended Data Fig. 1D,E). This divergence underscores the critical influence of sex on the molecular architecture of AD, revealing that the primary immune program engaged by the disease differs fundamentally between males and females.

Next, we examined whether this transcriptional signature was also detectable under non-pathological conditions. To this end, we analyzed transcriptomic data obtained from the PHG of age-matched control individuals (n = 42; 19 females and 23 males) (Supplementary Data Table 1). Functional analysis of sex-associated differentially

expressed genes revealed no significant enrichment of interferon-related pathways (Extended Data Fig. 1F,G). A weak enrichment of the IFN-I response was detected in GSEA (Extended Data Fig. 1H), suggesting that although females may exhibit slightly higher baseline interferon tone, it is the presence of AD pathology that amplifies these sex-dependent differences in a robust and functionally significant manner.

Together, these findings show that female AD patients exhibit a sustained upregulation of IFN-I in the parahippocampal gyrus. This persistent immune activation may contribute to the greater vulnerability and higher prevalence of AD observed in women, highlighting sex-specific neuroimmune mechanisms as potential contributors to disease susceptibility and progression.

**Sex-Specific Differences in Amyloid Pathology and Neurodegeneration in APP/PS1 Mice**

To evaluate whether the APPswe/PSEN1dE9 (APP/PS1) mouse model recapitulates the sex-specific features of human AD, we analyzed male and female transgenic mice at 6 months of age (Fig. 2A). Immunohistochemical staining with the anti–amyloid-β (Aβ) antibody 6E10 revealed significantly higher plaque burden in both the cortex and hippocampus of female mice compared with males (Fig. 2B). Thioflavin T staining confirmed these observations, showing more extensive fibrillar amyloid deposits in females (Fig. 2C). Consistently, immunostaining with the OC antibody, which recognizes β-sheet-rich Aβ conformations, revealed greater signal in female brains (Fig. 2D), further validating increased pathological aggregation. Notably, plaque size was similar between sexes (Extended Data Fig. 2A-C), suggesting that the sex difference primarily reflects increased plaque number or density rather than altered morphology.

Building on these findings, we next examined other neuropathological features

associated with AD (58–62). Immunostaining for c-FOS, a marker of neuronal activity,

revealed reduced expression in the dentate gyrus of female mice (Fig. 2E), indicating

potential functional impairment. To assess inhibitory network integrity, we quantified

perineuronal nets (PNNs), specialized extracellular matrix structures that stabilize

synaptic connectivity (63). The proportion of parvalbumin-positive interneurons

surrounded by PNNs in the CA1 region was significantly lower in females, as visualized

by *Wisteria floribunda* agglutinin (WFA) staining (Fig. 2F,D). In parallel, we observed

increased axonal damage in the female hippocampus, as evidenced by elevated

phosphorylated neurofilament (pNF) staining (Fig. 2G). These converging

histopathological alterations suggest that female APP/PS1 mice experience more severe

neuronal dysfunction and structural disruption.

To determine whether these sex-specific differences extended to the

transcriptional level, we first characterized the global molecular profile of APP/PS1 mice

relative to wild-type controls. Bulk RNA-Seq of hippocampal tissue revealed pronounced

upregulation of transcripts related to the model's transgenes (*App*, *Prnp*) (64) alongside

microglial activation markers (*Clec7a*, *Trem2*, *Cst7, Itgax*) (65–67) (Extended Data Fig.

2E and Supplementary Data Table 3). Functional enrichment analysis indicated

significant activation of pathways including "Cholesterol Homeostasis", "Allograft

Rejection", and "Inflammatory Response" (Extended Data Fig. 2F), confirming that the

APP/PS1 model exhibits the expected immune–metabolic alterations associated with AD.

Building on this baseline characterization, we next examined whether the

transcriptional response to Aβ pathology differed between sexes. PCA revealed limited

separation between wild-type and APP/PS1 males, whereas APP/PS1 females diverged

markedly from wild-type females (Fig. 2H), indicating more extensive transcriptional

remodeling in females at this age (six month old). This suggests that female brains may

undergo a more profound molecular response to Aβ pathology, potentially contributing

to their heightened vulnerability.

These data indicate that female APP/PS1 mice exhibit greater amyloid deposition,

more pronounced neurodegeneration, and broader transcriptomic alterations than males.

Importantly, this model recapitulates the sex-specific vulnerability observed in human

Alzheimer's disease, making it a valuable platform for investigating the biological

mechanisms underlying these differences.

**Enhanced Neuropathology in Female APP/PS1 Mice Is Associated with Increased
Interferon Signaling**

To investigate potential molecular correlates of the more pronounced AD-like

histopathology observed in female APP/PS1 mice, we profiled gene expression of bulk

hippocampal tissue. In line with the differences found in AD patient´s (Fig. 1), differential

gene expression analysis revealed a significant upregulation of inflammatory genes in

females, including IFN-I-stimulated transcripts such as *Irf7*, *B2m*, *Psmb8*, *Psmb9*, *Ifit3*,

*Oas1a*, and *Rtp4* (Fig. 2I and Supplementary Data Table 3).

Given the observed transcriptional signature, we examined glial activation using

GFAP, a marker of astrogliosis and neuroinflammation (58, 68). Immunostaining

revealed higher densities of GFAP-positive astrocytes in the hippocampi of female mice

(Fig. 2J), with similar increases observed in the cortex (Extended Data Fig. 2G). While

this suggested enhanced neuroinflammation in females, GFAP is a broad marker and does

not reveal which specific inflammatory pathways are engaged. To further dissect the

molecular nature of immune activation, we performed functional analysis on significantly

differentially expressed genes using the "hallmark" gene sets collection (56). This

revealed a selective overrepresentation of interferon-related pathways in females,

including "Interferon Alpha Response" and "Interferon Gamma Response" (Fig. 2K),

indicating that the inflammatory profile in females is characterized by a pronounced

activation of antiviral IFN-I signaling. GSEA further supported these findings, revealing

a significant enrichment of the interferon alpha response gene set in APP/PS1 females

(Extended Data Fig. 2H).

To validate these transcriptomic findings at the protein level, we assessed the

expression of Ifitm3, a key effector of the interferon response. Immunohistochemistry

revealed significantly increased numbers of Ifitm3-positive cells in the hippocampus of

females compared with males (Fig. 2L), with similar results observed in the cortex

(Extended Data Fig. 2I).

Overall, our data indicate that the greater neurodegeneration and amyloid

pathology observed in female APP/PS1 mice is accompanied by a selectively heightened

interferon response, a feature that only emerged upon sex-stratified analysis, suggesting

interferon-related mechanisms as a potential driver of sex-specific vulnerability in AD.

**Interferon Induction by Viral-Mimetic Stimulation Is Sufficient to Elicit Alzheimer-**

**Like Histopathology**

Given the previously observed correlation between elevated interferon signaling and

worsened neuropathological features in female APP/PS1 mice, we next examined

whether interferon activation alone is sufficient to induce similar pathological changes.

To test this hyphotesis, we administered a single intraperitoneal dose of the viral mimetic

polyinosinic:polycytidylic acid [poly(I:C), 12 mg/kg] to 3-month-old wild-type C57BL/6

mice and analyzed brain tissue 24- and 72-hours post-injection (Fig. 3A).

First, we assessed the molecular response to poly(I:C) by profiling the

transcriptome of acutely isolated cortical microglia from adult mice 24 hours after

injection. PCA demonstrated clear separation between control and treated samples,

indicating a robust transcriptional shift (Extended Data Fig. 3A). Differential expression

analysis revealed strong upregulation of interferon-related genes, including *B2m*, *Psmb8*,

*Psmb9*, *Oas1a*, *Irf7*, *Gbp4*, *Rtp4*, *Stat2*, and *Ifit3* (Fig. 3B). Functional analysis confirmed

significant activation of pathways involved in interferon signaling and antiviral

responses, such as "Interferon Alpha Response", "Interferon Gamma Response", and

"Complement" (Fig. 3C, Extended Data Fig. 3B and Supplementary Data Table 4).

We next validated these transcriptional changes at the protein level by examining

Ifitm3 expression in relation to APP/PS1 pathology. Immunohistochemistry revealed

increased hippocampal Ifitm3 labeling in poly(I:C)-treated mice at 24- and 72-hours post-

injection, reaching levels similar to those observed in APP/PS1 mice (Fig. 3D). A similar

increase was noted in the cortex (Extended Data Fig. 3C), suggesting a shared interferon-

related molecular signature between acute poly(I:C) stimulation and chronic APP/PS1

pathology.

Given the marked interferon response, we assessed neuroinflammatory responses

by quantifying GFAP-positive astrocytes. Poly(I:C)-treated mice showed marked

astrogliosis in both the hippocampus (Fig. 3H) and cortex (Extended Data Fig. 2D),

mirroring the elevated GFAP levels observed in APP/PS1 mice. These findings indicate

that poly(I:C) is sufficient to elicit glial reactivity and inflammatory signaling in the brain.

To determine whether this interferon-driven inflammatory response could also

trigger structural and functional changes characteristic of Alzheimer's disease, we next

analyzed key histopathological features. At 72 hours post poly(I:C) treatment, reduced

cFOS immunostaining in the dentate gyrus suggested decreased neuronal activity,

mirroring the hypoactivity observed in APP/PS1 mice (Fig. 3E). We further evaluated

inhibitory circuit integrity by assessing PNNs. Poly(I:C) treatment reduced the proportion

of parvalbumin-positive interneurons surrounded by PNNs, with a comparable decrease

detected in APP/PS1 animals (Fig. 3F). This was accompanied by a global reduction in

WFA-labeled PNN structures (Extended Data Fig. 3E), indicating disruption of

extracellular matrix components critical for circuit stability. Finally, we assessed

neurodegeneration-related pathology by measuring pNF accumulation. Poly(I:C)

administration led to increased pNF immunoreactivity in the hippocampus, reaching

levels comparable to those observed in APP/PS1 mice (Fig. 3G), further supporting a link

between interferon signaling and neuronal injury.

Together, these findings reveal that acute activation of the interferon pathway via

viral-mimetic challenge is sufficient to elicit hallmark Alzheimer-related

histopathological alterations in healthy mice. These results establish a mechanistic link

between interferon responses, neuroinflammation, and neurodegeneration, supporting a

causative role for interferon activation in the pathogenesis of Alzheimer's-like features.

**Genetic Elevation of Interferon Signaling Enhances Alzheimer's-Related**

**Neuropathology**

We recently showed that *Rela* targeting in microglia leads to transcriptional

reprogramming towards an interferon-responsive state (Bhojwani-Cabrera et al.,

submitted manuscript). Taking advantage of this model and to assess the impact of

chronic interferon upregulation on AD progression, we generated a gain-of-function

model by crossing APP/PS1 mice with a tamoxifen-inducible, microglia-specific *Rela*

knockout line (Cx3cr1::CreERT2; Rela$^{fl/fl}$), hereafter referred to as AD hiIFN (Fig. 4A).

Mice were sacrificed at 6 months, allowing us to evaluate the impact of sustained

microglia-driven interferon upregulation during a critical window of Alzheimer's

pathology progression.

Efficient deletion of *Rela* in microglia was confirmed by qPCR performed on

fluorescence-activated cell sorted (FACS) microglia, showing significant reduction in

exons 1–3 and 6–7 expression, with a more pronounced decrease in exons 6–7 (Fig. 4B).

This genetic manipulation resulted in increased expression of interferon-related genes,

including *Irf7*, and *Ifitm3* (Fig. 4C).

Bulk RNA sequencing of hippocampal tissue revealed robust upregulation of

interferon-associated genes such as *B2m*, *Psmb9*, *Mx1, Irf7*, *Rtp4*, *Oas1a*, and *Ifit3* in AD

hiIFN mice compared to AD controls (Fig. 4D). PCA confirmed clear transcriptional

separation between genotypes, supporting a distinct molecular signature induced by

chronic interferon activation (Extended Data Fig. 4A). Gene ontology enrichment

analysis further highlighted significant overrepresentation of interferon and viral defense

pathways, including "Interferon Alpha Response" and "Interferon Gamma Response"

(Fig. 4E and Supplementary Data Table 5). Gene set enrichment analysis using the

"hallmark" gene set collection (56) confirmed enrichment of "Interferon Alpha

Response" pathway in AD hiIFN mice (Extended Data Fig. 4B).

To determine whether this transcriptional program was influenced by sex, we

compared interferon responses across male and female mice in the AD hiIFN model.

Analysis of the top 15 significantly upregulated interferon-stimulated genes AD hiIFN

revealed that both sexes reached comparable levels of interferon-related transcripts (Extended Data Fig. 4C). Based on this observation, both male and female animals were included in the analysis. Protein-level validation by Ifitm3 immunohistochemistry confirmed elevated interferon activity in AD hiIFN mice, with increased labeling of Ifitm3 in the hippocampus (Fig. 4F) and cortex (Extended Data Fig. 4D).

Having confirmed sustained interferon activation at both the transcriptomic and protein levels we next assessed the consequences on key histopathological features. GFAP immunostaining revealed markedly exacerbated astrogliosis in AD hiIFN mice, with higher GFAP labeling in hippocampus (Fig. 4G) and, nearly to significance, in cortex (Extended Data Fig. 4E). In the dentate gyrus, AD hiIFN mice showed a more pronounced reduction of c-FOS expression than AD mice, suggesting further impairment of neuronal activity (Fig. 4H). Structural alterations in inhibitory interneuron networks were also apparent, with a decreased proportion of parvalbumin-positive interneurons enwrapped by perineuronal nets in CA1 (Fig. 4I) and a global reduction in WFA-labeled structures that approached significance (Extended Data Fig. 4F). Finally, axonal integrity was impaired, shown by increased phosphorylated neurofilament staining in hippocampus (Fig. 4J).

Interestingly, despite these exacerbated signs of neurodegeneration and gliosis, amyloid plaque burden was reduced in AD hiIFN mice. Quantification with 6E10 (Fig. 4K), Thioflavin T (Fig. 4L), and OC antibodies (Fig. 4M) demonstrated lower plaque density in both cortex and hippocampus, with largely comparable plaque size distribution (Extended Data Fig. 4G-I).

These data identify IFN-I signaling as a disease-modifying mechanism in AD. Chronic upregulation of the IFN-I axis exacerbates neuroinflammation and

neurodegeneration, uncoupling plaque load from neuronal damage and underscoring a

pathogenic role for interferon-driven processes in disease progression. These findings

highlight IFN-I signaling as a potential therapeutic target in AD.

**Reduction of Interferon Signaling Via STING Antagonism Ameliorates**

**Alzheimer's-Associated Histopathology**

Having demonstrated that elevated IFN-I signaling worsens AD pathology, we next

explored whether attenuating this pathway could yield neuroprotective effects. To address

this, APP/PS1 mice received intraperitoneal injections of C-176, a selective STING

antagonist that prevents Cys91 palmitoylation and thereby blocks downstream interferon

production (69), in alternate days from 3 to 6 months of age, matching the treatment

window used in the IFN-I gain-of-function model (Fig. 5A). These animals are hereafter

termed AD loIFN.

To validate the immunomodulatory effect of C-176, we first treated BV2

microglial cultures with poly(I:C) in the presence or absence of the compound. C-176

reduced expression of *Irf7* and *Ifitm3* while increasing levels of the anti-inflammatory

cytokine *Il-10*, supporting its efficacy in suppressing interferon signaling in vitro

(Extended Data Fig. 5A).

We next evaluated the transcriptional effects of targeting IFN-I signaling *in vivo*.

Given the sex differences observed in IFN-I activation across previous experiments, we

first assessed whether the transcriptional effects of C-176 were comparable between male

and female APP/PS1 mice. To this end, we assessed the top 15 significantly

downregulated interferon-stimulated genes in AD loIFN model. This analysis revealed a

robust suppression of interferon-related transcripts exclusively in C-176–treated females,

while gene expression in males remained largely unchanged (Extended Data Fig. 5B).

These findings suggest that the efficacy of interferon pathway inhibition is sex-dependent, likely due to the already low interferon tone observed in untreated AD males. Based on these results, transcriptomic analyses were focused on female mice, in whom C-176 produced a measurable molecular response.

Accordingly, differential gene expression analysis of hippocampal tissue from treated versus untreated female APP/PS1 mice revealed significant downregulation of key interferon-stimulated genes, including *Psmb8*, *Psmb9*, *Irf7*, *Irf9*, *Rtp4*, Oas1a, and *Ifit3* (Fig. 5B). Functional analysis indicated a loss of interferon- and viral defense–associated "hallmark" gene sets in the treated group, including "Interferon Gamma Response" and "Interferon Alpha Response" (Fig. 5C and Supplementary Data Table 6). GSEA further confirmed significant negative enrichment of the "Interferon Alpha Response" pathway elicited by C-176 treatment (Extended Data Fig. 5C). Analysis of male-specific datasets confirmed a minimal transcriptional response to C-176, with no significant changes in interferon-related genes or pathways detected (Extended Data Fig. 5D,E).

Consistent with overall transcriptional changes, protein-level validation by immunohistochemistry showed reduced numbers of Ifitm3-positive cells in the hippocampus (Fig. 5D) and cortex (Extended Data Fig. 5F) of treated mice.

Critically, these transcriptional and protein-level effects coincided with improvements in AD-related histopathological features. GFAP immunostaining revealed a significant reduction in astrogliosis in hippocampus (Fig. 5E) with a similar trend observed in cortex (Extended Data Fig. 5G). Neuronal activity, measured by c-FOS expression in the dentate gyrus, was partially restored following C-176 treatment (Fig. 5F). The integrity of inhibitory circuits was also preserved, as reflected by an increased

proportion of parvalbumin-positive interneurons surrounded by PNNs (Fig. 5G) and a

concomitant rise in WFA-labeled PNN structures (Extended Data Fig. 5H). Axonal

integrity was likewise improved, as evidenced by a decrease in pNF staining in the

hippocampus (Fig. 5H).

Notably, despite these improvements in neuroinflammation and neuronal

preservation, amyloid pathology remained largely unchanged. Immunostaining with

6E10 revealed no significant differences in plaque burden in the hippocampus, with only

a modest, non-significant increase in the cortex (Fig. 5I). Thioflavin T staining confirmed

the absence of changes in plaque load across brain regions, whereas OC immunostaining

showed no significant difference in the hippocampus but a significant increase in plaque

burden in the cortex of treated mice (Fig.s 5J,K). Importantly, across all three markers,

plaque size showed a consistent trend toward reduction in treated animals (Extended Data

Fig. 5I–K), suggesting that IFN-I pathway inhibition may preferentially impact plaque

maturation and compaction rather than initial deposition.

In summary, pharmacological inhibition of IFN-I signaling via STING

antagonism mitigates key AD-related histopathological features, dampening

neuroinflammation and preserving neuronal integrity independently of plaque load and

pointing to IFN-I pathway suppression as a promising therapeutic approach for modifying

disease trajectory.

**Downregulation of Type I Interferon Signaling Preserves Cognitive Function in**

**Female APP/PS1 Mice**

Given our findings that heightened IFN-I signaling worsens Alzheimer pathology, while

its pharmacological inhibition alleviates key histological hallmarks, we next investigated

whether modulation of IFN-I activity also influences cognitive performance (Fig. 6A).

Specifically, we aimed to establish whether IFN-I signaling serves as a functional

mediator linking neuroimmune dysregulation to behavioral impairments in AD.

We first confirmed group-specific IFN-I activity by transcriptomic profiling of

hippocampal tissue. Comparison between female and male APP/PS1 mice revealed

selective upregulation of ISGs in females, including *Irf7*, *Irf9*, *Psbm8*, *Ifit3*, *Mx1, Rtp4*,

and *B2m*. These transcriptional changes were all attenuated by C-176 treatment in females

(Fig. 6B), indicating sex-dependent regulation of interferon signaling and its

pharmacological modulation.

Behavioral testing revealed domain-specific impairments in APP/PS1 mice, with

a clear sex bias emerging across tasks of increasing complexity. In the elevated plus maze

(EPM), no significant differences were observed between APP/PS1 males and their wild-

type counterparts. In contrast, APP/PS1 females spent significantly more time in the open

arms compared to wild-type females (Fig. 6C), consistent with altered anxiety-like

behavior. Similarly, Y-maze testing revealed impaired spontaneous alternation in females

- indicative of working memory deficits - while male APP/PS1 mice again performed

comparably to controls (Fig. 6D). However, in the novel object location test (NOLT),

which places higher demands on spatial memory, both male and female transgenic mice

failed to recognize the novel location (Fig. 6E). These findings parallel the sex-specific

pattern of IFN-I activation observed at the molecular level and support a functional link

between interferon dysregulation and cognitive decline.

To test this hypothesis directly, we evaluated whether suppression of IFN-I

signaling could ameliorate cognitive impairments in female APP/PS1 mice. Consistent

with transcriptomic evidence of ISG downregulation in the hippocampus (Fig. 6B), C-

176 treatment led to behavioral improvements across multiple domains. In the EPM,

treated mice spent significantly less time in the open arms compared to untreated APP/PS1 females, resulting in anxiety-like behavior levels comparable to those observed in control mice (Fig. 6F). In the Y-maze, C-176–treated females showed improved spontaneous alternation, reaching performance levels comparable to wild-type controls and indicative of restored working memory (Fig. 6G). Finally, in the more demanding NOLT, treated mice successfully discriminated the novel location, in contrast to their untreated counterparts (Fig. 6H), suggesting recovery of spatial memory function.

Notably, performance in the Open Field test was unaffected by genotype or treatment, indicating no major alterations in locomotor activity or basal anxiety (Extended Data Fig. 6A-E). Similarly, no significant differences were detected in the T-maze (Extended Data Fig. 6B-F), contextual fear memory (Extended Data Fig. 6C-G) or cued fear memory performance (Extended Data Fig. 6D-H), suggesting that not all cognitive domains are equally influenced by interferon signaling.

Together, these data demonstrate that interferon signaling not only shapes molecular and histopathological features of AD but also contributes to cognitive dysfunction. Importantly, the ability of STING antagonism to reverse behavioral deficits highlights IFN-I modulation as a promising avenue for therapeutic intervention aimed at preserving cognitive function in AD.

## Discussion

Alzheimer's disease (AD) affects women disproportionately, both in prevalence and progression rate (1, 3, 70–72). While demographic factors such as increased lifespan contribute to this disparity, emerging evidence suggests that biological sex fundamentally shapes the neurobiological response to AD pathology (12, 73). In this study, we show that interferon signaling is selectively and robustly upregulated in the brains of female AD patients and that this sex bias in immune activation corresponds with heightened vulnerability to neurodegeneration in animal models. Through targeted manipulation of the interferon axis *in vivo*, we establish a causal relationship between interferon activity and hallmark features of AD pathology, including neuronal dysfunction, gliosis, and cognitive decline.

Our transcriptomic analyses of the parahippocampal gyrus from AD patients revealed significant enrichment of type I interferon response in females, along with increased type II interferon. These pathways, well-characterized mediators of antiviral immunity and inflammation, have been increasingly implicated in neurodegenerative contexts (74–84). This observation builds on prior reports of sex differences in peripheral immune responses (85–90) and extends them to the human brain, suggesting that female-specific amplification of innate immunity may underlie increased AD susceptibility (91–96).

To explore the functional relevance of these observations, we used the APP/PS1 mouse model of amyloidosis. Female transgenic mice exhibited more severe neuropathology than males, including greater amyloid burden, increased glial activation, and enhanced neuronal and axonal pathology. These histopathological findings were mirrored by transcriptomic alterations, with female hippocampi displaying stronger

upregulation of interferon-stimulated genes and antiviral defense pathways. While sex

differences in AD models have been reported before (97–101), our work provides a

mechanistic framework linking female-biased IFN-I activation to downstream

neuropathological consequences.

Importantly, inducing IFN-I signaling in otherwise healthy wild-type mice - via

systemic administration of the viral mimetic poly(I:C) - was sufficient to reproduce

several AD-like features (102–104), including reduced neuronal activity (cFOS) (103,

104), degradation of perineuronal nets (PNNs) (105), astrogliosis (106), and axonal

damage (107). These findings align with prior demonstrations that chronic IFN-I impairs

neuronal function and plasticity; for instance, Baruch et al. showed that IFN-I drives

synaptic loss and cognitive decline in aging mice, while Blank et al. reported suppression

of hippocampal neurogenesis and long-term potentiation by IFN-I (108–112). Our

observation that both cFOS-positive neurons and PNNs decline following IFN-I

activation supports the idea that innate immune overactivation directly disrupts neuronal

circuits (80, 108, 113–115). These results extend the role of IFN-I in the CNS beyond

antiviral defense, positioning it as a key regulator of neural integrity.

The use of poly(I:C) to mimic viral activation also intersects with the long-

standing "infectious hypothesis" of AD, which posits that the disease may be driven, at

least in part, by host immune responses to chronic or latent infections (30, 34, 110).

Several studies have identified herpesvirus DNA in postmortem AD brains (116–119),

and recent transcriptomic analyses have revealed molecular signatures of past viral

exposure, including SARS CoV 2–related signals (120–123). Although causality remains

unresolved, our data show that even sterile mimetics of viral RNA can trigger key aspects

of AD pathology, reinforcing the notion that persistent immune stimulation can act as a primary driver of neurodegeneration.

In this context, it is noteworthy that population-based studies have found associations between vaccination against pathogens such as influenza and herpes zoster, and a reduced incidence of dementia, including AD (124–128). Although mechanisms remain speculative, one hypothesis is that vaccines reduce the frequency or severity of viral reactivation and associated IFN-I bursts (114, 124, 129). Alternatively, they may establish a more regulated immune tone less prone to maladaptive activation (74, 75, 127). Our data provides mechanistic support for these epidemiological associations, suggesting that modulation of IFN-I activity may mediate the protective effects of immunization.

To further assess the consequences of sustained IFN-I activation, we developed a genetic gain-of-function model in which deletion of *Rela* in microglia reprogrammed these cells toward an IFN-I-responsive phenotype (Bhojwani-Cabrera et al., submitted manuscript). In that study, *Rela* knockout mice with heightened IFN-I pathway activation exhibited cognitive impairment, consistent with previous reports linking increased interferon signaling to cognitive decline (79, 83, 109, 112, 130, 131). When crossed with APP/PS1 mice, this manipulation led to exacerbated neuroinflammation, axonal pathology, and reduced neuronal activity. Interestingly, amyloid plaque burden was reduced in these animals, indicating that IFN-I signaling can drive neurotoxicity through mechanisms that are not strictly dependent on plaque accumulation (79, 80, 84). This reflects a broader conceptual shift in the field - from viewing AD primarily as a disorder of protein accumulation to recognizing the role of maladaptive immune responses (132–134). Our findings parallel results from microglia-specific *C9orf72* knockout mice, which

also showed robust IFN-I signature accompanied by synaptic loss and reduced amyloid deposition (135). However, the case of Ifitm3 - a key IFN-I effector - underscores the complexity: although it enhances γ-secretase activity and amyloidogenesis (136), its upregulation in *Rela* cKO brains coincided with fewer plaques, highlighting that its effect may depend on the overall inflammatory context. Together, these observations suggest that the neurotoxic impact of IFN-I responses may outweigh their influence on plaque dynamics.

A key upstream trigger of IFN-I signaling in AD is activation of the cGAS–STING pathway, which senses cytosolic DNA such as mitochondrial DNA released by cellular stress - a process exacerbated near amyloid plaques (137–140). This pathway is aberrantly activated in both AD models and human brain tissue, promoting IFN-I responses and neuroinflammation (113, 141–144). Notably, our transcriptomic analyses revealed increased p53 pathway activity in AD females compared to males, a relevant finding given that p53 can amplify cGAS–STING signaling in response to DNA damage–associated stress (145, 146). Consistent with these findings, we targeted STING using C-176 in APP/PS1 mice. This treatment reduced IFN-I-stimulated gene expression, dampened gliosis, preserved neuronal activity and axonal integrity, and improved cognitive performance - particularly in females with higher baseline IFN-I levels. Notably, these improvements occurred despite minimal changes in plaque burden, reinforcing the idea that cGAS–STING–driven IFN-I responses contribute to neurodegeneration independently of amyloid accumulation.

This dissociation between plaque deposition and clinical outcome mirrors observations in "resilient" individuals - older adults with high amyloid load who remain cognitively intact (147, 148). Postmortem analyses of such individuals often reveal low glial activation and minimal inflammation (149, 150). Similarly, in C-176 treated mice,

IFN-I signaling was suppressed and cognitive function preserved despite persistent

plaques. These data strengthen the argument that the immune system - especially the IFN-

I axis - is a key determinant of whether amyloidosis remains subclinical or progresses to

overt neurodegeneration.

Although these results nominate the IFN-I axis as a compelling therapeutic target,

translation to the clinic demands scrutiny. C-176 has demonstrated efficacy in modulating

IFN-I signaling *in vivo* in murine models, yet its activity is confined to the murine STING

isoform, with no evidence supporting efficacy against the human variant (69, 151). By

contrast, compounds such as H-151 exhibit greater selectivity toward human STING and

represent more promising candidates for clinical development (69). Importantly,

preclinical data in murine models support the blood–brain barrier permeability and CNS

penetrance of these inhibitors, providing a promising pharmacokinetic basis for further

investigation (113, 152–154). Nonetheless, their pharmacokinetic properties in humans

remain to be characterized. Although IFN-I plays an essential role in host defense and

tumor surveillance, and systemic inhibition carries potential risks (115, 129, 155, 156),

several IFN-I-targeting agents - such as JAK inhibitors and anti IFNAR antibodies - are

already in clinical use for autoimmune disorders, providing a strong foundation for

repurposing (157–159). Patient stratification based on peripheral or cerebrospinal

biomarkers of IFN-I activity could further refine candidate selection and enhance

therapeutic precision.

These considerations underscore the need for precision medicine approaches that

account for interindividual variability in immune tone and response. Notably, our findings

emphasize the importance of sex as a biological variable in both disease progression and

therapeutic efficacy. Historically, females have been underrepresented in neuroscience

research, and many animal models fail to account for sex-based differences (11, 21, 22,

24). Here, the inclusion of both sexes revealed critical differences in pathology and a

female-specific therapeutic response to C-176 that would have been missed in a unisex

design. Moreover, the observation that epithelial–mesenchymal transition emerged as the

dominant pathway in males, whereas interferon alpha response predominated in females,

highlights fundamentally different disease-associated programs that may each warrant

tailored intervention. Future studies should address the potential pathogenic role of

epithelial–mesenchymal transition in male AD, as understanding these sex-specific

signatures will be essential for developing stratified therapeutic strategies in which sex

and immune phenotype guide clinical decision-making.

In conclusion, our study reveals that IFN-I signaling is selectively elevated in the

female AD brain and that this pathway actively contributes to neuronal and cognitive

dysfunction, independently of amyloid burden. By integrating human transcriptomics

with mechanistic animal models, we identify IFN-I as a central, sex-biased effector of

AD pathogenesis. These findings suggest that modulating IFN-I responses may offer

therapeutic benefit, particularly in patient populations characterized by heightened innate

immune activation. Future research should aim to identify biomarkers of IFN-I tone,

evaluate the safety of long-term IFN-I modulation, and determine whether early

intervention in IFN-I-dominant disease trajectories can prevent the transition from

asymptomatic amyloidosis to neurodegenerative decline.

## Methods

**Sex as a biological variable**

Both male and female mice were included in all experimental groups unless otherwise stated, and sex was a central consideration in the study design. Experimental cohorts were balanced by sex, and all transcriptomic, histological, and behavioral analyses were stratified accordingly. This allowed for direct comparison of AD–related phenotypes and immune signaling between male and female animals. Similarly, human transcriptomic datasets from the The Mount Sinai Brain Bank (MSBB) study included 94 AD patients (72 females and 32 males) and 34 age-matched healthy controls (17 females and 17 males), and differential expression analyses were performed to identify sex-specific molecular signatures. Collectively, these approaches ensured that sex was rigorously integrated as a biological variable throughout the study.

**Mice**

APP/PS1 (64) and Cx3cr1$^{CreER}$ (160) mice are available at public repositories (RRID:MMRRC_034832-JAX and RRID:IMSR_JAX:021160, respectively). RelA$^{fl/fl}$ mice which harbor loxP sites flanking exons 5-8 of the *Rela* gene, have been previously described (MGI:3775205) (161) and were kindly provided by Dr. Angel Barco (Instituto de Neurociencias). Cx3cr1$^{CreER}$ mice were crossed with RelA$^{fl/fl}$ mice (161) resulting in the line referred as Cx3cr1$^{CreER}$-Rela$^{fl/fl}$. These mice were crossed with APP/PS1 mice to generate Cx3cr1$^{CreER}$-Rela$^{fl/fl}$-APP/PS1. In the experiments involving tamoxifen-induced knockouts, we used tamoxifen-treated Rela$^{fl/fl}$-APP/PS1-littermates as controls. All transgenic lines were maintained on a C57BL/6J genetic background.

**BV2 microglial cell line**

Mouse microglial BV2 cells were cultured in high-glucose Dulbecco's modified Eagle's

medium (DMEM) (Thermo Fisher Scientific, 41965047) supplemented with 10% heat-

inactivated fetal bovine serum (FBS) (Thermo Fisher Scientific, 10270106) and 1%

penicillin-streptomycin (Sigma-Aldrich, P4333). Cells were maintained at 37°C in a

humidified atmosphere with 5% $CO_2$ and plated at a density of $3 \times 10^5$ cells per well in

6-well plates for stimulation assays.

***In vivo* animal treatments**

According to each experiment, animals were treated following ethical guides as described

previously. To induce Cre recombination in Cx3cr1$^{CreER}$ mice, tamoxifen (TMX) (Sigma-

Aldrich, T5648) was dissolved at 25 mg/mL in corn oil (Sigma-Aldrich, C8267) by

heating to 50°C for 1 hour under light-protected conditions. Three-month-old mice

received two intraperitoneal injections of tamoxifen (75 mg/kg) on alternate days. All

downstream experiments involving TMX were conducted at least one month after TMX

administration. Polyinosinic:polycytidylic acid high molecular weight (poly(I:C))

(Sigma-Aldrich, P1530-25MG) was prepared in sterile, endotoxin-free water and

administered intraperitoneally at a dose of 12 mg/kg. Mice were sacrificed at 24- or 72-

hours post-injection for transcriptomic and histological analyses. The STING inhibitor C-

176 (MedChemExpress, HY-112906) was prepared at 1.5 mg/mL in a corn oil:ethanol

(90:10) solution. Mice received intraperitoneal injections of 10 mg/kg on alternate days

from 3 to 6 months of age, corresponding to the window of APP/PS1 pathology

development.

**Immunohistochemistry and image analysis**

Mice were deeply anesthetized and transcardially perfused with PBS, followed by 4%

paraformaldehyde (PFA) (PFA, w/v, dissolved in 0.1 M phosphate buffer, pH 7.4). Brains

were post-fixed overnight and cryoprotected in 30% (w/v) sucrose for 48h. Coronal

sections (50 μm) were obtained using a cryotome and stored in antifreeze solution (30%

ethylene glycol, 30% glycerol, 30% distilled water, 10% PBS) at −20°C until processing.

Briefly, free-floating sections were washed 3 times in PBS (5 min each) and blocked in

5% Newborn calf serum (NCS) (Sigma-Aldrich, N4762) with 0.3% Triton X-100 (Sigma-

Aldrich, T8787), followed by overnight incubation at 4°C with primary antibodies. The

following primary antibodies were used: Ifitm3 (Proteintech, 11714-1-AP, 1:500), 6E10

(anti-β-Amyloid, 1–16) (BioLegend, 803001, 1:1000), c-FOS (Synaptic Systems,

226308, 1:1000), phosphorylated neurofilament (pNF) (Biolegend, SMI 31P, 1:500),

Parvalbumin (Synaptic Systems, 195004 1:1000), GFAP (Sigma-Aldrich, G3893, 1:500),

and OC (Sigma-Aldrich, AB2286, 1:5000). On the following day, slides were rinsed 3

times in PBS, incubated with fluorophore-conjugated secondary antibodies, and rinsed

again 3 times in PBS. The following secondary antibodies were used: Donkey anti-rat

IgG H+L Alexa Fluor 488 (Abcam, ab150153), Donkey anti-rabbit IgG H+L Alexa Fluor

647 (Abcam, ab150067), Donkey anti-mouse IgG H+L Alexa Fluor Plus 594 (Thermo

Fisher Scientific, A32744) and Donkey anti-guinea pig IgG H+L Alexa Fluor 647

(Abcam, 706-605-148), all at 1:500 dilution. Finally, tissue sections were counterstained

with DAPI (Thermo Fisher Scientific, A32733) and mounted onto glass slides using

Fluoromount- G (Sigma-Aldrich, T5941). For thioflavin T staining, after secondary

antibody staining, sections were incubated with 0.01% Thioflavin T (Sigma-Aldrich,

T3516) in water for 5 minutes, followed by 50% ethanol and PBS washes. For the

immunodetection of perineuronal nets (PNNs), after secondary antibody staining, slices

were incubated overnight at 4 °C with a solution containing biotinylated *Wisteria*

*floribunda* Lectin (WFA) (Vector Laboratories, B-1355-2, 1:300). On the following day,

sections were rinsed 3 times in PBS (5 min each) at RT, incubated with a solution of red

fluorescent streptavidin (Alexa Fluor™ 555 conjugate) (Thermo Fisher Scientifics,

S32356, 1:300) and 5% NCS in PBS for 1h at RT, and rinsed again 3 times in PBS.

Images were acquired using a Zeiss Axio Scan.Z1 slide scanner (20× objective).

Quantification was performed using Arivis Vision4D software, employing standardized

macros for image thresholding and segmentation. Regions of interest (ROIs) were

manually drawn around the dentate gyrus, CA1, cortex, or the entire hippocampus.

Quantification metrics included: total number of cFOS$^+$ nuclei in the dentate gyrus; total

number of WFA$^+$ objects and the percentage of PV$^+$ interneurons surrounded by PNNs in

the CA1 region; number and total area of GFAP$^+$ astrocytic objects; total pNF$^+$ area in

CA1; total number and area of Ifitm3$^+$ cells in cortex and hippocampus; amyloid plaque

density, number, and average size of Thioflavin T$^+$, 6E10$^+$, and OC$^+$ amyloid deposits.

**FACS sorting of microglia**

Brains were collected and placed on ice in sterile phosphate-buffered saline (PBS). After

removal of the meninges, cortical hemispheres were mechanically dissociated in Dounce

buffer (15 mM HEPES, 0.5% glucose, HBSS 1x) using a tissue homogenizer (Fisher

Scientific, 10198611). Myelin was removed by centrifugation through a 25% isotonic

Percoll gradient at 800 x g for 15 minutes at 4°C. The resulting cell pellet was resuspended

in PBS, centrifuged, and processed for flow cytometry to isolate microglia as previously

described (Bhojwani-Cabrera et al., submitted manuscript). Briefly, Prior to antibody

staining, samples were blocked with Fc-block CD16/CD32 (Biolegend, 101320, 1:50) in

FACS Buffer sterile filtered (1% FCS, 2mM EDTA, 25mM HEPES in PBS) for 10 min

on ice and protected from light to block nonspecific binding. For cell surface staining,

cells were resuspended in the appropriate antibody cocktail and incubated for 30 min on

ice protected from light. Primary antibodies used were anti-mouse GFP (Aveslab, GFP-

1020, 1:500), anti-mouse Cd11b (BioLegend, 101235, 1:500), anti-mouse CD45

(BioLegend, 103106, 1:500). Samples were centrifuged and washed with FACS buffer.

Viability was assessed by staining with DAPI (1 μg/mL). Cells were sorted using a BD

FACS Aria III cell sorter (BD Bioscience). To sort the cells, a 85-micron nozzle with 4-

Way purity mode was used. The gating strategy involved selecting the cell population

based on forward scatter area (FSC-A) to discriminate cells by size, and side scatter area

(SSC-A) to separate them by granularity or internal complexity. Doublets were then

excluded by gating FSC-height (FSC-H) and FSC-A to isolate single cells. All final

analysis and data output were performed using FlowJo software (BD). For RNA-seq

experiments, microglia were isolated from the whole adult mouse cortex, yielding

200,000-300,000 cells. Sorted cells were collected directly into RLT buffer (Quiagen,

79216) for subsequent RNA extraction.

**Real-time quantitative PCR (RT-qPCR)**

To analyze gene expression in acutely FACS-isolated microglia from the adult mouse

hippocampus, RNA was extracted using the RNeasy Mini Kit (Qiagen, 74104) with on-

column DNase digestion (Thermo Fisher Scientific, 18047019). For BV2 cells, RNA was

isolated using TRIzol reagent (Thermo Fisher Scientific, A33251). For all samples, RNA

purity, integrity, and concentration were assessed by spectrophotometry (Nanodrop ND-

1000 spectrophotometer) and micro-capillary electrophoresis (2100 Bioanalyzer,

Agilent). Complementary DNA was synthesized using the RevertAid First Strand cDNA

Synthesis Kit (Thermo Fisher Scientific, EP0442) following the manufacturer's protocol,

and quantitative PCR was performed using EvaGreen Master Mix (Cultek, 755032) on a

QuantStudio 3 system (Applied Biosystems). Relative expression levels were analyzed

using the ΔΔCt method ((162), with Gapdh as the housekeeping gene for normalization.

Primer sequences used in RT–qPCR assays are listed in Supplementary Data Table 7.

**Human transcriptomic samples (MSBB Cohort)**

RNA-seq data from the parahippocampal gyrus (Brodmann area 36) were obtained from

the Mount Sinai Brain Bank (MSBB) via the AMP-AD Knowledge Portal (Synapse ID:

syn3159438; metadata: syn29855570; count data: syn27068754). We used processed raw

count matrices in which genes with fewer than 1 count per million (CPM) had already

been filtered out. For our analyses, we further subset the dataset to include only PHG

samples of AD cases aged 69 to 89 years at death. Resequenced samples were handled

by excluding unique sequencing batches; when biological replicates from the same

individual were available, those with higher RNA integrity (RIN and $RIN^2$ scores) were

preferentially retained. To reduce sex-related confounding effects, samples were

additionally filtered based on the expression of sex-linked genes, resulting in the removal

of one sample (see Extended Data Fig. 1). The final dataset included 33 female and 28

male AD cases. For additional analyses, samples were filtered using the same criteria,

selecting either control samples aged 69 to 89 years at death (17 females and 17 males)

or AD cases older than 69 years (72 females and 32 males).

**Mouse transcriptomic profiling by RNA-seq**

For the Alzheimer's disease-related samples (GSE304521), total RNA was extracted

from hippocampi, rapidly dissected on ice using fine forceps. For the poly(I:C)-related

samples (GSE304362), total RNA was extracted from acutely FACS-isolated microglia

from the adult brain. For all samples, total RNA was extracted using the RNeasy Mini Kit

(Qiagen, 74104), including on-column DNase digestion (Thermo Fisher Scientific,

18047019). RNA quality was assessed by micro-capillary electrophoresis (Agilent

TapeStation, Agilent), and only samples with a RNA Integrity Number (RIN) $\geq$ 7 were

used for library construction. Libraries were amplified by PCR, purified, and size-selected

to enrich for fragments compatible with Illumina sequencing, followed by quality

assessment using fluorometry (Qubit fluorometer, Thermo Fisher Scientific) and micro-

capillary electrophoresis (Agilent TapeStation, Agilent). Libraries were prepared by

Novogene Europe (Cambridge, UK) using a strand-specific protocol with poly-A

selection and sequenced on an Illumina NovaSeq X Plus platform (PE150). For the

Alzheimer's disease-related samples (GSE304521), RNA-seq libraries were constructed

using Novogene NGS Stranded RNA Library Prep Set (PT044). Raw paired-end RNA-

seq reads in FASTQ format were processed by Novogene Europe (Cambridge, UK) using

their standard bioinformatics pipeline. Briefly, adapter sequences, reads containing poly-

N, and low-quality bases were removed using fastp (163). Quality metrics such as Q20,

Q30, and GC content were also calculated during this step. All downstream analyses were

based on the resulting high-quality clean reads. Clean reads were aligned to the mm39

mouse reference genome using HISAT2 (v2.2.1; (164)). The reference genome index was

built using HISAT2, and gene model annotations were used to enable splice-aware

alignment with improved accuracy. Library sizes of primary mapped reads were

consistently above 45 million fragments. Gene-level quantification was performed using

featureCounts (v2.0.6; (165)). The resulting count tables were used for downstream

differential expression analysis. For the poly(I:C)-related samples (GSE304362), RNA-

seq libraries were constructed using Novogene NGS RNA Library Prep Set (PT042). Raw

paired-end data were processed in-house following a custom analysis pipeline. Briefly,

raw paired-end RNA-seq reads in FASTQ format were first subjected to quality control

using FastQC (v0.11.9; https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to

assess base quality and sequence content. Adapter sequences and low-quality bases were

then removed using Trim Galore (v0.6.7; https://zenodo.org/records/7598955) in paired-

end mode. Post-trimming quality was re-evaluated with FastQC to confirm adapter

removal and improved read quality. High-quality reads were aligned to the GRCm38

mouse reference genome

using HISAT2 (v2.2.1; (166))  with the --dta option to facilitate downstream transcript

assembly. Library sizes of primary mapped reads were consistently above 50 million

fragments. Alignment output in SAM format was converted and sorted into BAM format

using SAMtools (v1.13; (167)), and subsequently indexed. For genome browser

visualization, BAM files were converted to TDF format using IGVTools (v2.5.3), aligned

to the mm10 genome build, and visualized in IGV (168). Gene-level quantification was

performed using HTSeq-count (v0.13.5) with parameters --stranded=no, --type=exon,

and --order=pos. Gene annotations were obtained from Ensembl release 102

(Mus_musculus.GRCm38.102.chr.gtf). The resulting count tables were used for

downstream differential expression analysis.

**Differential gene expression analysis**

Differential expression analyses were conducted in R version 4.5.0 (R Core Team, 2025)

using DESeq2 (v1.48.1; (169)), considering genes with adjusted P value < 0.05 as

significantly differentially expressed. Batch correction was applied when needed using

the ComBat_seq() function from the sva package (v3.56.0, (170)). To facilitate the

detection of batch effects, outliers, and major sources of variance, Principal Component

Analyses were performed on variance-stabilized data (VST for datasets with >30

samples) or regularized log-transformed data (rlog for datasets with <30 samples) using

the plotPCA() function from the DESeq2 package. The top 500 most variable genes were

used for dimensionality reduction. The first two principal components were visualized

with ggplot2 (v3.5.2, (171)), and sample labels were added using geom_text_repel() from

ggrepel (v0.9.6, (172))to improve readability. Heatmaps were generated on gene

expression counts rlog-transformed using pheatmap (v1.0.13, (173)), with hierarchical

clustering applied to genes and samples ordered by experimental design. Rows were

scaled using z-scores, and styling was harmonized with the overall plot design. Volcano

plots were generated using ggplot2, where the $-\log_{10}$ adjusted p-value was plotted against

the $\log_2$ fold change. Genes were color-coded by significance (padj < 0.05 or < 0.1) and

classified as upregulated, downregulated, or not significant. Key interferon-alpha–related

genes were labeled using ggrepel. Optional axis cutoffs were applied to reduce the impact

of extreme outliers on plot scaling.

**Functional enrichment analysis**

Functional enrichment analyses were performed on significantly differentially expressed

genes as input (padj < 0.05) using Enrichr (174) and the MSigDB "hallmark" 2020 gene

set collection (56). Gene Set Enrichment Analysis (GSEA) was performed to identify

coordinated shifts in predefined gene sets, using the GSEA_4.4.0 desktop application

(57), with the following parameters: 1,000 phenotype permutations, No_Collapse, and a

fixed permutation seed (149). Ranked gene lists (.rnk files) were created from DESeq2

output using tidyverse tools (v2.0.0, (175))without additional filtering. Enrichment was

tested against the MSigDB "hallmark" gene sets for both human (v2025.1.Hs) and mouse

(v2025.1.Mm) (56) GSEA plots were recreated in R using ggplot2, cowplot (v1.2.0,

(176)), and gridExtra (v2.3, (177)) to maintain stylistic consistency across visual outputs.

## Mouse behavioral testing

All behavioral assays were conducted during the light phase under consistent illumination (~20 lux) in a dedicated testing suite. Mice were acclimated to the testing room for at least 30 minutes before each test. Behavior was recorded and analyzed using SMART 3.0 software (Panlab). We used adult mutant and control littermates of both sexes. For the open field (OF) test, mice were placed in the center of a 40 × 40 × 40 cm arena and allowed to explore freely for 15 minutes. Total distance traveled and time spent in the center zone were recorded as measures of locomotor activity and anxiety-like behavior, respectively. The *elevated plus maze* (EPM) consisted of two open and two closed arms (30 × 5 cm), elevated 50 cm above the floor. Mice were placed in the center facing an open arm and allowed to explore for 5 minutes. The percentage of time spent and number of entries into open arms were calculated. To evaluate spatial working memory spontaneous alternation was measured using the *Y-maze*. Briefly, the transparent symmetrical Y-maze consisted of three arms (30 × 5 × 15 cm). Three objects of similar size (~20 cm3) were situated surrounding the maze at ∼15–20 cm outside the walls and mice were allowed to freely explore the maze for 8 minutes. An alternation was defined as sequential entry into all three arms (triad). Percent alternation = (Number of alternations / [Total arm entries − 2]) × 100. To study short-term spatial memory, we used the *T-maze*. The apparatus consisted of three arms, two of them situated at 180° from each other and the third situated at 90° with respect to the other ones representing the stem arm of the T. All three arms were 45 cm long. In the training trial, one arm was closed (novel arm) and mice were placed in the stem arm of the T (home arm) and allowed to explore this arm and the other available arm (familiar arm) for 10 min, after which they were returned to the home cage. After an inter-trial interval of 1 h mice were placed in the stem arm of the T-maze and allowed to freely explore all three arms for 5 min (testing phase).

The time in the new arm∗100/total time in the two arms at 180° was the parameter

evaluated. In the *novel object location* (NOLT) memory task, mice were habituated to a

40 × 40 cm arena for 15 minutes. Different distal visual cues were allocated in the walls

of the arena. On day 2, two identical objects were presented during a 10-minute

familiarization phase. After a 24-hour delay, one object was relocated to a novel position

(test phase), and mice were allowed to explore for 5 minutes. The discrimination index

was calculated as: (Time_novel − Time_familiar) / Total_exploration_time. Associative

memory was assessed using *fear conditioning paradigm*. A 25 x 25 x 25 cm black

chamber system was used (Panlab, Spain).  On Day 1, animals were allowed to habituate

for 5 minutes. After habituation, a tone (90 dB) was presented for 30 seconds. During the

final 2 seconds of the tone, a 0.4 mA foot shock was administered for 1–2 seconds and

the mouse was allowed to remain 30 additional seconds in the chamber. This procedure

was repeated two times, and the mouse remained 2 additional minutes in the chamber

before removing it. On Day 2, the contextual memory test was conducted. The mouse

was placed in the chamber for 5 minutes, with no tone presented, and freezing behaviour

was recorded. Cued memory was assessed on day 3 in a novel context, consisting of a 5-

minute baseline followed by a 30-seconds tone presentation. Freezing was defined as the

absence of all movement except respiration for >1 second. Animals were tracked and

recorded with the Packwin V2.0.08 software (Panlab, Spain).

**Statistics**

Data are presented as mean ± standard error of the mean (SEM), unless otherwise

indicated. Statistical analyses were performed using GraphPad Prism (version 7) or R

(version 4.2 or later). For comparisons between two groups, unpaired two-tailed Student's

t tests were used. Comparisons involving more than two groups were analyzed using one-

way ANOVA followed by Bonferroni post hoc correction. For non-normally distributed

data, the Mann–Whitney U test was applied. Correlations were assessed using Pearson's

correlation coefficient. A *P* value less than 0.05 was considered statistically significant.

Specific statistical tests and sample sizes (n) for each experiment are reported in the figure

legends.

**Study approval**

The study was conducted in compliance with European Union Directive 2010/63/EU on

the protection of animals used for scientific purposes. All the protocols for animal

experimentation were approved by the Animal Welfare Committee at the Instituto de

Neurociencias, the CSIC Ethical Committee, and the Dirección General de Agricultura,

Ganadería y Pesca of Generalitat Valenciana. Mice were maintained under specific

pathogen-free (SPF) conditions at the Animal Facility of the Instituto de Neurociencias

(CSIC-UMH), on a 12 h light/12 h dark cycle (7:00 a.m. to 7:00 p.m.), at 20–24 °C and

40–60% controlled humidity, with *ad libitum* access to food and water.

**Data availability**

The transcriptomics data sets generated in this study can be accessed at the GEO public

repository using the accession number GSE304521 and GSE304362. In addition, human

RNA-seq data used in this study were obtained from the publicly available The Mount

Sinai Brain Bank (MSBB) study, hosted on the AD Knowledge Portal

(https://adknowledgeportal.org/). The dataset is accessible via Synapse

(https://www.synapse.org/): syn3159438; metadata: syn29855570; count data:

syn27068754). The results published here are in whole or in part based on data obtained

from the AD Knowledge Portal (https://adknowledgeportal.org/). These data were

generated from postmortem brain tissue collected through the Mount Sinai VA Medical

Center Brain Bank and were provided by Dr. Eric Schadt from Mount Sinai School of

Medicine. Processed data and code used for statistical analysis, gene expression

processing, and figure generation are available upon request from the corresponding

author.

## Author contributions

Conceptualization, V.L-L., and J.P.L-A.; methodology, V.L-L., M.G-G., G.I, A.E-C., A.B-C., A.G., and J.P.L-A.; investigation, V.L-L., M.G-G., and A.E-C.; data curation, V.L-L., M.G-G., G.I, and M.G-F.; formal analysis: V.L-L., M.G-G., and G.I.; writing – original draft, V.L-L., and J.L-A.; writing – review & editing, V.L-L., G.I., M.G-G., A.E-C., A.B-C., M.G-F., A.G., and J.L-A.; funding acquisition, J.P.L-A.; resources, J.P.L-A.; supervision, J.P.L-A. The order of authorship was determined by overall contributions and approved by the authors for the final manuscript.

## Acknowledgments

We thank Eduardo Soriano, Alberto Pascual, Jose V. Sánchez-Mut, and Berta L. Sánchez-Laorden for the critical reading of the manuscript. We are also grateful to members of the López-Atalaya laboratory for stimulating discussions throughout the course of this work. We thank the personnel of the Mouse Facility and the Microscopy and Omics Core Services at the Instituto de Neurociencias for their support and technical assistance. This study was supported by grant PID2021-129053OB-I00 funded by MCIU/AEI/10.13039/501100011033 and by ERDF, EU, and the Generalitat Valenciana, Conselleria d'Educació, Universitats i Ocupació (PROMETEO 2020/007) to J.P.L-A.; V.L-L. is the recipient of a 2021 neuroscience doctoral fellowship from the Fundación Tatiana Pérez de Guzmán el Bueno. G.I. holds a Formación de Personal Investigador (FPI) contract (CEX2021-001165-S-20-2), associated with "Severo Ochoa" grant CEX2021-001165-S from the MCIU/AEI/10.13039/501100011033. A.B-C. held a Formación de Personal Investigador (FPI) contract (PRE2019-090492), associated with grant RTI2018-102260-B-I00 awarded to J.P.L-A., and funded by MCIU/AEI/10.13039/501100011033 and by "ERDF A way of making Europe". M.G-F.

is the recipient of a Formación de Profesorado Universitario contract (FPU22/01375)

funded by the Spanish Ministry of Science, Innovation and Universities (MICIU). The

Institute of Neurosciences is a Severo Ochoa Center of Excellence (grant CEX2021-

001165-S, funded by MCIU/AEI/10.13039/501100011033). This work was also funded

by European Union NextGenerationEU/PRTR CNS2022 – 135391 ("Consolidación"

program) and the Fundación Ramón Areces (CIVP21A7024) both to A.G. The project

that gave rise to these results received the support of a fellowship from "la Caixa"

Foundation (ID 100010434). The fellowship code is LCF/BQ/DR21/11880016 to M.G.-

G.

# References

1. 2025 Alzheimer's disease facts and figures. *Alzheimer's & Dementia*. 2025;21(4). https://doi.org/10.1002/alz.70235.

2. Scheltens P, et al. Alzheimer's disease. *The Lancet*. 2021;397(10284):1577–1590.

3. Aguzzoli CS, et al. *World Alzheimer Report 2024 Global changes in attitudes to dementia Contributors: Survey translators*.

4. 2024 Alzheimer's disease facts and figures. *Alzheimer's and Dementia*. 2024;20(5):3708–3821.

5. Fang M, et al. Lifetime risk and projected burden of dementia. *Nat Med*. 2025;31(3):772–776.

6. Liu E, Zhang Y, Wang JZ. Updates in Alzheimer's disease: from basic research to diagnosis and therapies. *Transl Neurodegener*. 2024;13(1). https://doi.org/10.1186/s40035-024-00432-x.

7. Cummings JL, Osse AML, Kinney JW. Alzheimer's Disease: Novel Targets and Investigational Drugs for Disease Modification. *Drugs*. 2023;83(15):1387–1408.

8. Cummings J, et al. Alzheimer's disease drug development pipeline: 2023. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2023;9(2). https://doi.org/10.1002/trc2.12385.

9. Cui SS, Jiang QW, Chen S Di. Sex difference in biological change and mechanism of Alzheimer's disease: From macro- to micro-landscape. *Ageing Res Rev*. 2023;87. https://doi.org/10.1016/j.arr.2023.101918.

10. Aggarwal NT, Mielke MM. Sex Differences in Alzheimer's Disease. *Neurol Clin*. 2023;41(2):343–358.

11. Nebel RA, et al. Understanding the impact of sex and gender in Alzheimer's disease: A call to action. *Alzheimer's and Dementia*. 2018;14(9):1171–1183.

12. Lopez-Lee C, et al. Mechanisms of sex differences in Alzheimer's disease. *Neuron*. 2024;112(8):1208–1221.

13. Buckley RF, et al. Sex Differences in the Association of Global Amyloid and Regional Tau Deposition Measured by Positron Emission Tomography in Clinically Normal Older Adults. *JAMA Neurol*. 2019;76(5):542–551.

14. Rosenzweig N, et al. Sex-dependent APOE4 neutrophil–microglia interactions drive cognitive impairment in Alzheimer's disease. *Nat Med*. 2024;30(10):2990–3003.

15. Eikelboom WS, et al. Sex differences in neuropsychiatric symptoms in Alzheimer's disease dementia: a meta-analysis. *Alzheimers Res Ther*. 2022;14(1). https://doi.org/10.1186/s13195-022-00991-z.

16. Sohn D, et al. Sex Differences in Cognitive Decline in Subjects with High Likelihood of Mild Cognitive Impairment due to Alzheimer's disease. *Sci Rep*. 2018;8(1). https://doi.org/10.1038/s41598-018-25377-w.

17. Lin KA, et al. Marked gender differences in progression of mild cognitive impairment over 8 years. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2015;1(2):103–110.

18. Terstege DJ, Epp JR. Parvalbumin as a sex-specific target in Alzheimer's disease research – A mini-review. *Neurosci Biobehav Rev*. 2023;153. https://doi.org/10.1016/j.neubiorev.2023.105370.

19. Karp NA, Reavey N. Sex bias in preclinical research and an exploration of how to change the status quo. *Br J Pharmacol*. 2019;176(21):4107–4118.

20. Buckley RF, Gong J, Woodward M. A Call to Action to Address Sex Differences in
Alzheimer Disease Clinical Trials. *JAMA Neurol*. 2023;80(8):769–770.
21. Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. *Neurosci*
*Biobehav Rev*. 2011;35(3):565–572.
22. Shansky RM. Are hormones a "female problem" for animal research? *Science*
*(1979)*. 2019;364(6443):825–826.
23. Improving Bioscience Research Reporting: The ARRIVE Guidelines for Reporting
Animal Research. *Vet Clin Pathol*. 2012;41(1):27–31.
24. Clayton JA, Collins FS. Policy: NIH to balance sex in cell and animal studies.
*Nature*. 2014;509(7500):282–283.
25. Bale TL, Epperson CN. Sex as a Biological Variable: Who, What, When, Why, and
How. *Neuropsychopharmacology*. 2017;42(2):386–396.
26. Bracher-Smith M, et al. Machine learning in Alzheimer's disease genetics. *Nat*
*Commun*. 2025;16(1):6726.
27. Jansen IE, et al. Genome-wide meta-analysis identifies new loci and functional
pathways influencing Alzheimer's disease risk. *Nat Genet*. 2019;51(3):404–413.
28. Bellenguez C, et al. New insights into the genetic etiology of Alzheimer's disease
and related dementias. *Nat Genet*. 2022;54(4):412–436.
29. Shade LMP, et al. GWAS of multiple neuropathology endophenotypes identifies
new risk loci and provides insights into the genetic risk of dementia. *Nat Genet*.
2024;56(11):2407–2421.
30. Brown GC, Heneka MT. The endotoxin hypothesis of Alzheimer's disease. *Mol*
*Neurodegener*. 2024;19(1). https://doi.org/10.1186/s13024-024-00722-y.
31. Sarlus H, Heneka MT. Microglia in Alzheimer's disease. *Journal of Clinical*
*Investigation*. 2017;127(9):3240–3249.
32. Kinney JW, et al. Inflammation as a central mechanism in Alzheimer's disease.
*Alzheimer's and Dementia: Translational Research and Clinical Interventions*.
2018;4:575–590.
33. Heneka MT, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol*.
2015;14(4):388–405.
34. Brutkiewicz RR, et al. What would it take to prove that a chronic infection is a
causal agent in Alzheimer's disease? *Trends Neurosci*. [published online ahead of
print: 2025]. https://doi.org/10.1016/j.tins.2025.05.009.
35. Rubtsova K, Marrack P, Rubtsov A V. Sexual dimorphism in autoimmunity. *Journal*
*of Clinical Investigation*. 2015;125(6):2187–2193.
36. Jacobsen H, Klein SL. Sex Differences in Immunity to Viral Infections. *Front*
*Immunol*. 2021;12. https://doi.org/10.3389/fimmu.2021.720952.
37. Wilkinson NM, et al. Sex Differences in Immunity. *Annu Rev Immunol*. 2022;40:75–
94.
38. Dunn SE, Perry WA, Klein SL. Mechanisms and consequences of sex differences
in immune responses. *Nat Rev Nephrol*. 2024;20(1):37–55.
39. Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol*.
2016;16(10):626–638.
40. Mangold CA, et al. Sexually divergent induction of microglial-associated
neuroinflammation with hippocampal aging. *J Neuroinflammation*. 2017;14(1).
https://doi.org/10.1186/s12974-017-0920-8.

41. Wen H, et al. Altered immune response is associated with sex difference in vulnerability to Alzheimer's disease in human prefrontal cortex. *Brain Pathology*. 2025;35(3). https://doi.org/10.1111/bpa.13318.

42. Buckley RF. Sex differences in Alzheimer's disease risk: Are immune responses the key? *Brain*. 2022;145(10):3341–3342.

43. Villa A, et al. Sex-Specific Features of Microglia from Adult Mice. *Cell Rep*. 2018;23(12):3501–3511.

44. Meadows SM, et al. Hippocampal astrocytes induce sex-dimorphic effects on memory. *Cell Rep*. 2024;43(6). https://doi.org/10.1016/j.celrep.2024.114278.

45. Rurak GM, et al. Sex differences in developmental patterns of neocortical astroglia: A mouse translatome database. *Cell Rep*. 2022;38(5). https://doi.org/10.1016/j.celrep.2022.110310.

46. Guneykaya D, et al. Transcriptional and Translational Differences of Microglia from Male and Female Brains. *Cell Rep*. 2018;24(10):2773-2783.e6.

47. Kim GS, Uddin M. Sex-specific and shared expression profiles of vulnerability and resilience to trauma in brain and blood. *Biol Sex Differ*. 2020;11(1). https://doi.org/10.1186/s13293-020-00288-6.

48. Kundakovic M, Tickerhoof M. Epigenetic mechanisms underlying sex differences in the brain and behavior. *Trends Neurosci*. 2024;47(1):18–35.

49. Mottron L, et al. Sex differences in brain plasticity: A new hypothesis for sex ratio bias in autism Understanding the links between sex/gender and autism Dr Meng-Chuan Lai. *Mol Autism*. 2015;6(1). https://doi.org/10.1186/s13229-015-0024-1.

50. Garrett JE, Wellman CL. Chronic stress effects on dendritic morphology in medial prefrontal cortex: sex differences and estrogen dependence. *Neuroscience*. 2009;162(1):195–207.

51. Aguzzoli CS, et al. *World Alzheimer Report 2024 Global changes in attitudes to dementia Contributors: Survey translators*.

52. Fang M, et al. Lifetime risk and projected burden of dementia. *Nat Med*. 2025;31(3):772–776.

53. 2024 Alzheimer's disease facts and figures. *Alzheimer's and Dementia*. 2024;20(5):3708–3821.

54. Moutinho S. Women twice as likely to develop Alzheimer's disease as men — but scientists do not know why. *Nat Med*. 2025;31(3):704–707.

55. 2025 Alzheimer's disease facts and figures. *Alzheimer's & Dementia*. 2025;21(4). https://doi.org/10.1002/alz.70235.

56. Liberzon A, et al. The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Syst*. 2015;1(6):417–425.

57. Subramanian A, et al. *Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles*. 2005.

58. Serrano-Pozo A, et al. Neuropathological alterations in Alzheimer disease. *Cold Spring Harb Perspect Med*. 2011;1(1). https://doi.org/10.1101/cshperspect.a006189.

59. Hendrickx A, et al. Epigenetic Regulations of Immediate Early Genes Expression Involved in Memory Formation by the Amyloid Precursor Protein of Alzheimer Disease. *PLoS One*. 2014;9(6):e99467-.

60. Crapser JD, et al. Microglia facilitate loss of perineuronal nets in the Alzheimer's disease brain. *EBioMedicine*. 2020;58. https://doi.org/10.1016/j.ebiom.2020.102919.

61. Dan L, Zhang Z. Alzheimer's disease: an axonal injury disease? *Front Aging Neurosci*. 2023;15. https://doi.org/10.3389/fnagi.2023.1264448.

62. Corbett BF, et al. ΔFosB Regulates Gene Expression and Cognitive Dysfunction in
a Mouse Model of Alzheimer's Disease. *Cell Rep*. 2017;20(2):344–355.
63. Fawcett JW, Oohashi T, Pizzorusso T. The roles of perineuronal nets and the
perinodal extracellular matrix in neuronal function. *Nat Rev Neurosci*. 2019;20(8):451–
465.
64. Jankowsky JL, et al. Mutant presenilins specifically elevate the levels of the 42
residue β-amyloid peptide in vivo: evidence for augmentation of a 42-specific γ
secretase. *Hum Mol Genet*. 2004;13(2):159–170.
65. Keren-Shaul H, et al. A Unique Microglia Type Associated with Restricting
Development of Alzheimer's Disease. *Cell*. 2017;169(7):1276-1290.e17.
66. Sala Frigerio C, et al. The Major Risk Factors for Alzheimer's Disease: Age, Sex,
and Genes Modulate the Microglia Response to Aβ Plaques. *Cell Rep*.
2019;27(4):1293-1306.e6.
67. Mathys H, et al. Temporal Tracking of Microglia Activation in Neurodegeneration at
Single-Cell Resolution. *Cell Rep*. 2017;21(2):366–380.
68. Pelkmans W, et al. Astrocyte biomarkers GFAP and YKL-40 mediate early
Alzheimer's disease progression. *Alzheimer's and Dementia*. 2024;20(1):483–493.
69. Haag SM, et al. Targeting STING with covalent small-molecule inhibitors. *Nature*.
2018;559(7713):269–273.
70. Subramaniapillai S, et al. Sex and gender differences in cognitive and brain
reserve: Implications for Alzheimer's disease in women. *Front Neuroendocrinol*.
2021;60. https://doi.org/10.1016/j.yfrne.2020.100879.
71. Rosende-Roca M, et al. Exploring sex differences in Alzheimer's disease: a
comprehensive analysis of a large patient cohort from a memory unit. *Alzheimer's*
*Research and Therapy* . 2025;17(1). https://doi.org/10.1186/s13195-024-01656-9.
72. Boccalini C, et al. Sex differences in the association of Alzheimer's disease
biomarkers and cognition in a multicenter memory clinic study. *Alzheimer's Research*
*and Therapy* . 2025;17(1). https://doi.org/10.1186/s13195-025-01684-z.
73. Moutinho S. Women twice as likely to develop Alzheimer's disease as men — but
scientists do not know why. *Nat Med*. 2025;31(3):704–707.
74. Hammond TR, Marsh SE, Stevens B. Immune Signaling in Neurodegeneration.
*Immunity*. 2019;50(4):955–974.
75. Heppner FL, Ransohoff RM, Becher B. Immune attack: the role of inflammation in
Alzheimer disease. *Nat Rev Neurosci*. 2015;16(6):358–372.
76. Lopez-Atalaya JP, Bhojwani-Cabrera AM. Type I interferon signalling and
interferon-responsive microglia in health and disease. *FEBS Journal*. [published online
ahead of print: 2025]. https://doi.org/10.1111/febs.70126.
77. Roy E, Cao W. Glial interference: impact of type I interferon in neurodegenerative
diseases. *Mol Neurodegener*. 2022;17(1). https://doi.org/10.1186/s13024-022-00583-3.
78. Wang Q, et al. Brain derived β-interferon is a potential player in Alzheimer's
disease pathogenesis and cognitive impairment. *Alzheimer's Research and Therapy* .
2024;16(1). https://doi.org/10.1186/s13195-024-01644-z.
79. Roy ER, et al. Concerted type I interferon signaling in microglia and neural cells
promotes memory impairment associated with amyloid β plaques. *Immunity*.
2022;55(5):879-894.e6.
80. Roy ER, et al. Type I interferon response drives neuroinflammation and synapse
loss in Alzheimer disease. *Journal of Clinical Investigation*. 2020;130(4):1912–1930.

81. Tesseur I, et al. Deficiency in neuronal TGF-β signaling promotes
neurodegeneration and Alzheimer's pathology. *Journal of Clinical Investigation*.
2006;116(11):3060–3069.
82. Kapoor M, Chinnathambi S. TGF-β1 signalling in Alzheimer's pathology and
cytoskeletal reorganization: a specialized Tau perspective. *J Neuroinflammation*.
2023;20(1). https://doi.org/10.1186/s12974-023-02751-8.
83. Minter MR, et al. Deletion of the type-1 interferon receptor in APPSWE/PS1ΔE9
mice preserves cognitive function and alters glial phenotype. *Acta Neuropathol*
*Commun*. 2016;4(1):72.
84. Sanchez-Garcia MA, et al. *Inactivation of the PHD3-FOXO3 axis blunts the type I*
*interferon response in microglia and ameliorates Alzheimer's disease progression*.
2025.
85. Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol*.
2016;16(10):626–638.
86. Rubtsova K, Marrack P, Rubtsov A V. Sexual dimorphism in autoimmunity. *Journal*
*of Clinical Investigation*. 2015;125(6):2187–2193.
87. Ghosh S, Klein RS. Sex Drives Dimorphic Immune Responses to Viral Infections.
*The Journal of Immunology*. 2017;198(5):1782–1790.
88. Giefing-Kröll C, et al. How sex and age affect immune responses, susceptibility to
infections, and response to vaccination. *Aging Cell*. 2015;14(3):309–321.
89. Wilkinson NM, et al. Sex Differences in Immunity. *Annu Rev Immunol*. 2022;40:75–
94.
90. Markle JG, Fish EN. SeXX matters in immunity. *Trends Immunol*. 2014;35(3):97–
104.
91. Wen H, et al. Altered immune response is associated with sex difference in
vulnerability to Alzheimer's disease in human prefrontal cortex. *Brain Pathology*.
2025;35(3). https://doi.org/10.1111/bpa.13318.
92. Buckley RF. Sex differences in Alzheimer's disease risk: Are immune responses
the key? *Brain*. 2022;145(10):3341–3342.
93. Mangold CA, et al. Sexually divergent induction of microglial-associated
neuroinflammation with hippocampal aging. *J Neuroinflammation*. 2017;14(1).
https://doi.org/10.1186/s12974-017-0920-8.
94. Fyfe I. Immune responses influence sex differences in Alzheimer disease. *Nat Rev*
*Neurol*. 2024;20(12):691.
95. Coales I, et al. Alzheimer's disease-related transcriptional sex differences in
myeloid cells. *J Neuroinflammation*. 2022;19(1). https://doi.org/10.1186/s12974-022-
02604-w.
96. Lopez-Lee C, et al. Tlr7 drives sex differences in age- and Alzheimer's disease–
related demyelination. *Science (1979)*. 2024;386(6725).
https://doi.org/10.1126/science.adk7844.
97. Stephen TL, et al. APOE genotype and sex affect microglial interactions with
plaques in Alzheimer's disease mice. *Acta Neuropathol Commun*. 2019;7(1).
https://doi.org/10.1186/s40478-019-0729-z.
98. Mifflin MA, et al. Sex differences in the IntelliCage and the Morris water maze in the
APP/PS1 mouse model of amyloidosis. *Neurobiol Aging*. 2021;101:130–140.
99. Sala Frigerio C, et al. The Major Risk Factors for Alzheimer's Disease: Age, Sex,
and Genes Modulate the Microglia Response to Aβ Plaques. *Cell Rep*.
2019;27(4):1293-1306.e6.

100. Yang J-T, et al. Sex Differences in Neuropathology and Cognitive Behavior in
APP/PS1/tau Triple-Transgenic Mouse Model of Alzheimer's Disease. *Neurosci Bull*.
2018;34(5):736–746.
101. Jiao S-S, et al. Sex Dimorphism Profile of Alzheimer's Disease-Type Pathologies
in an APP/PS1 Mouse Model. *Neurotox Res*. 2016;29(2):256–266.
102. Serrano-Pozo A, et al. Neuropathological alterations in Alzheimer disease. *Cold*
*Spring Harb Perspect Med*. 2011;1(1). https://doi.org/10.1101/cshperspect.a006189.
103. Corbett BF, et al. ΔFosB Regulates Gene Expression and Cognitive Dysfunction
in a Mouse Model of Alzheimer's Disease. *Cell Rep*. 2017;20(2):344–355.
104. Hendrickx A, et al. Epigenetic Regulations of Immediate Early Genes Expression
Involved in Memory Formation by the Amyloid Precursor Protein of Alzheimer Disease.
*PLoS One*. 2014;9(6):e99467-.
105. Crapser JD, et al. Microglia facilitate loss of perineuronal nets in the Alzheimer's
disease brain. *EBioMedicine*. 2020;58. https://doi.org/10.1016/j.ebiom.2020.102919.
106. Pelkmans W, et al. Astrocyte biomarkers GFAP and YKL-40 mediate early
Alzheimer's disease progression. *Alzheimer's and Dementia*. 2024;20(1):483–493.
107. Dan L, Zhang Z. Alzheimer's disease: an axonal injury disease? *Front Aging*
*Neurosci*. 2023;15. https://doi.org/10.3389/fnagi.2023.1264448.
108. Deczkowska A, Baruch K, Schwartz M. Type I/II Interferon Balance in the
Regulation of Brain Physiology and Pathology. *Trends Immunol*. 2016;37(3):181–192.
109. Baruch K, et al. Aging-induced type I interferon response at the choroid plexus
negatively affects brain function. *Science (1979)*. 2014;346(6205):89–93.
110. Schwartz M, Deczkowska A. Neurological Disease as a Failure of Brain-Immune
Crosstalk: The Multiple Faces of Neuroinflammation. *Trends Immunol*.
2016;37(10):668–679.
111. Mendoza-Fernández V, Andrew RD, Barajas-López C. Interferon-α inhibits long-
term potentiation and unmasks a long-term depression in the rat hippocampus. *Brain*
*Res*. 2000;885(1):14–24.
112. Blank T, et al. Brain Endothelial- and Epithelial-Specific Interferon Receptor Chain
1 Drives Virus-Induced Sickness Behavior and Cognitive Impairment. *Immunity*.
2016;44(4):901–912.
113. Gulen MF, et al. cGAS–STING drives ageing-related inflammation and
neurodegeneration. *Nature*. 2023;620(7973):374–380.
114. Crow YJ. CNS disease associated with enhanced type I interferon signalling.
*Lancet Neurol*. 2024;23(11):1158–1168.
115. Hu X, et al. Emerging role of STING signalling in CNS injury: inflammation,
autophagy, necroptosis, ferroptosis and pyroptosis. *J Neuroinflammation*. 2022;19(1).
https://doi.org/10.1186/s12974-022-02602-y.
116. Feng Y, et al. Human herpesvirus-associated transposable element activation in
human aging brains with Alzheimer's disease. *Alzheimer's and Dementia*. 2025;21(2).
https://doi.org/10.1002/alz.14595.
117. Huang SY, et al. Herpesvirus infections and Alzheimer's disease: a Mendelian
randomization study. *Alzheimers Res Ther*. 2021;13(1). https://doi.org/10.1186/s13195-
021-00905-5.
118. Eimer WA, et al. Alzheimer's Disease-Associated β-Amyloid Is Rapidly Seeded by
Herpesviridae to Protect against Brain Infection. *Neuron*. 2018;99(1):56-63.e3.

119. Romanescu C, Schreiner TG, Mukovozov I. The Role of Human Herpesvirus 6 Infection in Alzheimer's Disease Pathogenicity—A Theoretical Mosaic. *J Clin Med*. 2022;11(11). https://doi.org/10.3390/jcm11113061.

120. Crivelli L, et al. Changes in cognitive functioning after COVID-19: A systematic review and meta-analysis. *Alzheimer's and Dementia*. 2022;18(5):1047–1066.

121. Bruno F, et al. Alzheimer's disease as a viral disease: Revisiting the infectious hypothesis. *Ageing Res Rev*. 2023;91. https://doi.org/10.1016/j.arr.2023.102068.

122. Levine KS, et al. Virus exposure and neurodegenerative disease risk across national biobanks. *Neuron*. 2023;111(7):1086-1093.e2.

123. Hu Y, et al. Infections among individuals with multiple sclerosis, Alzheimer's disease and Parkinson's disease. *Brain Commun*. 2023;5(2). https://doi.org/10.1093/braincomms/fcad065.

124. Ecarnot F, et al. Dementia, infections and vaccines: 30 years of controversy. *Aging Clin Exp Res*. 2023;35(6):1145–1160.

125. Pomirchy M, et al. Herpes Zoster Vaccination and Dementia Occurrence. *JAMA*. 2025;333(23):2083–2092.

126. Schnier C, et al. Reduced dementia incidence after varicella zoster vaccination in Wales 2013–2020. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2022;8(1). https://doi.org/10.1002/trc2.12293.

127. Lehrer S. *Vaccination Reduces Risk of Alzheimer's Disease, Parkinson's Disease, and Other Neurodegenerative Disorders*. 2022.

128. Eyting M, et al. A natural experiment on the effect of herpes zoster vaccination on dementia. *Nature*. 2025;641(8062):438–446.

129. McNab F, et al. Type I interferons in infectious disease. *Nat Rev Immunol*. 2015;15(2):87–103.

130. Hosseini S, et al. Type I Interferon Receptor Signaling in Astrocytes Regulates Hippocampal Synaptic Plasticity and Cognitive Function of the Healthy CNS. *Cell Rep*. 2020;31(7). https://doi.org/10.1016/j.celrep.2020.107666.

131. Deczkowska A, et al. Mef2C restrains microglial inflammatory response and is lost in brain ageing in an IFN-I-dependent manner. *Nat Commun*. 2017;8(1). https://doi.org/10.1038/s41467-017-00769-0.

132. Heneka MT, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol*. 2015;14(4):388–405.

133. Sarlus H, Heneka MT. Microglia in Alzheimer's disease. *Journal of Clinical Investigation*. 2017;127(9):3240–3249.

134. Kinney JW, et al. Inflammation as a central mechanism in Alzheimer's disease. *Alzheimer's and Dementia: Translational Research and Clinical Interventions*. 2018;4:575–590.

135. Lall D, et al. C9orf72 deficiency promotes microglial-mediated synaptic loss in aging and amyloid accumulation. *Neuron*. 2021;109(14):2275-2291.e8.

136. Hur J-Y, et al. The innate immunity protein IFITM3 modulates γ-secretase in Alzheimer's disease. *Nature*. 2020;586(7831):735–740.

137. Jin M, et al. Tau activates microglia via the PQBP1-cGAS-STING pathway to promote brain inflammation. *Nat Commun*. 2021;12(1). https://doi.org/10.1038/s41467-021-26851-2.

138. Dhir A, et al. Mitochondrial double-stranded RNA triggers antiviral signalling in humans. *Nature*. 2018;560(7717):238–242.

139. Quan S, et al. The neuroimmune nexus: unraveling the role of the mtDNA-cGAS-
STING signal pathway in Alzheimer's disease. *Molecular Neurodegeneration* .
2025;20(1). https://doi.org/10.1186/s13024-025-00815-2.
140. Udeochu JC, et al. Tau activation of microglial cGAS–IFN reduces MEF2C-
mediated cognitive resilience. *Nat Neurosci*. 2023;26(5):737–750.
141. Huang Y, et al. Mechanism and therapeutic potential of targeting cGAS-STING
signaling in neurological disorders. *Mol Neurodegener*. 2023;18(1).
https://doi.org/10.1186/s13024-023-00672-x.
142. Xie X, et al. Activation of innate immune cGAS-STING pathway contributes to
Alzheimer's pathogenesis in 5×FAD mice. *Nat Aging*. 2023;3(2):202–212.
143. Chung S, et al. Blockade of STING activation alleviates microglial dysfunction and
a broad spectrum of Alzheimer's disease pathologies. *Exp Mol Med*. 2024;56(9):1936–
1951.
144. Carling GK, et al. Alzheimer's disease-linked risk alleles elevate microglial cGAS-
associated senescence and neurodegeneration in a tauopathy model. *Neuron*.
2024;112(23):3877-3896.e8.
145. Ghosh M, et al. Mutant p53 suppresses innate immune signaling to promote
tumorigenesis. *Cancer Cell*. 2021;39(4):494-508.e5.
146. Ghosh M, et al. p53 engages the cGAS/STING cytosolic DNA sensing pathway for
tumor suppression. *Mol Cell*. 2023;83(2):266-280.e6.
147. Rahimi J, Kovacs GG. Prevalence of mixed pathologies in the aging brain.
*Alzheimers Res Ther*. 2014;6(9). https://doi.org/10.1186/s13195-014-0082-1.
148. Aizenstein HJ, et al. Frequent amyloid deposition without significant cognitive
impairment among the elderly. *Arch Neurol*. 2008;65(11):1509–1517.
149. Barroeta-Espar I, et al. Distinct cytokine profiles in human brains resilient to
Alzheimer's pathology. *Neurobiol Dis*. 2019;121:327–337.
150. Perez-Nievas BG, et al. Dissecting phenotypic traits linked to human resilience to
Alzheimer's pathology. *Brain*. 2013;136(8):2510–2526.
151. Wang B, et al. The STING inhibitor C-176 attenuates MPTP-induced
neuroinflammation and neurodegeneration in mouse parkinsonian models. *Int
Immunopharmacol*. 2023;124:110827.
152. Kong L, et al. mtDNA-STING Axis Mediates Microglial Polarization via IRF3/NF-κB
Signaling After Ischemic Stroke. *Front Immunol*. 2022;13.
https://doi.org/10.3389/fimmu.2022.860977.
153. Zhang S, et al. Potential Therapeutic Value of the STING Inhibitors. *Molecules*.
2023;28(7). https://doi.org/10.3390/molecules28073127.
154. Decout A, et al. The cGAS–STING pathway as a therapeutic target in
inflammatory diseases. *Nat Rev Immunol*. 2021;21(9):548–569.
155. Zitvogel L, et al. Type I interferons in anticancer immunity. *Nat Rev Immunol*.
2015;15(7):405–414.
156. Hopfner K-P, Hornung V. Molecular mechanisms and cellular functions of cGAS–
STING signalling. *Nat Rev Mol Cell Biol*. 2020;21(9):501–521.
157. Schwartz DM, et al. JAK inhibition as a therapeutic strategy for immune and
inflammatory diseases. *Nat Rev Drug Discov*. 2017;16(12):843–862.
158. Tanaka Y, Tummala R. Anifrolumab, a monoclonal antibody to the type I
interferon receptor subunit 1, for the treatment of systemic lupus erythematosus: an
overview from clinical trials. *Mod Rheumatol*. 2021;31(1):1–12.

159. Jones SA, Morand EF. Targeting Interferon Signalling in Systemic Lupus
Erythematosus: Lessons Learned. *Drugs*. 2024;84(6):625–635.
160. Parkhurst CN, et al. Microglia promote learning-dependent synapse formation
through brain-derived neurotrophic factor. *Cell*. 2013;155(7):1596–1609.
161. Steinbrecher KA, et al. Loss of Epithelial RelA Results in Deregulated Intestinal
Proliferative/Apoptotic Homeostasis and Susceptibility to Inflammation. *The Journal of*
*immunology (1950)*. 2008;180(4):2588–2599.
162. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. *Methods*. 2001;25(4):402–408.
163. Chen S, et al. Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*.
Oxford University Press; 2018:i884–i890.
164. Kim D, et al. Graph-based genome alignment and genotyping with HISAT2 and
HISAT-genotype. *Nat Biotechnol*. 2019;37(8):907–915.
165. Liao Y, Smyth GK, Shi W. FeatureCounts: An efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics*. 2014;30(7):923–
930.
166. Kim D, et al. Graph-based genome alignment and genotyping with HISAT2 and
HISAT-genotype. *Nat Biotechnol*. 2019;37(8):907–915.
167. Danecek P, et al. Twelve years of SAMtools and BCFtools. *Gigascience*.
2021;10(2). https://doi.org/10.1093/gigascience/giab008.
168. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV):
High-performance genomics data visualization and exploration. *Brief Bioinform*.
2013;14(2):178–192.
169. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12).
https://doi.org/10.1186/s13059-014-0550-8.
170. Zhang Y, Parmigiani G, Johnson WE. ComBat-seq: Batch effect adjustment for
RNA-seq count data. *NAR Genom Bioinform*. 2020;2(3).
https://doi.org/10.1093/nargab/lqaa078.
171. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. Cham: Springer
International Publishing; 2016.
172. Slowikowski K. Automatically Position Non-Overlapping Text Labels with "ggplot2"
[R package ggrepel version 0.9.6] [Internet]. *CRAN: Contributed Packages*. 2024.
https://doi.org/10.32614/CRAN.PACKAGE.GGREPEL.
173. Kolde R. Pretty Heatmaps [R package pheatmap version 1.0.13] [Internet]. *CRAN:*
*Contributed Packages*. 2025. https://doi.org/10.32614/CRAN.PACKAGE.PHEATMAP.
174. Kuleshov M V., et al. Enrichr: a comprehensive gene set enrichment analysis web
server 2016 update. *Nucleic Acids Res*. 2016;44(1):W90–W97.
175. Wickham H, et al. Welcome to the Tidyverse. *J Open Source Softw*.
2019;4(43):1686.
176. Wilke CO. Streamlined Plot Theme and Plot Annotations for "ggplot2" [R package
cowplot version 1.2.0] [Internet]. *CRAN: Contributed Packages*. 2025.
https://doi.org/10.32614/CRAN.PACKAGE.COWPLOT.
177. Auguie B. Miscellaneous Functions for "Grid" Graphics [R package gridExtra
version 2.3] [Internet]. *CRAN: Contributed Packages*. 2017.
https://doi.org/10.32614/CRAN.PACKAGE.GRIDEXTRA.
1368
1369

# Figure legends

**Fig. 1. Enhanced interferon signaling correlates with female-biased Alzheimer's disease risk and prevalence.** (A) Diagram illustrating lifetime risk and prevalence of AD, with women showing higher risk (20% vs. 10%) and prevalence (61% vs. 39%) compared to men (Alzheimer's Facts and Figures 2024). (B) Cohort overview of RNA-sequencing data from the MSBB, comprising 34 female and 32 male AD patients (aged 60–90 years) for parahippocampal gyrus analysis. (C) Volcano plot of differential gene expression in female vs. male AD brains. Significantly upregulated genes (FDR < 0.05) are shown in orange, and genes with $0.05 \leq$ FDR $< 0.1$ are shown in dark blue. Interferon alpha signaling genes are highlighted. (D) Heatmap showing elevated expression of the top differentially expressed (FDR < 0.1) interferon-related genes in female AD samples compared to male samples. (E) Functional enrichment analysis of significantly upregulated genes (FDR < 0.05) in female AD brains showing enrichment of interferon-alpha and interferon-gamma response pathways. (F) GSEA plot showing coordinated upregulation of the interferon alpha response pathway in female AD brains.

**Fig. 2. Sex-specific amyloid pathology and interferon signaling in APP/PS1 mice.** (A) Schematic diagram of the APPswe/PSEN1dE9 (APP/PS1) mouse model, with experiments conducted at 6 months of age. (B) IHC with 6E10 antibody showing Aβ plaque density in cortex (CTX, $**P < 0.01$) and hippocampus (HC, $*P < 0.05$) of male and female mice, with higher burden in females (quantified graphs, Mann-Whitney test). (C) Thioflavin T staining of fibrillar Aβ deposits in CTX ($**P < 0.01$) and HC ($*P < 0.05$), with increased density in females (quantified graphs, Mann-Whitney test). (D) OC antibody staining of β-sheet-rich Aβ conformations in CTX and HC, showing greater signal in females (quantified graphs, $*P < 0.05$, Mann-Whitney test). (E) c-FOS immunostaining in dentate gyrus, indicating reduced neuronal activity in female mice (quantified graphs, $*P < 0.05$, Mann-Whitney test). (F) PNNs stained with PV/WFA in CA1, with lower proportion of PV+ interneurons in females (quantified graphs, $**P < 0.01$, Mann-Whitney test). (G) pNF staining for axonal damage in HC, with higher levels in females (quantified graphs, $*P < 0.05$, Mann-Whitney test). (H) PCA of hippocampal RNA-seq data, showing greater transcriptomic divergence in female APP/PS1 mice. (I) Volcano plot of differentially expressed genes in female vs. male hippocampus, highlighting upregulated interferon alpha genes (e.g., *Irf7*, *B2m*). (L) GFAP immunohistochemistry in HC, with increased cell density in females (quantified graphs, $*P < 0.05$, Mann-Whitney test). (K) Functional enrichment analysis indicating an enrichment in the interferon alpha and gamma response pathways in female APP/PS1 mice. (L) Ifitm3 immunohistochemistry in HC, with increased labeling in females (quantified graphs, $*P < 0.05$, Mann-Whitney test). Data from 6-month-old APP/PS1 mice. IHC: n = 12 (6 females, 6 males). RNA-seq: n = 16 (8 females, 8 males). Data represent mean ± SEM.

**Fig. 3. Poly(I:C)-induced interferon signaling elicits Alzheimer-like pathology**. (A) Schematic of experimental design with intraperitoneal poly(I:C) administration (12 mg/kg) to 3-month-old wild-type C57BL/6 mice, analyzed at 24 and 72 hours, and compared to APP/PS1 mice using microglia RNA-sequencing and immunohistochemistry (IHC). (B) Volcano plot of differentially expressed genes in hippocampus 24 hours post-poly(I:C), highlighting upregulated interferon alpha-related genes (e.g., *Irf7*, *Stat2*, *Ifit3*). (C) Functional enrichment analysis showing activation of interferon alpha/gamma pathways in poly(I:C)-treated mice. (D) Ifitm3

immunohistochemistry in hippocampus, with increased labeling in poly(I:C) 24h (*$P <$ 0.05), 72h (**$P < 0.01$), and APP/PS1 (**$P < 0.01$) compared to controls (quantified graphs, one-way ANOVA with Bonferroni post hoc test). (E) GFAP immunohistochemistry in hippocampus, with increased labeling in poly(I:C) 24h (**$P <$ 0.01), 72h (****$P < 0.0001$), and APP/PS1 (****$P < 0.0001$) compared to controls (quantified graphs, one-way ANOVA with Bonferroni post hoc test). (F) c-FOS immunostaining in dentate gyrus, showing reduced neuronal activity in poly(I:C) 72h (**$P < 0.01$), and APP/PS1(****$P < 0.0001$) vs. controls (quantified graphs, one-way ANOVA with Bonferroni post hoc test). (G) PNNs stained with PV/WFA in CA1, with lower proportion of PV+ interneurons in poly(I:C) 72h (****$P < 0.0001$), and APP/PS1 (*$P < 0.05$) vs. controls (quantified graphs, one-way ANOVA with Bonferroni post hoc test). (H) pNF staining for axonal damage in hippocampus, with higher levels in poly(I:C) 72h (*$P < 0.05$), and APP/PS1 (**$P < 0.01$) vs. controls (quantified graphs, one-way ANOVA with Bonferroni post hoc test). Data from wild-type and APP/PS1 mice. IHC: n = 8 (control), 4 (Poly(I:C) 24h), 4 (Poly(I:C) 72h), and 6 (APP/PS1). Poly(I:C) groups included only males; other groups were sex-balanced (3 females, 3 males for APP/PS1; 4 females, 4 males for controls). RNA-seq: n = 4 (control), 3 (Poly(I:C) 24h). Data represent mean ± SEM.

**Fig. 4. Enhanced interferon signaling exacerbates Alzheimer's pathology.** (A) Schematic of APP/PS1 (AD) crossed with microglia-specific *Rela* knockout (AD hiIFN) model, with tamoxifen induction at 3 months and analysis at 6 months using qPCR, bulk-RNAseq and immunohistochemistry (IHC). (B) qPCR showing *Rela* deletion efficiency in microglia (exons 1-3 and 6-7) in AD hiIFN vs. AD (quantified graphs, **$P < 0.01$, Mann-Whitney test). (C) Microglia qPCR showing increased expression of interferon-related genes (e.g., *Irf7, Ifitm3)* in AD hiIFN hippocampus (quantified graphs, **$P < 0.01$, Mann-Whitney test). (D) Volcano plot of differentially expressed genes in AD hiIFN vs. APP/PS1 controls, highlighting upregulated interferon alpha genes. (E) Functional enrichment analysis showing interferon alpha/gamma response pathways in AD hiIFN. (F) Ifitm3 immunohistochemistry in hippocampus, with higher labeling in AD hiIFN vs. AD (quantified graphs, **$P < 0.01$, Mann-Whitney test). (G) GFAP immunohistochemistry for astrogliosis in hippocampus, showing increased density in AD hiIFN vs AD (quantified graphs, **$P < 0.01$, Mann-Whitney test). (H) c-FOS immunohistochemistry in dentate gyrus, indicating reduced neuronal activity in AD hiIFN vs. AD (quantified graphs, *$P < 0.05$, Mann-Whitney test). (I) PNNs stained with PV/WFA in CA1, with lower proportion of PV+ interneurons in AD hiIFN vs. AD (quantified graphs, *$P < 0.05$, Mann-Whitney test). (J) pNF staining for axonal damage in hippocampus, with higher levels in AD hiIFN vs. AD (quantified graphs, **$P < 0.01$, Mann-Whitney test). (K) 6E10 immunohistochemistry for Aβ plaque density, showing reduced burden in AD hiIFN vs. AD both in cortex (CTX) and hippocampus (HC) (quantified graphs, *$P < 0.05$, Mann-Whitney test). (L) Thioflavin T staining for fibrillar Aβ deposits, with lower density in AD hiIFN vs. AD in CTX (*$P < 0.05$) and HC (**$P < 0.01$), (quantified graphs, Mann-Whitney test). (M) OC staining for β-sheet-rich Aβ conformations, with reduced signal in AD hiIFN vs. AD in CTX and HC, (quantified graphs, **$P < 0.01$, Mann-Whitney test). Data from 6-month-old AD hiIFN and APP/PS1 mice. IHC: n = 6 per group (3 females, 3 males). RNA-seq: n = 8 per group (4 females, 4 males). Data represent mean ± SEM.

**Fig. 5. STING inhibition alleviates Alzheimer's pathology in APP/PS1 mice.** (A) Schematic of experimental design with STING inhibitor C-176 (10 mg/kg) administration

to 3-month-old APP/PS1 mice (AD loIFN), treated until 6 months, analyzed using bulk RNA-seq and immunohistochemistry (IHC). (B) Volcano plot of differentially expressed genes in hippocampus of treated vs untreated females, highlighting downregulated interferon alpha genes. (C) Functional enrichment analysis showing reduced interferon alpha/gamma response pathways in untreated vs. treated females. (D) Ifitm3 immunohistochemistry in hippocampus, with lower labeling in treated AD loIFN vs. untreated AD mice (quantified graphs, *$P < 0.05$, Mann-Whitney test). (E) GFAP immunohistochemistry for astrogliosis in hippocampus, showing decreased density in treated vs. untreated mice (quantified graphs, *$P < 0.05$, Mann-Whitney test). (F) c-FOS immunohistochemistry in dentate gyrus, indicating increased neuronal activity in treated vs. untreated mice (quantified graphs, ***$P < 0.001$, Mann-Whitney test). (G) PNNs stained with PV/WFA in CA1, with higher proportion of PV+ interneurons in treated AD loIFN vs. untreated AD mice (quantified graphs, **$P < 0.01$, Mann-Whitney test). (H) pNF staining for axonal damage in hippocampus, with lower levels in treated vs. untreated mice (quantified graphs, **$P < 0.01$, Mann-Whitney test). (I) 6E10 immunohistochemistry for Aβ plaque density, showing non-significant changes in AD loiIFN vs. AD both in cortex (CTX) and hippocampus (HC) (quantified graphs, ns, $P > 0.05$, Mann-Whitney test). (J) Thioflavin T staining for fibrillar Aβ deposits, with no changes in treated vs. untreated mice both in CTX and HC quantified graphs, ns, $P > 0.05$, Mann-Whitney test). (K) OC staining for β-sheet-rich Aβ conformations, with increased signal in AD loIFN vs. AD in CTX (*$P < 0.05$) and non-significant changes in HC (ns, $P > 0.05$) (quantified graphs, Mann-Whitney test). Data from 6-month-old AD loIFN and APP/PS1 mice. IHC: n = 6 per group (3 females, 3 males). RNA-seq: n = 8 AD (females), 4 AD loIFN (females). Data represent mean ± SEM.

**Fig. 6. Downregulation of Interferon Signaling Preserves Cognitive Performance in Female APP/PS1 Mice.** (A) Schematic of experimental design for cognitive testing in male and female APP/PS1 mice (AD) and female APP/PS1 mice treated with C-176 (AD loIFN) from 3 to 6 months. (B) Heatmap of the top 15 upregulated interferon-alpha–related genes in hippocampal tissue from female AD mice, showing elevated expression compared to both male AD mice and C-176–treated female AD mice. (C) Elevated Plus Maze (EPM) results, showing increased time in open arms in female AD mice compared to WT females (****$P < 0.0001$), with no differences in males (ns, $P > 0.05$) (quantified graphs, ***$P < 0.001$, one-way ANOVA with Bonferroni post hoc test). (D) Y-Maze spontaneous alternation, revealing reduced alternation in female AD mice compared to WT (*$P < 0.05$), with no differences in males (ns, $P > 0.05$) (quantified graphs, one-way ANOVA with Bonferroni post hoc test). (E) Novel Object Location Test (NOLT) results, indicating impaired spatial memory in both male and female AD mice compared to WT (quantified graphs, ***$P < 0.001$, ****$P < 0.0001$, ns $P > 0.05$, one-way ANOVA with Bonferroni post hoc test). (F) EPM in AD C-176-treated females, indicating normalized time in open arms compared to treated WT females (ns, $P > 0.05$) and demonstrating cognitive improve compared to untreated AD females (*$P < 0.05$) (quantified graphs, ***$P < 0.001$, one-way ANOVA with Bonferroni post hoc test). (G) Y-maze in AD C-176-treated females, indicating normalized time in open arms compared to treated WT females (ns, $P > 0.05$) (quantified graphs, *$P < 0.05$, one-way ANOVA with Bonferroni post hoc test). (H) NOLT in C-176-treated female AD mice (AD loIFN), showing restored discrimination of novel object location compared to untreated AD females (quantified graphs, ***$P < 0.001$, ns $P > 0.05$, one-way ANOVA with Bonferroni post hoc test). Data from 6-month-old WT, AD, and AD loIFN mice. n = 5 (male WT), 6 (male AD), 5

1518    (female WT), 5 (female AD), 5 (female WT + vehicle), 5 (female WT + C-176), 5 (female

1519    AD + vehicle), and 5 (female AD + C-176). Data represent mean ± SEM.

**Figure 1**

**Figure 2**

**Figure 3**

**Figure 4**

**Figure 5**

**Figure 6**

# Supplementary figures

**Extended Data Fig. 1. Increased interferon signaling in Alzheimer's disease is sex-specific and absent in controls.** (A) PCA of AD PHG RNA-seq data from the MSBB cohort (including samples from individuals aged 60–90 at death), with one misannotated male sample (excluded from analyses) clustering with female samples. (B) Heatmap of canonical sex chromosome gene expression in PHG from AD patients aged 60 to 90 years, confirming accurate sex annotation except for the misannotated male sample (bold). n = 32 (male AD), 34 (female AD). (C–E) Functional enrichment of differentially expressed genes (FDR < 0.05) in PHG tissue of AD patients (60 to 90 years). (C) Upregulated pathways in AD versus age-matched control samples. (D) Upregulated pathways in male AD versus age-matched male controls. (E) Upregulated pathways in female AD versus age-matched female controls. Female AD samples exhibit stronger interferon signatures, whereas in male AD samples the enrichment of epithelial–mesenchymal transition and inflammatory response pathways are prominent. (F) Volcano plot of differential gene expression in PHG of control individuals (female vs. male, aged from 60 to 90 years), showing no significant changes in interferon-related genes in females (gray, non-significant). n = 23 (male AD), 19 (female AD). (G) Functional enrichment analysis of differentially expressed genes (FDR < 0.05) in the PHG from control individuals (female vs. male, aged from 60 to 90 years), showing no significant activation of interferon alpha or gamma response pathways. n = 23 (male AD), 19 (female AD). (H) GSEA plot of the interferon alpha response pathway in control individuals (female vs. male, aged from 60 to 90 years). n = 23 (male AD), 19 (female AD).

**Extended Data Fig. 2. Sex-dependent neuropathological and transcriptomic changes in APP/PS1 Mice.** (A) Quantification of Aβ plaque size by 6E10 immunohistochemistry in the cortex (CTX) and hippocampus (HC) of 6-month-old male and female APP/PS1 mice, showing no significant differences in plaque size between sexes (ns, $P > 0.05$, Mann-Whitney test). (B) Quantification of fibrillar Aβ deposit size by Thioflavin T staining in CTX and HC, confirming no significant differences in deposit size between sexes (ns, $P > 0.05$, Mann-Whitney test). (C) Quantification of β-sheet-rich Aβ conformation size by OC antibody staining in CTX and HC, showing no significant differences between sexes (ns, $P > 0.05$, Mann-Whitney test). (D) Quantification of WFA-positive structure number in CA1, showing no significant differences between (ns, $P > 0.05$, Mann-Whitney test). (E) Volcano plot of differential gene expression in HC contrasting APP/PS1 vs control, highlighting top 15 significant genes (green, $P < 0.05$ in APP/PS1; dark grey, $P < 0.05$ in control; gray, non-significant). (F) Functional enrichment analysis from HC tissue of mice showing enriched pathways for APP/PS1 mice contrasted against control mice, where no interferon pathway was highlighted. (G) Quantification of GFAP immunohistochemistry density in the cortex, showing increased astrogliosis in female APP/PS1 mice (**$P \leq 0.01$), Mann-Whitney test). (H) GSEA plot of the interferon alpha response pathway in the hippocampi of female vs. male APP/PS1 mice. (I) Quantification of IFITM3 immunohistochemistry intensity in the cortex,

showing increased labeling in female APP/PS1 (**$P \leq 0.01$), Mann-Whitney test). Data from 6-month-old APP/PS1 mice. IHC: n = 7 (control), 4 (Poly(I:C) 24h), 4 (Poly(I:C) 72h), and 6 (APP/PS1). IHC: n = 12 (6 females, 6 males). RNA-seq: n = 8 AD females, 8 AD males, 8 WT females, 8 WT males. Data represent mean ± SEM.

**Extended Data Fig. 3. Poly(I:C)-induced interferon signaling and neuropathological changes in wild-type mice.** (A) PCA of hippocampal RNA-seq data from 3-month-old wild-type C57BL/6 mice treated with poly(I:C) (12 mg/kg) and analyzed at 24 hours, compared to untreated controls, showing transcriptomic divergence among the two groups. (B) GSEA plot of the interferon alpha response pathway in the HC of poly(I:C)-treated mice (24 hours) vs. untreated controls, confirming significant enrichment of interferon alpha signaling. (C) Quantification of IFITM3 immunohistochemistry intensity in the cortex, showing increased labeling in poly(I:C)-treated mice at 24 hours (****$P < 0.0001$), and 72 hours (*$P < 0.05$), and in APP/PS1 mice (*$P < 0.05$) compared to controls (Mann-Whitney test). (D) Quantification of GFAP immunohistochemistry intensity in the cortex, indicating no significant changes in astrogliosis in poly(I:C)-treated mice at 24 hours (ns, $P > 0.05$) and 72 hours (ns, $P > 0.05$), and an increased in APP/PS1 mice (****$P < 0.0001$) compared to controls (Mann-Whitney test). (E) Quantification of WFA-positive structure number in CA1, showing no significant changes in poly(I:C)-treated mice at 24 hours (ns, $P > 0.05$), and fewer structures at 72 hours (**$P < 0.01$) and APP/PS1 mice (*$P < 0.05$) compared to controls (Mann-Whitney test). Data from 3-month-old wild-type C57BL/6 mice (RNA-seq at 24 hours, immunohistochemistry at 24 and 72 hours) and 6-month-old APP/PS1 mice. IHC: n = 7 (control), 4 (Poly(I:C) 24h), 4 (Poly(I:C) 72h), and 6 (APP/PS1). Poly(I:C) groups included only males; other groups were sex-balanced (3 females, 3 males for APP/PS1; 3 females, 4 males for controls). RNA-seq: n = 4 (control), 3 (Poly(I:C) 24h). Data represent mean ± SEM.

**Extended Data Fig. 4. Neuropathological and transcriptomic effects of enhanced interferon signaling in APP/PS1 mice.** (A) PCA of hippocampal RNA-seq data from 6-month-old APP/PS1 (AD) and microglia-specific *Rela* knockout (AD hiIFN) mice, showing transcriptomic divergence between the two groups. (B) GSEA plot of the interferon alpha response pathway in the hippocampus of AD hiIFN vs. AD mice, confirming significant enrichment of interferon alpha signaling. (C) Heatmap of interferon-alpha-related gene expression in hippocampal tissue from male and female AD and AD hiIFN mice, not highlighting a sex-biased transcriptomic profile. (D) Quantification of IFITM3 immunohistochemistry intensity in the cortex, showing higher labeling in AD hiIFN vs. AD mice (**$P < 0.01$, Mann-Whitney test). (E) Quantification of GFAP immunohistochemistry intensity in the cortex, indicating nearly significant increase in astrogliosis in AD hiIFN vs. AD mice ($P = 0.06$, Mann-Whitney test). (F) Quantification of 6E10 immunohistochemistry for Aβ plaque size in the cortex (CTX) and hippocampus (HC), showing reduced plaque size in AD hiIFN vs. AD mice hippocampus (*$P < 0.05$) and no significant changes in cortex (ns, $P > 0.05$) (Mann-Whitney test). (G) Quantification of Thioflavin T staining for fibrillar Aβ deposit size in CTX (ns, $P > 0.05$) and HC (ns, $P > 0.05$), showing no significant changes in AD hiIFN

vs. AD mice (Mann-Whitney test). (H) Quantification of OC staining for β-sheet-rich Aβ
conformation size in CTX (ns, $P > 0.05$) and HC (ns, $P > 0.05$), showing no significant
changes in AD hiIFN vs. AD mice (Mann-Whitney test). Data from 6-month-old AD
hiIFN and APP/PS1 mice. IHC: n = 6 per group (3 females, 3 males). RNA-seq: n = 8 per
group (4 females, 4 males). Data represent mean ± SEM.

**Extended Data Fig. 5. Effects of STING inhibition on BV2 microglial cells and**
**neuropathology in APP/PS1 mice.** (A) Expression of interferon-stimulated genes (*Irf7*,
*Ifitm3*, *Stat2*) and the anti-inflammatory cytokine *Il10* in BV2 microglial cells treated with
vehicle (Veh) or STING inhibitor C-176, with or without poly(I:C) stimulation (ns, $P >$
0.05, \*\*$P < 0.01$, \*\*\*$P < 0.001$, \*\*\*\*$P < 0.0001$, Mann-Whitney test). (B) Heatmap of
interferon-alpha-related gene expression in hippocampal tissue from male and female 6-
month-old APP/PS1 (AD) and C-176-treated APP/PS1 (AD loIFN) mice, highlighting a
higher transcriptomic profile switch in females. (C) GSEA plot of the interferon alpha
response pathway from HC tissue of treated vs. untreated female AD mice, confirming
reduced enrichment of interferon signaling in treated females. (D) Volcano plot of
differential gene expression in the hippocampus of male AD loIFN vs. untreated AD
mice, showing no significant changes in interferon-related genes (gray, non-significant).
(E) Functional enrichment analysis from HC tissueof male AD loIFN vs. untreated AD
mice, showing no significant reduction of the interferon alpha or gamma response
pathways. (F) Quantification of IFITM3 immunohistochemistry intensity in the cortex,
showing lower labeling in untreated AD mice vs. AD loIFN (\*$P < 0.05$, Mann-Whitney
test). (G) Quantification of GFAP immunohistochemistry intensity in the cortex,
indicating nearly significance decrease of astrogliosis in AD loIFN vs. untreated AD mice
($P = 0,09$, Mann-Whitney test). (H) Quantification of WFA-positive structure number in
CA1, showing an increase in AD loIFN vs. untreated AD mice (\*$P > 0.05$, Mann-Whitney
test). (I) Quantification of 6E10 immunohistochemistry for Aβ plaque size in the cortex
(CTX) and hippocampus (HC), showing no significant changes in AD loIFN vs. untreated
AD mice (ns, $P > 0.05$, Mann-Whitney test). (J) Quantification of Thioflavin T staining
for fibrillar Aβ deposit size in CTX and HC, showing no significant changes in AD loIFN
vs. untreated AD mice (ns, $P > 0.05$, Mann-Whitney test). (K) Quantification of OC
staining for β-sheet-rich Aβ conformation size in CTX ($P > 0.05$) and HC (ns, $P > 0.05$),
showing no significant changes in plaques size in AD loIFN vs. untreated AD mice
(Mann-Whitney test). Data from BV2 cell cultures and 6-month-old AD loIFN and
APP/PS1 mice. qPCR: n = 4 per group. IHC: n = 6 per group (3 females, 3 males). RNA-
seq: n = 8 AD (females), 4 AD loIFN (females). Data represent mean ± SEM.

**Extended Data Fig. 6. Behavioral tasks with no significant differences by sex or**
**STING inhibition in APP/PS1 mice.** (A) Percentage of distance traveled in the center
zone of the open field test in 6-month-old male and female WT and APP/PS1 (AD) mice,
showing reduced center exploration in AD females compared to males and WT controls
($P = 0.05$, two-way ANOVA with Tukey's post hoc test). (B) Percentage of time spent
exploring in the T-maze test, showing no significant differences between sexes or
genotypes (\*\*\*$P < 0.001$, \*\*\*\*$P < 0.0001$, two-way ANOVA with Tukey's post hoc

test.) (C) Percentage of freezing during the contextual phase of fear conditioning, showing no significant differences across groups (ns, $P > 0.05$, two-way ANOVA with Tukey's post hoc test). (D) Percentage of freezing during the auditory cue in cued fear conditioning, also showing no significant differences (ns, $P > 0.05$, two-way ANOVA with Tukey's post hoc test). (E) Percentage of distance traveled in the center zone of the open field test in female AD mice treated with vehicle or C-176, showing a non-significant trend toward increased center exploration in C-176–treated mice ($P = 0.11$, two-way ANOVA with Tukey's post hoc test). (F) Percentage of time spent exploring in the T-maze test in female AD mice, showing no difference between vehicle- and C-176–treated groups (ns, $P > 0.05$, two-way ANOVA with Tukey's post hoc test). (G) Percentage of freezing during contextual fear conditioning in female AD mice, showing no effect of C-176 treatment (ns, $P > 0.05$, two-way ANOVA with Tukey's post hoc test). (H) Percentage of freezing during cued fear conditioning in female AD mice, showing no effect of C-176 treatment (ns, $P > 0.05$, two-way ANOVA with Tukey's post hoc test). All behavioral tests were performed in 6-month-old APP/PS1 mice. n = 5 (male WT), 6 (male AD), 5 (female WT), 5 (female AD), 5 (female WT + vehicle), 5 (female WT + C-176), 5 (female AD + vehicle), and 5 (female AD + C-176). Data represent mean ± SEM.

**Supplemental Figure 1**

**Supplemental Figure 2**

**Supplemental Figure 3**

**Supplemental Figure 5**

**Supplemental Figure 6**