INDITEK-2.0: A Bayesian inverse eco-evolutionary modelling framework for reconstructing Phanerozoic biodiversity

Pedro Cermeno[1,*] (orcid.org/0000-0002-3902-3475), Carmen García-Comas[1] (orcid.org/0000-0001-8054-3918), Gloria Herrero Gascón[1](orcid.org/0009-0007-1089-5230), Michael J. Benton[2] (orcid.org/0000-0002-4323-1824), Thomas Bodin[1] (orcid.org/0000-0002-7393-6112), Khushboo Gurung[3] (orcid.org/0000-0001-8074-7339), Benjamin Mills[3] (orcid.org/0000-0002-9141-0931), Alexandre Pohl[4] (orcid.org/0000-0003-2328-351X)

[1]Institut de Ciències del Mar, Consejo Superior de Investigaciones Científicas, Barcelona, 08003, Spain
[2]School of Earth Sciences, University of Bristol, Bristol BS8 1RL, UK
[3]School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK
[4]Biogéosciences, UMR 6282 CNRS, Université de Bourgogne, 6 Boulevard Gabriel, 21000 Dijon, France.

*Corresponding author: Pedro Cermeno (pedrocermeno@icm.csic.es)

Running headline: Reconstructing biodiversity's lost history

Acknowledgements

This work was funded by research grant PID2023-152076NB-I00 from the Spanish government. We thank Lewis Jones and Thiago Rangel for comments on an early version of the manuscript. T.B. acknowledges the support from the Spanish Ministry of Science and Innovation ATR2023-143517/AEI/10.13039/501100011033. A.P. acknowledges the support of the French *Agence Nationale de la Recherche* (ANR) under reference ANR-22-CE01-0003 (project ECO-BOOST). Calculations were performed using HPC resources from DNUM CCUB (*Centre de Calcul de l'Université de Bourgogne*) and the SMART HPC infrastructure at the Institut de Ciències del Mar. P.C., C.G.C, G.H.G. and T.B. acknowledge the 'Severo Ochoa Centre of Excellence' accreditation (CEX2019-000928-S).

Data and code availability statement

The Python code for the INDITEK-2.0 eco-evolutionary model and the MCMC algorithm is available on GitHub: https://github.com/INDITEK-eco-evo-Model/indicios2.0

Conflict of interest statement

The authors have no concflict of interest to declare.

Author contributions

P.C. and C.G.C. conceived and designed the research and wrote the manuscript. P.C. and C.G.C developed the INDITEK-2.0 eco-evolutionary model. G.H.G. conducted model simulations. K.G. and B.M. performed the SCION and FLORA simulations. A.P. performed the FOAM and cGENIE simulations. C.G.C, G.H.G., and T.B. implemented the Bayesian inverse modelling framework. All authors contributed to the development of the model and revised the final manuscript.

Abstract

1. Mechanistic eco-evolutionary models are powerful tools for understanding how life diversified over geological timescales, yet estimating their parameters remains a key challenge. The high computational requirements of these models make probabilistic inversion methods unfeasible.

2. We address this challenge with INDITEK-2.0, a new, cost-efficient eco-evolutionary model that integrates a Bayesian inverse modelling framework for probabilistic estimation of parameters.

3. To validate this framework, we conducted a proof-of-concept study. We first generated synthetic biodiversity data for marine invertebrates using INDITEK-2.0 with known parameter values. To mimic field data, some realistic random Gaussian errors were added to the data. We then used our Bayesian inverse modelling framework to recover these original (true) parameter values from the dataset of current biodiversity distributions after 500 million years of diversification. Our solution is a probability distribution on the parameter space, and we show that the method successfully recovered the target parameter values with a relatively low degree of uncertainity.

4. INDITEK-2.0's ability to probabilistically infer underlying eco-evolutionary processes provides a new modelling tool for reconstructing the evolutionary history of biodiversity across Earth's ecosystems and taxonomic groups.

Keywords: Mechanistic eco-evolutionary modelling, biodiversity history, Bayesian inference, inverse modelling, parameter estimation.

1. Introduction

Mechanistic modelling is emerging as a powerful complementary tool to the fossil record and molecular phylogenetics for reconstructing biodiversity history (Beaugrand, 2023; Hagen, 2023; Lovelace et al., 2020; Rangel et al., 2018). By simulating eco-evolutionary processes under dynamic environmental conditions, mechanistic eco-evolutionary models offer a unique opportunity to explore the drivers and patterns of taxonomic diversification, bridging gaps and testing hypotheses that are difficult to address with traditional methods alone. They notably permit to go beyond the observation of temporal correlations between environmental change and biodiversity dynamics and, through numerical experiments, to address causality and identify the mechanisms underlying observed changes. Mechanistic eco-evolutionary models have been developed to date at two levels of biological organization, at the individual level and the population level. Individual-based eco-evolutionary models provide a framework for simulating complex ecological and evolutionary processes, with trait variation and demographic stochasticity modelled at the level of individual organisms, each represented as a unique entity with its own set of traits (e.g., size, age, genotype, phenotype) (Garwood et al., 2019; Malchow et al., 2021; Xuereb et al., 2021). By incorporating genetic and phenotypic trait variation, individual-based eco-evolutionary models can predict how populations respond to environmental changes, including habitat fragmentation, climate change, or the introduction of invasive species. Population-based models, on the other hand, abstract the details of individual organisms into population or species-level traits that influence species' survival, reproduction, range-shifts, and interaction with the environment and other organisms (Beaugrand, 2023; Hagen et al., 2021; Rangel et al., 2018). The distribution and frequency of these traits within the population can change over time through natural selection, genetic drift, and dispersal, reflecting the evolutionary processes of speciation, extirpation, and extinction at play.

To capture the details of eco-evolutionary processes, individual- and population-based eco-

evolutionary models must operate at sufficiently high spatial and temporal resolutions. High-

resolution modelling represents habitat heterogeneity and species variability more realistically than

coarser models. However, this level of detail creates computational challenges when reconstructing

evolutionary histories over vast spatial and temporal scales. This is a problem for both the Earth

system models that provide palaeoenvironmental conditions and the eco-evolutionary models

themselves. Likewise, in a mechanistic eco-evolutionary model, each discrete entity or state

variable, whether an individual organism or a population of organisms, is governed by its own set of

differential equations that describe its state changes over time. These equations account for growth,

reproduction, death, migration, interactions with other entities and the environment, genetic drift,

heritability, speciation, and extinction. The need to solve a separate set of equations for each

discrete entity rapidly escalates the computational demands of these models.

Beyond current computational limitations, estimating eco-evolutionary model parameters

(evolutionary rates, environmental dependencies, carrying capacities) requires detailed knowledge

of the very biodiversity history we aim to reconstruct, which creates a circularity problem. Inverse

modelling offers a solution: it compares model simulations with empirical data (e.g. present-day

biodiversity, fossil data). By finding the simulation that best fits the empirical data, inverse

modelling can effectively estimate the parameters that most likely generated the observed patterns

(Tarantola, 2005). Probabilistic inference, particularly Bayesian methods employing Markov chain

Monte Carlo (MCMC) algorithms, provides a robust methodological framework to accomplish this.

MCMC algorithms explore the multidimensional parameter space by taking random walks guided

by the shape of the posterior distribution. By recording the visited locations within the parameter

space, MCMC builds a representative sample of the posterior distribution. While effective, this

method requires thousands of model simulations for adequate parameter space exploration, adding a

significant layer of computational requirements to these already demanding models. Consequently,

applying Bayesian inversion to these models requires a trade-off between model complexity and computational feasibility.

To address this trade-off, Cermeno et al. developed INDITEK, a cost-efficient eco-evolutionary model that simplifies mechanisms while still capturing essential broad-scale, macroevolutionary dynamics (Cermeño et al., 2022). Previous models use a bottom-up approach, modelling biodiversity by individually tracking individuals or species. In contrast, INDITEK utilizes a top-down approach, treating biodiversity as an aggregate property at a taxonomic level greater than the species level (e.g., class, phylum). This methodological shift moves the analytical focus from individual- or species-level traits to broader attributes at class or phylum level, making biodiversity itself—rather than individuals or species—the state variable. This strategic approach overcomes major computational hurdles by implicitly modelling the phylogenetic divversification process, allowing the simulation of over 500 million years of global marine invertebrate diversity in under thirty seconds on a standard computer. This article presents INDITEK-2.0, an upgraded version of the INDITEK model. INDITEK-2.0 integrates a Bayesian inverse modelling framework, which enables probabilistic estimation of its parameters. We first conduct a proof-of-concept study to assess the framework's reliability using synthetic community data. We then evaluate its applicability to macroevolutionary questions, concluding with a discussion of model limitations and future directions.

2. INDITEK-2.0 theoretical framework

2.1. Parameterizing evolutionary rates

As noted, in our model, biodiversity is an aggregate property of the community implicitly reflecting the cumulative effects of ecological and evolutionary processes at or below the species level. These processes give rise to spatial and temporal differences in diversification dynamics, whose

parameterization is critical to reconstructing and understanding the evolutionary history of

biodiversity.

The metabolic theory of ecology provides a theoretical basis, with substantial empirical support, for

understanding how temperature affects evolutionary rates by altering the energy available for

growth, reproduction, and survival (Allen et al., 2002; Brown et al., 2004). A greater energy input

to the system (or a higher ambient temperature) implies faster metabolic rates and, consequently,

shorter generation times (Rohde, 1992). Higher turnover rates allow for quicker accumulation of

genetic changes, potentially increasing the rate of evolution (Lanfear et al., 2010). This means that

adaptive traits can become prevalent in shorter time spans, allowing populations to respond to

selection pressures more rapidly (**Fig. 1**). The metabolic theory of ecology uses the Boltzmann

factor, a concept from statistical thermodynamics, to explain how the rates of biological processes,

like metabolism, growth and reproduction accelerate with temperature (Gillooly et al., 2002). Its

core idea – that probabilities can exponentially change with energy and temperature – can be

applied to parameterize how temperature affects speciation rate through its impacts on biological

rates. Alternatively, another quantitative expression for parameterising the sensitivity of metabolic

rates to temperature changes is the Q10 coefficient, a dimensionless parameter taken from the field

of ecophysiology (Mundim et al., 2020). The simplicity of the Q10 model – offering a numerical

value to represent how metabolic rates change at a different rate with every 10°C change – makes it

particularly suitable to eco-evolutionary modelling (**Box 1**).

Fig. 1. Parameterization of evolutionary rates. Speciation rate (λ) is a positive function of temperature (T), food availability (F), and, in terrestrial systems, water availability (not shown). A reduction in food supply rate (-ΔF) directly lowers the carrying capacity (K), implicitly driving an increase in the extinction rate (μ) in systems near K. The difference between λ and μ determines the net diversification rate, ρ_{potential}. The actual, or realized, net diversification rate (ρ_{realized}), is calculated by multiplying ρ_{potential} by the diversity-dependent limitation term (1-d/K). Diversity-dependent limitations feedback on evolutionary rates by implicitly decreasing λ, increasing μ, or both as d approaches K (dotted arrows) .

Our parameterization allows species to evolve their thermal tolerance range—the limits that influence their speciation rate—to track the prevailing climate conditions across each geological time interval (**Box 1**). For example, a temperature that is considered an extreme heat event in the Ordovician period might be a normal temperature in the much warmer Eocene epoch. This approach prevents the model from assuming a universal thermal tolerance for organisms across the entire Phanerozoic, which would be biologically unrealistic. It ensures that the influence of temperature on speciation rate is assessed based on deviations from the norm of that time, thus providing a more accurate and biologically meaningful representation of the relationship between temperature and speciation rate.

Food supply also plays a pivotal role in controlling speciation rates by affecting population dynamics and energetics (Bambach, 1993; Martin, 1996; Vermeij, 1977). For example, abundant food resources can support larger populations, which tend to maintain higher genetic diversity (Hague & Routman, 2016). Larger, genetically diverse populations provide a broader repertoire for natural selection to act upon, potentially accelerating the speciation process (**Fig. 1**). Similarly, for terrestrial ecosystems, water availability, primarily through precipitation, is a fundamental control on net primary productivity (NPP), which in turn determines the available food supply. As a key component of photosynthesis, water is a limiting resource in many ecosystems, directly influencing the amount of biomass produced and thus acting as a primary driver of speciation rates. As with temperature, quantifying the combined effect of food and water availability on speciation rate

necessitates the use of generalized mathematical models. These models, ranging from simple linear relationships to more complex nonlinear dynamics, provide a quantitative framework for representing the mechanisms that govern this relationship (**Box 1**). For example, a Michaelis-Menten equation provides a useful quantitative model for describing the dependence of speciation rates on both food and water availability. This approach is analogous to how chemical reaction rates depend on substrate concentration in biochemistry. It can capture the threshold-like behaviour of resource-driven productivity, where speciation rates increase rapidly with increasing resources in limited environments but plateau in resource-rich conditions where other factors become limiting.

Parameterizing the rate of extinction is considerably more difficult because there is no clear, well-established theoretical framework that comprehensively explains the mechanistic drivers behind it. Adaptation is a fundamental evolutionary process allowing species to survive in changing environments. However, significant changes in environmental conditions can outpace a species' ability to adapt, resulting in a mismatch between its physiological requirements and the new environmental conditions (Hallam, 1998; Peters & Foote, 2002; Saupe et al., 2014). When these changes occur rapidly, species may not have enough time to evolve new traits and cope with the new conditions. For example, it has been hypothesized that the latitudinal biodiversity gradient, characterized by the increase in species richness towards the equator, might result from greater environmental instability at higher latitudes (Chown & Gaston, 2000; Willig et al., 2003). This reasoning posits a direct linkage between the rate of environmental change and the rate of species extinction (Kaiho, 2022; Song et al., 2021). This parameterization requires, however, higher spatial and temporal resolution environmental data than is currently available (see ***Section 4. Model limitations and future directions***). While future Earth system model simulations may provide the necessary inputs, here, we assume a constant background extinction rate for the current model.

2.2. Diversity dependence of net diversification rate

Simulating the effect of ecological interactions, such as competition or predation on diversification

rates requires accounting for species interactions, which is not possible in a model operating above

the species level. One way to incorporate the effect of biotic factors without getting into the

complexities of species interactions is by using the concept of carrying capacity (K) for species

richness (Storch & Okie, 2019). The parameter K is intricately tied to the balance between the

available resources in the system and the species that depend on these resources for survival. On

evolutionary time scales, the dynamic interplay among resource availability and the strategies

species employ to exploit and compete for these resources establishes K, the maximum number of

taxonomic units an ecosystem can support (Bambach, 1983; Mondal & Harries, 2016). In the

context of evolutionary models, K can be used to approximate the effect of ecological interactions

(or biotic factors) on evolutionary rates. As diversity (d) approaches K, the strengthening of

ecological interactions slows down the diversification process until equilibrium diversity is reached

(Levinton, 1979; MacArthur & Wilson, 1967). These diversity-dependent ecological interactions

feedback on evolutionary rates by implicitly decreasing λ, increasing μ, or both (Rabosky, 2009;

Rineau et al., 2022) (**Fig. 1**).

In our model, the K of an ecosystem is directly tied to its net primary production (NPP), which we

use as a proxy for food supply. We reconstructed marine and terrestrial NPP coupling a long-term

carbon cycle model (SCION) (Mills et al., 2021) with an Earth system model of intermediate

complexity (cGENIE) (Ridgwell et al., 2007) (see **Supporting information**) and established a

global minimum K (K_{min}) and maximum K (K_{max}) value for each realm (marine and terrestrial). The

actual K for a specific location and time was then calculated as a linear function of the NPP at that

specific point, with a range that encompassed the values between K_{min} (lowest NPP) and K_{max}

(highest NPP) (see **BOX 1** for details). This approach hinges on the idea that NPP sets a

fundamental limit on the available chemical energy (or food) to support consumer populations

(Cárdenas & Harries, 2010; Martin, 1996). Accordingly, when a decrease in NPP occur due to

environmental shifts, the ecosystem's K is consequently lowered. In response to this resource

limitation, biodiversity is reduced mimicking a local extinction event to align with the new, lower K

(**Fig. 1**). In this way, and according to theory, our model links biodiversity directly to the dynamics

of energy input to ecosystems (Coelho et al., 2025; Hawkins et al., 2003; Wright, 1983). Ultimately,

K at a given location and time is determined by both NPP and the values of K_{min} and K_{max}. Yet,

these latter values should not be static as in the current model; rather, they are expected to change

over geological time, driven by evolutionary innovation (see *4. Caveats and future directions*).

Box 1. INDITEK-2.0's model parameterization

To reconstruct biodiversity, we employ a logistic differential equation that describes the change in local diversity over time:

$$\frac{\partial d}{\partial t} = (\lambda - \mu)\left(1 - \frac{d}{K}\right) \tag{1}$$

where d is diversity, t is time, λ is the speciation rate, μ is the background extinction rate, and K is the system's carrying capacity. K varies as a linear function of the net primary production (NPP), ranging between K_{min} and K_{max} corresponding to the 1 and 99 percentiles of the NPP throughout the Phanerozoic.

We model λ according to the following equation:

$$\lambda = \lambda_{max} - (\lambda_{max} - \lambda_{min})\,(1 - Q_{temp}\,Q_{food}), \tag{2}$$

where λ_{min} and λ_{max} are the range of the maximum potential values of λ. Specifically, λ_{max} is the maximum potential speciation rate under optimal environmental conditions. Q_{temp} and Q_{food} are dimensionless factors, ranging from 0 (maximum limitation) to 1 (no limitation), that modulate λ as a function of temperature and NPP or food supply, respectively.

Q_{temp} is calculated using the equation:

$$Q_{temp} = \frac{Q10^{T-Tmin}}{Q10^{Tmax-Tmin}} \tag{3}$$

where Q10 is a coefficient that quantifies the sensitivity of λ to temperature, T is the sea surface temperature (°C) at a specific location and time, and T_{min} and T_{max} represent the 1 and 99 percentiles, respectively, of the T frequency distribution within each time interval.

Q_{food} is modelled using a Michaelis-Menten formulation:

$$Q_{food} = \frac{F}{(k_{food} + F)} \tag{4}$$

where F (g C m^{-2} yr^{-1}) is marine export production, that is, the fraction of net primary production (NPP) that escapes the sunlit surface layer, at a given location and time. The parameter k_{food} (g C m^{-2} yr^{-1}) is the value of F at which λ is half of its λ_{max}. The same type of Michaelis-Menten parameterization can be used for modelling water availability in terrestrial systems as its effect on λ is similarly dependent on its influence on NPP.

In this study, the background extinction rate (μ) is a fixed value that represents the consistent, low-level rate of diversity loss that occurs even in an ideal environment. It is a fundamental component of the potential net diversification rate ($\rho_{potential}$), defined as:

$$\rho_{potential} = \lambda - \mu \tag{5}$$

However, this potential rate is not always realized. The realized net diversification rate ($\rho_{realized}$), which is what is actually observed, is constrained by ecological interactions that intensify as d approaches the system's K:

$$\rho_{realized} = \rho_{potential} \left(1 - \frac{d}{K}\right) \tag{6}$$

As d nears K, $\rho_{realized}$ declines toward zero, a reduction implicitly caused by a decrease in λ, an increase in μ, or a combination of both (**Fig. 1**). Crucially, K is not a fixed value; it is directly linked to NPP (or food availability). Consequently, a decrease in NPP due to environmental shifts lowers the system's K. This resource limitation triggers a local extinction to align the value of d with the new, reduced K.

Finally, mass extinctions are simulated by imposing negative net diversification rates. Genus-level marine invertebrate fossil diversity curves are used to determine the percentage of diversity loss, starting time, and duration of each mass extinction (Alroy, 2010; Sepkoski, 2002; Zaffos et al., 2017). To obtain a more robust estimate, independent of biases in individual databases, fossil diversity curves from the various sources are averaged into a single dataset.

The Phyton code for the INDITEK-2.0 model is available at:

https://github.com/INDITEK-eco-evo-Model/indicios2.0

2.3. Tracking diversification in space and time

Plate tectonics, through the fragmentation and coalescence of continental land masses, profoundly influenced the spatial patterns of biodiversity by creating geographic barriers to species dispersal and spurring provinciality (Pellissier et al., 2018; Valentine et al., 1978; Valentine & Moores, 1970; Zaffos et al., 2017). The shifting of continents and oceans also moved marine and terrestrial habitats across bioclimatic regions, further influencing diversification dynamics. However, previous eco-evolutionary models lack the capability to continuously track the environmental history and biodiversity of habitats as they moved with plates, precluding us from establishing a quantitative, mechanistic link between plate tectonics, global climate evolution, and macroevolutionary dynamics. To mechanistically account for the interplay of plate tectonics and climate on macroevolutionary dynamics, INDITEK implemented a Lagrangian approach (Cermeno et al. 2022). This approach allows the model to track habitats—represented as discrete palaeogeographic points by the plate tectonic model—as they move across Earth's surface. In doing so, it accounts for the contribution of continuously-evolving environmental conditions and the legacy of past biodiversity changes. A Lagrangian approach contrasts with the simulation of biodiversity 'snapshots' (Pohl et al., 2023) and provides a more realistic view of how biodiversity evolved in space and time, capturing the continuous nature of macroevolutionary processes. Currently, integrated modelling frameworks are used to reconstruct Earth's palaeoenvironmental conditions. For example, plate tectonic and landscape evolution models provide detailed insights into the planet's palaeogeographic and physiographic evolution (Kocsis & Scotese, 2021; Merdith et al., 2021; Salles et al., 2023; Scotese & Wright, 2018; Vérard et al., 2015). Likewise, coupled Earth system models and ocean-atmosphere general circulation models reconstruct palaeoclimatic and palaeoenvironmental conditions (Gurung et al., 2024; Mills et al., 2021; Ridgwell et al., 2007; Scotese, 2021). By simulating key environmental variables, such as air/sea temperature,

precipitation, and primary production in land and ocean, these models collectively provide the

essential context for the unfolding of eco-evolutionary processes (**Fig. 2a**).

Fig. 2. A schematic of the Bayesian inverse modelling framework. (a) Global reconstructions of key environmental variables for two distinct time points: 440 Ma (million years ago) and 80 Ma. The top panels show reconstructed air-sea temperatures, while the bottom panels display marine export production (EP) and land net primary production (NPP). The black stepped outlines show the positions of continents. (b) Flow diagram of the Bayesian inverse modelling process. The method uses a Metropolis-Hastings Markov chain Monte Carlo (MCMC) algorithm to estimate model parameters by iteratively comparing simulated biodiversity to a calibration dataset. The residual sum of squares (RSS), the likelihood function, quantifies the discrepancy between the modelled and observed biodiversity (or calibration) data, allowing the algorithm to find the best-fitting parameter values.

2.4. Bayesian inverse eco-evolutionary modelling

INDITEK-2.0 uses a Bayesian inverse modelling framework with a Metropolis-Hastings Markov chain Monte Carlo (MCMC) algorithm to efficiently explore the parameter space (Hastings, 1970). The MCMC algorithm is used to generate a series of samples from the posterior probability distribution of the model parameters, which allows us to find the most probable parameter values and quantify their uncertainty. In each iteration, the algorithm proposes a new parameter set by randomly perturbing the current set. It then calculates the Residual Sum of Squares (RSS) as a measure of fit to compare the model's simulated biodiversity and the calibration data (e.g., present-day and fossil biodiversity data) (**Fig. 2b**). New parameter sets are accepted if they reduce the RSS or according to a probabilistic criterion. This criterion allows for the occasional acceptance of less-optimal sets to avoid getting stuck in a local minimum. This iterative process creates a random walk through the parameter space that eventually converges on the posterior distribution, which is the probabilistic solution. After a sufficient "burn-in" period, the collected samples of parameter values approximate this distribution, providing a comprehensive representation of the probability of different parameter values given the observational data and our prior assumptions. This approach ensures that our parameter estimates are not single point values but a range of plausible values, reflecting the uncertainties inherent in the empirical data and the model processes. The model's parallelized architecture allows multiple independent chains to run simultaneously, which significantly increases the coverage of the sampled parameter space without requiring additional computational time.

2.5. Proof-of-concept design

To assess our approach and demonstrate its resolving potential, we performed a synthetic test, or proof-of-concept. First, we selected the target parameter values based on the results of Cermeño et al. (2022) for the continental shelves (**Table 1**). Due to a lack of prior information, we assigned uniform priors with relatively broad range limits to all parameters except for the Q10 parameter; for

290 this parameter, we used a Gaussian distribution, as ample literature for marine invertebrates reports

291 a constrained range of values (1–4), with a higher probability around the mean (1.5–2.5)

292 (Brockington & Clarke, 2001; R. E. Rangel & Sorte, 2022).

Table 1. INDITEK-2.0 model parameters: prior distributions and posterior results.

Parameter	Units	True value	Prior distribution	Posterior Mean	95% CI
K_{min}	#genera	19	Uniform (1, 150)	18.81	[18.19, 19.6]
K_{max}	#genera	161	Uniform (80, 1000)	162	[154.63, 170.63]
λ_{min}	Myr^{-1}	0.001	Uniform (0.0001, 0.05)	0.0019	[0.0005, 0.0028]
λ_{max}	Myr^{-1}	0.035	Uniform (0.001, ∞)	0.0349	[0.0344, 0.0352
μ	Myr^{-1}	0.001	n/a	n/a	n/a
Q10	dimensionless	1.75	Gaussian (1 – 4)	1.731	[1.621, 1.833]
k_{food}	dimensionless	1.5	n/a	n/a	n/a

With these parameter values, we ran the model to generate a synthetic dataset representing present-

day biodiversity distributions on a 0.5° x 0.5° spatial grid. In order to mimic observational and

theoretical errors and to enable efficient parameter exploration with the Metropolis–Hastings

MCMC algorithm, we added a random Gaussian error with a standard deviation of 2 genera to the

synthetic dataset. We then applied our inversion approach to this dataset. Six Markov chains were

used to explore the 5-dimensional parameter space and to sample the posterior probability

distribution. Each chain was initialized within a range of ±20% of the reference parameter values,

with starting points selected using a Latin hypercube sampling strategy to ensure broad coverage of

the parameter space. At each iteration, one forward simulation was performed, and the resulting

output was compared to the synthetic dataset. Model fit was quantified using the residual sum of

squares (RSS), calculated as the squared difference between the simulated and synthetic data,

normalized by the variance of the added noise.

3. Results

The MCMC chains for all five parameters: K_{max}, K_{min}, λ_{max}, λ_{min}, and Q10 converged toward a stable

mean over the 2000 interative simulations after successful exploration of the posterior distribution

(**Fig. 3**). The rate of convergence varied among parameters. K_{max}, K_{min}, and λ_{max} showed relatively

rapid convergence, with chains stabilizing within 500 to 750 iterations. In contrast, Q10 and

particularly λ_{min} showed slower convergence, with some chains for λ_{min} taking nearly 1500 iterations

to mix around the original, true value. The convergence of all chains confirms the stability and

reliability of the parameter estimates.

Fig. 3. Markov chain trajectories for parameter inference. *Evolution of five model parameters*

(K_{max}, K_{min}, λ_{max}, λ_{min}, and Q10) over 2000 iterations for each of the six independent Markov chains

(labeled 1 through 6). The horizontal dashed red line indicates the true value for each parameter,

while the horizontal dashed blue line indicates the mean value of the posterior distribution after the

burn-in period (from iteration 200 onward).

Our analysis further confirms the accuracy of the model, as the true parameter values were always

contained within the 65% credible interval provided by the posterior solution (**Table 1**, **Fig. 4**). This

demonstrates that the MCMC method not only found a stable solution but also accurately estimated

the range of plausible parameter values, reliably capturing the true underlying values within this

interval. The two-dimensional joint posterior distributions correctly identified the relationships

between parameter pairs (**Fig. 4**). The bivariate correlations among the model parameters offered

key insights into the structure of the parameter space and potential interdependencies. A strong

positive correlation was observed between λ_{min} and Q10, suggesting that the MCMC algorithm

compensated for a lower λ_{min} by exploring higher values for Q10. Conversely, K_{max} and K_{min}, the

upper and lower bounds of the carrying capacity showed, respectively, a negative and positive

correlation with λ_{max}. This might indicate a trade-off in how the model fits the synthetic biodiversity

data. Specifically, to explain biodiversity patterns with a very high λ_{max}, the model must constrain

K_{max} to prevent a runaway increase that would overshoot the data. In contrast, a high λ_{max} is

positively linked to K_{min}, indicating that rapid diversity growth is necessary to explain a high

baseline level of biodiversity. In essence, the model balances a high-speed diversification process

against both K_{max} and K_{min} to reproduce the synthetic biodiversity data. Other parameter pairs

showed weak or negligible correlation, as indicated by their generally more circular joint posterior

distributions, suggesting that the MCMC algorithm explored these parameters with a relatively high

degree of independence (**Fig. 4**).

Fig. 4. Validation of the INDITEK-2.0's Bayesian inverse modelling framework. *The diagonal*

panels show the one-dimensional posterior distributions for each of the five inferred parameters

(K_{max}, K_{min}, λ_{max}, λ_{min}, and Q10), represented as histograms of all pooled MCMC samples after the

burn-in period. The dashed red line indicates the true value for each parameter, while the dashed

blue line shows the mean value of the posterior distribution after the burn-in. The off-diagonal

panels display the joint posterior distributions and correlations between parameter pairs. The

colour gradient denotes the density of MCMC samples. The red square included on each heatmap

represents the true value used to generate the synthetic biodiversity data. The location of this red

square relative to the densest region of the heatmap is a key indicator of the framework's success in

recovering the original parameter values.

We found a strong and regionally balanced match between the true, synthetic biodiversity data and

the estimated biodiversity data derived from the mean of the probabilistic solutions (**Fig. 5**),

indicating the model's high performance in parameter estimation. The minimal differences between

the two biodiversity distribution maps do not detract from the model's overall accuracy.

Fig. 5. Marine invertebrate biodiversity data. (a) *Global distribution of marine invertebrate diversity generated using the INDITEK model with predefined (true) parameter values.* **(b)** *Global distribution of marine invertebrate diversity derived from parameters estimated using the INDITEK-2.0's Bayesian inversion method.*

Our approach generated an ensemble of plausible biodiversity histories rather than a single definitive one. Each history, with its assigned probability, allowed us to explicitly quantify the uncertainty in our reconstructions. To illustrate this uncertainty, we tracked the biodiversity of three representative modern regions; the Indo-Pacific, Caribbean, and Mediterranean submerged habitats, throughout the simulations (**Fig. 6**). These regions, now among the most striking marine biodiversity hotspots worldwide, began to develop from the Late Triassic onwards, likely due to the relatively high long-term environmental stability that prevailed during the Mesozoic and Cenozoic eras (Cermeño et al. 2022). Our analysis was designed to test the Bayesian inverse modelling approach, so the variability in our reconstructions was relatively low and mostly stemmed from the error we deliberately introduced to the synthetic data (**Fig. 6**). We anticipate a higher variability in biodiversity reconstructions when model simulations are calibrated with true, field observational data. The analysis of this variability will be crucial in future studies to pinpoint which macroevolutionary patterns and drivers are most robustly supported by real-world evidence.

Fig. 6. Macroevolutionary history of marine biodiversity hotspots. (upper panels) Temporal evolution of marine invertebrate diversity (median number of genera across all tracked palaeogeographic points) from the Late Triassic (~200 Ma) to the present day (0 Ma) for the Indo-Pacific, Caribbean, and Mediterranean regions. Each panel displays multiple histories, corresponding to the different probabilistic solutions from the model. (lower panels) Geographic positions of the tracked palaeogeographic points for each biodiversity hotspot (colour dots) at specific time intervals. Each map shows the continental configuration (dark grey) and continental shelves (light grey) at the indicated time. The orange dashed lines represent the approximate positions of the 25°C isolines (isotherms).

4. Caveats and future directions

Our mechanistic modelling tool provides a virtual window into a profoundly complex evolutionary history. Evolutionary processes are shaped by many factors, many of which are beyond the current possibilities of the model. Recognizing these limitations is crucial to adequately interpreting the model's results and devising future implementations.

i. The extreme environmental changes associated with mass extinctions pose a significant challenge for modelling due to the involvement of multiple stressors, often interacting in nonlinear ways. For example, the Ordovician mass extinction (ca. 445 million years ago) is

thought to have been initiated by global glaciation, which was followed by a rapid warming

period. These climate fluctuations led to rapid changes in sea level and ocean anoxia with

dramatic consequences for marine life (Pohl et al., 2023). Similarly, the end-Permian mass

extinction is thought to have been caused by massive volcanic eruptions, leading to severe

climatic shifts, ocean acidification, and widespread anoxia (Hülse et al., 2021). Modelling mass

extinctions requires sophisticated parameterizations to capture nonlinear dynamics and feedback

among Earth's evolving systems (geosphere, atmosphere, hydrosphere, and biosphere). These

systems are intrinsically interconnected, with changes in one domain often leading to cascading

effects in others (Hull, 2015; Junium et al., 2022; Schobben et al., 2019). The difficulty in

implementing parameterizations that capture these nonlinear dynamics and feedback loops in

palaeo-Earth system models currently poses significant challenges in recreating the

environmental scenarios responsible for mass extinctions.

ii. As discussed in *Section 2.1. Parameterizing evolutionary rates*, adaptation allows species to

persist in changing environments. However, rapid environmental changes can outpace the rate of

evolutionary adaptation, preventing species from developing the necessary traits to cope with

new conditions. This reasoning posits a direct linkage between the rate of environmental change

and the rate of species extinction (Kaiho, 2022; Song et al., 2021). Implementing this

parameterization is currently limited by lack of palaeoenvironmental reconstructions with

sufficient spatial and temporal resolution. Our current model uses environmental inputs from

reconstructions with a temporal resolution of 30 time slices each encompassing 10–20 million-

year and a spatial resolution of 36 x 36 (latitude, longitude) equal area grid. Extinction events,

however, are not uniform across vast geographic areas or over long timescales; instead, they are

often localized and triggered by relatively rapid environmental shifts. As a result, existing

coarse-resolution reconstructions may smooth out these crucial, fine-scale variations, resulting in

an inaccurate estimation of the extinction rate (μ). Therefore, the successful application of this

parameterization is contingent on the availability of high-resolution environmental

reconstructions that can accurately capture the nature of extinction dynamics. Current efforts are

being made to increase these resolutions to every million years and 72 x 72 equal area grid.

iii. Extinction selectivity is influenced by countless biological, ecological, and environmental

variables, often interacting unpredictably. The ecological niches, life history traits, and

geographic distributions of species significantly influence their resilience or vulnerability to

environmental perturbations (Cole & Hopkins, 2021; Payne et al., 2016; Pinsky et al., 2019).

These species-specific attributes cannot be implemented into a model operating above the

species level. However, it is feasible to implement extinction selectivity at a higher taxonomic

level by adjusting specific parameter values associated to a class or phylum known for their

overall sensitivity or resistance to particular stressors. For example, assigning different values to

ectotherms and endotherms for the thermal dependence of speciation rate can reflect their

differential responses to temperature changes. Insects' short generation times, and, in many cases,

ability to go dormant can confer resistance to environmental stressors. Their evolutionary

success could be modelled and parameterized by, for example, lowering their sensitivity to mass

extinctions.

iv. The emergence of innovative evolutionary traits and co-evolutionary relationships has been

pivotal in creating novel ecological opportunities and driving major increases in biodiversity

(Benton et al., 2015; Nitecki, 1990; Vermeij, 2017). At or below the species level, mechanistic

eco-evolutionary models can be designed to simulate how new traits emerge and evolve. These

models can include mechanisms for trait evolution (e.g., trait diffusion), niche construction, and

ecological interactions, allowing them to capture the interplay between organisms and their

environments that drives speciation (Hagen, 2023). To account for the effect of evolutionary

innovations and co-evolutionary relationships on macroevolutionary dynamics, future work

could involve using fossil data and molecular phylogenetic analyses to identify specific time

points that mark key innovations or the start of co-evolutionary histories. We could then estimate model parameters for the periods before and after these critical time points, allowing the eco-evolutionary model to represent discrete, biologically-justified shifts in evolutionary dynamics and carrying capacities. For example, evolutionary innovation can expand the ecospace and thus the amount of diversity supported per unit resource supply, which is parameterized as an increase in K_{min} and K_{max}.

5. Conclusions

Our study presents INDITEK-2.0, a cost-efficient mechanistic eco-evolutionary model designed to explore the deep-time history of biodiversity. Initially tested on synthetic communities of marine invertebrates, the model is extensible to other marine and terrestrial animal groups, microorganisms and plants, provided that 1) comprehensive and reliable calibration datasets are available, and 2) the model's parameter-dependent processes realistically represent their evolutionary dynamics.

INDITEK-2.0 is built on two innovative approaches that allow us to trace the ancient origins of today's biodiversity patterns. First, unlike traditional models that use a fixed geographic grid, INDITEK-2.0 employs a Lagrangian approach. This method tracks specific parcels of the Earth's surface as they move over millions of years due to plate tectonics. This dynamic tracking is essential for accurately linking modern regions of interest, such as biodiversity hotspots, to their ancient, non-stationary geographic locations and environmental conditions. Second, INDITEK-2.0 utilizes a Bayesian inverse modelling framework to probabilistically estimate model parameters, thereby generating hundreds to thousands of plausible biodiversity histories, each with a statistically quantified probability. By integrating these two methods, INDITEK-2.0 offers a powerful and novel way to investigate the link between Earth's surface dynamics and biodiversity history.

INDITEK-2.0 allows us to target a specific taxonomic group and reconstruct its biodiversity history by iteratively refining model outputs based on comparisons with both present-day biodiversity and

fossil data. While probabilistic methods have long been used to detect diversification rate shifts

from time-calibrated molecular phylogenies (Bouckaert et al., 2019; Morlon et al., 2024; Rabosky,

2014; Stadler, 2011), applying these methods to spatially-explicit mechanistic eco-evolutionary

models offers the ability to glean novel insights into the mechanisms driving these shifts. This is a

key distinction. By interrogating the model, we can directly investigate how, for example,

continental drift and global climate evolution influenced macroevolutionary dynamics, providing an

explicit and realistic representation of the environmental forces that shaped biodiversity over

millions of years. Ultimately, integrating mechanistic eco-evolutionary models with probabilistic

parameter estimation provides a robust, uncertainty-aware framework for reconstructing

biodiversity's lost history. This approach represents a significant step toward uncovering the origins

of the astonishing biodiversity we see today.

Supporting information

Earth system model simulations

This section describes the methods used to simulate the Late Ordovician and Late Cretaceous

palaeoenvironmental conditions shown in the figure 2 of the manuscript. To date, no numerical

model offers a coupled representation of climate-carbon-cycle equilibrium, primary productivity on

land and marine biogeochemistry in a spatially-resolved, 3D ocean – which together constitute the

boundary conditions needed for Phanerozoic-scale mechanistic eco-evolutionary model simulations.

This limitation arises from the contrasting time scales at which the underlying mechanisms occur.

Carbon cycle equilibrium, on the one hand, results from the balance between (silicate, carbonate

and kerogen) weathering on land and (carbonate and organic carbon) burial in the ocean, and

operates at the million-year time scale. Oceanic circulation and biogeochemistry, on the other hand,

reach equilibrium much faster, in a few 10s of kilo-years. As a result of this major difference in the

time constants of the different components of the climate system, the long-term carbon cycle and

the ocean-atmosphere system have traditionally been simulated using very different types of

models. Long-term carbon cycle models can be run for 100s of million years. They can be used to

simulate the evolution of the atmospheric concentration in greenhouse gases (CO_2 in particular) and

oxygen, and changes in oceanic nutrient inventories (Mills et al., 2021). Their rapid integration time

relies on parameterizations and spatial abstraction – generally representing the ocean and

atmosphere as a single box (hence value). Earth system models, on the other hand, offer a spatially-

resolved simulation of climate and ocean biogeochemistry, but their resulting computational cost

usually does not permit to run them for more than a few 10s of kyrs (Pohl et al., 2022).

To overcome this limitation, we here propose to couple a long-term carbon cycle model (SCION)

(Mills et al., 2021) with an Earth system model of intermediate complexity (cGENIE) (Ridgwell et

al., 2007). In detail, we propose to first simulate palaeoclimatic conditions using an ocean-

atmosphere general circulation model (e.g., FOAM) (Jacob, 1997). Palaeoclimatic conditions

simulated at various atmospheric CO_2 levels can then be used to calculate the long-term climate-

carbon-cycle equilibrium using SCION, together with primary productivity on land using the

vegetation module FLORA (Gurung et al., 2022, 2024). Atmospheric CO_2 and O_2 concentrations

and oceanic nutrient (phosphate) inventories simulated in SCION can be used as boundary

conditions in cGENIE, the latter model permitting in turn to simulate ocean temperatures and

biogeochemistry (including marine export production) in a 3D ocean. During all steps, the same

continental reconstructions are used in every model. The final result is a self-consistent set of

palaeoenvironmental conditions both on land and in the ocean – notably temperatures and primary

productivity that are two key drivers of speciation and extinction rates.

References

Allen, A. P., Brown, J. H., & Gillooly, J. F. (2002). Global biodiversity, biochemical kinetics, and the energetic-equivalence rule. *Science*, *297*(5586), 1545–1548. https://doi.org/10.1126/science.1072380

Alroy, J. (2010). The shifting balance of diversity among major marine animal groups. *Science*, *329*(5996), 1191–1194. https://doi.org/10.1126/science.1189910

Bambach, R. K. (1983). Ecospace utilization and guilds in marine communities through the Phanerozoic. *Biotic Interactions in Recent and Fossil Benthic Communities*, 719–746. https://doi.org/10.1007/978-1-4757-0740-3_15

Bambach, R. K. (1993). Seafood through time: Changes in biomass, energetics, and productivity in the marine ecosystem. *Paleobiology*, *3*(N/A), 372–397. https://doi.org/10.1017/S0094837300000336

Beaugrand, G. (2023). Towards an Understanding of Large-Scale Biodiversity Patterns on Land and in the Sea. *Biology*, *12*(3), 339. https://doi.org/10.3390/biology12030339

Benton, M. J., Donoghue, P. C. J., Asher, R. J., Friedman, M., Near, T. J., & Vinther, J. (2015). Constraints on the timescale of animal evolutionary history. In *Palaeontologia Electronica* (Vol. 18, Issue 1). https://doi.org/10.26879/424

Bouckaert, R., Vaughan, T. G., Barido-Sottani, J., Duchêne, S., Fourment, M., Gavryushkina, A., Heled, J., Jones, G., Kühnert, D., De Maio, N., Matschiner, M., Mendes, F. K., Müller, N. F., Ogilvie, H. A., Du Plessis, L., Popinga, A., Rambaut, A., Rasmussen, D., Siveroni, I., … Drummond, A. J. (2019). BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS Computational Biology*, *15*(4). https://doi.org/10.1371/journal.pcbi.1006650

Brockington, S., & Clarke, A. (2001). The relative influence of temperature and food on the metabolism of a marine invertebrate. *Journal of Experimental Marine Biology and Ecology*, *258*(1). https://doi.org/10.1016/S0022-0981(00)00347-6

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology*, *85*(7). https://doi.org/10.1890/03-9000

Cárdenas, A. L., & Harries, P. J. (2010). Effect of nutrient availability on marine origination rates throughout the Phanerozoic eon. *Nature Geoscience*, *3*(6), 430–434. https://doi.org/10.1038/ngeo869

Cermeño, P., García-Comas, C., Pohl, A., Williams, S., Benton, M., Le Gland, G., Muller, R. D., Ridgwell, A., & Vallina, S. (2022). Post-extinction recovery of the Phanerozoic oceans and biodiversity hotspots. *Nature*, *607*, 507–511. 10.1038/s41586-022-04932-6

Chown, S. L., & Gaston, K. J. (2000). Areas cradles and museums: The latitudinal gradient in species richness. In *Trends in Ecology and Evolution* (Vol. 15, Issue 8). https://doi.org/10.1016/S0169-5347(00)01910-8

Coelho, M. T. P., Barreto, E., Diniz-Filho, J. A. F., Rangel, T. F., Bohdalková, E., Storch, D., Jang, Y.-T., Pontarp, M., Cassemiro, F., de Araujo, M. L., Hull, P. M., & Graham, C. H. (2025). Consistent energy-diversity relationships in terrestrial vertebrates. *Science*, *389*(6755), 53–57. https://doi.org/10.1126/science.adu2590

Cole, S. R., & Hopkins, M. J. (2021). Selectivity and the effect of mass extinctions on disparity and functional ecology. *Science Advances*, *7*(19). https://doi.org/10.1126/sciadv.abf4072

Garwood, R. J., Spencer, A. R. T., & Sutton, M. D. (2019). REvoSim: Organism-level simulation of macro and microevolution. *Palaeontology*, *62*(3), 339–355. https://doi.org/10.1111/pala.12420

Gillooly, J. F., Charnov, E. L., West, G. B., Savage, V. M., & Brown, J. H. (2002). Effects of size and temperature on developmental time. *Nature*, *417*(6884), 70–73. https://doi.org/10.1038/417070a

Gurung, K., Field, K. J., Batterman, S. A., Goddéris, Y., Donnadieu, Y., Porada, P., Taylor, L. L., & Mills, B. J. W. (2022). Climate windows of opportunity for plant expansion during the Phanerozoic. *Nature Communications*, *13*(1). https://doi.org/10.1038/s41467-022-32077-7

Gurung, K., Field, K. J., Batterman, S. A., Poulton, S. W., & Mills, B. J. W. (2024). Geographic range of plants drives long-term climate change. *Nature Communications*, *15*(1). https://doi.org/10.1038/s41467-024-46105-1

Hagen, O. (2023). Coupling eco-evolutionary mechanisms with deep-time environmental dynamics to understand biodiversity patterns. *Ecography*, *2023*(4), e06132. https://doi.org/10.1111/ecog.06132

Hagen, O., Flück, B., Fopp, F., Cabral, J. S., Hartig, F., Pontarp, M., Rangel, T. F., & Pellissier, L. (2021). gen3sis: A general engine for eco-evolutionary simulations of the processes that shape Earth's biodiversity. *PLoS Biology*, *19*(7). https://doi.org/10.1371/journal.pbio.3001340

Hague, M. T. J., & Routman, E. J. (2016). Does population size affect genetic diversity? A test with sympatric lizard species. *Heredity*, *116*(1), 92–98. https://doi.org/10.1038/hdy.2015.76

Hallam, A. (1998). Mass extinctions in Phanerozoic time. *Geological Society Special Publication*, *140*. https://doi.org/10.1144/GSL.SP.1998.140.01.18

Hastings, W. K. (1970). Monte carlo sampling methods using Markov chains and their applications. *Biometrika*, *57*(1), 97–109. https://doi.org/10.1093/biomet/57.1.97

Hawkins, B. A., Field, R., Cornell, H. V., Currie, D. J., Guégan, J. F., Kaufman, D. M., Kerr, J. T., Mittelbach, G. G., Oberdorff, T., O'Brien, E. M., Porter, E. E., & Turner, J. R. G. (2003). Energy, water, and broad-scale geographic patterns of species richness. In *Ecology* (Vol. 84, Issue 12). https://doi.org/10.1890/03-8006

Hull, P. (2015). Life in the aftermath of mass extinctions. In *Current Biology* (Vol. 25, Issue 19). https://doi.org/10.1016/j.cub.2015.08.053

Hülse, D., Lau, K. V., van de Velde, S. J., Arndt, S., Meyer, K. M., & Ridgwell, A. (2021). End-Permian marine extinction due to temperature-driven nutrient recycling and euxinia. *Nature Geoscience*, *14*(11). https://doi.org/10.1038/s41561-021-00829-7

Jacob, R. L. (1997). Low frequency variability in a simulated atmosphere-ocean system. In *Dissertation*.

Junium, C. K., Zerkle, A. L., Witts, J. D., Ivany, L. C., Yancey, T. E., Liu, C., & Claire, M. W. (2022). Massive perturbations to atmospheric sulfur in the aftermath of the Chicxulub impact. *Proceedings of the National Academy of Sciences of the United States of America*, *119*(14). https://doi.org/10.1073/pnas.2119194119

Kaiho, K. (2022). Relationship between extinction magnitude and climate change during major marine and terrestrial animal crises. *Biogeosciences*, *19*(14), 3369–3380. https://doi.org/10.5194/bg-19-3369-2022

Kocsis, Á. T., & Scotese, C. R. (2021). Mapping paleocoastlines and continental flooding during the Phanerozoic. In *Earth-Science Reviews* (Vol. 213, p. 103463). Elsevier B.V. https://doi.org/10.1016/j.earscirev.2020.103463

Lanfear, R., Ho, S. Y. W., Love, D., & Bromham, L. (2010). Mutation rate is linked to diversification in birds. *Proceedings of the National Academy of Sciences of the United States of America*, *107*(47), 20423–20428. https://doi.org/10.1073/pnas.1007888107

Levinton, J. S. (1979). A theory of diversity equilibrium and morphological evolution. *Science*, *204*(4390). https://doi.org/10.1126/science.204.4390.335

Lovelace, D. M., Hartman, S. A., Mathewson, P. D., Linzmeier, B. J., & Porter, W. P. (2020). Modeling Dragons: Using linked mechanistic physiological and microclimate models to explore environmental, physiological, and morphological constraints on the early evolution of dinosaurs. *PLoS ONE*, *15*(5). https://doi.org/10.1371/journal.pone.0223872

MacArthur, R. H., & Wilson, E. O. (1967). *The Theory of Island Biogeography.* Princeton University Press.

Malchow, A. K., Bocedi, G., Palmer, S. C. F., Travis, J. M. J., & Zurell, D. (2021). RangeShiftR: an R package for individual-based simulation of spatial eco-evolutionary dynamics and species' responses to environmental changes. *Ecography*, *44*(10), 1443–1452. https://doi.org/10.1111/ecog.05689

Martin, R. E. (1996). Secular increase in nutrient levels through the Phanerozoic: Implications for productivity, biomass, and diversity of the marine biosphere. *Palaios*, *11*(3), 209–219. https://doi.org/10.2307/3515230

Merdith, A. S., Williams, S. E., Collins, A. S., Tetley, M. G., Mulder, J. A., Blades, M. L., Young, A., Armistead, S. E., Cannon, J., Zahirovic, S., & Müller, R. D. (2021). Extending full-plate

tectonic models into deep time: Linking the Neoproterozoic and the Phanerozoic. In *Earth-*
Science Reviews (Vol. 214, p. 103477). https://doi.org/10.1016/j.earscirev.2020.103477

Mills, B. J. W., Donnadieu, Y., & Goddéris, Y. (2021). Spatial continuous integration of Phanerozoic
global biogeochemistry and climate. In *Gondwana Research* (Vol. 100, pp. 73–86).
https://doi.org/10.1016/j.gr.2021.02.011

Mondal, S., & Harries, P. J. (2016). Phanerozoic trends in ecospace utilization: The bivalve
perspective. In *Earth-Science Reviews* (Vol. 152, pp. 106–118).
https://doi.org/10.1016/j.earscirev.2015.10.005

Morlon, H., Andréoletti, J., Barido-Sottani, J., Lambert, S., Perez-Lamarque, B., Quintero, I.,
Senderov, V., & Veron, P. (2024). Phylogenetic Insights into Diversification. *Annual Review of*
Ecology, Evolution, and Systematics, *55*(1), 1–21. https://doi.org/10.1146/annurev-ecolsys-
102722-020508

Mundim, K. C., Baraldi, S., Machado, H. G., & Vieira, F. M. C. (2020). Temperature coefficient
(Q10) and its applications in biological systems: Beyond the Arrhenius theory. In *Ecological*
Modelling (Vol. 431). https://doi.org/10.1016/j.ecolmodel.2020.109127

Nitecki, M. H. (1990). *Evolutionary innovations*. The University of Chicago Press.

Payne, J. L., Bush, A. M., Chang, E. T., Heim, N. A., Knope, M. L., & Pruss, S. B. (2016).
Extinction intensity, selectivity and their combined macroevolutionary influence in the fossil
record. *Biology Letters*, *12*(10). https://doi.org/10.1098/rsbl.2016.0202

Pellissier, L., Heine, C., Rosauer, D. F., & Albouy, C. (2018). Are global hotspots of endemic
richness shaped by plate tectonics? *Biological Journal of the Linnean Society*, *123*(1).
https://doi.org/10.1093/biolinnean/blx125

Peters, S. E., & Foote, M. (2002). Determinants of extinction in the fossil record. *Nature*,
416(6879), 420–424. https://doi.org/10.1038/416420a

Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., & Sunday, J. M. (2019). Greater
vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, *569*(7754).
https://doi.org/10.1038/s41586-019-1132-4

Pohl, A., Ridgwell, A., Stockey, R. G., Thomazo, C., Keane, A., Vennin, E., & Scotese, C. R.
(2022). Continental configuration controls ocean oxygenation during the Phanerozoic. *Nature*,
608(7923), 523–527. https://doi.org/10.1038/s41586-022-05018-z

Pohl, A., Stockey, R. G., Dai, X., Yohler, R., Le Hir, G., Hülse, D., Brayard, A., Finnegan, S., &
Ridgwell, A. (2023). Why the Early Paleozoic was intrinsically prone to marine extinction.
Science Advances, *9*(35). https://doi.org/10.1126/sciadv.adg7679

Rabosky, D. L. (2009). Ecological limits and diversification rate: Alternative paradigms to explain
the variation in species richness among clades and regions. *Ecology Letters*, *12*(8).
https://doi.org/10.1111/j.1461-0248.2009.01333.x

Rabosky, D. L. (2014). Automatic detection of key innovations, rate shifts, and diversity-
dependence on phylogenetic trees. *PLoS ONE*, *9*(2).
https://doi.org/10.1371/journal.pone.0089543

Rangel, R. E., & Sorte, C. J. B. (2022). Local-scale thermal history influences metabolic response
of marine invertebrates to warming. *Marine Biology*, *169*(10). https://doi.org/10.1007/s00227-
022-04110-2

Rangel, T. F., Edwards, N. R., Holden, P. B., Diniz-Filho, J. A. F., Gosling, W. D., Coelho, M. T. P.,
Cassemiro, F. A. S., Rahbek, C., & Colwell, R. K. (2018). Modeling the ecology and evolution
of biodiversity: Biogeographical cradles, museums, and graves. *Science*, *361*(6399).
https://doi.org/10.1126/science.aar5452

Ridgwell, A., Hargreaves, J. C., Edwards, N. R., Annan, J. D., Lenton, T. M., Marsh, R., Yool, A., &
Watson, A. (2007). Marine geochemical data assimilation in an efficient Earth system model of
global biogeochemical cycling. *Biogeosciences*, *4*(1), 87–104. https://doi.org/10.5194/bg-4-87-
2007

Rineau, V., Smyčka, J., & Storch, D. (2022). Diversity dependence is a ubiquitous phenomenon
across Phanerozoic oceans. *Science Advances*, *8*(43). https://doi.org/10.1126/sciadv.add9620

Rohde, K. (1992). Latitudinal Gradients in Species Diversity: The Search for the Primary Cause.
Oikos, *65*(3), 514. https://doi.org/10.2307/3545569

Salles, T., Husson, L., Lorcery, M., & Hadler Boggiani, B. (2023). Landscape dynamics and the
Phanerozoic diversification of the biosphere. *Nature*, *624*(7990).
https://doi.org/10.1038/s41586-023-06777-z

Saupe, E. E., Hendricks, J. R., Portell, R. W., Dowsett, H. J., Haywood, A., Hunter, S. J., &
Lieberman, B. S. (2014). Macroevolutionary consequences of profound climate change on
niche evolution in marine molluscs over the past three million years. *Proceedings of the Royal
Society B: Biological Sciences*, *281*(1795). https://doi.org/10.1098/rspb.2014.1995

Schobben, M., Van De Schootbrugge, B., & Wignall, P. B. (2019). Interpreting the carbon isotope
record of mass extinctions. *Elements*, *15*(4). https://doi.org/10.2138/gselements.15.5.331

Scotese, C. R. (2021). An atlas of phanerozoic paleogeographic maps: The seas come in and the
seas go out. In *Annual Review of Earth and Planetary Sciences* (Vol. 49, pp. 679–728).
https://doi.org/10.1146/annurev-earth-081320-064052

Scotese, C. R., & Wright, N. (2018). *PALEOMAP Paleodigital Elevation Models (PaleoDEMS) for
the Phanerozoic PALEOMAP Project, https://www.earthbyte.org/paleodem-resource-scotese-
and-wright-2018/.*

Sepkoski, J. J. J. (2002). A compendium of fossil marine animal genera. Edited by David Jablonski
and Michael Foote. *Bulletins of American Paleontology*, *363*(363), 1–560.

Song, H., Kemp, D. B., Tian, L., Chu, D., Song, H., & Dai, X. (2021). Thresholds of temperature
change for mass extinctions. *Nature Communications*, *12*(1). https://doi.org/10.1038/s41467-

021-25019-2

Stadler, T. (2011). Mammalian phylogeny reveals recent diversification rate shifts. *Proceedings of*
the National Academy of Sciences of the United States of America, *108*(15).
https://doi.org/10.1073/pnas.1016876108

Storch, D., & Okie, J. G. (2019). The carrying capacity for species richness. *Global Ecology and*
Biogeography, *28*(10), 1519–1532. https://doi.org/10.1111/geb.12987

Tarantola, A. (2005). Inverse Problem Theory and Methods for Model Parameter Estimation. In
Inverse Problem Theory and Methods for Model Parameter Estimation.
https://doi.org/10.1137/1.9780898717921

Valentine, J. W., Foin, T. C., & Peart, D. (1978). A provincial model of Phanerozoic marine
diversity. *Paleobiology*, *4*(1), 55–66. https://doi.org/10.1017/S0094837300005686

Valentine, J. W., & Moores, E. M. (1970). Plate-tectonic regulation of faunal diversity and sea level:
A model. *Nature*, *228*(5272), 657–659. https://doi.org/10.1038/228657a0

Vérard, C., Hochard, C., Baumgartner, P. O., Stampfli, G. M., & Liu, M. (2015). 3D
palaeogeographic reconstructions of the Phanerozoic versus sea-level and Sr-ratio variations.
Journal of Palaeogeography, *4*(1), 64–84. https://doi.org/10.3724/SP.J.1261.2015.00068

Vermeij, G. J. (1977). The mesozoic marine revolution: Evidence from snails, predators and
grazers. *Paleobiology*, *3*(3), 245–258. https://doi.org/10.1017/S0094837300005352

Vermeij, G. J. (2017). How the Land Became the Locus of Major Evolutionary Innovations. *Current*
Biology, *27*(20). https://doi.org/10.1016/j.cub.2017.08.076

Willig, M. R., Kaufman, D. M., & Stevens, R. D. (2003). Latitudinal Gradients of Biodiversity:
Pattern, Process, Scale, and Synthesis. In *Annual Review of Ecology, Evolution, and*
Systematics (Vol. 34). https://doi.org/10.1146/annurev.ecolsys.34.012103.144032

Wright, D. H. (1983). Species-Energy Theory: An Extension of Species-Area Theory. *Oikos*, *41*(3).
https://doi.org/10.2307/3544109

Xuereb, A., Rougemont, Q., Tiffin, P., Xue, H., & Phifer-Rixey, M. (2021). Individual-based eco-
evolutionary models for understanding adaptation in changing seas. In *Proceedings of the*
Royal Society B: Biological Sciences (Vol. 288, Issue 1962).
https://doi.org/10.1098/rspb.2021.2006

Zaffos, A., Finnegan, S., & Peters, S. E. (2017). Plate tectonic regulation of global marine animal
diversity. *Proceedings of the National Academy of Sciences of the United States of America*,
114(22), 5653–5658. https://doi.org/10.1073/pnas.1702297114