Caenorhabditis elegans **Uses Canonical and Non-canonical Hippo signaling**

Linh Huynh[1], Razan A. Fakieh[1,2], C'Brionne Hendrix[1], Reid Powell[1], David J. Reiner[1,3,§]

[1] College of Medicine, Texas A&M University, Houston TX, 77030, USA

[2] Clinical Laboratory Sciences Department, College of Applied Medical Sciences, Imam Abdulrahman bin

Faisal University, Dammam 34212, Kingdom of Saudi Arabia.

[3] Institute of Biosciences and Technology, Texas A&M Health Science Center, Houston TX, 77030, USA

16   [§] To whom correspondence should be addressed: dreiner@tamu.edu

41   Key words: Wts, Yki, Hpo, Sterile 20, MST1, MST2, Hippo, Warts, Yap, TEAD, MIG-15, GCK-2

ABSTRACT

Hippo signaling is a conserved regulator of tissue homeostasis across metazoans. The Ste20 family kinase Hippo/MST activates the NDR family kinase Warts/LATS to inhibit the transcriptional coactivator Yorkie/YAP/TAZ and its transcription factor partner Scalloped/TEAD. In *Caenorhabditis elegans*, cell lineages and organ sizes are largely invariant, and classical Hippo phenotypes such as tissue overgrowth are absent. Nevertheless, WTS-1, YAP-1, and the TEAD-like transcription factor EGL-44 form a conserved core module required for larval development past the L2 stage. Crucially, a direct role for Hippo signaling remains unestablished. To address this question, we generated a fluorescently tagged endogenous YAP-1 as a live biomarker of pathway activity. Upon WTS-1 loss, endogenous YAP-1 translocated from cytosol to nucleus in epithelium and intestine. Tissue-specific depletion revealed that intestinal, but not epithelial, WTS-1 is essential for progression past L2. The duplicated Hippo-related kinases CST-1 and CST-2 repressed YAP-1 nuclear localization in the epithelium but not intestine, indicating that intestinal WTS-1 functions without CST-1/2. The Ste20 kinase MIG-15, orthologous to *Drosophila* Misshapen and mammalian MAP4K4/6/7/8, was redundant with CST-1/2 for larval progression. Yet deficient MIG-15 uniquely increased YAP-1 abundance without driving nuclear localization. By contrast, the Ste20 kinase GCK-2, orthologous to *Drosophila* Happyhour and mammalian MAP4K1/2/3/5, had no detectable role. Our findings establish *C. elegans* as a model for Hippo signaling, with a canonical cascade active in the epithelium and noncanonical inputs controlling WTS-1 in the intestine. In this context, YAP-1/EGL-44 outputs are repurposed from growth control to non-proliferative developmental functions.

INTRODUCTION

The Hippo signaling pathway was first discovered in *Drosophila* loss-of-function screens that identified tumor suppressor genes required to prevent tissue overgrowth (JUSTICE *et al.* 1995; TAPON *et al.* 2002; HARVEY *et al.* 2003; WU *et al.* 2003). Subsequent studies revealed that the downstream transcriptional coactivator Yorkie (Yki) is essential to avoid the converse phenotype of tissue undergrowth (HUANG *et al.* 2005). The canonical pathway consists of the Ste20-family (from yeast Sterile 20) kinase Hippo (Hpo; mammalian Mst1/2) phosphorylating and activating the NDR-family kinase Warts (Wts; mammalian LATS1/2), which in turn phosphorylates and represses Yki/YAP/TAZ (HUANG *et al.* 2005; OH AND IRVINE 2010). Phosphorylated Yki/YAP/TAZ is excluded from the nucleus, preventing it from associating with the transcription factor Scalloped (Sd; mammalian TEAD1-4) to drive expression of growth-promoting transcriptional client genes (VASSILEV *et al.* 2001; WU *et al.* 2008; ZHANG *et al.* 2008), reviewed in (HALDER AND JOHNSON 2011). In summary, the canonical Hippo cascade is Hippo/Mst → Warts/LATS ⊣ YAP/TAZ+TEAD.

Genetic studies in *Drosophila* and mammalian cells revealed an absolute requirement for Wts/LATS in pathway function, but a partial or context-dependent requirement for Hippo/MST1/2. These observations prompted the discovery of other Ste20-family kinases, but outside the Hippo subfamily, that act redundantly or in parallel to Hippo in specific contexts (MENG *et al.* 2015; ZHENG *et al.* 2015), highlighting the complexity of inputs upstream of Wts/LATS. In most metazoans, the Hippo pathway maintains tissue homeostasis by allowing organs to expand or regress in response to developmental or environmental cues, and its dysregulation contributes to tumorigenesis. Upstream kinases and scaffolds act as tumor suppressors, while YAP/TAZ and TEAD can serve as oncogenic drivers when overactivated (MOROISHI *et al.* 2015; PEARSON *et al.* 2021; HAN *et al.* 2024; HARVEY AND TANG 2025).

A distinctive feature of the Hippo pathway that distinguishes it from other pathways is the absence of a conventional extracellular signaling ligand and transmembrane receptor. Instead, the upstream Hippo kinase is activated by diverse structural elements of the cell: the cytoskeleton, apicobasal cell polarity, and substrate-cell/cell-cell adhesion (MARTIN-BELMONTE AND PEREZ-MORENO 2011). Attenuation of these structures accompany epithelial-to-mesenchymal transition, and hence activation of YAP/TAZ signaling is a major driver of tumor progression (KAPOOR *et al.* 2014; LIN *et al.* 2023). Triggering of downstream YAP activity also suppresses anoikis (FRISCH *et al.* 2013).

Yet in the nematode worm *Caenorhabditis elegans*, development is largely invariant and mosaic: most cell fates are determined by lineage rather than by cell-cell signaling. This rigid developmental program underlies its consistent organ size and stereotyped embryogenesis, unlike the plastic growth observed in flies or vertebrates (HARIHARAN 2015; ROTHMAN AND JARRIAULT 2019). Consequently, canonical Hippo phenotypes – tissue overgrowth or undergrowth – have not been observed in *C. elegans*, and conventional forward genetic screens have not recovered mutants in Hippo, Warts, YAP, or TEAD.

Nonetheless, molecular conservation suggested that a core Warts→YAP/TEAD module might be present.

Warts is conserved as WTS-1, whose loss causes early larval arrest with compromised intestinal integrity

under nutrient stress (CAI *et al.* 2009; KANG *et al.* 2009). YAP-1 and the TEAD ortholog EGL-44 physically

interact and are negatively regulated by WTS-1 (IWASA *et al.* 2013), and WTS-1-dependent arrest

requires both YAP-1 and EGL-44 (LEE *et al.* 2019). Similarly, NFM-1, the ortholog of Merlin/NF2, the well-

established co-regulator of Wts in *Drosophila* and a tumor suppressor in mammals, acts in parallel to

WTS-1 to support intestinal polarity and developmental progression (LEE *et al.* 2019). Together, these

findings suggest that while the molecular framework of WTS-1→YAP-1/EGL-44 is conserved, it does not

drive proliferative phenotypes as in other metazoans.

In *C. elegans,* the role of the namesake Hippo kinase in *C. elegans* remains unclear. Two nearly

identical Ste20-family kinases, CST-1 and CST-2, share homology with Hippo but differ at the C-terminus:

CST-2 lacks three exons encoding the canonical SARAH coiled-coil domain, a motif characteristic of

Hippo, Salvador, and RASSF proteins that mediates homo- and -hetero-dimerization and pathway

activation (KARCHUGINA *et al.* 2021). RNAi depletion of both genes induces autophagy and affects aging

(LEHTINEN *et al.* 2006; WILKINSON *et al.* 2015), and double deletion causes mild neural defects but no

arrest, with no clear connection to the WTS-1→YAP-1/EGL-44 axis. These findings have led to the

proposal that *C. elegans* lacks Hippo input to WTS-1 (YANG AND HATA 2013).

This background raises a central question: does *C. elegans* employ Hippo-related kinases to activate

WTS-1 repression of YAP-1, or has regulation of the pathway diverged from canonical mechanisms? To

address this, we developed a fluorescent YAP-1 reporter to directly monitor pathway activity and assess

the contributions of CST-1/2 and other Ste20 kinases to WTS-1 regulation in two distinct tissues. We

validated this tool by showing that WTS-1 cell autonomously represses YAP-1 in both epithelium and

intestine. In the epithelium, Hippo-like proteins CST-1/2, like WTS-1, repress YAP-1, indicating likely

conservation of the canonical Hippo signaling pathway. In contrast, in the intestine repression of YAP-1

upstream of WTS-1 may occur independently of or in parallel to CST-1/2. These findings highlight both

the conservation and divergence of Hippo signaling mechanisms in *C. elegans*.

RESULTS

WTS-1 is required for larval development and represses YAP-1 and EGL-44/TEAD to promote

developmental progression. A prior study reported that the *wts-1(ok753)* deletion allele causes early

larval arrest (KANG *et al.* 2009). To more comprehensively characterize the terminal phenotype and

expand available genetic tools, we analyzed two additional alleles: *wts-1(tm4081)*, a deletion not

available during the original study, and *wts-1(re436)*, a gene disruption (see below). We also developed

a conditional depletion strategy using an auxin-inducible degron (AID*) to acutely remove endogenous

WTS-1 protein (**Fig. S1A**).

As expected, animals homozygous for *wts-1(ok753)* or *wts-1(tm4081)* exhibited early larval arrest.

Both alleles were outcrossed and rebalanced with a fluorescently tagged *tmC18* balancer (DEJIMA *et al.*

2018) covering the *wts-1* locus on Chromosome I. Homozygotes segregated from heterozygous mothers

arrested at the L2 stage (**Fig. 1A,B**; **S1A**). Arrested animals continued to move and pump for several

days, indicating they were alive (**Fig. 1C**). Differential interference contrast (DIC) microscopy revealed

no overt structural defects in L2-arrested animals of either genotype. However, both locomotion and

pumping declined by Day 3, suggesting reduced physiological health. This finding is consistent with

earlier observations of loss of intestinal integrity and cytoskeletal disorganization in *wts-1(ok753)* animals

subjected to prolonged starvation (KANG *et al.* 2009). We speculate that sustained WTS-1 loss

compromises animal resilience, perhaps including intestinal integrity. This differs from arrest caused by

loss of the small GTPase RHEB-1 or mTORC1, where animals remain viable and can live a full *C. elegans*

lifespan after hatching (DUONG *et al.* 2020).

Knockout consortium mutants are heavily mutagenized and can harbor background mutations. SO

using CRISPR/Cas9-dependent genome editing, we introduced a STOP-IN cassette (WANG *et al.* 2018)

into the 5' end of the *wts-1* locus to generate *wts-1(re436)*, a 43 bp insertion that also removes the A of

the ATG start codon (**Fig. S1A**). While *wts-1(re436)* mutants arrested, many progressed beyond L2,

suggesting that this allele is not a complete loss-of-function (**Fig. 1B**). To assess residual gene activity,

we compared *wts-1(re436)* and *wts-1(ok753)* animals grown on *control(RNAi)* versus *wts-1(RNAi)*. *wts-*

1(ok753) animals showed mildly enhanced arrest on *wts-1(RNAi)*, with some animals arresting in L1 (**Fig.**

S1B), likely due to depletion of maternally contributed gene product. These results suggest that *wts-*

1(ok753) is a strong loss-of-function allele, and that a full null phenotype may manifest as L1 or L1/L2

arrest. *wts-1(re436)* animals exhibited a more severe phenotype when exposed to *wts-1(RNAi)* than to

control(RNAi), confirming that *re436* mutants retain partial function. As *wts-1* has an 81 nt 5′ UTR,

transcriptional initiation is likely unaffected by *re346*. We speculate that the mutant transcript may

undergo cryptic translation initiation downstream of the disrupted ATG.

To enable conditional chemical-genetic depletion of endogenous WTS-1, we used CRISPR/Cas9 to

insert an *mTurquoise2::2xMyc::AID** tag at the 5′ end of the *wts-1* coding sequence, using the same guide

RNA as the STOP-IN strategy (mTurquois2, or mT2, is a blue fluorescent protein). This minimal AID*

degron (44 amino acids) is recognized by the TIR1 E3 ubiquitin ligase, which mediates degradation of

AID*-tagged proteins in the presence of auxin (indole-3-acetic acid; IAA; (ZHANG *et al.* 2015). The

insertion was determined to be error-free by Sanger sequencing. However, we were unable to visualize

blue-tagged WTS-1 beyond very faint signal from the intestinal brush border and we have struggled to

detect Myc tags in our lab (Wu and Reiner, unpublished results).

AID*::WTS-1+somatic TIR1 embryos hatched on auxin plates showed delayed development relative

to vehicle-treated controls at Day 3. However, by Day 4, these animals had still reached adulthood,

indicating developmental delay rather than arrest (**Fig. 1D–I**), indicating depletion was incomplete. In

contrast, double depletion with both auxin treatment and *wts-1(RNAi)* caused robust L2 arrest at both

Day 3 and Day 4, closely resembling the phenotype of strong loss-of-function alleles (**Fig. 1D, J–N**).

Expression of TIR1 in the germline did not confer arrest in auxin-treated AID*::WTS-1 animals (**Fig. S1C**),

suggesting that WTS-1 is not essential for embryonic development prior to the maternal-to-zygotic

transition in gene expression. Additionally, animals expressing somatic TIR1 but lacking the AID* tag

developed normally in the presence of auxin (**Fig. S1D–F**), confirming that auxin itself does not impair

development.

WTS-1 is required to repress YAP-1 and EGL-44/TEAD for development progression past beyond

L2. Previous work by Lee and colleagues demonstrated that the *wts-1* mutant arrest depends on the

activity of YAP-1 and the TEAD transcription factor EGL-44 (LEE *et al.* 2019). This observation is

consistent with the conserved role of Warts/LATS kinases in repressing YAP/TAZ and TEAD in *Drosophila*

and mammals (*i.e.* Hippo/Mst → Warts/LATS ⊣ YAP/TAZ+TEAD) (HANSEN *et al.* 2015; DASGUPTA AND

MCCOLLUM 2019). We tested whether RNAi directed against *yap-1* or *egl-44* could suppress the L2 arrest

of *wts-1(tm4081)* animals. *yap-1(RNAi)* fully reversed while *egl-44(RNAi)* partially reversed the arrest

(**Fig. 1O**). Similarly, both *yap-1(RNAi)* and *egl-44(RNAi)* suppressed the developmental delay caused by

auxin treatment in AID*::WTS-1 animals (**Fig. S1G–J**). These results corroborate the findings of Lee *et*

al. using new genetic alleles and validate the use of conditional WTS-1 depletion. We conclude that WTS-

1 represses YAP-1 and EGL-44/TEAD to permit developmental progression, a mechanism that is

conserved among metazoans.

Generation of tagged endogenous YAP-1 for use as a reporter of upstream signaling. The

transcriptional coactivator *Drosophila* Yorkie (Yki) and its human orthologs Yap and TAZ are well

established reporters of upstream Wts/LATS activity. Their subcellular localization reflects regulation by

upstream kinases, particularly the Warts serine/threonine kinase (LATS1/2 in mammals). In this

conserved mechanism, phosphorylation of YAP by Warts prevents nuclear entry by promoting

cytoplasmic retention; inactivation of Warts permits dephosphorylation and nuclear translocation of YAP

(HANSEN *et al.* 2015; DASGUPTA AND MCCOLLUM 2019).

Hata and colleagues (IWASA *et al.* 2013) identified the *C. elegans* ortholog of YAP-1 and generated a

high-copy integrated translational reporter (*yap-1*p>*yap-1::gfp*). GFP expression from this transgene is

broad and YAP-1::GFP signal is predominantly cytoplasmic, with pronounced nuclear exclusion,

particularly in epithelial tissues. WTS-1 and the 14-3-3 protein FTT-2 are predicted inhibitors of YAP-1

nuclear localization (HANSEN *et al.* 2015; DASGUPTA AND MCCOLLUM 2019). Consistent with this

prediction, RNAi-mediated depletion of *wts-1* or *ftt-2* resulted in nuclear accumulation of YAP-1::GFP,

supporting an evolutionarily conserved role for Warts-mediated inhibition of YAP in *C. elegans* (IWASA *et*
al. 2013)*.*

In contrast, RNAi of other putative upstream components failed to cause nuclear accumulation of
YAP-1::GFP. These negative results include depletion of *cst-1* and *cst-2*, which encode predicted Hippo-
like kinases resembling mammalian MST1/2, the eponymous initiators of the pathway. These negative
results raise doubts that the Hippo pathway is conserved in *C. elegans* upstream of Warts and Yap (IWASA
et al. 2013). Indeed, the "Hippo pathway" in *C. elegans* has been debated as to whether it is a conserved
signaling module, a collection of disconnected components, or ignored in the discussion altogether
(HALDER AND JOHNSON 2011; YANG AND HATA 2013). Similar ambiguity surrounds the *C. elegans*
"Hedgehog" pathway, which retains a GLI transcription factor ortholog but lacks most core pathway
elements, but in the case of Hedgehog the conclusion is the pathway has been mostly lost during
evolution (BURGLIN AND KUWABARA 2006).

We reasoned that the high expression level of transgenic YAP-1::GFP may reduce sensitivity to
perturbation of upstream signal. Possible explanations include buffering of phosphorylation due to
excessive protein levels, a lower signal-to-noise ratio in detection of fluorescence, insufficient RNAi
knockdown to overcome the threshold required for YAP-1 nuclear entry, or potential redundancy of the
duplicated *cst-1* and *cst-2* genes (though these duplicated genes are mostly identical at the DNA level,
see below).

To generate a more sensitive and physiologically relevant reporter of endogenous YAP-1 dynamics,
we used CRISPR/Cas9-dependent genome editing to insert sequences encoding a short linker,
mNeonGreen (a photostable, yellow-shifted fluorescent protein; (SHANER *et al.* 2013) and a 2x FLAG
epitope at the 3' end of the *yap-1* coding sequence, generating *yap-1(re269[yap-1::mNG::2xflag])*; **Fig.**
S2A). The resulting edited *yap-1* was error-free by Sanger sequencing. Immunoblotting with anti-FLAG
antibody confirmed the predicted tagged protein size of 80.6 kDA (**Fig. S2B**). YAP-1::mNG::2xFLAG was
crossed with blue-fluorescent histone marker HIS-72::mT2 and imaged, revealing apparently ubiquitous
expression relative to the wild type and low background autofluorescence from unedited controls.
Representative images of epithelium and intestine are shown (**Fig. S2C**).
230

WTS-1 represses YAP-1 nuclear translocation in epithelium and intestine. In all systems studied to
date, Warts/LATS1/2 kinases repress YAP nuclear translocation. To test whether this regulatory
mechanism is conserved in *C. elegans*, we used our endogenously tagged YAP-1::mNG reporter to
monitor subcellular localization in response to WTS-1 depletion.

We constructed a reporter strain expressing YAP-1::mNG (yellow, imaged at 514 nm) and HIS-
72::mTurquoise2 (blue, imaged at 445 nm) for nuclear reference. This strain also harbored a *rrf-3* loss-

of-function mutation to enhance sensitivity to RNAi (SIMMER *et al.* 2002). Despite this sensitization, *wts-*

1(RNAi) did not cause overt developmental delay or lethality in this strain.

However, imaging revealed that *wts-1(RNAi)* robustly induced nuclear accumulation of YAP-1::mNG

in both epithelial cells (**Fig. 2A-C**) and intestinal cells (**Fig. 2D-F**), compared to *control(RNAi)*. These

results were recapitulated using the AID*::WTS-1 system: addition of auxin induced nuclear translocation

of YAP-1::mNG in epithelial (**Fig. S2D-F**) and intestinal cells (**Fig. S2G-I**), relative to vehicle-treated

controls.

Unexpectedly, we found that the mNG::2xFLAG knock-in into the C-terminus of YAP-1 suppressed

the lethality associated with *wts-1* deletion. Double mutants of genotype *wts-1(tm4081)*; *yap-1(re269[yap-*

1::mNG::2xflag]) developed normally, showing no arrest or delay, and were fully viable and fertile (**Fig.**

S2J-L), in stark contrast to *wts-1* single mutants (**Fig. 1**). Nevertheless, YAP-1::mNG strongly

accumulated in the nucleus in both epithelial and intestinal cells of *wts-1(tm4081)* mutants (**Fig. S2M-P'**).

Lee *et al*. had showed that *wts-1(RNAi)* in an *rrf-3* mutant background conferred arrest, but the

endogenous *yap-1* gene in their experiment was unaltered (KANG *et al.* 2009).

Notably, a previous study used an exogenous *yap-1*p>*yap-1::gfp* transgene (IWASA *et al.* 2013), which

would not reveal defects in co-transcriptional activity caused by C-terminal tagging. Yet this defect was

unexpected, as key functional domains of YAP-1, including the TEAD binding domain (TBD) and WW

domains, are located at the N-terminus and central region of the protein, respectively, across diverse

species (HILMAN AND GAT 2011; YANG AND HATA 2013; MESROUZE *et al.* 2021).

We conclude that YAP-1::mNG is a reliable reporter of upstream WTS-1 activity, with nuclear

localization reflecting loss of WTS-1-mediated inhibition. However, the C-terminally tagged YAP-1::mNG

does not appear to function as a transcriptional coactivator; while some YAP/TAZ proteins have a C-

terminal PDZ-recognition motif, these are not present in insect YAP proteins nor in YAP-1 (HILMAN AND

GAT 2011). Since *yap-1* deletion mutants are superficially wild type (LEE *et al.* 2019; APKEN AND

OECKINGHAUS 2021), the transcriptional defect caused by C-terminal tagging of YAP-1 would remain

unnoticed unless assayed in a *wts-1* mutant background, where L2 arrest is reversed by loss of YAP-1

function.

WTS-1 functions cell autonomously to repress YAP-1 in epithelium and intestine. A key test for of

cell-cell signaling function is cell autonomy: does a protein function in the tissue manifesting the mutant

phenotype or another tissue? Previous results showed that the early larval arrest of *wts-1(ok753)* animals

was rescued by transgenic expression of *wts-1(+)* expressed by an intestinal promoter, demonstrating

that intestinal expression of WTS-1 is sufficient to support development (KANG *et al.* 2009). Using our

conditional chemical-genetic degron, we used tissue-specific expression of the cofactor for AID*-auxin to

test the converse question: in which tissue is WTS-1 necessary to support normal developmental

progression?

We expressed TIR1, the E3 ubiquitin ligase substrate recognition protein for AID*, specifically in

intestinal or epithelial tissues and treated AID*::WTS-1 with auxin vs. vehicle as described (ASHLEY *et al.*

2021). In AID*::WTS-1 animals expressing epithelial TIR1 and grown on *wts-1(RNAi)* to increase

robustness, addition of auxin caused mild growth delay relative to vehicle controls (**Fig. 3A-C**). The same

outcome occurred without RNAi, though with reduced severity (**Fig. S3A-C**). Degradation efficacy was

monitored using degradation of a tissue-specific AID*-tagged nuclear BFP reporter as an internal control.

In epithelial TIR1 animals not treated with *wts-1(RNAi)*, auxin induced BFP degradation and nuclear

accumulation of YAP-1::mNG in the epithelium (**Fig. 3G-I'**), but not in the intestine (**Fig. 3I-J'**).

In the converse experiment, intestinal TIR1 expression with auxin treatment in *wts-1(RNAi)* animals

caused complete developmental arrest (**Fig. 3D-F**), with weaker effects without RNAi (**Fig. S3D-F**). Auxin

treatment also induced intestinal BFP degradation and nuclear translocation of YAP-1::mNG in the

intestine, but not in the epithelium (**Fig. 3K-L", I-J"**).

These results confirm tissue-specific degradation via the AID*-TIR1 system and show that WTS-1 is

required in the intestine for developmental progression and repression of YAP-1::mNG. This

complements prior rescue studies of *wts-1(Δ)* with intestinally driven *wts-1(+)* (KANG *et al.* 2009). We also

find that WTS-1 represses YAP-1::mNG in epithelium, and that epithelial WTS-1 depletion causes a

modest growth delay, though whether this is related to intestinal effects remains unclear.

CST-1/2, the putative *C. elegans* Hippo ortholog, represses YAP-1::mNG in epithelium but not

intestine. In *Drosophila* and mammals, Hippo (Hpo) and MST1/2 are Ste20-family kinases (DAN *et al.*

2001; DELPIRE 2009) that regulate growth via activation of Wts and hence repression of YAP/TEAD. While

conserved in many species, a clear Hippo ortholog function in *C. elegans* had not been established.

C. elegans encodes two Hippo-like kinases, CST-1 and CST-2, are roughly equivalently expressed

from tandemly duplicated genes on chromosome X (**Fig. 4A**; mean FPKM of 32 and 24 for *cst-1* and *cst-*

2, respectively). These proteins are 100% identical through most of their sequence, including the kinase

domain, but differ in the final 13 amino acids of exon 9. *cst-2* CDS (coding DNA sequence) stops at the

end of exon 9, while *cst-1* contains three additional exons that are not present in *cst-2*. Importantly, CST-

2 lacks the C-terminal SARAH coiled-coil domain present in all Hpo-related proteins, including CST-1

(**Fig. S4A**), which is essential in other systems for Hippo signaling via homo- and hetero-dimerization

(KARCHUGINA *et al.* 2021). Thus, CST-2 may be regulated differently, act independently, or be a non-

functional byproduct of genomic duplication.

100% DNA identity extends through all shared exons and upstream sequences of *cst-1* and *cst-2*,

even extending several hundred bp upstream of the ATG initiator methionine codon (**Fig. S4B,C**).

Examined related Caenorhabditid nematodes encode only a single CST-1 ortholog with a SARAH

domain, suggesting a recent duplication event in *C. elegans*.

The *cst-1-cst-2* genomic region was previously deleted as a validation of the transposon-mediated

MosDEL deletion technology (Frokjaer-Jensen *et al.* 2010). In our hands, mutants for the resulting

lesion, *basDf1*, are viable, grow slightly slowly, and defective for locomotion. We deleted the *cst-1/2* locus

– from exon 1 of *cst-2* to intron 11 of *cst-1* – using CRISPR/Cas9-dependent genome editing, creating

the *cst-1/2(reDf4[re452])* allele (**Fig. 4A**). Deletion of *cst-1/2*, like *basDf1*, conferred slow growth and mild

locomotion defects, but was not arrested.

The absence of lethality from Δ*cst-1/2* mutants, in contrast to Δ*wts-1*, leads us to speculate that CST-

1/2 does not recapitulate the role of WTS-1 throughout the animal. Yet in the *reDf4* mutant, YAP-1::mNG

translocated to epithelial nuclei (**Fig. 4B-D**), similar to WTS-1 depletion, suggesting CST-1 and possibly

CST-2 activate WTS-1 in epithelium.

Strikingly, Δ*cst-1/2* did not cause nuclear YAP-1::mNG translocation in intestine (**Fig. 4E-G**). Given

that intestinal *wts-1(+)* rescues *wts-1(ok753)* and that intestinal WTS-1 degradation causes arrest, we

propose that CST-1/2 is not necessary for the intestinal WTS-1-dependent event, only the epithelial WTS-

1-dependent event. This observation is consistent with studies in flies and mammals, where in certain

contexts Hippo/Mst1/2 mutants showed less penetrant phenotypes than Warts/LATS mutants, likely due

to kinase redundancy at the level of Ste20 family kinases Hippo/Mst1/2 (Meng *et al.* 2015; Zheng *et al.*

2015).

Deletion/depletion of both CST-1/2 and Ste20-family kinase MIG-15 cause WTS-1-like L2 arrest.

Previous studies identified Ste20-family kinases from the GCK-I and GCK-IV subfamilies functioning

redundantly with Hippo/MST1/2 to activate Wts/LATS1/2 in flies and mammalian cells (Meng *et al.* 2015;

Zheng *et al.* 2015). Specifically, these are the GCK-I and GCK-IV subfamilies of the Ste20 family (Dan

et al. 2001; Delpire 2009). Both consist of N-terminal Ste20 kinase domains and C-terminal CNH (Citron

homology) domains, yet they are distinct subfamilies in metazoans. The GCK-I subfamily includes *C.*

elegans GCK-2*,* fly Happyhour, and mammalian MAP4K1,2,3,5. The GCK-IV subfamily includes *C.*

elegans MIG-15*,* fly Misshapen, and mammalian MAP4K4,6,7,8. We previously showed that in *C.*

elegans GCK-2 promotes vulval 2° fate and MIG-15 promotes vulval 3° fate (Shin *et al.* 2018; Fakieh

and Reiner 2025).

We tested whether these Ste-20-family GCK-I and GCK-IV subfamily kinases function redundantly

with CST-1/2 to activate WTS-1. In AID*::mNG::2xHA::MIG-15 animals with somatic TIR1, auxin

treatment of MIG-15-depleted animals caused mild growth delay (**Fig. 5A-D, I-J**). With same experiment

in a *cst-1/2(Δ)* background, auxin-induced MIG-15 depletion caused robust L2 arrest, phenocopying *wts-*

1 mutants (**Fig. 5E-H, I-J**). Also, as with deficient *wts-1*, MIG-15 and CST-1/2 deficient and arrested

animals continued movement and feeding (**Fig. 5K**). This arrest phenotype was reversed by *yap-1(RNAi)*

and to a lesser extent by *egl-44(RNAi)* (**Fig. 5L-O**). One interpretation is that MIG-15 also contributes to

WTS-1/YAP-1 regulation. Alternatively, the function of MIG-15 could be entirely in parallel to CST-1/2;

RNAi of *yap-1* or *egl-44* may suppress effects of mutant *cst-1/2* to retore growth, without impacting the

downstream consequences of mutant *mig-15*.

To test the Ste20-family GCK-I subfamily member GCK-2, we generated animals with *cst-1/2(Δ)*,

AID*::MIG-15, and somatically expressed TIR1 with and without *gck-2(re483)*. Animals arrested upon

addition of auxin regardless of whether *gck-2* was deleted (**Fig. S5A-J**). Triple mutant *gck-2 cst-1/2 mig-*

15 animals continued feeding (**Fig. S5K**), like *cst-1/2 mig-15* mutants or *wts-1* mutants, above. A STOP-

IN allele, *gck-2(re427)*, edited into the *cst-1/2(Δ)* YAP-1::mNG animal, failed to send YAP-1::mNG into

the nucleus (**Fig. 5SL-O**). Thus, GCK-2 does not appear to contribute to this developmental event, though

we cannot rule out GCK-2 in other Hippo-regulated signaling events in the animal.

Ste20-family kinase MIG-15 represses YAP-1::mNG levels but not nuclear translocation. Given the

synthetic L2 arrest in auxin+AID*::MIG-15-depleted *cst-1/2(Δ)* animals, we examined YAP-1::mNG

localization in these backgrounds. Our MIG-15 tag thus far is with sequences encoding mNeonGreen

(FAKIEH AND REINER 2025), which would interfere with visualization of YAP-1::mNG nuclear translocation.

Consequently, we used CRISPR/Cas9-dependent genome editing to insert mTurquoise2::2xMyc::AID*

at the 5' end of *mig-15* in animals expressing somatic TIR1 and YAP-1::mNG (**Fig. S6A**). MIG-15

depletion by auxin caused a protruding-vulva phenotype found in all *mig-15* mutants (**Fig. S6B-C**),

consistent with strong depletion.

Surprisingly, auxin+AID*::MIG-15 depletion with *cst-1/2(+)* increased total YAP-1::mNG levels but did

not induce nuclear localization. In epithelial cells, we observed perinuclear aggregation of YAP-1::mNG

(**Fig. 6A-C**), potentially indicating localization to Golgi. In intestine, MIG-15 depletion increased YAP-

1::mNG expression and weak nuclear localization, but less than that observed in *wts-1* (**Fig. 6D-G**).

We expected that combined MIG-15 depletion and *cst-1/2(Δ)* would cause robust YAP-1::mNG

nuclear localization. Instead, nuclear levels resembled those in MIG-15 depletion alone (**Fig. 6H-K**).

While this result was unexpected, the synthetic L2 arrest and rescue by *yap-1(RNAi)* and *egl-44(RNAi)*

suggest that MIG-15 and CST-1/2 converge on WTS-1 or YAP-1 regulation. However, they may regulate

YAP-1 via different mechanisms or in parallel pathways, as shown in our model of these results (**Fig. 7**).

DISCUSSION

In this study, we established that *C. elegans* epithelium possesses the molecular wiring of a canonical

Hippo→Warts ⊣ YAP/TEAD pathway, while also revealing unexpected divergence in its tissue-specific

regulation, specific in intestine. Using an endogenous YAP-1 fluorescent reporter, we showed that WTS-
1 cell-autonomously excludes YAP-1 from the nucleus in both epithelia and intestine. Tissue-specific
depletion of WTS-1 demonstrated that WTS-1 cell-autonomously prevents animal arrest in intestine. At
the same time, our data show that the upstream Hippo-related kinases CST-1/2 contribute to YAP-1
regulation only in the epithelium, while intestinal regulation proceeds through a distinct, CST-1/2-
independent or CST-1/2-parallel mechanism. We identified MIG-15, the nematode ortholog of *Drosophila*
Msn and mammalian MAP4K4,6,7,8 kinases – implicated in activation of fly and mammalian Wts/LATS
in parallel to Hippo/MST1/2 – as redundant with CST-1/2 developmental progression past the L2 stage
and reliant on inhibition of YAP-1/EGL-44. Unexpectedly, MIG-15 represses of abundance of YAP-1
protein, perhaps through blocking a degradative process, but not nuclear translocation. Together, these
results establish a hybrid picture: a canonical Hippo-like cascade in the epithelium *vs.* noncanonical,
parallel inputs to WTS-1 in the intestine.
388
Divergence of Hippo inputs in *C. elegans*. The presence of one Hippo kinase with a canonical SARAH-
domain (CST-1) and another Hippo kinase without (CST-2), coupled with the redundant CST-1/2 kinases
not being essential like WTS-1, has fueled uncertainty about whether a *bona fide* Hippo pathway operates
in *C. elegans*. Our findings clarify this issue. CST-1/2 act as functional Hippo-like inputs in the epithelium,
supporting the model of conservation of the canonical cascade. However, in intestine regulation appears
to rely on unknown, alternate kinases. In flies and mammals, MAP4K family members (orthologous to
GCK-2 and MIG-15 in worms) act redundantly with Hippo to activate LATS/WTS (MENG *et al.* 2015;
ZHENG *et al.* 2015). Our analysis shows that MIG-15, but not GCK-2, alters YAP-1 protein levels in
nematodes, suggesting that these kinases may have retained ancillary regulatory functions while losing
the ability to control YAP-1 nuclear localization. This divergence highlights the evolutionary plasticity of
Hippo inputs across cell types.
Alternatively, MIG-15 and GCK-2 may contribute to phosphorylation of WTS-1 and repression of YAP-
1 in other tissues. Epithelium and intestine were assayed because they are large tissues with nuclei
readily imaged via confocal microscopy during development. Yet many other tissues or individual cells
undergo myriad developmental events while forming the mature animal. For example, EGL-44 regulates
identity of a small set of neurons (WU *et al.* 2001), but we would not know if deficiency of WTS-1 or CST-
1/2 impacted nuclear translocation of YAP-1 during a key developmental window in the life of these
neurons.
407
Functional outcomes without proliferation. A striking feature of the *C. elegans* Hippo pathway is the
absence of proliferative mutant phenotypes. In other animals, Hippo signaling governs tissue size,
regeneration, and tumorigenesis; in nematodes, cell lineages and organ sizes, and indeed animal size,

is mostly fixed. Our findings suggest that while the molecular wiring is conserved, the transcriptional

outputs of YAP-1/EGL-44 may have shifted. Many YAP/TEAD target genes in *C. elegans* are linked to

pathogen responses and gut barrier integrity (MA *et al.* 2020). We hypothesize that YAP-1/EGL-44 has

been uncoupled from growth-control genes – either through changes in regulatory DNA elements or

altered DNA-binding specificity of EGL-44 – and thus repurposed to support stress and barrier functions.

In this sense, *C. elegans* provides a model for Hippo signaling stripped of its growth-control role,

emphasizing instead its contributions to tissue robustness.

Interpreting the arrest phenotype

Loss of WTS-1 causes developmental arrest accompanied by defective intestinal integrity. This arrest

could reflect the requirement of an EGL-44/TEAD target gene expression for developmental progression,

while other genes are required for tissue integrity. Alternatively, perhaps arrest is a secondary

consequence of developmental defects, say perhaps a developmental checkpoint in response to loss of

gut integrity, or an intestinal defect that results in starvation. Analogies may be drawn to the mTORC1

pathway, where stress of reduced anabolism can cause programmed diapause at a variety of stages

(DUONG *et al.* 2020; SUN *et al.* 2025). Unexpectedly, hyperactivation of YAP-1/EGL-44 upon loss of WTS-

1 reduces, rather than enhances, robustness of gut integrity. This finding contrasts with the presumed

protective induction of YAP/TEAD in response to pathogenesis/barrier disruption (MA *et al.* 2020),

suggesting a feedback loop in which excessive pathway activity destabilizes epithelial function. As with

Hippo-mediated growth in other systems, balanced homeostasis may be critical in the gut.

Tissue- and process-specific roles. Our work complements previous studies showing that RNAi

depletion of *cst-1/2* induces autophagy and influences lifespan (LEHTINEN *et al.* 2006; WILKINSON *et al.*

2015), phenotypes not easily explained through WTS-1 or YAP-1. Similarly, Hippo-related roles in

neuronal development have been reported in *C. elegans*, unattached to WTS-1 or YAP-1/EGL-44 (FENG

et al. 2017; JOSEPHSON *et al.* 2017; LEE *et al.* 2018). While our WTS-1 null animals did not exhibit

locomotion defects, *cst-1/2* deletions were Uncoordinated and slightly undergrown, suggesting that CST-

1/2 may act independently of WTS-1 in sculpting the nervous system. Future experiments will be needed

to determine whether YAP-1 or EGL-44 contribute to these locomotory phenotypes.

Contributions of this work. Our results extend prior studies in several important ways. First, we used a

novel endogenous YAP-1 reporter to directly monitor Hippo activity *in vivo*, validating WTS-1-dependent

repression across two tissues. Second, we employed tissue-specific depletion strategies to show that

WTS-1 regulation is cell-autonomous. Third, we clarified the contributions of CST-1/2, identifying them

as functional Hippo-like kinases in the epithelium but perhaps not intestine. Finally, we assessed functions

of Ste20-family orthologs, showing that GCK-2 does not contribute to YAP-1 regulation while MIG-15

influences YAP-1 stability but not localization. Together, these analyses position *C. elegans* as a unique

model in which Hippo signaling is preserved at the level of WTS-1 inhibition of YAP-1 but diverges in its

upstream regulation and biological outputs.

Broader implications. The field of Hippo signaling has been dominated by studies of over-proliferation,

regeneration and tumorigenesis. *C. elegans* provides a contrasting model in which conserved pathway

components are retained but uncoupled from growth control. Instead, Hippo signaling contributes to

epithelial robustness, stress responses, and possibly neural development. These findings underscore the

evolutionary flexibility of Hippo signaling and open new directions for understanding how signaling

pathways are rewired to meet the developmental logic of different organisms.

METHODS

***C. elegans* handling and genetics.** All *C. elegans* strains were derived from the N2 Bristol wild type and

grown on NG agar plates seeded with the OP50 strain *E. coli* at 20˚C unless otherwise noted. Growth,

culturing and nomenclature were generally as described (BRENNER 1974; HORVITZ *et al.* 1979; TULI *et al.*

2018). Wormbase was used extensively during these studies (STERNBERG *et al.* 2024).

CRISPR/Cas9 mediated genome editing. Guide RNAs were that maximally satisfied a combination of

three approaches. First, where possible, G and not T nucleotides were selected at positions -1/-2, or

GCGG and not T-TT nucleotides at positions -1/-2/-3/-4 (KAPOOR *et al.* 2014; FARBOUD AND MEYER 2015).

Second, we considered strong predicted specificity and efficiency scores using the CRISPOR

(http://crispor.tefor.net/) algorithm, which incorporates the original MIT specificity score. Third, we

considered strong predicted efficiency scores using the WU-CRISPR (http://crisprdb.org/wu-crispr/)

algorithm.

We formulated mixes for microinjection as described previously (FAKIEH *et al.* 2022). Using a

dedicated RNase-free bench, we mixed concentrations as final volume per 20 µL: 1 µl of 5 µg/µl stock

Streptococcus pyogenes Cas9 (PNAbio, #CP01) for 0.25 µg/µl final concentration; 1µl of 2 µg/µl stock

universal tracrRNA (IDT) for 0.1 µg/µl final concentration; 1.4 µl each of 0.4 µg/µl stocks of *dpy-10* co-

CRISPR and gene-specific crRNAs for 0.028 µg/µl crRNA final concentration each. Not yet at final

volume, this mix was incubated at 37°C for 15 min. After incubation, *dpy-10* ssODN repair oligo was

added as 3.3 µl of 20 µM stock for final concentration of 3.3 µM; gene-specific column-purified PCR repair

template was denatured and renatured as described (GHANTA AND MELLO 2020), and added from stock

of ≤300 ng/µl to a final concentration of 100 ng/µl (total final concentration of PCR product no greater

than 2 µg/20 µl for reasons of viscosity). As this point, commercial nuclease-free water was added up to

20 µl. (Concentrations before the incubation were calculated per 20 µl, but 20 µl volume was not reached

until the final addition of water.)

Adult animals with no more than a single row of embryos in each uterus side were microinjected as

described (KADANDALE *et al.* 2009). We used co-CRISPR to generate dominant Rol mutations in F1

animals harboring the *dpy-10(cn64*gf*)* marker (ARRIBERE *et al.* 2014), which were picked singly or in pools

of two to plates. After F1 Rols had laid embryos, the single or pooled parents were picked to PCR tubes,

lysed for PCR and detected via triplex PCR. Non-Dpy/Rol F2s were picked singly, incubated overnight to

obtain genetic material, lysed and PCR detected to generate homozygous lines. Knockins were

sequenced from points in flanking DNA outside that subjected to homology-directed recombination repair.

Only expected knockins were used for analysis.

RNA interference. Plates for bacterially mediated RNAi interference (RNAi) were made with NGM agar

supplemented with 1 mM IPTG and 50 µg/mL carbenicillin. Plates were spotted with 80 µl of HT115

bacteria culture expressing dsRNA as described (TIMMONS *et al.* 2001). L4 P0s were picked to 24 hr-

seeded RNAi plates, grown overnight, then transferred the next day to fresh RNAi plates for synchronized

laying of embryos. The P0s were removed 24 hours later and F1 animals were analyzed at the time

appropriate for the experiments. Positive control for efficacy of RNAi plates was *pop-1(RNAi),* where we

expect 95-100% embryonic or L1 lethality of the experiment is discarded. For negative control with used

luciferase(RNAi) as previously described (SHIN *et al.* 2018), to engage the RNAi machinery with a

sequence not encoded in the *C. elegans* genome. RNAi was performed at 20˚C.

Pumping assays. Benchtop mechanical clickers were uses to count contractions of the pharynx

("pumping") per 60 sec. Observations used a Nikon SMZ18 stereo microscope.

Auxin treatment. Auxin (IAA; Indole-3-acetic acid, 98+%, Thermo Scientific) storage stock solution was

prepared at 400 mM in ethanol and stored up to a month at 4˚C. For experiments, stock auxin was diluted

to 16 mM working auxin stock by diluting storage 400 mM auxin with filtered Milli-Q water. For final

concentration of 1 mM auxin in plates, 500 µl of 16 mM auxin was added onto 8 ml NGM agar plates or

RNAi plates and allowed to diffuse. For vehicle control, 500 µl of 4% ethanol was similarly added to

plates.

Detection of tagged endogenous YAP-1::mNG::2xFLAG. Mixed-stage animals were washed from 3-

4 full but not starved plates using M9 buffer and lysed in 4% SDS loading buffer with boiling at 90˚C for

2 min. Samples were run on a 4-15% SDS-PAGE gel (Bio-Rad, #5671084) and transferred to Immobilon-

P Membrane, PVDF (EMD Millipore, IPVH00010). Mouse monoclonal anti-Flag antibody (Sigma-Aldrich,

F1804) and mouse monoclonal anti-α-tubulin antibody (Sigma-Aldrich, T6199) were diluted 1:2000 in

blocking solution (6% w/v non-fat dry milk in PBST). HRP conjugated goat anti-mouse secondary

antibody (Sigma-Aldrich, 12-349) was diluted 1:5000 in blocking solution. Chemiluminescent detection

was performed using ECL reaction (Thermo Fisher Scientific 32106) and detected via film processor,

SRX-101A (Konica Minolta) on X-ray film (Phenix).

Imaging. Animals used for live imaging were mounted on a 5% agar pad on a glass slide in 5 μl of 2

mg/ml tetramisole in M9 buffer with cover slip. Confocal microscopy used a Nikon Ti2 microscope with a

Yokogawa CSU-W1 Spinning Disk, 405 nm, 445 nm, 514 nm lasers, and a Photometrics Prime BSI

camera, and NIS Elements Advanced Research software, version 5.42. Quantification of fluorescence

intensity were performed by ImageJ (Fiji) with a customized script code (deposited at:

https://github.com/ReidTPowell/2025_Genetics_Reiner). In 3-channel confocal

photomicrographs analyzed in ImageJ with the script running, the user identifies the center of a

nucleus in the DIC channel and placed a point using the "multi-point tool." The script draws a

uniform circle for each point of interest and measures fluorescence intensity. This confers

precision in area of circle from sample to sample.

10 cm NG plate photomicrographs were captured using a Nikon Eclipse Ni microscope, 4x objective,

ANDOR Zyla sCMOS camera and NIS Elements Advanced Research software, version 4.30, with plate

and animals on the stage with sufficient working distance.

Sequence analysis. Protein sequences were accessed via Wormbase (wormbase.org; (STERNBERG *et*

al. 2024) and Alliance of Genome Resources (https://www.alliancegenome.org/). We used the A isoform

for both *cst-1* and *cst-2*, thus excluding the two-codon minority splice addition to exon 9 for both. *mig-15*

isoform A was selected as the reference and most abundant isoform. Uniprot accession numbers were

Q13188 for human MST1/STK3, Q13043 for human MST2/STK4, Q8T0S6 for *Drosophila* Hpo. Protein

sequences were aligned with Clustal Omega (https://www.ebi.ac.uk/jdispatcher/msa/).

Statistical analysis. Animal experiments presented in the same graph were scored concurrently.

Statistical analyses were performed by two-tail unpaired Student's t-test or Fisher's Exact test (GraphPad

Prism 10).

Software/website

GraphPad Prism was used for statistical analysis and making graphs. BioRender

(https://www.biorender.com/) was used to generate illustrations. Exon-Intron Graphic Maker was used for

making schematic gene structures http://wormweb.org/exonintron. AI software was used only for text

editing purposes.

Data availability statement.

Strains, plasmids and sequence files are available upon request. All data supporting the conclusions of

the study and required for reproduction of experiments are present within the text, tables and figures.

Conflicts of interest.

None.

Acknowledgements. We thank L. Vergara in the Center for Advanced Imaging for technical assistance,

members of the Reiner lab for helpful discussions and technical expertise, J. Bembenek (Wayne State

University) for the strain harboring HIS-72::mT2, M. Bastiani (University of Utah) for *basDf1*, Y. Hata

(Tokyo Medical and Dental University) and B. Grant (Rutgers University) for the strain harboring

*ihIs35[yap-1*p>*yap-1::GFP], and* S. Mitani (Japanese National Bioresource Project) for *wts-1(tm4081)*.

Some strains were provided by the *Caenorhabditis* Genetics Center, which is funded by the NIH Office

of Research Infrastructure Programs (P40 OD010440). Wormbase was used extensively. Our CRISPR

protocol was based on an amalgam protocol generously shared by Dr. Andy Golden (NIDDK).

Funding. This work was funded by NIH grants R35GM144237 and R03CA289854 to D.J.R.

Supplementary Tables.

Strains used are shown in Table S1.

Oligonucleotides in Table S2.

Guide RNAs in Table S3.

Plasmids in Table S4.

Figure Legends

Figure 1. Loss of WTS-1 confers YAP-1/YAP and EGL-44/TEAD-dependent developmental arrest.

(A) Schematic of experimental design for scoring *wts-1* mutant developmental stages. **(B)** Quantification

of arrest stages of homozygous *wts-1* mutant compared to the heterozygote on Day 3. △1 = *wts-*

1(tm4081). △2 = *wts-1(ok753)*. * = *wts-1(re436[STOP-IN*]). **(C)** Pumping rate/60s of *wts-1(tm4081)*

homozygote vs. heterozygote. **(D)** Schematic of protocol for auxin treatment of **E-H)** *wts-*

1(re419[mTurq2::2xMyc::AID*::wts-1]); *ieSi57[eft-3p::TIR1::mRuby]* animals grown on vehicle or 1 mM

auxin, scored on Day 3 or Day 4. **I)** Quantification of developmental stage for animals in **E-H**. **(J-M)** The

same animals as above grown on vehicle or 1 mM auxin and *ctrl(RNAi)* vs *wts-1(RNAi)*, scored on Day

3 or Day 4. *ctrl(RNAi)* = luciferase sequences with no homology in the *C. elegans* genome (Shin *et al.*

2018). **(N)** Quantification of developmental stage for animals in **J-M**. **(O)** Quantification of developmental

stages on Day 3 of *wts-1(tm4081)* treated with *ctrl(RNAi)*, *yap-1(RNAi)* or *egl-44(RNAi)*. ****=P<0.0001,

***=P<0.001, **=P<0.01, ns = not significant (Fisher's exact test).

Figure 2: YAP-1::mNG functions as a reporter for WTS-1 activity in epithelium and intestine. (A-C)

Confocal and DIC photomicrographs and quantification of lateral epithelium; animals are oriented side-

up. Animal genotype is *rrf-3(re390[STOP-IN])*; *his-72(erb77[his-72::linker::mTurquoise2])*; *yap-*

1(re269[yap-1::mNG::2xflag]) to confer hypersensitivity to RNAi. **(A-B")** Cytoplasm-to-nuclear

translocation of endogenous YAP-1::mNG grown on *ctrl(RNAi)* vs. *wts-1(RNAi)* bacteria, respectively. **(C)**

Quantification of fluorescence intensity in **A-B"** of nuclear YAP-1::mNG in arbitrary units (A.U.). N =

multiple nuclei from 9 and 10 animals, respectively. **(D-F)** Confocal and DIC photomicrographs and

quantification of intestine, left-right mid-animal plane; animals are oriented side-up. Animals are depicted

at lower magnification to show multiple intestinal nuclei. **(D-E")** Cytoplasm-to-nuclear translocation of

endogenous YAP-1::mNG grown on *ctrl(RNAi)* vs. *wts-1(RNAi)* bacteria. Animal genotype is *rrf-*

3(re390[STOP-IN]); *his-72(erb77[his-72::linker::mTurquoise2])*; *yap-1(re269[yap-1::mNG::2xflag])* to

confer hypersensitivity to RNAi. **(F)** Quantification of fluorescence intensity in **D-E"** of nuclear YAP-

1::mNG as A.U.. N = multiple nuclei from 4 and 5 animals, respectively. Epithelium scale bars = 20 μm,

pop out scale bars for intestine = 10 μm. **** = P<0.0001 (*t*-test).

Figure 3. Effects of tissue-specific depletion of AID*::WTS-1. (A,B) Bright field photomicrographs of

animals with AID*::WTS-1 depletion: global RNAi with epithelial-specific AID* depletion via epithelial-

specific expression of TIR1. Genotype is *wts-1(re419[mT2::2xMyc::AID*::wts-1])*; *reSi2[col-*

10p>TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR]. **(A)** Animals grown on vehicle+*ctrl(RNAi)* vs. **(B)**

auxin+*wts-1(RNAi)*. **(C)** Quantification of **(A** vs. **B)**. **(D,E)** Bright field photomicrographs of animals

AID*::WTS-1 depletion: global RNAi with intestinal-specific AID* depletion via intestinal-specific

expression of TIR1. Genotype is *wts-1(re419[mT2::2xMyc::AID*::wts-1])*; *reSi12[ges-*

1p>TIR1::F2A::mTagBFP2::AID::NLS::tbb-2 3'UTR]. **(D)** Animals grown on vehicle+*ctrl(RNAi)* vs. **(E)**

auxin+*wts-1(RNAi)*. **(F)** Quantification of **(D** vs. **E)**. Scale bars = 1 mm. ****=P<0.0001 (Fisher's exact

test). **(G-N')** Effects on YAP-1::mNG localization of depletion of WTS-1 in epithelium vs. intestine. **(G-J')**

Confocal and DIC photomicrographs of epithelium (**G-G",I-I"**) vs. intestine (**H-H',J-J'**) with YAP-1::mNG,

epithelial-specific TIR1+AID*::BFP and AID*::WTS-1, no RNAi. Genotype is *wts-*

1(re419[mT2::2xMyc::AID*::wts-1]); *reSi2[col-10p>TIR1::F2A::mTagBFP2::AI*D::NLS::tbb-2 3'UTR]*;

yap-1(re269[yap-1::mNG::2xFlag]). **(G-H')** vehicle-treated animal vs. **(I-J')** auxin-treated. **(G' vs. I')** Note

auxin-dependent depletion of nuclear BFP internal control in epithelium. **(K-N')** Confocal and DIC

photomicrographs of intestine (**K-K",M-M"**) vs epithelium (**L-L',N-N'**) with YAP-1::mNG, intestinal-

specific TIR1+AID*::BFP and AID*::WTS-1, no RNAi. Genotype is *wts-1(re419[mT2::2xMyc::AID*::wts-*

1]); *reSi12[ges-1p>TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR]*; *yap-1(re269[yap-1::mNG::2xFlag])*.

(K-L') vehicle-treated animal vs. **(M-N')** auxin-treated. **(K' vs. M')** Note auxin-dependent depletion o

nuclear BFP internal control in intestine. Scale bars in epithelial confocal images: 20 µm. Scale bars in

intestinal confocal images = 10 µm.

Figure 4: The CST-1/2 putative Hippo orthologs inhibit YAP-1::mNG in epithelium but not intestine.

(A) Schematic of *cst-1* and *cst-2* genomic region on Chromosome X. The red line indicates the

reDf4[re452] deletion, which removes *cst-1*, *cst-2* and seven ncRNA genes in between. Lines above the

chromosome map indicate conserved protein domains from protein products. CST-2 lacks the C-terminal

SARAH (<u>S</u>av/<u>RASS</u>F/<u>H</u>po) coiled-coiled dimerization domain present in the C-terminus of *cst-1* and

Hippo proteins in other metazoans. **(B-D)** Confocal photomicrographs, DIC photomicrographs and

quantification of lateral epithelium (animals are oriented side-up). Animal genotypes are *i.e.* wild type vs.

cst-1/2 deletion: *his-72(erb77[his-72::linker::mT2])*; *yap-1(re269[yap-1::mNG::2xflag]) vs. his-*

72(erb77[his-72::linker::mT2]); *cst-1/2(reDf4[re452]) yap-1(re269[yap-1::mNG::2xflag])*. **D)** Quantification

of fluorescence intensity in **B-C"** of nuclear YAP-1::mNG in arbitrary units (A.U.). N = multiple nuclei from

9 and 10 animals, respectively. **(E-G)** Confocal photomicrographs, DIC photomicrographs, and

quantification of intestine, left-right mid-animal intestinal plane (animals are oriented side-up). Animals

are depicted at lower magnification to show multiple intestinal nuclei. Animal genotypes are *his-*

72(erb77[his-72::linker::mT2]); *yap-1(re269[yap-1::mNG::2xflag])* vs. *his-72(erb77[his-72::linker::mT2])*;

cst-1/2(reDf4[re452]) yap-1(re269[yap-1::mNG::2xflag]), *i.e.* wild type vs. *cst-1/2* deletion. N = multiple

nuclei from 4 animals each. Epithelium scale bars = 20 µm, pop out scale bars = 10 µm. ****=P<0.0001,

ns = not significant (*t*-test)

Figure 5. CST-1/2 functions redundantly with MIG-15 to control developmental progression. (A-

H) Bright-field photomicrographs of depleted AID*::MIG-15 with or without deleted *cst-1/2*, scored on day

3 (**A,B,E,F**) and day 4 (**C,D,G,H**) to illustrate degree of growth delay/arrest. Genotypes are *ieSi57[eft-*

3p>TIR1::mRuby::unc-54 3'UTR + Cbr-unc-119(+)]; unc-119(ed3); mig-15(re264[AID*::mNG::2xHA::mig-

15]) (**A-D**)*vs. ieSi57[eft-3p>TIR1::mRuby::unc-54 3'UTR + Cbr-unc-119(+)]; unc-119(ed3); mig-*

15(re264[AID*::mNG::2x::mig-15]) cst-1/2(reDf1[re484]) (**E-H**). (**B,D** and **F,H**) Treatment with 1 mM auxin

vs. vehicle on other plates. Depletion of *AID*::MIG-15* causes mild growth delay and locomotion defects

(**A/C** vs **B/D**), as published for *mig-15* mutants. Deletion of *cst-1/2* causes mild growth delay and

locomotion defects (**A/C** vs **E/G**), as we observed on plates mutant for *basDf1* or *cst-1/2(reDf1[re484])*

(**I,J**) Quantification of arrest at Day 3 and Day 4, respectively, reveals the strong arrest of *cst-1/2 mig-15*

double mutants relative to single mutants. Animal assays shown in (**E,F,G,H**) and their quantification in

(**I,J**), in columns with N of 143 and 135, respectively, are recapitulated in **Fig. S5**, to contrast the effects

with and without mutant *gck-2*. All were scored concurrently. (**K**) Pumping rate of control *vs.* double

mutant on Day 1 and Day 3 indicates arrest, not lethality. N=10 animals for each group. Animals also

continue to swim. (**L-N**) Growth arrest phenotype of AID*::MIG-15 depletion combined with *cst-1/2*

deletion was reversed by *yap-1(RNAi)* or *egl-44(RNAi)*. (**O**) quantification of **L-N**. ****=P<0.0001,

***=P<0.001, **=P<0.01, ns= not significant. Scale bar = 1 mm.

Figure 6. MIG-15 represses YAP-1 levels but does not restrict nuclear translocation. (A-B')

Confocal and DIC photomicrographs of YAP-1::mNG in lateral epithelium. Animal genotype is *ieSi57[eft-*

3p>TIR1::mRuby::unc-54   3'UTR   +   Cbr-unc-119(+)];   unc-119(ed3);   mig-

15(re435[mT2::2xMyc::AID::mig-15)   yap-1(re269[yap-1::mNG::2xflag])* on vehicle *vs.* auxin,

respectively. (**C**) Quantification of YAP-1::mNG intensity in AID*::MIG-15 animals with and without auxin.

N = multiple nuclei from 10 and 9 animals, respectively. (**D-E'**) Confocal and confocal photomicrographs

of YAP-1 in midline intestine. Animal genotype is *ieSi57[eft-3p>TIR1::mRuby::unc-54 3'UTR + Cbr-unc-*

119(+)]; unc-119(ed3); mig-15(re435[mT2::2xMyc::AID::mig-15) yap-1(re269[yap-1::mNG::2xflag])* on

vehicle vs. auxin. Fluorescence intensity was measured for nucleus (**F'**) and cytoplasm (**G'**), respectively.

N = multiple nuclei from 9, 10 animals, respectively.(**H-J'**) Confocal and confocal photomicrographs of

YAP-1 in midline intestine. Animal genotype is *ieSi57[eft-3p::TIR1::mRuby::unc-54 3'UTR + Cbr-unc-*

119(+)]; unc-119(ed3); mig-15(re435[mT2::2xMyc::AID::mig-15) yap-1(re269[yap-1::mNG::2xflag]) cst-*

1/2(re479) on vehicle vs. auxin. Fluorescence intensity was measured for nucleus (**H'**) and cytoplasm

(**I'**), respectively. N=10-18 animals. Scale bars = 10 µm. ****=P<0.0001, ***=P<0.001. (*t*-test)

Figure 7. A model for Hippo and Warts signaling in *C. elegans*. (A) In epithelium a canonical Hippo

pathway functions, with potentially redundant CST-1 and CST-2 phosphorylating and activating WTS-1.

WTS-1 in turn phosphorylates and inhibits YAP-1, keeping it out of the nucleus. Disruption of CST-1/2 or

WTS-1 results in nuclear translocation of YAP-1, as observed in other systems. Rather than contributing

to activation of WTS-1 to repress YAP-1, MIG-15 inhibits YAP-1 protein levels. (**B**) In Intestine a non-

canonical Hippo pathway functions. Unknown upstream inputs regulate WTS-1 to phosphorylate and

retain YAP-1 out of the nucleus. Loss of WTS-1 causes nuclear translocation of YAP-1 but also L2 arrest,

indicated as "Growth." CST-1/2 and MIG-15 act redundantly to maintain growth.

REFERENCES

Apken, L. H., and A. Oeckinghaus, 2021 The RAL signaling network: Cancer and beyond. Int Rev Cell Mol Biol 361**:** 21-105.

Arribere, J. A., R. T. Bell, B. X. Fu, K. L. Artiles, P. S. Hartman *et al.*, 2014 Efficient marker-free recovery of custom genetic modifications with CRISPR/Cas9 in Caenorhabditis elegans. Genetics 198**:** 837-846.

Ashley, G. E., T. Duong, M. T. Levenson, M. A. Q. Martinez, L. C. Johnson *et al.*, 2021 An expanded auxin-inducible degron toolkit for Caenorhabditis elegans. Genetics 217.

Brenner, S., 1974 The genetics of Caenorhabditis elegans. Genetics 77**:** 71-94.

Burglin, T. R., and P. E. Kuwabara, 2006 Homologs of the Hh signalling network in C. elegans. WormBook**:** 1-14.

Cai, Q., W. Wang, Y. Gao, Y. Yang, Z. Zhu *et al.*, 2009 Ce-wts-1 plays important roles in Caenorhabditis elegans development. FEBS Lett 583**:** 3158-3164.

Dan, I., N. M. Watanabe and A. Kusumi, 2001 The Ste20 group kinases as regulators of MAP kinase cascades. Trends Cell Biol 11**:** 220-230.

Dasgupta, I., and D. McCollum, 2019 Control of cellular responses to mechanical cues through YAP/TAZ regulation. J Biol Chem 294**:** 17693-17706.

Dejima, K., S. Hori, S. Iwata, Y. Suehiro, S. Yoshina *et al.*, 2018 An Aneuploidy-Free and Structurally Defined Balancer Chromosome Toolkit for Caenorhabditis elegans. Cell Rep 22**:** 232-241.

Delpire, E., 2009 The mammalian family of sterile 20p-like protein kinases. Pflugers Arch 458**:** 953-967.

Duong, T., N. R. Rasmussen, E. Ballato, F. S. Mote and D. J. Reiner, 2020 The Rheb-TORC1 signaling axis functions as a developmental checkpoint. Development 147.

Fakieh, R., T. Duong, Y. Wu, N. Rasmussen and D. Reiner, 2022 A simple strategy for addition of degron tags to endogenous genes harboring prior insertions of fluorescent protein. MicroPubl Biol 2022.

Fakieh, R. A., and D. J. Reiner, 2025 RAP-2 and CNH-MAP4 Kinase MIG-15 confer resistance in bystander epithelium to cell-fate transformation by excess Ras or Notch activity. Proc Natl Acad Sci U S A 122**:** e2414321121.

Farboud, B., and B. J. Meyer, 2015 Dramatic enhancement of genome editing by CRISPR/Cas9 through improved guide RNA design. Genetics 199**:** 959-971.

Feng, G., Z. Zhu, W. J. Li, Q. Lin, Y. Chai *et al.*, 2017 Hippo kinases maintain polarity during directional cell migration in Caenorhabditis elegans. EMBO J 36**:** 334-345.

Frisch, S. M., M. Schaller and B. Cieply, 2013 Mechanisms that link the oncogenic epithelial-mesenchymal transition to suppression of anoikis. J Cell Sci 126**:** 21-29.

Frokjaer-Jensen, C., M. W. Davis, G. Hollopeter, J. Taylor, T. W. Harris *et al.*, 2010 Targeted gene deletions in C. elegans using transposon excision. Nat Methods 7**:** 451-453.

Ghanta, K. S., and C. C. Mello, 2020 Melting dsDNA Donor Molecules Greatly Improves Precision Genome Editing in Caenorhabditis elegans. Genetics 216**:** 643-650.

Halder, G., and R. L. Johnson, 2011 Hippo signaling: growth control and beyond. Development 138**:** 9-22.

Han, H., Z. Huang, C. Xu, G. Seo, J. An *et al.*, 2024 Functional annotation of the Hippo pathway somatic mutations in human cancers. Nat Commun 15**:** 10106.

Hansen, C. G., T. Moroishi and K. L. Guan, 2015 YAP and TAZ: a nexus for Hippo signaling and beyond. Trends Cell Biol 25**:** 499-513.

Hariharan, I. K., 2015 Organ Size Control: Lessons from Drosophila. Dev Cell 34**:** 255-265.

Harvey, K. F., C. M. Pfleger and I. K. Hariharan, 2003 The Drosophila Mst ortholog, hippo, restricts growth and cell proliferation and promotes apoptosis. Cell 114**:** 457-467.

Harvey, K. F., and T. T. Tang, 2025 Targeting the Hippo pathway in cancer. Nat Rev Drug Discov.

Hilman, D., and U. Gat, 2011 The evolutionary history of YAP and the hippo/YAP pathway. Mol Biol Evol 28**:** 2403-2417.

Horvitz, H. R., S. Brenner, J. Hodgkin and R. K. Herman, 1979 A uniform genetic nomenclature for the nematode Caenorhabditis elegans. Mol Gen Genet 175**:** 129-133.

Huang, J., S. Wu, J. Barrera, K. Matthews and D. Pan, 2005 The Hippo signaling pathway coordinately
regulates cell proliferation and apoptosis by inactivating Yorkie, the Drosophila Homolog of YAP.
Cell 122**:** 421-434.

Iwasa, H., S. Maimaiti, H. Kuroyanagi, S. Kawano, K. Inami *et al.*, 2013 Yes-associated protein homolog,
YAP-1, is involved in the thermotolerance and aging in the nematode Caenorhabditis elegans.
Exp Cell Res 319**:** 931-945.

Josephson, M. P., R. Aliani, M. L. Norris, M. E. Ochs, M. Gujar *et al.*, 2017 The Caenorhabditis elegans
NF2/Merlin Molecule NFM-1 Nonautonomously Regulates Neuroblast Migration and Interacts
Genetically with the Guidance Cue SLT-1/Slit. Genetics 205**:** 737-748.

Justice, R. W., O. Zilian, D. F. Woods, M. Noll and P. J. Bryant, 1995 The Drosophila tumor suppressor
gene warts encodes a homolog of human myotonic dystrophy kinase and is required for the
control of cell shape and proliferation. Genes Dev 9**:** 534-546.

Kadandale, P., I. Chatterjee and A. Singson, 2009 Germline transformation of Caenorhabditis elegans by
injection. Methods Mol Biol 518**:** 123-133.

Kang, J., D. Shin, J. R. Yu and J. Lee, 2009 Lats kinase is involved in the intestinal apical membrane
integrity in the nematode Caenorhabditis elegans. Development 136**:** 2705-2715.

Kapoor, A., W. Yao, H. Ying, S. Hua, A. Liewen *et al.*, 2014 Yap1 activation enables bypass of oncogenic
Kras addiction in pancreatic cancer. Cell 158**:** 185-197.

Karchugina, S., D. Benton and J. Chernoff, 2021 Regulation of MST complexes and activity via SARAH
domain modifications. Biochem Soc Trans 49**:** 675-683.

Lee, H., J. Kang, S. Ahn and J. Lee, 2019 The Hippo Pathway Is Essential for Maintenance of Apicobasal
Polarity in the Growing Intestine of Caenorhabditis elegans. Genetics 213**:** 501-515.

Lee, H., J. Kang and J. Lee, 2018 Involvement of YAP-1, the Homolog of Yes-Associated Protein, in the
Wnt-Mediated Neuronal Polarization in Caenorhabditis elegans. G3 (Bethesda) 8**:** 2595-2602.

Lehtinen, M. K., Z. Yuan, P. R. Boag, Y. Yang, J. Villen *et al.*, 2006 A conserved MST-FOXO signaling
pathway mediates oxidative-stress responses and extends life span. Cell 125**:** 987-1001.

Lin, W. H., L. M. Cooper and P. Z. Anastasiadis, 2023 Cadherins and catenins in cancer: connecting
cancer pathways and tumor microenvironment. Front Cell Dev Biol 11**:** 1137013.

Ma, Y. C., Z. S. Yang, L. Q. Ma, R. Shu, C. G. Zou *et al.*, 2020 YAP in epithelium senses gut barrier loss
to deploy defenses against pathogens. PLoS Pathog 16**:** e1008766.

Martin-Belmonte, F., and M. Perez-Moreno, 2011 Epithelial cell polarity, stem cells and cancer. Nat Rev
Cancer 12**:** 23-38.

Meng, Z., T. Moroishi, V. Mottier-Pavie, S. W. Plouffe, C. G. Hansen *et al.*, 2015 MAP4K family kinases
act in parallel to MST1/2 to activate LATS1/2 in the Hippo pathway. Nat Commun 6**:** 8357.

Mesrouze, Y., F. Bokhovchuk, M. Meyerhofer, C. Zimmermann, P. Fontana *et al.*, 2021 Study of the TEAD-
binding domain of the YAP protein from animal species. Protein Sci 30**:** 339-349.

Moroishi, T., C. G. Hansen and K. L. Guan, 2015 The emerging roles of YAP and TAZ in cancer. Nat Rev
Cancer 15**:** 73-79.

Oh, H., and K. D. Irvine, 2010 Yorkie: the final destination of Hippo signaling. Trends Cell Biol 20**:** 410-
417.

Pearson, J. D., K. Huang, M. Pacal, S. R. McCurdy, S. Lu *et al.*, 2021 Binary pan-cancer classes with
distinct vulnerabilities defined by pro- or anti-cancer YAP/TEAD activity. Cancer Cell 39**:** 1115-
1134 e1112.

Rothman, J., and S. Jarriault, 2019 Developmental Plasticity and Cellular Reprogramming in
Caenorhabditis elegans. Genetics 213**:** 723-757.

Shaner, N. C., G. G. Lambert, A. Chammas, Y. Ni, P. J. Cranfill *et al.*, 2013 A bright monomeric green
fluorescent protein derived from Branchiostoma lanceolatum. Nat Methods 10**:** 407-409.

Shin, H., R. E. W. Kaplan, T. Duong, R. Fakieh and D. J. Reiner, 2018 Ral Signals through a MAP4
Kinase-p38 MAP Kinase Cascade in C. elegans Cell Fate Patterning. Cell Rep 24**:** 2669-2681
e2665.

Simmer, F., M. Tijsterman, S. Parrish, S. P. Koushika, M. L. Nonet *et al.*, 2002 Loss of the putative RNA-
directed RNA polymerase RRF-3 makes C. elegans hypersensitive to RNAi. Curr Biol 12**:** 1317-
1319.
Sternberg, P. W., K. Van Auken, Q. Wang, A. Wright, K. Yook *et al.*, 2024 WormBase 2024: status and
transitioning to Alliance infrastructure. Genetics 227.
Sun, C. L., C. Xu, O. Itani, E. L. Christensen, H. Vijay *et al.*, 2025 Biased regulation of protein synthesis
and hypoxic death by a conditional raptor mutation. Curr Biol 35**:** 2567-2582 e2565.
Tapon, N., K. F. Harvey, D. W. Bell, D. C. Wahrer, T. A. Schiripo *et al.*, 2002 salvador Promotes both cell
cycle exit and apoptosis in Drosophila and is mutated in human cancer cell lines. Cell 110**:** 467-
478.
Timmons, L., D. L. Court and A. Fire, 2001 Ingestion of bacterially expressed dsRNAs can produce
specific and potent genetic interference in Caenorhabditis elegans. Gene 263**:** 103-112.
Tuli, M. A., A. Daul and T. Schedl, 2018 Caenorhabditis nomenclature. WormBook 2018**:** 1-14.
Vassilev, A., K. J. Kaneko, H. Shu, Y. Zhao and M. L. DePamphilis, 2001 TEAD/TEF transcription factors
utilize the activation domain of YAP65, a Src/Yes-associated protein localized in the cytoplasm.
Genes Dev 15**:** 1229-1241.
Wang, H., H. Park, J. Liu and P. W. Sternberg, 2018 An Efficient Genome Editing Strategy To Generate
Putative Null Mutants in Caenorhabditis elegans Using CRISPR/Cas9. G3 (Bethesda) 8**:** 3607-
3616.
Wilkinson, D. S., J. S. Jariwala, E. Anderson, K. Mitra, J. Meisenhelder *et al.*, 2015 Phosphorylation of
LC3 by the Hippo kinases STK3/STK4 is essential for autophagy. Mol Cell 57**:** 55-68.
Wu, J., A. Duggan and M. Chalfie, 2001 Inhibition of touch cell fate by egl-44 and egl-46 in C. elegans.
Genes Dev 15**:** 789-802.
Wu, S., J. Huang, J. Dong and D. Pan, 2003 hippo encodes a Ste-20 family protein kinase that restricts
cell proliferation and promotes apoptosis in conjunction with salvador and warts. Cell 114**:** 445-
456.
Wu, S., Y. Liu, Y. Zheng, J. Dong and D. Pan, 2008 The TEAD/TEF family protein Scalloped mediates
transcriptional output of the Hippo growth-regulatory pathway. Dev Cell 14**:** 388-398.
Yang, Z., and Y. Hata, 2013 What is the Hippo pathway? Is the Hippo pathway conserved in
Caenorhabditis elegans? J Biochem 154**:** 207-209.
Zhang, L., F. Ren, Q. Zhang, Y. Chen, B. Wang *et al.*, 2008 The TEAD/TEF family of transcription factor
Scalloped mediates Hippo signaling in organ size control. Dev Cell 14**:** 377-387.
Zhang, L., J. D. Ward, Z. Cheng and A. F. Dernburg, 2015 The auxin-inducible degradation (AID) system
enables versatile conditional protein depletion in C. elegans. Development 142**:** 4374-4384.
Zheng, Y., W. Wang, B. Liu, H. Deng, E. Uster *et al.*, 2015 Identification of Happyhour/MAP4K as
Alternative Hpo/Mst-like Kinases in the Hippo Kinase Cascade. Dev Cell 34**:** 642-655.

Treatment: vehicle   auxin   vehicle   auxin
Genotype: +   +   cst-1/2(△)   cst-1/2(△)

TIR1 + AID::mig-15*

I Day 3

% developmental stage

Treatment: vehicle   auxin   vehicle   auxin
             214   199   143   135
Genotype: +   +   cst-1/2(△) cst-1/2(△)

TIR1 + AID::mig-15*

J Day 4

Treatment: vehicle   auxin   vehicle   auxin
             214   199   143   135
Genotype: +   +   cst-1/2(△) cst-1/2(△)

TIR1 + AID::mig-15*

- adult
- L4
- L3
- L2
- L1

K

Day 1   Day 3

pumping rate / 60s

vehicle
auxin

Genotype: TIR1 + *AID*::mig-15, cst-1/2(△)

+ auxin

ctrl(RNAi)   *yap-1(RNAi)*   *egl-44(RNAi)*

Day 3

TIR1 + AID::mig-15, cst-1/2(△)*

O Day 3

% developmental stage

RNAi: ctrl   yap-1   egl-44
           119   143   127

Genotype: TIR1 + *AID*::mig-15, △cst-1/2

+ auxin

- adult
- L4
- L3
- L2
- L1

Figure 7. A model for Hippo and Warts signaling in *C. elegans*. (A) In epithelium a canonical Hippo pathway functions, with potentially redundant CST-1 and CST-2 phosphorylating and activating WTS-1. WTS-1 in turn phosphorylates and inhibits YAP-1, keeping it out of the nucleus. Disruption of CST-1/2 or WTS-1 results in nuclear translocation of YAP-1. Rather than contributing to activation of WTS-1 to repress YAP-1, MIG-15 inhibits YAP-1 protein levels. **(B)** In intestine a non-canonical Hippo pathway functions. Unknown upstream inputs regulate WTS-1 to inhibit and retain YAP-1 out of the nucleus. Loss of WTS-1 causes nuclear translocation of YAP-1 but also L2 arrest, indicated as "Growth." CST-1/2 and MIG-15 act redundantly to maintain growth via unknown mechanisms, since their loss does not cause tagged YAP-1 to enter the nucleus.

A

B

C

AID*::WTS-1 + germline expressed TIR1

D vehicle **E** 1 mM auxin

somatic TIR1 alone

auxin + *ctrl(RNAi)*   auxin + *yap-1(RNAi)*   auxin + *egl-44(RNAi)*

G   **H**   **I**

AID*::WTS-1 + somatic TIR1

J

AID*::WTS-1 + somatic TIR1 + YAP-1::mNG

Genotype: +   *wts-1(tm4081)*

yap-1::mNG::2xflag

L

ns

% developmental stage

- adult
- L4
- L3
- L2
- L1

Genotype: +   *wts-1(tm4081)*

yap-1::mNG::2xflag

	Epithelium	YAP-1::mNG	Intestine	YAP-1::mNG
Wild type	M	M'	N	N'
wts-1(tm4081)	O	O'	P	P'

Huynh Fig S4A

CLUSTAL O(1.2.4) multiple sequence alignment

```
CeCST-1      --MPPST-----DSSRRNSEEGSSDGFKLDSSALNKPPEEVFDIVGKLGEGSYGSVHKAI    53
DmHpo        MSEPEVTSVVDMKSPNISSSCSFFKLKKLSEESLLQPPEKVFDIMYKLGEGSYGSVYKAV    60
HsMST1/STK3  -----------ME----QPPAPKSKLKKLSEDSLTKQPEEVFDVLEKLGEGSYGSVFKAI    45
HsMst2/STK4  --------METVQ----LRNPPRRQLKKLDEDSLTKQPEEVFDVLEKLGEGSYGSVYKAI    48
                          .         .  **...:* : **:***:: **********.**:

CeCST-1      HRESGHVLAIKKVPVDTDLQEIIKEISIMQQCKSKYVVKYYGSYFKHSDLWIVMEYCGAG    113
DmHpo        HKESSSIVAIKLVPVESDLHEIIKEISIMQQCDSPYVVRYYGSYFKQYDLWICMEYCGAG    120
HsMST1/STK3  HKESGQVVAIKQVPVESDLQEIIKEISIMQQCDSPYVVKYYGSYFKNTDLWIVMEYCGAG    105
HsMst2/STK4  HKETGQIVAIKQVPVESDLQEIIKEISIMQQCDSPHVVKYYGSYFKNTDLWIVMEYCGAG    108
             *.*:. .:*** ***::**:**********.* :**:******: **** *******

CeCST-1      SISDIMRARRKPLSEQEISAVLRDTLKGLQYLHDLKKIHRDIKAGNILLNTDGIAKLADF    173
DmHpo        SVSDIMRLRKKTLTEDEIATILSDTLQGLVYLHLRRKIHRDIKAANILLNTEGYAKLADF    180
HsMST1/STK3  SVSDIIRLRNKTLIEDEIATILKSTLKGLEYLHFMRKIHRDIKAGNILLNTEGHAKLADF    165
HsMst2/STK4  SVSDIIRLRNKTLTEDEIATILQSTLKGLEYLHFMRKIHRDIKAGNILLNTEGHAKLADF    168
             *.***:* *.* * *.:**::* .**:** ***  :*******.******:* ******

CeCST-1      GVAGQLTDTMAKRNTVIGTPFWMAPEVIEEIGYDTKADIWSLGITAIEMAEGRPPYSDIH    233
DmHpo        GVAGQLTDTMAKRNTVIGTPFWMAPEVIEEIGYDCVADIWSLGITALEMAEGKPPYGEIH    240
HsMST1/STK3  GVAGQLTDTMAKRNTVIGTPFWMAPEVIQEIGYNCVADIWSLGITSIEMAEGKPPYADIH    225
HsMst2/STK4  GVAGQLTDTMAKRNTVIGTPFWMAPEVIQEIGYNCVADIWSLGITAIEMAEGKPPYADIH    228
             ******************************:****:  *********::*****:***.:**

CeCST-1      PMRAIFMIPTKPPPTFKKPEEWSSEFNDFIRSCLIKKPEERKTALRLCEHTFIKNAPGCD    293
DmHpo        PMRAIFMIPQKPPPSFREPDRWSTEFIDFVSKCLVKEPDDRATATELLEHEFIRNAKHRS    300
HsMST1/STK3  PMRAIFMIPTNPPPTFRKPELWSDDFTDFVKKCLVKNPEQRATATQLLQHPFIKNAKPVS    285
HsMst2/STK4  PMRAIFMIPTNPPPTFRKPELWSDNFTDFVKQCLVKSPEQRATATQLLQHPFVRSAKGVS    288
             ********* :***:*::*: ** :* **: .**:*.*::* ** .* :* *::.*   .

CeCST-1      IMQLMIQDAQEKAILGQAPMA------AS-----------SGNDATLL----SEGMSTMI    332
DmHpo        ILKPMLEETCAIREQQRANRSFGGVLAASQAKSLAT--QENGMQQHITDNAFMEDPGTLV    358
HsMST1/STK3  ILRDLITEAMEIKAKRHEEQQ---------RELEE-EEENSDEDEL-------DSHTMV    327
HsMst2/STK4  ILRDLINEAMDVKLKRQESQQ---------REVDQDDEENSEEDEM-------DSGTMV    331
             *:: :: ::         :                  .. : :      . *::

CeCST-1      D--GG--------ESTLVQHKDNYVTA-----QSLR----------------SQME---    357
DmHpo        PEKFGEYQQSSASDATMIAHAEQGVDEGTLGPGGLRNLSKAAAPAAASSAASPLDMPAVD    418
HsMST1/STK3  KTSV------------------ESV-----GTMRATSTM------SEGA--QTMI---    351
HsMst2/STK4  RAVG------------------DEM-----GTVRVASTM------TDGA--NTMI---    355
                                       :*                         *

CeCST-1      -------SLRIGGEIPKS-AYSSSKNGNSPRVQPPGHTASASDPSKNQPFAQ----DGTG    405
DmHpo        SGTMVELESNLGSTMVINSDSDDSTTAKNNDDQKPR---NRYRPQFLEHFDRKNAGDGRG    475
HsMST1/STK3  EHNSTMLESDLG-TMVINSEDEEEEDGTMKRNATSP---QVQRPSFMDYFDKQDFKNKSH    407
HsMst2/STK4  EHDDT-LPSQLG-TMVINAEDEE-EEGTMKRRDETM---QPAKPSFLEYFEQKEKENQIN    409
                          :*    .   .. ..       .   *. : * :   :
```

```
CeCST-1      PNF-------------------------------QLGTSE-----S----SY--------   417
DmHpo        DEKPIATEYSPAAAEQQQQQQQQQQQQQQQDEQHLASGANDLNNWEHNMEMQFQQISAINQ   535
HsMST1/STK3  E-----------------------------------------NCNQNM-----------   414
HsMst2/STK4  S-----------------------------------------FGKSV-----------   415

CeCST-1      -----------KDASYNMMNTE-------------------------------A----E   430
DmHpo        YGLQQHQQQQQVLMAYPLMNEQLIALNNQPNLLLSNAAPMGQQGIPAAAPAQPPPAYQNQ   595
HsMST1/STK3  -----------HEPF--------------------PMS----------------KN   423
HsMst2/STK4  -----------PGP------------------------------------------L   419

CeCST-1      YENRFQRAVVDGDFEFLRNITLDELIRRKESLDSEMEEEIRELQRRYKTKRQPILDVIEI   490
DmHpo        HMHTQSHAYVEGEFEFLKFLTFDDLNQRLCNIDHEMELEIEQLNKKYNAKRQPIVDAMNA   655
HsMST1/STK3  VFPDNWKVPQDGDFDFLKNLSLEELQMRLKALDPMMEREIEELRQRYTAKRQPILDAMDA   483
HsMst2/STK4  KNSSDWKIPQDGDYEFLKSWTVEDLQKRLLALDPMMEQEIEEIRQKYQSKRQPILDAIEA   479
                    :    :*:::**:  :.::*   *    :*  ** **.::.::* :*****:*.::

CeCST-1      KKRLQ----------  495
DmHpo        KRKRQQNINNNLIKI  670
HsMST1/STK3  KKRRQQN--------  490
HsMst2/STK4  KKRRQQNF-------  487
             *::  *
```

Huynh Fig. S4B

***cst-1* genomic seq annot:**

Lt gray: UTR

Yellow: exons

Green: alternating exons

Blue: two-codon alternative splice in intron 8.

Purple: exon 9 sequence that is IDENTICAL with *cst-2* exon 9.

```
>Unspliced + UTR + upstream + downstream (8069bp)
atttttcaaaaatatatacacccataatatgaaaaagtattttcaactatcaataaacac
cgagaaaagcggaatatcaaaatttttatgaattaggggtgggaggcaattgttaaaatg
cggaaaatattgaattgctgatgaattcaagattcggcgattcccgaaaataccgaatat
tttcaaactcggcaatctggaatagcagacatttagtagggaaacgttgtatgtcttccc
aaaaatatttattacaatgaactattgtttgtttgtttgtttttgaattttatttgttca
aaattatcgattgtcaaattgccaaaaaattttgatattccgcttttctcggtgtttatt
gatagttgaaaatacttttcatattatgggtgtatatatttttgaaaaatcaattaaca
gttctttaaagttaaatagattgaaattaaagaataacttggaaattgtttttgaattta
tgacatctctttgcttgaaaaaatggatgaaacatgctttataaaatcaactagaataat
tgtccaaactgacaaaaagcaatcaactactatgcaccataacaaataaagtctactgac
aagtaaagatagtaaacatacgtgaatcaggtgatgactacccaacaactctcataataa
aacagtaccgcaaaacagcaccaggaaggacacagttagcaatacgatattccgcattca
aacatcctgacgtgtgaagggtcaagtcttatgaaacgagcgcagtacacacttcgcact
ctttcccgtccttttcccccctttcgcctcgaccacacactgctcaaataacgtagagtga
gccgagggcattacgaatgctcgtcgaatgtagcgcctgcggtcggttatattaacttct
tttttttatcacggtacaacttatctgaaatagagtactgtttttgagttgttctctgg
tctttgatagtttgatagtttcaatgggaatatttaattctcacactttcaaaaaggata
tttcaatcttaaatgaaatgctaacattgcgaaatagaattgaaaatcatgatacttcaa
tctctttcatgaaaaataaggttcttctaaatttgtattgtttttaaaatatgctcaggtc
tatctcaaagatttcataaattataaaatttattaattagttttttctcggctttttcaac
cattttaccaatgaaatcttaatctaaaaacgttggactttgtcggattacttttttttc
caaaagacaattgccgtagtcaattattcattcttcgggccatcggggccacatgggtgg
aacccccgcagacgtcttttcgcttttcaaaacgacgtccaaatcgattcgcctgtctttct
cttctcactcttctcgcctcccttcagaatatgaccatacatgcactgttgacgttctaa
ttgaacaagcagagtgcatgtgcaaccaagagaacgcagagaaacatgcgactaatgtgt
cctcaaggataggtttttatcttcttcttcttctttcttcttctttcttcgattttcttgc
cccatagttctcgtcggtcggacggacaacatctgctcttcatcttacttatatctcatt
gggtgttcattcctatactcacacaagatcttacaacactttttttataaatccttcaact
tgtacacagttaaaggtggagtagcgcaagttggaaaattgtttaaaactactcttatgg
tgtcaaaatgactaaatattaaacaaataaaaatgctgaacagtttttgaaaggacgact
ttcaaaaagtccaacaattggcaaaaactgagtttattttcaattttcaaaaaaaaatca
gataaaatttagaacatctttcagaaacttttgctaaaatatttgattattgtggtgcc
gcatgcatgttttttgagcaatttcagcactgacgttactccacttttaaatttcagcaaa
gttaatttcattaatttcag
```
cgtcctcccg`ATG`CCACCGTCTACAGACAGCTCGCGGCGG AATTCGGAAGAGGGCTCAAGgtttgtcagacacaaaacaaaaaaaaagagttactcttcc aaaataaacattctcttttgtttttagCGATGGATTTAAATTGGATTCGTCGGCGCTTAAC AAACCACCAGAAGAAGTTTTCGACATTGTTGGAAAGCTGGGCGAAGGgtttgttatttca attctatcggattgaaaaattccaaattttcagATCATATGGAAGCGTACATAAGGCAAT TCACAGAGAGTCAGGTCATGTGTTAGCCATCAAAAAAGTGCCAGTGGATACTGATCTTCA

```
AGAAATTATCAAGGAAATCAGTATAATGCAACAATGTAAAAGTAAATATGTTGTCAAATA
CTATGGATCGTATTTCAAACATTCGGATCTATGGgtgagattgtttagataaattttaag
attttaatgctgtgcccaagtgggactaaataaaagaaattgaacaaaaattatattcgt
tttgtgaattagagttaaaaacagtaaagttctgattttttccacagaaaaaaatgtagct
taaagtggaaataacattcaaaagtattcattcaaatgcatgccaattgaaattcaaaga
ttttattccctacccctagtcgaaaatcaagaagaaactggactaacaaaaaaactaat
ttaaagATTGTGATGGAATACTGTGGAGCTGGGTCGATCTCCGATATAATGCGTGCCAGG
CGGAAACCGTTATCCGAACAAGAAATCAGTGCAGTTTTGCGCGATACTCTCAAGGGATTG
CAATATTTGCATGACTTGAAGAAGATTCATAGAGATATAAAGgttagaaactgtttttttt
ttgaaaaaaaaactgtattttcagGCTGGAAACATTCTACTCAATACCGATGGAATTGCA
AAGCTTGCGGATTTCGGTGTCGCTGGACAGTTGACTGATACAATGGCAAAAAGAAATACA
GTCATTGGAACGCCATTCTGGATGGCCCCTGAAGTTATCGAAGAAATCGGATATGACACA
AAAGCTGATATATGGTCACTGGGAATAACAGCTATCGAAATGGCAGAGGGACGGCCTCCT
TATTCAGATATTCATCCGATGCGAGCAATCTTCATGATTCCAACTAAACCACCACCTACG
TTTAAAAAACCTGAAGAATGGTCGTCCGAGTTCAATGATTTCATAAGAAGCTGTTTAATT
AAAAAGCCAGAAGAAAGAAAAACGGCACTCCGATTGTGTGAGgtacgtatggcttaaaaa
aaaagaaatttcccaataagatttatccaaaaaatattgaaattctcaaatattcgttat
atttgttaaaactgtgaaaaaaggtacatccacctaaaattcaaaatttcgacaacttttt
ttttgtcgcagggactagaaattaattttcaatcatgtagtatttctttttttctattttt
taaagtttctagtcggtagtcagctaaacttgtgttctaatcggctgaaaatatgtccat
gtcagcaaaaaatgcgagaaaagcgataaactttcacgtttgactacaaaatgatgtcga
taaaagcttgaaatctataaaaacaaacaatccaaaaaacgctagaactcaacataattc
tctactcggtattcagaggcaacttcctgccacactttgacaacttatttagtgccacta
aaataagttgccaaattgctggcaggaagttgccgagttgttggcaggaagttgccacgt
agtggcagtaagtgggaaaaaacatctaaatgttggcaggaatgtgcaaagtttggcag
aaagttgccaaaagttggcaggaagttgcctttaaatgccgaggagcgttctttcccact
tgctgccactacgtggcaactcatttaaaaactgccaagatatagacgggggccaaatttt
catatttcctacaactactctcaaccgaaaatagaagatttcttataatttgacaacccc
ttcggagtaatttttaactttttacatagaattttctactattttttatacacttttttg
ttgttttctgtcatagtattactaatactgctgaaaaacaaatgatacacagtttgtgaa
cattgctcacttcatagatattttttcagCACACATTCATCAAAAATGCACCAGGTTGCGA
TATTATGCAGTTGATGATCCAGGATGCTCAAGAAAAAGCTATACTAGGACAAGCACCAAT
GgttgacaaaaaatttcattagaaaacgaaaatttgattgtttttttcagGCAGCAAGCAG
CGGAAATGATGCAACGTTGCTAAGCGAGGGAATGTCCACTATGATTGACGGTGGAGAGTC
TACGTTAGTTCAACACAAAGgtattatgatagttttgaatatgtaatttgatcatatgtg
tcgtgggaaagtgcaaaactcacgcacctaatttgcactactttgtagctcgatatttag
tagtaacgtgaagcaggacacagggagaaattacaaaattacctagtgaaatagaactca
caatttcagACAACTATGTCACTGCTCAAAGTCTTCGAAGTCAAATGGAAAGCTTGAGAA
TTGGTGGAGAAATTCCTAAATCGGCATATAgttcgttttttcatgagcgttaaattgcat
tcaccgatatagttccagGCTCGTCGAAAAATGGCAACTCACCGCGCGTTCAACCACCGG
GTCACACCGCAAGCGCTTCAGACCCTTCAAAAAACCAACCGTTTGCTCAAGATGGAACCG
GTCCAAACTTTCAATTGGGCACCAGCGAGAGCAGTTACAAAGATGCTAGTTATAATATGA
gttagttatttgaaagaaaaccaattataaaatgatgatctatgagaaaaaaagctagc
ataaaaaacgttattcagctcgaatgactttgacaaaaaaatgaaaaattattcaggcat
tttcacaaacctaaaattttattaaaattgaaatgtcataccccttattctagccatgca
tgaaatgttagctgattaagattttgcctgtaaattctaaaaaatgaaaaatgtcactac
taatgtttttgtggaaaatttgtattgcataagttttgaaacttagaaagacatattttga
aatcttgaattgaatcacagttaaaaaacacctactctttcagTGAACACGGAAGCCGAA
TACGAAAATAGATTTCAACGGGCAGTGGTCGATGGAGATTTCGAATTTgtaagtgtattt
```

cgtaattttttctaataacgaaatagctgcatttag**TTACGAAACATCACGCTAGACGAAC
TGATTCGAAGGAAAGAAAGTTTGGATTCGGAGATGGAAGAAGAAATACGAGAGTTGCAGA
GAAG**gtaaaatcttttttttttaattggttgaatttgaggaaatttaattcaaacacctc
aaaaagccggaaattgagatttttaacaaaaaatcctaagcgaattttctagctggattt
tttgtgagtttactgatttaaacatctaaaggacagttttatgaatcgcgaactactgct
atttataattttaagaaatagtgatgctgaaattttgctcagaattttttctttgatg
cttaaaattccattctcaaatttaggatatttgaagcttttggcatcgaatgttttcaaa
atgagccaacttttctctttttttcttacattttggagtcaagaaaattgatattaaaat
ccgcaacagtggcaattcacgattaaagtcaaatactaaaaaaagaacaaaatttcag**AT
ACAAAACAAAAAGGCAACCAATACTGGATGTTATAGAAATCAAAAAACGACTTCAA**TGAc
ggcttcgatacatgcttgtttaataataattttttttgtcaatctgcaaatatatgaaacc
tattaattaactagattgtattttaaatgtcatatttattctaaattttgttagatcaga
agcaagagcaaactgaacttgttgtgttttgcttagcttttttcatgctaaaataaatctg
gtcaattttaatttttttgggaaggtttgttctgaaggcggatattttatcagtgactaa
caaggaacaatgggagtgacggatataattttttcgaataacggaaatagataaaatttca
cttacaattgaaaaagctgcaatgagtcagagcgatagatggttgatctcgatgagtcta
gttttttcttatttgtttgcctaccgcggagttcacgtagcgaccaattaatatttttagg
taaattttgcatggcatcctagtagaaacaaaaaaccacattctgagtgctttttgcatt
acgactgccgttttgaatcgtaagatttatggcttatgacggacgtccttttagctttag
tctgttataattgtcatattgggtctctgctgccgttgggaagcagatatatttcgtcaa
ctactagccgagagggcgtttgctttttttttttcaagaagaggcaaatatacacttcga
ggcgaatgtttctcaactgtcaggtaatatttgcgaaaccgaaatatatttaaggagtga
aggaagcgaagattgccatcgaacaaaagttcatccgctatcagcttatcaacaatgttc
gcacttattctttctttgtgtctttttcaaggtaaactattctttcagaattatagtatt
ttctgaaaaatgtttccaatttctgaaacagttttttttttgattgttataaaatattttt
aggaacactttcatatgtaattggaggcggtgtacaagcaaatccaagcggatgcaacgt
atgtggtgatagccctcagtggagtggatggacagaatggtcttcctgttctgccgcttt
tggaagtaggtcttctattttttgaattttttgctatagcagtttttgcgaaaacctgacac
atcaatttaaagaaacattcagttttactggtcccgccacaatttcagtgatctttatga
aatgtaaaattctacagtgcacagagattcttggatttgaaatagaatttccaacacctg
gagtcacatttccgctatatgtacatcttcataataataatttcaatttcagcgccagtc
caaactcgcactagactttgtccatccggcaactgccaaggaggttctacaacagaatca
aaaccatgcgtcctttatgatccacaaccaactcaaccacaatggggagcttggggaggt
tggagtagttgcaggttagcaatattttgctttaatctgtttattattgttttgttttct
ttattgccaaactgataagcctaacttttctttttgcagtgccacctgcggcggtggtac
gatgacgagaagtcgtgtttgcaataacggatgctcaacctgccaatgtgtcggagctgc
tgccgagtcgcaagcttgcaatgctcaaccgtgctgcacctggactgctgtgagaacaaa
ttgagttcaacttctcacttaataactttttgtttcagtggtcttcctggtcggcgtgctc
agtaacctgcggatctggaggatcaatcaccagatctaggcaatgtagctgtggatcggg
agtaagtttaaaatttataaagctaaacgttttttattcaatataacatttataactgttt
gtttcagtgtactggaggctccgttgaacaagagccatgccctcaacaagctgcttgccc
gtgcacaacatgcaatcaaccaccaccaccatgcaatacttgtaatacccaaccggttgt
cattgtgactccagctccatgcacaacttgttaccaaccaccagcgtgctcaacctgcgg
gcatgcacaacctttctatgatccatacggaaatggaagaaagaagagaatgattacagt
atctggaaattcgacaagtgcttaactgtattgaaacttgtgtacatttatgcaatattt
tgcacattccgaagcaatataaataaagtattgagcaattattatttctatccactacag
cagaatcacagaaccactaaaaaaacaaa

>Unspliced + UTR + upstream + downstream (6994bp)
taattgatttttcaaaaatatatacacccataatatgaaaaagtattttcaactatcaat
aaacaccgagaaaagcggaatatcaaaattttttggcaatttgacaatcgataattttga
acaaataaaattcaaaaacaaacaaacaaacaatagttcattgtaataaatattttttggg
aagacatacaacgtttccctactaaatgtctgctattccagattgccgagtttgaaaata
ttcggtattttcgggaatcgccgaatcttgaattcatcagcaattcaatattttccgcat
tttaacaattgcctccaccccctaattcataaaaattttgatattccgcttttctcggtg
tttattgatagttgaaaatacttttcatattatgggtgtatatattttgaaaaatcaa
ttaacagttctttaaagttaaatagattgaaattaaagaataacttggaaattgtttttg
aatttatgacatctctttgcttgaaaaaatggatgaaacatgctttataaaatcaactag
aataattgtccaaactgacaaaaagcaatcaactactatgcaccataacaaataaagtct
actgacaagtaaagatagtaaacatacgtgaatcaggtgatgactacccaacaactctca
taataaaacagtaccgcaaaacagcaccaggaaggacacagttagcaatacgatattccg
cattcaaacatcctgacgtgtgaagggtcaagtcttatgaaacgagcgcagtacacactt
cgcactctttcccgtccttttccccctttcgcctcgaccacacactgctcaaataacgta
gagtgagccgagggcattacgaatgctcgtcgaatgtagcgcctgcggtcggttatatta
acttcttttttttttatcacggtacaacttatctgaaatagagtactgtttttgagttgtt
ctctggtctttgatagtttgatagtttcaatgggaatatttaattctcacactttcaaaa
aggatatttcaatcttaaatgaaatgctaacattgcgaaatagaattgaaaatcatgata
cttcaatctctttcatgaaaaataaggttcttctaaatttgtattgtttttaaaatatgct
caggtctatctcaaagatttcataaattataaaatttattaattagtttttctcggcttt
ttcaaccattttaccaatgaaatcttaatctaaaaacgttggactttgtcggattacttt
tttttccaaaagacaattgccgtagtcaattattcattcttcgggccatcggggccacat
gggtggaaccccgcagacgtcttttcgctttcaaaacgacgtccaaatcgattcgcctgt
ctttctcttctcactcttctcgcctcccttcagaatatgaccatacatgcactgttgacg
ttctaattgaacaagcagagtgcatgtgcaaccaagagaacgcagagaaacatgcgacta
atgtgtcctcaaggataggtttttatcttcttcttcttctttcttcttctttcttcgattt
tcttgccccatagttctcgtcggtcggacggacaacatctgctcttcatcttacttatat
ctcattgggtgttcattcctatactcacacaagatcttacaacacttttttataaatcct
tcaacttgtacacagttaaaggtggagtagcgcaagttggaaaattgtttaaaactactc
ttatggtgtcaaaatgactaaatattaaacaaataaaaatgctgaacagtttttgaaagg
acgactttcaaaaagtccaacaattggcaaaaactgagtttattttcaattttcaaaaaa
aaatcagataaaatttagaacatctttcagaaactttttgctaaaatatttgattattgt
ggtgccgcatgcatgttttttgagcaatttcagcactgacgttactccacttttaaatttc
agcaaagttaatttcattaatttcagcgtcctcccg**ATG**CCACCGTCTACAGACAGCTCG
CGGCGGAATTCGGAAGAGGGCTCAAGgtttgtcagacacaaaacaaaaaaaaaagagttac
tcttccaaaataaacattctctttttgtttttagCGATGGATTTAAATTGGATTCGTCGGCG
CTTAACAAACCACCAGAAGAAGTTTTCGACATTGTTGGAAAGCTGGGCGAAGGgtttgtt
atttcaattctatcggattgaaaaattccaaattttcagATCATATGGAAGCGTACATAA
GGCAATTCACAGAGAGTCAGGTCATGTGTTAGCCATCAAAAAAGTGCCAGTGGATACTGA
TCTTCAAGAAATTATCAAGGAAATCAGTATAATGCAACAATGTAAAAGTAAATATGTTGT

CAAATACTATGGATCGTATTTCAAACATTCGGATCTATGGgtgagattgtttagataaat
tttaagattttaatgctgtgcccaagtgggactaaataaaagaaattgaacaaaaattat
attcgttttgtgaattagagttaaaaacagtaaagttctgattttttccacagaaaaaaat
gtagcttaaagtggaaataacattcaaaagtattcattcaaatgcatgccaattgaaatt
caaagattttattccctacccctagtcgaaaatcaagaagaaaactggactaacaaaaaa
actaatttaaagATTGTGATGGAATACTGTGGAGCTGGGTCGATCTCCGATATAATGCGT
GCCAGGCGGAAACCGTTATCCGAACAAGAAATCAGTGCAGTTTTGCGCGATACTCTCAAG
GGATTGCAATATTTGCATGACTTGAAGAAGATTCATAGAGATATAAAGgttagaaactgt
ttttttttgaaaaaaaaactgtattttcagGCTGGAAACATTCTACTCAATACCGATGGA
ATTGCAAAGCTTGCGGATTTCGGTGTCGCTGGACAGTTGACTGATACAATGGCAAAAAGA
AATACAGTCATTGGAACGCCATTCTGGATGGCCCCTGAAGTTATCGAAGAAATCGGATAT
GACACAAAAGCTGATATATGGTCACTGGGAATAACAGCTATCGAAATGGCAGAGGGACGG
CCTCCTTATTCAGATATTCATCCGATGCGAGCAATCTTCATGATTCCAACTAAACCACCA
CCTACGTTTAAAAAACCTGAAGAATGGTCGTCCGAGTTCAATGATTTCATAAGAAGCTGT
TTAATTAAAAAGCCAGAAGAAAGAAAAACGGCACTCCGATTGTGTGAGgtacgtatggct
taaaaaaaagaaatttcccaataagatttatccaaaaaatattgaaattctcaaatatt
cgttatatttgttaaaactgtgaaaaaaggtacatccacctaaaattcaaaatttcgaca
acttttttttgtcgcagggactagaaattaattttcaatcatgtagtatttctttttct
attttttaaagtttctagtcggtagtcagctaaacttgtgttctaatcggctgaaaatat
gtccatgtcagcaaaaaatgcgagaaaagcgataaactttcacgtttgactacaaaatga
tgtcgataaaagcttgaaatctataaaaacaaacaatccaaaaaacgctagaactcaaca
taattctctactcggtattcagaggcaacttcctgccacactttgacaacttatttagtg
ccactaaaataagttgccaaattgctggcaggaagttgccgagttgttggcaggaagttg
ccacgtagtggcagtaagtgggaaaaaacatctaaatgttggcaggaatgtgcaaaagtt
tggcagaaagttgccaaaagttggcaggaagttgcctttaaatgccgaggagcgttcttt
cccacttgctgccactacgtggcaactcatttaaaaactgccaagatatagacggggcca
aattttcatatttcctacaactactctcaaccgaaaatagaagatttcttataatttgac
aaccccttcggagtaatttttttaactttttacatagaattttctactattttttatacact
tttttgttgttttctgtcatagtattactaatactgctgaaaacaaatgatacacagtt
tgtgaacattgctcacttcatagatatttttcagCACACATTCATCAAAAATGCACCAGG
TTGCGATATTATGCAGTTGATGATCCAGGATGCTCAAGAAAAAGCTATACTAGGACAAGC
ACCAATGgttgacaaaaaatttcattagaaaacgaaaatttgattgtttttttcagGCAGC
AAGCAGCGGAAATGATGCAACGTTGCTAAGCGAGGGAATGTCCACTATGATTGACGGTGG
AGAGTCTACGTTAGTTCAACACAAAGgtattatgatagttttgaatatgtaatttgatca
tatgtgtcgtgggaaagtgcaaaactcacgcacctaatttgcactactttgtagctcgat
atttagtagtaacgtgaagcaggacacagggagaaattacaaaattacctagtgaaatag
aactcacaatttcagACAACTATGTCACTGCTCAAAGTCTTCGAAGTCAAATGGAAAGCT
TGAGAATTGGTGGAGAAATTCCTAAATCGGCATATAgttcgttttttcatgagcgttaaa
ttgcattcaccgatatagttccagGCTCGTCGAAAAATGGCAACTCACCGCGCGTTCAAC
CACCGGGTCACACCGCAAGCGCTTCAGACCCTTCAAAAAACCAACCGTTTGCTCAAGATG
GAACCGGTCCAAACTTTCATCCACTATCTCAACACTTTGTTTTGGTTTTTATAATTTAAa
agcctagctaatatcctaatctgtttacctgatataatagttttcaaatttaccttctgg
aaaacttatatattactatactgtcgaaacggatgccgcacatccctttgctgtaaaaat
tgccgagttcctaactctttgtccgcaaaactcacaaatagtctatgagataatgcatgt
tctcttctgaacctttcaaatcaactatagctggtaagtactctttcaattagttcagct
tcttttcttttctcaccaactactatctcgactccgacctatttagatcggaaccaattc
tctcccgatcacttcaacatatgaaaaagacgaatttgtgagttagagagtatactcgaa
atagtaacattgatttcccccattttttcgactttcttttcttcgctatcgtgtgtactta
tcgcatgtaacatttgctgattagctgactgaccacggcttcaagtagcaaatttatttc

```
tagcacgtgaggttgcatagatgagtaccttcgttagctttgctttcaggaactcaaaat
tgaaaagagaaacatgtggtgaagcagcggtaggtgaaactctttttctaagctattgca
agattttctcaagatcacacgcctagtatgcagggtgagaatgccggatgcattgaaaat
aaatgcaacaagttgcagctatgattagctttttttaatgattcgactgctttcagaatg
gctttatattgcaccagctctttgagggatctttgaataattttttcaggttgaaaagc
gaaagatgagacgtttatccgagatgatatctgattgggaacgaccgtcaagaagttcga
tcagtaacatgtatctttatgcgagaagagaaagcgtctacattgcaaggaagaagttgg
ccggtctgaaagtaataatttgagctttaaaaacaaacacactataaagactgaaaataa
tccgttttttaatacaaaactgtttaaaaaaatttaaaaaagattcatacttgttgttgtt
ctagaatgttcatggttcaatttagattagctcaagtatgattttaaatattttttgtta
tatccagattgaaaatacgtagaatattcaaagaaagacattgaaatttgaattacctgc
aaaatacgcatgtttcagccacttttggtaactcaacgctaaattttcataaaaatttc
catatcctttgaaacatccggaatctaaaaaaaaacaagaaaaaaacggaaaatattgga
actataccaaaataaaattaccgaattttcttcaactgtccgattttccaggactgggcg
ctggacctgttagcaaaactgaaatccaagcagggaatggcgattcgtcgagagactcgt
gcaaccaagttggttgcaactgtaatgggtaactgaaactattctttgacattaaaacta
aaccaaatacattgcagtggtttttctcgtctgctggcttccgttttttcacattgaacat
gatcaaaatttacaaattgatattcaacgtttggtcagcggacctcgaaatttggttcca
ttggtttaccgcactggggtatttgaattcgtcactcacttttttcatttactcaacaat
caatccagtaagcattttttatatcaaatttgatagtaattatagaaataattataataca
taattggctatataataaaaaagtttagtcgaaaaagattatatttttaattttttcaagt
tcaatttgcgttgtataaaacattaattcatggaattgagacaaggtaaaaccgacaacg
ttttccaaaattttgacatgtctgtattccctctataggcattgcgccagcttaactcaa
tttgacaagaaatagtaaacccgcttagcaaactaaatatactgaaaactgtactgtata
atccattcaacacctcctttttttcagaaattccgtcattcattccgccgcttacttggct
tccgtcgatcttctcgacgtcagagagaaaaatcgtggatgttaccgccacgagatagca
gcaggtgagcctcatttattcgtccatagctttttaaatgtttcataacagaggtcctgca
aaatgtggatgcggtttccataaaatcgtgaaaa
```

L

STOP-IN = *re427*

gck-2

Δ = *re483*

A

B Treatment:

| vehicle | auxin |

Genotype: somatic TIR1 + AID*::MIG-15 + YAP-1::mNG

C

Genotype: somatic TIR1 + AID*::MIG-15 + YAP-1::mNG

SUPPLEMENTARY TABLES

Table S1: Strain

Strain	Genotype
CA1200	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II; *unc-119(ed3)* III
DV3709	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II; *unc-119(ed3)* III; *mig-15(re264[AID*::mNG::2xHA::mig-15])* X
DV3948	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II; *unc-119(ed3)* III; *mig-15(re264[AID*::mNG::2xHA::mig-15]) cst-1/2(reDf1[re484])* X
DV3969	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II; *unc-119(ed3)* III; *gck-2(re483)* V; *mig-15(re264[AID*::mNG::2xHA::mig-15]) cst-1/2(reDf1[re484])* X
DV3611	*yap-1(re269[yap-1::mNG::2xflag])* X
DV4205	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II; *unc-119(ed3)* III; *yap-1(re269[yap-1::mNG::2xflag])* X
DV4130	*rrf-3(re390[STOP-IN*])* II; *his-72 (erb77[his-72::linker::mTurquoise2])* III, *yap-1(re269[yap-1::mNG::2xflag]* X
DV4186	*his-72(erb77[his-72::linker::mTurquoise2])* III, *yap-1(re269[yap-1::mNG::2xflag] cst-1/2(reDf4[re452])* X
DV4217	*wts-1(tm4081)/tmC18[dpy-5(tmIs1236[myo-2p>mCherry])]* I
DV4214	*wts-1(re419[mTurqoise2::2xMyc::AID*::wts-1])* I; *ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II; *yap-1(re269[yap-1::mNG::2xflag]* X

DV4288	*his-72(erb77[his-72::linker::mTurquoise2]) III; gck-2(re427[STOP-IN*]) V; yap-1(re269[yap-1::mNG::2xflag]) cst-1/2(reDf4[re452]) X*
DV4290	*wts-1(re419[mTurqoise2::2xMyc::AID*::wts-1]) I; ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)] II; unc-119(ed3) III*
DV4292	*wts-1(ok753)/tmC18[dpy-5(tmIs1236[myo-2p>mCherry])] I*
DV4278	*wts-1(tm4081) I; yap-1(re269[yap-1::mNG::2xflag]) X*
DV4297	*wts-1(re436[STOP-IN*])/tmC18[dpy-5(tmIs1200[myo-2p>Venus]) I*
DV4296	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)] II; unc-119(ed3) III; mig-15(re435[mTurquoise2::2xmyc::AID*::mig-15) yap-1(re269[yap-1::mNG::2xflag]) X*
DV4337	*wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1]) I; reSi2[col-10p>TIR1::F2A::mTaqBFP2::AID::NLS] II*
DV4338	*wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1]) I; reSi12[ges-1p>TIR1::F2A::mTaqBFP2::AID*::NLS] II*
DV4339	*wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1]) I; reSi2[col-10p>TIR1::F2A::mTaqBFP2::AID*::] II; yap-1(re269[yap-1::mNG::2xflag]) X*
DV4340	*wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1]) I; reSi12[ges-1p>TIR1::F2A::mTaqBFP2::AID*::NLS] II; yap-1(re269[yap-1::mNG::2xflag]) X*
DV4341	*wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1]) I; ieSi68[sun-1p>TIR1::mRuby + Cbr-unc-119(+)] II*

DV4384	*ieSi57[eft-3p>TIR1::mRuby + Cbr-unc-119(+)]* II*; unc-119(ed3)* III*; mig-15(re435[mTurquoise2::2xmyc::AID*::mig-15) yap-1(re269[yap-1::mNG::2xflag]) cst-1/2(re479)* X	

Table S2: Oligonucleotide

Name	Oligonucleotide sequence	Use
LH12	GATATTGCCGAGCACACATGG	genotyping triplex *rrf-3(re390)*
DJR881-RV1	GCTTATCACTTAGTCACCTCTGCTC	genotyping triplex *rrf-3(re390)*
LH13	GATGTGAAACTTGATGCGAACTCG	genotyping triplex *rrf-3(re390)*
LH11 repair *rrf-3*	AATTTCAGTCTAAAGTTGACGAGGTAAACTGGATCAGGGAAGTTTGTCCAGAGCAGAGGTGACTAAGTGATAAGCTAGCAGGTGGGGATGGTGCCACGAGCTGCGTACGAAGATAAA	repair template for *rrf-3(re390)*
LH60	CTTACTTTCTATTCAAGGTTTTATTTCAAACGTAGGACACC	genotyping triplex *wts-1(tm4081)*
LH61	TGCTGTAGAAGTCGTTCAACATGTTGTTCC	genotyping triplex *wts-1(tm4081)*
LH62	TGTCTTCGCAGACACCCATAATCAGC	genotyping *triplex* *wts-1(tm4081)*

LH200	AAAGCGGATGTCATAATGAAACAACAGGTAGG	genotyping triplex *wts-1(ok753)*
LH201	GTGAAAGCCGAACGCGATATTCTGG	genotyping triplex *wts-1(ok753)*
LH202	AGCTGCAACTTCAGGAGGAAGC	genotyping triplex *wts-1(ok753)*
LH131	CGGGAAGCAAAGAGACCACATAACG	genotyping triplex *wts-1(re436[STOP IN])* and *wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1])*
LH133	AACCAAAACCTACTGAATTGTGATGGGC	genotyping triplex *wts-1(re436[STOP IN])* and *wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1])*
DJR881-RV1	GCTTATCACTTAGTCACCTCTGCTC	genotyping triplex *wts-1(re436[STOP IN])*
LH134 repair wts-1 stop-in	GTGTTCGGCAGGAGGCCCTCCTTTGCATTGTATTCCCGGGAAGTTTGTCCAGAGCAGAGGTGACTAAGTGATAAGCTAGCTGCGACCTGCTGCCCCTGGTACTACTCCAAAT	repair template for *wts-1(re436[STOP IN])*

LH132 RV1_wts-1-AID	CGGTAAAGAGTTCTTCTCCTTTGGAGACC	genotyping triplex *wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1])*
LH100 FW_repair mTurq2-wts-1	GGAGGCCCTCCTTTGCATTGTATTCCCCTGGAATGGTCTCTAAGGGAGAAGAACTCTTTACC	primer for amplifying fragment #1 repair template for *wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1])*
LH101 RV_repair mTurq2-wts-1	ATCAGAAGATGCACCATTTGGAGTAGTACCTGGAGCTGCAGGTCGCATacCtGCgCCgCTTGCTCCtGATCCaGCaCCCTTCACG	primer for amplifying fragment #2 repair template for *wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1])*
CB003 Deletion Repair Template	CAAAACAAAGTGTTGAGATAGTGGATGAAAGTTTGCAGTGGCAATTCACGATTAAAGTCAAATACTAAAA	repair template for *cst-1/2(reDf1)*, *cst-1/2(reDf4)* and *cst-1/2(re479)*
CB004 CST FW	GGCATCCGTTTCGACAGTATAG	genotyping triplex *cst-1/2(reDf1)*, *cst-1/2(reDf4)* and *cst-1/2(re479)*

CB005 CST RV-1	GCCACTGCAAACTTTCATCC	genotyping triplex *cst-1/2(reDf1)*, *cst-1/2(reDf4) and cst-1/2(re479)*
CB006 CST RV-2	TATGTGTCGTGGGAAAGTGC	genotyping triplex *cst-1/2(reDf1)*, *cst-1/2(reDf4) and cst-1/2(re479)*
RF21	TTAGCAACGATAACAATCGAGGAGTGGGATCAATGGTGCACCATT ATCAAGGAGCATCGGGAGCCTCAG	primer for amplifying fragment #1 repair template for YAP-1 tag
RF22	AGAGAAAGAGAGTGGTTATTCTGCTGATTAGACATTTACGAGGCTC CACGCTTGTCG	primer for amplifying fragment #2 repair template for YAP-1 tag
RF26 FW	CACACGAATTCGACCAGTATCTACA	genotyping *yap-1(re269[yap-1::mNG::2xflag])*
RF27 RV	CATCGACCATAGCGGCTTG	genotyping *yap-1(re269[yap-1::mNG::2xflag])*

RF28 RV	GTATTGGAAACGGGGTTGAGC	genotyping *yap-1(re269[yap-1::mNG::2xflag])*
LH004v2 GCK-2 Repair Template	GTGATTCACTGGCAGCTGTCAAAGTAGTCAAACTCGATCGCAACCAATGTCACCCCGAAACTCTGGCGATAC	repair template for *gck-2(re483)*
LH140 GCK-2 STOP-IN Repair Template	TGATTCACTGGCAGCTGTCAAAGTAGTCAAACTCGGGGAAGTTTGTCCAGAGCAGAGGTGACTAAGTGATAAGCTAGCAGGCGGGCGACAATTTTGCAGTAATCCAGCAGGAGATT	repair template for *gck-2(re427[STOP IN)*
LH80 GCK-2 FW	GGGTAATGAGTGCCGATGTAATCAAACG	genotyping triplex *gck-2(re483)*
LH81 GCK-2 RV-1	GAAGGCGATTTGAAGTTCCGAAAGTGG	genotyping triplex *gck-2(re483)*
LH83 GCK-2 RV-2	AGTTTCGGGGTGACATTGGTTGC	genotyping triplex *gck-2(re483)*
LH141	GGTAATGAGTGCCGATGTAATCAAACG	genotyping triplex *gck-2(re427[STOP IN)*
DJR881 stop-in RV1	GCTTATCACTTAGTCACCTCTGCTC	genotyping triplex *gck-2(re427[STOP IN)*
LH143	GCCGCAGTACTCCATAACAATCC	genotyping triplex *gck-2(re427[STOP IN)*

LH148_FW primer of AID-mig-15 repair template	TTGCTCCAAGCCGCTCACAGCACCCAAAACCATGTCAATGGTCTCCAAAGGAGAAGAACTC	primer for amplifying fragment #1 repair template for *mig-15(re435[mTurquoise2::2xmyc::AID*::mig-15])*
LH149_RV primer of AID-mig-15 repair template	AGAATTCAAATCAATCTCGTCGAGTCCTGATGACGACATACCTGCGCCGCTTGCTCCTGATCCAGCACCCTTCACG	primer for amplifying fragment #2 repair template for *mig-15(re435[mTurquoise2::2xmyc::AID*::mig-15])*
RF15_FW_AID-MIG-15	CATTGTGTGGTTTAAGTGTCGGC	genotyping triplex *mig-15(re435)*
LH151_RV1_AID-MIG-15	GTGCAAATGAATTTAAGGGTGAGTTTTCC	genotyping triplex *mig-15(re435)*
LH152_RV2_AID-MIG-15	CGTGTTGTTTGGTGTAGTTTCGC	genotyping triplex *mig-15(re435)*
*dpy-10(cn64*gf*)* ssODN repair template	CACTTGAACTTCAATACGGCAAGATGAGAATGACTGGAAACCGTACCGCATGCGGTGCCTATGGTAGCGGAGCTTCACATGGCTTCAGACCAACAGCCTAT	ssODN repair template for *dpy-10(cn64*gf*)*

Table S3: CRISPR guide RNA

RNA	Sequence	Use
LH10 rrf-3 crRNA-1	CGAGGUAAACUGGAUCAAGGGUUUUAGAGC UAUGCUGUUUUG	crRNA for *rrf-3(re390)*
LH130 crRNA rev wts-1	GGCAGCAGGUCGCAUUCCAGGUUUUAGAGC UAUGCUGUUUUG	crRNA for *wts-1(re436)* and *wts-1(re419)*
CB001 cst-2 crRNA	UAGUGGAUGAAAGUUUGGACGUUUUAGAGC UAUGCUGUUUUG	crRNA#1 for *cst-1/2(reDf1), cst-1/2(reDf4)* and *cst-1/2(re479)*
CB002 cst-1 crRNA	AUCGUGAAUUGCCACUGUUGGUUUUAGAGC UAUGCUGUUUUG	crRNA#2 for *cst-1/2(reDf1), cst-1/2(reDf4) and cst-1/2(re479)*
RF20	CAATCGAGGAGTGGGATCAAGUUUUAGAGC UAUGCUGUUUUG	crRNA for C term tag *yap-1(re269[yap-1::mNG::2xflag])*

LH005 GCK-2 crRNA-1	CAAAGUAGUCAAACUCGAGGGUUUUAGAGC UAUGCUGUUUUG	crRNA for *gck-2(re483) and gck-2(re427)*
LH006 GCK-2 crRNA-2	GUGACAUUGGUUGCGAUCGGGUUUUAGAG CUAUGCUGUUUUG	crRNA for *gck-2(re483)*
crRNA guide for mig-15 5' end	GTCGAGTCCTGATGACGACAGUUUUAGAGC UAUGCUGUUUUG	crRNA for tag *mig-15(re435[mTurquoise2::2 xmyc::AID*::mig-15])*
dpy-10 crRNA	GCUACCAUAGGCACCACGAGGUUUUAGAGC UAUGCUGUUUUG	crRNA for *dpy-10*
tracrRNA	AACAGCAUAGCAAGUUAAAAUAAGGCUAGU CCGUUAUCAACUUGAAAAAGUGGCACCGAG UCGGUGCUUUUUUU	universal

Table S4: Plasmids

Plasmid	Use	Note
pNR pBS mNG::2xFLAG	tag *yap-1(re269[yap-1::mNG::2xflag])*	sequence verified
pYW2 pBS mTurq2::2xMYC::AID*	tag *wts-1(re419[mTurqoise2::2xmyc::AID*::wts-1])* and *mig-15(re435[mTurquoise2::2xmyc::AID*::mig-15])*	sequence verified
pREW2	*luciferase(RNAi)*	sequence verified
I-5K13	*wts-1(RNAi)* (T20F10.1)	sequence verified
X-5M19	*yap-1(RNAi)* RNAi (F13E6.4)	sequence verified
X-11E16	*egl-44(RNAi)* (F28B12.2a)	sequence verified
I-1K04	*pop-1(RNAi).* (W10C8.2)	sequence verified