# Expanding diversity of virome associated with ammonium-oxidizing archaea and related lineages

Kaiyang Zheng[1], Yantao Liang[1,2,3*], Fabai Wu[4*], Jiaxiang Li[1], Mario López-Pérez[5], David Paez-Espino[6], Andrew McMinn[1,3,7*], Min Wang[1,2,3,8*]

[1] College of Marine Life Sciences, MoE Key Laboratory of Evolution & Marine Biodiversity, Institute of Evolution and Marine Biodiversity, Frontiers Science Centre for Deep Ocean Multispheres and Earth System, Centre for Ocean Carbon Neutrality, Ocean University of China, Qingdao, China.

[2] Haide College, Ocean University of China, Qingdao, China.

[3] UMT-OUC Joint Centre for Marine Studies, Qingdao, China.

[4] Faculty of Science, Eastern Institute of Technology, Ningbo, China

[5] Evolutionary Genomics Group, División de Microbiología, Universidad Miguel Hernández, San Juan, Spain

[6] Ancilia Biosciences, Co. Ltd., Alexandria Launchlabs, New York, USA.

[7] Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia.

[8] The Affiliated Hospital of Qingdao University, Qingdao, China.

* Correspondence:

Yantao Liang, Email: **liangyantao@ouc.edu.cn**

Fabai Wu, Email: **fabaiwu@eitech.edu.cn**

Andrew McMinn, Email: **andrew.mcminn@utas.edu.au**

Min Wang, Email: **mingwang@ouc.edu.cn**

## Abstract

*Nitrososphaeria*, an archaeal class containing all ammonium-oxidizing archaea (AOA) and their closely related clades (e.g. *Ca. Caldarchaeales*), is a widely distributed archaeal lineage across diverse ecosystems. Due to the difficulty of isolation and cultivation, the viral diversity and evolution associated with non-marine AOA lineages and *Ca. Caldarchaeales* remain largely uncharacterized. Here, based on globally sampled genomes and metagenomes, we identified 67 and 232 species-level viruses or mobile genetic elements (MGEs) associated with AOA, including 104 sequences longer than 10 kb. Notably, 77 and 182 previously unrecognized species-level viruses and MGEs associated with *Ca. Caldarchaeales* were identified, including 84 sequences longer than 10 kb. AOA-associated viruses were classified into eight viral families, with five representing newly identified head-tailed lineages; *Caldarchaeales*-associated viruses were classified into eight families, including four novel head-tailed and one novel rod-shaped family. Phylogenetic inference revealed that the *Caldarchaeales*-associated rod-shaped viral family (*Demeterviridae*) shares a common ancestor with rod-shaped viruses of methanogens and they exhibit a unique phenomenon of major capsid gene duplication. Functional genomics analyses further revealed that these novel rod-shaped viruses encode virus-specific rare quinone biosynthesis genes and arsenic response transcriptional regulator genes, providing potential clues to their mechanisms of adaptation to hyperthermal environments. Collectively, our work uncovers the hidden genomic, phylogenetic and functional diversity of *Nitrososphaeria* viromes for the first time and closes a long-standing gap in archaeal virome research.

## Introduction

The *Nitrososphaeria* is an archaeal class within *Thermoprotei*, encompassing the previously defined *Thaumarchaeota* and *Augarchaeota*, both of which are important members of the *Thermoproteati* (previously known as TACK superphylum)[1,2]. In many aquatic ecosystems, *Nitrososphaeria* constitute up to 40% of microorganisms, contributing significantly to ammonia oxidation, thus heavily influencing nitrogen cycling [3-5]. *Nitrososphaeria* includes all ammonia-oxidizing archaea (AOA) [6] and utilizes ammonia oxidation pathways as their primary energy source [7,8]. Compared to ammonia-oxidizing bacteria, AOA thrive in low-ammonia conditions due to higher ammonium affinity, dominating nitrogen-limited environments as efficient "nitrogen thieves" [9-14]. The ancestors of AOA are considered to have originated in terrestrial environments and to have adapted aerobically after the Great Oxygenation Event [13,15,16]. Subsequently, these archaea became key microbial players across diverse ecosystems and make a substantial contribution to carbon fixation and ammonia oxidation [8,15,17-21]. As the sister clade of AOA, *Ca. Caldarchaeales* shares substantial genetic content involved in sulphur oxidation and ammonia oxidation, suggesting an ancestral ecological transition of AOA from hyperthermal to moderate environments [22]. Primarily inhabiting hydrothermal vents and hot springs, *Ca. Caldarchaeales* influences carbon and sulphur cycling, nutrient transformation and microbial community structures, playing critical ecological roles in extreme environments [23,24].

Viruses and mobile genetic elements (MGEs) exhibit complex morphological and genomic diversity, significantly shaping archaeal genomes and their ecological footprint [25,26]. Due to cultivation challenges, environmental metagenomics has become the primary method for exploring archaeal viral diversity. Recent studies using environmental metagenomes have greatly expanded viral diversity associated with the Asgard super-phylum, methanogens and DPANN super-phylum, addressing gaps in viral evolution and host-virus interactions [27-31]. Over 50 AOA-associated viral or MGE sequences from marine samples have so far been identified, revealing diverse viral elements that influence ammonia oxidation and host genomic organization [32-38]. However, the limited number of case studies prevents a comprehensive profiling of the diversity, evolution and functions of these archaeal viruses. The absence of viruses associated with extremophilic AOA lineages (e.g.,

*Nitrosocaldales* and *Nitrosotalea*) poses a challenge for resolving phylogenetic placements of AOA viruses within the virosphere. In particular, as a phylogenetic sister clade closely related to AOA and one of the main lineages constituting the TACK super-phylum, *Caldarchaeales*-associated viruses or MEGs remain underexplored and represent a major enigma within virology.

To address this knowledge gap, we systematically explored *Nitrososphaeria*-associated viruses or MGEs diversity based on extensive global genomic and metagenomic datasets. Unlike previous studies that focused only on viruses from a single AOA lineage [32-36], a comprehensive analysis of the *Nitrososphaeria*-associated virome was conducted, yielding systematic insights into its genomics, phylogenetics and virus–host interactions. Through the establishment of a genomic catalogue comprising 781 viral or MGE sequences (including 659 species-level sequences), we revealed ten previously unrecognized archaeal viral families, including a group of rod-shaped viruses associated with *Ca. Caldarchaeales*, thereby providing new insights into the archaeal virosphere.

**Results**

**Overview of the genomic catalogue of *Nitrososphaeria*-associated viruses or MGEs**

*Nitrososphaeria* genome database containing 2,137 genomes and metagenome-assembled genomes

(MAGs) was constructed from NCBI GenBank and IMG/M databases (Fig. 1a and Supplementary

Note 1)[39,40]. The virus or MGE database comprised over 15 million contigs from IMG/VR v.4,

together with 170 metagenome-assembled contigs originating from hyperthermal, abyssal/hadal

ocean, hypersaline, frozen, and acidic environments, including 17 from our published and

unpublished studies (Supplementary Dataset 1) [39,41-44]. By integrating CRISPR spacer matches and

protein-sharing analysis, 781 *Nitrososphaeria*-associated viruses or MGEs were identified, including

747 extrachromosomal and 34 integrated ones. These were grouped into 659 species-level viral

operative taxonomic units (vOTUs), comprising 160 vOTUs with and 499 MGE OTUs without

known viral capsid proteins (Fig. 1a, Supplementary Dataset 2). In addition, one vOTU and 39 MGE

OTUs associated with AOA and 76 vOTUs and 178 MGE OTUs associated with *Ca. Caldarchaeales*

were newly identified, respectively.

*Nitrososphaeria*-associated viruses or MGEs are widely distributed from tropical to polar

regions and across terrestrial to marine environments (Fig. 1b). Most of them originated from

geothermal, marine, soil and hydrothermal environments, aligning with habitats of AOA,

deeply-branched AOA-related lineage and *Ca. Caldarchaeales* (Fig. 1b) [7,20,22,23,45-48]. Overall, this

genome catalogue revealed novel viruses associated with nine new archaeal lineages within

*Nitrososphaeria*, including *Ca. Caldarchaeales*, *Nitrosocaldales*, *Nitrosotalea* and other

AOA-related lineages (also known as non-AO *Thaumarchaeota*), thereby expanding the host

spectrum from two to eleven lineages [32-38,49].

**Ten novel viral families are associated with *Nitrososphaeria***

*Nitrososphaeria*-associated viruses or MGEs exhibit distinct differences in genome content

compared with other viruses. The genome similarity network (sharing at least 20% of protein clusters)

yielded 216 groups, with *Nitrososphaeria* contributing to 83 of them (Fig. 2a). These groups

included 40 AOA-specific and 24 *Ca. Caldarchaeales*-specific groups, suggesting the distinctiveness

of *Nitrososphaeria*-associated viral or MGE genomes within the virosphere. At the

genus-to-subfamily viral cluster (VC) level, 95.3% of *Nitrososphaeria*-associated viruses or MGEs

formed 121 independent VCs outside of the known ones. However, some *Nitrososphaeria*-associated

viruses exhibited close genome-content relationships with other archaeal viruses. Ten shared

significant genomic similarity (≥20% shared protein clusters) with established ANME-1-associated

viral families (*Chaacviridae*, *Ekchuahviridae* and *Itzamnaviridae*) (Figs. 2a and 2b and

Supplementary Dataset 3) [29,30].

Proteome-based hierarchical clustering was employed to further characterize novel families from

the *Nitrososphaeria*-associated virome. Sixteen viral families were found to be associated with

*Nitrososphaeria*, including ten previously undescribed ones. Except for the 31 vOTUs that were

assigned to established viral families (Fig. 2a and Supplementary Dataset 2), 68 vOTUs were

classified into newly named viral families here, they include: one rod-shaped viral family

(*Demeterviridae*) and four head-tailed viral families (*Erebusviridae*, *Hypnosviridae*, *Morosviridae*

and *Thanatosviridae*), which were associated with *Ca. Caldarchaeales* and five head-tailed family

(*Aetherviridae*, *Alboranviridae*, *Apateviridae*, *Baleariviridae*, *Tyrrheniaviridae*) associated with

AOA (Fig. 3). In addition, 21 MGE families without known capsid proteins were identified,

indicating a more complex diversity of the *Nitrososphaeria*-associated virome (Fig. 3 and

Supplementary Dataset 2).

The core genomes of different viral families involve ORFs related to replication, structure,

packaging, and virus–host recognition. For *Demeterviridae*, in addition to the SIRV2-like capsid, the

trimeric autotransporter adhesin (Fig. 3b, Supplementary Dataset 4 and Extended Data 1), one of the

key elements of virus–host recognition, was also part of its core genome. For the other five

head-tailed viral families, their core genomes were primarily composed of ORFs associated with

essential viral functions, including replication, structure, packaging, and virus counter-defence

related proteins. The core genomes of these families tended to be clustered together within the

genome, suggesting their conserved functional modules.

**The *Nitrososphaeria*-associated viruses exhibit distinct phylogenetic placements within the *Adnaviria* and *Duplodnaviria***

Bayesian and Maximum-Likelihood inference with structural modelling was used to characterize the phylogenetic placements of *Nitrososphaeria*-associated viruses (Extended Data 2 and 3). Two *Caldarchaeales*-associated rod-shaped viruses, together with ANME-1-associated *Ahmunviridae*, formed a phylogenetic clade and identified as a sister group of *Demeterviridae* (Fig. 4a). In the structural similarity network, the capsid monomers and dimers of *Demeterviridae* both formed distinct groups (Figs. 5a and 5b, Extended Data 4). Unlike *Rudiviridae* and *Ahmunviridae*, the major capsid protein (MCP) genes of *Demeterviridae* appear to have undergone duplication events. Although the genomes of *Lipothrixviridae* and *Tristromaviridae* encode two MCP genes and form heterodimeric capsids[50], the MCPs encoded by *Demeterviridae* are more closely related to those of *Rudiviridae* and *Ahmunviridae* (Fig. 4a and Supplementary Figs. 1a and 1b). Considering that *Rudiviridae* and *Ahmunviridae* possess homodimeric capsids, the mode of capsid formation in *Demeterviridae* remains highly uncertain. Nevertheless, the occurrence of multiple copies of MCP genes in *Demeterviridae* represents a rare case within *Adnaviria*. Taken together, these lines of evidence indicate that *Demeterviridae* represent a novel family distinct from other rod-shaped viruses, characterized by unique genome content, phylogenetic placement and structural features of their capsid proteins.

*Nitrososphaeria*-associated head-tailed viruses exhibit typical polyphyletic characteristics. Most newly identified *Caldarchaeales*-associated viral families formed distinct monophyletic clades in phylogenetic trees of both MCP and terminase, such as *Morosviridae*, *Erebusviridae* and *Thanatosviridae* (Figs. 4b and 4e). The MCPs encoded by *Morosviridae* form three distinct structurally-similar-based groups (Fig. 5c and Extended Data 5), supporting the uniqueness observed in their phylogenetic inference. These lines of evidence indicate their novel phylogenetic placements within *Caudoviricetes*. Eight *Caldarchaeales*-associated vOTUs were assigned to *Ekchuahviridae*, which had previously been found only in association with ANME-1 [30]. *Ekchuahviridae* was phylogenetically clustered with other archaea viral families (*Ahpuchviridae*, *Halomagnusviridae* and *Haloferuviridae*), suggesting a possible common ancestor of these archaeal head-tailed viruses (Fig. 4c). Three AOA-associated viral families belong to established families, including *Thaspiviridae*,

*Yangangviridae* and *Kyanoviridae*. Although *Kyanoviridae* primarily infects cyanobacteria, previous

studies have also reported associations with other heterotrophic bacteria, such as *Ca. Pelagibacter*,

*Methylophilales* and *Ochrobactrum* [51-53]. Associations between *Kyanoviridae* and AOA were

supported by both phylogenetic evidence and the structural similarity of the MCP (Figs. 4d, 4f and

5c), thereby extending the host spectrum of this family to the archaeal domain for the first time.

*Aetherviridae* is a newly defined AOA-associated viral family in this work, although members of this

family have been described in a previous study. This family encodes a unique ammonia

monooxygenase subunit C (amoC) gene [35]. *Alboranviridae*, *Baleariviridae* and *Tyrrheniaviridae*

were described as "Marthavirus" in a previous study, representing a group of AOA-related viruses

discovered in the Mediterranean Sea [34]. Although these viruses were classified into different families

based on their genome content (Fig. 3a), the MCP and terminase of *Alboranviridae* and

*Baleariviridae* are phylogenetically intertwined and form a sister clade with *Aetherviridae*. This

suggests that *Aetherviridae*, *Alboranviridae*, and *Baleariviridae* may share a common ancestor (Figs.

4b and 4e).

**Novel auxiliary genes encoded by *Caldarchaeales*-associated rod-shaped viruses**

The protein functional diversity of *Nitrososphaeria*-associated viruses or MGEs and their potential

impacts on the host were further explored. Following the recommendations of Martin et al., potential

auxiliary viral genes (AVGs) encoded by these viruses or MGEs were screened [54]. A total of 299

auxiliary regulatory genes (AReGs), 190 auxiliary metabolic genes (AMGs) and 48 auxiliary

physiological genes (APGs) were obtained (Supplementary Dataset 4). For AMGs,

2-polyprenyl-3-methyl-5-hydroxy-6-metoxy-1,4-benzoquinol methylase (UbiG) was found to be

encoded by four vOTUs from the newly identified rod-shaped viral family, *Demeterviridae*

(Supplementary Dataset 4). To the best of our knowledge, CoQ is rare in archaea and there is

currently no evidence that *Nitrososphaeria* or archaeal viruses possess a complete CoQ biosynthesis

pathway [23,55]. Homologues of *Demeterviridae*-encoded UbiG-like protein are extremely rare in

viruses, with no high-similarity homologues found in the nr database (alignment identity and

coverage both ≥ 50%), highlighting its novelty. These virus-encoded UbiG-like proteins form a

distinct monophyletic clade that groups with distantly related archaeal homologues in the

phylogenetic tree. This indicates their connection to archaea while retaining a clear virus-specific

phylogenetic monophyly (Fig. 6a). This finding provides a rare case of archaeal viruses encoding

genes involved in quinone electron carrier biosynthesis [56], suggesting that *Demeterviridae*-encoded

UbiG may be involved in modulating quinone electron carriers on the host membrane. For AReGs,

*Demeterviridae* was found to encode an ArsR-like transcriptional regulator, which was only

previously reported from spindle-shaped ssDNA viruses and head-tailed viruses [49,57]. The

*Demeterviridae*-encoded ArsR-like proteins forms a well-supported, long-branched monophyletic

clade with a pronounced stem branch in the phylogenetic tree and is evolutionarily distant from other

bacteriophages and archaeal viruses. This suggests their unique phylogenetic placement (Fig. 6b).

Regulators of archaeal filamentous or rod-shaped viruses are predominantly represented by

RHH-fold proteins such as SvtR (*Rudiviridae*) and AvtR (*Lipothrixviridae*) [58-60]. This result suggests

that *Caldarchaeales*-associated rod-shaped viral genomes may employ a similar mechanism of

transcriptional regulation as spindle-shaped viruses [57], which may represent an unrecognized

mechanism within this viral realm.

## Viruses or MGEs serve as the primary vectors rather than donors or receptors in horizontal gene transfers

Virus–host linkages identified by spacer–protospacer matching showed lineage specificity, with no

cross-lineage protospacer sharing observed (Supplementary Fig. 2a and Supplementary Dataset 5).

After establishing virus-host linkages by spacer-protospacer matching, the direction of horizontal

gene transfers (HGTs) mediated by viruses or MGEs were explored based on phylogenetic

topologies [61]. A total of 2,509 HGT events between viruses or MGEs and corresponding

*Nitrososphaeria* lineages were identified (Supplementary Dataset 6 and Extended Data 6) [61]. For

AOA, 68.0% of transfers are bidirectional (n = 1,359), while 27.3% originate from archaea as donors

(n = 546). Only 4.7% of transfers involve viruses or MGEs as donors (n = 94). Similarly, in *Ca.*

*Caldarchaeales*, 56.5% of transfers are bidirectional (n = 183), 34% originate from archaea (n = 110)

and 9.6% involve viruses or MGEs as donors (n = 31) (Fig. 6c). This suggests that viruses or MGEs

serve as the primary vectors in these transfers, rather than donors or receptors.

Functionally, most HGTs are related to glycan biosynthesis and metabolism, replication and

repair and carbohydrate metabolism, accounting for 60.5% of the total HGTs (Fig. 6d). Within

glycan biosynthesis and metabolism, more than half of the HGTs (n = 283) are associated with

glycosylation and are linked to Group 1.1a (Supplementary Dataset 4 and Supplementary Dataset 6).

The related functional proteins include all dolichyl-phosphate-mannose–protein

mannosyltransferases, glycosyl transferases (GTs), sulfatases and N-acetyl sugar amidotransferases.

Since glycosylation or glycoprotein modification on the surface of archaeal viral capsids or

envelopes is involved in host interactions or enhances structural stability [62-64], this provides evidence

for the assimilation of host functional proteins by *Nitrososphaeria*-associated viruses or MGEs. In

addition, 87.8% of the HGTs related to viral structure were found to be associated with *Ca.*

*Caldarchaeales*. These structural proteins include tail-related proteins and MCPs from *Erebusviridae*

and *Morosviridae*, further confirming the interactions between these proposed viral families and *Ca.*

*Caldarchaeales*.

**Discussion**

In this study, we provide a deeper understanding of the diversity of *Nitrososphaeria*-associated

viruses or MGEs, broadening the scope from previously reported AOA-associated viruses and

identified hundreds of previously unrecognized *Caldarchaeales*-associated vOTUs for the first time.

This genomic catalogue expands the known number of *Nitrososphaeria*-associated viruses or MGEs

by ten-fold [32-36]. The mining of *Caldarchaeales*-associated viral or MGE genomes fills a critical

knowledge gap of archaeal viromes. Ten previously unknown viral families were identified,

including a novel rod-shaped family, *Demeterviridae*. The unique evolutionary placement and

functional characteristics of this family were elucidated through comparative genomic and

phylogenetic analyses, providing new insights into the archaeal virosphere and viral evolution.

The core genome of *Demeterviridae*, in addition to the SIRV2-like capsid and glycosyl

transferase, also contains genes encoding trimeric autotransporter adhesins and several right-handed

parallel β-helix repeat proteins (RHBH) (Fig. 3b, Supplementary Dataset 4 and Extending Data 1).

Trimeric autotransporter adhesins are key mediators of prokaryote–host recognition [65], but have not

previously been reported in viruses. Similarly, RHBH structures are widely found in various

enzymes and adhesion factors involved in carbohydrate processing or cell surface recognition [66].

Typically, proteins associated with virus–host recognition are subject to intense selective pressure in the ongoing "arms race" against the frequent alterations of host membrane receptors, and thus are rarely incorporated into the core viral genome [67-69]. The retention of these genes in the core genome of *Demeterviridae* may result from co-evolutionary constraints imposed by other core genes. In other words, these virus–host recognition–related genes may have become integrated into the virion structural and assembly framework [70-72], forming part of a "structural community" whose evolutionary flexibility has been correspondingly reduced.

The phylogenetic analysis of *Nitrososphaeria*-associated viruses showed their evolutionary relationships with other viruses. In the past, only the capsids of *Lipothrixviridae* and *Tristromaviridae* were known to be composed of heterodimers[60]. Here, it was found that some *Caldarchaeales*-associated rod-shaped viruses encode two copies of MCP genes (Figs. 4a and 5b, Supplementary Figs. 1a and 1b). With respect to capsid assembly strategy, homodimers may provide symmetry and repetitiveness, thereby contributing to the maintenance of capsid stability [73,74]. Heterodimers may afford greater structural flexibility and diversity, enabling adaptation to varying environmental stresses and host defence mechanisms [75,76]. Given that *Rudiviridae* and *Ahmunviridae* encode only a single-copy MCP gene [30,50,60], it is possible that in *Demeterviridae* only one copy is functional, while the other may have been inactivated under natural selection [77-79]. Nevertheless, the actual capsid assembly mechanism of these rod-shaped viruses remains unknown and these inferences still require direct observation for confirmation. Some *Caldarchaeales*-associated viruses have close relationships with ANME-1-associated viruses or MGEs originating from hydrothermal vents (e.g. Methanophagales virus PBV300, *Ahmunviridae*) (Figs. 2a, 4c and 5b) [30], implying viral host-switching or non-specific transfers occurred between both archaeal lineages, which is common in the evolution of viruses [80-82].

Viruses have been found to encode diverse AVGs that manipulate host metabolism, thereby creating favourable conditions for completing their life cycles and indirectly contributing to global biogeochemical cycles [83-86]. In archaea-associated viruses, such as those infecting methanogenic archaea, Asgard superphylum, halophilic archaea, virus-encoded metabolic genes have been identified that potentially modulate archaeal methane-oxidization, sulfur metabolism, nucleotide and amino acid metabolisms [27,87,88]. Regarding AOA-associated viruses or MGEs, previous studies have discovered that viruses may manipulate host ammonia oxidation metabolism by carrying the *amoC*

gene and participate in host vitamin synthesis [34,35]. Through the exploration of functional repertoires

for *Nitrososphaeria*-associated viruses, we provide new insights into the functional mechanisms of

archaeal viruses. An important example is provided by viral ArsR-like and UbiG-like protein, which

represent a rare AReG and AMG, respectively, encoded by a previously unrecognized rod-shaped

viral family. If the viral ArsR-like protein is functional in the archaeal virocell, it might be able to

hijack the arsenic response of *Ca. Caldarchaeales*, thereby facilitating viral replication within the

host cell. Considering that the habitat of *Ca. Caldarchaeales* is rich in metals and metalloids

(including arsenic) and exhibits highly active redox dynamics[24,89,90], this process may be essential for

the completion of the *Demeterviridae* life cycle. Phylogenetically, *Demeterviridae*-encoded ArsR

falls on a long-branch, virus-specific clade supports their viral origin, rather than recent horizontal

gene transferring from archaea.

The more unusual finding is the viral carriage of UbiG-like protein, an enzyme implicated in

CoQ biosynthesis. Quinones (e.g. ubiquinone, menaquinone) are key electron carriers in membrane

redox chains, yet they are considered rare in archaea and corresponding viruses [23,55,91]. The detection

of *Demeterviridae*-encoded UbiG-like protein suggests a potential viral contribution to the redox

balance of archaeal membranes, such as altering the local pool of quinone electron carriers [55,92]. They

might influence electron flux, respiration efficiency, reactive oxygen species (ROS) levels, or even

redox signaling [92-94]. In hyperthermal habitats of *Ca. Caldarchaeales*, control of archaeal quinone

pools may confer a fitness advantage to related viruses by maintaining a more favorable intracellular

environment for replication. On the other hand, the *Demeterviridae*-encoded UbiG-like protein,

together with other methyltransferases encoded by archaeal viruses, may perform similar functions in

counteracting host defense mechanisms, representing an exaptation of archaeal gene functions by

viruses [81]. Since *Ca. Caldarchaeales* has not been observed to encode UbiG-like proteins, the origin

of these genes in *Demeterviridae* remains unknown. Overall, the discovery of *Demeterviridae* not

only expands the boundaries of the archaeal virosphere and offers new insights into viral evolution,

but also provides direction in how viruses might adapt to extreme environments in terms of

transcriptional regulation and host metabolic reprogramming.

Our results reinforce the view that viruses or MGEs act predominantly as gene vectors. This

supports the "virus world" model, considering viruses or other selfish genetic elements as

intermediaries in gene exchange rather than as main sources of new genes [95]. The prevalence of

bidirectional HGTs highlights reciprocal coevolution between viruses or MGEs and archaeal hosts,

evoking the arms-race dynamics observed in archaeal virus–host systems [96,97]. The functional

enrichment of glycan biosynthesis and glycosylation among transferred genes implies that viruses

may co-opt host surface modification machinery to refine capsid stability or host recognition—a

phenomenon similarly observed in eukaryotic viruses adapting envelope glycosylation [98,99]. It is

hypothesized that the frequent HGT events between archaea and viruses may have driven the

diversification of archaeal envelope biochemistry and the emergence of niche specialization in

ammonia☐oxidizing lineages.

In summary, the establishment of *Nitrososphaeria*-associated viromic catalogue and our further

analysis provides critical knowledge for future in-depth research into virus-archaea interactions.

**Methods**

**Assembly of *Nitrososphaeria*-associated CRISPR database**

The *Nitrososphaeria* phylum comprises several major clades: AOA, other AOA-related lineages and

*Ca. Caldarchaeales*. To produce a comprehensive genomic dataset, reference genomes were

retrieved from GenBank and IMG/M [39,40], encompassing a total of 2,137 genomic assemblages.

Specifically, 1,830 assemblages (comprising 322,784 genomic contigs) were obtained from GenBank;

307 assemblages (containing 23,395 genomic contigs) were sourced from IMG/M.

To identify CRISPR spacers within these genomes, similar to previous studies [27,28], CRASS

v.0.3.8 and MinCED v.0.4.2 (CRISPR recognition tool, CRT) were employed using the following

parameters to detect spacers and discard those with low-complexity and short length [100,101]: word size

= 7, spacer length = 26 and a minimum of three directed repeats (DRs). This step resulted in the

identification of 37,485 *Nitrososphaeria*-associated spacers, which were subsequently used for the

detection of viruses or MGEs and downstream analyses.

The metagenome-derived spacers that are associated with *Nitrososphaeria* were detected to

further expand the diversity of these CRISPR spacers. First, a total of 170 metagenome-derived

assemblages (including 58 from the Malaspina expedition and 22 from our previous studies) from

diverse ecosystems were retrieved and downloaded from IMG/M and ENA (European Nucleotide

Archive) (Supplementary Dataset 1)[40,43]. These metagenomes were from 23 different studies,

focusing on extreme natural habitats and including 92,037,046 assembled contigs. These include four

studies from deep-sea, three two from sea ice, four studies from a hypersaline environment and one

study from an acidic environment. Second, the CRISPR spacers found in these contigs were

identified using the same approach described above, producing 1,828,641 spacers and 1,918,441

DRs. Third, all DRs were aligned to *Nitrososphaeria* genomes using MMseqs2 with parameters [102]:

word size = 7, pairwise alignment identity = 1, percentage coverage of DRs = 1, mismatches = 1.

This step produced 196,551 spacers derived from 170 metagenomes. Finally, this process generated a

*Nitrososphaeria*-associated spacer database containing 234,036 spacer sequences (Fig. 1a and

Extended Data 7).

As in previous studies [28], CRISPR-Cas cassette prediction and subtyping for archaeal contigs

and genomes were conducted using CRISPRcasIdentifier v1.0.0 [103], employing a combination of

Classification and Regression Trees (CART), Support Vector Machine (SVM) and Extremely

Randomized Trees (ERT) algorithms for classification and regression. The analysis was performed

with the following parameters: –mode ERT SVM CART, -hmm HMM1 HMM2 HMM3 HMM4

HMM5. Subtype assignment was determined through a consensus-based approach, where only

subtype classifications consistently predicted by all three models (CART, SVM and ERT) were

accepted. The final subtype was selected based on the highest mean probability and the lowest

standard deviation among the predictions (Supplementary Fig. 2b and Supplementary Dataset 7).

**Mining and filtering *Nitrososphaeria*-associated viruses or MGEs from IMG/VR (v.4)**

The IMG/VR (v.4) database, contigs from 170 metagenomes and archaeal genomes served as the

primary resources for mining *Nitrososphaeria*-associated viral or MGE genomes. At first, the

archaeal viral or MGE mining pipeline was validated using CRISPR spacers derived from archaeal

genomes and host predictions from the IMG/VR (v.4) database and other studies to minimize false

positives [33-36,38,104,105]. The 37,485 CRISPR spacers from *Nitrososphaeria* genomes were aligned to

IMG/VR (v.4) database using MMseqs2 with the parameters: word size = 7, pairwise alignment

identity = 0.9, percentage coverage of spacers = 0.9, mismatches = 1, resulting in 368 targeted.

Among these results, the hosts of 46 viruses or MGEs were also predicted to be *Nitrososphaeria* in

IMG/VR (v.4). Due to some viral contigs being assigned to non-*Nitrososphaeria* hosts in IMG/VR (v.4) metadata, the genome-content similarity between viruses or MGEs and hosts was applied to reduce the false positives: i) the open reading frames (ORFs) of these viral contigs were predicted by Prodigal v.2.6.3 and aligned to the NR database (2024.05.20) with 1e-5 as the E-value cut-off [106]; ii) the *Nitrososphaeria*-related hits were counted and normalized based on viral contig lengths; iii) for each genome, it was observed that at least 0.2 ORFs per 1-kb genome could be matched to *Nitrososphaeria* (Supplementary Dataset 2); iv) further validation was conducted to assess the applicability of this screening criterion to previously reported *Nitrososphaeria*-associated viruses or MGEs:

1) the 48 identified *Nitrososphaeria*-associated viral or MGE genomes were downloaded from GenBank or provided by data owner;

2) the 46 representative genomes were generated from 48 *Nitrososphaeria*-associated viral genomes through MMseqs2 with parameters [102]: pairwise alignment identity = 0.95, pairwise alignment coverage for shorter sequences = 0.9;

3) the 44 out of 46 viral genomes derived from GenBank meet this criterion. Two isolated spindle-shaped viruses (NSV4 and NSV5) failed to pass the test [37]. This indicates that the approach is also applicable to most previously reported *Nitrososphaeria*-associated viral or MGE genomes.

After validating the filtering approach, it was used as a cut-off for viral-host genome-content similarity to further filter the results obtained from spacer-protospacer matching. By applying this approach, 171 *Nitrososphaeria*-associated viruses or MGEs were filtered from 368 spacer-targeted viral or MGE candidates. Similar to previous studies [29,107], all relevant archaeal CRISPR-Cas systems were evaluated using CRISPRCasFinder with default parameters [108]. Among the identified CRISPR arrays, 91.2% (n = 301) exhibited an evidence level of 4, characterized by a repeat conservation index exceeding 70% and spacer identity below 8% (Supplementary Dataset 2). The remaining 29 viruses or MGEs, corresponding to evidence levels below 4 for CRISPR arrays, were retained for two reasons: 1) they passed genome-content similarity filtering; 2) in genome-wide analyses, they clustered with other viruses or MGEs possessing CRISPR arrays with an evidence level of 4.

In addition, IMG/VR (v.4) contains 1,457 viruses or MGEs that were predicted to be associated

with *Nitrososphaeria*, although these contigs were not retrieved through spacer-protospacer matching. The 1,457 viruses or MGEs predicted to be associated with *Nitrososphaeria* in IMG/VR (v.4) were tested. First, 699 representative contigs were clustered using MMSeq2 with parameters: pairwise alignment identity = 0.9, pairwise alignment coverage for shorter sequences = 0.9. Second, these contigs were tested by genome-content similarity calculation, resulting in 441 *Nitrososphaeria*-associated viruses or MGEs. Finally, 612 *Nitrososphaeria*-associated viruses or MGEs were identified and filtered from IMG/VR (v.4).The completeness of these viral or MGEs were evaluate using CheckV [109].

**Expanded mining of *Nitrososphaeria*-associated viruses or MGEs from 170 metagenomes**

The *Nitrososphaeria*-associated CRISPR spacer database was aligned to from 170 metagenomes using MMseqs2 with the same approach and parameters [102]. Contigs containing DRs and those shorter than 2-kb were removed before performing alignment. This resulted in 1,951 contigs matched by *Nitrososphaeria*-associated CRISPR spacers. To remove duplications, 1,951 contigs were aligned to previously identified *Nitrosospaherota*-associated viruses or MGEs set using MMseqs2 with same parameters same parameters [102]. This step generated 1,895 contigs, including 1,396 representative contigs clustered using the same approach described before. To recall as many as possible viruses or MGEs, these contigs were used to filter *Nitrososphaeria*-associated viruses or MGEs using three multiple viral identification pipelines, including geNomad, VirSorter2 and DeepVirFinder[110-112]. Multiple pipelines were employed for virus identification based on the following considerations: 1) geNomad demonstrates superior overall classification performance compared to other pipelines [111]; 2) VirSorter2 (VS2) is recommended when integrating multiple pipelines [113]; 3) DeepVirFinder (DVF), utilizing an alignment-free approach, is well-suited for identifying unique viral sequences [110]; 4) the identification results from each pipeline complement one another, enhancing recall while mitigating cellular contamination in the dataset[114-116]. For each pipeline, precision was prioritized over sensitivity; therefore, stringent virus detection strategies were adopted to minimize potential false positives: 1) geNomad was used to detect viral or MGE contigs using conserved mode ("--conservative"), resulting in a more strict set of viruses with high confidence; 2) VS2 was used to detect viral contigs under high-confidence mode ('--high-confidence-only'), retaining results with

classification score ≥ 0.9 and classification score ≤ 0.7 with viral hallmark ≥ 1; 3) The DVF used to

detect viral contig with parameters: score ≥ 0.9 and p-value ≤ 0.01. To validate this hybrid viral

detection pipeline, 20% of sequences were randomly selected from previously identified

*Nitrososphaeria*-associated viruses or MGE and were used as a test set to evaluate the efficiency of

the approach. Based on the test, this viral or MGE identification approach achieved a 74%

recruitment rate, which was considered acceptable for further viral or MGE identification

(Supplementary Dataset 8).

Using the validated hybrid viral identification approach, 419 viral or MGE contigs were

identified from 1,396 metagenome-derived contigs. These contigs had already been targeted by

*Nitrososphaeria*-associated CRISPR spacers in previous steps. The viral-host genome-content

similarity was used to filter these contigs as described before, resulting 108

*Nitrososphaeria*-associated viral or MGEs. Relevant archaeal CRISPR-Cas systems of these contigs

were evaluated using CRISPRCasFinder with default parameters [108]. The completeness of these viral

or MGEs was evaluated using CheckV [109].

**Detection of viruses or MGEs integrated in *Nitrososphaeria* genomes**

Proviruses integrated into *Nitrososphaeria* genomes were identified using the "find-proviruses"

program in geNomad [111]. Integrases were first detected with the MMseqs2 plugin in geNomad [102],

applying an E-value cut-off of 1e-5. Additional parameters were set as follows: a minimum

gene-level score of 0.4 for flagging proviral genes using the conditional random field model; a

minimum total virus marker score of 12 for proviruses lacking integrases or not located at scaffold

edges; a minimum total virus marker score of 8 for proviruses encoding integrases or located at

scaffold edges; a maximum allowed distance of 10-kb between provirus boundaries and integrases

for boundary extension; and a maximum allowed distance of 5-kb between provirus boundaries and

tRNAs for boundary extension. This process resulted in the identification of 30

*Nitrososphaeria*-associated proviruses.

To remove redundancy, the 30 proviral or MGE genomes performed self-alignment and were

compared against previously identified *Nitrososphaeria*-associated viral or MGE genomes using the

same approach and parameters described above. This filtering step yielded 24 non-redundant

*Nitrososphaeria*-associated proviruses (Supplementary Dataset 2).

**Host assignments of *Nitrososphaeria*-associated viruses or MGEs**

The host assignments of *Nitrososphaeria*-associated viruses or MGEs were primarily based on: 1)

CRISPR-spacers-associated host genomes; 2) information provided by previous studies; 3)

genome-content similarity between viruses or MGEs and hosts. For *Nitrososphaeria*-associated

viruses or MGEs that match CRISPR spacers from multiple distinct hosts, similar with previous

study [29], the host of a virus or MGE targeted by CRISPR spacers was assigned based on the

taxonomy of the genome from which the spacer with the highest identity originated. For

*Nitrososphaeria*-associated viruses or MGEs that were identified based on oligonucleotide pattern

similarity, the most possible host genomes associated with these viruses or MGEs were determined

through alignment against *Nitrososphaeria* genomes using MMseqs2 with the parameters: 1e-5 as

the E-value, iteratively searched two times [102]. A host genome was linked to a viral or MGE genome

if they have the most shared ORFs. This approach was also applied in IMG/VR (v.4) [104,117]. For

*Nitrososphaeria*-associated viruses or MGEs reported in previous studies, some host information was

retrieved from related literature. For those viruses or MGEs with uncertain host information [35], the

related viral ORFs were subject to MMseqs2 against all *Nitrososphaeria* genomes with the same

approach. The 35 Marthaviruses derived from López-Pérez et al. (PRJNA484324) have been

authorized for use and are included in this study [34]. The archaeal host phylogenetic tree was

calculated based on 46 concatenated ribosomal proteins by IQ-Tree2 (Supplementary Note 2)[118]. The

archaeal lineages were determined based on their phylogenetic placements and previous studies

[2,8,21,22,24,119].

**Comparative genomics and family-/species-level demarcation of *Nitrososphaeria*-associated**

**viruses or MGEs**

Similar to previous studies [27,29,31], new families were assigned based on the genome-content

similarity network and the genome-wide proteomic tree. Using DNA virus sequences longer than

2-kb from NCBI RefSeq v.223 as reference genomes, vConTACT2 was applied (with default

parameters) to compute the genome-content-based similarity network of those

*Nitrososphaeria*-associated viruses or MGEs. *Nitrososphaeria*-associated viral or MGE genome pairs

that shared at least 20% protein clusters and encoded at least six genes were filtered from

genome-content-based similarity network, generating a subset of network, which was used to guide

taxonomic assignments. In the network, viruses not related to *Nitrososphaeria*-associated viruses or

MGEs were discarded to achieve a clear layout. Genome-wide proteomic tree of those genomes

included in the subset of network were calculated by ViPTree v.4. The viruses or MGEs were

assigned to the same family if they were represented by a cohesive group in the

genome-content-based similarity network and a monophyletic group in genome-wide proteomic tree

(with threshold 0.05) [29]. The network and tree were visualized by Cytoscape v.3.10.2 and iToL v.6,

respectively [120,121]. The nomenclature of the proposed viral family was referred to Cthulhu mythos

(Supplementary Note 3).

**Annotation of viral or MGEs genomes**

Viral or MGE species (vOTUs or MGE OTUs) were defined using 95% nucleotide identity threshold

over 90% of the full length using MMseqs2. The family-level taxonomic assignments of

*Nitrososphaeria*-associated viruses or MGEs based on reference taxonomic database were assigned

using 1) VITAP with custom database: this database includes all viral reference genomes collected in

ICTV VMR-MSL40, high-quality vOTUs with genus-level taxonomy collected in IMG/VR (v.4) and

all viral genomes collected in NCBI RefSeq v.223 [104,122,123]; 2) CAT with NCBI RefSeq v.223

database [123,124]; 3) geNomad with default parameters [111]. The CAT only assigned class/family level

units for 10 contigs; the geNomad and VITAP assigned class/family level units for 683 and 676

contigs, respectively. Hence, the taxonomic assignments were mainly based on geNomad and VITAP.

If geNomad or VITAP showed consistency at higher taxonomic levels (Phylum or Class), but one of

the pipelines produced annotations at lower taxonomic levels, then the results with the lower

taxonomic level annotations were used. The results with discrepancies between VITAP and geNomad

were compared against the NCBI RefSeq v.223 using MMseqs2 with default parameters and

subsequently subjected to manual verification. The consensus VCs generated from vConTACT2

were used to expanded and furtherly curated the viral taxonomic assignment [125]. The members of

each proposed family were furtherly expanded based on consensus vOTU.

ORFs encoded in *Nitrososphaeria*-associated viruses or MGEs were predicted by Prodigal

v.2.6.3 with metagenome mode and annotated using the KEGG Ortholog database (January, 2025) [106],

Pfam-A v.35, COG2020, Swiss-Prot database (January, 2025), VOG223 database and NR database

(August, 2025) [126-128]. Potential AVGs encoded by these viruses or MGEs with length over 10-kb

were screened based on proposal from Martin et al.[54]. The functional screening criteria excluded all

genes associated with viral core functions (replication, recombination, repair, transcription,

translation, packaging and lysis), nucleotide metabolism and one-carbon pool processes, nucleic acid

modification and immune evasion, structural and host recognition functions (including all CAZymes

and polysaccharide synthesis related fucntions), dNTP supply, as well as various proteases or

multifunctional enzymes. In addition, candidate genes were filtered by examining whether viral

genes were present in their flanking regions. Considering the incompleteness of viral contigs, once

an AVG was confirmed, its homologues were also considered as AVGs, even if no viral genes were

present in the flanking regions. The maximum-likelihood phylogenetic trees of ArsR and UbiG were

constructed. For ArsR, protein sequences from archaea, bacteria, bacteriophages and archaeal viruses

were retrieved from InterPro (PF01022). After merging these with the ArsR protein sequences

encoded by rod-shaped viruses, CD-HIT v. 4.8 was used to cluster the sequences from different

sources (at 0.5 pairwise identity with global alignment). Seqkit v.2.5 was then applied to randomly

sample 100 sequences from each category [129], or to include all sequences if fewer than 100 were

available. Multiple sequence alignment (MSA) was performed with MUSCLE5 [130] and trimAl v.1.4

was used to remove sites with more than 30% gaps [131]. Maximum-likelihood phylogenetic inference

of the MSAs was carried out with IQ-TREE2 [118], with 1,000 bootstraps and LG+F+R10 as the

best-fit substitution model (Extended Data 8). For UbiG, protein sequences from archaea, bacteria

and eukaryotes were retrieved from InterPro (IPR004951). These were merged with rod-shaped

virus-encoded UbiG-like sequences and maximum-likelihood phylogenetic inference was performed

using a similar procedure, with Blosum62+R10 as the best-fit substitution model (Extended Data 8).

The maximum-likelihood phylogenetic trees were visualized with iTOL v.6 [121].

**Determining of *Nitrososphaeria*-associated viral or MGE horizontal gene transfer events**

HGT events between *Nitrososphaeria*-associated viruses (vOTUs) and their archaeal hosts were

identified following an established pipeline [61]. After removal of proviral sequences, viral and

archaeal sequences were aligned and shared protein families (PFs) were identified. Potential

contamination was rigorously screened based on lineage, GC content, scaffold length and genomic

context, retaining only robust alignments (Supplementary Note 4 and Supplementary Dataset 9).

Filtered protein families underwent phylogenetic analysis using MUSCLE5 alignment, trimAl

v.1.4 trimming and IQ-TREE2 tree construction with the LG+F+R5 model. Trees were rooted via

minimal ancestral deviation and potential HGT events were classified based on the phylogenetic

relationships between viral and archaeal sequences (Extended Data 6). Transfer direction labels

("v-from-a", "v-to-a", "v-?-a" and "non-HGT") were assigned following predefined criteria

considering sister and cousin clade compositions. Iterative screening was performed with temporary

reassignment of labels to ensure accurate HGT detection across multiple rounds of analysis

(Supplementary Note 4).

**Phylogenetic inference of *Nitrososphaeria*-associated viruses or MGEs**

To investigate the evolutionary relationships among rod-shaped and head-tailed viruses associated

with *Nitrososphaeria* within the virosphere context, phylogenetic analyses were performed based on

key viral proteins, including the SIRV2-like MCPs, HK97-like MCPs and terminase proteins. For

rod-shaped viruses, MCP sequences from 33 viruses or MGEs were aligned with 34 viral RefSeqs

v.223 from the *Adnaviria* realm using MMSeqs2 (E-value ≤1e-5). Notably, some viruses encoded

adjacent duplicate MCP subunits, including Metallosphaera rod-shaped virus 1 and

Saccharolobussolfataricus rod-shaped virus 1, along with five uncultured viral genomes (UViGs).

Considering the fundamental rod-shaped viral MCP structural unit as either homodimers or

heterodimers, phylogenetic trees were inferred based on concatenated MSAs generated using

MUSCLE5 (ensemble mode). The DUF732 domain-containing protein (PF05305), identified via

FoldSeek structural alignment as the closest homolog [132], served as an outgroup for rooting the

rod-shaped viral MCP phylogeny. HK97-MCP sequences exhibited significant diversity,

encompassing multiple families from Pfam-A v.35 and VOG223 [127,133]; thus, three dominant domains

(PF05065, PF07068 and VOG01305) were selected and aligned against *Duplodnaviria* proteins from

RefSeqs v.223. Appropriate outgroups were chosen based on structural and domain-based alignments

(e.g., PF06673 and PF00130). Similarly, terminase proteins were analyzed by selecting the top three

domains (K06909, VOG00195 and PF03237) and employing P-loop containing nucleoside

triphosphate hydrolases (SSF52540) as an outgroup. All MSAs were refined using trimAl v.1.4 and

trees were inferred by ML with IQ-Tree2 and BI using MrBayes [118,134], supported by 1,000 bootstrap

(ML) and convergence diagnostics (BI) (Extended Data 2 and 3). The topological comparisons

between BI and ML were performed using Phylo.io [135].

**Structure analysis of *Nitrososphaeria*-associated viruses or MGEs**

To further assess the novelty of capsid proteins from *Nitrososphaeria*-associated viruses or MGEs,

the HK97-like and SIRV2-like capsid proteins identified in this study were used to perform structural

prediction. A total of 5,153 head-tailed viral genomes from RefSeqs v.223 and 44 rod-shaped

archaeal viral genomes from GenBank (length ≥ 2,000 bp) were retrieved from GenBank (as of

November 31, 2024). ORFs in these genomes were predicted using Prodigal v.2.6.3 [106]. HK97-like

and SIRV2-like capsid proteins were identified using HMM-based searches against specific HMM

profiles (PF12193, PF05065, VOG01305 and PF07068) with an E-value cut-off of 1e-5 [136].

Subsequently, the HK97-like capsid proteins were clustered using MMSeqs2 with thresholds of 0.9

alignment coverage and identity [102]. Structures of the HK97-like and SIRV2-like capsid proteins

were predicted using AlphaFold2 implemented in ColabFold (Extended Data 3 and 4) [137] and the

models with the highest predicted local distance difference test (pLDDT) scores were selected as

best-predicted models. Structural alignments were conducted through all-to-all comparisons using

FoldSeek with an identity cut-off of 0.5 [132] and reciprocal hits were retained. The structural similarity

network was built based on bitscores (top 10% hits for each sequence were selected) calculated from

FoldSeek by Cytoscape [120,132].

**Data availability**

All nucleotide and protein sequences of viral or mobile genetic elements, spacers from the CRISPR database, phylogenetic trees, structural prediction files are available in FigShare (https://figshare.com/s/a2e94d686028b674a905). The related metagenomes used in this study can be accessed via the URL provided in Supplementary Dataset 1 through IMG/M. Related genomes from IMG/VR database is available at https://img.jgi.doe.gov/cgi-bin/vr/main.cgi. Other information and metadata are provided as Supplementary Datasets.

**Code availability**

No custom code was generated and used.

# Reference

1. Nunoura, T., Takaki, Y., Kakuta, J., Nishi, S., Sugahara, J., Kazama, H., Chee, G.J., Hattori, M., Kanai, A., Atomi, H., et al. (2011). Insights into the evolution of Archaea and eukaryotic protein modifier systems revealed by the genome of a novel archaeal group. Nucleic Acids Res *39*, 3204-3223. 10.1093/nar/gkq1228.

2. Spang, A., Caceres, E.F., and Ettema, T.J.G. (2017). Genomic exploration of the diversity, ecology, and evolution of the archaeal domain of life. Science *357*. 10.1126/science.aaf3883.

3. Alfreider, A., Grimus, V., Luger, M., Ekblad, A., Salcher, M.M., and Summerer, M. (2018). Autotrophic carbon fixation strategies used by nitrifying prokaryotes in freshwater lakes. FEMS Microbiol Ecol *94*. 10.1093/femsec/fiy163.

4. Klotz, F., Kitzinger, K., Ngugi, D.K., Busing, P., Littmann, S., Kuypers, M.M.M., Schink, B., and Pester, M. (2022). Quantification of archaea-driven freshwater nitrification from single cell to ecosystem levels. ISME J *16*, 1647-1656. 10.1038/s41396-022-01216-9.

5. Li, Y., Chen, J., Lin, Y., Zhong, C., Jing, H., and Liu, H. (2024). Thaumarchaeota from deep-sea methane seeps provide novel insights into their evolutionary history and ecological implications. Microbiome *12*, 197. 10.1186/s40168-024-01912-y.

6. Brochier-Armanet, C., Boussau, B., Gribaldo, S., and Forterre, P. (2008). Mesophilic Crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. Nat Rev Microbiol *6*, 245-252. 10.1038/nrmicro1852.

7. Pester, M., Schleper, C., and Wagner, M. (2011). The Thaumarchaeota: an emerging view of their phylogeny and ecophysiology. Curr Opin Microbiol *14*, 300-306. 10.1016/j.mib.2011.04.007.

8. Zhong, H., Lehtovirta-Morley, L., Liu, J., Zheng, Y., Lin, H., Song, D., Todd, J.D., Tian, J., and Zhang, X.H. (2020). Novel insights into the Thaumarchaeota in the deepest oceans: their metabolism and potential adaptation mechanisms. Microbiome *8*, 78. 10.1186/s40168-020-00849-2.

9. Kuypers, M.M.M., Marchant, H.K., and Kartal, B. (2018). The microbial nitrogen-cycling network. Nat Rev Microbiol *16*, 263-276. 10.1038/nrmicro.2018.9.

10. Prosser, J.I., and Nicol, G.W. (2012). Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. Trends Microbiol *20*, 523-531. 10.1016/j.tim.2012.08.001.

11. Martens-Habbena, W., Berube, P.M., Urakawa, H., de la Torre, J.R., and Stahl, D.A. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. Nature *461*, 976-979.

10.1038/nature08465.

12.    Stahl, D.A., and de la Torre, J.R. (2012). Physiology and diversity of ammonia-oxidizing archaea. Annu Rev Microbiol *66*, 83-101. 10.1146/annurev-micro-092611-150128.

13.    Walker, C.B., de la Torre, J.R., Klotz, M.G., Urakawa, H., Pinel, N., Arp, D.J., Brochier-Armanet, C., Chain, P.S., Chan, P.P., Gollabgir, A., et al. (2010). Nitrosopumilus maritimus genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchaea. Proc Natl Acad Sci U S A *107*, 8818-8823. 10.1073/pnas.0913533107.

14.    Könneke, M., Schubert, D.M., Brown, P.C., Hügler, M., Standfest, S., Schwander, T., Schada von Borzyskowski, L., Erb, T.J., Stahl, D.A., and Berg, I.A. (2014). Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway for CO2 fixation. Proceedings of the National Academy of Sciences *111*, 8239-8244.

15.    Ren, M., Feng, X., Huang, Y., Wang, H., Hu, Z., Clingenpeel, S., Swan, B.K., Fonseca, M.M., Posada, D., Stepanauskas, R., et al. (2019). Phylogenomics suggests oxygen availability as a driving force in Thaumarchaeota evolution. ISME J *13*, 2150-2161. 10.1038/s41396-019-0418-8.

16.    Abby, S.S., Kerou, M., and Schleper, C. (2020). Ancestral Reconstructions Decipher Major Adaptations of Ammonia-Oxidizing Archaea upon Radiation into Moderate Terrestrial and Marine Environments. mBio *11*. 10.1128/mBio.02371-20.

17.    Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., and Oakley, B.B. (2005). Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. Proc Natl Acad Sci U S A *102*, 14683-14688. 10.1073/pnas.0506625102.

18.    Nelkner, J., Huang, L., Lin, T.W., Schulz, A., Osterholz, B., Henke, C., Blom, J., Puhler, A., Sczyrba, A., and Schluter, A. (2023). Abundance, classification and genetic potential of Thaumarchaeota in metagenomes of European agricultural soils: a meta-analysis. Environ Microbiome *18*, 26. 10.1186/s40793-023-00479-9.

19.    Biggs-Weber, E., Aigle, A., Prosser, J.I., and Gubry-Rangin, C. (2020). Oxygen preference of deeply-rooted mesophilic thaumarchaeota in forest soil. Soil Biology and Biochemistry *148*, 107848. https://doi.org/10.1016/j.soilbio.2020.107848.

20.    Aylward, F.O., and Santoro, A.E. (2020). Heterotrophic Thaumarchaea with Small Genomes Are Widespread in the Dark Ocean. mSystems *5*. 10.1128/mSystems.00415-20.

21.    Zhang, R.Y., Wang, Y.R., Liu, R.L., Rhee, S.K., Zhao, G.P., and Quan, Z.X. (2024). Metagenomic characterization of a novel non-ammonia-oxidizing Thaumarchaeota from hadal sediment. Microbiome *12*, 7. 10.1186/s40168-023-01728-2.

22. Hua, Z.S., Qu, Y.N., Zhu, Q., Zhou, E.M., Qi, Y.L., Yin, Y.R., Rao, Y.Z., Tian, Y., Li, Y.X., Liu, L., et al. (2018). Genomic inference of the metabolism and evolution of the archaeal phylum Aigarchaeota. Nat Commun *9*, 2832. 10.1038/s41467-018-05284-4.

23. Beam, J.P., Jay, Z.J., Schmid, M.C., Rusch, D.B., Romine, M.F., Jennings Rde, M., Kozubal, M.A., Tringe, S.G., Wagner, M., and Inskeep, W.P. (2016). Ecophysiology of an uncultivated lineage of Aigarchaeota from an oxic, hot spring filamentous 'streamer' community. ISME J *10*, 210-224. 10.1038/ismej.2015.83.

24. Buessecker, S., Palmer, M., Lai, D., Dimapilis, J., Mayali, X., Mosier, D., Jiao, J.Y., Colman, D.R., Keller, L.M., St John, E., et al. (2022). An essential role for tungsten in the ecology and evolution of a previously uncultivated lineage of anaerobic, thermophilic Archaea. Nat Commun *13*, 3773. 10.1038/s41467-022-31452-8.

25. Prangishvili, D., Bamford, D.H., Forterre, P., Iranzo, J., Koonin, E.V., and Krupovic, M. (2017). The enigmatic archaeal virosphere. Nat Rev Microbiol *15*, 724-739. 10.1038/nrmicro.2017.125.

26. Al-Shayeb, B., Schoelmerich, M.C., West-Roberts, J., Valentin-Alvarado, L.E., Sachdeva, R., Mullen, S., Crits-Christoph, A., Wilkins, M.J., Williams, K.H., Doudna, J.A., and Banfield, J.F. (2022). Borgs are giant genetic elements with potential to expand metabolic capacity. Nature *610*, 731-736. 10.1038/s41586-022-05256-1.

27. Medvedeva, S., Sun, J., Yutin, N., Koonin, E.V., Nunoura, T., Rinke, C., and Krupovic, M. (2022). Three families of Asgard archaeal viruses identified in metagenome-assembled genomes. Nat Microbiol *7*, 962-973. 10.1038/s41564-022-01144-6.

28. Rambo, I.M., Langwig, M.V., Leao, P., De Anda, V., and Baker, B.J. (2022). Genomes of six viruses that infect Asgard archaea from deep-sea sediments. Nat Microbiol *7*, 953-961. 10.1038/s41564-022-01150-8.

29. Medvedeva, S., Borrel, G., Krupovic, M., and Gribaldo, S. (2023). A compendium of viruses from methanogenic archaea reveals their diversity and adaptations to the gut environment. Nat Microbiol *8*, 2170-2182. 10.1038/s41564-023-01485-w.

30. Laso-Perez, R., Wu, F., Cremiere, A., Speth, D.R., Magyar, J.S., Zhao, K., Krupovic, M., and Orphan, V.J. (2023). Evolutionary diversification of methanotrophic ANME-1 archaea and their expansive virome. Nat Microbiol. 10.1038/s41564-022-01297-4.

31. Sakai, H.D., Nur, N., Kato, S., Yuki, M., Shimizu, M., Itoh, T., Ohkuma, M., Suwanto, A., and Kurosawa, N. (2022). Insight into the symbiotic lifestyle of DPANN archaea revealed by cultivation and genome analyses. Proc Natl Acad Sci U S A *119*. 10.1073/pnas.2115449119.

32. Krupovic, M., Spang, A., Gribaldo, S., Forterre, P., and Schleper, C. (2011). A thaumarchaeal provirus testifies

for an ancient association of tailed viruses with archaea. Biochem Soc Trans *39*, 82-88. 10.1042/BST0390082.

33. Krupovic, M., Makarova, K.S., Wolf, Y.I., Medvedeva, S., Prangishvili, D., Forterre, P., and Koonin, E.V. (2019). Integrated mobile genetic elements in Thaumarchaeota. Environ Microbiol *21*, 2056-2078. 10.1111/1462-2920.14564.

34. Lopez-Perez, M., Haro-Moreno, J.M., de la Torre, J.R., and Rodriguez-Valera, F. (2019). Novel Caudovirales associated with Marine Group I Thaumarchaeota assembled from metagenomes. Environ Microbiol *21*, 1980-1988. 10.1111/1462-2920.14462.

35. Ahlgren, N.A., Fuchsman, C.A., Rocap, G., and Fuhrman, J.A. (2019). Discovery of several novel, widespread, and ecologically distinct marine Thaumarchaeota viruses that encode amoC nitrification genes. ISME J *13*, 618-631. 10.1038/s41396-018-0289-4.

36. Zhou, Y., Zhou, L., Yan, S., Chen, L., Krupovic, M., and Wang, Y. (2023). Diverse viruses of marine archaea discovered using metagenomics. Environ Microbiol *25*, 367-382. 10.1111/1462-2920.16287.

37. Kim, J.G., Kim, S.J., Cvirkaite-Krupovic, V., Yu, W.J., Gwak, J.H., Lopez-Perez, M., Rodriguez-Valera, F., Krupovic, M., Cho, J.C., and Rhee, S.K. (2019). Spindle-shaped viruses infect marine ammonia-oxidizing thaumarchaea. Proc Natl Acad Sci U S A *116*, 15645-15650. 10.1073/pnas.1905682116.

38. Liu, Y., Brandt, D., Ishino, S., Ishino, Y., Koonin, E.V., Kalinowski, J., Krupovic, M., and Prangishvili, D. (2019). New archaeal viruses discovered by metagenomic analysis of viral communities in enrichment cultures. Environ Microbiol *21*, 2002-2014. 10.1111/1462-2920.14479.

39. Nayfach, S., Roux, S., Seshadri, R., Udwary, D., Varghese, N., Schulz, F., Wu, D., Paez-Espino, D., Chen, I.M., Huntemann, M., et al. (2021). A genomic catalog of Earth's microbiomes. Nat Biotechnol *39*, 499-509. 10.1038/s41587-020-0718-6.

40. Chen, I.A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek, P., Ritter, S., Varghese, N., Seshadri, R., et al. (2021). The IMG/M data management and analysis system v.6.0: new tools and advanced capabilities. Nucleic Acids Res *49*, D751-D763. 10.1093/nar/gkaa939.

41. Sharma, S., Lee, M., Reinmann, C.S., Pumneo, J., Cutright, T.J., and Senko, J.M. (2020). Impact of acid mine drainage chemistry and microbiology on the development of efficient Fe removal activities. Chemosphere *249*, 126117. 10.1016/j.chemosphere.2020.126117.

42. Zhang, X., Xu, W., Liu, Y., Cai, M., Luo, Z., and Li, M. (2018). Metagenomics Reveals Microbial Diversity and Metabolic Potentials of Seawater and Surface Sediment From a Hadal Biosphere at the Yap Trench. Front Microbiol *9*, 2402. 10.3389/fmicb.2018.02402.

43.     Acinas, S.G., Sanchez, P., Salazar, G., Cornejo-Castillo, F.M., Sebastian, M., Logares, R., Royo-Llonch, M., Paoli, L., Sunagawa, S., Hingamp, P., et al. (2021). Deep ocean metagenomes provide insight into the metabolic architecture of bathypelagic microbial communities. Commun Biol *4*, 604. 10.1038/s42003-021-02112-2.

44.     Gao, C., Liang, Y., Jiang, Y., Paez-Espino, D., Han, M., Gu, C., Wang, M., Yang, Y., Liu, F., Yang, Q., et al. (2022). Virioplankton assemblages from challenger deep, the deepest place in the oceans. iScience *25*, 104680. 10.1016/j.isci.2022.104680.

45.     Karner, M.B., DeLong, E.F., and Karl, D.M. (2001). Archaeal dominance in the mesopelagic zone of the Pacific Ocean. Nature *409*, 507-510. 10.1038/35054051.

46.     Schattenhofer, M., Fuchs, B.M., Amann, R., Zubkov, M.V., Tarran, G.A., and Pernthaler, J. (2009). Latitudinal distribution of prokaryotic picoplankton populations in the Atlantic Ocean. Environ Microbiol *11*, 2078-2093. 10.1111/j.1462-2920.2009.01929.x.

47.     Pester, M., Rattei, T., Flechl, S., Grongroft, A., Richter, A., Overmann, J., Reinhold-Hurek, B., Loy, A., and Wagner, M. (2012). amoA-based consensus phylogeny of ammonia-oxidizing archaea and deep sequencing of amoA genes from soils of four different geographic regions. Environ Microbiol *14*, 525-539. 10.1111/j.1462-2920.2011.02666.x.

48.     Vetriani, C., Jannasch, H.W., MacGregor, B.J., Stahl, D.A., and Reysenbach, A.L. (1999). Population structure and phylogenetic characterization of marine benthic Archaea in deep-sea sediments. Appl Environ Microbiol *65*, 4375-4384. 10.1128/AEM.65.10.4375-4384.1999.

49.     Liu, Y., Demina, T.A., Roux, S., Aiewsakun, P., Kazlauskas, D., Simmonds, P., Prangishvili, D., Oksanen, H.M., and Krupovic, M. (2021). Diversity, taxonomy, and evolution of archaeal viruses of the class Caudoviricetes. PLoS Biol *19*, e3001442. 10.1371/journal.pbio.3001442.

50.     Goulet, A., Blangy, S., Redder, P., Prangishvili, D., Felisberto-Rodrigues, C., Forterre, P., Campanacci, V., and Cambillau, C. (2009). Acidianus filamentous virus 1 coat proteins display a helical fold spanning the filamentous archaeal viruses lineage. Proc Natl Acad Sci U S A *106*, 21155-21160. 10.1073/pnas.0909893106.

51.     Zhao, Y., Temperton, B., Thrash, J.C., Schwalbach, M.S., Vergin, K.L., Landry, Z.C., Ellisman, M., Deerinck, T., Sullivan, M.B., and Giovannoni, S.J. (2013). Abundant SAR11 viruses in the ocean. Nature *494*, 357-360. 10.1038/nature11921.

52.     Buchholz, H.H., Bolanos, L.M., Bell, A.G., Michelsen, M.L., Allen, M.J., and Temperton, B. (2022). A Novel and Ubiquitous Marine Methylophage Provides Insights into Viral-Host Coevolution and Possible Host-Range Expansion in Streamlined Marine Heterotrophic Bacteria. Appl Environ Microbiol *88*, e0025522.

10.1128/aem.00255-22.

53.  Decewicz, P., Golec, P., Szymczak, M., Radlinska, M., and Dziewit, L. (2020). Identification and Characterization of the First Virulent Phages, Including a Novel Jumbo Virus, Infecting Ochrobactrum spp. Int J Mol Sci *21*. 10.3390/ijms21062096.

54.  Martin, C., Emerson, J.B., Roux, S., and Anantharaman, K. (2025). A call for caution in the biological interpretation of viral auxiliary metabolic genes. Nat Microbiol *10*, 2122-2129. 10.1038/s41564-025-02095-4.

55.  Elling, F.J., Becker, K.W., Konneke, M., Schroder, J.M., Kellermann, M.Y., Thomm, M., and Hinrichs, K.U. (2016). Respiratory quinones in Archaea: phylogenetic distribution and application as biomarkers in the marine environment. Environ Microbiol *18*, 692-707. 10.1111/1462-2920.13086.

56.  Guerra, R.M., and Pagliarini, D.J. (2023). Coenzyme Q biochemistry and biosynthesis. Trends Biochem Sci *48*, 463-476. 10.1016/j.tibs.2022.12.006.

57.  Frols, S., Gordon, P.M., Panlilio, M.A., Schleper, C., and Sensen, C.W. (2007). Elucidating the transcription cycle of the UV-inducible hyperthermophilic archaeal virus SSV1 by DNA microarrays. Virology *365*, 48-59. 10.1016/j.virol.2007.03.033.

58.  Guilliere, F., Peixeiro, N., Kessler, A., Raynal, B., Desnoues, N., Keller, J., Delepierre, M., Prangishvili, D., Sezonov, G., and Guijarro, J.I. (2009). Structure, function, and targets of the transcriptional regulator SvtR from the hyperthermophilic archaeal virus SIRV1. J Biol Chem *284*, 22222-22237. 10.1074/jbc.M109.029850.

59.  Peixeiro, N., Keller, J., Collinet, B., Leulliot, N., Campanacci, V., Cortez, D., Cambillau, C., Nitta, K.R., Vincentelli, R., Forterre, P., et al. (2013). Structure and function of AvtR, a novel transcriptional regulator from a hyperthermophilic archaeal lipothrixvirus. J Virol *87*, 124-136. 10.1128/JVI.01306-12.

60.  Krupovic, M., Kuhn, J.H., Wang, F., Baquero, D.P., Dolja, V.V., Egelman, E.H., Prangishvili, D., and Koonin, E.V. (2021). Adnaviria: a New Realm for Archaeal Filamentous Viruses with Linear A-Form Double-Stranded DNA Genomes. J Virol *95*, e0067321. 10.1128/JVI.00673-21.

61.  Irwin, N.A.T., Pittis, A.A., Richards, T.A., and Keeling, P.J. (2022). Systematic evaluation of horizontal gene transfer between eukaryotes and viruses. Nat Microbiol *7*, 327-336. 10.1038/s41564-021-01026-3.

62.  Gebhard, L.J., Vershinin, Z., Alarcon-Schumacher, T., Eichler, J., and Erdmann, S. (2023). Influence of N-Glycosylation on Virus-Host Interactions in Halorubrum lacusprofundi. Viruses *15*. 10.3390/v15071469.

63.  Quemin, E.R., Pietila, M.K., Oksanen, H.M., Forterre, P., Rijpstra, W.I., Schouten, S., Bamford, D.H., Prangishvili, D., and Krupovic, M. (2015). Sulfolobus Spindle-Shaped Virus 1 Contains Glycosylated Capsid Proteins, a Cellular Chromatin Protein, and Host-Derived Lipids. J Virol *89*, 11681-11691.

10.1128/JVI.02270-15.

64. Wang, F., Cvirkaite-Krupovic, V., Kreutzberger, M.A.B., Su, Z., de Oliveira, G.A.P., Osinski, T., Sherman, N., DiMaio, F., Wall, J.S., Prangishvili, D., et al. (2019). An extensively glycosylated archaeal pilus survives extreme conditions. Nat Microbiol *4*, 1401-1410. 10.1038/s41564-019-0458-x.

65. Kiessling, A.R., Malik, A., and Goldman, A. (2020). Recent advances in the understanding of trimeric autotransporter adhesins. Med Microbiol Immunol *209*, 233-242. 10.1007/s00430-019-00652-3.

66. Burnim, A.A., Dufault-Thompson, K., and Jiang, X. (2024). The three-sided right-handed beta-helix is a versatile fold for glycan interactions. Glycobiology *34*. 10.1093/glycob/cwae037.

67. de Leeuw, M., Baron, M., Ben David, O., and Kushmaro, A. (2020). Molecular Insights into Bacteriophage Evolution toward Its Host. Viruses *12*. 10.3390/v12101132.

68. Yehl, K., Lemire, S., Yang, A.C., Ando, H., Mimee, M., Torres, M.T., de la Fuente-Nunez, C., and Lu, T.K. (2019). Engineering Phage Host-Range and Suppressing Bacterial Resistance through Phage Tail Fiber Mutagenesis. Cell *179*, 459-469 e459. 10.1016/j.cell.2019.09.015.

69. Taslem Mourosi, J., Awe, A., Guo, W., Batra, H., Ganesh, H., Wu, X., and Zhu, J. (2022). Understanding Bacteriophage Tail Fiber Interaction with Host Surface Receptor: The Key "Blueprint" for Reprogramming Phage Host Range. Int J Mol Sci *23*. 10.3390/ijms232012146.

70. Zheng, K., Liang, Y., Paez-Espino, D., Zou, X., Gao, C., Shao, H., Sung, Y.Y., Mok, W.J., Wong, L.L., Zhang, Y.Z., et al. (2023). Identification of hidden N4-like viruses and their interactions with hosts. mSystems *8*, e0019723. 10.1128/msystems.00197-23.

71. Champeimont, R., Laine, E., Hu, S.W., Penin, F., and Carbone, A. (2016). Coevolution analysis of Hepatitis C virus genome to identify the structural and functional dependency network of viral proteins. Sci Rep *6*, 26401. 10.1038/srep26401.

72. Krupovic, M., and Koonin, E.V. (2017). Multiple origins of viral capsid proteins from cellular ancestors. Proc Natl Acad Sci U S A *114*, E2401-E2410. 10.1073/pnas.1621061114.

73. Tubiana, T., Boulard, Y., and Bressanelli, S. (2017). Dynamics and asymmetry in the dimer of the norovirus major capsid protein. PLoS One *12*, e0182056. 10.1371/journal.pone.0182056.

74. Hadden, J.A., Perilla, J.R., Schlicksup, C.J., Venkatakrishnan, B., Zlotnick, A., and Schulten, K. (2018). All-atom molecular dynamics of the HBV capsid reveals insights into biological function and cryo-EM resolution limits. Elife *7*. 10.7554/eLife.32478.

75. Mueller, K.L. (2016). Structural insights into capsid flexibility. Science *354*, 1387-1389.

10.1126/science.354.6318.1387-p.

76. Budean, D., Almeida-Hernández, Y., Pandey, J., Pérez, J.M., Sánchez García, E., and Haglund, E. (2024). Exploring Chemokine Homodimer Stability: Structural Insights into CXC and CC Interfaces. Biophysica *4*, 561-572.

77. Brennan, G., Stoian, A.M.M., Yu, H., Rahman, M.J., Banerjee, S., Stroup, J.N., Park, C., Tazi, L., and Rothenburg, S. (2023). Molecular Mechanisms of Poxvirus Evolution. mBio *14*, e0152622. 10.1128/mbio.01526-22.

78. Monzon, S., Varona, S., Negredo, A., Vidal-Freire, S., Patino-Galindo, J.A., Ferressini-Gerpe, N., Zaballos, A., Orviz, E., Ayerdi, O., Munoz-Gomez, A., et al. (2024). Monkeypox virus genomic accordion strategies. Nat Commun *15*, 3059. 10.1038/s41467-024-46949-7.

79. Pope, W.H., Jacobs-Sera, D., Russell, D.A., Peebles, C.L., Al-Atrache, Z., Alcoser, T.A., Alexander, L.M., Alfano, M.B., Alford, S.T., Amy, N.E., et al. (2011). Expanding the diversity of mycobacteriophages: insights into genome architecture and evolution. PLoS One *6*, e16329. 10.1371/journal.pone.0016329.

80. Krupovic, M., Dolja, V.V., and Koonin, E.V. (2019). Origin of viruses: primordial replicators recruiting capsids from hosts. Nat Rev Microbiol *17*, 449-458. 10.1038/s41579-019-0205-6.

81. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2022). The logic of virus evolution. Cell Host Microbe *30*, 917-929. 10.1016/j.chom.2022.06.008.

82. Hwang, Y., Roux, S., Coclet, C., Krause, S.J.E., and Girguis, P.R. (2023). Viruses interact with hosts that span distantly related microbial domains in dense hydrothermal mats. Nat Microbiol *8*, 946-957. 10.1038/s41564-023-01347-5.

83. Hurwitz, B.L., and U'Ren, J.M. (2016). Viral metabolic reprogramming in marine ecosystems. Curr Opin Microbiol *31*, 161-168. 10.1016/j.mib.2016.04.002.

84. Tian, F., Wainaina, J.M., Howard-Varona, C., Dominguez-Huerta, G., Bolduc, B., Gazitua, M.C., Smith, G., Gittrich, M.R., Zablocki, O., Cronin, D.R., et al. (2024). Prokaryotic-virus-encoded auxiliary metabolic genes throughout the global oceans. Microbiome *12*, 159. 10.1186/s40168-024-01876-z.

85. Kieft, K., Zhou, Z., Anderson, R.E., Buchan, A., Campbell, B.J., Hallam, S.J., Hess, M., Sullivan, M.B., Walsh, D.A., Roux, S., and Anantharaman, K. (2021). Ecology of inorganic sulfur auxiliary metabolism in widespread bacteriophages. Nat Commun *12*, 3503. 10.1038/s41467-021-23698-5.

86. Huang, X., Zhou, Z., Liu, H., Li, Y., Ge, T., Tang, X., He, Y., Ma, B., Xu, J., Anantharaman, K., and Li, Y. (2024). Soil nutrient conditions alter viral lifestyle strategy and potential function in phosphorous and nitrogen

metabolisms. Soil Biology and Biochemistry *189*, 109279. https://doi.org/10.1016/j.soilbio.2023.109279.

87.     Wang, L., Wang, Y., Huang, X., Ma, R., Li, J., Wang, F., Jiao, N., and Zhang, R. (2022). Potential metabolic and genetic interaction among viruses, methanogen and methanotrophic archaea, and their syntrophic partners. ISME Communications *2*, 50. 10.1038/s43705-022-00135-2.

88.     Zhou, Y., Gutiérrez-Preciado, A., Liu, Y., Moreira, D., Yakimov, M.M., López-García, P., and Krupovic, M. (2025). Nested parasitism in hypersaline environments: viruses and virus satellites of haloarchaea and their nanosized cellular symbionts. bioRxiv, 2025.2002.2015.638363. 10.1101/2025.02.15.638363.

89.     Colman, D.R., Keller, L.M., Arteaga-Pozo, E., Andrade-Barahona, E., St Clair, B., Shoemaker, A., Cox, A., and Boyd, E.S. (2024). Covariation of hot spring geochemistry with microbial genomic diversity, function, and evolution. Nat Commun *15*, 7506. 10.1038/s41467-024-51841-5.

90.     Yin, Z., Ye, L., and Jing, C. (2022). Genome-Resolved Metagenomics and Metatranscriptomics Reveal that Aquificae Dominates Arsenate Reduction in Tengchong Geothermal Springs. Environ Sci Technol *56*, 16473-16482. 10.1021/acs.est.2c05764.

91.     Yuan, L., and Ju, F. (2023). Potential Auxiliary Metabolic Capabilities and Activities Reveal Biochemical Impacts of Viruses in Municipal Wastewater Treatment Plants. Environ Sci Technol *57*, 5485-5498. 10.1021/acs.est.2c07800.

92.     Turunen, M., Olsson, J., and Dallner, G. (2004). Metabolism and function of coenzyme Q. Biochim Biophys Acta *1660*, 171-199. 10.1016/j.bbamem.2003.11.012.

93.     Bekker, M., Alexeeva, S., Laan, W., Sawers, G., Teixeira de Mattos, J., and Hellingwerf, K. (2010). The ArcBA two-component system of Escherichia coli is regulated by the redox state of both the ubiquinone and the menaquinone pool. J Bacteriol *192*, 746-754. 10.1128/JB.01156-09.

94.     Sarewicz, M., and Osyczka, A. (2015). Electronic connection between the quinone and cytochrome C redox pools and its role in regulation of mitochondrial electron transport and redox signaling. Physiol Rev *95*, 219-243. 10.1152/physrev.00006.2014.

95.     Koonin, E.V., Dolja, V.V., Krupovic, M., and Kuhn, J.H. (2021). Viruses Defined by the Position of the Virosphere within the Replicator Space. Microbiol Mol Biol Rev *85*, e0019320. 10.1128/MMBR.00193-20.

96.     Alvarez, L.M., and Peng, X. (2025). Redefining paradigms in the archaeal virus-host arms race. Preprint. 10.1101/2025.04.20.649705.

97.     Lopez-Beltran, A., Botelho, J., and Iranzo, J. (2024). Dynamics of CRISPR-mediated virus-host interactions in the human gut microbiome. ISME J *18*. 10.1093/ismejo/wrae134.

98. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2021). The healthy human virome: from virus-host symbiosis to disease. Curr Opin Virol *47*, 86-94. 10.1016/j.coviro.2021.02.002.

99. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2015). Origins and evolution of viruses of eukaryotes: The ultimate modularity. Virology *479-480*, 2-25. 10.1016/j.virol.2015.02.039.

100. Skennerton, C.T., Imelfort, M., and Tyson, G.W. (2013). Crass: identification and reconstruction of CRISPR from unassembled metagenomic data. Nucleic Acids Res *41*, e105. 10.1093/nar/gkt183.

101. Bland, C., Ramsey, T.L., Sabree, F., Lowe, M., Brown, K., Kyrpides, N.C., and Hugenholtz, P. (2007). CRISPR recognition tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. BMC Bioinformatics *8*, 209. 10.1186/1471-2105-8-209.

102. Steinegger, M., and Soding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. Nat Biotechnol *35*, 1026-1028. 10.1038/nbt.3988.

103. Padilha, V.A., Alkhnbashi, O.S., Shah, S.A., de Carvalho, A., and Backofen, R. (2020). CRISPRcasIdentifier: Machine learning for accurate identification and classification of CRISPR-Cas systems. Gigascience *9*. 10.1093/gigascience/giaa062.

104. Camargo, A.P., Nayfach, S., Chen, I.A., Palaniappan, K., Ratner, A., Chu, K., Ritter, S.J., Reddy, T.B.K., Mukherjee, S., Schulz, F., et al. (2023). IMG/VR v4: an expanded database of uncultivated virus genomes within a framework of extensive functional, taxonomic, and ecological metadata. Nucleic Acids Res *51*, D733-D743. 10.1093/nar/gkac1037.

105. Ahlgren, N.A., Chen, Y., Needham, D.M., Parada, A.E., Sachdeva, R., Trinh, V., Chen, T., and Fuhrman, J.A. (2017). Genome and epigenome of a novel marine Thaumarchaeota strain suggest viral infection, phosphorothioation DNA modification and multiple restriction systems. Environ Microbiol *19*, 2434-2452. 10.1111/1462-2920.13768.

106. Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010). Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics *11*, 119. 10.1186/1471-2105-11-119.

107. Wu, Z., Liu, S., and Ni, J. (2024). Metagenomic characterization of viruses and mobile genetic elements associated with the DPANN archaeal superphylum. Nat Microbiol. 10.1038/s41564-024-01839-y.

108. Couvin, D., Bernheim, A., Toffano-Nioche, C., Touchon, M., Michalik, J., Neron, B., Rocha, E.P.C., Vergnaud, G., Gautheret, D., and Pourcel, C. (2018). CRISPRCasFinder, an update of CRISRFinder, includes a portable version, enhanced performance and integrates search for Cas proteins. Nucleic Acids Res *46*, W246-W251.

10.1093/nar/gky425.

109.    Nayfach, S., Camargo, A.P., Schulz, F., Eloe-Fadrosh, E., Roux, S., and Kyrpides, N.C. (2021). CheckV assesses the quality and completeness of metagenome-assembled viral genomes. Nat Biotechnol *39*, 578-585. 10.1038/s41587-020-00774-7.

110.    Ren, J., Song, K., Deng, C., Ahlgren, N.A., Fuhrman, J.A., Li, Y., Xie, X., Poplin, R., and Sun, F. (2020). Identifying viruses from metagenomic data using deep learning. Quant Biol *8*, 64-77. 10.1007/s40484-019-0187-4.

111.    Camargo, A.P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain, P.S.G., Nayfach, S., and Kyrpides, N.C. (2023). Identification of mobile genetic elements with geNomad. Nat Biotechnol. 10.1038/s41587-023-01953-y.

112.    Guo, J., Bolduc, B., Zayed, A.A., Varsani, A., Dominguez-Huerta, G., Delmont, T.O., Pratama, A.A., Gazitua, M.C., Vik, D., Sullivan, M.B., and Roux, S. (2021). VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses. Microbiome *9*, 37. 10.1186/s40168-020-00990-y.

113.    Hegarty, B., Riddell, V.J., Bastien, E., Langenfeld, K., Lindback, M., Saini, J.S., Wing, A., Zhang, J., and Duhaime, M. (2024). Benchmarking informatics approaches for virus discovery: caution is needed when combining in silico identification methods. mSystems *9*, e0110523. 10.1128/msystems.01105-23.

114.    Li, R., Wang, Y., Hu, H., Tan, Y., and Ma, Y. (2022). Metagenomic analysis reveals unexplored diversity of archaeal virome in the human gut. Nat Commun *13*, 7978. 10.1038/s41467-022-35735-y.

115.    Chibani, C.M., Shah, S.A., Schmitz, R.A., and Nayfach, S. (2024). Inaccurate viral prediction leads to overestimated diversity of the archaeal virome in the human gut. Nat Commun *15*, 5976. 10.1038/s41467-024-49902-w.

116.    Wang, Y., Li, R., and Ma, Y. (2024). Reply to: Inaccurate viral prediction leads to overestimated diversity of the archaeal virome in the human gut. Nat Commun *15*, 5977. 10.1038/s41467-024-49903-9.

117.    Paez-Espino, D., Eloe-Fadrosh, E.A., Pavlopoulos, G.A., Thomas, A.D., Huntemann, M., Mikhailova, N., Rubin, E., Ivanova, N.N., and Kyrpides, N.C. (2016). Uncovering Earth's virome. Nature *536*, 425-430. 10.1038/nature19094.

118.    Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol *37*, 1530-1534. 10.1093/molbev/msaa015.

119.    Parks, D.H., Rinke, C., Chuvochina, M., Chaumeil, P.A., Woodcroft, B.J., Evans, P.N., Hugenholtz, P., and Tyson, G.W. (2017). Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree

of life. Nat Microbiol *2*, 1533-1542. 10.1038/s41564-017-0012-7.

120.    Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res *13*, 2498-2504. 10.1101/gr.1239303.

121.    Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. Nucleic Acids Res *52*, W78-W82. 10.1093/nar/gkae268.

122.    Zheng, K., Sun, J., Liang, Y., Kong, L., Paez-Espino, D., McMinn, A., and Wang, M. (2025). VITAP: a high precision tool for DNA and RNA viral classification based on meta-omic data. Nat Commun *16*, 2226. 10.1038/s41467-025-57500-7.

123.    Pruitt, K.D., Tatusova, T., Brown, G.R., and Maglott, D.R. (2012). NCBI Reference Sequences (RefSeq): current status, new features and genome annotation policy. Nucleic Acids Res *40*, D130-135. 10.1093/nar/gkr1079.

124.    von Meijenfeldt, F.A.B., Arkhipova, K., Cambuy, D.D., Coutinho, F.H., and Dutilh, B.E. (2019). Robust taxonomic classification of uncharted microbial sequences and bins with CAT and BAT. Genome Biol *20*, 217. 10.1186/s13059-019-1817-x.

125.    Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J.H., Roux, S., Adriaenssens, E.M., Brister, J.R., Kropinski, A.M., Krupovic, M., Lavigne, R., et al. (2019). Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. Nat Biotechnol *37*, 632-639. 10.1038/s41587-019-0100-8.

126.    Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., and Ogata, H. (2020). KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics *36*, 2251-2252. 10.1093/bioinformatics/btz859.

127.    Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer, E.L.L., Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., et al. (2021). Pfam: The protein families database in 2021. Nucleic Acids Res *49*, D412-D419. 10.1093/nar/gkaa913.

128.    Galperin, M.Y., Wolf, Y.I., Makarova, K.S., Vera Alvarez, R., Landsman, D., and Koonin, E.V. (2021). COG database update: focus on microbial diversity, model organisms, and widespread pathogens. Nucleic Acids Res *49*, D274-D281. 10.1093/nar/gkaa1018.

129.    Shen, W., Sipos, B., and Zhao, L. (2024). SeqKit2: A Swiss army knife for sequence and alignment processing. Imeta *3*, e191. 10.1002/imt2.191.

130.    Edgar, R.C. (2022). Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence

homology and phylogeny. Nat Commun *13*, 6968. 10.1038/s41467-022-34630-w.

131. Capella-Gutierrez, S., Silla-Martinez, J.M., and Gabaldon, T. (2009). trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics *25*, 1972-1973. 10.1093/bioinformatics/btp348.

132. Kim, W., Mirdita, M., Levy Karin, E., Gilchrist, C.L.M., Schweke, H., Soding, J., Levy, E.D., and Steinegger, M. (2025). Rapid and sensitive protein complex alignment with Foldseek-Multimer. Nat Methods. 10.1038/s41592-025-02593-7.

133. Grazziotin, A.L., Koonin, E.V., and Kristensen, D.M. (2017). Prokaryotic Virus Orthologous Groups (pVOGs): a resource for comparative genomics and protein family annotation. Nucleic Acids Res *45*, D491-D498. 10.1093/nar/gkw975.

134. Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S., Larget, B., Liu, L., Suchard, M.A., and Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol *61*, 539-542. 10.1093/sysbio/sys029.

135. Robinson, O., Dylus, D., and Dessimoz, C. (2016). Phylo.io: Interactive Viewing and Comparison of Large Phylogenetic Trees on the Web. Mol Biol Evol *33*, 2163-2166. 10.1093/molbev/msw080.

136. Mistry, J., Finn, R.D., Eddy, S.R., Bateman, A., and Punta, M. (2013). Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res *41*, e121. 10.1093/nar/gkt263.

137. Mirdita, M., Schutze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022). ColabFold: making protein folding accessible to all. Nat Methods *19*, 679-682. 10.1038/s41592-022-01488-1.

**Acknowledgements**

We thank Jilu Han for his help with the installation and debugging of the software used in this study. We thank the support of the high-performance servers of the Center for High Performance Computing and System Simulation, Laoshan Laboratory (Qingdao), the High-Performance Biological Supercomputing Center at the Ocean University of China, the Marine Big Data Center of the Institute for Advanced Ocean Study of the Ocean University of China and the IEMB-1, a high-performance computing cluster operated by the Institute of Evolution and Marine Biodiversity. This study was supported by the National Natural Science Foundation of China (No. 42120104006, 42176111 and 42306111), the Ocean Negative Carbon Emissions (ONCE), 2024 Graduate Self-directed Research Project (2024ZZKY, 202461036) and the Fundamental Research Funds for the Central Universities (202172002, 201812002, 202461036 andrew McMinn).

**Author contributions**

Conceptualization: K.Z. and Y.L.; Analysis: K.Z.; Database: D.P.-E.; Writing manuscript: K.Z.; Review and editing: Y.L., A.M. and M.W.; Computational resource: D.P.-E., M.L-P and M.W.; Funding acquisition and supervision: Y.L., A.M. and M.W.

**Competing interests**

The authors declare no competing interests for the manuscript "Expanding diversity of virome associated with ammonium-oxidizing archaea and related lineages".

# Figure legends

Figure 1. The expanding diversity of *Nitrososphaeria*-associated viruses or mobile genetic elements (MGEs). a, The workflow for mining, filtering, and identifying viruses or MGEs primarily included the prediction of archaeal CRISPR systems, viruses or MGEs mining, genome filtering, and the definition of viral or MGE operational taxonomic units (vOTUs or MGE OTUs). Only genomes encoding detectable capsid or related proteins were considered viruses; others were classified as MGEs. b, Biogeographic distribution of *Nitrososphaeria*-associated viral or MGE genomes. In the map, nodes of different colors indicate their putative archaeal hosts: *Ca. Caldarchaeales*, ammonia-oxidizing archaea (AOA), deeply-branched AOA-related archaea (DB-AO-related), and deeply-branched *Nitrososphaeria* archaea (D-B *Nitrososphaeria*). c, The number of vOTUs or MGE OTUs with distinct putative morphologies were originated from different habitats and were associated with diverse *Nitrososphaeria* lineages. Bar colours represent different archaeal lineages. d, Phylogenetic tree of *Nitrososphaeria* genomes. This tree was constructed based on concatenated alignments of 46 ribosomal proteins (totalling 5,963 amino acid positions) using the maximum likelihood method with the LG+C60+F model as recommended by Spang et al. (see Methods). Reference genomes from *Thermoprotei* were used as the outgroup to root the tree. Tree annotation includes the habitats and lineages of these archaeal genomes. Virion icons to the right of the tree indicate the numbers of viruses or MGEs with different possible morphologies (at the OTU level) associated with each archaeal lineage.

Figure 2. The similarity of *Nitrososphaeria*-associated viruses or mobile genetic elements (MGEs) based on their genome contents. a, Genome-content similarity network of *Nitrososphaeria*-associated viruses or MGEs within the context of the global virosphere (sequences from NCBI RefSeq v. 223). For clarity, only connected component involving *Nitrososphaeria*-associated viruses or MGEs are shown. Each node represents a viral or MGE contig and is coloured according to its host lineage. *Nitrososphaeria*-associated viruses or MGEs clustering with established viral families are annotated adjacent to their respective groups. The pie charts illustrate the number of *Nitrososphaeria*-specific viral clusters (VCs) generated from this network. b, Similarity network between *Nitrososphaeria*-associated virome. Nodes represent viral or MGE genomes and are coloured according to their associated archaeal lineages. Each distinct super viral cluster (sVC) as a connected component may consist of one or more

VCs. The percentage of shared proteins between genomes is indicated by the thickness of the edges connecting the nodes. The shapes of the nodes denote the potential morphologies of these viruses or MGEs.

Figure 3. Assignments of new families within *Nitrososphaeria*-associated viruses or mobile genetic elements (MGEs). a, Virus proteomic hierarchy clustering tree used for the delineation of viral families. A distance threshold of 0.05 and a minimum of six shared proteins among members were used as the criteria for defining families (see Methods). The names of the proposed families are indicated on the left side of the tree, with different virion icons representing the possible morphologies of these families. Tree annotation denotes host lineage and virion morphology. b, Genome organisation diagrams representing each new viral family. Boxes denote individual open reading frames (ORFs), with the colour gradient indicating the sharing rate of each ORF among members within the family. ORFs with a sharing rate greater than 0.9 are considered to encode viral hallmark proteins (VHPs), which are indicated by asterisks below the corresponding ORFs. The numbers inside the genome diagrams denote the number of members in each family and the average rate of shared protein clusters among the members. Abbreviations: MT/GW, Marine Trench and Ground Water; MBG-A, Marine Beneath Group A; D-B-AOA-related, deeply-branched AOA-related archaea; D-B *Nitrososphaeria*, deeply-branched *Nitrososphaeria* archaea.

Figure 4. Phylogenetic inferences of *Nitrososphaeria*-associated viruses or mobile genetic elements (MGEs) based on viral hallmark proteins (VHPs) through Bayesian inference (BI) and Maximum Likelihood (ML) approach. a, The trees display the topologies inferred by BI based on VHPs. *Nitrososphaeria*-associated viruses or MGEs are highlighted with bold red branches. To the right of each tree are shown: 1) BI convergence indices, including minimum effective sample size (minESS), potential scale reduction factor (PSRF), and average standard deviation of split frequencies (ASDSF); 2) indices for comparing BI and ML topologies, including normalized Robinson–Foulds (RF) distance based on ML and the number of clades shared between BI and ML. Posterior probabilities from BI are indicated on the tree branches with colored nodes, alongside ML bootstrap values (MLB). Names of viral families (in bold red font) and other host groups are annotated beside the corresponding branches. Tree annotations indicate the *Nitrososphaeria* lineages associated with these viruses/MGEs. a, BI tree based on the SIRV2-like capsid protein (domain accession in Pfam: PF12193). DUF732 (domain accession in Pfam: PF05305 and in VOG223: VOG14611) was used as the outgroup to root the tree. Leaves corresponding to rod-shaped viruses with heterodimeric capsids are highlighted with red squares. b, BI tree based on the HK97-like

capsid protein (domain accession in Pfam: PF05065), rooted using L_lactis_ph-MCP (domain accession in Pfam: PF06673) as the outgroup. c, BI tree based on the HK97-like capsid protein (domain accession in VOGdb: VOG01305), rooted using C1_1 (domain accession in Pfam: PF00130) as the outgroup. d, BI tree based on the HK97-like capsid protein (domain accession in Pfam: PF07068), rooted using L_lactis_ph-MCP (domain accession in Pfam: PF06673) as the outgroup. e, BI tree based on the terminase large subunit protein (domain accession in KEGG: K06909), rooted using P-loop NTPase (domain accession in SUPERFAMILY: SSF52540) as the outgroup. f, BI tree based on the terminase large subunit protein (domain accession in Pfam: PF03237), rooted using P-loop NTPase (domain accession in SUPERFAMILY: SSF52540) as the outgroup. Abbreviations: MT/GW, Marine Trench and Ground Water; MBG-A, Marine Beneath Group A; D-B-AOA-related, deeply-branched AOA-related archaea; D-B *Nitrososphaeria*, deeply-branched *Nitrososphaeria* archaea.

Figure 5. Capsid structure features of *Nitrososphaeria*-associated rod-shaped and head-tailed viruses. a, Three-dimensional structural prediction of archaeal rod-shaped viral capsids and their structure-based similarity network. Nodes represent viral capsids and are color-coded according to their viral families. Edge width indicates the degree of structural similarity between capsids (only identities ≥ 0.5 are shown). Representative structures from each cluster are displayed adjacent to their respective groups. a, predicted structures of viral capsid protein monomers and their clustering network; b, predicted structures of viral capsid protein dimers and their clustering network. Possible heterodimers are indicated with black arrows. Genome IDs and corresponding per-residue measures of local confidence (pLDDT) are indicated below each structure. c, Three-dimensional structural prediction of archaeal HK97-like viral capsids and their structure-based similarity network.

Figure 6. *Nitrososphaeria*-associated rod-shaped viruses encode unique auxiliary genes and exhibit a distinctive landscape of horizontal gene transfer (HGT) with their hosts. a, Maximum-likelihood (ML) phylogenetic tree of the *Demeterviridae*-encoded ArsR-like transcriptional regulator. Branches associated with archaeal viruses and bacteriophages are highlighted in purple and black, respectively, and are shown in bold. The names of archaea-associated viral families are annotated alongside the corresponding branches on the tree. The tree annotation includes proteins of cellular origin (from InterPro), derived from archaeal or bacterial genomes. For clarity, only clades or branches related to viruses were retained, while the others were pruned. Representative structures and their per-residue measures of local confidence (pLDDT) are displayed below the phylogenetic tree. b, Maximum-likelihood (ML)

phylogenetic tree of the *Demeterviridae*-encoded UbiG-like protein. c, Proportion of HGT events with different transfer directions among various *Nitrososphaeria* lineages. Different colours of the sectors indicate the direction of HGT, and the numbers within them represent the number of observations. c, c, The number of HGTs with distinct directions were associated from different functional catagories and were linked with diverse *Nitrososphaeria* lineages. Bar colours represent the direction of HGT and *Nitrososphaeria* lineages, respectively. The numbers of observations were indicated beside bars. Abbreviations: HTV27, Halorubrum tailed virus 27; SSV5, Sulfolobus spindle-shaped virus 5; phiCDIF1296T, Peptoclostridium phage phiCDIF1296T; MT/GW, Marine Trench and Ground Water; MBG-A, Marine Beneath Group A; D-B-AOA-related, deeply-branched AOA-related archaea; D-B *Nitrososphaeria*, deeply-branched *Nitrososphaeria* archaea.

**a**

**1. CRISPR prediction**

2,137 genomes and *Nitrososphaeria* MAGs from GenBank and IMG/M
170 metagenomic samples

MinCED, CRASS, CRISPR-Cas-Finder, CRISPR-Cas-Identifier

MMSeqs2

55,995 DRs from *Nitrososphaeria*-associated contigs
234,036 spacers of *Nitrososphaeria*

geNomad, VirSorter2, DeepVirFinder

**2.1. Proviral prediction**
IMG/VR (v.4)
**2.2. Viral/MGE prediction**
**2.3. Viral identification**

MMSeqs2

geNomad, MMSeqs2

**3. Genome filtering**

499 vOTUs without predicted capsid-related proteins (possible viruses and other MGEs)

MMSeqs2

781 of *Nitrososphaeria*-associated viruses/MGEs

MMSeqs2
MMSeqs2

**4. vOTU definition**
659 vOTUs

160 vOTUs with predicted capsid-related proteins (viruses, expanded by sVCs)

**b**

- *Ca. Caldarchaeales*  ● AOA  ● D-B AOA-related  ● D-B *Nitrososphaeria*

**c**

D-B *Nitrososphaeria* (7)
pSL4-affiliation (3)
Geothermal spring (202)
Head-tailed (526)
*Ca. Caldarchaeales* (231)
Hydrothermal vents (75)
D-B AOA-related (50)
MBG-A (15)
*Nitrosocaldales* (12)
Seawater (192)
MT/GW (61)
Rod-shaped (51)
Group 1.1b (57)
Freshwater (57)
*Nitrosotalea* (10)
Spindle-shaped (16)
Mine wastwater (2)
Icosahedral (15)
Group 1.1a (162)
Soil (80)

**d**

*Thermoprotei*
D-B *Nitrososphaeria*
pSL4-affiliation

*Ca. Caldarchaeales*
Non-AO *Nitrososphaeria*

D-B AOA-related
MBG-A
*Nitrosocaldales*
MT/GW
Group 1.1b
*Nitrosotalea*
Group 1.1a

AOA

Tree scale: 0.2

Morphology
⬡ Head-tailed
▭ Rod-shaped
⬡ Spindle-shaped
⬡ Icosahedral

Boostraps
● ≥ 90
● ≥ 70

Habitats
■ Hydrothermal vents
■ Marine
■ Geothermal springs
■ Soil
■ Freshwater

**a**

VC_9/62: *Marthavirus*

VC_106: *Itzamnaviridae* (*Methanobacteriota*)

VC_15: *Ekchuahviridae* (*Methanobacteriota*)

VC_109: *Ahmunviridae* (*Methanobacteriota*)

*Nitrososphaeria*-specific VCs

Non-specific VCs

**Host lineages**

AOA

ANME-1

*Ca. Caldarchaeales*

*Pseudomonadota*

Other *Nitrososphaeria*

*Cyanobacteriota*

Other Hosts

**b**

sVC_0003  sVC_0012  sVC_0036  sVC_0002  sVC_0015  sVC_0039  sVC_0019  sVC_0024  sVC_0044  sVC_0004  sVC_0035

sVC_0070

sVC_0052  sVC_0047  sVC_0041  sVC_0040  sVC_0038  sVC_0037  sVC_0034

sVC_0022  sVC_0058

sVC_0032  sVC_0001  sVC_0028  sVC_0025  sVC_0045  sVC_0031  sVC_0023  sVC_0020

sVC_0008  sVC_0030  sVC_0043  sVC_0027  sVC_0029  sVC_0013  sVC_0014  sVC_0010  sVC_0068  sVC_0054  sVC_0051  sVC_0049

sVC_0061

sVC_0046  sVC_0042  sVC_0033  sVC_0007  sVC_0057

sVC_0032  sVC_0026  sVC_0021  sVC_0005  sVC_0069  sVC_0067  sVC_0061  sVC_0053  sVC_0018  sVC_0066  sVC_0060  sVC_0059

**Host lineage**

AOA

Group 1.1a

*Nitrosotales*

Group 1.1b

MT/GW

*Nitrosocaldales*

Non-AO *Nitros.*

MBG-A

D-B AOA-related

*Ca. Caldarchaeales*

pSL4-affiliation

D-B *Nitrososphaeria*

**Morphology**

Head-tail

Filamentous

Icosahedral

Spindle

**Shared protein clusters (%)**

97          40

80          20

**a**

Host lineage

*Hypnosviridae* (sV_0043)
*Ekchuahviridae* (sV_0044)
*Yangangviridae* (sV_0069)
***Thanatosviridae*** (sV_0005)
***Aetherviridae*** (sV_0013)
***Alboranviridae*** (sV_0070)
***Baleariviridae*** (sV_0070)
***Tyrrheniaviridae*** (sV_0070)
***Apateviridae*** (sV_0003)
*Kyanoviridae* (sV_0003)
*Chaacviridae* (sV_0017)
*Thaspiviridae* (sV_0068)
*Itzamnaviridae* (sV_0042)
***Erebusviridae*** (sV_0002)
***Morosviridae*** (sV_0002)
***Demeterviridae*** (sV_0012)

Host lineage

- Group 1.1a
- Group 1.1b
- MT/GW
- *Nitrosocaldales*
- MBG-A
- D-B AOA-related
- *Ca. Caldarchaeales*
- *Nitrosotales*
- pSL4-affiliation
- D-B *Nitrososphaeria*

**b**

***Demeterviridae***
3300042910_000011
(14/0.29)

Trimeric autotrans-porter adhesin
Sulfolobus virus coat protein

***Alboranviridae***
Marthavirus-1-C1
(9/0.24)

***Hypnosviridae***
3300028891_000102
(4/0.24)

***Thanatosviridae***
3300005298_000040
(2/0.58)

DNA replication related proteins
Tail-related proteins

***Baleariviridae***
Marthavirus-3
(18/0.31)

DNA replication related proteins

***Aetherviridae***
3300006752_000072
(2/0.74)

Head decoration protein
Terminase
Tail-related proteins
Major capsid protein

***Morosviridae***
Ga0197142_1000541
(8/0.36)

Portal protein
Capsid related protein
Tail-related protein

***Erebusviridae***
Ga0209493_1000412
(5/0.29)

Tail tube protein

***Apateviridae***
3300025082_001358
(4/0.26)

DNA replication related proteins
Transcriptional regulator protein
DNA polymerase
Recombination protein

***Tyrrheniaviridae***
Marthavirus-22
(2/0.28)

DNA helicase
m6A methyl-transferase
Minor head protein

Sharing rate: ≥ 0.75    0.5    0.25    0

**a**

Family
- *Rudiviridae*
- *Ahmunviridae*
- *\*Byatisviridae*
- Other Nitrososphaeria-associated viruses

(1) pLDDT = 92.5
(2) pLDDT = 90.3
(3) pLDDT = 90.1
(4) pLDDT = 94.2
(5) pLDDT = 90.6

Identity
- ≥ 50
- ≥ 65
- ≥ 75
- ≥ 85
- ≥ 95

(1) IMGVR_UViG_3300025089_000008; (2) Sulfolobus islandicus rod-shaped virus 2; (3) Acidianus rod-shaped virus 2;
(4) IMGVR_UViG_3300014911_000547; (5) IMG_3300025089|Ga0209394_1003436

**b**

Family
- *Rudiviridae*
- *Ahmunviridae*
- *\*Demeterviridae*
- Other Nitrososphaeria-associated viruses

(1) pLDDT = 91.5
(3) pLDDT = 73.2
(4) pLDDT = 62.9
(5) pLDDT = 83.6
(6) pLDDT = 75.5

Identity
- ≥ 50
- ≥ 65
- ≥ 75
- ≥ 85
- ≥ 95

(1) IMG_3300025089|Ga0209394_1001498; (3) Sulfolobus islandicus rod-shaped virus 2 ; (4) Saccharolobus solfataricus
rod-shaped virus 1; (5) Methanophagales virus PBV300; (6) IMG_3300025089|Ga0209394_1001595

**c**

GCA_963578755.1|
CAUZDV010000001.1
pLDDT = 78.6

Escherichia phage T4
pLDDT = 81.9

Acinetobacter phage Acj9
pLDDT = 82.3

Aeromonas phage AhSzw-1
pLDDT = 57.8

Serratia phage phiMAM1
pLDDT = 78.9

IMGVR_UViG_3300017593_000761
pLDDT = 82.5

IMGVR_UViG_3300021495_000002
pLDDT = 83.1

Marthavirus-1-C1
pLDDT = 82.4

Sinorhizobium phage phiM9
pLDDT = 81.3

Aeromonas phage Ah1
pLDDT = 81.3

IMGVR_UViG_3300028675_000006
pLDDT = 72.2

IMGVR_UViG_3300033490_000008
pLDDT = 73.2

IMG_3300009598|
Ga0105154_1000501
pLDDT = 73.2

IMGVR_UViG_3300011323_000033
pLDDT = 76.3

Halorubrum virus HRTV-29
pLDDT = 77.8

Vibrio phage nt-1
pLDDT = 81.3

Aeromonas phage CC2
pLDDT = 83.0

Family
- *Kyanoviridae* (AOA)
- *\*Morosviridae*
- *\*Alboranviridae*
- *\*Baleariviridae*
- *Ekchuahviridae*
- *Kyanoviridae* (Cyanobacteria)
- *Straboviridae*
- *Ackermannviridae*
- *Pootjesviridae*
- *Haloferuviridae*
- Other *Nitrososphaeria*-associated viruses
- Other viral RefSeqs

Identity
- ≥ 50
- ≥ 65
- ≥ 75
- ≥ 85
- ≥ 95