A global viromic survey reveals unprecedented viral diversity within ammonium-oxidizing archaea and related lineages

Kaiyang Zheng[1], Yantao Liang[1,2,3*], Fabai Wu[4*], Hao Yu[1], David Paez-Espino[5], Mario López-Pérez[6], Sungeun Lee[7], Graeme W. Nicol[7], Christina Hazard[7], Andrew McMinn[1,3,8*], Min Wang[1,2,3*]

[1] College of Marine Life Sciences, MoE Key Laboratory of Evolution & Marine Biodiversity, Institute of Evolution and Marine Biodiversity, Frontiers Science Centre for Deep Ocean Multispheres and Earth System, Centre for Ocean Carbon Neutrality, Ocean University of China, Qingdao, China.

[2] Haide College, Ocean University of China, Qingdao, China.

[3] UMT-OUC Joint Centre for Marine Studies, Qingdao, China.

[4] Faculty of Science, Eastern Institute of Technology, Ningbo, China

[5] Ancilia Biosciences, Co. Ltd., Alexandria Launchlabs, New York, USA.

[6] Evolutionary Genomics Group, División de Microbiología, Universidad Miguel Hernández, San Juan, Spain

[7] Laboratoire d'Ecologie Microbienne, Universite Claude Bernard Lyon 1, Villeurbanne, France

[8] Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia.

* Correspondence:

Yantao Liang, Email: **liangyantao@ouc.edu.cn**

Fabai Wu, Email: **fabaiwu@eitech.edu.cn**

Andrew McMinn, Email: **andrew.mcminn@utas.edu.au**

Min Wang, Email: **mingwang@ouc.edu.cn**

Abstract

Nitrososphaeria, an archaeal class containing all ammonium-oxidizing archaea (AOA) and their closely related clades (e.g. *Ca. Caldarchaeales*), is a widely distributed archaeal lineage across diverse ecosystems. Due to the difficulty of isolation and cultivation, the viral diversity and evolution associated with non-marine AOA lineages and *Ca. Caldarchaeales* remain largely uncharacterized. Here, based on globally sampled genomes and metagenomes, we assembled a genomic catalogue associated with AOA, including 283 species-level viruses and 415 species-level mobile genetic elements (MGEs). Notably, 259 previously unrecognized species-level viruses or MGEs associated with *Ca. Caldarchaeales* were identified. Across the two major *Nitrososphaeria* lineages, we uncovered an expansion of archaeal viral families: 11 families associated with AOA, including seven new head-tailed lineages, one of which comprises a mega archaeal virus with a genome size exceeding 220 kb; ten families associated with *Ca. Caldarchaeales*, including four novel head-tailed groups and two previously unknown rod-shaped family (*Demeterviridae* and *Gaiaviridae*). Phylogenetic inference showed that rod-shaped viruses associated with *Ca. Caldarchaeales* and methanogens are phylogenetically intertwined and likely share a common ancestor with *Hoswirudivirus*, a *Sulfolobales*-associated rod-shaped virus characterized by two copies of the coat gene. Through structural modelling, a rare fusion of the coat protein in *Demeterviridae* was observed, in which a single open reading frame within a gene island adjacent to a glycosyl transferase appears capable of forming a complete capsomer. This provides new insights into the origin and evolution of *Adnaviria* viral realm. Functional genomics analyses further revealed that these novel rod-shaped viruses encode virus-specific rare quinone biosynthesis genes and arsenic response transcriptional regulator genes, providing potential clues to their mechanisms of adaptation to hyperthermal environments. Collectively, our work uncovers the hidden genomic, evolution and functional diversity of *Nitrososphaeria* viromes for the first time and closes a long-standing gap in archaeal virome research.

Highlights

1. First genomic catalogue of viruses and MGEs linked to *Nitrososphaeria*.

2. Discovery of diverse novel archaeal viral families.

3. New insights into rod-shaped virus evolution and archaeal virus–host adaptation.

Introduction

The *Nitrososphaeria* is an archaeal class within *Thermoprotei*, encompassing the previously defined *Thaumarchaeota* and *Augarchaeota*, both of which are important members of the *Thermoproteati* (previously known as TACK superphylum)[1,2]. In many aquatic ecosystems, *Nitrososphaeria* constitute up to 40% of microorganisms, contributing significantly to ammonia oxidation, thus heavily influencing nitrogen cycling [3-5]. *Nitrososphaeria* includes all ammonia-oxidizing archaea (AOA) [6] and utilizes ammonia oxidation pathways as their primary energy source [7,8]. Compared to ammonia-oxidizing bacteria, AOA thrive in low-ammonia conditions due to higher ammonium affinity, dominating nitrogen-limited environments as efficient "nitrogen thieves" [9-14]. The ancestors of AOA are considered to have originated in terrestrial environments and to have adapted aerobically after the Great Oxygenation Event [13,15,16]. Subsequently, these archaea became key microbial players across diverse ecosystems and make a substantial contribution to carbon fixation and ammonia oxidation [8,15,17-21]. As the sister clade of AOA, *Ca. Caldarchaeales* shares substantial genetic content involved in sulphur oxidation and ammonia oxidation, suggesting an ancestral ecological transition of AOA from hyperthermal to moderate environments [22]. Primarily inhabiting hydrothermal vents and hot springs, *Ca. Caldarchaeales* influences carbon and sulphur cycling, nutrient transformation and microbial community structures, playing critical ecological roles in extreme environments [23,24].

Viruses and mobile genetic elements (MGEs) exhibit complex morphological and genomic diversity, significantly shaping archaeal genomes and their ecological footprint [25,26]. Due to cultivation challenges, environmental metagenomics has become the primary method for exploring archaeal viral diversity. Recent studies using environmental metagenomes have greatly expanded viral diversity associated with the Asgard super-phylum, methanogens and DPANN super-phylum, addressing gaps in viral evolution and host-virus interactions [27-31]. Over 100 AOA-associated viral or MGE sequences from marine or soil samples have so far been identified from metagenomic survey, revealing diverse viral elements that influence ammonia oxidation and host genomic organization [32-40]. However, these case studies were all based on samples from single geographic regions rather than systematic global surveys, which prevents a comprehensive profiling of the diversity, evolution,

and functions of these archaeal viruses. The absence of viruses associated with extremophilic AOA

lineages (e.g., *Nitrosocaldales* and *Nitrosotalea*) poses a challenge for resolving phylogenetic

placements of AOA viruses within the virosphere. In particular, as a phylogenetic sister clade closely

related to AOA and one of the main lineages constituting the TACK super-phylum,

Caldarchaeales-associated virome remain underexplored and represent a major enigma within

virology.

To address this knowledge gap, we systematically explored *Nitrososphaeria*-associated viruses

or MGEs diversity based on extensive global genomic and metagenomic datasets. Unlike previous

studies that focused on AOA-associated viruses from only single geographic regions[32-36,39,40], our

comprehensive analysis of the *Nitrososphaeria*-associated virome fills a critical gap in the

understanding of its genomics, phylogenetics, and functional repertoire. Through the establishment

of a genomic catalogue comprising hundreds of viral and MGE sequences, we identified nine

previously unrecognized archaeal viral families, including two group of rod-shaped viruses

associated with *Ca. Caldarchaeales* that exhibit unique capsomer features and metabolic potential.

These culture-independent virus–archaea associations remain to be empirically verified as infections,

yet they offer a testable data foundation for subsequent establishment of *Nitrososphaeria*–virus

culture systems, thereby providing new insights into the evolution of archaeal virosphere.

Results

Overview of the genomic catalogue of *Nitrososphaeria*-associated viruses or MGEs

Nitrososphaeria genome database containing 2,137 genomes and metagenome-assembled genomes (MAGs) was constructed from NCBI GenBank and IMG/M databases (Supplementary Fig. 1 and Supplementary Note 1)[41,42]. The virus or MGE database comprised over 15 million contigs from IMG/VR v.4, together with 170 metagenome-assembled contigs originating from hyperthermal, abyssal/hadal ocean, hypersaline, frozen, and acidic environments, including 17 from our published and unpublished studies (Supplementary Dataset 1) [41,43-46]. The integration of viral and MGE sequences from our pipeline and previous studies enabled the establishment of a genomic catalogue comprising 820 *Nitrososphaeria*-associated viruses or MGEs (Supplementary Fig. 1 and Dataset 2). These were grouped into 698 species-level operative taxonomic units (vOTUs), comprising 286 vOTUs with and 407 MGE OTUs without known viral capsid proteins. In addition, 13 vOTU and 27 MGE OTUs associated with AOA and 103 vOTUs and 151 MGE OTUs associated with *Ca. Caldarchaeales* were newly identified, respectively.

Nitrososphaeria-associated viruses or MGEs are widely distributed from tropical to polar regions and across terrestrial to marine environments (Fig. 1a). Most of these regions have not been covered by previous studies focusing on AOA-associated viruses[32-36,39,40]. These viruses or MGEs predominately originated from geothermal, marine, soil and hydrothermal environments, aligning with habitats of AOA, deeply-branched AOA-related lineage and *Ca. Caldarchaeales* (Fig. 1b) [7,20,22,23,47-50]. This genome catalogue revealed novel viruses associated with nine previously unrecognized archaeal lineages within *Nitrososphaeria* (Fig. 1c), including *Ca. Caldarchaeales*, *Nitrosocaldales*, *Nitrosotalea*, and other deeply branched AOA-related lineages (also known as non-AO *Thaumarchaeota*), thereby expanding the number of lineages known to host these viruses from two to eleven [32-38,51].

The expansion of *Nitrososphaeria* virome diversity revealed numerous previously unrecognized archaeal viral families

Nitrososphaeria-associated viruses or MGEs exhibit distinct differences in genome content compared with other viruses. The genome similarity network (sharing at least 20% of protein clusters) yielded 67 *Nitrososphaeria*-associated viral communities (super viral clusters, sVCs), with 90% of these communities containing only *Nitrososphaeria*-associated viruses or MGEs (Fig. 2). AOA-associated sVCs and *Caldarchaeales*-associated sVCs show no overlap, indicating clear differences between the viromes of these two main *Nitrososphaeria* lineages. Notably, both exhibit overlaps with the sVCs associated with non-AO *Nitrososphaeria*, which represents a transitional group (n = 6), suggesting the possible presence of transitional lineages within the *Nitrososphaeria*-associated viromes. In addition, some *Nitrososphaeria*-associated viruses exhibited close genome-content relationships with other archaeal or bacterial viruses (e.g. ANME-1 and *Bacteroidota*), suggesting that these *Nitrososphaeria*-associated viruses or MGEs may have undergone host switching or engaged in gene exchange with viruses of other prokaryotes. (Figs. 2 and Supplementary Dataset 3) [29,30].

Proteome-based hierarchical clustering tree was employed to further characterize novel families based on current viral taxonomic framework [52]. Twenty-two viral families (or family-level lineage) were found to be associated with *Nitrososphaeria*, including 14 previously undescribed ones. Except for the 22 vOTUs that were assigned to established viral families (Figs. 2, 3 and Supplementary Dataset 2), 108 vOTUs were proposed to be newly named viral families here, they include: two rod-shaped viral families (*Demeterviridae* and *Gaiaviridae*) and four head-tailed viral families (*Erebusviridae*, *Hypnosviridae*, *Morosviridae* and *Nyxviridae*), which were associated with *Ca. Caldarchaeales* and seven head-tailed family (*Apateviridae*, *Alboranviridae*, *Baleariviridae*, *Charonviridae*, *Thanatosviridae*, *Freyrviridae* and *Peleviridae*) associated with AOA (Fig. 3). In addition, nine MGE families without known capsid proteins were generated, indicating a more complex genomic diversity of the *Nitrososphaeria*-associated virome (Fig. 3 and Supplementary Dataset 2).

The genomes of *Nitrososphaeria*-associated viral families vary in length, and the proteins they encode exhibit diverse functional potentials (Fig. 4). *Apateviridae* contains a member with a genome

length of 228 kb (vOTU_0015), which encodes 34 viral core functional genes (Supplementary Dataset 4), offering clues to the existence of previously unrecognized archaeal "jumbo" viruses [53]. The vOTU_0015 encodes a variety of metabolic genes, notably carrying 16 Radical SAM–associated proteins, suggesting that these functions may play important roles in the survival of this virus. *Jordviridae* represent a family of temperate archaeal viruses that encode integrase and a series of Ig-like domain–containing protein, whose functions may be similar with host-recognition proteins found in other archaeal viruses [29]. *Nyxviridae* encode TFII-like proteins (transcription factor II), a class of transcription factors typical of archaea and eukaryotes, suggesting that these viruses may hijack the archaeal transcription machinery [28,54]. *Erebusviridae* encode MCM-like proteins (minichromosome maintenance proteins), a core DNA replication factor widely conserved in archaea and eukaryotes, suggesting that these viruses may hijack the archaeal DNA replication machinery [55,56]. *Demeterviridae* represent a rod-shaped viral lineage encoding multiple SIRV2-like MCPs and harboring UbiG-like proteins (2-polyprenyl-3-methyl-5-hydroxy-6-methoxy-1,4-benzoquinol methyltransferase) and ArsR-like proteins, highlighting the unique quinone metabolic potential and transcriptional regulatory mechanisms of this group of archaeal rod-shaped viruses [57].

The distinct evolutionary placements of *Nitrososphaeria*-associated viruses within the *Adnaviria* and *Duplodnaviria*

Bayesian (BI) and Maximum-Likelihood (ML) inference with structural modelling was used to characterize the evolution of *Nitrososphaeria*-associated viruses (Extended Data 1). Sequence-based phylogenetic inference of SIRV2-like capsomers supports a common ancestry for these rod-shaped viruses. *Sulfolobales*-associated rod-shaped viruses (*Rudiviridae*), the ANME-1–associated rod-shaped viral family (PBV300, *Ahmunviridae*), D-B-*Nitrososphaeria*-associated rod-shaped viruses (vOTU_0179), and the two *Caldarchaeales*-associated rod-shaped viral families (*Demeterviridae* and *Gaiaviridae*) each form well-supported monophyletic clades, with both posterior probabilities and maximum-likelihood bootstrap values exceeding 0.9 (90) (Fig. 5a). However, the posterior probabilities and bootstrap values supporting the nested branching relationships among these rod-shaped viral families are all below 0.7 (70), indicating that their evolutionary relationships remain unresolved. In the structural similarity network, with the exception

of vOTU_0179, all SIRV2-like capsomers formed a single community (structural similarity ≥ 0.3),

indicating pronounced structural similarity among them (Fig. 5b and Extended Data 2). vOTU_0179

lacks one N-terminal α-helix, whereas all other monomers possess the typical two N-terminal

α-helices and three C-terminal α-helices. The monomer encoded by vOTU_0179 and one of the

monomers encoded by PBV300 (GenBank accession: UYL64970.1) are highly similar in their

C-terminal structures, both containing an additional eight β-sheets, suggesting that their capsids

likely share a common origin. However, vOTU_0179 was not observed to encode a second

SIRV2-like capsid protein as seen in PBV300, its capsomer is likely composed of *Rudiviridae*-like

homodimers.

Some *Nitrososphaeria*-associated rod-shaped viruses were observed to possess unique capsid

protein–encoding potential. Through structure-based functional annotation, we found that several

rod-shaped viruses encode two copies of SIRV2-like capsid proteins (Fig. 5c). Similar to

Hoswirudivirus, the two capsid protein copies are located adjacent to each other and to a

glycosyltransferase gene in the genome, raising the possibility that only one of the capsid proteins

has a structural function [58]. Unexpectedly, we also identified certain rod-shaped viruses (including

PBV300 and members from *Demeterviridae*) that appear to additionally encode a single gene

capable of forming a complete capsomer fold (Fig. 5c and Supplementary Fig. 2), which is likewise

located adjacent to a glycosyltransferase. This finding suggests the possibility that, in these viruses, a

complete capsomer may be assembled from a single monomer. In addition, potential capsid protein

fusion and cleavage events were observed in vOTU_0179. Although the SIRV2-like capsid protein

encoded by vOTU_0179 lacks one N-terminal α-helix, fusion with the last eight residues of the

adjacent gene product could generate a complete SIRV2-like capsid protein. Near the SIRV2-like

capsid protein gene in vOTU_0179, a neighboring gene encodes a hypothetical protein with three

N-terminal α-helices and eight C-terminal β-sheets, highly similar to the C-terminal structures of the

SIRV2-like capsid proteins encoded by vOTU_0179 and PBV300. This finding suggests that

SIRV2-like capsids may undergo dynamic fusion or cleavage events, exhibiting previously

unrecognized functional polymorphism.

Nitrososphaeria-associated head-tailed viruses exhibit typical polyphyletic characteristics and

generally form well-supported monophyletic clades in phylogenetic tree of major capsid protein

(MCP) (Fig. 6a). The results of phylogenetic inference are consistent with the observations from the

HK97-like MCP structural similarity clustering network (Fig. 6b and Extended Data 3). Eight newly

identified viral families form three *Nitrososphaeria*-specific monophyletic clades, including four

families associated with *Ca. Caldarchaeales* (*Morosviridae*, *Nyxviridae*, *Hypnosviridae*, and

Erebusviridae), one family associated with *Nitrosocaldales* (*Peleviridae*), one family associated with

Nitrososphaerales (*Freyrviridae*), two families associated with *Nitrosopumilales* (*Alboranviridae*

and *Baleariviridae*), and one family associated with deep-branching AOA-related *Nitrososphaeria*

(*Jordviridae*). In contrast, the remaining three AOA-related viral families (*Thanatosviridae*,

Charonviridae, and *Apateviridae*) are phylogenetically nested with bacterial viruses, suggesting that

some viruses may have undergone host switching during their evolution.

Novel auxiliary genes encoded by *Caldarchaeales*-associated rod-shaped viruses

Following the recommendations of Martin et al., potential auxiliary viral genes (AVGs) encoded by

these viruses or MGEs were screened [59]. A total of 376 auxiliary regulatory genes (AReGs), 235

auxiliary metabolic genes (AMGs) and 88 auxiliary physiological genes (APGs) were obtained

(Supplementary   Dataset   4).   For   AMGs,

2-polyprenyl-3-methyl-5-hydroxy-6-metoxy-1,4-benzoquinol methylase (UbiG) was found to be

encoded by four vOTUs from the newly identified rod-shaped viral family, *Demeterviridae*

(Supplementary Dataset 4). These virus-encoded UbiG-like proteins exhibit structural similarity to

their cellular counterparts, all featuring the typical Rossmann-like fold (a β-sheet of seven strands

surrounded by α-helices) (Fig. 7a). Molecular docking analyses showed that the virus-encoded

UbiG-like proteins have affinity for 2,6-dihydroxybenzoic acid, with the benzene ring inserted into

the SAM-binding pocket. Homologues of *Demeterviridae*-encoded UbiG-like protein are extremely

rare in viruses, with no high-similarity homologues found in the nr database (alignment identity and

coverage both ≥ 50%), highlighting its novelty. They form a distinct monophyletic clade that groups

with distantly related archaeal homologues in the phylogenetic tree. This indicates their connection

to archaea while retaining a clear virus-specific phylogenetic monophyly (Fig. 7a). To the best of our

knowledge, CoQ is rare in archaea and there is currently no evidence that archaea and related viruses

possess aromatic quinone biosynthesis pathway [23,57]. Whether the functions of these virus-encoded

UbiG-like proteins extend to the synthesis of menaquinone (MK), the major respiratory quinone in

archaea [57], remains unknown. However, this finding provides a rare example of archaeal viruses

encoding genes potentially related to quinone electron-carrier metabolism. [60]. For AReGs,

Demeterviridae was found to encode an ArsR-like transcriptional regulator, which was only

previously reported from spindle-shaped ssDNA viruses and head-tailed viruses [51,61]. The

Demeterviridae-encoded ArsR-like proteins forms a well-supported, long-branched monophyletic

clade with a pronounced stem branch in the phylogenetic tree and is evolutionarily distant from other

bacteriophages and archaeal viruses (Fig. 7b). Regulators of archaeal filamentous or rod-shaped

viruses are predominantly represented by RHH-fold proteins such as SvtR (*Rudiviridae*) and AvtR

(*Lipothrixviridae*) [62-64]. This result suggests that *Caldarchaeales*-associated rod-shaped viral

genomes may employ a transcriptional regulatory mechanism similar to that of spindle-shaped

viruses [61], providing clues to a previously unrecognized mechanism within this viral realm that

awaits further validation.

Viruses or MGEs serve as the primary vectors rather than donors or receptors in horizontal gene transfers

Virus–host linkages identified by spacer–protospacer matching showed lineage specificity, with no

cross-lineage protospacer sharing observed (Supplementary Fig. 3a and Supplementary Dataset 5).

After establishing virus-host linkages by spacer-protospacer matching, the direction of horizontal

gene transfers (HGTs) mediated by viruses or MGEs were explored based on phylogenetic

topologies [65]. A total of 2,509 possible HGT events between viruses or MGEs and corresponding

Nitrososphaeria lineages were identified (Supplementary Dataset 6 and Extended Data 4) [65]. For

AOA, 68.0% of transfers are bidirectional (n = 1,359), while 27.3% originate from archaea as donors

(n = 546). Only 4.7% of transfers involve viruses or MGEs as donors (n = 94). Similarly, in *Ca.*

Caldarchaeales, 56.5% of transfers are bidirectional (n = 183), 34% originate from archaea (n = 110)

and 9.6% involve viruses or MGEs as donors (n = 31) (Fig. 7c). This suggests that viruses or MGEs

serve as the primary vectors in these transfers, rather than donors or receptors.

Functionally, most HGTs are related to glycan biosynthesis and metabolism, replication and

repair and carbohydrate metabolism, accounting for 60.5% of the total HGTs (Fig. 7d). Within

glycan biosynthesis and metabolism, more than half of the HGTs (n = 283) are associated with

glycosylation and are linked to *Nitrosopumilales* (Supplementary Dataset 4 and Supplementary Dataset 6). The related functional proteins include all dolichyl-phosphate-mannose–protein mannosyltransferases, glycosyl transferases (GTs), sulfatases and N-acetyl sugar amidotransferases. Since glycosylation or glycoprotein modification on the surface of archaeal viral capsids or envelopes is involved in host interactions or enhances structural stability [66-68], it raises the possibility that *Nitrososphaeria*-associated viruses or MGEs have incorporate host-derived functional protein modules. In addition, 87.8% of the HGTs related to viral structure were found to be associated with *Ca. Caldarchaeales*. These structural proteins include tail-related proteins and MCPs from *Erebusviridae* and *Morosviridae*, further confirming the interactions between these proposed viral families and *Ca. Caldarchaeales*.

Discussion

In this study, we provide a deeper understanding of the diversity of *Nitrososphaeria*-associated viruses or MGEs, broadening the scope from previously reported AOA-associated viruses and identified hundreds of previously unrecognized *Caldarchaeales*-associated vOTUs for the first time. The virus–archaea associations identified in this study have not yet been validated through isolation experiments, but they provide valuable guidance for establishing future cultivation systems for *Nitrososphaeria* viruses, particularly those associated with *Ca. Caldarchaeales*. This genomic catalogue expands the known number of *Nitrososphaeria*-associated viruses or MGEs by over seven-fold [32-36]. The mining of *Caldarchaeales*-associated viral or MGE genomes fills a critical knowledge gap of archaeal viromes. Nine previously unknown viral families were identified, including a novel rod-shaped family and an archaeal "jumbo" virus. The distinct evolutionary placement and functional characteristics of this family were elucidated through comparative genomic and phylogenetic analyses, providing new insights into the archaeal virosphere and viral evolution.

Through phylogenetic inference and structural modelling, we revealed the evolutionary relationships among archaeal rod-shaped viruses, providing new evidence for the phylogeny and taxonomy of the *Adnaviria* viral realm. In the past, only the capsomers of *Rudiviridae* were known to be composed of homodimers[64]. Here, it was found that some *Caldarchaeales*-associated rod-shaped viruses encode multiple copies of SIRV2-like MCP genes (Figs. 4 and 5c). Given the pronounced

sequence homology between the MCPs encoded by *Demeterviridae* and those of *Rudiviridae*, and the fact that no *Rudiviridae* with homodimeric capsomers have yet been identified [30,64,69], it is possible that in *Demeterviridae* only one MCP copy is functional, while the other may have been inactivated through natural selection [70-72]. *Rudiviridae*-like viruses with a single functional MCP subunit likely represent a more ancestral state, whereas viruses encoding two related MCPs evolved through duplication and divergence [73]. The observation of a fusion gene encoding two SIRV2-like MCPs further advances this hypothesis: the ancestral capsomer of rod-shaped viruses may have been composed entirely of a single protein, which subsequently gave rise to homodimers and, later, heterodimers during evolution. Through structural modelling of the MCP encoded by vOTU_0179 and its adjacent genes, we found that this MCP appears to be undergoing ongoing cleavage, potentially generating additional protein products (Fig. 5c). Because its structure is incomplete, this MCP may no longer be functional. The MCP encoded by vOTU_0179 is structurally highly similar to one of the MCPs encoded by *Ahmunviridae* PBV300 (GenBank accession: UYL64970.1). PBV300 may retain functionality in only a single MCP (GenBank accession: UYL64989.1), thereby forming a homodimeric capsomer. However, because PBV300 also encodes a fusion gene containing two MCP subunits (GenBank accession: UYL64987.1), the bona fide capsid architecture of these rod-shaped viruses remains unknown. Taken together, our evidence indicates that *Nitrososphaeria*-associated rod-shaped viruses, *Rudiviridae*, and *Ahmunviridae* are highly similar in the evolutionary and structural characteristics of their capsid proteins. In addition, comparative genomics analyses revealed pronounced genomic synteny among these rod-shaped viruses (Supplementary Fig. 4). This raises the need to reconsider whether the placement of *Ahmunviridae* within *Maximonvirales* as a lineage parallel to *Ligamenvirales* is justified, or whether *Ahmunviridae* should instead be incorporated into *Ligamenvirales*.

Viruses have been found to encode diverse AVGs that manipulate host metabolism, thereby creating favourable conditions for completing their life cycles and indirectly contributing to global biogeochemical cycles [74-77]. In archaea-associated viruses, such as those infecting methanogenic archaea, Asgard superphylum, halophilic archaea, virus-encoded metabolic genes have been identified that potentially modulate archaeal methane-oxidization, sulphur metabolism, nucleotide and amino acid metabolisms [27,78,79]. Regarding AOA-associated viruses or MGEs, previous studies have discovered that viruses may manipulate host ammonia oxidation metabolism by carrying the

amoC gene and participate in host vitamin synthesis [34,35]. Through the exploration of functional

repertoires for *Nitrososphaeria*-associated viruses, we provide new insights into the functional

mechanisms of archaeal viruses. An important example is provided by viral ArsR-like and UbiG-like

protein, which represent a rare AReG and AMG, respectively, encoded by a previously unrecognized

rod-shaped viral family. If the viral ArsR-like protein is functional in the archaeal virocell, it might

be able to hijack the arsenic response of *Ca. Caldarchaeales*, thereby facilitating viral replication

within the host cell. Considering that the habitat of *Ca. Caldarchaeales* is rich in metals and

metalloids (including arsenic) and exhibits highly active redox dynamics[24,80,81], this process may be

essential for the completion of the *Demeterviridae* life cycle. Phylogenetically,

Demeterviridae-encoded ArsR falls on a long-branch, virus-specific clade supports their viral origin,

rather than recent horizontal gene transferring from archaea.

The more unusual finding is the viral carriage of UbiG-like protein, an enzyme implicated in

CoQ biosynthesis. Quinones (e.g. ubiquinone, menaquinone) are key electron carriers in membrane

redox chains, yet they are considered rare in archaea and corresponding viruses [23,57,82]. The detection

of *Demeterviridae*-encoded UbiG-like protein suggests a potential viral contribution to the redox

balance of archaeal membranes, such as altering the local pool of quinone electron carriers [57,83]. They

might influence electron flux, respiration efficiency, reactive oxygen species (ROS) levels, or even

redox signaling [83-85]. In hyperthermal habitats of *Ca. Caldarchaeales*, control of archaeal quinone

pools may confer a fitness advantage to related viruses by maintaining a more favorable intracellular

environment for replication. On the other hand, the *Demeterviridae*-encoded UbiG-like protein,

together with other methyltransferases encoded by archaeal viruses, may perform similar functions in

counteracting host defense mechanisms, representing an exaptation of archaeal gene functions by

viruses [86]. Since *Ca. Caldarchaeales* has not been observed to encode UbiG-like proteins, the origin

of these genes in *Demeterviridae* remains unknown. Overall, the discovery of *Demeterviridae* not

only expands the boundaries of the archaeal virosphere and offers new insights into viral evolution,

but also provides direction in how viruses might adapt to extreme environments in terms of

transcriptional regulation and host metabolic reprogramming.

Our results reinforce the view that viruses or MGEs act predominantly as gene vectors. This

supports the "virus world" model, considering viruses or other selfish genetic elements as

intermediaries in gene exchange rather than as main sources of new genes [87]. The prevalence of

bidirectional HGTs highlights reciprocal coevolution between viruses or MGEs and archaeal hosts, evoking the arms-race dynamics observed in archaeal virus–host systems [88,89]. The functional enrichment of glycan biosynthesis and glycosylation among transferred genes implies that viruses may co-opt host surface modification machinery to refine capsid stability or host recognition—a phenomenon similarly observed in eukaryotic viruses adapting envelope glycosylation [90,91]. It is hypothesized that the frequent HGT events between archaea and viruses may have driven the diversification of archaeal envelope biochemistry and the emergence of niche specialization in ammonia□oxidizing lineages.

Methods

Assembly of *Nitrososphaeria*-associated CRISPR database

The *Nitrososphaeria* phylum comprises several major clades: AOA, other AOA-related lineages and *Ca. Caldarchaeales*. To produce a comprehensive genomic dataset, reference genomes were retrieved from GenBank and IMG/M [41,42], encompassing a total of 2,137 genomic assemblages. Specifically, 1,830 assemblages (comprising 322,784 genomic contigs) were obtained from GenBank; 307 assemblages (containing 23,395 genomic contigs) were sourced from IMG/M.

To identify CRISPR spacers within these genomes, similar to previous studies [27,28], CRASS v.0.3.8 and MinCED v.0.4.2 (CRISPR recognition tool, CRT) were employed using the following parameters to detect spacers and discard those with low-complexity and short length [92,93]: word size = 7, spacer length = 26 and a minimum of three directed repeats (DRs). This step resulted in the identification of 37,485 *Nitrososphaeria*-associated spacers, which were subsequently used for the detection of viruses or MGEs and downstream analyses.

The metagenome-derived spacers that are associated with *Nitrososphaeria* were detected to further expand the diversity of these CRISPR spacers. First, a total of 170 metagenome-derived assemblages (including 58 from the Malaspina expedition and 22 from our previous studies) from diverse ecosystems were retrieved and downloaded from IMG/M and ENA (European Nucleotide Archive) (Supplementary Dataset 1)[42,45]. These metagenomes were from 23 different studies, focusing on extreme natural habitats and including 92,037,046 assembled contigs. These include four studies from deep-sea, three two from sea ice, four studies from a hypersaline environment and one

study from an acidic environment. Second, the CRISPR spacers found in these contigs were

identified using the same approach described above, producing 1,828,641 spacers and 1,918,441

DRs. Third, all DRs were aligned to *Nitrososphaeria* genomes using MMseqs2 with parameters [94]:

word size = 7, pairwise alignment identity = 1, percentage coverage of DRs = 1, mismatches = 1.

This step produced 196,551 spacers derived from 170 metagenomes. Finally, this process generated a

Nitrososphaeria-associated spacer database containing 234,036 spacer sequences (Supplementary

Fig. 1 and Extended Data 5).

As in previous studies [28], CRISPR-Cas cassette prediction and subtyping for archaeal contigs

and genomes were conducted using CRISPRcasIdentifier v1.0.0 [95], employing a combination of

Classification and Regression Trees (CART), Support Vector Machine (SVM) and Extremely

Randomized Trees (ERT) algorithms for classification and regression. The analysis was performed

with the following parameters: –mode ERT SVM CART, -hmm HMM1 HMM2 HMM3 HMM4

HMM5. Subtype assignment was determined through a consensus-based approach, where only

subtype classifications consistently predicted by all three models (CART, SVM and ERT) were

accepted. The final subtype was selected based on the highest mean probability and the lowest

standard deviation among the predictions (Supplementary Fig. 3b and Supplementary Dataset 7).

Mining and filtering *Nitrososphaeria*-associated viruses or MGEs from IMG/VR (v.4)

The IMG/VR (v.4) database, contigs from 170 metagenomes and archaeal genomes served as the

primary resources for mining *Nitrososphaeria*-associated viral or MGE genomes. At first, the

archaeal viral or MGE mining pipeline was validated using CRISPR spacers derived from archaeal

genomes and host predictions from the IMG/VR (v.4) database and other studies to minimize false

positives [33-36,38,96,97]. The 37,485 CRISPR spacers from *Nitrososphaeria* genomes were aligned to

IMG/VR (v.4) database using MMseqs2 with the parameters: word size = 7, pairwise alignment

identity = 0.9, percentage coverage of spacers = 0.9, mismatches = 1, resulting in 368 targeted.

Among these results, the hosts of 46 viruses or MGEs were also predicted to be *Nitrososphaeria* in

IMG/VR (v.4). Due to some viral contigs being assigned to non-*Nitrososphaeria* hosts in IMG/VR

(v.4) metadata, the genome-content similarity between viruses or MGEs and hosts was applied to

reduce the false positives: i) the open reading frames (ORFs) of these viral contigs were predicted by

Prodigal v.2.6.3 and aligned to the NR database (2024.05.20) with 1e-5 as the E-value cut-off [98]; ii)

the *Nitrososphaeria*-related hits were counted and normalized based on viral contig lengths; iii) for

each genome, it was observed that at least 0.2 ORFs per 1-kb genome could be matched to

Nitrososphaeria (Supplementary Dataset 2); iv) further validation was conducted to assess the

applicability of this screening criterion to previously reported *Nitrososphaeria*-associated viruses or

MGEs:

1) the 87 identified *Nitrososphaeria*-associated viral or MGE genomes were downloaded from

GenBank or provided by data owner;

2) the 46 representative genomes were generated from 48 *Nitrososphaeria*-associated viral genomes

through MMseqs2 with parameters [94]: pairwise alignment identity = 0.95, pairwise alignment

coverage for shorter sequences = 0.9;

3) the 83 out of 85 viral genomes derived from GenBank meet this criterion. Two isolated

spindle-shaped viruses (NSV4 and NSV5) failed to pass the test [37]. This indicates that the

approach is also applicable to most previously reported *Nitrososphaeria*-associated viral or

MGE genomes.

After validating the filtering approach, it was used as a cut-off for viral-host genome-content

similarity to further filter the results obtained from spacer-protospacer matching. By applying this

approach, 171 *Nitrososphaeria*-associated viruses or MGEs were filtered from 368 spacer-targeted

viral or MGE candidates. Similar to previous studies [29,99], all relevant archaeal CRISPR-Cas systems

were evaluated using CRISPRCasFinder with default parameters [100]. Among the identified CRISPR

arrays, 91.2% (n = 301) exhibited an evidence level of 4, characterized by a repeat conservation

index exceeding 70% and spacer identity below 8% (Supplementary Dataset 2). The remaining 29

viruses or MGEs, corresponding to evidence levels below 4 for CRISPR arrays, were retained for

two reasons: 1) they passed genome-content similarity filtering; 2) in genome-wide analyses, they

clustered with other viruses or MGEs possessing CRISPR arrays with an evidence level of 4.

In addition, IMG/VR (v.4) contains 1,457 viruses or MGEs that were predicted to be associated

with *Nitrososphaeria*, although these contigs were not retrieved through spacer-protospacer matching.

The 1,457 viruses or MGEs predicted to be associated with *Nitrososphaeria* in IMG/VR (v.4) were

tested. First, 699 representative contigs were clustered using MMSeq2 with parameters: pairwise

alignment identity = 0.9, pairwise alignment coverage for shorter sequences = 0.9. Second, these

contigs were tested by genome-content similarity calculation, resulting in 441

Nitrososphaeria-associated viruses or MGEs. Finally, 612 *Nitrososphaeria*-associated viruses or

MGEs were identified and filtered from IMG/VR (v.4).The completeness of these viral or MGEs

were evaluate using CheckV [101].

Expanded mining of *Nitrososphaeria*-associated viruses or MGEs from 170 metagenomes

The *Nitrososphaeria*-associated CRISPR spacer database was aligned to from 170 metagenomes

using MMseqs2 with the same approach and parameters [94]. Contigs containing DRs and those shorter

than 2-kb were removed before performing alignment. This resulted in 1,951 contigs matched by

Nitrososphaeria-associated CRISPR spacers. To remove duplications, 1,951 contigs were aligned to

previously identified *Nitrosospaherota*-associated viruses or MGEs set using MMseqs2 with same

parameters same parameters [94]. This step generated 1,895 contigs, including 1,396 representative

contigs clustered using the same approach described before. To recall as many as possible viruses or

MGEs, these contigs were used to filter *Nitrososphaeria*-associated viruses or MGEs using three

multiple viral identification pipelines, including geNomad, VirSorter2 and DeepVirFinder[102-104].

Multiple pipelines were employed for virus identification based on the following considerations: 1)

geNomad demonstrates superior overall classification performance compared to other pipelines [103]; 2)

VirSorter2 (VS2) is recommended when integrating multiple pipelines [105]; 3) DeepVirFinder (DVF),

utilizing an alignment-free approach, is well-suited for identifying unique viral sequences [102]; 4) the

identification results from each pipeline complement one another, enhancing recall while mitigating

cellular contamination in the dataset[106-108]. For each pipeline, precision was prioritized over

sensitivity; therefore, stringent virus detection strategies were adopted to minimize potential false

positives: 1) geNomad was used to detect viral or MGE contigs using conserved mode

("--conservative"), resulting in a more strict set of viruses with high confidence; 2) VS2 was used to

detect viral contigs under high-confidence mode ('--high-confidence-only'), retaining results with

classification score $\geq$ 0.9 and classification score $\leq$ 0.7 with viral hallmark $\geq$ 1; 3) The DVF used to

detect viral contig with parameters: score $\geq$ 0.9 and p-value $\leq$ 0.01. To validate this hybrid viral

detection pipeline, 20% of sequences were randomly selected from previously identified

Nitrososphaeria-associated viruses or MGE and were used as a test set to evaluate the efficiency of

the approach. Based on the test, this viral or MGE identification approach achieved a 74%

recruitment rate, which was considered acceptable for further viral or MGE identification

(Supplementary Dataset 8).

Using the validated hybrid viral identification approach, 419 viral or MGE contigs were

identified from 1,396 metagenome-derived contigs. These contigs had already been targeted by

Nitrososphaeria-associated CRISPR spacers in previous steps. The viral-host genome-content

similarity was used to filter these contigs as described before, resulting 108

Nitrososphaeria-associated viral or MGEs. Relevant archaeal CRISPR-Cas systems of these contigs

were evaluated using CRISPRCasFinder with default parameters [100]. The completeness of these viral

or MGEs was evaluated using CheckV [101].

Detection of viruses or MGEs integrated in *Nitrososphaeria* genomes

Proviruses integrated into *Nitrososphaeria* genomes were identified using the "find-proviruses"

program in geNomad [103]. Integrases were first detected with the MMseqs2 plugin in geNomad [94],

applying an E-value cut-off of 1e-5. Additional parameters were set as follows: a minimum

gene-level score of 0.4 for flagging proviral genes using the conditional random field model; a

minimum total virus marker score of 12 for proviruses lacking integrases or not located at scaffold

edges; a minimum total virus marker score of 8 for proviruses encoding integrases or located at

scaffold edges; a maximum allowed distance of 10-kb between provirus boundaries and integrases

for boundary extension; and a maximum allowed distance of 5-kb between provirus boundaries and

tRNAs for boundary extension. This process resulted in the identification of 30

Nitrososphaeria-associated proviruses.

To remove redundancy, the 30 proviral or MGE genomes performed self-alignment and were

compared against previously identified *Nitrososphaeria*-associated viral or MGE genomes using the

same approach and parameters described above. This filtering step yielded 24 non-redundant

Nitrososphaeria-associated proviruses (Supplementary Dataset 2).

Host assignments of *Nitrososphaeria*-associated viruses or MGEs

441 The host assignments of *Nitrososphaeria*-associated viruses or MGEs were primarily based on: 1)

CRISPR-spacers-associated host genomes; 2) information provided by previous studies; 3) genome-content similarity between viruses or MGEs and hosts. For *Nitrososphaeria*-associated viruses or MGEs that match CRISPR spacers from multiple distinct hosts, similar with previous study [29], the host of a virus or MGE targeted by CRISPR spacers was assigned based on the taxonomy of the genome from which the spacer with the highest identity originated. For *Nitrososphaeria*-associated viruses or MGEs that were identified based on oligonucleotide pattern similarity, the most possible host genomes associated with these viruses or MGEs were determined through alignment against *Nitrososphaeria* genomes using MMseqs2 with the parameters: 1e-5 as the E-value, iteratively searched two times [94]. A host genome was linked to a viral or MGE genome if they have the most shared ORFs. This approach was also applied in IMG/VR (v.4) [96,109]. For *Nitrososphaeria*-associated viruses or MGEs reported in previous studies, some host information was retrieved from related literature. For those viruses or MGEs with uncertain host information [35], the related viral ORFs were subject to MMseqs2 against all *Nitrososphaeria* genomes with the same approach. The 35 Marthaviruses derived from López-Pérez et al. (PRJNA484324) have been authorized for use and are included in this study [34]. The archaeal host phylogenetic tree was calculated based on 46 concatenated ribosomal proteins by IQ-Tree2 (Supplementary Note 2)[110]. The archaeal lineages were determined based on their phylogenetic placements and previous studies [2,8,21,22,24,111,112].

Comparative genomics and family-/species-level demarcation of *Nitrososphaeria*-associated viruses or MGEs

For species-level unit assignments, viral or MGE species (vOTUs or MGE OTUs) were defined using 95% nucleotide identity threshold over 90% of the full length using MMseqs2. For family-level unit assignments, similar to previous studies [27,29,31], new families were assigned based on the genome-content similarity network and the genome-wide proteomic tree. Using DNA virus sequences longer than 2-kb from NCBI RefSeq v.223 as reference genomes, vConTACT2 was applied (with default parameters) to compute the genome-content-based similarity network of those *Nitrososphaeria*-associated viruses or MGEs. *Nitrososphaeria*-associated viral or MGE genome pairs that shared at least 20% protein clusters and encoded at least six genes were filtered from

genome-content-based similarity network, generating a subset of network, which was used to guide taxonomic assignments. In the network, viruses not related to *Nitrososphaeria*-associated viruses or MGEs were discarded to achieve a clear layout. Genome-wide proteomic tree of those genomes included in the subset of network were calculated by ViPTree v.4. The viruses or MGEs were assigned to the same family if they were represented by a cohesive group in the genome-content-based similarity network and a monophyletic group in genome-wide proteomic tree (with threshold 0.05) [29]. The network and tree were visualized by Cytoscape v.3.10.2 and iToL v.6, respectively [113,114]. The nomenclature of the proposed viral family was referred to Greek or Norse mythology (Supplementary Note 3).

Annotation of viral or MGEs genomes

Viral or MGE species (vOTUs or MGE OTUs) were defined using 95% nucleotide identity threshold over 90% of the full length using MMseqs2. The family-level taxonomic assignments of *Nitrososphaeria*-associated viruses or MGEs based on reference taxonomic database were assigned using 1) VITAP with custom database: this database includes all viral reference genomes collected in ICTV VMR-MSL40, high-quality vOTUs with genus-level taxonomy collected in IMG/VR (v.4) and all viral genomes collected in NCBI RefSeq v.223 [96,115,116]; 2) CAT with NCBI RefSeq v.223 database [116,117]; 3) geNomad with default parameters [103]. The CAT only assigned class/family level units for 10 contigs; the geNomad and VITAP assigned class/family level units for 683 and 676 contigs, respectively. Hence, the taxonomic assignments were mainly based on geNomad and VITAP. If geNomad or VITAP showed consistency at higher taxonomic levels (Phylum or Class), but one of the pipelines produced annotations at lower taxonomic levels, then the results with the lower taxonomic level annotations were used. The results with discrepancies between VITAP and geNomad were compared against the NCBI RefSeq v.223 using MMseqs2 with default parameters and subsequently subjected to manual verification. The consensus VCs generated from vConTACT2 were used to expanded and furtherly curated the viral taxonomic assignment [118]. The members of each proposed family were furtherly expanded based on consensus vOTU.

ORFs encoded in *Nitrososphaeria*-associated viruses or MGEs were predicted by Prodigal v.2.6.3 with metagenome mode and annotated using the KEGG Ortholog database (January, 2025) [98],

Pfam-A v.35, COG2020, Swiss-Prot database (January, 2025), VOG223 database and NR database

(August, 2025), and HHpred [119-122]. Potential AVGs encoded by these viruses or MGEs with length

over 10-kb were screened based on proposal from Martin et al.[59]. The functional screening criteria

excluded all genes associated with viral core functions (replication, recombination, repair,

transcription, translation, packaging and lysis), nucleotide metabolism and one-carbon pool

processes, nucleic acid modification and immune evasion, structural and host recognition functions

(including all CAZymes and polysaccharide synthesis related functions), dNTP supply, as well as

various proteases or multifunctional enzymes. In addition, candidate genes were filtered by

examining whether viral genes were present in their flanking regions. Considering the

incompleteness of viral contigs, once an AVG was confirmed, its homologues were also considered

as AVGs, even if no viral genes were present in the flanking regions. The maximum-likelihood

phylogenetic trees of ArsR and UbiG were constructed. For ArsR, protein sequences from archaea,

bacteria, bacteriophages and archaeal viruses were retrieved from InterPro (PF01022). After merging

these with the ArsR protein sequences encoded by rod-shaped viruses, CD-HIT v. 4.8 was used to

cluster the sequences from different sources (at 0.5 pairwise identity with global alignment). Seqkit

v.2.5 was then applied to randomly sample 100 sequences from each category [123], or to include all

sequences if fewer than 100 were available. Multiple sequence alignment (MSA) was performed

with MUSCLE5 [124] and trimAl v.1.4 was used to remove sites with more than 30% gaps [125].

Maximum-likelihood phylogenetic inference of the MSAs was carried out with IQ-TREE2 [110], with

1,000 bootstraps and LG+F+R10 as the best-fit substitution model (Extended Data 8). For UbiG,

protein sequences from archaea, bacteria and eukaryotes were retrieved from InterPro (IPR004951).

These were merged with rod-shaped virus-encoded UbiG-like sequences and maximum-likelihood

phylogenetic inference was performed using a similar procedure, with Blosum62+R10 as the best-fit

substitution model (Extended Data 6). The maximum-likelihood phylogenetic trees were visualized

with iTOL v.6 [114].

Determining of *Nitrososphaeria*-associated viral or MGE horizontal gene transfer events

HGT events between *Nitrososphaeria*-associated viruses (vOTUs) and their archaeal hosts were

identified following an established pipeline [65]. After removal of proviral sequences, viral and

archaeal sequences were aligned and shared protein families (PFs) were identified. Potential contamination was rigorously screened based on lineage, GC content, scaffold length and genomic context, retaining only robust alignments (Supplementary Note 4 and Supplementary Dataset 9).

Filtered protein families underwent phylogenetic analysis using MUSCLE5 alignment, trimAl v.1.4 trimming and IQ-TREE2 tree construction with the LG+F+R5 model. Trees were rooted via minimal ancestral deviation and potential HGT events were classified based on the phylogenetic relationships between viral and archaeal sequences (Extended Data 4). Transfer direction labels ("v-from-a", "v-to-a", "v-?-a" and "non-HGT") were assigned following predefined criteria considering sister and cousin clade compositions. Iterative screening was performed with temporary reassignment of labels to ensure accurate HGT detection across multiple rounds of analysis (Supplementary Note 4).

Phylogenetic inference of *Nitrososphaeria*-associated viruses or MGEs

To investigate the evolutionary relationships among rod-shaped and head-tailed viruses associated with *Nitrososphaeria* within the virosphere context, phylogenetic analyses were performed based on key viral proteins, including the SIRV2-like MCPs and HK97-like MCPs. For rod-shaped viruses, SIRV2-like MCP sequences were aligned with viral genomes of *Adnaviria* realm from GenBank using MMSeqs2 (E-value $\leq$ 1e-5). Some viruses within *Rudiviridae* (*Hoswirudivirus*) encoded adjacent duplicate MCP subunits, proteins adjacent to these SIRV2-like MCPs were annotated using HHpred to determine whether additional distantly related SIRV2-like MCPs are present. SIRV2-like MCPs directly adjacent to glycosyl transferases were used to generate homodimer sequences, aligned using MUSCLE5 (ensemble mode) to generate MSAs. Sites with more than 30% gaps were removed using trimAl v.1.4 [125]. The trimmed MSAs were subsequently used for BI (300 million generations) and ML (3000 bootstraps) phylogenetic inference based on MrBayes and IQ-TREE2, respectively [110,126]. The DUF732 domain-containing protein (PF05305) with three α helices (similar with C-terminal α helices of SIRV2-like MCP), identified via FoldSeek structural alignment as the closest homolog [127], served as an outgroup for rooting the rod-shaped viral MCP phylogeny.

HK97-like MCP sequences exhibited significant diversity, encompassing multiple families from Pfam-A v.35, CDD, and VOG223 [120,128,129]. Thus, a structure-guided approach was employed to

perform phylogenetic inference of HK97-like MCPs. First, putative HK97-like MCP sequences were collected through domain annotation. Second, structural modelling of these sequences was performed using AlphaFold2 implemented via ColabFold. Third, all MCPs were aligned under the guidance of structural information using Reseek and MUSCLE5 to generate a structure-based multiple sequence alignment (MSA)[130]. Fourth, the resulting MSA was used to build a hidden HMM with HMMER3 [131], which was then applied to search for homologs among all *Nitrososphaeria*-associated viral proteins (E-value ≤ 1e-5). Fifth, based on all identified *Nitrososphaeria*-associated viral HK97-like MCPs, BLASTp searches were conducted against RefSeq v.223 to retrieve viral homologs using thresholds of 0.5 for both pairwise coverage and sequence identity. Sixth, all newly identified HK97-like MCP sequences, together with archaeal Dodecin sequences as an outgroup [86], were subjected to a second round of structural modelling using AlphaFold2, and a second structure-based MSA was constructed following the same procedure. Finally, this MSA was used for phylogenetic inference using the same methods.

Structure modelling and molecular docking for viral proteins

All structural modelling was performed using AlphaFold2 implemented via ColabFold with default parameters. The models with the highest pLDDT scores were selected as the best-predicted structures. Foldseek was used to perform all-vs-all structural alignments for SIRV2-like MCPs and HK97-like MCPs separately (E-value ≤ 1e-5; identity ≤ 0.3) [127]. The resulting alignments were visualized as networks in Cytoscape [113], using bitscore as the edge weight. For molecular docking of UbiG, the following steps were applied: 1) structural data of a potential substrate containing an aromatic ring and phenolic hydroxyl groups (2,6-dihydroxybenzoic acid) were retrieved from PubChem [132]; 2) docking simulations were performed using HDOCK [133], with the search space covering the entire β-sheet core region; and 3) docking results were visualized using PyMOL [134]. For molecular docking of ArsR, the following steps were conducted: 1) ArsR monomers were assembled into the biologically active dimeric form; 2) A-DNA was generated from a random genomic fragment of *Demeterviridae* (vOTU_0592) using Web 3DNA 2.0; 3) docking simulations were performed using HDOCK, with the search space covering the entire cluster of C-terminal α-helices; and 4) docking results were visualized using PyMOL.

Data availability

All nucleotide and protein sequences of viral or mobile genetic elements, spacers from the CRISPR

database, phylogenetic trees, structural prediction files are available in FigShare

(https://figshare.com/s/a2e94d686028b674a905). The related metagenomes used in this study can be

accessed via the URL provided in Supplementary Dataset 1 through IMG/M. Related genomes from

IMG/VR database is available at https://img.jgi.doe.gov/cgi-bin/vr/main.cgi. Other information and

metadata are provided as Supplementary Datasets.

Code availability

No custom code was generated and used.

Reference

1. Nunoura, T., Takaki, Y., Kakuta, J., Nishi, S., Sugahara, J., Kazama, H., Chee, G.J., Hattori, M., Kanai, A., Atomi, H., et al. (2011). Insights into the evolution of Archaea and eukaryotic protein modifier systems revealed by the genome of a novel archaeal group. Nucleic Acids Res *39*, 3204-3223. 10.1093/nar/gkq1228.

2. Spang, A., Caceres, E.F., and Ettema, T.J.G. (2017). Genomic exploration of the diversity, ecology, and evolution of the archaeal domain of life. Science *357*. 10.1126/science.aaf3883.

3. Alfreider, A., Grimus, V., Luger, M., Ekblad, A., Salcher, M.M., and Summerer, M. (2018). Autotrophic carbon fixation strategies used by nitrifying prokaryotes in freshwater lakes. FEMS Microbiol Ecol *94*. 10.1093/femsec/fiy163.

4. Klotz, F., Kitzinger, K., Ngugi, D.K., Busing, P., Littmann, S., Kuypers, M.M.M., Schink, B., and Pester, M. (2022). Quantification of archaea-driven freshwater nitrification from single cell to ecosystem levels. ISME J *16*, 1647-1656. 10.1038/s41396-022-01216-9.

5. Li, Y., Chen, J., Lin, Y., Zhong, C., Jing, H., and Liu, H. (2024). Thaumarchaeota from deep-sea methane seeps provide novel insights into their evolutionary history and ecological implications. Microbiome *12*, 197. 10.1186/s40168-024-01912-y.

6. Brochier-Armanet, C., Boussau, B., Gribaldo, S., and Forterre, P. (2008). Mesophilic Crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. Nat Rev Microbiol *6*, 245-252. 10.1038/nrmicro1852.

7. Pester, M., Schleper, C., and Wagner, M. (2011). The Thaumarchaeota: an emerging view of their phylogeny and ecophysiology. Curr Opin Microbiol *14*, 300-306. 10.1016/j.mib.2011.04.007.

8. Zhong, H., Lehtovirta-Morley, L., Liu, J., Zheng, Y., Lin, H., Song, D., Todd, J.D., Tian, J., and Zhang, X.H. (2020). Novel insights into the Thaumarchaeota in the deepest oceans: their metabolism and potential adaptation mechanisms. Microbiome *8*, 78. 10.1186/s40168-020-00849-2.

9. Kuypers, M.M.M., Marchant, H.K., and Kartal, B. (2018). The microbial nitrogen-cycling network. Nat Rev Microbiol *16*, 263-276. 10.1038/nrmicro.2018.9.

10. Prosser, J.I., and Nicol, G.W. (2012). Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. Trends Microbiol *20*, 523-531. 10.1016/j.tim.2012.08.001.

11. Martens-Habbena, W., Berube, P.M., Urakawa, H., de la Torre, J.R., and Stahl, D.A. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. Nature *461*, 976-979. 10.1038/nature08465.

12. Stahl, D.A., and de la Torre, J.R. (2012). Physiology and diversity of ammonia-oxidizing archaea. Annu Rev Microbiol *66*, 83-101. 10.1146/annurev-micro-092611-150128.

13. Walker, C.B., de la Torre, J.R., Klotz, M.G., Urakawa, H., Pinel, N., Arp, D.J., Brochier-Armanet, C., Chain, P.S., Chan, P.P., Gollabgir, A., et al. (2010). Nitrosopumilus maritimus genome reveals unique mechanisms for nitrification and autotrophy in globally distributed marine crenarchea. Proc Natl Acad Sci U S A *107*, 8818-8823. 10.1073/pnas.0913533107.

14. Könneke, M., Schubert, D.M., Brown, P.C., Hügler, M., Standfest, S., Schwander, T., Schada von Borzyskowski, L., Erb, T.J., Stahl, D.A., and Berg, I.A. (2014). Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway for CO2 fixation. Proceedings of the National Academy of Sciences *111*, 8239-8244.

15. Ren, M., Feng, X., Huang, Y., Wang, H., Hu, Z., Clingenpeel, S., Swan, B.K., Fonseca, M.M., Posada, D., Stepanauskas, R., et al. (2019). Phylogenomics suggests oxygen availability as a driving force in Thaumarchaeota evolution. ISME J *13*, 2150-2161. 10.1038/s41396-019-0418-8.

16. Abby, S.S., Kerou, M., and Schleper, C. (2020). Ancestral Reconstructions Decipher Major Adaptations of

Ammonia-Oxidizing Archaea upon Radiation into Moderate Terrestrial and Marine Environments. mBio *11*. 10.1128/mBio.02371-20.

17. Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., and Oakley, B.B. (2005). Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. Proc Natl Acad Sci U S A *102*, 14683-14688. 10.1073/pnas.0506625102.

18. Nelkner, J., Huang, L., Lin, T.W., Schulz, A., Osterholz, B., Henke, C., Blom, J., Puhler, A., Sczyrba, A., and Schluter, A. (2023). Abundance, classification and genetic potential of Thaumarchaeota in metagenomes of European agricultural soils: a meta-analysis. Environ Microbiome *18*, 26. 10.1186/s40793-023-00479-9.

19. Biggs-Weber, E., Aigle, A., Prosser, J.I., and Gubry-Rangin, C. (2020). Oxygen preference of deeply-rooted mesophilic thaumarchaeota in forest soil. Soil Biology and Biochemistry *148*, 107848. https://doi.org/10.1016/j.soilbio.2020.107848.

20. Aylward, F.O., and Santoro, A.E. (2020). Heterotrophic Thaumarchaea with Small Genomes Are Widespread in the Dark Ocean. mSystems *5*. 10.1128/mSystems.00415-20.

21. Zhang, R.Y., Wang, Y.R., Liu, R.L., Rhee, S.K., Zhao, G.P., and Quan, Z.X. (2024). Metagenomic characterization of a novel non-ammonia-oxidizing Thaumarchaeota from hadal sediment. Microbiome *12*, 7. 10.1186/s40168-023-01728-2.

22. Hua, Z.S., Qu, Y.N., Zhu, Q., Zhou, E.M., Qi, Y.L., Yin, Y.R., Rao, Y.Z., Tian, Y., Li, Y.X., Liu, L., et al. (2018). Genomic inference of the metabolism and evolution of the archaeal phylum Aigarchaeota. Nat Commun *9*, 2832. 10.1038/s41467-018-05284-4.

23. Beam, J.P., Jay, Z.J., Schmid, M.C., Rusch, D.B., Romine, M.F., Jennings Rde, M., Kozubal, M.A., Tringe, S.G., Wagner, M., and Inskeep, W.P. (2016). Ecophysiology of an uncultivated lineage of Aigarchaeota from an oxic, hot spring filamentous 'streamer' community. ISME J *10*, 210-224. 10.1038/ismej.2015.83.

24. Buessecker, S., Palmer, M., Lai, D., Dimapilis, J., Mayali, X., Mosier, D., Jiao, J.Y., Colman, D.R., Keller, L.M., St John, E., et al. (2022). An essential role for tungsten in the ecology and evolution of a previously uncultivated lineage of anaerobic, thermophilic Archaea. Nat Commun *13*, 3773. 10.1038/s41467-022-31452-8.

25. Prangishvili, D., Bamford, D.H., Forterre, P., Iranzo, J., Koonin, E.V., and Krupovic, M. (2017). The enigmatic archaeal virosphere. Nat Rev Microbiol *15*, 724-739. 10.1038/nrmicro.2017.125.

26. Al-Shayeb, B., Schoelmerich, M.C., West-Roberts, J., Valentin-Alvarado, L.E., Sachdeva, R., Mullen, S., Crits-Christoph, A., Wilkins, M.J., Williams, K.H., Doudna, J.A., and Banfield, J.F. (2022). Borgs are giant genetic elements with potential to expand metabolic capacity. Nature *610*, 731-736. 10.1038/s41586-022-05256-1.

27. Medvedeva, S., Sun, J., Yutin, N., Koonin, E.V., Nunoura, T., Rinke, C., and Krupovic, M. (2022). Three families of Asgard archaeal viruses identified in metagenome-assembled genomes. Nat Microbiol *7*, 962-973. 10.1038/s41564-022-01144-6.

28. Rambo, I.M., Langwig, M.V., Leao, P., De Anda, V., and Baker, B.J. (2022). Genomes of six viruses that infect Asgard archaea from deep-sea sediments. Nat Microbiol *7*, 953-961. 10.1038/s41564-022-01150-8.

29. Medvedeva, S., Borrel, G., Krupovic, M., and Gribaldo, S. (2023). A compendium of viruses from methanogenic archaea reveals their diversity and adaptations to the gut environment. Nat Microbiol *8*, 2170-2182. 10.1038/s41564-023-01485-w.

30. Laso-Perez, R., Wu, F., Cremiere, A., Speth, D.R., Magyar, J.S., Zhao, K., Krupovic, M., and Orphan, V.J. (2023). Evolutionary diversification of methanotrophic ANME-1 archaea and their expansive virome. Nat Microbiol. 10.1038/s41564-022-01297-4.

31. Sakai, H.D., Nur, N., Kato, S., Yuki, M., Shimizu, M., Itoh, T., Ohkuma, M., Suwanto, A., and Kurosawa, N. (2022). Insight into the symbiotic lifestyle of DPANN archaea revealed by cultivation and genome analyses.

Proc Natl Acad Sci U S A *119*. 10.1073/pnas.2115449119.

32.   Krupovic, M., Spang, A., Gribaldo, S., Forterre, P., and Schleper, C. (2011). A thaumarchaeal provirus testifies for an ancient association of tailed viruses with archaea. Biochem Soc Trans *39*, 82-88. 10.1042/BST0390082.

33.   Krupovic, M., Makarova, K.S., Wolf, Y.I., Medvedeva, S., Prangishvili, D., Forterre, P., and Koonin, E.V. (2019). Integrated mobile genetic elements in Thaumarchaeota. Environ Microbiol *21*, 2056-2078. 10.1111/1462-2920.14564.

34.   Lopez-Perez, M., Haro-Moreno, J.M., de la Torre, J.R., and Rodriguez-Valera, F. (2019). Novel Caudovirales associated with Marine Group I Thaumarchaeota assembled from metagenomes. Environ Microbiol *21*, 1980-1988. 10.1111/1462-2920.14462.

35.   Ahlgren, N.A., Fuchsman, C.A., Rocap, G., and Fuhrman, J.A. (2019). Discovery of several novel, widespread, and ecologically distinct marine Thaumarchaeota viruses that encode amoC nitrification genes. ISME J *13*, 618-631. 10.1038/s41396-018-0289-4.

36.   Zhou, Y., Zhou, L., Yan, S., Chen, L., Krupovic, M., and Wang, Y. (2023). Diverse viruses of marine archaea discovered using metagenomics. Environ Microbiol *25*, 367-382. 10.1111/1462-2920.16287.

37.   Kim, J.G., Kim, S.J., Cvirkaite-Krupovic, V., Yu, W.J., Gwak, J.H., Lopez-Perez, M., Rodriguez-Valera, F., Krupovic, M., Cho, J.C., and Rhee, S.K. (2019). Spindle-shaped viruses infect marine ammonia-oxidizing thaumarchaea. Proc Natl Acad Sci U S A *116*, 15645-15650. 10.1073/pnas.1905682116.

38.   Liu, Y., Brandt, D., Ishino, S., Ishino, Y., Koonin, E.V., Kalinowski, J., Krupovic, M., and Prangishvili, D. (2019). New archaeal viruses discovered by metagenomic analysis of viral communities in enrichment cultures. Environ Microbiol *21*, 2002-2014. 10.1111/1462-2920.14479.

39.   Lee, S., Sieradzki, E.T., Nicol, G.W., and Hazard, C. (2023). Propagation of viral genomes by replicating ammonia-oxidising archaea during soil nitrification. ISME J *17*, 309-314. 10.1038/s41396-022-01341-5.

40.   Lee, S., Hazard, C., and Nicol, G.W. (2024). Activity of novel virus families infecting soil nitrifiers is concomitant with host niche differentiation. ISME J *18*. 10.1093/ismejo/wrae205.

41.   Nayfach, S., Roux, S., Seshadri, R., Udwary, D., Varghese, N., Schulz, F., Wu, D., Paez-Espino, D., Chen, I.M., Huntemann, M., et al. (2021). A genomic catalog of Earth's microbiomes. Nat Biotechnol *39*, 499-509. 10.1038/s41587-020-0718-6.

42.   Chen, I.A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek, P., Ritter, S., Varghese, N., Seshadri, R., et al. (2021). The IMG/M data management and analysis system v.6.0: new tools and advanced capabilities. Nucleic Acids Res *49*, D751-D763. 10.1093/nar/gkaa939.

43.   Sharma, S., Lee, M., Reinmann, C.S., Pumneo, J., Cutright, T.J., and Senko, J.M. (2020). Impact of acid mine drainage chemistry and microbiology on the development of efficient Fe removal activities. Chemosphere *249*, 126117. 10.1016/j.chemosphere.2020.126117.

44.   Zhang, X., Xu, W., Liu, Y., Cai, M., Luo, Z., and Li, M. (2018). Metagenomics Reveals Microbial Diversity and Metabolic Potentials of Seawater and Surface Sediment From a Hadal Biosphere at the Yap Trench. Front Microbiol *9*, 2402. 10.3389/fmicb.2018.02402.

45.   Acinas, S.G., Sanchez, P., Salazar, G., Cornejo-Castillo, F.M., Sebastian, M., Logares, R., Royo-Llonch, M., Paoli, L., Sunagawa, S., Hingamp, P., et al. (2021). Deep ocean metagenomes provide insight into the metabolic architecture of bathypelagic microbial communities. Commun Biol *4*, 604. 10.1038/s42003-021-02112-2.

46.   Gao, C., Liang, Y., Jiang, Y., Paez-Espino, D., Han, M., Gu, C., Wang, M., Yang, Y., Liu, F., Yang, Q., et al. (2022). Virioplankton assemblages from challenger deep, the deepest place in the oceans. iScience *25*, 104680. 10.1016/j.isci.2022.104680.

47.   Karner, M.B., DeLong, E.F., and Karl, D.M. (2001). Archaeal dominance in the mesopelagic zone of the Pacific Ocean. Nature *409*, 507-510. 10.1038/35054051.

48. Schattenhofer, M., Fuchs, B.M., Amann, R., Zubkov, M.V., Tarran, G.A., and Pernthaler, J. (2009). Latitudinal distribution of prokaryotic picoplankton populations in the Atlantic Ocean. Environ Microbiol *11*, 2078-2093. 10.1111/j.1462-2920.2009.01929.x.

49. Pester, M., Rattei, T., Flechl, S., Grongroft, A., Richter, A., Overmann, J., Reinhold-Hurek, B., Loy, A., and Wagner, M. (2012). amoA-based consensus phylogeny of ammonia-oxidizing archaea and deep sequencing of amoA genes from soils of four different geographic regions. Environ Microbiol *14*, 525-539. 10.1111/j.1462-2920.2011.02666.x.

50. Vetriani, C., Jannasch, H.W., MacGregor, B.J., Stahl, D.A., and Reysenbach, A.L. (1999). Population structure and phylogenetic characterization of marine benthic Archaea in deep-sea sediments. Appl Environ Microbiol *65*, 4375-4384. 10.1128/AEM.65.10.4375-4384.1999.

51. Liu, Y., Demina, T.A., Roux, S., Aiewsakun, P., Kazlauskas, D., Simmonds, P., Prangishvili, D., Oksanen, H.M., and Krupovic, M. (2021). Diversity, taxonomy, and evolution of archaeal viruses of the class Caudoviricetes. PLoS Biol *19*, e3001442. 10.1371/journal.pbio.3001442.

52. Adriaenssens, E., Kropinski, A.M., Turner, D., Krupovic, M., Millard, A., Dutilh, B.E., Oksanen, H.M., Lood, C., Reyes, A., and Moraru, C. (2022). Taxonomy proposal 2021: Abolish the order Caudovirales and the families Myoviridae, Siphoviridae and Podoviridae (Caudoviricetes). In ICTV Online: International Committee on Taxonomy of Viruses (ICTV), (International Committee on Taxonomy of Viruses (ICTV)).

53. Krupovic, M., Cvirkaite-Krupovic, V., Iranzo, J., Prangishvili, D., and Koonin, E.V. (2018). Viruses of archaea: Structural, functional, environmental and evolutionary genomics. Virus Res *244*, 181-193. 10.1016/j.virusres.2017.11.025.

54. Weynberg, K.D., Allen, M.J., Ashelford, K., Scanlan, D.J., and Wilson, W.H. (2009). From small hosts come big viruses: the complete genome of a second Ostreococcus tauri virus, OtV-1. Environ Microbiol *11*, 2821-2839. 10.1111/j.1462-2920.2009.01991.x.

55. Kelman, L.M., O'Dell, W.B., and Kelman, Z. (2020). Unwinding 20 Years of the Archaeal Minichromosome Maintenance Helicase. J Bacteriol *202*. 10.1128/JB.00729-19.

56. Krupovic, M., Gribaldo, S., Bamford, D.H., and Forterre, P. (2010). The evolutionary history of archaeal MCM helicases: a case study of vertical evolution combined with hitchhiking of mobile genetic elements. Mol Biol Evol *27*, 2716-2732. 10.1093/molbev/msq161.

57. Elling, F.J., Becker, K.W., Konneke, M., Schroder, J.M., Kellermann, M.Y., Thomm, M., and Hinrichs, K.U. (2016). Respiratory quinones in Archaea: phylogenetic distribution and application as biomarkers in the marine environment. Environ Microbiol *18*, 692-707. 10.1111/1462-2920.13086.

58. Wang, F., Baquero, D.P., Beltran, L.C., Su, Z., Osinski, T., Zheng, W., Prangishvili, D., Krupovic, M., and Egelman, E.H. (2020). Structures of filamentous viruses infecting hyperthermophilic archaea explain DNA stabilization in extreme environments. Proc Natl Acad Sci U S A *117*, 19643-19652. 10.1073/pnas.2011125117.

59. Martin, C., Emerson, J.B., Roux, S., and Anantharaman, K. (2025). A call for caution in the biological interpretation of viral auxiliary metabolic genes. Nat Microbiol *10*, 2122-2129. 10.1038/s41564-025-02095-4.

60. Guerra, R.M., and Pagliarini, D.J. (2023). Coenzyme Q biochemistry and biosynthesis. Trends Biochem Sci *48*, 463-476. 10.1016/j.tibs.2022.12.006.

61. Frols, S., Gordon, P.M., Panlilio, M.A., Schleper, C., and Sensen, C.W. (2007). Elucidating the transcription cycle of the UV-inducible hyperthermophilic archaeal virus SSV1 by DNA microarrays. Virology *365*, 48-59. 10.1016/j.virol.2007.03.033.

62. Guilliere, F., Peixeiro, N., Kessler, A., Raynal, B., Desnoues, N., Keller, J., Delepierre, M., Prangishvili, D., Sezonov, G., and Guijarro, J.I. (2009). Structure, function, and targets of the transcriptional regulator SvtR from the hyperthermophilic archaeal virus SIRV1. J Biol Chem *284*, 22222-22237. 10.1074/jbc.M109.029850.

63. Peixeiro, N., Keller, J., Collinet, B., Leulliot, N., Campanacci, V., Cortez, D., Cambillau, C., Nitta, K.R., Vincentelli, R., Forterre, P., et al. (2013). Structure and function of AvtR, a novel transcriptional regulator from a hyperthermophilic archaeal lipothrixvirus. J Virol *87*, 124-136. 10.1128/JVI.01306-12.

64. Krupovic, M., Kuhn, J.H., Wang, F., Baquero, D.P., Dolja, V.V., Egelman, E.H., Prangishvili, D., and Koonin, E.V. (2021). Adnaviria: a New Realm for Archaeal Filamentous Viruses with Linear A-Form Double-Stranded DNA Genomes. J Virol *95*, e0067321. 10.1128/JVI.00673-21.

65. Irwin, N.A.T., Pittis, A.A., Richards, T.A., and Keeling, P.J. (2022). Systematic evaluation of horizontal gene transfer between eukaryotes and viruses. Nat Microbiol *7*, 327-336. 10.1038/s41564-021-01026-3.

66. Gebhard, L.J., Vershinin, Z., Alarcon-Schumacher, T., Eichler, J., and Erdmann, S. (2023). Influence of N-Glycosylation on Virus-Host Interactions in Halorubrum lacusprofundi. Viruses *15*. 10.3390/v15071469.

67. Quemin, E.R., Pietila, M.K., Oksanen, H.M., Forterre, P., Rijpstra, W.I., Schouten, S., Bamford, D.H., Prangishvili, D., and Krupovic, M. (2015). Sulfolobus Spindle-Shaped Virus 1 Contains Glycosylated Capsid Proteins, a Cellular Chromatin Protein, and Host-Derived Lipids. J Virol *89*, 11681-11691. 10.1128/JVI.02270-15.

68. Wang, F., Cvirkaite-Krupovic, V., Kreutzberger, M.A.B., Su, Z., de Oliveira, G.A.P., Osinski, T., Sherman, N., DiMaio, F., Wall, J.S., Prangishvili, D., et al. (2019). An extensively glycosylated archaeal pilus survives extreme conditions. Nat Microbiol *4*, 1401-1410. 10.1038/s41564-019-0458-x.

69. Goulet, A., Blangy, S., Redder, P., Prangishvili, D., Felisberto-Rodrigues, C., Forterre, P., Campanacci, V., and Cambillau, C. (2009). Acidianus filamentous virus 1 coat proteins display a helical fold spanning the filamentous archaeal viruses lineage. Proc Natl Acad Sci U S A *106*, 21155-21160. 10.1073/pnas.0909893106.

70. Brennan, G., Stoian, A.M.M., Yu, H., Rahman, M.J., Banerjee, S., Stroup, J.N., Park, C., Tazi, L., and Rothenburg, S. (2023). Molecular Mechanisms of Poxvirus Evolution. mBio *14*, e0152622. 10.1128/mbio.01526-22.

71. Monzon, S., Varona, S., Negredo, A., Vidal-Freire, S., Patino-Galindo, J.A., Ferressini-Gerpe, N., Zaballos, A., Orviz, E., Ayerdi, O., Munoz-Gomez, A., et al. (2024). Monkeypox virus genomic accordion strategies. Nat Commun *15*, 3059. 10.1038/s41467-024-46949-7.

72. Pope, W.H., Jacobs-Sera, D., Russell, D.A., Peebles, C.L., Al-Atrache, Z., Alcoser, T.A., Alexander, L.M., Alfano, M.B., Alford, S.T., Amy, N.E., et al. (2011). Expanding the diversity of mycobacteriophages: insights into genome architecture and evolution. PLoS One *6*, e16329. 10.1371/journal.pone.0016329.

73. Krupovic, M., Dolja, V.V., and Koonin, E.V. (2019). Origin of viruses: primordial replicators recruiting capsids from hosts. Nat Rev Microbiol *17*, 449-458. 10.1038/s41579-019-0205-6.

74. Hurwitz, B.L., and U'Ren, J.M. (2016). Viral metabolic reprogramming in marine ecosystems. Curr Opin Microbiol *31*, 161-168. 10.1016/j.mib.2016.04.002.

75. Tian, F., Wainaina, J.M., Howard-Varona, C., Dominguez-Huerta, G., Bolduc, B., Gazitua, M.C., Smith, G., Gittrich, M.R., Zablocki, O., Cronin, D.R., et al. (2024). Prokaryotic-virus-encoded auxiliary metabolic genes throughout the global oceans. Microbiome *12*, 159. 10.1186/s40168-024-01876-z.

76. Kieft, K., Zhou, Z., Anderson, R.E., Buchan, A., Campbell, B.J., Hallam, S.J., Hess, M., Sullivan, M.B., Walsh, D.A., Roux, S., and Anantharaman, K. (2021). Ecology of inorganic sulfur auxiliary metabolism in widespread bacteriophages. Nat Commun *12*, 3503. 10.1038/s41467-021-23698-5.

77. Huang, X., Zhou, Z., Liu, H., Li, Y., Ge, T., Tang, X., He, Y., Ma, B., Xu, J., Anantharaman, K., and Li, Y. (2024). Soil nutrient conditions alter viral lifestyle strategy and potential function in phosphorous and nitrogen metabolisms. Soil Biology and Biochemistry *189*, 109279. https://doi.org/10.1016/j.soilbio.2023.109279.

78. Wang, L., Wang, Y., Huang, X., Ma, R., Li, J., Wang, F., Jiao, N., and Zhang, R. (2022). Potential metabolic and genetic interaction among viruses, methanogen and methanotrophic archaea, and their syntrophic partners.

ISME Communications *2*, 50. 10.1038/s43705-022-00135-2.

79. Zhou, Y., Gutiérrez-Preciado, A., Liu, Y., Moreira, D., Yakimov, M.M., López-García, P., and Krupovic, M. (2025). Nested parasitism in hypersaline environments: viruses and virus satellites of haloarchaea and their nanosized cellular symbionts. bioRxiv, 2025.2002.2015.638363. 10.1101/2025.02.15.638363.

80. Colman, D.R., Keller, L.M., Arteaga-Pozo, E., Andrade-Barahona, E., St Clair, B., Shoemaker, A., Cox, A., and Boyd, E.S. (2024). Covariation of hot spring geochemistry with microbial genomic diversity, function, and evolution. Nat Commun *15*, 7506. 10.1038/s41467-024-51841-5.

81. Yin, Z., Ye, L., and Jing, C. (2022). Genome-Resolved Metagenomics and Metatranscriptomics Reveal that Aquificae Dominates Arsenate Reduction in Tengchong Geothermal Springs. Environ Sci Technol *56*, 16473-16482. 10.1021/acs.est.2c05764.

82. Yuan, L., and Ju, F. (2023). Potential Auxiliary Metabolic Capabilities and Activities Reveal Biochemical Impacts of Viruses in Municipal Wastewater Treatment Plants. Environ Sci Technol *57*, 5485-5498. 10.1021/acs.est.2c07800.

83. Turunen, M., Olsson, J., and Dallner, G. (2004). Metabolism and function of coenzyme Q. Biochim Biophys Acta *1660*, 171-199. 10.1016/j.bbamem.2003.11.012.

84. Bekker, M., Alexeeva, S., Laan, W., Sawers, G., Teixeira de Mattos, J., and Hellingwerf, K. (2010). The ArcBA two-component system of Escherichia coli is regulated by the redox state of both the ubiquinone and the menaquinone pool. J Bacteriol *192*, 746-754. 10.1128/JB.01156-09.

85. Sarewicz, M., and Osyczka, A. (2015). Electronic connection between the quinone and cytochrome C redox pools and its role in regulation of mitochondrial electron transport and redox signaling. Physiol Rev *95*, 219-243. 10.1152/physrev.00006.2014.

86. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2022). The logic of virus evolution. Cell Host Microbe *30*, 917-929. 10.1016/j.chom.2022.06.008.

87. Koonin, E.V., Dolja, V.V., Krupovic, M., and Kuhn, J.H. (2021). Viruses Defined by the Position of the Virosphere within the Replicator Space. Microbiol Mol Biol Rev *85*, e0019320. 10.1128/MMBR.00193-20.

88. Alvarez, L.M., and Peng, X. (2025). Redefining paradigms in the archaeal virus-host arms race. Preprint. 10.1101/2025.04.20.649705.

89. Lopez-Beltran, A., Botelho, J., and Iranzo, J. (2024). Dynamics of CRISPR-mediated virus-host interactions in the human gut microbiome. ISME J *18*. 10.1093/ismejo/wrae134.

90. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2021). The healthy human virome: from virus-host symbiosis to disease. Curr Opin Virol *47*, 86-94. 10.1016/j.coviro.2021.02.002.

91. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2015). Origins and evolution of viruses of eukaryotes: The ultimate modularity. Virology *479-480*, 2-25. 10.1016/j.virol.2015.02.039.

92. Skennerton, C.T., Imelfort, M., and Tyson, G.W. (2013). Crass: identification and reconstruction of CRISPR from unassembled metagenomic data. Nucleic Acids Res *41*, e105. 10.1093/nar/gkt183.

93. Bland, C., Ramsey, T.L., Sabree, F., Lowe, M., Brown, K., Kyrpides, N.C., and Hugenholtz, P. (2007). CRISPR recognition tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. BMC Bioinformatics *8*, 209. 10.1186/1471-2105-8-209.

94. Steinegger, M., and Soding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. Nat Biotechnol *35*, 1026-1028. 10.1038/nbt.3988.

95. Padilha, V.A., Alkhnbashi, O.S., Shah, S.A., de Carvalho, A., and Backofen, R. (2020). CRISPRcasIdentifier: Machine learning for accurate identification and classification of CRISPR-Cas systems. Gigascience *9*. 10.1093/gigascience/giaa062.

96. Camargo, A.P., Nayfach, S., Chen, I.A., Palaniappan, K., Ratner, A., Chu, K., Ritter, S.J., Reddy, T.B.K.,

Mukherjee, S., Schulz, F., et al. (2023). IMG/VR v4: an expanded database of uncultivated virus genomes within a framework of extensive functional, taxonomic, and ecological metadata. Nucleic Acids Res *51*, D733-D743. 10.1093/nar/gkac1037.

97. Ahlgren, N.A., Chen, Y., Needham, D.M., Parada, A.E., Sachdeva, R., Trinh, V., Chen, T., and Fuhrman, J.A. (2017). Genome and epigenome of a novel marine Thaumarchaeota strain suggest viral infection, phosphorothioation DNA modification and multiple restriction systems. Environ Microbiol *19*, 2434-2452. 10.1111/1462-2920.13768.

98. Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010). Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics *11*, 119. 10.1186/1471-2105-11-119.

99. Wu, Z., Liu, S., and Ni, J. (2024). Metagenomic characterization of viruses and mobile genetic elements associated with the DPANN archaeal superphylum. Nat Microbiol. 10.1038/s41564-024-01839-y.

100. Couvin, D., Bernheim, A., Toffano-Nioche, C., Touchon, M., Michalik, J., Neron, B., Rocha, E.P.C., Vergnaud, G., Gautheret, D., and Pourcel, C. (2018). CRISPRCasFinder, an update of CRISRFinder, includes a portable version, enhanced performance and integrates search for Cas proteins. Nucleic Acids Res *46*, W246-W251. 10.1093/nar/gky425.

101. Nayfach, S., Camargo, A.P., Schulz, F., Eloe-Fadrosh, E., Roux, S., and Kyrpides, N.C. (2021). CheckV assesses the quality and completeness of metagenome-assembled viral genomes. Nat Biotechnol *39*, 578-585. 10.1038/s41587-020-00774-7.

102. Ren, J., Song, K., Deng, C., Ahlgren, N.A., Fuhrman, J.A., Li, Y., Xie, X., Poplin, R., and Sun, F. (2020). Identifying viruses from metagenomic data using deep learning. Quant Biol *8*, 64-77. 10.1007/s40484-019-0187-4.

103. Camargo, A.P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain, P.S.G., Nayfach, S., and Kyrpides, N.C. (2023). Identification of mobile genetic elements with geNomad. Nat Biotechnol. 10.1038/s41587-023-01953-y.

104. Guo, J., Bolduc, B., Zayed, A.A., Varsani, A., Dominguez-Huerta, G., Delmont, T.O., Pratama, A.A., Gazitua, M.C., Vik, D., Sullivan, M.B., and Roux, S. (2021). VirSorter2: a multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses. Microbiome *9*, 37. 10.1186/s40168-020-00990-y.

105. Hegarty, B., Riddell, V.J., Bastien, E., Langenfeld, K., Lindback, M., Saini, J.S., Wing, A., Zhang, J., and Duhaime, M. (2024). Benchmarking informatics approaches for virus discovery: caution is needed when combining in silico identification methods. mSystems *9*, e0110523. 10.1128/msystems.01105-23.

106. Li, R., Wang, Y., Hu, H., Tan, Y., and Ma, Y. (2022). Metagenomic analysis reveals unexplored diversity of archaeal virome in the human gut. Nat Commun *13*, 7978. 10.1038/s41467-022-35735-y.

107. Chibani, C.M., Shah, S.A., Schmitz, R.A., and Nayfach, S. (2024). Inaccurate viral prediction leads to overestimated diversity of the archaeal virome in the human gut. Nat Commun *15*, 5976. 10.1038/s41467-024-49902-w.

108. Wang, Y., Li, R., and Ma, Y. (2024). Reply to: Inaccurate viral prediction leads to overestimated diversity of the archaeal virome in the human gut. Nat Commun *15*, 5977. 10.1038/s41467-024-49903-9.

109. Paez-Espino, D., Eloe-Fadrosh, E.A., Pavlopoulos, G.A., Thomas, A.D., Huntemann, M., Mikhailova, N., Rubin, E., Ivanova, N.N., and Kyrpides, N.C. (2016). Uncovering Earth's virome. Nature *536*, 425-430. 10.1038/nature19094.

110. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol *37*, 1530-1534. 10.1093/molbev/msaa015.

111. Parks, D.H., Rinke, C., Chuvochina, M., Chaumeil, P.A., Woodcroft, B.J., Evans, P.N., Hugenholtz, P., and

Tyson, G.W. (2017). Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life. Nat Microbiol *2*, 1533-1542. 10.1038/s41564-017-0012-7.

112. Alves, R.J.E., Minh, B.Q., Urich, T., von Haeseler, A., and Schleper, C. (2018). Unifying the global phylogeny and environmental distribution of ammonia-oxidising archaea based on amoA genes. Nat Commun *9*, 1517. 10.1038/s41467-018-03861-1.

113. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res *13*, 2498-2504. 10.1101/gr.1239303.

114. Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. Nucleic Acids Res *52*, W78-W82. 10.1093/nar/gkae268.

115. Zheng, K., Sun, J., Liang, Y., Kong, L., Paez-Espino, D., McMinn, A., and Wang, M. (2025). VITAP: a high precision tool for DNA and RNA viral classification based on meta-omic data. Nat Commun *16*, 2226. 10.1038/s41467-025-57500-7.

116. Pruitt, K.D., Tatusova, T., Brown, G.R., and Maglott, D.R. (2012). NCBI Reference Sequences (RefSeq): current status, new features and genome annotation policy. Nucleic Acids Res *40*, D130-135. 10.1093/nar/gkr1079.

117. von Meijenfeldt, F.A.B., Arkhipova, K., Cambuy, D.D., Coutinho, F.H., and Dutilh, B.E. (2019). Robust taxonomic classification of uncharted microbial sequences and bins with CAT and BAT. Genome Biol *20*, 217. 10.1186/s13059-019-1817-x.

118. Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J.H., Roux, S., Adriaenssens, E.M., Brister, J.R., Kropinski, A.M., Krupovic, M., Lavigne, R., et al. (2019). Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. Nat Biotechnol *37*, 632-639. 10.1038/s41587-019-0100-8.

119. Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., and Ogata, H. (2020). KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics *36*, 2251-2252. 10.1093/bioinformatics/btz859.

120. Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer, E.L.L., Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., et al. (2021). Pfam: The protein families database in 2021. Nucleic Acids Res *49*, D412-D419. 10.1093/nar/gkaa913.

121. Galperin, M.Y., Wolf, Y.I., Makarova, K.S., Vera Alvarez, R., Landsman, D., and Koonin, E.V. (2021). COG database update: focus on microbial diversity, model organisms, and widespread pathogens. Nucleic Acids Res *49*, D274-D281. 10.1093/nar/gkaa1018.

122. Zimmermann, L., Stephens, A., Nam, S.Z., Rau, D., Kubler, J., Lozajic, M., Gabler, F., Soding, J., Lupas, A.N., and Alva, V. (2018). A Completely Reimplemented MPI Bioinformatics Toolkit with a New HHpred Server at its Core. J Mol Biol *430*, 2237-2243. 10.1016/j.jmb.2017.12.007.

123. Shen, W., Sipos, B., and Zhao, L. (2024). SeqKit2: A Swiss army knife for sequence and alignment processing. Imeta *3*, e191. 10.1002/imt2.191.

124. Edgar, R.C. (2022). Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. Nat Commun *13*, 6968. 10.1038/s41467-022-34630-w.

125. Capella-Gutierrez, S., Silla-Martinez, J.M., and Gabaldon, T. (2009). trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics *25*, 1972-1973. 10.1093/bioinformatics/btp348.

126. Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S., Larget, B., Liu, L., Suchard, M.A., and Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol *61*, 539-542. 10.1093/sysbio/sys029.

127. Kim, W., Mirdita, M., Levy Karin, E., Gilchrist, C.L.M., Schweke, H., Soding, J., Levy, E.D., and Steinegger, M.

(2025). Rapid and sensitive protein complex alignment with Foldseek-Multimer. Nat Methods. 10.1038/s41592-025-02593-7.

128. Grazziotin, A.L., Koonin, E.V., and Kristensen, D.M. (2017). Prokaryotic Virus Orthologous Groups (pVOGs): a resource for comparative genomics and protein family annotation. Nucleic Acids Res *45*, D491-D498. 10.1093/nar/gkw975.

129. Wang, J., Chitsaz, F., Derbyshire, M.K., Gonzales, N.R., Gwadz, M., Lu, S., Marchler, G.H., Song, J.S., Thanki, N., Yamashita, R.A., et al. (2023). The conserved domain database in 2023. Nucleic Acids Res *51*, D384-D388. 10.1093/nar/gkac1096.

130. Edgar, R.C., and Tolstoy, I. (2024). Muscle-3D: scalable multiple protein structure alignment. bioRxiv, 2024.2010.2026.620413. 10.1101/2024.10.26.620413.

131. Mistry, J., Finn, R.D., Eddy, S.R., Bateman, A., and Punta, M. (2013). Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res *41*, e121. 10.1093/nar/gkt263.

132. Kim, S., Chen, J., Cheng, T., Gindulyte, A., He, J., He, S., Li, Q., Shoemaker, B.A., Thiessen, P.A., Yu, B., et al. (2025). PubChem 2025 update. Nucleic Acids Res *53*, D1516-D1525. 10.1093/nar/gkae1059.

133. Yan, Y., Zhang, D., Zhou, P., Li, B., and Huang, S.Y. (2017). HDOCK: a web server for protein-protein and protein-DNA/RNA docking based on a hybrid strategy. Nucleic Acids Res *45*, W365-W373. 10.1093/nar/gkx407.

134. Janson, G., and Paiardini, A. (2021). PyMod 3: a complete suite for structural bioinformatics in PyMOL. Bioinformatics *37*, 1471-1472. 10.1093/bioinformatics/btaa849.

Acknowledgements

We thank Jilu Han for his help with the installation and debugging of the software used in this study. We thank the support of the high-performance servers of the Center for High Performance Computing and System Simulation, Laoshan Laboratory (Qingdao), the High-Performance Biological Supercomputing Center at the Ocean University of China, the Marine Big Data Center of the Institute for Advanced Ocean Study of the Ocean University of China and the IEMB-1, a high-performance computing cluster operated by the Institute of Evolution and Marine Biodiversity. This study was supported by the National Natural Science Foundation of China (No. 42120104006, 42176111 and 42306111), the Ocean Negative Carbon Emissions (ONCE), 2024 Graduate Self-directed Research Project (2024ZZKY, 202461036) and the Fundamental Research Funds for the Central Universities (202172002, 201812002, 202461036 andrew McMinn).

Author contributions

Conceptualization: K.Z. and Y.L.; Analysis: K.Z. and H.Y.; Database: D.P.-E.; Writing manuscript: K.Z.; Review and editing: Y.L., A.M., F.W. and M.W.; Computational resource: D.P.-E., M.L-P., S.L., G.W.N., C.H. and M.W.; Funding acquisition and supervision: Y.L., A.M. and M.W.

Competing interests

The authors declare no competing interests for the manuscript "A global viromic survey reveals unprecedented viral diversity within ammonium-oxidizing archaea and related lineages".

Figure legends

Figure 1. The expanding diversity of *Nitrososphaeria*-associated viruses or mobile genetic elements (MGEs). a, Biogeographic distribution of *Nitrososphaeria*-associated viral or MGE genomes. In the map, nodes of different colors indicate their putative archaeal hosts: *Ca. Caldarchaeales*, ammonia-oxidizing archaea (AOA), deeply-branched AOA-related archaea (DB-AO-related), and deeply-branched *Nitrososphaeria* archaea (D-B *Nitrososphaeria*). Sequences generated or reported in previous studies are indicated alongside the corresponding nodes. b, The number of vOTUs or MGE OTUs with distinct putative morphologies were originated from different habitats and were associated with diverse *Nitrososphaeria* lineages. Bar colours represent different archaeal lineages. c, Phylogenetic tree of *Nitrososphaeria* genomes. This tree was constructed based on concatenated alignments of 46 ribosomal proteins (totalling 5,963 amino acid positions) using the maximum likelihood method with the LG+C60+F model as recommended by Spang et al. and Alves et al. [2,112]. Reference genomes from *Thermoprotei* were used as the outgroup to root the tree. Tree annotation includes the habitats and lineages of these archaeal genomes. Virion icons to the right of the tree indicate the numbers of viruses or MGEs with different possible morphologies (at the OTU level) associated with each archaeal lineage.

Figure 2. The similarity of *Nitrososphaeria*-associated viruses or mobile genetic elements (MGEs) based on their genome contents. a, Genome-content similarity network of *Nitrososphaeria*-associated viruses or MGEs within the context of the global virosphere (sequences from NCBI RefSeq v. 223). For clarity, only connected component involving *Nitrososphaeria*-associated viruses or MGEs are shown. Each node represents a viral or MGE contig and is coloured according to its host lineage. *Nitrososphaeria*-associated viruses or MGEs clustering with established viral families are annotated adjacent to their respective groups. The pie charts illustrate the number of *Nitrososphaeria*-specific viral clusters (VCs) generated from this network. The percentage of shared proteins between genomes is indicated by the thickness of the edges connecting the nodes. The shapes of the nodes denote the potential morphologies of these viruses or MGEs. Each connected component is regarded as a super viral cluster (sVC), with asterisks indicating sVCs composed exclusively of *Nitrososphaeria*-associated viruses or MGEs.

Figure 3. Assignments of new families within *Nitrososphaeria*-associated viruses or mobile genetic elements

(MGEs). Virus proteomic hierarchy clustering tree used for the delineation of viral families. A distance threshold of 0.05 and a minimum of six shared proteins among members were used as the criteria for defining most families. *Charonviridae* were delineated by splitting from the *Kyanoviridae* clade, guided by the most recent common ancestor node. The names of the proposed families are indicated on the left side of the tree, with different virion icons representing the possible morphologies of these families. Clades corresponding to newly established viral families are highlighted on the tree with bold red branches. Tree annotation denotes host lineage and virion morphology. An asterisk is added before the names of viral families proposed in this study.

Figure 4. Genome organization diagrams of nine newly established viral families associated with newly expanded archaeal lineages. Arrows denote individual open reading frames (ORFs), with the colours indicating the function categories. Primary metabolism includes amino acid metabolism, carbohydrate metabolism, energy metabolism, lipid metabolism and nucleotide metabolism; secondary and specialized metabolism includes biosynthesis of other secondary metabolites, metabolism of terpenoids and polyketides, xenobiotics biodegradation and metabolism, metabolism of cofactors and vitamins and glycan biosynthesis and metabolism; genetic information processing includes replication and repair, transcription, translation, folding/sorting/degradation, information processing in viruses; cellular processes and community includes cell growth/death and cellular community. Relevant functional annotations are indicated above the corresponding ORFs. ORFs associated with viral core functions are highlighted with bold outlines. The corresponding archaeal lineages and their morphologies are indicated alongside each genome.

Figure 5. Evolution of *Nitrososphaeria*-associated rod-shaped viruses based on SIRV2-like capsomers through Bayesian inference (BI), Maximum Likelihood (ML) inference and structural modelling. a, BI tree based on the SIRV2-like capsid protein (domain accession in Pfam: PF12193). DUF732 (domain accession in Pfam: PF05305 and in VOG223: VOG14611) was used as the outgroup to root the tree. Posterior probabilities from BI are indicated on the tree branches with coloured nodes, alongside ML bootstrap values (MLB) with Blosum62+F+I+G4 as the best-fit substitutional model. The tree annotation indicates corresponding host archaeal lineages. *Nitrososphaeria*-associated rod-shaped viruses are shown as bold red branches. The names of the corresponding viral genera or families are labelled alongside the respective branches. BI convergence indices, including minimum effective sample size (minESS), potential scale reduction factor (PSRF), average standard deviation of split frequencies (ASDSF) and

max standard deviation of split frequencies (MSDSF). b, Structural modelling of archaeal rod-shaped viral capsomers and their structure-based similarity network. Nodes represent viral capsomers and are color-coded according to their viral families. Edge width indicates the degree of structural similarity between capsids (only identities $\geq$ 0.3 are shown). Representative structures from each cluster are displayed below the networks. c, Structural modelling of proteins encoded by open reading frames (ORFs) adjacent to SIRV2-like major capsid protein (MCP) genes in *Ahmunviridae*, *Demeterviridae*, and *Gaiaviridae*. Blue boxes indicate putative structural ORFs, while orange boxes represent ORFs with other functions. Functional annotations are shown above the boxes. An asterisk is added before the names of viral families proposed in this study.

Figure 6. Evolution of *Nitrososphaeria*-associated rod-shaped viruses based on HK97-like major capsid proteins (MCPs) through Bayesian inference (BI), Maximum Likelihood (ML) inference and structural modelling. a, BI tree based on the HK97-like MCP sequences, rooted using archaeal dodecin (domain accession in Pfam: PF07311) as the outgroup. Because HK97-like MCPs originate from multiple distinct domains and are difficult to align accurately at the sequence level, phylogenetic inference was performed using structure-guided multiple sequence alignments (MSAs) (see Methods). Posterior probabilities from BI are indicated on the tree branches with coloured nodes, alongside ML bootstrap values (MLB) with LG+F+R5 as the best-fit substitutional model. The tree annotation indicates corresponding host archaeal lineages. *Nitrososphaeria*-associated rod-shaped viruses are shown as bold red branches. The names of the corresponding viral genera or families are labelled alongside the respective branches. BI convergence indices, including minimum effective sample size (minESS), potential scale reduction factor (PSRF), average standard deviation of split frequencies (ASDSF) and max standard deviation of split frequencies (MSDSF). b, Structural modelling of archaeal head-tailed viral HK97-like MCPs and their structure-based similarity network. Nodes represent viral HK97-like MCPs and are color-coded according to their family-level taxonomic assignments: nodes with pink indicate members of newly proposed viral families in this study, nodes with grey represent members of previously established viral families, and nodes with light grey denote members lacking family-level taxonomic assignments. Each community is labeled with the corresponding viral family name above it (with the suffix "-*viridae*" omitted). Viral families newly proposed in this study are indicated by red labels with an asterisk prefix.

Figure 7. *Nitrososphaeria*-associated rod-shaped viruses encode unique auxiliary genes and exhibit a distinctive landscape of horizontal gene transfer (HGT) with their hosts. a, Maximum-likelihood (ML)

phylogenetic tree of the *Demeterviridae*-encoded UbiG-like protein, with Blosum62+R10 as the best-fit model. The tree annotation includes proteins of cellular origin (from InterPro), derived from archaeal or bacterial genomes. For clarity, only clades or branches related to viruses were retained, while the others were pruned. Representative structures and their per-residue measures of local confidence (pLDDT) are displayed below the phylogenetic tree. Molecular docking simulations showed that the benzene ring of 2,6-dihydroxybenzoic acid is inserted into the Rossmann-like fold pocket of UbiG encoded by vOTU_0592. b, ML phylogenetic tree of the *Demeterviridae*-encoded ArsR-like transcriptional regulator, with LG+F+R10 as the best-fit model. Representative structures and their per-residue measures of local confidence (pLDDT) are displayed below the phylogenetic tree. Molecular docking simulations showed the binding of A-DNA to the ArsR dimer encoded by vOTU_0592. c, Proportion of HGT events with different transfer directions among various *Nitrososphaeria* lineages. Different colours of the sectors indicate the direction of HGT, and the numbers within them represent the number of observations. d, The number of HGTs with distinct directions were associated from different functional catagories and were linked with diverse *Nitrososphaeria* lineages. Bar colours represent the direction of HGT and *Nitrososphaeria* lineages, respectively. The numbers of observations were indicated beside bars.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

a

Krupovic et. al., 2019 (Genomics)

Lee et. al., 2023 (metagenomics)

Krupovic et. al., 2011 (Genomics)

Lee et. al., 2024 (metagenomics)

Krupovic et. al., 2019 (Genomics)

Krupovic et. al., 2019 (Genomics)

Zhou et. al., 2022 (metagenomics)

Kim et. al., 2019 (isolate)

Ahlgren et. al., 2019 (metagenomcis)

Lopez-Perez et. al., 2019 (metagenomics)

Krupovic et. al., 2019 (Genomcis)

Krupovic et. al., 2019 (genomics)

● *Ca. Caldarchaeales*   ● AOA   ● D-B AOA-related   ● D-B *Nitrososphaeria*

b

Head-tailed (551)

Rod-shaped (51)

Spindle-shaped (16)

Icosahedral (15)

D-B *Nitrososphaeria* (9)

pSL4-affiliation (4)

Ca. Caldarchaeales (259)

D-B AOA-related (51)

MBG-A (15)

Nitrosocaldales (12)

MT/GW (63)

Nitrososphaerales (116)

Nitrosotalea (12)

Nitrosopumilales (169)

Geothermal spring (233)

Hydrothermal vents (76)

Seawater (225)

Freshwater (22)

Mine wastwater (2)

Soil (142)

c

Thermoprotei

Habitats Lineage Morphology

D-B *Nitrososphaeria*   6   1

pSL4-affiliation   3

Ca. Caldarchaeales   169   44

11   6

Non-AO *Nitrososphaeria*

D-B AOA-related   45   5

MBG-A (Group I.1c)   15

Tree scale: 0.2

AOA

Nitrosocaldales (HWCGIII, NC)   11   1

MT/GW   61

Nitrososphaerales (Group I.1b, NS)   89   6

Nitrosotalea (SAGMCG-1, NT)   10

Nitrosopumilales (Group I.1a, NP)   154   5

Morphology

⬡ Head-tailed

▭ Rod-shaped

⬡ Spindle-shaped

⬡ Icosahedral

Boostraps

● ≥ 90

● ≥ 70

Habitats

Hydrothermal vents

Marine

Geothermal springs

Soil

Freshwater

Cyanobacteria T7-like viruses

Cyanobacteria T4-like viruses

Pseudomonadota T7-like viruses

Pseudomonadota T4-like viruses

*002   *003   *004   *005

*006   007   008   009   010   *011   *012

Bacillota T4-like viruses

Bacteroidota T4-like viruses

ANME-1 Spindle-shaped viruses

ANME-1 head-tailed viruses

013   *014   *015   *016   *017   *018   *019   020   *021   *022 *023 *024 *025 *026

Myxococcota T4-like viruses

*027 *028 *029 *030 *031 *032 *033 *034   035   *036 *037   *038 *039 *040 *041 *042 *043 *044 *(045-047)

ANME-1 rod-shaped viruses

ANME-1 icosahedral viruses

*(048-061)

*(063-067)

60   7

Nitrososphaeria-specific clusters

Non-specific clusters

Host lineages

● AOA
● Ca. Caldarchaeales
● Other Nitrososphaeria
● ANME-1
● Pseudomonadota
● Cyanobacteriota
● Other Hosts

Morphology

○ Head-tail
▱ Filamentous
⬡ Icosahedral
△ Spindle

Shared protein clusters (%)

≥ 0.97
≥ 0.8
≥ 0.6
≥ 0.4
≥ 0.2

Host

0.05	

Itzamnaviridae (sVC_009)

Hypnosviridae (sVC_017)

Ekchuahviridae (sVC_010)

Chaacviridae (sVC_035)

Ahmunviridae (sVC_035)
Gaiaviridae (sVC_031)

Demeterviridae (sVC_004)

Peleviridae (sVC_012)
Freyrviridae (sVC_023)
Nyxviridae (sVC_043)
Erebusviridae (sVC_005)
Morosviridae (sVC_005)

Yangangviridae (sVC_015)

Alboranviridae (sVC_003)

Baleariviridae (sVC_003)

Jordviridae (sVC_002)

Thanatosviridae (sVC_020)

Apateviridae (sVC_001)
Chakrabarty-like (sVC_001)
Charonviridae (sVC_001)
Emdodeca-like (sVC_001)
Kyanoviridae (sVC_001)

Host lineage

- *Nitrosopumilales*
- *Nitrososphaerales*
- MT/GW
- *Nitrosocaldales*
- MBG-A
- D-B AOA-related
- *Ca. Caldarchaeales*
- D-B *Nitrososphaeria*

Morphologies

- Head-tailed
- Icosahedral
- Rod-shaped
- Spindle-shaped

Functions

Viral core function
- Viral structure
- Virus–host recognition
- Viral packaging
- Viral lysis

Other potential cellular functions
- Metabolism & Biosynthesis
- Genetic Information Processing
- Signal Transduction & Interaction
- Cell Structure, Transport & Motility
- Cell Growth, Death & Community

a

Tree scale: 0.5

HK97-fold

BI convergence indicies:
minESS: 738
PSRF: 1
ASDSF: 0.02
MSDSF: 0.19

Posterior probability
● ≥ 90
● ≥ 70

Host lineage
- Nitrosopumilales
- Nitrososphaerales
- MT/GW
- Nitrosocaldales
- MBG-A
- D-B AOA-related
- Ca. Caldarchaeales
- D-B Nitrososphaeria
- Methanophagales
- Pelagibacterales
- Nitrosomonadales

MLB:97
MLB:98
MLB:100
*Morosviridae
MLB:96
Fuxiviridae
MLB:61
*Peleviridae
*Nyxviridae
*Freyrviridae/*Jordviridae
MLB:87
Ekchuahviridae
MLB:89
MLB:90
*Hypnosviridae
*Thanatosviridae
MLB:100
Peduoviridae
MLB:100
*Alboranviridae/*Baleariviridae
MLB:100
Peduoviridae
MLB:99
*Erebusviridae
Yangangviridae
Peduoviridae
MLB:100
*Charonviridae
*Apateviridae
Emdodeca-like
Peduoviridae
Kyanoviridae

Host

b

*Charon./*Apate./Kyano./Peduo.A/Emdodeca-like

*Erebus./Yangang.

*Alboran./*Baleari

*Thanatos./Peduo.B

*Moros.

Ekchuah.A

*Jord./*Freyr./*Pele/*Nyx

Fuxi.

Ekchuah.B

*Hypnos.

Peduo.C

Identity
≥ 90
≥ 70
≥ 50
≥ 30

vOTU_0346
*Charonviridae

vOTU_0675
*Erebusviridae

vOTU_0755
Yangangviridae

vOTU_0759
*Alboranviridae

vOTU_0816
*Thanatosviridae

vOTU_0542
*Morosviridae

vOTU_0188
*Jordviridae

vOTU_0327
(35183..36037)
Ekchuahviridae (A)

vOTU_0327
(52541..53527)
Ekchuahviridae (B)

vOTU_0512
*Hypnosviridae

a

Bacteria
Archaea

Tree scale: 1 ⊢——⊣   Boostraps ≥ 90: ●

vOTU_0592
Demeterviridae
(pLDDT=94.7)

Affinity = -6.19 kca/mol

Thermoprotei_A0A662UAW1
Thermoprotei archaeon
(pLDDT=94.6)

Saprospiraceae_A0A521URU1
Saprospiraceae bacterium
(pLDDT=94.6)

b

Bacteria
Archaea

Fuselloviridae   *Kirjokansivirales*

Demeterviridae

Tree scale: 1 ⊢——⊣   Boostraps ≥ 90: ●

vOTU_0592
Demeterviridae
(pLDDT=92.3)

dimer
(pLDDT=86.8)

confidence score = 0.85

HTV27
Hafunaviridae
(residue 1-60, pLDDT=80.9)

SSV5
Fuselloviridae
(pLDDT=79.4)

phiCDIF1296T
unclassified *Caudoviricetes*
(pLDDT=79.8)

c

AOA

Nitrosopumilales — 79 / 375 / 982
Nitrososphaerales — 15 / 161 / 377
Nitrosotalea — 2
Nitrosocaldales — 8

Non-AO *Nitrososphaeria*

Ca. Caldarchaeales — 31 / 110 / 183
D-B AOA-related — 21 / 10 / 152
pSL4-affiliation — 3

● Bidirection   ● Archaeal donor   ● Viral donor

d

Carbohydrate metabolism 384
Replication and repair 570
Glycan biosynthesis and metabolism 565
viral structure 107
Metabolism of cofactors and vitamins 148
Unknown function 202
Nucleotide metabolism 190

v-?-a 1,323
v-from-a 589
v-to-a 95

Nitrosopumilales 501
Nitrosopumilales 1,262
Ca. Caldarchaeales 221
D-B AOA-related 173
pSL4-affiliation 3
Nitrosocaldales 4
Nitrosotalea 2