Viruses associated with *Nitrososphaerota* archaea: diversity, evolution and horizontal gene transfers

Kaiyang Zheng[1], Yantao Liang[1,2,3,4,5,6,7*], David Paez-Espino[8], Andrew McMinn[1,4,7,9*], Min Wang[1,2,3,4,5,6,7,10*]

[1] College of Marine Life Sciences, Ocean University of China, Qingdao, China.

[2] Haide College, Ocean University of China, Qingdao, China.

[3] MoE Key Laboratory of Evolution & Marine Biodiversity, Ocean University of China, Qingdao, China.

[4] Institute of Evolution and Marine Biodiversity, Ocean University of China, Qingdao, China.

[5] Frontiers Science Center for Deep Ocean Multispheres and Earth System, Ocean University of China, Qingdao, China.

[6] Center for Ocean Carbon Neutrality, Ocean University of China, Qingdao, China.

[7] UMT-OUC Joint Centre for Marine Studies, Qingdao, China.

[8] Ancilia Biosciences, Co. Ltd., Alexandria Launchlabs, New York, USA.

[9] Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia.

[10] The Affiliated Hospital of Qingdao University, Qingdao, China.

* Correspondence:

Yantao Liang, Email: liangyantao@ouc.edu.cn

Andrew McMinn, Email: andrew.mcminn@utas.edu.au

Min Wang, Email: mingwang@ouc.edu.cn

Abstract

Nitrososphaerota, an archaeal phylum mainly containing *Thaumarchaea* and *Aigarchaea*, is an ecological widespread archaeal lineage belonging to TACK super-phylum. However, viral diversity and evolution associated with this archaeal lineage, especially for extremophilic *Thaumarchaea* and *Aigarchaea*, remain uncharacterized. Here, we reveal the neglected diversity of the *Thaumarchaea* virome by identifying 152 species-level viral and mobile genetic element (MGE) sequences associated with extremophilic *Thaumarchaea*, which mainly colonize in hyperthermophilic environments, deep sea, and acidic soils. Importantly, we identified 263 previously unrecognized species-level viral/MGE sequences associated with *Aigarchaea*, including a novel rod-shaped viral family uniquely characterized by capsid heterodimers, thereby filling a long-standing gap in the viral diversity of the TACK super-phylum. Phylogenetic analysis revealed close evolutionary relationships between *Aigarchaea*-associated rod-shaped viruses and *Methanophagales* (ANME-1) rod-shaped viruses, suggesting that they may have evolved from a common rod-shaped viral ancestor. In contrast, *Aigarchaea*-associated head-tailed viruses formed three distinct monophyletic clades compared to other viruses, representing previously undiscovered evolutionary lineages within virosphere. Gene functional analysis identified 32 viral-encoded genes potentially involved in supplementing host metabolism, including, for the first time, genes that confer selective advantages to their extremophilic archaeal hosts, such as those involved in oxidative stress adaptation and antibiotic synthesis. Approximately 54% of 6,505 horizontal gene transfers (HGTs) occur bidirectionally between *Nitrososphaerota* and viruses/MGEs, highlight the significant role of viruses/MGEs in facilitating gene exchange within the archaeal communities. Collectively, this work sheds light on the previously unknown *Aigarchaea*-associated viromes, and unveils the underexplored diversity, evolution, functional repertoires, and HGTs of *Nitrososphearota*-associated viruses.

Introduction

The archaea phylum *Nitrososphaerota* within TACK super-phylum mainly contains *Thaumarchaea* and *Aigarchaea* [1,2]. In aquatic ecosystems, *Thaumarchaea* constitute up to 40% of microorganisms, contributing significantly to ammonia oxidation, thus heavily influencing nitrogen cycling [3-5]. *Thaumarchaea* including all ammonia-oxidizing archaea (AOA) [6], utilize ammonia oxidation pathways as their primary energy source [7,8]. Compared to ammonia-oxidizing bacteria, AOA thrive in low-ammonia conditions due to higher ammonium affinity, dominating nitrogen-limited environments as efficient "nitrogen thieves" [9-14]. Originating from terrestrial, non-ammonia-oxidizing ancestors, *Thaumarchaea* is thought to have adapted aerobically post-Great Oxygenation Event, becoming abundant in diverse environments and contributing substantially to carbon fixation and ammonia oxidation, even in extreme habitats [8,15-20]. As the sister clade of *Thaumarchaea*, *Aigarchaea* shares substantial genetic content involved in sulphur oxidation and amino oxidation, suggesting an ancestral ecological transition from thermal to moderate environments [21]. Primarily inhabiting hydrothermal vents and hot springs, *Aigarchaea* influence carbon and sulphur cycling, nutrient transformation, and microbial community structures, playing critical ecological roles in geothermal systems [22,23].

Viruses and mobile genetic elements (MGEs) significantly shape archaeal genomic and ecological landscapes, exhibiting high-level of morphological and genomic diversity relative to viruses from other life domains [24]. Due to cultivation challenges, environmental metagenomics has become the primary method for exploring archaeal viral diversity. Recent studies have greatly expanded viral diversity associated with the Asgard super-phylum, ANME-1, and DPANN super-phylum, addressing gaps in viral evolution and host-virus interactions [25-29]. Over 50 *Thaumarchaea*-associated viral/MGE sequences from marine samples have so far been identified, revealing diverse viral elements that influence ammonia oxidation

and host genomic integrity [30-36]. However, the limited number of case studies prevents

a comprehensive profiling of the genomic and functional diversity of these archaeal

viruses. In particular, the knowledge gap of viruses associated with extremophilic

Thaumarchaea lineages (e.g., *Nitrosocaldales* and *Nitrosotalea*) poses a significant

challenge for systematically resolving their phylogenetic placements within the

virosphere. Furthermore, as a phylogenetic sister clade closely related to

Thaumarchaea and one of the main lineages constituting the TACK super-phylum,

Aigarchaea-associated viruses/MEGs remain entirely underexplored to date,

representing a major enigma within virology.

To address this knowledge gap, we systematically explored *Thaumarchaea* and

Aigarchaea (collectively referred to as *Nitrososphaerota* in this study) associated

viral/MGE diversity using extensive global metagenomic datasets. Unlike previous

studies [30-34], our analysis of extensive global datasets enabled detailed taxonomy,

phylogeny, metabolic potential analyses, and host-virus interactions. Our research

established a genomic catalogue comprising 746 viral/MGE sequences (624 species-

level sequences) associated with *Nitrososphaerota*, spanning diverse extreme and

moderate ecosystems globally, and shedding light on the previously unknown

Aigarchaea-associated viromes.

Results

Overview of the genomic catalogue of *Nitrososphaerota*-associated viruses/MGEs

The mining of *Nitrososphaerota*-associated viruses/MGEs primarily utilized public
viral databases, *Nitrososphaerota* genomes, and metagenomic samples from
hydrothermal and deep-sea environments (Fig. 1a and Supplementary Note 1). A
comprehensive *Nitrososphaerota* genome database containing 2,137 genomes and
metagenome-assembled genomes (MAGs) was constructed from NCBI GenBank and
IMG/M databases [37,38]. Metagenomic datasets from extreme habitats, including
hyperthermal, abyssal/hadal ocean, hypersaline, frozen, and acidic environments,
were analyzed (Supplementary Dataset 1) [37,39-41]. Employing a hybrid identification
strategy integrating CRISPR spacer matches and protein-sharing analysis, a total of
746 *Nitrososphaerota*-associated viruses/MGEs were identified, including 712
extrachromosomal viruses/MGEs, and 34 integrated viruses/MGEs. These were
grouped into 624 species-level viral operative taxonomic units (vOTUs), comprising
153 vOTUs with known viral capsid proteins and 471 vOTUs lacking annotated
capsids (Fig. 1a, Supplementary Dataset 2). The pipeline prioritized specificity,
verifying 14 previously reported viruses/MGEs [31,32] (Fig. 1b), and newly identified
350 viral/MGE contigs (336 vOTUs) that were associated with *Nitrososphaerota*.
Most viruses/MGEs originated from geothermal, marine, soil, and hydrothermal
environments, aligning with *Thaumarchaea* and *Aigarchaea* habitats [7,19,22].

Identifying the host and habitat diversity of *Nitrososphaerota*-associated
viruses/MGEs

The 1,192 CRISPR-Cas cassettes were identified across 636 *Nitrososphaerota*
genomes (Supplementary Dataset 3) and these were used to describe associations
between *Nitrososphaerota* and viruses/MGEs. These cassettes were classified into
five distinct CRISPR types, whose spacer diversity reflected more extensive diversity

of *Nitrososphaerota* viromes. The spacer-sharing network has low viral coverage (only 336 viral/MGE sequences) (Supplementary Dataset 4), suggesting broader undiscovered viral diversity. Host-virus specificity varied among *Nitrososphaerota* genera: for example, spacers from *Caldarchaeum* demonstrated genus-level specificity, whereas other *Thaumarchaea* genera indicated more strict viral specificity (Fig. 2a). In contrast, spacers from *Aigarchaea* lineage, such as *Terraquivivens* and *Wolframiiraptoraceae*, showed broader sharing patterns, suggesting weaker genus-level specificity.

Nitrososphaerota-associated viruses/MGEs were retrieved from diverse ecosystems and linked to multiple archaeal lineages. They are widely distributed from tropical to polar regions and across terrestrial and marine environments (Fig. 2b). With regard to habitat, Group 1.1a and MT/GW (marine trench and ground water) - associated viruses/MGEs were mainly found in aquatic environments, while Group 1.1b-associated viruses/MGEs are predominantly from soil (Fig. 2c). Viruses/MGEs from geothermal and hydrothermal environments were primarily linked to *Caldarchaeales* and *Wolframiiraptoraceae*, aligning with their archaeal hosts' habitat preferences [21,42-45]. Of the host lineages, 746 of the identified viruses/MGEs spanned 11 *Nitrososphaerota* lineages (Fig. 2d and Supplementary Dataset 2) [2,20,21,23,46]. For the 352 *Thaumarchaea*-associated vOTUs, 287 linked to AOA, while 65 were associated with Non-AOA. These viruses/MGEs included 152 previously undescribed vOTUs associated with extremophilic *Thaumarchaea* lineages, comprising 63, 51, 15, 12, and 12 vOTUs related to MT/GW, Deeply-branching *Thaumarchaea* (D-B Thaum), Marine Beneath Group A (MBG-A), *Nitrosotalea*, and *Nitrosocaldales*, respectively. Of the 263 newly identified *Aigarchaea*-associated vOTUs, 210 linked to *Wolframiiraptoraceae*, 49 were linked to *Caldarchaeales*, and four were linked to pSL4-affiliation. With regard to morphology, head-tailed viruses, known as cosmopolitan prokaryotic viruses, are widespread across 11 *Nitrososphaerota* lineages (n = 447 vOTUs) (Fig. 2d and Supplementary Dataset 2), aligning with prior studies [25-27,30,32-34,47]. Additionally, 44 previously undescribed *Aigarchaea*-associated rod-shaped vOTUs were identified. These findings show that viruses/MGEs linked to nine

154 additional archaeal lineages in contrast to previous studies [30-36,48].

Genomic and phylogenetic diversity of viruses/MGEs associated with
Nitrososphaerota

Genome-content-based analysis was used to characterize relationships and diversity
among _Nitrososphaerota_-associated viruses/MGEs. Among these, 16 viral/MGE
genomes (six viral clusters) showed genome-wide linkages to other bacterial and
archaeal viruses (Supplementary Fig. 2 and Supplementary Dataset 5); ten genomes
shared significant genomic similarity (>20% shared protein clusters) with established
archaeal viral families (_Chaacviridae_, _Ekchuahviridae_, and _Itzamnaviridae_), which
typically infect methanogenic archaea with diverse morphologies [27,28]. However, most
Nitrososphaerota-associated viruses/MGEs (95.2%) exhibit unique genome content,
forming 119 independent viral clusters (Supplementary Fig. 2). These clusters belong
to 58 groups based on ≥20% protein-cluster sharing, including 31 _Thaumarchaea_-
specific and 25 _Aigarchaea_-specific groups (Fig. 3a). These reflect host lineage and
morphological specificity consistent with previous studies [27,47].

Twelve families encoding known capsid proteins were found to be associated
with _Nitrososphaerota_, including eight previously undescribed families. Of the
established viral families, three families (_Ekchuahviridae_, _Itzamnaviridae_,
Chaacviridae), initially identified from deep-sea hydrothermal vents linked to
ANME-1 archaea [28], were found to be also associated with _Aigarchaea_
(_Wolframiiraptoraceae_) (Supplementary Dataset 2 and Fig. 3b). Of the possible new
viral families, one rod-shaped family (_Byatisviridae_) and four head-tailed families
(_Fthagguaviridae_, _Sthaneeviridae_, _Ythogthaviridae_, and _Cthughaviridae_) were
established, which were associated with _Aigarchaea_ (_Wolframiiraptoraceae_ and
Caldarchaeales); two head-tailed families (_Ommogviridae_ and _Yebviridae_) were
established and these were associated with _Thaumarchaea_. Notably, _Ommogviridae_
was initially identified in a previous metagenomic study, characterized by their
nitrification genes (ammonia monooxygenase, _amoC_) [33]. Moreover, 18 MGE groups

which lacked capsid proteins were detected, forming independent groups in the genome-content-based network. These suggest that they have unique genomic features compared with other reference viral genomes (Supplementary Dataset 2, Supplementary Figs. 2 and 3).

This comprehensive genomic catalogue of *Nitrososphaerota*-associated viruses/MGEs enables systematic assessment of their phylogenetic placements using Bayesian (BI) and Maximum-Likelihood (ML) joint phylogenetic inference (Extended Data 3 and 4). For *Nitrososphaerota*-associated rod-shaped viruses (*Byatisviridae*), two genomes showed close relationships with ANME-1-associated *Ahmunviridae* viruses [28]. *Byatisviridae* members formed a distinct monophyletic sister clade to *Ahmunviridae*, suggesting divergence from a common ancestor (Fig. 4a). This is supported by the unique structural characteristics of their SIRV2-like fold capsid proteins (Supplementary Fig. 5a). Unlike *Rudiviridae* and *Ahmunviridae*, which share a common evolutionary origin, *Byatisviridae* appears to possess a capsid composed of heterodimers, implying that those heterodimeric capsids of *Aigarchaea*-associated rod-shaped viruses may have evolved from an ancestral homodimeric architecture (Fig. 4a, Supplementary Figs. 5b and 6). For head-tailed viruses, most *Aigarchaea*-associated viruses formed monophyletic clades in phylogenetic trees. Some *Thaumarchaea*-associated viruses are phylogenetically intertwined with other archaeal and bacterial viruses. They exhibit typical polyphyletic characteristics. *Fthagguaviridae* and *Cthughaviridae* are two *Aigarchaea*-associated viral families, forming two distinct monophyletic groups in the phylogenetic trees of capsid and terminase, respectively, which is further supported by capsid structural similarity networks (Figs. 4b, 4e, and Supplementary Fig. 7). This indicates that these viruses occupy entirely distinct evolutionary placements from other archaeal viruses and bacteriophages. *Ekchuahviridae* clustered within archaea-associated viruses (*Ahpuchviridae*, *Halomagnusviridae*, and *Haloferuviridae*), and is closely related to bacteriophages infecting *Actinomycetes* (Fig. 4c). *Yebviridae* likely shares a common ancestor with T4-like cyanophages (*Kyanoviridae*); and this is supported by capsid structural similarity networks (Figs. 4e, 4f, and Supplementary Fig. 7). Collectively,

an additional seven *Nitrososphaerota*-associated viral families are proposed and the

evolutionary placement of these archaeal viruses is elucidated.

Metabolic potential of *Nitrososphaerota*-associated viruses/MGEs

Due to the previously limited characterization of the diversity of *Nitrososphaerota*-

associated viruses/MGEs, our understanding of their metabolic capacities and

potential contributions to archaeal host metabolism remains largely incomplete. As

with other archaeal head-tailed viruses [26,27,47], these viruses/MGEs encode proteins

essential for core viral functions (DNA replication, transcription, packaging, and

structural components) and proteins counteracting host defences, including

restriction-modification systems and chromosome partitioning (Supplementary

Dataset 6, Supplementary Figs. 3 and 4). Notably, methyltransferases were

widespread (170 genomes, 140 vOTUs), occasionally coupled with endonucleases.

The presence of *parB* genes in 71 genomes (36 vOTUs) suggests the likelihood of

plasmid-like partitioning mechanisms in viral replication [49,50]. The newly identified

Aigarchaea-associated rod-shaped viral family, *Byatisviridae*, was found to uniquely

encode demethyl-ubiquinol-6 methylase (Supplementary Dataset 6), implying

involvement of the host's ubiquinone metabolism, a metabolic potential not observed

in other rod-shaped archaeal viruses[27,28,51].

Some virus-encoded proteins may serve as an auxiliary metabolic potential

reservoir for the hosts. A total of 32 distinct viral-encoded protein families have no

detectable homologs in host archaeal panproteomes, suggesting potential host

metabolic compensation (Fig. 5 and 6a, Supplementary Figs. 8, Supplementary

Dataset 7). Specific examples include 6-phosphogluconolactonase (PPP pathway)

encoded by *Aigarchaea*-associated (*Wolframiiraptoraceae*) viruses/MGEs [21,23], and

sdmt (sarcosine/dimethylglycine N-methyltransferase) in *Aigarchaea*-associated

(*Caldarchaeales*) viruses/MGEs. The viral/MGE-encoded *sdmt* gene potentially aids

the osmotic and oxidative stress tolerance of *Caldarchaeales* cells through betaine

production (Fig. 5) [52,53]. The presence of glucose-6-phosphate isomerase (*pgi1*)

encoded by a Group 1.1b-associated viruses suggests a novel glycolytic capability of

AOA (Fig. 5), previously reported only in non-AOA *Thaumarchaea* [20]. Additionally,

the detection of Isopenicillin-N synthase (*pcbc*) gene indicates that viruses may

potentially confer the metabolic capacity for β-lactam antibiotic production to archaea

[54,55]. For Group 1.1a archaea, virus-encoded *calS13* (capsular polysaccharide

biosynthesis) implies possible roles in biofilm formation, expanding archaeal

capabilities in extracellular polymeric substance production [56].

Horizontal gene transfers landscape between *Nitrososphaerota*-associated viruses/MGEs and their archaeal hosts

Owing to the previously insufficient exploration of the diversity of

Nitrososphaerota-associated viruses/MGEs, our understanding of horizontal gene

transfer (HGT) dynamics between these elements and their archaeal hosts remains

fragmentary. This gap is particularly pronounced for *Aigarchaea*, for which virus–host

associations related to protein metabolism and HGT have not been characterized yet.

We explored virus-host gene exchange, identifying 6,505 HGT events from 527

phylogenetic trees, using the approach established by Irwin et al. (Extended Data 5) [57].

Approximately 54% of gene transfers occur bidirectionally between *Nitrososphaerota*

and viruses/MGEs, 45% are from *Nitrososphaerota* to viruses/MGEs, and only 0.8%

are from viruses/MGEs to *Nitrososphaerota* (Fig. 6b, Supplementary Dataset 8). This

highlights the significant role of viruses and MGEs in facilitating gene exchange

within archaeal communities. With regard to different viral/MGE families,

MGE_F003 might facilitate extensive transfer of glycosylation-related genes within

Group 1.1a (Fig. 6c and Supplementary Dataset 8); MGE_F001 transferred gene for

the recombination protein RecA extensively within clades of *Thaumarchaea*;

Fthagguaviridae was primarily involved in structural-protein-associated transfers

within *Aigarchaea*; *Ekchuahviridae* and *Yangangviridae* mediated glycosyltransferase

and methyltransferase exchanges (Supplementary Dataset 8), respectively. This result

indicates that different viral families exhibit distinct preferences in the types of

metabolic functions involved in these HGT events.

HGT events mediated by these viruses/MGEs have the potential to indirectly

influence elemental cycling, prominently phosphorus (60%) and nitrogen cycles via

queC and *queD* genes (folate biosynthesis), encoding enzymes producing

orthophosphate, triphosphate, and ammonia-derived reactions (Fig. 6d and

Supplementary Dataset 8) [58,59]. Cobaltochelatase transfer events suggest viral roles in

cobalt metabolism, especially between Group 1.1a and *Yebviridae*. Further, HGTs

involving in carbon fixation, urea cycle, TCA cycle, purine synthesis, and DNA

synthesis highlight the viral contribution to carbon cycling. Additional transfers

related to sulphur, manganese, halogens, and copper metabolism were identified (Fig.

6d and Supplementary Dataset 8), suggesting that archaeal viruses/MGEs may

indirectly participate in extensive elemental cycling.

Discussion

These results significantly expand the current understanding of *Nitrososphaerota*-

associated viruses/MGEs, broadening the scope from previously reported

Thaumarchaea-associated viruses, and identified 263 previously unrecognized

Aigarchaea-associated viruses for the first time, expanding the known number of

Nitrososphaerota-associated viruses/MGEs by 10-times [30-34]. The mining of

Aigarchaea-associated viruses/MGEs viral genomes fills a critical knowledge gap of

archaeal viromes. Based on this catalogue, diverse *Nitrososphaerota*-associated viral

families were identified and their host associations, genomic features, phylogenetic

traits, functional potentials, and HGT landscape were described.

The phylogenetic analysis of *Nitrososphaerota*-associated viruses showed their

evolutionary relationships with other viruses. In the past, only the capsids of

Lipothrixviridae and *Tristromaviridae* were known to be composed of heterodimers[51].

Here, it was found that some *Aigarchaea*-associated rod-shaped viral capsids also

consist of heterodimers (Fig. 4a, Supplementary Figs. 5 and 6), thus updating our

understanding of rod-shaped archaeal viral evolution. With respect to capsid assembly

strategy, homodimers may provide symmetry and repetitiveness, thereby contributing

to the maintenance of capsid stability [60,61]. Heterodimers may afford greater structural

flexibility and diversity, enabling adaptation to varying environmental stresses and

host defence mechanisms [62,63]. The presence of both homodimers and heterodimers in

Byatisviridae suggests that these viruses are undergoing capsid diversification in

response to environmental stress. Some *Aigarchaea*-associated viruses have close

relationships with ANME-1 (*Methanobacteriota*)-associated viruses/MGEs

originating from hydrothermal vents (e.g. Methanophagales virus PBV300,

Ahmunviridae) (Supplementary Fig. 2)[28], implying viral host-switching or non-

specific transfers occurred between both archaeal lineages, which is common in the

evolution of viruses [64-66].

Viruses have been found to encode diverse metabolic genes that manipulate host

metabolism, thereby creating favorable conditions for completing their life cycles and

indirectly contributing to global biogeochemical cycles [67-70]. In archaea-associated

viruses, such as those infecting methanogenic archaea, Asgard superphylum,

halophilic archaea, viral-encoded metabolic genes have been identified that

potentially modulate archaeal methane-oxidization, sulfur metabolism, nucleotide and

amino acid metabolisms [25,71,72]. Regarding *Thaumarchaea*-associated viruses/MGEs,

previous studies have discovered that viruses may manipulate host ammonia oxidation

metabolism by carrying the *amoC* gene and participate in host vitamin synthesis [32,33].

Through the systematic exploration of *Nitrososphaerota*-associated viruses, we have

not only expanded the functional diversity of metabolic genes encoded by

Thaumarchaea-associated viruses/MGEs but also, more importantly, filled the gap in

the functional diversity of *Aigarchaea*-associated viruses/MGEs. Metagenomic

investigations of geothermal springs have revealed that *Aigarchaea* may possess the

pentose phosphate pathway (PPP) but lack two core enzymes [21]: glucose-6-phosphate

dehydrogenase and 6-phosphogluconolactonase. The 6-phosphogluconolactonase

encoded by *Aigarchaea*-associated viruses/MGEs may compensate for the missing

steps in the host's PPP (Fig. 5, Supplementary Figs. 8, Supplementary Dataset 6),

thereby supporting redox balance and cellular resilience. Additionally, we have

identified, for the first time, the presence of the *sdmt* gene in viruses, suggesting that viruses may assist hosts in synthesizing betaine through the glycine betaine methylation pathway, helping them cope with high osmotic environments [73,74]. As a type of methyltransferase, *sdmt* may also function similarly to other viral-encoded methyltransferases, playing a role in immune evasion during viral-host interactions [75-77]. In this context, this represents a typical example of the pervasive exaptation of host proteins by viruses [65]. Beyond encoding genes that potentially aid host environmental adaptation, archaeal viruses also carry antibiotic resistance genes (Fig. 5, Supplementary Figs. 8, Supplementary Dataset 6). Archaea are generally unaffected by β-lactam antibiotics, and previous studies have demonstrated that archaea can produce antibiotics [78]; however, no research has yet indicated that archaea are capable of producing β-lactam antibiotics. The presence of *pcbc* genes in related viruses may help archaeal hosts gain an advantage in interspecies competition with bacteria through production of β-lactam antibiotics, thereby providing more opportunities for viruses to complete their life cycles. In summary, the significantly expanded diversity of *Nitrososphaerota*-associated viruses/MGEs revealed in this study provides prior knowledge for future in-depth research into virus-archaea interactions.

Methods

Assembly of *Nitrososphaerota*-associated CRISPR database

The *Nitrososphaerota* phylum comprises two major clades: *Thaumarchaea* and its

sister clade, *Aigarchaea*. To produce a comprehensive genomic dataset, reference

genomes were retrieved from GenBank and IMG/M [37,38], encompassing a total of

2,137 genomic assemblages. Specifically, 1,830 assemblages (comprising 322,784

genomic contigs) were obtained from GenBank; 307 assemblages (containing 23,395

genomic contigs) were sourced from IMG/M.

To identify CRISPR spacers within these genomes, similar to previous studies

[25,26], CRASS v.0.3.8 and MinCED v.0.4.2 (CRISPR recognition tool, CRT) were

employed using the following parameters to detect spacers and discard those with

low-complexity and short length [79,80]: word size = 7, spacer length = 26, and a

minimum of three directed repeats (DRs). This step resulted in the identification of

37,485 *Nitrososphaerota*-associated spacers, which were subsequently used for the

detection of viruses/MGEs and downstream analyses.

The metagenome-derived spacers that are associated with *Nitrososphaerota*

were detected to further expand the diversity of these CRISPR spacers. First, a total of

153 metagenome-derived assemblages (including 58 from the Malaspina expedition)

from diverse ecosystems were retrieved and downloaded from IMG/M and ENA

(European Nucleotide Archive) (Supplementary Dataset 1)[38,41]. These metagenomes

were from 21 different studies, focusing on extreme natural habitats and including

92,037,046 assembled contigs. These include four studies from deep-sea, three two

from sea ice, four studies from a hypersaline environment, and one study from an

acidic environment. Second, the CRISPR spacers found in these contigs were

identified using the same approach described above, producing 1,828,641 spacers and

1,918,441 DRs. Third, all DRs were aligned to *Nitrososphaerota* genomes using

MMseqs2 with parameters [81]: word size = 7, pairwise alignment identity = 1,

percentage coverage of DRs = 1, mismatches = 1. This step produced 196,551 spacers

derived from 153 metagenomes. Finally, this process generated a *Nitrososphaerota*-associated spacer database containing 234,036 (Fig. 1 and Extended Data 1).

As in previous studies [26], CRISPR-Cas cassette prediction and subtyping for archaeal contigs and genomes were conducted using CRISPRcasIdentifier v1.0.0 [82], employing a combination of Classification and Regression Trees (CART), Support Vector Machine (SVM), and Extremely Randomized Trees (ERT) algorithms for classification and regression. The analysis was performed with the following parameters: –mode ERT SVM CART, -hmm HMM1 HMM2 HMM3 HMM4 HMM5. Subtype assignment was determined through a consensus-based approach, where only subtype classifications consistently predicted by all three models (CART, SVM, and ERT) were accepted. The final subtype was selected based on the highest mean probability and the lowest standard deviation among the predictions.

Mining and filtering *Nitrososphaerota*-associated viruses/MGEs from IMG/VR (v.4)

The IMG/VR (v.4) database, contigs from 153 metagenomes, and archaeal genomes served as the primary resources for mining *Nitrososphaerota*-associated viral/MGE genomes. At first, the archaeal viral/MGE mining pipeline was validated using CRISPR spacers derived from archaeal genomes and host predictions from the IMG/VR (v.4) database and other studies to minimize false positives [31-34,36,83,84]. The 37,485 CRISPR spacers from *Nitrososphaerota* genomes were aligned to IMG/VR (v.4) database using MMseqs2 with the parameters: word size = 7, pairwise alignment identity = 0.9, percentage coverage of spacers = 0.9, mismatches = 1, resulting in 368 targeted. Among these results, the hosts of 46 viruses/MGEs were also predicted to be *Nitrososphaerota* in IMG/VR (v.4). Due to some viral contigs being assigned to non-*Nitrososphaerota* hosts in IMG/VR (v.4) metadata, the genome-content similarity between viruses/MGEs and hosts was applied to reduce the false positives: i) the open reading frames (ORFs) of these viral contigs were predicted by Prodigal v.2.6.3 and aligned to the NR database (2024.05.20) with 1e-5 as the E-value cut-off [85]; ii) the

Nitrososphaerota-related hits were counted and normalized based on viral contig

lengths; iii) for each genome, it was observed that at least 20% of ORFs per 1-kb

genome could be matched to *Nitrososphaerota* (Supplementary Dataset 2); iv) further

validation was conducted to assess the applicability of this screening criterion to

previously reported *Nitrososphaerota*-associated viruses/MGEs:

1) the 13 identified *Nitrososphaerota*-associated viral/MGE genomes were retrieved

and downloaded from GenBank;

2) the 11 representative genomes were generated from 13 *Nitrososphaerota*-associated

viral genomes through MMseqs2 with parameters [81]: pairwise alignment identity

= 0.9, pairwise alignment coverage for shorter sequences = 0.9;

3) the 9 out of 11 viral genomes derived from GenBank meet this criterion. Two

isolated spindle-shaped viruses (NSV4 and NSV5) failed to pass the test [35]. This

indicates that the approach is also applicable to most previously reported

Nitrososphaerota-associated viral/MGE genomes.

After validating the filtering approach, it was used as a cut-off for viral-host genome-

content similarity to further filter the results obtained from spacer-protospacer

matching. By applying this approach, 171 *Nitrososphaerota*-associated viruses/MGEs

were filtered from 368 spacer-targeted viral/MGE candidates. Similar to previous

studies [27,47], all relevant archaeal CRISPR-Cas systems were evaluated using

CRISPRCasFinder with default parameters [86]. Among the identified CRISPR arrays,

91.2% (n = 301) exhibited an evidence level of 4, characterized by a repeat

conservation index exceeding 70% and spacer identity below 8% (Supplementary

Dataset 2). The remaining 29 viruses/MGEs, corresponding to evidence levels below

4 for CRISPR arrays, were retained for two reasons: 1) they passed genome-content

similarity filtering; 2) in genome-wide analyses, they clustered with other

viruses/MGEs possessing CRISPR arrays with an evidence level of 4.

In addition, IMG/VR (v.4) contains 1,457 viruses/MGEs that were predicted to be

associated with *Nitrososphaerota*, although these contigs were not retrieved through

spacer-protospacer matching. The 1,457 viruses/MGEs predicted to be associated with

Nitrososphaerota in IMG/VR (v.4) were tested. First, 699 representative contigs were

clustered using MMSeq2 with parameters: pairwise alignment identity = 0.9, pairwise

alignment coverage for shorter sequences = 0.9. Second, these contigs were tested by

genome-content similarity calculation, resulting in 441 *Nitrososphaerota*-associated

viruses/MGEs. Finally, 612 *Nitrososphaerota*-associated viruses/MGEs were

identified and filtered from IMG/VR (v.4).The completeness of these viral/MGEs

were evaluate using CheckV [87].

Expanded mining of *Nitrososphaerota*-associated viruses/MGEs from 153 metagenomes

The *Nitrososphaerota*-associated CRISPR spacer database was aligned to from 153

metagenomes using MMseqs2 with the same approach and parameters [81]. Contigs

containing DRs and those shorter than 2-kb were removed before performing

alignment. This resulted in 1,951 contigs matched by *Nitrososphaerota*-associated

CRISPR spacers. To remove duplications, 1,951 contigs were aligned to previously

identified *Nitrosospaherota*-associated viruses/MGEs set using MMseqs2 with same

parameters same parameters [81]. This step generated 1,895 contigs, including 1,396

representative contigs clustered using the same approach described before. To recall

as many as possible viruses/MGEs, these contigs were used to filter *Nitrososphaerota*-

associated viruses/MGEs using three multiple viral identification pipelines, including

geNomad, VirSorter2, and DeepVirFinder[88-90]. Multiple pipelines were employed for

virus identification based on the following considerations: 1) geNomad demonstrates

superior overall classification performance compared to other pipelines [89]; 2)

VirSorter2 (VS2) is recommended when integrating multiple pipelines [91]; 3)

DeepVirFinder (DVF), utilizing an alignment-free approach, is well-suited for

identifying unique viral sequences [88]; 4) the identification results from each pipeline

complement one another, enhancing recall while mitigating cellular contamination in

the dataset[92-94]. For each pipeline, precision was prioritized over sensitivity; therefore,

stringent virus detection strategies were adopted to minimize potential false positives:

1) geNomad was used to detect viral/MGE contigs using conserved mode ("--

conservative"), resulting in a more strict set of viruses with high confidence; 2) VS2

was used to detect viral contigs under high-confidence mode ('--high-confidence-

only'), retaining results with classification score ≥ 0.9, and classification score ≤ 0.7

with viral hallmark ≥ 1; 3) The DVF used to detect viral contig with parameters: score

≥ 0.9 and p-value ≤ 0.01. To validate this hybrid viral detection pipeline, 20% of

sequences were randomly selected from previously identified *Nitrososphaerota*-

associated viruses/MGE and were used as a test set to evaluate the efficiency of the

approach. Based on the test, this viral/MGE identification approach achieved a 74%

recruitment rate, which was considered acceptable for further viral/MGE

identification (Supplementary Dataset 9).

Using the validated hybrid viral identification approach, 419 viral/MGE contigs

were identified from 1,396 metagenome-derived contigs. These contigs had already

been targeted by *Nitrososphaerota*-associated CRISPR spacers in previous steps. The

viral-host genome-content similarity was used to filter these contigs as described

before, resulting 108 *Nitrososphaerota*-associated viral/MGEs. Relevant archaeal

CRISPR-Cas systems of these contigs were evaluated using CRISPRCasFinder with

default parameters [86]. The completeness of these viral/MGEs was evaluated using

CheckV [87].

Detection of viruses/MGEs integrated in *Nitrososphaerota* genomes

Proviruses integrated into *Nitrososphaerota* genomes were identified using the "find-

proviruses" program in geNomad [89]. Integrases were first detected with the MMseqs2

plugin in geNomad [81], applying an E-value cut-off of 1e-5. Additional parameters

were set as follows: a minimum gene-level score of 0.4 for flagging proviral genes

using the conditional random field model; a minimum total virus marker score of 12

for proviruses lacking integrases or not located at scaffold edges; a minimum total

virus marker score of 8 for proviruses encoding integrases or located at scaffold edges;

a maximum allowed distance of 10-kb between provirus boundaries and integrases for

boundary extension; and a maximum allowed distance of 5-kb between provirus

boundaries and tRNAs for boundary extension. This process resulted in the

identification of 30 *Nitrososphaerota*-associated proviruses.

To remove redundancy, the 30 proviral/MGE genomes performed self-alignment

and were compared against previously identified *Nitrososphaerota*-associated

viral/MGE genomes using the same approach and parameters described above. This

filtering step yielded 24 non-redundant *Nitrososphaerota*-associated proviruses

(Supplementary Dataset 2).

Host assignments of *Nitrososphaerota*-associated viruses/MGEs

The host assignments of *Nitrososphaerota*-associated viruses/MGEs were primarily

based on: 1) CRISPR-spacers-associated host genomes; 2) information provided by

previous studies; 3) genome-content similarity between viruses/MGEs and hosts. For

Nitrososphaerota-associated viruses/MGEs that match CRISPR spacers from multiple

distinct hosts, similar with previous study [27], the host of a virus/MGE targeted by

CRISPR spacers was assigned based on the taxonomy of the genome from which the

spacer with the highest identity originated. For *Nitrososphaerota*-associated

viruses/MGEs that were identified based on oligonucleotide pattern similarity, the

most possible host genomes associated with these viruses/MGEs were determined

through alignment against *Nitrososphaerota* genomes using MMseqs2 with the

parameters: 1e-5 as the E-value, iteratively searched two times [81]. A host genome was

linked to a viral/MGE genome if they have the most shared ORFs. This approach was

also applied in IMG/VR (v.4) [83,95]. For *Nitrososphaerota*-associated viruses/MGEs

reported in previous studies, some host information was retrieved from related

literature. For those viruses/MGEs with uncertain host information [33], the related viral

ORFs were subject to MMseqs2 against all *Nitrososphaerota* genomes with the same

approach. The 35 Marthaviruses derived from López-Pérez et al. study were excluded

from this study for two reasons [32]: 1) the related data (PRJNA484324) are still

embargoed when an attempt was made to access these genomes; 2) although the

proposed taxa 'Marthaviruses' has been documented in NCBI, the host information of

these viruses is labelled as bacteria. The archaeal host phylogenetic tree was

calculated based on 46 concatenated ribosomal proteins by IQ-Tree2 (Supplementary

Note 2)[96]. The archaeal lineages were determined based on their phylogenetic

placements and previous studies [2,8,20,21,23,46].

Comparative   genomics   and   family-/species-level   demarcation   of
***Nitrososphaerota*-associated viruses/MGEs**

Similar to previous studies [25,27,29], new families and vOTUs were assigned based on

the genome-content similarity network and the genome-wide proteomic tree. Using

DNA virus sequences longer than 2-kb from NCBI RefSeq v.223 as reference

genomes, vConTACT2 was applied (with default parameters) to compute the genome-

content-based similarity network of those *Nitrososphaerota*-associated viruses/MGEs.

Nitrososphaerota-associated viral/MGE genome pairs that shared at least 20% protein

clusters and encoded at least six genes were filtered from genome-content-based

similarity network, generating a subset of network, which was used to guide

taxonomic assignments. In the network, viruses not related to *Nitrososphaerota*-

associated viruses/MGEs were discarded to achieve a clear layout. Genome-wide

proteomic tree of those genomes included in the subset of network were calculated by

ViPTree v.4. The viruses/MGEs were assigned to the same family if they were

represented by a cohesive group in the genome-content-based similarity network and

a monophyletic group in genome-wide proteomic tree (with threshold 0.05) [27]. The

network and tree were visualized by Cytoscape v.3.10.2 and iToL v.6, respectively

[97,98]. The nomenclature of the proposed viral family was referred to Cthulhu mythos

(Supplementary Note 3).

Annotation of viral/MGEs genomes

Viral/MGE species (or vOTUs) were defined using 95% nucleotide identity threshold

over 90% of the full length using MMseqs2. The family-level taxonomic assignments

of *Nitrososphaerota*-associated viruses/MGEs were assigned using 1) VITAP with

custom database: this database includes all viral reference genomes collected in ICTV

VMR-MSL39, high-quality vOTUs with genus-level taxonomy collected in IMG/VR

(v.4), and all viral genomes collected in NCBI RefSeq v.223 [83,99,100]; 2) CAT with

NCBI RefSeq v.223 database [100,101]; 3) geNomad with default parameters [89]. The CAT

only assigned class/family level units for 10 contigs; the geNomad and VITAP

assigned class/family level units for 683 and 676 contigs, respectively. Hence, the

taxonomic assignments were mainly based on geNomad and VITAP. If geNomad or

VITAP showed consistency at higher taxonomic levels (Phylum or Class), but one of

the pipelines produced annotations at lower taxonomic levels, then the results with the

lower taxonomic level annotations were used. The results with discrepancies between

VITAP and geNomad were compared against the NCBI RefSeq v.223 using MMseqs2

with default parameters and subsequently subjected to manual verification. The

consensus VCs generated from vConTACT2 were used to expanded and furtherly

curated the viral taxonomic assignment [102]. The members of each proposed family

were furtherly expanded based on consensus vOTU.

ORFs encoded in *Nitrososphaerota*-associated viruses/MGEs were predicted by

Prodigal v.2.6.3 with metagenome mode[85], and annotated using the KEGG Ortholog

database (January, 2024), Pfam-A v.35, COG2020, Swiss-Prot database (January,

2024), and VOG223 database [103-105].The pan-proteomes of related host lineages were

aligned to KEGG Ortholog database with the same parameters. The metabolic

pathways involved in viruses and their hosts were visualized using iPath 3 [106].

Determining of *Nitrososphaerota*-associated viral/MGE horizontal gene transfer

events

HGT events between *Nitrososphaerota*-associated viruses (vOTUs) and their archaeal

hosts were identified following an established pipeline [57]. After removal of proviral

sequences, viral and archaeal sequences were aligned, and shared protein families

(PFs) were identified. Potential contamination was rigorously screened based on

lineage, GC content, scaffold length, and genomic context, retaining only robust

alignments (Supplementary Note 4 and Supplementary Dataset 10).

Filtered protein families underwent phylogenetic analysis using MUSCLE5

alignment, trimAl v.1.4 trimming, and IQ-TREE2 tree construction with the

LG+F+R5 model. Trees were rooted via minimal ancestral deviation, and potential

HGT events were classified based on the phylogenetic relationships between viral and

archaeal sequences (Extended Data 5). Transfer direction labels ("v-from-a", "v-to-a",

"v-?-a" and "non-HGT") were assigned following predefined criteria considering

sister and cousin clade compositions. Iterative screening was performed with

temporary reassignment of labels to ensure accurate HGT detection across multiple

rounds of analysis (Supplementary Note 4).

Phylogenetic inference of *Nitrososphaerota*-associated viruses/MGEs

To investigate the evolutionary relationships among rod-shaped and head-tailed

viruses associated with *Nitrososphaerota* within the virosphere context, phylogenetic

analyses were performed based on key viral proteins, including the SIRV2-like major

capsid protein (MCP), HK97-like MCP, and terminase proteins. For rod-shaped

viruses, MCP sequences from 33viruses/MGEs were aligned with 34 viral

RefSeqsv.223 from the *Adnaviria* realm using MMSeqs2 (E-value ≤1e-5). Notably,

some viruses encoded adjacent duplicate MCP subunits, including Metallosphaera

rod-shaped virus 1 and Saccharolobussolfataricus rod-shaped virus 1, along with five

uncultured viral genomes (UViGs). Considering the fundamental rod-shaped viral

MCP structural unit as either homodimers or heterodimers, phylogenetic trees were

inferred based on concatenated MSAs generated using MUSCLE5 (ensemble mode).

The DUF732 domain-containing protein (PF05305), identified via FoldSeek structural

alignment as the closest homolog [107], served as an outgroup for rooting the rod-shaped

viral MCP phylogeny. HK97-MCP sequences exhibited significant diversity,

encompassing multiple families from Pfam-A v.35 and VOG223 [104,108]; thus, three

dominant domains (PF05065, PF07068, and VOG01305) were selected and aligned

against *Duplodnaviria* proteins from RefSeqs v.223. Appropriate outgroups were

chosen based on structural and domain-based alignments (e.g., PF06673 and PF00130). Similarly, terminase proteins were analyzed by selecting the top three domains (K06909, VOG00195, and PF03237) and employing P-loop containing nucleoside triphosphate hydrolases (SSF52540) as an outgroup. All MSAs were refined using trimAl v.1.4, and trees were inferred by ML with IQ-Tree2 and BI using MrBayes [96,109], supported by 1,000 bootstrap (ML) and convergence diagnostics (BI) (Extended Data 3 and 4). The topological comparisons between BI and ML were performed using Phylo.io [110].

Structure analysis of *Nitrososphaerota*-associated viruses/MGEs

To further assess the novelty of capsid proteins from *Nitrososphaerota*-associated viruses/MGEs, the HK97-like and SIRV2-like capsid proteins identified in this study were used to perform structural prediction. A total of 5,153 head-tailed viral genomes from RefSeqs v.223 and 44 rod-shaped archaeal viral genomes from GenBank (length ≥ 2,000 bp) were retrieved from GenBank (as of November 31, 2024). ORFs in these genomes were predicted using Prodigal v.2.6.3 [85]. HK97-like and SIRV2-like capsid proteins were identified using HMM-based searches against specific HMM profiles (PF12193, PF05065, VOG01305, and PF07068) with an E-value cut-off of 1e-5 [111], resulting in 1,844 HK97-like capsids and 28 SIRV2-like capsids. Subsequently, the HK97-like capsid proteins were clustered using MMSeqs2 with thresholds of 0.9 alignment coverage and identity [81], yielding 468 representative sequences. Structures of the HK97-like and SIRV2-like capsid proteins were predicted using AlphaFold2 implemented in ColabFold (Extended Data 6 and 7) [112], and the models with the highest predicted local distance difference test (pLDDT) scores were selected as best-predicted models. Structural alignments were conducted through all-to-all comparisons using FoldSeek with an identity cut-off of 0.5 [107], and reciprocal hits were used to construct a structural similarity network. The structural similarity network was built based on bitscores (top 10% hits were selected) calculated from FoldSeek by Cytoscape[97,107].

Data availability

All nucleotide and protein sequences of viral or mobile genetic elements, spacers from the CRISPR database, phylogenetic trees, structural prediction files are available in FigShare (https://figshare.com/s/a2e94d686028b674a905). The related metagenomes used in this study can be accessed via the URL provided in Supplementary Dataset 1 through IMG/M. Related genomes from IMG/VR database is available at https://img.jgi.doe.gov/cgi-bin/vr/main.cgi. Other information and metadata are provided as Supplementary Datasets.

Code availability

No custom code was generated and used.

Reference

1. Nunoura, T., Takaki, Y., Kakuta, J., Nishi, S., Sugahara, J., Kazama, H., Chee, G.J., Hattori, M., Kanai, A., Atomi, H., et al. (2011). Insights into the evolution of Archaea and eukaryotic protein modifier systems revealed by the genome of a novel archaeal group. Nucleic Acids Res *39*, 3204-3223. 10.1093/nar/gkq1228.

2. Spang, A., Caceres, E.F., and Ettema, T.J.G. (2017). Genomic exploration of the diversity, ecology, and evolution of the archaeal domain of life. Science *357*. 10.1126/science.aaf3883.

3. Alfreider, A., Grimus, V., Luger, M., Ekblad, A., Salcher, M.M., and Summerer, M. (2018). Autotrophic carbon fixation strategies used by nitrifying prokaryotes in freshwater lakes. FEMS Microbiol Ecol *94*. 10.1093/femsec/fiy163.

4. Klotz, F., Kitzinger, K., Ngugi, D.K., Busing, P., Littmann, S., Kuypers, M.M.M., Schink, B., and Pester, M. (2022). Quantification of archaea-driven freshwater nitrification from single cell to ecosystem levels. ISME J *16*, 1647-1656. 10.1038/s41396-022-01216-9.

5. Li, Y., Chen, J., Lin, Y., Zhong, C., Jing, H., and Liu, H. (2024). Thaumarchaeota from deep-sea methane seeps provide novel insights into their evolutionary history and ecological implications. Microbiome *12*, 197. 10.1186/s40168-024-01912-y.

6. Brochier-Armanet, C., Boussau, B., Gribaldo, S., and Forterre, P. (2008). Mesophilic Crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. Nat Rev Microbiol *6*, 245-252. 10.1038/nrmicro1852.

7. Pester, M., Schleper, C., and Wagner, M. (2011). The Thaumarchaeota: an emerging view of their phylogeny and ecophysiology. Curr Opin Microbiol *14*, 300-306. 10.1016/j.mib.2011.04.007.

8. Zhong, H., Lehtovirta-Morley, L., Liu, J., Zheng, Y., Lin, H., Song, D., Todd, J.D., Tian, J., and Zhang, X.H. (2020). Novel insights into the Thaumarchaeota in the deepest oceans: their metabolism and potential adaptation mechanisms. Microbiome *8*, 78. 10.1186/s40168-020-00849-2.

9. Kuypers, M.M.M., Marchant, H.K., and Kartal, B. (2018). The microbial nitrogen-cycling network. Nat Rev Microbiol *16*, 263-276. 10.1038/nrmicro.2018.9.

10. Prosser, J.I., and Nicol, G.W. (2012). Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. Trends

Microbiol *20*, 523-531. 10.1016/j.tim.2012.08.001.

11.   Martens-Habbena, W., Berube, P.M., Urakawa, H., de la Torre, J.R., and Stahl,
D.A. (2009). Ammonia oxidation kinetics determine niche separation of
nitrifying Archaea and Bacteria. Nature *461*, 976-979. 10.1038/nature08465.

12.   Stahl, D.A., and de la Torre, J.R. (2012). Physiology and diversity of
ammonia-oxidizing archaea. Annu Rev Microbiol *66*, 83-101.
10.1146/annurev-micro-092611-150128.

13.   Walker, C.B., de la Torre, J.R., Klotz, M.G., Urakawa, H., Pinel, N., Arp, D.J.,
Brochier-Armanet, C., Chain, P.S., Chan, P.P., Gollabgir, A., et al. (2010).
Nitrosopumilus maritimus genome reveals unique mechanisms for nitrification
and autotrophy in globally distributed marine crenarchaea. Proc Natl Acad Sci
U S A *107*, 8818-8823. 10.1073/pnas.0913533107.

14.   Könneke, M., Schubert, D.M., Brown, P.C., Hügler, M., Standfest, S.,
Schwander, T., Schada von Borzyskowski, L., Erb, T.J., Stahl, D.A., and Berg,
I.A. (2014). Ammonia-oxidizing archaea use the most energy-efficient aerobic
pathway for CO2 fixation. Proceedings of the National Academy of Sciences
111, 8239-8244.

15.   Ren, M., Feng, X., Huang, Y., Wang, H., Hu, Z., Clingenpeel, S., Swan, B.K.,
Fonseca, M.M., Posada, D., Stepanauskas, R., et al. (2019). Phylogenomics
suggests oxygen availability as a driving force in Thaumarchaeota evolution.
ISME J *13*, 2150-2161. 10.1038/s41396-019-0418-8.

16.   Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., and Oakley, B.B.
(2005). Ubiquity and diversity of ammonia-oxidizing archaea in water
columns and sediments of the ocean. Proc Natl Acad Sci U S A *102*, 14683-
14688. 10.1073/pnas.0506625102.

17.   Nelkner, J., Huang, L., Lin, T.W., Schulz, A., Osterholz, B., Henke, C., Blom,
J., Puhler, A., Sczyrba, A., and Schluter, A. (2023). Abundance, classification
and genetic potential of Thaumarchaeota in metagenomes of European
agricultural soils: a meta-analysis. Environ Microbiome *18*, 26.
10.1186/s40793-023-00479-9.

18.   Biggs-Weber, E., Aigle, A., Prosser, J.I., and Gubry-Rangin, C. (2020).
Oxygen preference of deeply-rooted mesophilic thaumarchaeota in forest soil.
Soil Biology and Biochemistry *148*, 107848.
https://doi.org/10.1016/j.soilbio.2020.107848.

19.   Aylward, F.O., and Santoro, A.E. (2020). Heterotrophic Thaumarchaea with
Small Genomes Are Widespread in the Dark Ocean. mSystems *5*.
10.1128/mSystems.00415-20.

20.   Zhang, R.Y., Wang, Y.R., Liu, R.L., Rhee, S.K., Zhao, G.P., and Quan, Z.X.
(2024). Metagenomic characterization of a novel non-ammonia-oxidizing

Thaumarchaeota from hadal sediment. Microbiome *12*, 7. 10.1186/s40168-
023-01728-2.

21.   Hua, Z.S., Qu, Y.N., Zhu, Q., Zhou, E.M., Qi, Y.L., Yin, Y.R., Rao, Y.Z., Tian,
Y., Li, Y.X., Liu, L., et al. (2018). Genomic inference of the metabolism and
evolution of the archaeal phylum Aigarchaeota. Nat Commun *9*, 2832.
10.1038/s41467-018-05284-4.

22.   Beam, J.P., Jay, Z.J., Schmid, M.C., Rusch, D.B., Romine, M.F., Jennings Rde,
M., Kozubal, M.A., Tringe, S.G., Wagner, M., and Inskeep, W.P. (2016).
Ecophysiology of an uncultivated lineage of Aigarchaeota from an oxic, hot
spring filamentous 'streamer' community. ISME J *10*, 210-224.
10.1038/ismej.2015.83.

23.   Buessecker, S., Palmer, M., Lai, D., Dimapilis, J., Mayali, X., Mosier, D., Jiao,
J.Y., Colman, D.R., Keller, L.M., St John, E., et al. (2022). An essential role
for tungsten in the ecology and evolution of a previously uncultivated lineage
of anaerobic, thermophilic Archaea. Nat Commun *13*, 3773. 10.1038/s41467-
022-31452-8.

24.   Prangishvili, D., Bamford, D.H., Forterre, P., Iranzo, J., Koonin, E.V., and
Krupovic, M. (2017). The enigmatic archaeal virosphere. Nat Rev Microbiol
15, 724-739. 10.1038/nrmicro.2017.125.

25.   Medvedeva, S., Sun, J., Yutin, N., Koonin, E.V., Nunoura, T., Rinke, C., and
Krupovic, M. (2022). Three families of Asgard archaeal viruses identified in
metagenome-assembled genomes. Nat Microbiol *7*, 962-973. 10.1038/s41564-
022-01144-6.

26.   Rambo, I.M., Langwig, M.V., Leao, P., De Anda, V., and Baker, B.J. (2022).
Genomes of six viruses that infect Asgard archaea from deep-sea sediments.
Nat Microbiol *7*, 953-961. 10.1038/s41564-022-01150-8.

27.   Medvedeva, S., Borrel, G., Krupovic, M., and Gribaldo, S. (2023). A
compendium of viruses from methanogenic archaea reveals their diversity and
adaptations to the gut environment. Nat Microbiol *8*, 2170-2182.
10.1038/s41564-023-01485-w.

28.   Laso-Perez, R., Wu, F., Cremiere, A., Speth, D.R., Magyar, J.S., Zhao, K.,
Krupovic, M., and Orphan, V.J. (2023). Evolutionary diversification of
methanotrophic ANME-1 archaea and their expansive virome. Nat Microbiol.
10.1038/s41564-022-01297-4.

29.   Sakai, H.D., Nur, N., Kato, S., Yuki, M., Shimizu, M., Itoh, T., Ohkuma, M.,
Suwanto, A., and Kurosawa, N. (2022). Insight into the symbiotic lifestyle of
DPANN archaea revealed by cultivation and genome analyses. Proc Natl Acad
Sci U S A *119*. 10.1073/pnas.2115449119.

30.   Krupovic, M., Spang, A., Gribaldo, S., Forterre, P., and Schleper, C. (2011). A

753 thaumarchaeal provirus testifies for an ancient association of tailed viruses
754 with archaea. Biochem Soc Trans *39*, 82-88. 10.1042/BST0390082.

755 31. Krupovic, M., Makarova, K.S., Wolf, Y.I., Medvedeva, S., Prangishvili, D.,
756 Forterre, P., and Koonin, E.V. (2019). Integrated mobile genetic elements in
757 Thaumarchaeota. Environ Microbiol *21*, 2056-2078. 10.1111/1462-
758 2920.14564.

759 32. Lopez-Perez, M., Haro-Moreno, J.M., de la Torre, J.R., and Rodriguez-Valera,
760 F. (2019). Novel Caudovirales associated with Marine Group I
761 Thaumarchaeota assembled from metagenomes. Environ Microbiol *21*, 1980-
762 1988. 10.1111/1462-2920.14462.

763 33. Ahlgren, N.A., Fuchsman, C.A., Rocap, G., and Fuhrman, J.A. (2019).
764 Discovery of several novel, widespread, and ecologically distinct marine
765 Thaumarchaeota viruses that encode amoC nitrification genes. ISME J *13*,
766 618-631. 10.1038/s41396-018-0289-4.

767 34. Zhou, Y., Zhou, L., Yan, S., Chen, L., Krupovic, M., and Wang, Y. (2023).
768 Diverse viruses of marine archaea discovered using metagenomics. Environ
769 Microbiol *25*, 367-382. 10.1111/1462-2920.16287.

770 35. Kim, J.G., Kim, S.J., Cvirkaite-Krupovic, V., Yu, W.J., Gwak, J.H., Lopez-
771 Perez, M., Rodriguez-Valera, F., Krupovic, M., Cho, J.C., and Rhee, S.K.
772 (2019). Spindle-shaped viruses infect marine ammonia-oxidizing
773 thaumarchaea. Proc Natl Acad Sci U S A *116*, 15645-15650.
774 10.1073/pnas.1905682116.

775 36. Liu, Y., Brandt, D., Ishino, S., Ishino, Y., Koonin, E.V., Kalinowski, J.,
776 Krupovic, M., and Prangishvili, D. (2019). New archaeal viruses discovered
777 by metagenomic analysis of viral communities in enrichment cultures. Environ
778 Microbiol *21*, 2002-2014. 10.1111/1462-2920.14479.

779 37. Nayfach, S., Roux, S., Seshadri, R., Udwary, D., Varghese, N., Schulz, F., Wu,
780 D., Paez-Espino, D., Chen, I.M., Huntemann, M., et al. (2021). A genomic
781 catalog of Earth's microbiomes. Nat Biotechnol *39*, 499-509. 10.1038/s41587-
782 020-0718-6.

783 38. Chen, I.A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M.,
784 Hajek, P., Ritter, S., Varghese, N., Seshadri, R., et al. (2021). The IMG/M data
785 management and analysis system v.6.0: new tools and advanced capabilities.
786 Nucleic Acids Res *49*, D751-D763. 10.1093/nar/gkaa939.

787 39. Sharma, S., Lee, M., Reinmann, C.S., Pumneo, J., Cutright, T.J., and Senko,
788 J.M. (2020). Impact of acid mine drainage chemistry and microbiology on the
789 development of efficient Fe removal activities. Chemosphere *249*, 126117.
790 10.1016/j.chemosphere.2020.126117.

791 40. Zhang, X., Xu, W., Liu, Y., Cai, M., Luo, Z., and Li, M. (2018). Metagenomics

Reveals Microbial Diversity and Metabolic Potentials of Seawater and Surface
Sediment From a Hadal Biosphere at the Yap Trench. Front Microbiol *9*, 2402.
10.3389/fmicb.2018.02402.

41. Acinas, S.G., Sanchez, P., Salazar, G., Cornejo-Castillo, F.M., Sebastian, M.,
Logares, R., Royo-Llonch, M., Paoli, L., Sunagawa, S., Hingamp, P., et al.
(2021). Deep ocean metagenomes provide insight into the metabolic
architecture of bathypelagic microbial communities. Commun Biol *4*, 604.
10.1038/s42003-021-02112-2.

42. Karner, M.B., DeLong, E.F., and Karl, D.M. (2001). Archaeal dominance in
the mesopelagic zone of the Pacific Ocean. Nature *409*, 507-510.
10.1038/35054051.

43. Schattenhofer, M., Fuchs, B.M., Amann, R., Zubkov, M.V., Tarran, G.A., and
Pernthaler, J. (2009). Latitudinal distribution of prokaryotic picoplankton
populations in the Atlantic Ocean. Environ Microbiol *11*, 2078-2093.
10.1111/j.1462-2920.2009.01929.x.

44. Pester, M., Rattei, T., Flechl, S., Grongroft, A., Richter, A., Overmann, J.,
Reinhold-Hurek, B., Loy, A., and Wagner, M. (2012). amoA-based consensus
phylogeny of ammonia-oxidizing archaea and deep sequencing of amoA genes
from soils of four different geographic regions. Environ Microbiol *14*, 525-
539. 10.1111/j.1462-2920.2011.02666.x.

45. Vetriani, C., Jannasch, H.W., MacGregor, B.J., Stahl, D.A., and Reysenbach,
A.L. (1999). Population structure and phylogenetic characterization of marine
benthic Archaea in deep-sea sediments. Appl Environ Microbiol *65*, 4375-
4384. 10.1128/AEM.65.10.4375-4384.1999.

46. Parks, D.H., Rinke, C., Chuvochina, M., Chaumeil, P.A., Woodcroft, B.J.,
Evans, P.N., Hugenholtz, P., and Tyson, G.W. (2017). Recovery of nearly
8,000 metagenome-assembled genomes substantially expands the tree of life.
Nat Microbiol *2*, 1533-1542. 10.1038/s41564-017-0012-7.

47. Wu, Z., Liu, S., and Ni, J. (2024). Metagenomic characterization of viruses
and mobile genetic elements associated with the DPANN archaeal
superphylum. Nat Microbiol. 10.1038/s41564-024-01839-y.

48. Liu, Y., Demina, T.A., Roux, S., Aiewsakun, P., Kazlauskas, D., Simmonds, P.,
Prangishvili, D., Oksanen, H.M., and Krupovic, M. (2021). Diversity,
taxonomy, and evolution of archaeal viruses of the class Caudoviricetes. PLoS
Biol *19*, e3001442. 10.1371/journal.pbio.3001442.

49. Breier, A.M., and Grossman, A.D. (2007). Whole-genome analysis of the
chromosome partitioning and sporulation protein Spo0J (ParB) reveals
spreading and origin-distal sites on the Bacillus subtilis chromosome. Mol
Microbiol *64*, 703-718. 10.1111/j.1365-2958.2007.05690.x.

50. Chu, C.H., Wu, C.T., Lin, M.G., Yen, C.Y., Wu, Y.Z., Hsiao, C.D., and Sun,
Y.J. (2024). Insights into the molecular mechanism of ParABS system in
chromosome partition by HpParA and HpParB. Nucleic Acids Res *52*, 7321-
7336. 10.1093/nar/gkae450.

51. Krupovic, M., Kuhn, J.H., Wang, F., Baquero, D.P., Dolja, V.V., Egelman, E.H.,
Prangishvili, D., and Koonin, E.V. (2021). Adnaviria: a New Realm for
Archaeal Filamentous Viruses with Linear A-Form Double-Stranded DNA
Genomes. J Virol *95*, e0067321. 10.1128/JVI.00673-21.

52. Nyyssola, A., Kerovuo, J., Kaukinen, P., von Weymarn, N., and Reinikainen, T.
(2000). Extreme halophiles synthesize betaine from glycine by methylation. J
Biol Chem *275*, 22196-22201. 10.1074/jbc.M910111199.

53. Lee, Y.R., Lin, T.S., Lai, S.J., Liu, M.S., Lai, M.C., and Chan, N.L. (2016).
Structural Analysis of Glycine Sarcosine N-methyltransferase from
Methanohalophilus portucalensis Reveals Mechanistic Insights into the
Regulation of Methyltransferase Activity. Sci Rep *6*, 38071.
10.1038/srep38071.

54. Chen, S., Dai, Y., Ke, J., Luo, Y., Wang, C., Hao, Y., Zhang, A., Han, J., and
Xiang, H. (2024). Halocin H4 is activated through cleavage by halolysin
HlyR4. Appl Environ Microbiol *90*, e0228423. 10.1128/aem.02284-23.

55. Torres, M.D.T., Wan, F., and de la Fuente-Nunez, C. (2024). Discovery of
antibiotics in the archaeome using deep learning. bioRxiv.
10.1101/2024.11.15.623859.

56. Wang, P., Li, M., Dong, L., Zhang, C., and Xie, W. (2022). Comparative
Genomics of Thaumarchaeota From Deep-Sea Sponges Reveal Their Niche
Adaptation. Front Microbiol *13*, 869834. 10.3389/fmicb.2022.869834.

57. Irwin, N.A.T., Pittis, A.A., Richards, T.A., and Keeling, P.J. (2022).
Systematic evaluation of horizontal gene transfer between eukaryotes and
viruses. Nat Microbiol *7*, 327-336. 10.1038/s41564-021-01026-3.

58. Thiaville, J.J., Kellner, S.M., Yuan, Y., Hutinet, G., Thiaville, P.C.,
Jumpathong, W., Mohapatra, S., Brochier-Armanet, C., Letarov, A.V.,
Hillebrand, R., et al. (2016). Novel genomic island modifies DNA with 7-
deazaguanine derivatives. Proc Natl Acad Sci U S A *113*, E1452-1459.
10.1073/pnas.1518570113.

59. Gaur, R., and Varshney, U. (2005). Genetic analysis identifies a function for
the queC (ybaX) gene product at an initial step in the queuosine biosynthetic
pathway in Escherichia coli. J Bacteriol *187*, 6893-6901.
10.1128/JB.187.20.6893-6901.2005.

60. Tubiana, T., Boulard, Y., and Bressanelli, S. (2017). Dynamics and asymmetry
in the dimer of the norovirus major capsid protein. PLoS One *12*, e0182056.

10.1371/journal.pone.0182056.

61. Hadden, J.A., Perilla, J.R., Schlicksup, C.J., Venkatakrishnan, B., Zlotnick, A., and Schulten, K. (2018). All-atom molecular dynamics of the HBV capsid reveals insights into biological function and cryo-EM resolution limits. Elife *7*. 10.7554/eLife.32478.

62. Mueller, K.L. (2016). Structural insights into capsid flexibility. Science *354*, 1387-1389. 10.1126/science.354.6318.1387-p.

63. Budean, D., Almeida-Hernández, Y., Pandey, J., Pérez, J.M., Sánchez García, E., and Haglund, E. (2024). Exploring Chemokine Homodimer Stability: Structural Insights into CXC and CC Interfaces. Biophysica *4*, 561-572.

64. Krupovic, M., Dolja, V.V., and Koonin, E.V. (2019). Origin of viruses: primordial replicators recruiting capsids from hosts. Nat Rev Microbiol *17*, 449-458. 10.1038/s41579-019-0205-6.

65. Koonin, E.V., Dolja, V.V., and Krupovic, M. (2022). The logic of virus evolution. Cell Host Microbe *30*, 917-929. 10.1016/j.chom.2022.06.008.

66. Hwang, Y., Roux, S., Coclet, C., Krause, S.J.E., and Girguis, P.R. (2023). Viruses interact with hosts that span distantly related microbial domains in dense hydrothermal mats. Nat Microbiol *8*, 946-957. 10.1038/s41564-023-01347-5.

67. Hurwitz, B.L., and U'Ren, J.M. (2016). Viral metabolic reprogramming in marine ecosystems. Curr Opin Microbiol *31*, 161-168. 10.1016/j.mib.2016.04.002.

68. Tian, F., Wainaina, J.M., Howard-Varona, C., Dominguez-Huerta, G., Bolduc, B., Gazitua, M.C., Smith, G., Gittrich, M.R., Zablocki, O., Cronin, D.R., et al. (2024). Prokaryotic-virus-encoded auxiliary metabolic genes throughout the global oceans. Microbiome *12*, 159. 10.1186/s40168-024-01876-z.

69. Kieft, K., Zhou, Z., Anderson, R.E., Buchan, A., Campbell, B.J., Hallam, S.J., Hess, M., Sullivan, M.B., Walsh, D.A., Roux, S., and Anantharaman, K. (2021). Ecology of inorganic sulfur auxiliary metabolism in widespread bacteriophages. Nat Commun *12*, 3503. 10.1038/s41467-021-23698-5.

70. Huang, X., Zhou, Z., Liu, H., Li, Y., Ge, T., Tang, X., He, Y., Ma, B., Xu, J., Anantharaman, K., and Li, Y. (2024). Soil nutrient conditions alter viral lifestyle strategy and potential function in phosphorous and nitrogen metabolisms. Soil Biology and Biochemistry *189*, 109279. https://doi.org/10.1016/j.soilbio.2023.109279.

71. Wang, L., Wang, Y., Huang, X., Ma, R., Li, J., Wang, F., Jiao, N., and Zhang, R. (2022). Potential metabolic and genetic interaction among viruses, methanogen and methanotrophic archaea, and their syntrophic partners. ISME Communications *2*, 50. 10.1038/s43705-022-00135-2.

72. Zhou, Y., Gutiérrez-Preciado, A., Liu, Y., Moreira, D., Yakimov, M.M., López-
García, P., and Krupovic, M. (2025). Nested parasitism in hypersaline
environments: viruses and virus satellites of haloarchaea and their nanosized
cellular symbionts. bioRxiv, 2025.2002.2015.638363.
10.1101/2025.02.15.638363.

73. D'Souza-Ault, M.R., Smith, L.T., and Smith, G.M. (1993). Roles of N-
acetylglutaminylglutamine amide and glycine betaine in adaptation of
Pseudomonas aeruginosa to osmotic stress. Appl Environ Microbiol *59*, 473-
478. 10.1128/aem.59.2.473-478.1993.

74. Arumugam, M.K., Paal, M.C., Donohue, T.M., Jr., Ganesan, M., Osna, N.A.,
and Kharbanda, K.K. (2021). Beneficial Effects of Betaine: A Comprehensive
Review. Biology (Basel) *10*. 10.3390/biology10060456.

75. Seong, H.J., Roux, S., Hwang, C.Y., and Sul, W.J. (2022). Marine DNA
methylation patterns are associated with microbial community composition
and inform virus-host dynamics. Microbiome *10*, 157. 10.1186/s40168-022-
01340-w.

76. Takahashi, M., Hiraoka, S., Matsumoto, Y., Shibagaki, R., Ujihara, T., Maeda,
H., Seo, S., Nagasaki, K., Takeuchi, H., and Matsuzaki, S. (2025). Host-
encoded DNA methyltransferases modify the epigenome and host tropism of
invading phages. iScience *28*, 112264. 10.1016/j.isci.2025.112264.

77. Rani, A., Alam, A., Ahmad, F., P, M., Saurabh, A., Zarin, S., Mitra, D.K.,
Hasnain, S.E., and Ehtesham, N.Z. (2022). Mycobacterium tuberculosis
Methyltransferase Rv1515c Can Suppress Host Defense Mechanisms by
Modulating Immune Functions Utilizing a Multipronged Mechanism. Front
Mol Biosci *9*, 906387. 10.3389/fmolb.2022.906387.

78. Kumar, V., and Tiwari, S.K. (2017). Halocin HA1: An archaeocin produced by
the haloarchaeon Haloferax larsenii HA1. Process Biochemistry *61*, 202-208.
https://doi.org/10.1016/j.procbio.2017.06.010.

79. Skennerton, C.T., Imelfort, M., and Tyson, G.W. (2013). Crass: identification
and reconstruction of CRISPR from unassembled metagenomic data. Nucleic
Acids Res *41*, e105. 10.1093/nar/gkt183.

80. Bland, C., Ramsey, T.L., Sabree, F., Lowe, M., Brown, K., Kyrpides, N.C.,
and Hugenholtz, P. (2007). CRISPR recognition tool (CRT): a tool for
automatic detection of clustered regularly interspaced palindromic repeats.
BMC Bioinformatics *8*, 209. 10.1186/1471-2105-8-209.

81. Steinegger, M., and Soding, J. (2017). MMseqs2 enables sensitive protein
sequence searching for the analysis of massive data sets. Nat Biotechnol *35*,
1026-1028. 10.1038/nbt.3988.

82. Padilha, V.A., Alkhnbashi, O.S., Shah, S.A., de Carvalho, A., and Backofen, R.

(2020). CRISPRcasIdentifier: Machine learning for accurate identification and

classification of CRISPR-Cas systems. Gigascience *9*.

10.1093/gigascience/giaa062.

83. Camargo, A.P., Nayfach, S., Chen, I.A., Palaniappan, K., Ratner, A., Chu, K.,

Ritter, S.J., Reddy, T.B.K., Mukherjee, S., Schulz, F., et al. (2023). IMG/VR

v4: an expanded database of uncultivated virus genomes within a framework

of extensive functional, taxonomic, and ecological metadata. Nucleic Acids

Res *51*, D733-D743. 10.1093/nar/gkac1037.

84. Ahlgren, N.A., Chen, Y., Needham, D.M., Parada, A.E., Sachdeva, R., Trinh,

V., Chen, T., and Fuhrman, J.A. (2017). Genome and epigenome of a novel

marine Thaumarchaeota strain suggest viral infection, phosphorothioation

DNA modification and multiple restriction systems. Environ Microbiol *19*,

2434-2452. 10.1111/1462-2920.13768.

85. Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., and Hauser,

L.J. (2010). Prodigal: prokaryotic gene recognition and translation initiation

site identification. BMC Bioinformatics *11*, 119. 10.1186/1471-2105-11-119.

86. Couvin, D., Bernheim, A., Toffano-Nioche, C., Touchon, M., Michalik, J.,

Neron, B., Rocha, E.P.C., Vergnaud, G., Gautheret, D., and Pourcel, C. (2018).

CRISPRCasFinder, an update of CRISRFinder, includes a portable version,

enhanced performance and integrates search for Cas proteins. Nucleic Acids

Res *46*, W246-W251. 10.1093/nar/gky425.

87. Nayfach, S., Camargo, A.P., Schulz, F., Eloe-Fadrosh, E., Roux, S., and

Kyrpides, N.C. (2021). CheckV assesses the quality and completeness of

metagenome-assembled viral genomes. Nat Biotechnol *39*, 578-585.

10.1038/s41587-020-00774-7.

88. Ren, J., Song, K., Deng, C., Ahlgren, N.A., Fuhrman, J.A., Li, Y., Xie, X.,

Poplin, R., and Sun, F. (2020). Identifying viruses from metagenomic data

using deep learning. Quant Biol *8*, 64-77. 10.1007/s40484-019-0187-4.

89. Camargo, A.P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain,

P.S.G., Nayfach, S., and Kyrpides, N.C. (2023). Identification of mobile

genetic elements with geNomad. Nat Biotechnol. 10.1038/s41587-023-01953-

y.

90. Guo, J., Bolduc, B., Zayed, A.A., Varsani, A., Dominguez-Huerta, G.,

Delmont, T.O., Pratama, A.A., Gazitua, M.C., Vik, D., Sullivan, M.B., and

Roux, S. (2021). VirSorter2: a multi-classifier, expert-guided approach to

detect diverse DNA and RNA viruses. Microbiome *9*, 37. 10.1186/s40168-

020-00990-y.

91. Hegarty, B., Riddell, V.J., Bastien, E., Langenfeld, K., Lindback, M., Saini,

J.S., Wing, A., Zhang, J., and Duhaime, M. (2024). Benchmarking informatics

987 approaches for virus discovery: caution is needed when combining in silico

988 identification methods. mSystems *9*, e0110523. 10.1128/msystems.01105-23.

989 92. Li, R., Wang, Y., Hu, H., Tan, Y., and Ma, Y. (2022). Metagenomic analysis

990 reveals unexplored diversity of archaeal virome in the human gut. Nat

991 Commun *13*, 7978. 10.1038/s41467-022-35735-y.

992 93. Chibani, C.M., Shah, S.A., Schmitz, R.A., and Nayfach, S. (2024). Inaccurate

993 viral prediction leads to overestimated diversity of the archaeal virome in the

994 human gut. Nat Commun *15*, 5976. 10.1038/s41467-024-49902-w.

995 94. Wang, Y., Li, R., and Ma, Y. (2024). Reply to: Inaccurate viral prediction leads

996 to overestimated diversity of the archaeal virome in the human gut. Nat

997 Commun *15*, 5977. 10.1038/s41467-024-49903-9.

998 95. Paez-Espino, D., Eloe-Fadrosh, E.A., Pavlopoulos, G.A., Thomas, A.D.,

999 Huntemann, M., Mikhailova, N., Rubin, E., Ivanova, N.N., and Kyrpides, N.C.

1000 (2016). Uncovering Earth's virome. Nature *536*, 425-430.

1001 10.1038/nature19094.

1002 96. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D.,

1003 von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and

1004 Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol

1005 Evol *37*, 1530-1534. 10.1093/molbev/msaa015.

1006 97. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D.,

1007 Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software

1008 environment for integrated models of biomolecular interaction networks.

1009 Genome Res *13*, 2498-2504. 10.1101/gr.1239303.

1010 98. Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent

1011 updates to the phylogenetic tree display and annotation tool. Nucleic Acids

1012 Res *52*, W78-W82. 10.1093/nar/gkae268.

1013 99. Zheng, K., Sun, J., Liang, Y., Kong, L., Paez-Espino, D., McMinn, A., and

1014 Wang, M. (2025). VITAP: a high precision tool for DNA and RNA viral

1015 classification based on meta-omic data. Nat Commun *16*, 2226.

1016 10.1038/s41467-025-57500-7.

1017 100. Pruitt, K.D., Tatusova, T., Brown, G.R., and Maglott, D.R. (2012). NCBI

1018 Reference Sequences (RefSeq): current status, new features and genome

1019 annotation policy. Nucleic Acids Res *40*, D130-135. 10.1093/nar/gkr1079.

1020 101. von Meijenfeldt, F.A.B., Arkhipova, K., Cambuy, D.D., Coutinho, F.H., and

1021 Dutilh, B.E. (2019). Robust taxonomic classification of uncharted microbial

1022 sequences and bins with CAT and BAT. Genome Biol *20*, 217.

1023 10.1186/s13059-019-1817-x.

1024 102. Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J.H., Roux, S., Adriaenssens,

1025 E.M., Brister, J.R., Kropinski, A.M., Krupovic, M., Lavigne, R., et al. (2019).

Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. Nat Biotechnol *37*, 632-639. 10.1038/s41587-019-0100-8.

103.   Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., and Ogata, H. (2020). KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics *36*, 2251-2252. 10.1093/bioinformatics/btz859.

104.   Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer, E.L.L., Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., et al. (2021). Pfam: The protein families database in 2021. Nucleic Acids Res *49*, D412-D419. 10.1093/nar/gkaa913.

105.   Galperin, M.Y., Wolf, Y.I., Makarova, K.S., Vera Alvarez, R., Landsman, D., and Koonin, E.V. (2021). COG database update: focus on microbial diversity, model organisms, and widespread pathogens. Nucleic Acids Res *49*, D274-D281. 10.1093/nar/gkaa1018.

106.   Darzi, Y., Letunic, I., Bork, P., and Yamada, T. (2018). iPath3.0: interactive pathways explorer v3. Nucleic Acids Res *46*, W510-W513. 10.1093/nar/gky299.

107.   Kim, W., Mirdita, M., Levy Karin, E., Gilchrist, C.L.M., Schweke, H., Soding, J., Levy, E.D., and Steinegger, M. (2025). Rapid and sensitive protein complex alignment with Foldseek-Multimer. Nat Methods. 10.1038/s41592-025-02593-7.

108.   Grazziotin, A.L., Koonin, E.V., and Kristensen, D.M. (2017). Prokaryotic Virus Orthologous Groups (pVOGs): a resource for comparative genomics and protein family annotation. Nucleic Acids Res *45*, D491-D498. 10.1093/nar/gkw975.

109.   Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S., Larget, B., Liu, L., Suchard, M.A., and Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol *61*, 539-542. 10.1093/sysbio/sys029.

110.   Robinson, O., Dylus, D., and Dessimoz, C. (2016). Phylo.io: Interactive Viewing and Comparison of Large Phylogenetic Trees on the Web. Mol Biol Evol *33*, 2163-2166. 10.1093/molbev/msw080.

111.   Mistry, J., Finn, R.D., Eddy, S.R., Bateman, A., and Punta, M. (2013). Challenges in homology search: HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res *41*, e121. 10.1093/nar/gkt263.

112.   Mirdita, M., Schutze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022). ColabFold: making protein folding accessible to all. Nat Methods *19*, 679-682. 10.1038/s41592-022-01488-1.

Acknowledgements

We thank Jilu Han for his help with the installation and debugging of the software used in this study. We thank the support of the high-performance servers of the Center for High Performance Computing and System Simulation, Laoshan Laboratory (Qingdao), the High-Performance Biological Supercomputing Center at the Ocean University of China, the Marine Big Data Center of the Institute for Advanced Ocean Study of the Ocean University of China, and the IEMB-1, a high-performance computing cluster operated by the Institute of Evolution and Marine Biodiversity. This study was supported by the National Natural Science Foundation of China (No. 42120104006, 42176111, and 42306111), the Ocean Negative Carbon Emissions (ONCE), 2024 Graduate Self-directed Research Project (2024ZZKY, 202461036), and the Fundamental Research Funds for the Central Universities (202172002, 201812002, 202461036, Andrew McMinn).

Author contributions

Conceptualization: K.Z. and Y.L.; Analysis: K.Z.; Database: D.P.-E.; Writing manuscript: K.Z.; Review and editing: Y.L., A.M., and M.W.; Computational resource: D.P.-E. and M.W.; Funding acquisition and supervision: Y.L., A.M., and M.W.

Competing interests

The authors declare no competing interests for the manuscript "Viruses associated with *Nitrososphaerota* archaea: diversity, evolution and horizontal gene transfers".

 Figure legends

Figure 1 The establishment of *Nitrososphaerota*-associated viruses and mobile genetic

elements (MGEs). a, The workflow for mining, filtering, and identifying

viruses/MGEs primarily included the prediction of archaeal CRISPR systems,

viral/MGE mining, genome filtering, and the definition of viral operational

taxonomic units (vOTUs). Only genomes encoding detectable capsid or related

proteins were considered viruses; others were classified as MGEs. b, The data

sources for *Nitrososphaerota*-associated viruses/MGEs included in this genomic

catalogue. Due to the pipeline's prioritization of specificity, not all previously

reported *Nitrososphaerota*-associated viral/MGE genomes were included. c,

Habitat sources of Nitrososphaerota-associated viruses/MGEs in this genomic

catalogue at the vOTU level.

Figure 2 Overview of the diversity of the genome catalogue of *Nitrososphaerota*-

associated viruses and mobile genetic elements (MGEs). a, Spacer-sharing

network among *Nitrososphaerota* genomes. Nodes represent archaeal genomes

and are colored by distinct archaeal lineages. An edge is generated between two

genomes if they share spacer sequences, suggesting that their CRISPR systems

may target the same viral/MGE species. Bold edges indicate that the shared

spacer sequences have corresponding protospacers present in the

Nitrososphaerota-associated viral/MGE catalogue generated in this study. b,

Biogeographic distribution of *Nitrososphaerota*-associated viral/MGE genomes.

In the map, nodes of different colors indicate their putative archaeal hosts:

Aigarchaea, ammonia-oxidizing archaea (AOA), non-ammonia-oxidizing

Thaumarchaea (Non-AO Thaum), and deeply-branching *Nitrososphaerota*

archaea (D-B NitroA). c, The number of viral operational taxonomic units

(vOTUs) from each habitat in the *Nitrososphaerota*-associated viral/MGE

catalogue. Bar colors represent different archaeal lineages, including Group 1.1a,

 Nitrosotales, Group 1.1b, Marine Trench and Ground Water (MT/GW),

Nitrosocaldales, Marine Beneath Group A (MBG-A), deeply-branching

Thaumarchaea (D-B Thaum), *Wolframiiraptoraceae*, *Caldarchaeales*, pSL4-

affiliation, and D-B NitroA. Virion icons to the right of the bar plot illustrate the

numbers of viruses/MGEs with different possible morphologies across habitats. d,

Phylogenetic tree of *Nitrososphaerota* archaeal genomes. This tree was

constructed based on concatenated alignments of 46 ribosomal proteins (totaling

5,963 amino acid positions) using the maximum likelihood method with the

LG+C60+F model as recommended by Spang et al.[2] (see Methods). Reference

genomes from *Thermoprotei* were used as the outgroup to root the tree. Tree

annotation includes the habitats and lineages of these archaeal genomes. Virion

icons to the right of the tree indicate the numbers of viruses/MGEs with different

possible morphologies (at the vOTU level) associated with each archaeal lineage.

Figure 3 Similarity of *Nitrososphaerota*-associated viruses and mobile genetic

elements (MGEs) based on their genome contents. a, Genome-content similarity

network of *Nitrososphaerota*-associated viruses/MGEs. Nodes represent

viral/MGE genomes and are coloured according to their associated archaeal

lineages. Each distinct super viral cluster (sVC) may consist of one or more viral

clusters (VCs). The percentage of shared proteins between genomes is indicated

by the thickness of the edges connecting the nodes. The shapes of the nodes

denote the potential morphologies of these viruses/MGEs. b, Viral proteomic tree

used for the delineation of viral families. A distance threshold of 0.05 and a

minimum of six shared proteins among members were used as the criteria for

defining families [27] (see Methods). The names of the proposed viral families or

MGE families are indicated on the right side of the tree, with different virion

icons representing the possible morphologies of these families. Bar plots indicate

the percentage of shared proteins within each family. Tree annotations, from left

to right, include genome length, G+C content (%), coding density (%), habitats,

host lineage, and morphology. Abbreviations: MT/GW, Marine Trench and

Ground Water; MBG-A, Marine Beneath Group A; D-B Thaum, deeply-

branching Thaumarchaea; D-B NitroA, deeply-branching Nitrososphaerota
archaea.

Figure 4 Phylogenetic inferences of *Nitrososphaerota*-associated viruses and mobile
genetic elements (MGEs) based on viral hallmark proteins (VHPs) through
Bayesian inference (BI) and Maximum Likelihood (ML) joint approach. a, The
trees display the topologies inferred by BI based on VHPs. *Nitrososphaerota*-
associated viruses/MGEs are highlighted with bold red branches. To the right of
each tree are shown: 1) BI convergence indices, including minimum effective
sample size (minESS), potential scale reduction factor (PSRF), and average
standard deviation of split frequencies (ASDSF); 2) indices for comparing BI
and ML topologies, including normalized Robinson–Foulds (RF) distance based
on ML and the number of clades shared between BI and ML. Posterior
probabilities from BI are indicated on the tree branches with colored nodes,
alongside ML bootstrap values (MLB). Names of viral families (in bold red font)
and other host groups are annotated beside the corresponding branches. Tree
annotations indicate the *Nitrososphaerota* lineages associated with these
viruses/MGEs. a, Bayesian tree based on the SIRV2-like capsid protein (domain
accession in Pfam: PF12193). DUF732 (domain accession in Pfam: PF05305)
was used as the outgroup to root the tree. Leaves corresponding to rod-shaped
viruses with heterodimeric capsids are highlighted with red squares. b, Bayesian
tree based on the HK97-like capsid protein (domain accession in Pfam:
PF05065), rooted using L_lactis_ph-MCP (domain accession in Pfam: PF06673)
as the outgroup. c, Bayesian tree based on the HK97-like capsid protein (domain
accession in VOGdb: VOG01305), rooted using C1_1 (domain accession in
Pfam: PF00130) as the outgroup. d, Bayesian tree based on the HK97-like capsid
protein (domain accession in Pfam: PF07068), rooted using L_lactis_ph-MCP
(domain accession in Pfam: PF06673) as the outgroup. e, Bayesian tree based on
the terminase large subunit protein (domain accession in KEGG: K06909),
rooted using P-loop NTPase (domain accession in SUPERFAMILY: SSF52540)
as the outgroup. f, Bayesian tree based on the terminase large subunit protein

(domain accession in Pfam: PF03237), rooted using P-loop NTPase (domain
accession in SUPERFAMILY: SSF52540) as the outgroup. Abbreviations:
MT/GW, Marine Trench and Ground Water; MBG-A, Marine Beneath Group A;
D-B Thaum, deeply-branching *Thaumarchaea*; D-B NitroA, deeply-branching
Nitrososphaerota archaea.

Figure 5 Genes encoded by *Nitrososphaerota*-associated viruses and mobile genetic
elements (MGEs) may be involved in metabolic compensation for their archaeal
hosts. Hexagons of different colours represent viruses/MGEs associated with
different *Nitrososphaerota* lineages. If a particular metabolic gene is found only
in viral/MGE genomes but is absent from the panproteome of the corresponding
archaeal lineage, it is considered as compensation from viruses/MGEs and is
highlighted with an asterisk. These metabolic genes were assigned to 11 different
metabolic categories based on the KEGG database (Supplementary Dataset 7).
Different metabolic categories based on KEGG are indicated in bold next to the
corresponding modules.

Figure 6 Metabolic potential and horizontal gene transfer (HGT) landscape of
Nitrososphaerota-associated viruses and mobile genetic elements (MGEs). a,
Metabolic genes encoded by viral/MGE genomes but not detected in their
corresponding hosts. These metabolic genes were assigned to 11 different
metabolic categories based on the KEGG database (Supplementary Dataset 7). b,
Proportion of HGT events with different transfer directions among various
archaeal lineages. c, Relationships among HGT event direction, archaeal lineages,
and viral/MGE families. The number of events is indicated in parentheses. d,
Relationships among HGT event direction, viral function, cellular function, and
elements-related metabolism.

a

2,137 genomes and Nitrososphaerota MAGs from GenBank and IMG/M → 1. CRISPR prediction ← 158 metagenomic samples from thermal springs/hydrothermal vents/deep sea

MinCED, CRASS, CRISPR-Cas-Finder, CRISPR-Cas-Identifier

MMSeqs2

55,995 DRs from Nitrososphaerota-associated contigs → 234,036 spacers of Nitrososphaerota

geNomad, VirSorter2, DeepVirFinder

2.1. Proviral prediction ← IMG/VR (v.4) → 2.2. Viral/MGE prediction ← 2.3. Viral identification

MMSeqs2

geNomad, MMSeqs2

3. Genome filtering

471 vOTUs without predicted capsid-related proteins (possible viruses and other MGEs)

MMSeqs2

746 of Nitrososphaerota-associated viruses/MGEs

MMSeqs2

MMSeqs2

153 vOTUs with predicted capsid-related proteins (viruses, expanded by sVCs)

4. vOTU definition → 624 vOTUs

b

Number of *Nitrososphaerota*-associated viral/MGE vOTUs

- Kim et al., study: 5
- Krupovic et al., study: 1
- Ahlgren et. al. study: 3
- Zhou et al., study: 5
- IMG/VR v.4 (Confirmed from this study): 275
- Metagenomes-derived (Recalled from this study): 336

c

Habitats

- Geothermal springs: 233
- Marine: 190
- Soil: 103
- Hydrothermal vents: 76
- Freshwater: 22
- Sweage: 2

a

sVC_0003 sVC_0012 sVC_0036 sVC_0002 sVC_0015 sVC_0039 sVC_0019 sVC_0024 sVC_0044 sVC_0004 sVC_0035

Host lineage

Group 1.1a — Group 1.1b
Nitrosocaldales — D-B Thaum
Caldarchaeales — D-B NitroA
Nitrosotales — MT/GW
MBG-A — *Wolframiiraptoraceae*
pSL4-affiliation

sVC_0052 sVC_0047 sVC_0041 sVC_0040 sVC_0038 sVC_0037 sVC_0034

sVC_0008 sVC_0030 sVC_0043 sVC_0027 sVC_0032 sVC_0001 sVC_0028 sVC_0025 sVC_0045 sVC_0031 sVC_0023 sVC_0020

Morphology

◯ Head-tail
◇ Filamentous
⬡ Icosahedral
△ Spindle

sVC_0029 sVC_0013 sVC_0014 sVC_0010

sVC_0068 sVC_0054 sVC_0051 sVC_0049 sVC_0046 sVC_0042 sVC_0033 sVC_0007 sVC_0069 sVC_0067 sVC_0061 sVC_0053

Shared protein clusters (%)

— 97   — 40
— 80   — 20
— 60

sVC_0032 sVC_0026 sVC_0021 sVC_0005 sVC_0018 sVC_0066 sVC_0060 sVC_0059 sVC_0022 sVC_0058 sVC_0061 sVC_0057

b

I. II. III. IV. V. Tree scale: 0.1 |—| scale 0.05

(proposed) Family | Morphology | Average shared PCs (%)

Ekchuahviridae
sVC_0044

MGE_F005
sVC_0019

Sthaneeviridae
sVC_0043

MGE_F008
sVC_0045

MGE_F015
sVC_0039

Itzamnaviridae
sVC_0042

MGE_F003
sVC_0001

MGE_F010
sVC_0024

MGE_F001
sVC_0015

MGE_F006
sVC_0003

Yebviridae
sVC_0003

MGE_F014
sVC_0035

Chaac-Thaspi
sVC_0017/00068

MGE_F017
sVC_0061

MGE_F012
sVC_0008

MGE_F019
sVC_0051

Byatisviridae
sVC_0012

MGE_F011
sVC_0021

Ommogviridae
sVC_0013

Ythogthaviridae
sVC_0005

MGE_F009
sVC_0032

MGE_F016
sVC_0027

Yangangviridae
sVC_0069

MGE_F002
sVC_0036

MGE_F007
sVC_0004

MGE_F004
sVC_0025

MGE_F018
sVC_0054

MGE_F013
sVC_0026

Cthughaviridae
sVC_0002

Fthagguaviridae
sVC_0002

I. Genome length (kb)
10 — 55 — 100

II. G+C content (%)
25 — 45 — 60

III. Coding density (%)
75 — 85 — 95

IV. Habitats
Hydrothermal vents
Marine
Geothermal springs
Soil
Freshwater

V. Host lineage
Group 1.1a
Group 1.1b
MT/GW
Nitrosocaldales
MBG-A
D-B Thaum
Wolframiiraptoraceae
Caldarchaeales
D-B NitroA

Morphology
Head-tailed
Spindle-shaped
Rod-shaped
Icosahedral

Pentose and glucuronate interconversions

5-Dehydro-4-deoxy-D-glucuronate

(4S)-4,6-Dihydroxy-2,5-dioxohexanoate

2-Dehydro-3-deoxy-D-gluconate

2-Dehydro-3-deoxy-D-gluconate-6P

kduD

Pentose phosphate pathway; Starch and sucrose metabolism; Carbon metabolism

D-Ribose 5-phosphate   PRPP

D-Ribulose 5-phosphate   CO2

6-Phospho-D-gluconate

D-Glucono-1,5-lactone 6-phosphate

pgl

H6PD

β-D-Glucose 6-phosphate

pgi1

α-D-Glucose 6-phosphate

pgi1

D-Fructose 6-phosphate

Lipoarabinomannan (LAM) biosynthesis

α12mnt   *aftB*

Glycosylphosphatidylinositol (GPI)-anchor biosynthesis

PIGO

1-Phosphatidyl-D-myo-inositol

Photophosphorylation

ATP   ADP + P

petF

Fatty acid metabolism

Malonyl-[acp]

ACP

Malonyl-CoA

ACCd

Acetyl-CoA   Glyoxylate

(4R)-4-Hydroxy-2-oxoglutarate   Proline

P4HA

4-Hydroxyproline

Arginine and proline metabolism

3-Sulfinyl-pyruvate   L-Cysteine

Pyruvate   *dcyD*   D-Cysteine   L-Valine   δ-(L-2-Aminoadipyl)-L-cysteinyl-D-valine

PCBC   Isopenicillin

Penicillin and cephalosporin biosynthesis

Penicillin

Cysteine, methionine, D-Amino acid metabolism

Glycine, serine and threonine metabolism

Choline

Betaine aldehyde   *sdmt*

Dimethylglycine   Sarcosine

Porphyrin, cofactors metabolism

5-Amino levulinate   Glycine   G

Precorrin 8X   Precorrin 2   Uroporphyrinogen III

Hydrogenobyrinate   Sirohydrochlorin   Cobalt-precorrin 6B

Hydrogenobyrinate diamide   *cbiT*   Cobalt-precorrin 7

Cob(II)yrinate diamide   *cobS*   Coenzyme B12   7,8-Dihydroneopterin 3'-triphosphate

α-Ribazole   *GCH1*

Dimethylbenzimidazole   GTP

FMN   Riboflavin

Glycosaminoglycan biosynthesis

NDST1   [Protein]-L-serine

HS2ST1   Chain elongation   *CHSY*

HS3ST2   Chondroitin

Dermatan   Glycosaminoglycan degradation

PAP   PAP

Sulfur metabolism

Sulfide

Sulfite

PAPS

Sulfate   *cysD*   Purine metabolism   ADP   ApppppA

Adenylyl sulfate   ATP

3'-Phosphoadenylyl sulfate   Sulfazecin

MM 42842

Monobactam biosynthesis

Arginine biosynthesis

Glutamine

NH3

Glutamate   *argA*   N-Acetyl-L-glutamate   Urea cycle

Steroid hormone biosynthesis

16α- Hydroxyestrone

SULT1E1

Estrone   Estrone 3-sulfate

Glutathione biosynthesis

5-Oxoproline   Glutathione

gshA

L-Glutamate   L-γ-Glutamylcysteine

Glyoxylate and dicarboxylate metabolism

N-Formylderivatives

dmfA2

Formate

CO2

Glycerophospholipid metabolism

CDP-glycerol

Acyl-CoA

sn-Glycerol 3-phosphate   *tagD*

CDP-diacylglycerol

1,2-Diacyl-sn-glycerol 3-phosphate

Phosphatidyl-L-serine   *psd*

Phosphatidylethanolamine

Glycerophospholipid metabolism

Triacylglycerol

lip   1,2-Diacyl-sn-glycerol

lip   Monoglyceride

fatty acid

Pyrimidine metabolism

dUMP   dCMP

dut   *gp42*

dUDP   dUTP   hm5dCMP

hm5dCDP

Staurosporine biosynthesis

L-Tryptophan

Rebeccamycin   *prnA*

7-Chloro-L-tryptophan

Pyrrolnitrin

Lipopolysaccharide metabolism

KDO2-lipid A   ADP-L-glycero-β-D-manno-heptose

inner core oligosaccharide   *waaF*

Mature LPS   *waaJ*

Polyketide sugar unit biosynthesis

D-Glucose 1-phosphate

dTDP-D-glucuronate   dTDP-glucose   dTDP-4-amino-4,6-dideoxy-D-glucose

calS13

dTDP-4-oxo-2-deoxy-β-L-xylose

Steroid hormone biosynthesis

7,8-Dihydroneopterin 3'-triphosphate

6-Carboxy-5,6,7,8-tetrahydropterin

7-Carboxy-7-carbaguanine   *queE*

7-Aminomethyl-7-carbaguanine

Wolframiiraptoraceae   *Caldarchaeales*   Group 1.1b   Group 1.1a   ★ Compensation from viruses/MGEs