Beak elongation as key innovation in shorebird adaptive radiation

Michel Baguette[1,2*†], Glenn Le Floch[1†], Eleanor Lacassagne[1], Vincent Bels[1], Nicolas Schtickzelle[3]

1. Muséum National d'Histoire Naturelle, Institut de Systématique, Évolution, Biodiversité, UMR 7205 Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France

2. CNRS, Station d'Écologie Théorique et Expérimentale, UAR 2029, Moulis, France

3. UCLouvain, Earth and Life Institute, Louvain-la-Neuve, Belgium

* Corresponding author

† Co-first authors

Abstract

Adaptive radiations often hinge on the emergence of key innovations - traits that unlock new ecological opportunities and drive lineage diversification. In shorebirds (Charadriiformes), variation in beak length has long been associated with resource partitioning, yet its role as a functional innovation has remained untested. Here, we integrate behavioral, morphological, and phylogenetic data from 29 Western Palearctic shorebird species to test whether beak elongation acted as key innovation. We show that species with longer than expected beaks (relative to body size) consistently exhibit distinct foraging and food transport mechanisms. These morphological-behavioral syndromes occur across independent lineages, suggesting convergent evolution driven by ecological constraints. Phylogenetic analyses reveal independent origins of beak elongation, reinforcing its role as a repeated evolutionary solution. Our findings support a functionally integrated view of adaptation, where morphological change enables novel behaviors and ecological exploitation, catalyzing adaptive radiation. This study provides a testable framework for identifying key innovations and underscores the importance of coordinated trait evolution in shaping biodiversity.

One sentence summary

Beak elongation in shorebirds drives behavioral innovation and opens access to previously unreachable ecological niches.

Introduction

Adaptive radiation refers to the evolutionary emergence of phenotypic diversity within a lineage undergoing diversification (*1,2*). This process often depends on the availability of novel ecological niches, which may be accessed through the emergence of a key innovation - that is, a phenotypic modification enabling individuals to exploit previously inaccessible ecological opportunities (*3*). While the role of key innovations in driving adaptive radiation has been widely acknowledged, Miller et al. (*4*) argue that the concept has gradually shifted in meaning. Specifically, they note a conflation between lineage diversification (e.g., *5*) and trait-dependent ecological transitions (e.g., *6*). In response, Miller et al. (*4*) advocate for a return to a more ecologically grounded definition of key innovations. They propose two complementary approaches to operationalizing this concept, by elucidating the mechanisms through which phenotypic traits alter the interaction between organisms and their environments, and/or demonstrating that these trait changes lead to measurable shifts in the organism's ecological niche.

In this study, we adopt the second of the two approaches proposed by Miller et al. (*4*), focusing on identifying phenotypic changes that may have facilitated adaptive radiation in shorebirds. Shorebirds represent a relatively homogeneous group of approximately 215 species (*7*), encompassing two of the three suborders within the order Charadriiformes: Scolopaci (e.g., shanks, tattlers, sandpipers, godwits, snipes, curlews) and Charadrii (e.g., plovers, avocets, oystercatchers, lapwings). A recent time-calibrated phylogeny - integrating genetic, morphological, and paleontological data - provides a robust framework for understanding the remarkable species diversity found within the Charadriiformes (*8*). Shorebirds constitute one of the most prominent avian radiations, not only in terms of species richness but also with respect to morphological, behavioral, and biogeographical variation (*7,9*). They are frequently cited as a textbook example of resource partitioning, wherein contrasting morphological traits - particularly beak and leg structures - facilitate the localization, capture, and ingestion of food items. During the non-breeding season, most shorebirds predominantly forage along land - water interfaces in freshwater, brackish, or marine environments, where they feed primarily on invertebrates, but also on small vertebrates and plant material. It is widely assumed that interspecific variation in beak length allows different shorebird species to access prey located at different sediment or water column depths, while variation in leg length permits foraging across a range of water depths. However, the extent to which these trait divergences represent true key innovations - as opposed to post hoc "just-so stories" (*10*) grounded more in narrative plausibility than in rigorous empirical testing of adaptive hypotheses (*11*) - remains an open question.

In a previous study, we described and classified the food acquisition behaviors of 26 western Palearctic shorebird species during migration and wintering periods (Figure 1 in [*12*]). Among these, 8 species belonged to the suborder Charadrii and 18 to Scolopaci. This work revealed the existence of a behavioral syndrome associated with food acquisition, characterized by covariation among stereotyped behaviors used during each of the three fundamental steps of foraging: food detection, capture, and transport. Furthermore, our findings suggested a phylogenetic component underlying this behavioral covariation. We also demonstrated that these three behavioral steps constitute ecological performances in the sense (*13*), i.e., quantifiable traits that reflect how effectively an individual performs tasks critical to its fitness.

In the present study, we expand our behavioral dataset by incorporating information from three additional Scolopaci species. In parallel, we compiled morphological data related to body structure (body mass, body length, and the size of musculoskeletal appendages), sensory organs (eye diameter and the number of sensory pits on the beak - serving as proxies for visual and tactile foraging abilities, respectively), as well as beak and tongue lengths. To develop a standardized morphological baseline, we first combined the body structure traits (excluding beak length) to construct a global body size index for each of the 29 species, using a phylogenetically informed approach. We then quantified allometric relationships between this index and key morphological traits involved in food acquisition - specifically, sensory structures, beak, and tongue - representing their roles in food detection, capture, and transport, respectively. Using these relationships, we calculated Relative-to-Expected (RTE) trait values, defined as species-specific deviations of a given morphological trait from its predicted value based on the global body size index. To explore potential differences in evolutionary trajectories between the two shorebird suborders, we compared RTE trait values between Charadrii and Scolopaci. Lastly, we investigated whether, and in what manner, the three components of the food acquisition behavioral syndrome - locomotion, prey capture, and transport - are associated with deviations in RTE beak length across species.

Materials and Methods

Phenotypic traits

We collected data on the food acquisition behavior of 29 species in wintering or migration sites (12,44), which represents ca. 75% of the total shorebird fauna of the western Palearctic (47, except for vagrants), and ca. 90% of the species using land - water interfaces during migration and overwintering (47, except for vagrants) . Locomotion behaviors (how individuals move around their feeding habitats), capture behaviors (how individuals catch and locate their prey) and transport behaviors (how individuals move food particles from the tip of the beak to the pharynx) were categorized based on a recent repertoire (12). There are five capture behaviors (pecking, probing, pecking-probing, sweeping, routing), three locomotion behaviors (run-stop-run, continuous walking, continuous walking-continuous swimming), and three transport behaviors (lingual, ballistic and ballistic-surface tension).

Aggregated data at the species level were used for morphological variables (beak length, body size, weight, wing length, tarsus length, length of the longest toe, tongue length) and sensory organs (diameter of the eye, number of sensory pits on the beak). Morphological data come from Cramp & Simmons (47), except tongue lengths, which were published by Burton (26), diameters of the eye that come from (21) and the number of sensory pits on the beak that were measured by one of us (EL) on the available specimens of the Compared Anatomy Lab of the Muséum National d'Histoire Naturelle (Paris). The values used are averages between sexes, and when values from different geographical origins were available, we used those measured in the localities closest to our behavior observation sites. Four variables were available for a subset of species only: eye diameter (22 species), number of sensory pits (9 species) and tongue length (19 species). Two variables were transformed before analysis: weight was cubic root transformed to account for its scaling as a volume, not a length; tongue length was expressed as a proportion of the beak length and this tongue/beak length ratio was ln transformed to make its distribution symmetric.

Statistical analyses

First, we summarized the variation among the 29 species in five morphological traits (body size, wing length, tarsus length, length of the longest toe, and cubic root of weight) using a phylogeny constrained Principal Component Analysis (PCA; *phyl.pca* function from *phytools* R package) (14,15) to control for the effect of evolutionary relatedness among species. Phylogenetic distances used in this PCA were inferred using a pruned tree restricted to our study species from the Charadriiformes time tree of Černý & Natale (8). The first two Principal Components (PCs) were used to place the species in a phylomorphospace. PC1 summarizes a very high proportion (80%) of the morphological variation and can be interpreted as a phylogenetically controlled global body size index because it is highly and positively correlated with all the five morphological traits.

Second, we used this phylogenetically controlled global body size index to correct morphological traits of interest (eye diameter, number of sensory pits, beak length, and tongue/beak length ln-ratio) for allometry. A linear regression between the trait and the body index was estimated (*lm* function from *stats* R package), and a new Relative To Expected (RTE) trait variable was created to express how much a species possesses a trait value that differs from its expected value given its body size index; RTE trait for a species was computed as the ln transformed ratio between its residual value and its predicted value, both from the allometric regression. We tested whether species from the two suborders significantly differ for these four RTE traits using two sample t-tests.

Third, we checked for specific relationships among morphological traits and behaviors associated to the three main steps of food acquisition (locomotion, capture and transport) using appropriate linear models according to the nature of the response variable: linear regression (*lm* function from *stats* R package) for a continuous normal variable, binomial logistic regression (*glm* function from *stats* R package) for a two levels discrete variable, and binomial logistic regression (*vglm* function from *VGAM* R package) for an ordered multilevel discrete variable.

Results

We constructed a phylomorphospace (Figure 1) by plotting species scores along the first two axes of a phylogenetically constrained Principal Component Analysis (pPCA) (*14, 15*), which summarizes interspecific variation in five morphological traits across the 29 studied shorebird species. The first principal component (PC1), which accounts for 80% of the total morphological variation, was interpreted as a phylogenetically controlled global body size index. Along this axis, species range from the smallest plovers and sandpipers to the largest taxa, such as the avocet, stint, godwit, and curlews, with the largest species being 3.8 times larger than the smallest.

Fig. 1. Shorebird phylomorphospace.

We used the phylogenetic tree of the 29 shorebirds species (**A**) to perform a phylogenetic constrained Principal Component Analysis (pPCA) on five morphological traits and create a phylomorphospace (**B**) using its first two principal components (PC1 and PC2). PC1 summarizes 80% of the variance (**C**) and can be interpreted as a phylogenetically constrained global body size index, being highly and positively correlated with the five traits (**D**) and (**E**). 14 species are represented on the phylomorphospace (**B**) at the same scale (bird silhouettes from phylopic.org; detailed credits in the R script provided in the supplementary material).

Among the traits analyzed, eye diameter increases significantly with body size. After correcting for this allometric relationship, relative eye diameter (RTE) is significantly larger in Charadrii than in Scolopaci (Figure 2A). In contrast, the number of sensory pits on the beak decreases significantly with body size and, when corrected for allometry, is significantly higher in Scolopaci than in Charadrii (Figure 2B). A negative correlation between the two RTE-corrected sensory traits indicates that species with fewer sensory pits tend to have relatively larger eyes (Figure 3). Beak length scales positively with body size, and once corrected for allometry, species in Charadrii have significantly shorter beaks than those in Scolopaci (Figure 2C). Additionally, the ratio of tongue length to beak length shows a significant negative scaling relationship with body size: larger species possess proportionally shorter tongues, although not necessarily in absolute length. However, after correcting for this allometric effect, no significant difference in tongue-to-beak length ratio is observed between the two suborders (Figure 2D).

Fig. 2. Allometric relationships.
We use allometric relations (first line, with linear regression statistics) with the phylogenetically controlled global body size index to correct four morphological traits for the species body size by computing their RTE (Relative-to-Expected) counterparts. We then check whether these RTE traits differ between the two suborders, Charadrii and Scolopaci (second line, with t-test statistics). The four morphological traits are the eye diameter (A), the number of sensory pits on the beak (B), the beak length (C) and the tongue length in proportion of the beak, expressed as the ln-ratio of tongue over beak length (D). Sample size for each trait is indicated in panel titles.

Fig. 3. Relationship between eye diameter and sensory pits.
Species with a smaller eyes have more sensory pits on their beak. Both variables are corrected for their allometric relation with body size, being expressed as Relative-to-Expected (RTE) values. Statistics refer to a linear regression.

Relative-to-Expected (RTE) beak length is a significant predictor of behavioral strategies employed during food acquisition. Species with longer than expected beaks (relative to their body size index) are more likely to exhibit: (1) continuous walking rather than stop-run-stop locomotion (Figure 4A); (2) probing rather than pecking as a prey capture technique, with intermediate species exhibiting a combination of both behaviors (Figure 4B); and (3) ballistic rather than lingual mechanisms for food transport, with transitional species employing both surface tension and ballistic strategies (Figure 4C). In contrast, the relative tongue length (corrected for beak length) does not significantly influence the mode of food transport behavior (Figure 4D).

Fig. 4. Relationship between beak/tongue length and food acquisition behaviors.
Locomotion (A), prey capture (B) and food transport (C) behaviors used by a species significantly depends on how longer than expected its beak is given its body size index. Transport behavior is however not influenced by how longer its tongue (relative to beak length) is (D). Beak length and the ration tongue/beak length are corrected for their allometric relation with body size, being expressed as Relative-to-Expected (RTE) values. Continuously walking (CW) and stop-run-stop (SRS) are the main two locomotor modes used by shorebirds (*12,20*). Statistics refer to binomial (A) , cumulative (B) or multinomial (C and D) logistic regressions.

Discussion

Outside the breeding season, shorebird species tend to aggregate in large numbers within suitable foraging habitats, predominantly along coastal and shoreline environments (e.g.,*7,16*). On overwintering grounds, food biomass is limited (*17*); moreover, for prey items to be considered profitable, they must accessible, detectable, manageable (i.e., handleable and ingestible), and digestible (*16*). Consequently, interspecific food partitioning within shorebird communities is commonly observed (e.g., *18*). Recent advances in stable isotope analysis and DNA metabarcoding have provided important insights into the dietary composition of co-occurring shorebird species (*18,19*). These studies reveal that species often exhibit local and temporal dietary specializations, frequently relying on specific prey types, and developing cues and behaviors tailored to exploiting those prey effectively. A substantial body of research grounded in optimal foraging theory has demonstrated that foraging shorebirds tend to maximize net energy gain by optimizing prey selection, handling, and ingestion strategies (e.g.,*7,16*). However, despite these insights into behavioral decision-making, questions remain regarding the evolutionary processes that have shaped the morphological and behavioral traits underlying such ecological specialization.

Sensory organs play a critical role in the successful detection of food. Shorebirds employ two primary strategies for locating prey or plant material: visual detection (hunting by sight) and tactile detection (hunting by touch), with some species potentially combining both approaches (e.g., *12,16,20-24*). In the present study, we use eye diameter and the number of sensory pits on the beak as proxies for a species' relative investment in visual and tactile prey detection, respectively. Across the 29 species in our dataset, eye diameter scales positively with the global body size index. Notably, six of the eight Charadrii species exhibit eye diameters that are - sometimes substantially - larger than expected based on this allometric relationship. According to the dated phylogeny by Černý and Natale (*8*), our Charadrii sample includes five species from the Charadriidae family (plovers and lapwing), which represents an early-diverging clade within the suborder, and three species from more recently diverged lineages. The six species with larger-than-expected eye diameters comprise the five Charadriidae representatives and the oystercatcher, suggesting that enlarged eye size is a retained trait within the early-diverging clade.

In contrast, the number of sensory pits on the beak shows a significant negative relationship with the global body size index. However, sensory pit data are lacking for many species, particularly within Charadrii, for which data are available for only two species: the lapwing (representing the ancient clade) and the oystercatcher (a representative of the more recently diverged clade). Despite these data limitations, the observed differences in sensory organ morphology between Charadrii and Scolopaci are broadly consistent with their known prey capture strategies. Most Charadrii, especially those from the ancient clade, exhibit larger than expected eye diameters and rely predominantly on visual prey detection. In contrast, Scolopaci species, which locate prey by touch or through a combination of tactile and visual cues, possess more sensory pits on their beaks than expected given their body size. Furthermore, although based on a limited

sample, our findings suggest a potential trade-off in sensory investment: species with larger-than-expected eye diameters tend to exhibit a lower than expected number of sensory pits, implying a specialization towards either visual or tactile foraging, but rarely both

In our dataset, beak length exhibits a positive allometric relationship with the global body size index, indicating that larger shorebird species generally possess longer beaks. However, after correcting for this allometric effect, species belonging to the suborder Charadrii display significantly shorter than expected beak lengths compared to those in Scolopaci. This pattern is primarily driven by the five species within the family Charadriidae (plovers and lapwing), the early-diverging clade within Charadrii. In contrast, species from the more recently diverged (Haematopodidae + Recurvirostridae) clade (e.g., oystercatchers, avocets, and stints) exhibit relative beak lengths comparable to those of Scolopaci species.

The tongue plays a crucial role in the evolution of food transport mechanisms in Sauropsida, including birds (25). Among shorebirds, tongue length varies markedly, ranging from nearly equal to beak length to less than one-third of it (26). To account for this diversity, we expressed tongue length as a proportion of beak length and found that this ratio scales negatively with body size. These findings suggest that tongue elongation is subject to an upper constraint that prevents proportional increases in length relative to the beak, possibly due to biomechanical or physiological trade-offs. Such constraints may be linked to the evolution of specific food transport mechanisms, such as ballistic transport (25,27,28) or surface tension-based transport (25,29-31).

In previous work, we decomposed shorebird food acquisition behavior into three sequential stages: foraging, feeding, and swallowing (12). Foraging encompasses locomotion and prey capture - respectively focused on locating and seizing food items - while feeding refers to the manipulation and transport of food toward the beak. Swallowing marks the final stage of food acquisition, when food is ingested into the pharynx. Locomotion, capture, and transport are three performances (13) that can be further subdivided into three to four stereotyped behaviors, following Tinbergen's hierarchical model of behavioral complexity (32). These behaviors are readily observable and quantifiable under field conditions (12).

We previously identified a behavioral syndrome in shorebirds characterized by consistent associations among stereotyped behaviors of each of these performances (12). Pecking is considered the plesiomorphic prey capture behavior in shorebirds (33,34), and all five Charadriidae species in our dataset exhibit a behavioral profile characterized by stop-run-stop locomotion, pecking as the primary capture strategy, and lingual transport of food particles (12). In contrast, species from the Scolopaci suborder and from the more recently diverged Charadrii clade (Haematopodidae + Recurvirostridae) employ a mixed capture strategy combining pecking and probing, use continuous walking as their primary locomotion behavior, and - except for one species - utilize a combination of ballistic and surface tension-based food transport strategies.

Here, we demonstrate that this behavioral syndrome aligns closely with differences in relative beak length between the ancient and more derived clades. Species with shorter than expected beaks relative to body size consistently employ the behavioral combination of stop-run-stop locomotion, pecking capture, and lingual transport. Conversely, species with longer than expected beaks are more likely to exhibit probing or combined pecking-probing behaviors, continuous walking, and employ ballistic or surface tension transport mechanisms. These findings support the hypothesis that morphological divergence in beak length has co-evolved with functional behavioral adaptations that optimize food acquisition strategies within distinct ecological contexts.

In particular, the convergence of morphological and behavioral phenotypes observed between long-beaked Scolopaci and the recently diverged long-beaked Charadrii further corroborates the notion that beak elongation has been a key driver facilitating the evolution of novel locomotion, prey capture, and food transport behaviors.The beak, serving as the primary grasping organ in birds and often analogized to the primate hand (35), plays a fundamental role in food acquisition (36). However, studies examining the relationship between beak morphology and feeding ecology across modern birds as a whole have yielded inconclusive results (37). Two principal factors likely account for this lack of a clear relationship: first, similar trophic resources may be exploited via markedly different beak morphologies; second, beak morphology may be shaped by selective pressures unrelated to feeding, such as communication, nest construction, preening, or thermoregulation (38). These confounding factors can be mitigated by focusing analyses at lower taxonomic levels (37), as we do here, and as previously demonstrated by Olsen (38), who found feeding ecology to be the primary driver of beak diversification within Anseriformes.

Our data set encompasses the vast majority of species that utilize land-water interfaces during migration and overwintering in the Western Palearctic. During this period, when many individuals concentrate in small patches of suitable habitat, resources become particularly difficult to access, intensifying competition and highlighting functional differences among species. Moreover, much of the diversification within shorebird genera is thought to have occurred in response to Quaternary climatic oscillations (7), whereas the evolutionary emergence of elongated beaks is likely far older. It is noteworthy that an ancestor resembling a shorebird, existing around the Cretaceous–Paleogene boundary approximately 65 million years ago, is widely considered foundational to the radiation of modern birds (e.g., 39–42). This so-called "transitional shorebird" is characterized by a short beak (39, 40). From this perspective, we propose that the elongation of the beak - by granting access to a broader array of food resources - constitutes a key innovation in shorebirds, thereby constraining the subsequent evolution of other phenotypic traits such as body design, sensory organs, and foraging behaviors. Consequently, the adaptive radiation of modern shorebird species and their phenotypic diversification may have been initiated by this evolutionary lengthening of the beak. Although the proportion of species included in our study may seem limited compared to the ca. 215 recognized shorebird species worldwide (7), here we capture both the ecological context in which competitive interactions are strongest and the evolutionary context in which morphological innovations have long preceded more recent taxonomic diversification.

One might question whether this hypothesis is merely a "just-so" story, as criticized by Gould and Lewontin (*11*) for many adaptive explanations. The classical hypothetico-deductive experimental approach cannot be applied here, since historical evolutionary events cannot be recreated or directly tested (*43*). As in paleontology, where trait evolution is inferred from fossils and comparative data, only correlative approaches are feasible to assess historical scenarios. Although correlative, our results are supported by several indirect lines of evidence. Firstly, the repeated association of beak elongation with distinct behavioral syndromes across independent lineages suggests convergent selection pressures rather than coincidental trait combinations. Secondly, these morphological-behavioral syndromes correspond to ecological differentiation, with longer-beaked species foraging at greater water depths (*44*), consistent with character displacement reducing interspecific competition. Comparative diet analyses could provide further insight into niche differentiation linked to beak length, but the pronounced dietary plasticity of shorebirds (*18, 45, 46*) limits the effectiveness of such studies. This plasticity also constrains the use of species assemblage analyses and niche modeling aimed at detecting adaptive peaks in beak length variation.

Altogether, our findings provide empirical support for the idea that morphological innovations such as beak elongation can drive the emergence of novel behavioral repertoires, thereby opening new ecological opportunities. This functional integration between structure and behavior highlights a broader principle in evolutionary ecology: innovations are rarely isolated events, but rather part of coordinated phenotypic transitions. While the concept of "key innovations" is often invoked post hoc, our results point to a specific sequence - elongation of the beak, followed by shifts in sensory investment and feeding mechanics - as plausible evolutionary pathway underpinning niche differentiation in shorebirds. This underscores the need to consider behavioral and sensory traits in tandem with morphology when evaluating the drivers of adaptive radiation. Furthermore, the convergence observed between unrelated long-beaked lineages suggests strong selective constraints and offers a rare example of replicated evolutionary solutions within a single avian order.

References

1. Schluter D. *The Ecology of Adaptive Radiation* (Oxford Univ. Press, Oxford, 2000).

2. Gavrilets S., Losos J. B. Adaptive radiation: contrasting theory with data. *Science* **323**, 732–737 (2009).

3. Stroud J. T., Losos J. B. Ecological opportunity and adaptive radiation. *Annu. Rev. Ecol. Evol. Syst.* **47**, 507–532 (2016).

4. Miller A. H., Stroud J. T., Losos J. B. The ecology and evolution of key innovations. *Trends Ecol. Evol.* **38**, 122–131 (2023).

5. Simões M., Breitkreuz L., Alvarado M., Baca S., Heins L., Herzog K., Lieberman BS. The evolving theory of evolutionary radiations. *Trends Ecol. Evol.* **31**, 27–34 (2016).

6. Baguette M., Bertrand J. A., Stevens V. M., Schatz B. Why are there so many bee-orchid species? Adaptive radiation by intra-specific competition for mnesic pollinators. *Biol. Rev.* **95**, 1630–1663 (2020).

7. Colwell M. A. *Shorebird Ecology, Conservation and Management* (Univ. of California Press, Berkeley, 2010).

8. Černý D., Natale R. Comprehensive taxon sampling and vetted fossils help clarify the time tree of shorebirds (Aves, Charadriiformes). *Mol. Phylogenet. Evol.* **177**, 107620 (2022).

9. Van Tuinen M., Waterhouse G. J., Dyke G. J. Avian molecular systematics on the rebound: a fresh look at modern shorebird phylogenetic relationships. *J. Avian Biol.* **35**, 191–194 (2004).

10. Kipling R. *Just So Stories* (Macmillan & Co. Ltd, London, 1902).

11. Gould S. J., Lewontin R. C. The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proc. R. Soc. Lond. B Biol. Sci.* **205**, 581–598 (1979).

12. Baguette M., Le Floch G., Hannier L., Kirchhoff F., Schtickzelle N., Stevens V.M., Bels V. Repertoire of food acquisition behaviors in Western Palearctic shorebirds (Aves, Charadriiformes). *Front. Ethol.* **3**, 1351994 (2024).

13. Irshick D. J., Higham T. E. *Animal Athletes: An Ecological and Evolutionary Approach* (Oxford Univ. Press, Oxford, 2016).

14. Revell L. J. Size correction and principal components for interspecific comparative studies. *Evolution* **63**, 3258–3268 (2009).

15. Revell L. J. phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

16. van der Kam J., Ens B., Piersma T., Zwarts L. *Shorebirds: An Illustrated Behavioural Ecology* (KNNV Publishers, Utrecht, 2004).

17. Lourenço P. M., Granadeiro J. P., Catry T. Low macroinvertebrate biomass suggests limited food availability for shorebird communities in intertidal areas of the Bijagós archipelago (Guinea-Bissau). *Hydrobiologia* **816**, 197–212 (2018).

18. Catry T., Lourenço P.M., Lopes R.J., Carneiro C., Alves J.A., Costa J., Rguibi-Idrissi H., Bearhop S., Piersma T., Granadeiro J.P. Structure and functioning of intertidal food webs along an avian flyway: A comparative approach using stable isotopes. *Funct. Ecol.* **30**, 468–478 (2016).

19. Correia E., Granadeiro J.P., Santos B., Regalla A., Mata V.A., Catry T. Trophic ecology of a migratory shorebird community at a globally important non-breeding site: Combining DNA metabarcoding and conventional techniques. *Mar. Ecol. Prog. Ser.* **705**, 127–144 (2023).

20. Barbosa A., Moreno E. Evolution of foraging strategies in shorebirds: An ecomorphological approach. *Auk* **116**, 712–725 (1999).

21. Thomas T., Székely R. F., Cuthill I. C. Eye size, foraging methods and the timing of foraging in shorebirds. *Funct. Ecol.* **20**, 157–165 (2006).

22. Lourenço P. M., Catry T., Piersma T., Granadeiro J. P. Comparative feeding ecology of shorebirds wintering at Banc d'Arguin, Mauritania. *Estuaries Coasts* **39**, 855–865 (2016).

23. du Toit C. *Mechanosensory Structures in the Beaks of Probe-Foraging Birds in Relation to Their Foraging Ecology* (PhD thesis, Univ. of Cape Town, 2021).

24. Martin G. R. *The Sensory Ecology of Birds* (Oxford Univ. Press, Oxford, 2017).

25. Bels V., Pallandre J.-P., Pelle E., Kirchhoff F., Gastebois G., Davenport J., Baguette M. Food transport in Reptilia: A comparative viewpoint. *Philos. Trans. R. Soc. B* **378**, 20220542 (2023).

26. Burton P. J. K. *Feeding and Feeding Apparatus in Waders* (British Museum of Natural History, London, 1974).

27. Baussart S., Korsoun L., Libourel P. A., Bels V. Ballistic food transport in toucans. *J. Exp. Zool. A Ecol. Genet. Physiol.* **311**, 465–474 (2009).

28. Harte M., et al. Ballistic food transport in birds: The example of *Casuarius casuarius*. *Comput. Methods Biomech. Biomed. Eng.* **15**, 137–139 (2011).

29. Rubega M. A., Obst B. S. Surface-tension feeding in phalaropes: Discovery of a novel feeding mechanism. *Auk* **110**, 169–178 (1993).

30. Rubega M. A. Surface tension prey transport in shorebirds: How widespread is it? *Ibis* **139**, 488–493 (1997).

31. Prakash M., Quéré D., Bush J. W. M. Surface tension transport of prey by feeding shorebirds: The capillary ratchet. *Science* **320**, 931–934 (2008).

32. Tinbergen N. *The Study of Instinct* (Pygmalion Press, 2020).

33. Zweers G. A. Transformation of avian feeding mechanisms: A deductive method. *Acta Biotheor.* **39**, 15–36 (1991).

34. Zweers G. A., Gerritsen A. F. C. Transitions from pecking to probing mechanisms in waders. *Neth. J. Zool.* **47**, 167–207 (1997).

35. Bhullar B. A., Hanson M., Fabbri M., Pritchard A., Bever G.S., Hoffman E. How to make a bird skull: Major transitions in the evolution of the avian cranium, paedomorphosis, and the beak as a surrogate hand. *Integr. Comp. Biol.* **56**, 389–403 (2016).

36. Xu Y., Price M., Que P., Zhang K., Sheng S., He X., Wen Z., Wang B. Ecological predictors of interspecific variation in bird bill and leg lengths on a global scale. *Proc. R. Soc. Lond. B Biol. Sci.* **290**, 20231387 (2023).

37. Navalón G., Bright J. A., Marugán-Lobón J., Rayfield E. J. The evolutionary relationship among beak shape, mechanical advantage, and feeding ecology in modern birds. *Evolution* **73**, 422–435 (2019).

38. Olsen A. M. Feeding ecology is the primary driver of beak shape diversification in waterfowl. *Funct. Ecol.* **31**, 1985–1995 (2017).

39. Feduccia A. Explosive evolution in tertiary birds and mammals. *Science* **267**, 637–638 (1995).

40. Feduccia A. 'Big Bang' for Tertiary birds? *Trends Ecol. Evol.* **18**, 172–176 (2003).

41. Feduccia A. Avian extinction at the end of the Cretaceous: Assessing the magnitude and subsequent explosive radiation. *Cretac. Res.* **50**, 1–15 (2015).

42. Brusatte S. L., O'Connor J. K., Jarvis E. D. The origin and diversification of birds. *Curr. Biol.* **25**, R888–R898 (2015).

43. Losos J. B. Ecological character displacement and the study of adaptation. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 5693–5695 (2000).

44. Hernandez--Possémé L., Le Floch G., Bels V., Baguette M. Character displacement in shorebirds. In review.

45. Angarita-Báez J. A., Carlos C. J. Systematic review of migratory shorebirds carnivorous diet in their non-breeding period in the Americas. *Waterbirds* **47**, 1–16 (2024).

46. Gliesch L., et al. Specialist shorebirds respond to prey and habitat availability through trophic plasticity. *Food Webs* **36**, e00300 (2023).

47. Cramp S., Simmons K. E. L. *Handbook of the Birds of Europe, the Middle East and North Africa. Vol. III: Waders to Gulls* (Oxford Univ. Press, Oxford, 1983).

Acknowledgments

We warmly thank Raphaël Cornette and Frédéric Legendre for helpful methodological advice, and Virginie M. Stevens, Delphine Legrand and Hervé Philippe for constructive discussions and daily support. Lucyle Hernandez -- Possémé granted access to unpublished results. This manuscript is publication BRCXXX of the Biodiversity Research Centre at UCLouvain, to which N.S. is affiliated.

Funding: Include all funding sources, including grant numbers, complete funding agency names, and recipient's initials. Each funding source should be listed in a separate paragraph such as:
Internship UMR 7205 (EL)
Doctoral fellowship ED226, MNHN (GLF)
Internal grants UMR 7205 (MB, VB)
Agence Nationale de la Recherche grant TULIP 10-LABX-0041 (MB)

Author contributions:
      Conceptualization: MB
      Study design MB, GLF, VB, NS
      Data Acquisition: MB, GLF, EL
      Methodology: MB, NS, VB
      Data Analysis: NS
      Visualization: NS
      Supervision: MB
      Writing - original draft: MB
      Writing - review & editing: MB, GLF, EL, VB, NS

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data and code are available in the supplementary material.

Supplementary Materials