Contrasting genomic trajectories of *Bartonellaceae* symbionts of planthoppers

Mingjie Ma[1,2 *], Anna Michalik[3], Junchen Deng[2,4], Yi Hu[1], Piotr Łukasik[2 *]

1. Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences, Beijing Normal University, Beijing 100875, China

2. Institute of Environmental Sciences, Faculty of Biology, Jagiellonian University, Kraków, Poland

3. Department of Developmental Biology and Morphology of Invertebrates, Institute of Zoology and Biomedical Research, Faculty of Biology, Jagiellonian University, Kraków, Poland

4. Doctoral School of Exact and Natural Sciences, Jagiellonian University, Kraków, Poland

* to whom correspondence should be addressed. E-mail: mamingjie@mail.bnu.edu.cn; p.lukasik@uj.edu.pl

Abstract

Symbioses with microorganisms have shaped the nutritional biology and evolution of many insects. For example, several ant clades have adapted to nutrient-poor diets through symbiosis with a specific clade of bacteria in the family *Bartonellaceae* (Hyphomicrobiales), notorious for also including virulent vertebrate pathogens. Here we show that *Bartonellaceae* phylogenetically placed within the clade that has only encompassed ant symbionts to date – *Candidatus* genus Tokpelaia – have established as symbionts in four different clades of planthoppers (Insecta: Hemiptera: Fulgoromorpha). Genome size and contents indicate different levels of integration of

these strains into the planthopper host biology and their diverse roles. Symbionts infecting one of

the clades have some of the largest genomes among *Bartonellaceae*, at ca. 2 Mb, two others are

under 700 kb, and the fourth is reduced to barely 158 kb. The planthopper-associated *Tokpelaia*

strains with larger genomes, similarly to ant symbionts, encode multiple amino acid and vitamin

biosynthesis genes, complementing the degraded nutritional capabilities of their hosts' ancient

heritable endosymbionts. Strikingly, the smallest *Tokpelaia* genome lacks any genes linked to

essential amino acid biosynthesis, in contrast to all other known insect-associated bacteria with

genomes of comparable size. We identified a single vitamin biosynthesis gene and iron-sulfur

cluster assembly genes as its only putative contributions to the host biology.

Our results broaden the host spectrum of non-pathogenic *Bartonellaceae*, indicating that they

have contributed to nutrition and symbiotic consortium function in diverse diet-restricted host

clades. They also highlight an unexpectedly broad range of evolutionary outcomes for this

important bacterial group.

Key words: nutritional endosymbiont, amino acid biosynthesis, vitamins, Auchenorrhyncha,

Rhizobiales, *Bartonella*

41 Significance

Many insects rely on bacterial partners to overcome nutrient-poor diets, yet the diversity and

evolutionary trajectories and outcomes for these associations remain unclear. Our discovery of

Bartonellaceae bacteria in planthoppers improves the understanding of the host range of these

broadly relevant bacteria best-known as vertebrate pathogens. The finding that these planthopper

symbionts generally have the capacity to produce essential amino acids and vitamins relevant to

hosts and likely form long-term symbiotic associations agree with the expectations for sap-sucking

hemipteran symbionts. However, the breadth of sizes and functions of the newly assembled

genomes substantially expand the known range of states for *Bartonellaceae*, and their

evolutionary trajectories. In particular, the identification of a strain with an ultra-reduced genome

of only 158kb with no amino acid biosynthetic functions contrasts with all other known insect-

symbiotic bacteria with genomes in a similar size range. This study also indicates how symbionts

may enable the degeneration, functional loss, and evolution of other bacteria residing in the same

host.

1. Introduction

Microbial symbionts can play dramatically important and highly diverse roles in the biology of

their animal hosts, shaping their life history traits, ecological niches, biological interactions, and

evolutionary processes (Moran 2007; Douglas 2014; Lukasik and Kolasa 2024). In particular, host

long-term evolutionary trajectories have repeatedly been shaped by the microbial contributions to

the host nutrition (Moran 2007; Sudakaran, et al. 2017; Brentassi and de la Fuente 2024).

Microbial symbionts frequently help address challenges related to the digestion of recalcitrant

foods, the neutralization of toxic compounds, and the acquisition of limiting nutrients that animals

require but cannot make on their own. For example, plant-based foods typically have low nitrogen

contents and specifically lack essential amino acids as well as vitamins, and many herbivores rely

on symbiotic bacteria for nitrogen recycling, upgrading, and sometimes fixation, and for the

biosynthesis of lacking nutrients (Hansen and Moran 2014). Insects provide many independent

examples of associations between heritable symbiont acquisitions and shifts to limiting diets

(Cornwallis, et al. 2023).

Ants (Hymenoptera: Formicidae) are often quoted as an example of a clade that has

undergone extensive dietary diversification, including multiple independent transitions from more

balanced omnivorous diets to herbivory (Russell, et al. 2009; Smith, et al. 2023). These nutritional

changes have typically been enabled by bacterial symbionts, most often in the family

Bartonellaceae (Alphaproteobacteria: Hyphomicrobiales) (Russell, et al. 2009; Bisch, et al. 2018).

Genomics and experiments have confirmed *Bartonellaceae* roles in herbivorous genera

Cephalotes (Hu, et al. 2018), *Dolichoderus* (Bisch, et al. 2018), and *Tetraponera* (Ma, et al. 2025),

where they help recycle nitrogenous waste and supplement the host with nitrogen-rich

compounds, especially amino acids. On the other hand, *Bartonellaceae* were also found in

carnivores, and in the predatory ant *Harpegnathos saltator* they encode genes related to protein

degradation, amino acid transport, and riboflavin biosynthesis, apparently assisting the hosts with

the digestion of protein-rich diets and using amino acids as energy source, in addition to

provisioning vitamins (Neuvonen, et al. 2016; Bisch, et al. 2018). Phylogenomic analyses have

shown that the functionally diverse ant symbionts form a monophyletic clade – *Candidatus* genus

Tokpelaia – that is sister to the genus *Bartonella* that contains notorious human and mammalian

pathogens (Segers, et al. 2017; Wagner and Dehio 2019), but also symbionts of bees and other

arthropods (Bisch, et al. 2018; Ma, et al. 2025).

Planthoppers (Hemiptera: Fulgoromorpha) showcase a very different nutritional history and

symbiotic make-up. They have ancestrally fed on phloem sap, and to counteract its imbalanced

nutritional contents, have repeatedly formed symbiotic associations with microorganisms that

provide the lacking nutrients (Bennett and Moran 2013; Bell-Roberts, et al. 2019; Brentassi and

de la Fuente 2024). Specifically, this 263-million-year-old infraorder (Deng, et al. 2025) has

ancestrally associated with a Bacteroidetes symbiont *Candidatus* Sulcia muelleri (further referred

to as *Sulcia*) and a Betaproteobacteria symbiont *Ca.* Vidania fulgoroidea (thereafter: *Vidania*)

(Urban and Cryan 2012; Deng, et al. 2023). These two symbionts, transmitted strictly maternally

and co-diversifying with their planthopper hosts (Moran, et al. 2005), jointly synthesize the 10

essential amino acids. In most planthoppers, *Sulcia* produces 3 [leucine, valine and isoleucine],

and *Vidania* 7 [methionine, histidine, tryptophan, threonine, lysine, arginine and phenylalanine]

(Michalik, et al. 2021; Michalik, et al. 2025). However, in several planthopper clades, these

microbes have been replaced by Hypocreales fungi (Siehl, et al. 2024; Michalik, et al. 2025), or

else, joined by additional microbes that established residence within host tissues alongside the

ancient endosymbionts (Michalik, et al. 2023; Brentassi and de la Fuente 2024). These additional

symbionts, representing the genera *Sodalis*, *Asenophonus*, *Serratia*, and others, can complement

the amino acid biosynthesis pathways of the ancestral symbionts and frequently encode genes

involved in the biosynthesis of B vitamins (Bennett and Mao 2018; Michalik, et al. 2021; Michalik,

et al. 2024). However, we do not have a good understanding of how the nutritional functions vary

among symbiont or planthopper clades, how they have evolved, and how they complement the

functions encoded by other members of the symbiotic partnership. As the genomics-based

characterization of these more recently acquired symbionts in Auchenorrhyncha has been largely

limited to a few species carrying *Sodalis* (Koga, et al. 2013; McCutcheon, et al. 2019; Michalik,

et al. 2021), we know little about other microbes forming relationships with these sap feeders.

In our recent metagenomics study spanning 149 species of taxonomically diverse

planthoppers from different geographic areas (Deng, et al. 2025; Michalik, et al. 2025), in some

of the surveyed species we identified, for the first time, *Bartonellaceae* symbionts. This included

a strain infecting *Saccharosydne* sp., a Delphacidae planthopper, which had lost *Sulcia* and

whose *Vidania* had the smallest genome of any known bacteria, barely 50 kilobases, reduced to

provisioning a single amino acid, phenylalanine (Michalik, et al. 2025). We thus postulated that in

this and other systems, the nutritional functions lost from the ancient symbionts' genomes may

have been taken over by the *Bartonellaceae*. This conclusion has prompted a closer investigation

into the nature and significance of the associations between *Bartonellaceae,* their planthopper

hosts, and co-residing other microbes.

Specifically, we asked about *Bartonellaceae* distribution across planthopper clades and the

phylogenetic relationships among the newly identified strains and other members of the family,

including mammalian pathogens and ant symbionts. We also asked about the genomic

characteristics and contents of these newly identified strains, particularly in relation to nutritional

provisioning that had been identified as a key function of previously studied *Tokpelaia* symbionts

of ants. Finally, we asked how these microbes fit into multi-partner symbioses that provide hosts

with essential nutrients. We addressed these questions by exploring the extensive recent

metagenomics data for diverse global planthoppers, and through comparative genomics analyses

of the newly identified planthopper-associated and previously published *Bartonellaceae* strains.

2. Results

2.1. The distribution and phylogenetic relationships of planthopper-associated *Bartonellaceae*

Fig. 1. *Bartonellaceae* symbiont distribution across planthoppers, their representation within metagenomic assemblies, and genomic characteristics. (A) The genome-based phylogeny of 149

planthopper species representing 19 families, modified from Deng, et al. (2025), with species harboring *Bartonellaceae* indicated with red font. Clades where these symbionts were not detected were collapsed, with the number of species in these clades shown. **(B)** Taxon-annotated GC content-coverage plots for metagenomic assemblies of six planthopper species where *Bartonellaceae* were identified. Colored blobs represent contigs corresponding to the identified genomic fragments of different symbiotic bacteria. The size of each blob represents contig size. **(C)** Circular diagrams of *Tokpelaia* genomes from six species, showing gene positions on forward and reverse strands and basic genome characteristics. Note that only two genomes were circularized - most comprise multiple contigs, connected for the ease of illustration. The inner and outer circles represent coding sequences (CDS), and the functional categories are shown in green. tRNA, rRNA, and tmRNA genes are shown in red, blue, and purple, respectively. The dark gray and light gray in the middle circle represent different contigs. The size of the first five circles correlates with genome size, the sixth (OECLID) is magnified three times.

The systematic survey of 149 phylogenetically diverse planthopper species (Deng, et al. 2025; Michalik, et al. 2025) using 16S rRNA sequences reconstructed from metagenomes using phyloFlash (Gruber-Vodicka, et al. 2020), revealed *Bartonellaceae* in six species from four planthopper families, all classified to the genus level only. They included three members of the family Cixiidae: two representatives of the genus *Bothriocera* (BOTSP1, BOTSP2) and *Oecleus* sp. (OECLEU); a Dictyopharidae species *Lyncides* sp. (LYNSP); a Delphacidae planthopper *Saccharosydne* sp. (SACSP1), and a Kinnaridae species *Oeclidius* sp. (OECLID) (Fig. 1A, Tab. S1). This, combined with the phylogenetic distance among *Bartonellaceae*-hosting planthoppers (see below) indicated at least four independent host-microbe associations – one in each family. In turn, the Cixiidae clade that included three *Bartonellaceae*-hosting species also contained two species with no traces of this bacterium, and based on currently available data, we cannot distinguish among scenarios such as independent infections, an ancestral gain followed by subsequent losses in some clades, or a facultative association.

The metagenome assemblies for these six planthopper species contained a set of contigs,

between 1 and 344, with significant sequence similarity to the genome of *Candidatus* Tokpelaia

hoelldoblerii (*Bartonellaceae*) from a predatory ant *Harpegnathos saltator* (Neuvonen, et al. 2016),

and forming a distinct cluster in GC contents and coverage space (Fig. 1B, Tab. S2, S3). The

Bartonellaceae symbionts from Cixiidae planthoppers (BOTSP1, BOTSP2, and OECLEU) all

possessed relatively large genomes, ranging from 1.96 to 2.21 Mb across 17-344 contigs,

predicted to encode 2,114–2,181 proteins, and with the coding density from 78.5% to 84.84%

(Fig. 1C). Their genomic GC content values were relatively high, between 51.6% and 52.34%.

The *Bartonellaceae* symbionts of SACSP1 and LYNSP had genomes approximately three-fold

smaller, 0.68 Mb (15 contigs - SACSP1) and 0.63 Mb (circular - LYNSP), with 647 and 608

predicted protein-coding genes, respectively. These reduced genomes also exhibited markedly

higher coding densities—90.57% for SACSP1 and 93.2% for LYNSP—as well much lower GC

contents, 23.62% and 21.93%, respectively (Fig. 1C, Tab. S2). In turn, for the Kinnaridae strain

OECLID, we have identified a single circularly mapping contig of 158 kb, with a GC content of

16.86% and a coding density of 88.48%, encoding predicted 164 genes that included the large

majority of those conserved in other extremely reduced bacterial genomes (McCutcheon, et al.

2024). Despite the low sequence similarity to any previously known *Bartonellaceae* (84.13 - 86.39%

at 16S rRNA gene, amino acid identity across phylogenetically informative genes: 47.03 - 51.47%,

Tab. S4, S5) amino acid sequences of protein-coding genes primarily resembled *Bartonellaceae*

among all NCBI records. No further contigs within the assembly could be classified to the same

genome.

Fig. 2. Phylogenetic placement of *Bartonellaceae* symbionts of planthoppers. (A) Maximum likelihood

analysis of *Bartonellaceae* strains with sequenced genomes, based on 68 homologous single-copy genes.

The purple bars represent the genome size, and the blue bars - the GC content. **(B)** Maximum likelihood

phylogeny of full-length 16S rRNA gene sequences of *Bartonellaceae* associated with planthoppers, ants,

and selected other hosts, derived from metagenomes and the NCBI database. In both panels, the *Tokpelaia*

clade is indicated with green boxes; bold black labels indicate ant-associated, and bold red – planthopper-

associated strains. Bootstrap support values above 50% are shown by the respective nodes. The very long

branch leading to the OECLID strain was truncated in both figures.

Phylogenomic analysis of the planthopper-associated *Bartonellaceae* reliably placed them

among previously characterized ant symbionts forming the genus *Tokpelaia* (Fig. 2A, S1).

Specifically, the planthopper strains form two well-supported subclades within *Tokpelaia*. One of
them, on a relatively short branch, comprises symbionts of Cixiidae planthoppers (BOTSP1,
BOTSP2, and OECLEU) and forms a sister clade with strains of *Tetraponera* ants. The second
clade, monophyletic but with very long branches among strains, comprises symbionts of LYNSP,
SACSP1, and OECLID. These small-genome strains show a closer relationship to smallest-
genome symbionts of *Dolichoderus* ants, with strong support but long branches. This indicates
an accelerated rate of sequence evolution, which combined with a strong AT bias, could have
affected the accuracy of phylogenomic reconstructions through systematic problems such as long
branch attraction, especially for the extremely reduced OECLID (Susko and Roger 2021). 16S
rRNA gene-based phylogeny, including a much broader sampling of *Tokpelaia* sequences,
produced a topology broadly consistent with the genome-based analysis, again placing the
planthopper-associated symbionts among ant-associated *Tokpelaia* strains and forming two
separate clades (Fig. 2B, S2).

The genome synteny comparisons did not support phylogenetics-based conclusions (Fig. S3).
The highly fragmented assemblies of two of Cixiidae symbiont genomes prevented comparisons
within that clade, and there was little similarity among strains comprising the second clade.
Surprisingly, large syntenic regions were detected between BOTSP2 and SACSP1, despite them
representing different clades. Also, symbionts from BOTSP1 and BOTSP2 shared limited but
distinct regions of synteny with *Tokpelaia* from *Tetraponera* ants.

213 2.2. Genomic features and metabolic functions in *Tokpelaia*
214 symbionts from planthoppers and ants

The genomic contents comparisons among planthopper and ant symbionts revealed large
differences among bacterial strains and host clades (Fig. 3). Generally, *Tokpelaia* strains devote

a large portion of their genome contents to the metabolism of carbohydrates, amino acids,

cofactors and vitamins, nucleotides, and energy metabolism, as well as to membrane transport.

Interestingly, Cixiidae symbiont strains consistently encode much greater numbers of

carbohydrate metabolism genes than others. In turn, the numbers of amino acid and

cofactor/vitamin biosynthesis genes were relatively consistent except for the genome-reduced

planthopper symbiont clade and a single ant strain (JR021), where they were substantially

reduced. Interestingly, secondary metabolism genes are consistently much less abundant in

planthopper symbionts than in ant symbionts.

As the number of translation-related genes is comparable across all but the tiniest genome,

their relative proportion negatively correlates with the genome size. In the tiniest *Tokpelaia*-

OECLID genome, genetic information processing genes make up 61.8% of the total gene number,

as despite it losing many genes in that category, much greater losses occurred in other categories.

This included the elimination of all genes involved in essential amino acid biosynthesis, with only

two non-essential amino acid metabolism genes (*cysE* and *cysK*) and one vitamin biosynthesis

gene (*cobA*) remaining.

Fig. 3. Relative abundance of KEGG pathways in *Tokpelaia* genomes. For each genome and each

functional category, the numbers in the squares represent the number of genes in that category, and the

background colors represent the proportion of the genes in that category relative to all annotated genes in

the genome.

Comparative analysis of nutrient biosynthesis pathways across the sequenced planthopper

Tokpelaia genomes indicated a strong similarity among symbionts of Cixiidae (BOTSP1, BOTSP2,

and OECLEU) and those of herbivorous *Dolichoderus* and *Tetraponera* ants (Fig 4), whereas

strains from LYNSP and SACSP1 had substantially reduced biosynthetic repertoire. Most

strikingly, the highly reduced OECLID retained very little nutrient biosynthetic capacity.

Specifically, amino acid biosynthesis pathway comparisons revealed that the Cixiidae

planthopper and *Dolichoderus* and *Tetraponera* ant symbionts retain very similar gene sets,

spanning nearly complete pathways for the biosynthesis of non-essential and essential amino

acids with the exception of alanine-synthesizing transaminase (*alaA*) and asparagine synthase

(*asnB*). *Tokpelaia* of BOTSP2 is the only one missing prephenate dehydratase (*pheA2*), which

catalyzes the decarboxylation of prephenate to phenylpyruvate in the phenylalanine pathway. The

Tokpelaia of OECLEU lacks three additional genes: *aspB* and *glyA* for aspartate and glycine

synthesis, and *argF* involved in arginine biosynthesis. Notably, however, many of the genes in

Tokpelaia genomes of BOTSP1, BOTSP2, and OECLEU scattered across pathways, appeared

pseudogenized (Fig. 4). In contrast, *Tokpelaia* of LYNSP and SACSP1 have both lost most amino

acid biosynthesis genes. The LYNSP strain lacks the complete pathways for any essential amino

acid, retaining six of nine genes for lysine biosynthesis and complete pathways for four non-

essential amino acids: glutamate, aspartate, glycine, and serine. The SACSP1 strain retains

complete pathways for three essential amino acids (lysine, histidine and threonine) and three non-

essential amino acids (glutamate, aspartate, and glycine), along with partial pathways for leucine,

isoleucine, and valine. This reduced biosynthetic capacity resembles strain JR021, a culturable

member of conserved multi-partite microbiota of a herbivorous *Cephalotes* ant (Hu, et al. 2018).

The highly reduced OECLID strain encoded only two genes involved in the biosynthesis of one

non-essential amino acid, cysteine.

Fig. 4. Summary of *Tokpelaia* biosynthetic functions. Genes associated with nitrogen cycling and

cofactor biosynthesis in the genomes of planthopper (green) and ant (orange)-associated *Tokpelaia*. Each

bar represents a single gene with an abbreviation at the bottom. Genes are classified as present (green or

orange) and absent (light gray); putative pseudogenes, with disrupted open reading frames, are placed in

a black frame. A cladogram based on the phylogenomic relationships of *Tokpelaia* is shown on the left, with

labels in the same order as in Fig. 2A (where full host names are provided). UrD: Uric acid degradation,

UD: Urea degradation.

Besides amino acids, the *Tokpelaia* symbionts display a broad vitamin biosynthetic capability,

with substantial variation among strains, and again a substantial similarity among Cixiidae,

Dolichoderus, and *Tetraponera* symbionts. For thiamine biosynthesis, BOTSP1 and BOTSP2

symbionts encode the same set of seven genes as *Dolichoderus* strains, whereas all other strains

encode a more reduced set. All *Tokpelaia* strains except OECLID symbiont retain genes for

riboflavin (vitamin B2) biosynthesis, only missing *yigB* gene, with OECLEU additionally lacking

ribD. For CoA and pyridoxal-5P, *Tokpelaia* symbionts of Cixiidae planthoppers encode similar

gene sets as those of *Tetraponera* ants, with *Dolichoderus* and LYNSP-SACSP1 symbionts

encoding fewer. In turn, the set of four biotin (vitamin B7) biosynthesis genes is also highly

conserved among *Tokpelaia* strains (with a single gene loss in LYNSP), but all planthopper

symbionts encode a set of additional genes involved in the production of this vitamin. Interestingly,

BOTSP1's *Tokpelaia* is the only one retaining the full pathway from purines to folic acid, with all

other strains having experienced gene losses. Finally, cobalamin, a key coenzyme involved in the

metabolism of branched-chain amino acids and methionine, has the longest synthesis pathway

of 22 steps, all but one of which (*cbiK*) is encoded by *Tetraponera* and BOTSP1 symbionts. Here,

BOTSP2 and one of the *Dolichoderus* strains lost more genes, and all others have a more

substantially reduced set. The highly reduced OECLID symbiont only encodes one vitamin

biosynthesis gene, *cobA*, in the cobalamin biosynthesis pathway.

Similarly, *Tokpelaia* from Cixiidae planthoppers and most ants (*Dolichoderus*, *Tetraponera*,

and *Harpegnathos saltator*) harbor a complete urease gene cluster (ureABCDEFGJ) for urea

degradation. However, the LYNSP strain lacks two accessory components (ureE and ureJ), and

SACSP1 has lost the entire gene cluster, same as the *Cephalotes* ant symbiont JR021 (Fig. 4).

On the other hand, while the pathway for uric acid degradation is retained in *Tokpelaia* from

Dolichoderus and *Tetraponera* ants, BOTSP1 is the only planthopper whose symbiont retains the

entire pathway, with partial or complete losses in other species (Fig. 4). Again, none of the

nitrogen recycling genes were found in *Tokpelaia*-OECLID.

The close inspection of the OECLID symbiont gene set revealed, outside of genetic

information processing genes, two substantial functions that were retained (Fig. S4, Tab. S7). The

first is energy metabolism, including components of the electron transport chain (17 genes related

to cytochrome c oxidase, ATP synthase, and electron transfer), the second is iron–sulfur cluster

assembly. The only two amino acid (cysteine) biosynthesis-related genes present in this genome

are also involved in iron-sulfur cluster assembly (Fig. S5).

303   2.3. The degeneration of ancient symbionts is often associated with

304   the *Tokpelaia* acquisition

Fig. 5. Genomic complementarity in nutritional provisioning among planthopper symbionts. Amino acid and B vitamin biosynthesis gene distribution among genomes of different symbionts that were identified in metagenomes of six *Tokpelaia*-hosting planthopper species. Genes are classified as functional genes, absent (light gray), and pseudogenes (black).

Within metagenomic assemblies of the six *Tokpelaia*-hosting planthopper species we also identified other bacteria (Fig. 1B, Tab. S2) (Michalik, et al. 2025). Specifically, within all studied insects, we identified rRNA sequences of *Vidania*, and in five of them (all except SACSP1), of *Sulcia* – two ancestral symbionts of planthoppers. BOTSP1, BOTSP2 and OECLID also contained the facultative symbiont *Wolbachia*. LYNSP carried *Rhodanobacter*, whereas SACSP1 hosted *Sodalis* as an additional symbiont (Fig. 1B). For most of these symbionts, we recovered genomic bins with acceptable completeness and contamination statistics (Tab. S2). Intriguingly, in Cixiidae species, *Vidania* and *Sulcia* genomes were fragmented and apparently incomplete, unlike in the three other *Tokpelaia*-hosting species and the vast majority of previously studied planthoppers (Michalik, et al. 2025). Further, in BOTSP2, *Wolbachia* was not successfully reconstructed through binning; we identified matching contigs via BLASTn, but because of high contamination scores for this set, as computed by CheckM (72.81%), we decided to exclude it from analyses.

The ancient nutritional endosymbionts' genome organization and functions varied among the studied species (Fig. 5). Despite its fragmentation in Cixiidae, *Sulcia* nutritional functions were conserved in the five planthoppers that retained it, always comprising full pathways for valine, leucine, and isoleucine biosynthesis. In contrast, *Vidania* genome retained ancestral organization and the ability to produce seven essential amino acids in only two species, LYNSP and OECLID. In the third, SACSP1, *Vidania* genome was reduced to provisioning phenylalanine (Michalik, et al. 2025). In turn, in all three Cixiidae species, the unusually fragmented *Vidania* strains only retained a small and variable subset of genes scattered across biosynthetic pathways. In Cixiidae and SACSP1, the gaps in conserved essential amino acid biosynthesis pathways are mostly filled by

the genes encoded by other symbionts (Fig. 5). Specifically, in Cixiidae, the largely complete

biosynthetic gene sets encoded in *Tokpelaia* genomes seem to fully compensate for the *Vidania*

gene losses. With a few exceptions (lysine and phenylalanine), this is also the case in SACSP1;

additionally, in this species, 17 genes overlapping with those of *Tokpelaia* are encoded by *Sodalis*.

In LYNSP, the conserved gene sets of *Sulcia* and *Vidania* are almost entirely duplicated by

Rhodanobacter, with *Tokpelaia* providing further scattered functions. Finally, in OECLID, two

Wolbachia strains encode a set of biosynthetic genes that largely overlap with those of ancient

symbionts.

Species differed substantially in terms of B vitamin and cofactor biosynthesis gene sets and

their distribution across symbiont genomes (Fig. 5). In Cixiidae species, *Tokpelaia* encodes large

numbers of genes across different pathways, with *Wolbachia* duplicating these functions in

BOTSP1. In SACSP1 and LYNSP, *Sodalis* and *Rhodanobacter*, respectively, encode the greatest

number of genes, with *Tokpelaia* duplicating many of them. Finally, in OECLID, its two *Wolbachia*

genomes encode multiple functions, and *Tokpelaia* contributes one gene. Across all these

species, few vitamin biosynthetic pathways appear complete, but we note that the assembly

fragmentation may have prevented the detection of some genes, and they may also be encoded

within the host genomes.

3. Discussion

When famously writing about 'a veritable fairyland of insect symbiosis', Paul Buchner (1965)

referred to the huge variety of bacterial symbionts of Auchenorrhyncha that accompany, and

sometimes have replaced, their ancient, degraded nutritional endosymbionts. Of these, the

enterobacterial genus *Sodalis* – apparently, the most broadly distributed across the taxonomic

diversity of insects – may be the most widespread but also best understood in terms of genomic

evolution and nutritional contributions to the host biology (McCutcheon, et al. 2019; Renoz, et al.

2023). Through our research, the family *Bartonellaceae* joins the ranks of Auchenorrhynchan co-

symbionts, showing patterns of the distribution, evolution, and function not unlike *Sodalis*, but

casting a new perspective on microbial symbioses of hemipterans, ants, and other insects.

359 3.1. Bartonellaceae host range broadens

Bartonellaceae are primarily known as pathogens of mammals, often vectored by blood-

sucking insects (Engel, et al. 2011). As an example, *Bartonella bacilliformis* causes the

devastating and often lethal Carrion disease in humans (Pons, et al. 2016), while *Bartonella*

henselae causes persistent and usually asymptomatic infections in cats, vectored by cat fleas

(Chomel, et al. 1996). The discovery of the bee gut symbiont *Bartonella apis* (Segers, et al. 2017)

and the identification of more divergent strains colonizing diverse ants, classified to the

Candidatus genus *Tokpelaia* (Russell, et al. 2009; Bisch, et al. 2018; Ma, et al. 2025), have

changed the understanding of the biological spectrum of this bacterial group. Our discovery of

planthopper-associated *Bartonellaceae*, fitting right within the *Tokpelaia* clade and with at least

four independent original host association events, further broadens the list of confirmed hosts.

The plant-based diet of many *Bartonellaceae*-hosting ants and planthoppers, but also the

mutualistic interactions among many ants and hemipterans, including planthoppers (Ivens, et al.

2017; Bourgoin, et al. 2023), suggest potential avenues for parallel acquisition of related microbes

by these groups, or their exchange among groups. However, we already know that

Bartonellaceae host list is longer: for example, a divergent clade commonly colonizes stored-

product mites (Xiong, et al. 2024), and *Bartonellaceae* 16S rRNA sequences were identified in a

scorpion and in heteropteran bugs (Matsuura, et al. 2012; Martinez, et al. 2019). We expect that

a systematic survey across diverse arthropods will reveal these bacteria in a substantially broader

range of hosts. At the same time, as shown recently, systematic screens across related species

and within species, microscopy, transcriptomics, and/or experiments, can clarify the nature of their associations with diverse hosts (Xiong, et al. 2024; Ma, et al. 2025).

Overall, our findings support the view of *Bartonellaceae* as relatively broadly distributed across animals, and perhaps a wider range of environments. We suspect that some strains are relatively versatile and capable of synthesizing a wide range of nutrients, which enable them to move across and colonize new host lineages and then potentially evolve into more specialized associations. Some of these, including the mammal-pathogenic *Bartonella* clade, may retain the ability to switch hosts, while others may become strictly heritable, promoting genomic reduction and the loss of functions not essential to the specific host. The latter evolutionary trajectory agrees with the emerging consensus on the evolution of some other key insect-symbiotic bacteria, as exemplified by the enterobacterial genus *Sodalis* (McCutcheon, et al. 2019). It includes strains such as *Sodalis ligni* that live in dead wood (Tlaskal, et al. 2021)*; Sodalis praecaptivus* isolated from a human wounds following impalement on a branch (Clayton, et al. 2012); and diverse strains that have independently established heritable nutritional symbiotic relationships with a wide variety of insect lineages (Renoz, et al. 2023). Such heritable associations typically lead to rampant genomic changes, increased evolutionary rates, and rapid gene losses. Some of the degenerating symbionts may reach a stable state where gene loss slows to nearly a halt – as evidenced by high similarity among genomes of *Buchnera*, *Blochmannia*, or *Baumannia*, all likely derived from *Sodalis* tens or hundreds of millions years ago (McCutcheon, et al. 2019).

3.2. Planthopper symbionts represent contrasting genomic evolutionary outcomes

Variation in planthopper-associated *Tokpelaia* genome size and contents indicates their different lifestyles and varying effects on the host biology, paralleling the range of associations of *Sodalis*-allied symbionts.

On the one hand, the three *Tokpelaia* strains infecting Cixiidae planthoppers have the largest

genomes (~2Mb) among the *Tokpelaia* strains studied so far (0.96-1.9Mb), retaining a broad

range of amino acid and vitamin biosynthesis genes. This suggests a substantial contribution to

the host nutrition, comparable to that of ant-associated *Tokpelaia* (Bisch, et al. 2018; Ma, et al.

2025). Further, these strains' positions on short branches in the *Tokpelaia* phylogeny and high

GC contents suggest proximity to the putative ancestral state of other associations with diverse

ants and planthoppers. This resembles the versatile *Sodalis praecaptivus*, whose function-rich,

high-GC genome is much larger than most insect-symbiotic *Sodalis* strains, and consistently falls

near the base of *Sodalis* phylogenies, among variable-length branches representing insect-

symbiotic strains that departed from this ancestral state (McCutcheon, et al. 2019).

On the other hand, the *Tokpelaia* genomes of LYNSP and SACSP1 planthoppers are

substantially reduced (632-677 Kb), to the level comparable with *Sodalis*-derived long-term

obligate nutritional symbionts of aphids (B*uchnera*, 412-671 kb), carpenter ants (*Blochmannia*,

705-792 kb), sharpshooter leafhoppers (B*aumannia*, 633-759 kb), or tsetse flies (Wigglesworthia,

~700 kb) (Bennett, et al. 2015; Bing, et al. 2017; Chong, et al. 2019). Despite the massive variation

in their age of association, tissue localization, transmission mechanisms, and biosynthetic

capacity, all these gamma-proteobacterial symbionts provide deficient nutrients that make them

virtually indispensable to their hosts (Feldhaar, et al. 2007; McCutcheon, et al. 2009; Bing, et al.

2017). We suspect that LYNSP and SACSP1 *Tokpelaia* strains play comparable roles in

planthoppers. As both strains have lost multiple nutritional genes that are retained by other

symbionts, their specific situation may particularly resemble that of *Baumannia* (Wu, et al. 2006),

or *Buchnera* from *Cinara* aphids (Manzano-Marin, et al. 2016), or endobacterial symbionts of

mealybugs (Husnik and McCutcheon 2016): all of them are complementary in their amino acid

biosynthetic capacity to other symbionts residing within the same host insect.

In our view, the most striking discovery is the tiny, 158-kb genome of the *Tokpelaia* symbiont

of the Kinnaridae planthopper *Oeclidius* sp. (OECLID), comparable in size to the most reduced

known insect endosymbionts: *Sulcia*, *Vidania*, *Nasuia*, *Hodgkinia*, *Tremblaya*, or *Carsonella*

(McCutcheon, et al. 2024). All these bacteria colonize sap-sucking hemipterans, where they have

well-defined roles as providers of essential amino acids and sometimes vitamins. This does not

appear to be the case with *Tokpelaia*-OECLID – even if, as shown in a mealybug, one nutrition-

related biosynthetic gene (*cobA* in the current case) might potentially be enough to justify long-

term symbiosis (Garber, et al. 2024). But in fact, OECLID symbiont's genetic repertoire resembles

much more closely the endosymbionts of certain protists. Specifically, the 158-kb genome of a

Verrucomicrobium, *Ca.* Organicella extenuata, symbiont of an *Euplotes* protist inhabiting a

hypersaline Antarctic lake, also retains the capacity for genetic information processing but not for

the biosynthesis of amino acids or vitamins (Williams, et al. 2021), with fatty acid synthesis and

iron-sulfur (Fe-S) cluster assembly identified as its likely contribution to the host biology. Fe-S

cluster assembly has not been identified as critical to sap-sucking insects' biology. Thus, the

nature of the association, significance, and evolutionary history of this unusual planthopper-

associated bacterium that conflicts with the current understanding of insect heritable

endosymbioses remains a mystery and, unfortunately, it cannot be resolved based on a single

mid-coverage metagenome alone. To further complicate conclusions, from OECLID metagenome,

we also reconstructed a low-coverage rRNA sequence representing an Entomophthorales fungus

with 96.6% identity to entomopathogens *Pandora delphacis* and *Furia pieris*, and we cannot

exclude the possibility that *Tokpelaia*-OECLID actually associates with the planthopper's

pathogen (Matsuura, et al. 2018; Siehl, et al. 2024).

3.3. *Bartonellaceae* significance in the biology of planthoppers and

their other symbionts

We know that heritable nutritional endosymbionts in diverse insect systems produce essential

nutrients deficient in their hosts' diets, thus being indispensable to the hosts. At the same time,

they drive the evolution of hosts' dependence on symbiont nutritional contributions, even as these

symbionts degrade and loose efficiency – a process aptly described as 'the evolutionary rabbit

hole' (Bennett and Moran 2015). Further, they drive the complementary loss of functions by their

co-symbionts (McCutcheon, et al. 2009; Husnik and McCutcheon 2016), forcing the hosts to

manage an increasingly specialized symbiotic consortium. We think that these general statements

accurately describe the biology and significance of most planthopper-associated *Tokpelaia*

strains that we have identified, which in all cases, belong to multi-partite symbiotic consortia with

complementary biosynthetic capabilities (Michalik, et al. 2021). Specifically, in a delphacid

SACSP1, the *Bartonellaceae* symbiont largely fills up the nutritional gap resulting from *Sulcia* loss

and *Vidania* genome reduction to the smallest-known state, while in Cixiidae, they seem to

compensate for the apparent degeneration of *Vidania.* While such complementarity cannot be

seen in LYNSP, and in OECLID it is limited to a single gene, it is tempting to think of *Tokpelaia*

as the common driver of the functional loss by ancient symbioses. At the same time, other

symbionts are likely to be driving the gene loss in *Tokpelaia.* We propose that through alternative

evolutionary routes, symbiotic complexes involving *Tokpelaia* have reached, or are on a path

towards, complementarity in the biosynthesis of largely the same set of nutrients as most other

planthoppers. Depending on the order of arrival and the genomic properties of bacteria that were

in planthoppers before, the independently acquired *Bartonellaceae* have assumed different

positions and roles in the symbiotic consortium.

We are not aware of any major ecological transitions in planthopper lineages represented by

Tokpelaia-infected species. Such changes, exemplified by Typhlocybinae leafhoppers that

switched to feeding on parenchymal cell cytoplasm (Buchner 1965; Kobiałka, et al. 2025) and in

Derbidae and Achilidae planthoppers that switched to feeding on fungi during nymphal stages

(Michalik, et al. 2025), seem to have enabled major changes and functional losses in their
microbiota.

However, in all presented systems, achieving the full understanding of host-microbe
associations would require a much broader set of data than what we had available - a combination
of within-species microbiome surveys, genomic comparisons among closely related species,
microscopy, and, ideally, experiments. Unfortunately, these approaches are not easy to
implement on unidentified non-model species sampled on different continents years ago, which
was the case here. Nevertheless, this study demonstrates that a broad genomics survey alone
can help discover new biological states and processes that change our understanding of biology
and evolution of significant clades of organisms. In the age of genomics, such broad surveys
should become an important tool for the discovery and characterization of biodiversity and
biological interactions that shape it.

488 4. Methods

489 4.1. Study material and data

For the study, we used a set of metagenomic datasets for 149 taxonomically diverse
planthoppers, presented in our previous work that focused on phylogenomic relationships of these
planthoppers and comparative genomics of their *Sulcia* and *Vidania* symbionts (Deng, et al. 2025;
Michalik, et al. 2025). Briefly, insects were wild-caught from their natural habitats around the world
between 1996 and 2022. Following identifications by taxonomic experts, the DNA extracted from
dissected abdominal tissue of single individuals, or in a few cases, dissected bacteriome tissue,
was used for metagenomic library preparation and sequencing on Illumina platforms. Collection
metadata, reagents, and sequencing platforms for the entire metagenomic collection are provided

in the abovementioned publications. For the six specimens within which we detected *Tokpelaia*,

listed in Table S1, abdominal tissue was used for DNA extraction using the Bio-Trace DNA

Purification Kit (Eurx, Poland), and sequencing libraries were prepared and sequenced by

Novogene Ltd. on NovaSeq 6000 platform using a 300-cycle kit.

502   4.2. Bioinformatics analysis

To detect *Bartonellaceae* within the sequenced metagenomes, we systematically examined

16S rRNA sequences assembled using phyloFlash v3.4.2 (Gruber-Vodicka, et al. 2020),

validating all those without close matches to recognized bacteria through phylogenetics and

blastn comparisons against NCBI databases. In all cases where these data suggested

Hyphomicrobiales infections, we followed with the analysis of metagenomic assemblies obtained

using Megahit v1.2.9 (maximum k-mer size = 255, min contig size= 1000) (Li, et al. 2015). To

obtain the draft genomes of *Bartonellaceae* and other bacteria present in these samples, we

binned the metagenomic assembly data using the metaWRAP pipeline (Uritskiy, et al. 2018),

incorporating three binning tools: CONCOCT, MaxBin, and metaBAT. The resulting bins were

refined with metaWRAP's "Bin_refinement" module, applying a minimum completeness threshold

of 70% and a maximum contamination limit of 10%. To avoid the omission of genomic contigs,

assemblies were further examined using BLASTn and BLASTx searches against the NCBI

database, and separately against published and unpublished genomes of different insect-

symbiotic bacteria. Only contigs consistent with the bin classification, and meeting criteria for read

coverage and GC content were retained (Tables S2, S3). Bin completeness and contamination

were evaluated using CheckM v1.0.12 (Parks, et al. 2015), and taxonomic classification was

performed using the Genome Taxonomy Database via GTDB-Tk v2.1.122 (Chaumeil, et al. 2019).

Average   amino   acid   identity   (AAI)   was   calculated   using   CompareM   v0.1.2

(https://github.com/dparks1134/CompareM).

Contigs of bacteria other than *Sulcia* and *Vidania* were annotated with Prokka v1.14.6

(Seemann 2014), and gene functions were assigned to KEGG Orthology (KO) pathways using

KofamScan (Aramaki, et al. 2020) and the online tool eggNOG-mapper (Tab. S6) (Cantalapiedra,

et al. 2021). Based on these annotations, metabolic pathways of the symbiotic bacteria were

manually reconstructed using the KEGG database. Genomes were visualized using Proksee

(Grant, et al. 2023). Genome synteny analysis was performed using Promer tool for Mummer

v3.07, with default settings (Kurtz, et al. 2004). Pseudofinder v1.1.0 (Syberg-Olsen, et al. 2022)

software was used to predict pseudogenes in the genomes of symbiotic bacteria. In turn, all

confirmed *Sulcia* and *Vidania* contigs and genomes were annotated using a custom pipeline

(Łukasik, et al. 2018) adopted for planthopper symbionts recently (Michalik, et al. 2025). The

pipeline first extracted all Open Reading Frames (ORFs) and translated them into amino acid

sequences. These ORFs were then searched using HMMER v3.3.1 (Eddy 2011) (hmmer.org)

against a custom database containing manually curated protein-coding and rRNA genes from

previously characterized *Sulcia* and *Vidania* genomes. Ribosomal RNA genes were detected with

nhmmer implemented in HMMER v3.3.1, and tRNA genes were identified with tRNAscan-SE

v2.0.7 (Chan, et al. 2021) Based on the relative length to the reference genes, protein-coding

genes were classified as functional (>90%), putative pseudogenes (>60%), or pseudogenes

(<60%). Any ORFs longer than 300 bp with no significant similarity to any reference genes were

further examined by querying UniProt (The UniProt 2021) and NCBI databases and compared

carefully to the top hits. Genes lacking any annotations were marked as "hypothetical".

4.3. Phylogenetic analyses

To investigate the evolutionary origin of *Bartonellaceae* bacteria associated with

planthoppers, we compared phyloFlash-reconstructed 16S rRNA sequences against top hits of

insect and environmental origin, especially published ant-associated *Tokpelaia* sequences. All

sequences were aligned using MAFFT v7.526 (Rozewicki, et al. 2019), with divergent regions

removed using Gblocks v0.91b (Talavera and Castresana 2007). Phylogenetic analysis was

performed using RAxML v8.2.12 (Stamatakis 2014) under the GTRGAMMA model, with *Brucella*

pseudintermedia (accession number: CP099968) designated as the outgroup. The analysis

included 1000 bootstrap replicates and was visualized using iTOL (Letunic and Bork 2021).

To clarify the evolutionary relationships among the newly identified strains and other

Bartonellaceae, we conducted a genome-wide phylogenetic analysis that included all currently

published *Tokpelaia* or *Bartonella* genomes representing the major recognized clades. The

proteins were clustered into families using OrthoFinder v2.5.5 (Emms and Kelly 2019), resulting

in a data set of 68 single-copy orthologs. And then, proteins from each family were aligned using

MAFFT v7.526 (Rozewicki, et al. 2019), with divergent and ambiguously aligned blocks trimmed

using Gblocks v0.91b (Talavera and Castresana 2007) . We concatenated the resulting

alignments. Prottest 3.4.2 (Guindon and Gascuel 2003; Darriba, et al. 2011) was used to select

the optimal amino acid substitution model, JTT+I+G+F, according to AIC and BIC criterions.

Phylogenetic analysis was performed with RAxML v8.2.12 (Stamatakis 2014) using the best

model and 1000 bootstrap replicates, and results were visualized and annotated using iTOL

(Letunic and Bork 2021). All images were adjusted using Adobe Illustrator CC 2019.

5. Data availability

564 All raw data and genomic assemblies used in comparisons have been uploaded to NCBI

565 databases. BioProject and BioSample accession numbers are listed in Tables S1-S2.

566 Supplementary Tables were deposited in the public Figshare repository under the DOI:

567 10.6084/m9.figshare.30010978.

6. Acknowledgements

This study was supported by the Polish National Science Centre grant 2018/30/E/NZ8/00880 to P.Ł. and the China Scholarship Council grant no. 202406040164 to M.M.

7. Author Contributions

M.M., A.M., and P.Ł. conceived the study. J.D. and A.M. provided bioinformatics support and additional *Vidania* and *Sulcia* genome annotations. M.M., A.M., J.D., Y.H., and P.Ł. analyzed the data and wrote the manuscript. All authors reviewed and agreed on the manuscript.

8. References

Aramaki T, et al. 2020. KofamKOALA: KEGG ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics 36: 2251–2252. doi: 10.1093/bioinformatics/btz859

Bell-Roberts L, Douglas AE, Werner GDA 2019. Match and mismatch between dietary switches and microbial partners in plant sap-feeding insects. Proc. Biol. Sci. 286: 20190065. doi: 10.1098/rspb.2019.0065

Bennett GM, Mao M 2018. Comparative genomics of a quadripartite symbiosis in a planthopper host reveals the origins and rearranged nutritional responsibilities of anciently diverged bacterial lineages. Environ. Microbiol. 20: 4461–4472. doi: 10.1111/1462-2920.14367

Bennett GM, McCutcheon JP, McDonald BR, Moran NA 2015. Lineage-specific patterns of genome deterioration in obligate symbionts of sharpshooter leafhoppers. Genome Biol. Evol. 8: 296–301. doi: 10.1093/gbe/evv159

Bennett GM, Moran NA 2015. Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. Proc. Natl. Acad. Sci. USA 112: 10169–10176. doi: 10.1073/pnas.1421388112

Bennett GM, Moran NA 2013. Small, smaller, smallest: the origins and evolution of ancient dual symbioses in a Phloem-feeding insect. Genome Biol. Evol. 5: 1675–1688. doi: 10.1093/gbe/evt118

Bing X, et al. 2017. Unravelling the relationship between the tsetse fly and its obligate symbiont *Wigglesworthia*: transcriptomic and metabolomic landscapes reveal highly integrated physiological networks. Proc. R. Soc. B 284. doi: 10.1098/rspb.2017.0360

Bisch G, et al. 2018. Genome evolution of *Bartonellaceae* symbionts of ants at the opposite ends
of the trophic scale. Genome Biol. Evol. 10: 1687–1704. doi: 10.1093/gbe/evy126

Bourgoin T, et al. 2023. When cockroaches replace ants in trophobiosis: a new major life-trait
pattern of Hemiptera planthoppers behaviour disclosed when synthesizing photographic data.
Diversity 15: 356. doi: 10.3390/d15030356

Brentassi ME, de la Fuente D 2024. Obligate mutualistic heritable symbiosis in sap-feeding
insects: an intricate relationship in nature. Symbiosis 92: 159–185. doi: 10.1007/s13199-023-
00965-1

Buchner P. 1965. Endosymbiosis of animals with plant microorganisms. Interscience Publishers,
Inc, New York.

Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J 2021. eggNOG-mapper
v2: functional annotation, orthology assignments, and domain prediction at the metagenomic
scale. Mol. Biol. Evol. 38: 5825–5829. doi: 10.1093/molbev/msab293

Chan PP, Lin BY, Mak AJ, Lowe TM 2021. tRNAscan-SE 2.0: improved detection and functional
classification of transfer RNA genes. Nucleic Acids Res. 49: 9077–9096. doi:
10.1093/nar/gkab688

Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH 2019. GTDB-Tk: a toolkit to classify genomes
with the Genome Taxonomy Database. Bioinformatics 36: 1925–1927. doi:
10.1093/bioinformatics/btz848

Chomel BB, et al. 1996. Experimental transmission of *Bartonella henselae* by the cat flea. J. Clin.
Microbiol. 34: 1952–1956. doi: 10.1128/jcm.34.8.1952-1956.1996

Chong RA, Park H, Moran NA 2019. Genome evolution of the obligate endosymbiont *Buchnera*
aphidicola. Mol. Biol. Evol. 36: 1481–1489. doi: 10.1093/molbev/msz082

Clayton AL, et al. 2012. A novel human-infection-derived bacterium provides insights into the
evolutionary origins of mutualistic insect-bacterial symbioses. PLoS Genet. 8: e1002990. doi:
10.1371/journal.pgen.1002990

Cornwallis CK, et al. 2023. Symbioses shape feeding niches and diversification across insects.
Nat. Ecol. Evol.: 1022–1044. doi: 10.1038/s41559-023-02058-0

Darriba D, Taboada GL, Doallo R, Posada D 2011. ProtTest 3: fast selection of best-fit models of
protein evolution. Bioinformatics 27: 1164–1165. doi: 10.1093/bioinformatics/btr088

Deng J, et al. 2023. Genome comparison reveals inversions and alternative evolutionary history
of nutritional endosymbionts in planthoppers (Hemiptera: Fulgoromorpha). Genome Biol. Evol. 15:
evad120. doi: 10.1093/gbe/evad120

Deng J, et al. 2025. Phylogenomic insights into the relationship and the evolutionary history of
planthoppers (Insecta: Hemiptera: Fulgoromorpha). Syst. Entomol. 50: 495–518. doi:
10.1111/syen.12666

Douglas AE 2014. Symbiosis as a general principle in eukaryotic evolution. Cold Spring Harbor
Perspect. Biol. 6: a016113. doi: 10.1101/cshperspect.a016113

Eddy SR 2011. Accelerated profile HMM searches. PLoS Comp. Biol. 7: e1002195. doi:
10.1371/journal.pcbi.1002195

Emms DM, Kelly S 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 20: 238. doi: 10.1186/s13059-019-1832-y

Engel P, et al. 2011. Parallel evolution of a type IV secretion system in radiating lineages of the
host-restricted bacterial pathogen *Bartonella*. PLoS Genet. 7: e1001296. doi:
10.1371/journal.pgen.1001296

Feldhaar H, et al. 2007. Nutritional upgrading for omnivorous carpenter ants by the endosymbiont
Blochmannia. BMC Biol. 5: 48–59. doi: 10.1186/1741-7007-5-48

Garber AI, et al. 2024. Retention of an endosymbiont for the production of a single molecule.
Genome Biol. Evol. 16: evae075. doi: 10.1093/gbe/evae075

Grant JR, et al. 2023. Proksee: in-depth characterization and visualization of bacterial genomes.
Nucleic Acids Res. 51: W484–W492. doi: 10.1093/nar/gkad326

Gruber-Vodicka HR, Seah BKB, Pruesse E 2020. phyloFlash: rapid small-subunit rRNA profiling
and targeted assembly from metagenomes. mSystems 5: e00920–00920. doi:
10.1128/msystems.00920-20

Guindon S, Gascuel O 2003. A simple, fast, and accurate algorithm to estimate large phylogenies
by maximum likelihood. Syst. Biol. 52: 696–704. doi: 10.1080/10635150390235520

Hansen AK, Moran NA 2014. The impact of microbial symbionts on host plant utilization by
herbivorous insects. Mol. Ecol. 23: 1473–1496. doi: 10.1111/mec.12421

Hu Y, et al. 2018. Herbivorous turtle ants obtain essential nutrients from a conserved nitrogen-
recycling gut microbiome. Nat. Commun. 9: 964–978. doi: 10.1038/s41467-018-03357-y

Husnik F, McCutcheon JP 2016. Repeated replacement of an intrabacterial symbiont in the
tripartite nested mealybug symbiosis. Proc. Natl. Acad. Sci. USA 113: E5416–5424. doi:
10.1073/pnas.1603910113

Ivens ABF, Gadau A, Kiers ET, Kronauer DJC 2017. Can social partnerships influence the
microbiome? Insights from ant farmers and their trophobiont mutualists. Mol. Ecol. 27: 1898–1914.
doi: 10.1111/mec.14506

Kobiałka M, Świerczewski D, Walczak M, Urbańczyk W 2025. Extremely distinct microbial
communities in closely related leafhopper subfamilies: Typhlocybinae and Eurymelinae
(Cicadellidae, Hemiptera). mSystems 10: e00603–00625. doi: 10.1128/msystems.00603-25

Koga R, Bennett GM, Cryan JR, Moran NA 2013. Evolutionary replacement of obligate symbionts
in an ancient and diverse insect lineage. Environ. Microbiol. 15: 2073–2081. doi: 10.1111/1462-
2920.12121

Kurtz S, et al. 2004. Versatile and open software for comparing large genomes. Genome Biol. 5:
R12. doi: 10.1186/gb-2004-5-2-r12

Letunic I, Bork P 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Res. 49: W293–W296. doi: 10.1093/nar/gkab301

Li D, Liu C-M, Luo R, Sadakane K, Lam T-W 2015. MEGAHIT: an ultra-fast single-node solution
for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics 31:
1674–1676. doi: 10.1093/bioinformatics/btv033

Lukasik P, Kolasa MR 2024. With a little help from my friends: the roles of microbial symbionts in
insect populations and communities. Philos. Trans. R. Soc. Lond., Ser. B: Biol. Sci. 379:
20230122. doi: 10.1098/rstb.2023.0122

Łukasik P, et al. 2018. Multiple origins of interdependent endosymbiotic complexes in a genus of
cicadas. Proc. Natl. Acad. Sci. USA 115: E226–E235. doi: 10.1073/pnas.1712321115

Ma M, Zhu B, Zhang D, Łukasik P, Hu Y 2025. Symbiotic solutions for colony nutrition: conserved
nitrogen recycling within the bacterial pouch of *Tetraponera* ants. bioRxiv. doi:
10.1101/2025.03.13.643113

Manzano-Marin A, Simon JC, Latorre A 2016. Reinventing the wheel and making it round again:
evolutionary convergence in *Buchnera*-*Serratia* symbiotic consortia between the distantly related
Lachninae aphids *Tuberolachnus salignus* and *Cinara cedri*. Genome Biol. Evol. 8: 1440–1458.
doi: 10.1093/gbe/evw085

Martinez AJ, et al. 2019. Angiosperm to Gymnosperm host-plant switch entails shifts in microbiota
of the *Welwitschia* bug, *Probergrothius angolensis* (Distant, 1902). Mol. Ecol. 28: 5172–5187. doi:
10.1111/mec.15281

Matsuura Y, Kikuchi Y, Meng XY, Koga R, Fukatsu T 2012. Novel clade of alphaproteobacterial
endosymbionts associated with stinkbugs and other arthropods. Appl. Environ. Microbiol. 78:
4149–4156. doi: 10.1128/AEM.00673-12

Matsuura Y, et al. 2018. Recurrent symbiont recruitment from fungal parasites in cicadas. Proc.
Natl. Acad. Sci. USA 115: E5970–E5979. doi: 10.1073/pnas.1803245115

McCutcheon JP, Boyd BM, Dale C 2019. The life of an insect endosymbiont from the cradle to
the grave. Curr. Biol. 29: R485–R495. doi: 10.1016/j.cub.2019.03.032

McCutcheon JP, Garber AI, Spencer N, Warren JM 2024. How do bacterial endosymbionts work
with so few genes? PLoS Biol. 22: e3002577. doi: 10.1371/journal.pbio.3002577

McCutcheon JP, McDonald BR, Moran NA 2009. Convergent evolution of metabolic roles in
bacterial co-symbionts of insects. Proc. Natl. Acad. Sci. USA 106: 15394–15399.

Michalik A, et al. 2021. Alternative transmission patterns in independently acquired nutritional
cosymbionts of Dictyopharidae planthoppers. mBio 12: e0122821. doi: 10.1128/mBio.01228-21

Michalik A, Diego CF, Szklarzewicz T, Stroiński A, Łukasik P 2024. Facultatively intrabacterial
localization of a planthopper endosymbiont as an adaptation to its vertical transmission.
mSystems 9: e00634–00624. doi: 10.1128/msystems.00634-24

Michalik A, et al. 2025. The tiniest genomes shrink much further: extreme reductive evolution in
planthopper symbionts. bioRxiv. doi: 10.1101/2025.05.16.654412

Michalik A, et al. 2023. Variable organization of symbiont-containing tissue across planthoppers
hosting different heritable endosymbionts. Front. Physiol. 14: 1135346. doi:
10.3389/fphys.2023.1135346

Moran NA 2007. Symbiosis as an adaptive process and source of phenotypic complexity. Proc.
Natl. Acad. Sci. USA 104: 8627–8633.

Moran NA, Tran P, Gerardo NM 2005. Symbiosis and insect diversification: an ancient symbiont
of sap-feeding insects from the bacterial phylum Bacteroidetes. Appl. Environ. Microbiol. 71:
8802–8810. doi: 10.1128/AEM.71.12.8802-8810.2005

Neuvonen MM, et al. 2016. The genome of Rhizobiales bacteria in predatory ants reveals urease
gene functions but no genes for nitrogen fixation. Sci. Rep. 6: 39197. doi: 10.1038/srep39197

Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW 2015. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome
Res. 25: 1043–1055. doi: 10.1101/gr.186072.114

Pons MJ, Gomes C, Del Valle-Mendoza J, Ruiz J 2016. Carrion's disease: more than a sand fly-
vectored illness. PLoS Pathog. 12: e1005863. doi: 10.1371/journal.ppat.1005863

Renoz F, Arai H, Pons I 2023. The genus *Sodalis* as a resource for understanding the multifaceted
evolution of bacterial symbiosis in insects. Symbiosis 92: 187–208. doi: 10.1007/s13199-023-
00966-0

Rozewicki J, Li S, Amada KM, Standley DM, Katoh K 2019. MAFFT-DASH: integrated protein
sequence and structural alignment. Nucleic Acids Res. 47: W5–W10. doi: 10.1093/nar/gkz342

Russell JA, Moreau CS, Goldman-Huertas B, Fujiwara M, Lohman DJ 2009. Bacterial gut
symbionts are tightly linked with the evolution of herbivory in ants. Proc. Natl. Acad. Sci. USA 106:
21236–21241. doi: 10.1073/pnas.0907926106

Seemann T 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30: 2068–2069.
doi: 10.1093/bioinformatics/btu153

Segers FH, Kesnerova L, Kosoy M, Engel P 2017. Genomic changes associated with the
evolutionary transition of an insect gut symbiont into a blood-borne pathogen. ISME J. 11: 1232–
1244. doi: 10.1038/ismej.2016.201

Siehl R, Vyhnal K, Goffredi SK 2024. Friendly fungi: tropical insect families form partnerships with
intracellular fungi related to pathogens. iScience 27: 110674. doi: 10.1016/j.isci.2024.110674

Smith EJ, Vizueta J, Younger MA, Mullen SP, Traniello JFA 2023. Dietary diversity, sociality, and
the evolution of ant gustation. Front. Ecol. Evol. 11. doi: 10.3389/fevo.2023.1175719

Stamatakis A 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30: 1312–1313. doi: 10.1093/bioinformatics/btu033

Sudakaran S, Kost C, Kaltenpoth M 2017. Symbiont acquisition and replacement as a source of
ecological innovation. Trends Microbiol. 25: 375–390. doi: 10.1016/j.tim.2017.02.014

Susko E, Roger AJ 2021. Long branch attraction biases in phylogenetics. Syst. Biol. 70: 838–843.
doi: 10.1093/sysbio/syab001

Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F 2022. Pseudofinder:
detection of pseudogenes in prokaryotic genomes. Mol. Biol. Evol. 39: msac153. doi:
10.1093/molbev/msac153

Talavera G, Castresana J 2007. Improvement of phylogenies after removing divergent and
ambiguously aligned blocks from protein sequence alignments. Syst. Biol. 56: 564–577. doi:
10.1080/10635150701472164

The UniProt C 2021. UniProt: the universal protein knowledgebase in 2021. Nucleic Acids Res.
49: D480–D489. doi: 10.1093/nar/gkaa1100

Tlaskal V, Pylro VS, Zifcakova L, Baldrian P 2021. Ecological divergence within the
enterobacterial genus *Sodalis*: from insect symbionts to inhabitants of decomposing deadwood.
Front. Microbiol. 12: 668644. doi: 10.3389/fmicb.2021.668644

Urban JM, Cryan JR 2012. Two ancient bacterial endosymbionts have coevolved with the
planthoppers (Insecta: Hemiptera: Fulgoroidea). BMC Evol. Biol. 12: 87. doi: 10.1186/1471-2148-
12-87

Uritskiy GV, DiRuggiero J, Taylor J 2018. MetaWRAP—a flexible pipeline for genome-resolved
metagenomic data analysis. Microbiome 6: 158–171. doi: 10.1186/s40168-018-0541-1

Wagner A, Dehio C 2019. Role of distinct type-IV-secretion systems and secreted effector sets in
host adaptation by pathogenic *Bartonella* species. Cell. Microbiol. 21: e13004. doi:
10.1111/cmi.13004

Williams TJ, et al. 2021. Genome analysis of a Verrucomicrobial endosymbiont with a tiny genome
discovered in an Antarctic Lake. Front. Microbiol. 12: 674758. doi: 10.3389/fmicb.2021.674758

Wu D, et al. 2006. Metabolic complementarity and genomics of the dual bacterial symbiosis of
sharpshooters. PLoS Biol. 4: e188. doi: 10.1371/journal.pbio.0040188

Xiong Q, Sopko B, Klimov PB, Hubert J 2024. A novel *Bartonella*-like bacterium forms an
interdependent mutualistic symbiosis with its host, the stored-product mite *Tyrophagus*
putrescentiae. mSystems 9: e0082923. doi: 10.1128/msystems.00829-23
771