Title

Bruce suppresses autophagy-regulated caspase activity and wing tissue growth in *Drosophila*

Authors

Natsuki Shinoda[a,1], Yutaro Hama[b], Nozomi Hanawa[a] and Masayuki Miura[a,c,1]

Affiliations & Footnotes

[a]Department of Genetics, Graduate School of Pharmaceutical Sciences, The University of

Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[b]Institute for Genetic Medicine, Hokkaido University, Kita 15, Nishi 7, Kita-ku, Sapporo,

Hokkaido, 060-0815, Japan

[c]Laboratory for Cell Vigor Regulation, National Institute for Basic Biology, Nishigonaka 38,

Myodaiji, Okazaki, Aichi 444-8585, Japan

[1]Corresponding authors

Corresponding Authors

Natsuki Shinoda: Department of Genetics, Graduate School of Pharmaceutical Sciences,

University of Tokyo 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. Telephone: +81-3-

5841-4863. E-mail: f-shinoda.natsuki@g.ecc.u-tokyo.ac.jp

Masayuki Miura: Laboratory for Cell Vigor Regulation, National Institute for Basic Biology,

Nishigonaka 38, Myodaiji, Okazaki, Aichi 444-8585, Japan. Telephone: +81-564-55-7650. E-

mail: miura@nibb.ac.jp

Abstract

Caspases are cysteine-aspartic proteases that mediate both lethal and non-lethal cellular

outcomes, including the promotion of tissue growth. However, the mechanisms underlying the

differential regulation of these activities remain unclear. We have previously shown that among

the two *Drosophila* executioner caspases, Dcp-1 and Drice, Dcp-1 promotes tissue growth in a

non-lethal manner, independent of canonical apoptotic signaling. Herein, we demonstrated that

overexpressed Dcp-1, but not Drice, was activated without canonical apoptosome components.

TurboID-based proximity labeling revealed distinct proximal proteomes, among which Sirtuin

1, an Atg8a deacetylase, which promotes autophagy, was specifically required for Dcp-1

activation. Autophagy-related genes, including Bcl-2 family members *Debcl* and *Buffy*, are

required for Dcp-1 activation. Structure-based prediction using AlphaFold3 further identified

Bruce, an autophagy-regulated inhibitor of apoptosis, as a Dcp-1-specific regulator acting

outside the apoptosome-mediated pathway. Physiologically, Bruce suppresses wing tissue

growth. These findings indicate that non-lethal Dcp-1 activity is governed by the autophagy-

Bruce axis, enabling distinct non-lethal functions independent of cell death.

Introduction

Caspases are cysteine-aspartic acid proteases that serve as the central executioners of apoptosis. In *Drosophila*, core caspase-mediated signaling for apoptosis is well conserved (Nakajima and Kuranaga, 2017; Shinoda and Miura, 2024). Upon apoptotic stimulation, the pro-apoptotic genes *rpr*, *hid*, and *grim* (*RHG*), which antagonize DIAP-1, an inhibitor of apoptosis protein (IAP), are transcriptionally upregulated. The degradation of DIAP-1 allows the initiator caspase Dronc and its activator Dark to assemble into the apoptosome, thereby triggering a proteolytic cascade that activates the executioner caspases Dcp-1 and Drice, ultimately leading to apoptosis. In addition to their role in apoptosis, caspases perform diverse non-cell death functions collectively termed caspase-dependent non-lethal cellular processes (Aram et al., 2017). However, the mechanisms that differentially regulate lethal and non-lethal caspase activity remain largely unknown.

Functional specificity appears to arise at the level of the executioner caspases. Of the two executioner caspases in *Drosophila*, Drice is more potent than Dcp-1 in inducing cell death in wing imaginal discs (Florentin and Arama, 2012). In contrast, Dcp-1 is required for autophagy triggered by starvation or proteasome dysfunction in the ovaries (Choutka et al., 2017; DeVorkin et al., 2014; Hou et al., 2008). We have previously demonstrated that Dcp-1, but not Drice, specifically promotes imaginal tissue growth independent of apoptosome and cell death (Shinoda et al., 2019). Mechanistically, we showed that cleavage-mediated inactivation of Acinus, a positive regulator of autophagy (Haberman et al., 2010; Nandi et al., 2014), which limits tissue growth independently of autophagy (Tyra et al., 2020), partly promotes tissue growth (Shinoda et al., 2019). However, the upstream regulatory mechanisms that selectively regulate Dcp-1 activity remain unknown.

Proximity labeling has emerged as a powerful approach for comprehensively analyzing molecular interactions, including protein-protein interactions (Qin et al., 2021). TurboID

enables the rapid biotinylation of lysine residues on neighboring proteins within 10 minutes in

cultured cells (Branon et al., 2018). Streptavidin-based enrichment of these biotinylated

proteins, followed by mass spectrometry, allows the proteome-wide identification of proteins

near a target of interest. We previously generated *Drosophila* lines in which TurboID was

knocked in at the C-termini of Dronc, Dcp-1, or Drice (Shinoda et al., 2019). In wing imaginal

discs, TurboID-based labeling has revealed distinct proximity patterns for Dcp-1 and Drice,

suggesting that each executioner caspase is associated with a unique set of neighboring proteins

(Shinoda et al., 2019). We have shown that the differences in substrate specificity between Dcp-

1 and Drice were consistent with their distinct proximity profiles in the ovary, as exemplified

by BubR1 (Shinoda et al., 2023). Moreover, overexpression of *Fasciclin 3*, a Drice-proximal

protein identified in adult brains, induces non-lethal caspase activation in a Dronc-dependent

manner (Muramoto et al., 2025). These findings indicate that caspase-proximal proteins can

function not only as potential substrates but also as regulatory factors.

In this study, we investigated the regulatory mechanisms underlying Dcp-1 activation.

We found that the overexpression of *Dcp-1* induced a wingless phenotype. Using this

phenotype as a functional readout, we found that Dcp-1 activation specifically requires its

proximal protein Sirtuin 1 (Sirt1), Bcl-2 family members Debcl and Buffy, and components of

the autophagy pathway. Notably, the autophagy-regulated giant IAP Bruce specifically

interacts with and regulates Dcp-1 activity without affecting apoptosome-mediated core

apoptotic signaling, thereby suppressing wing imaginal tissue growth. Together, these findings

reveal that Bruce acts upstream to diverge the regulatory pathways of Dcp-1 and Drice, enabling

Dcp-1 to mediate distinct non-lethal functions through autophagy, independent of cell death.

Results

Overexpression of Dcp-1, but not Drice, induces apoptosis

Structurally, the executioner caspases, Dcp-1 and Drice, are similar, each comprising catalytic P20 and P10 domains and a highly disordered N-terminal prodomain (Figure 1A). First, we assessed the ability of two executioner caspases to trigger apoptosis. Using the *WP-Gal4* driver, which is expressed in the wing pouch region of the wing imaginal disc (Obata et al., 2014), overexpression of *Dcp-1* resulted in a wingless phenotype whereas the overexpression of *Drice* did not (Figure 1B). In wing imaginal discs, *Dcp-1* overexpression induces nuclear condensation (Figure 1C), indicative of apoptotic cell death. These results suggest that *Dcp-1* overexpression is sufficient to induce apoptosis, consistent with our previous observations in olfactory receptor neurons (Muramoto et al., 2025). Next, we examined the apoptotic potential of fluorescently tagged caspases in *Drosophila* S2 cells expressing mNeonGreen (mNG)-fused Dcp-1 or Drice. Dcp-1 expression caused a marked reduction in mNG intensity (Figure 1–figure supplement 1A–C) and a decreased number of mNG-positive cells (Figure 1–figure supplement 1A, D). These effects required Dcp-1 catalytic activity, as the mutation of the catalytic cysteine to glycine restored both mNG intensity and cell number (Figure 1–figure supplement 1A–D). In contrast, Drice expression neither reduced mNG intensity nor decreased the number of mNG-positive cells, resembling the catalytically inactive mutant (Figure 1–figure supplement 1A–D). Taken together, these results demonstrate that Dcp-1, but not Drice, is activated upon overexpression and induces apoptosis in multiple cell types.

Dcp-1 activation is independent of the apoptosome composed of Dronc and Dark

Executioner caspases are typically activated by cleavage mediated by the initiator caspase, Dronc, which is activated through the assembly of the Dark-dependent apoptosome (Figure 1D) (Nakajima and Kuranaga, 2017; Shinoda and Miura, 2024). We previously proposed that Dcp-

1 can be activated independently of Dark and Dronc (Shinoda et al., 2019). To test this

hypothesis further, we analyzed genetic modifiers of the Dcp-1-induced wingless phenotype.

A heterozygous mutation in the *H99* deficiency, which encompasses the pro-apoptotic *reape*r,

hid, and *grim* (*RHG*) genes, rescued the wingless phenotype caused by *Dcp-1* overexpression

(Figure 1E, G). Similarly, expression of the caspase inhibitor *p35* or knockdown of *RHG* genes

also rescued this phenotype (Figure 1F, G), suggesting that Dcp-1 induces apoptosis in a

catalytic activity-dependent manner under the control of RHG proteins. In contrast, knockdown

of other core apoptotic components, including *Dark*, *Dronc*, and *Drice*, or overexpression of

the *dominant-negative form of Dronc* (*Dronc^{DN}*) failed to suppress the wingless phenotype

(Figure 1F, G). These findings support our earlier observation that Dcp-1 can be activated

independent of the canonical apoptosome composed of Dronc and Dark to mediate non-lethal

functions (Shinoda et al., 2019). Collectively, these results indicate that, at least under

overexpression conditions, Dcp-1 activation proceeds independently of the apoptosome, in

contrast to that of Drice.

Dcp-1 and Drice show distinct expression patterns among tissues

Proximal proteins of caspases have emerged as potential regulators of executioner caspase

activity (Muramoto et al., 2025). We have previously demonstrated that the proximal protein

patterns of Dcp-1 and Drice differ even within the same tissue, such as wing imaginal discs

(Shinoda et al., 2019). Based on this, we sought to identify Dcp-1-specific regulatory factors

among its proximal proteins. First, we examined the tissue distribution of *Drosophila* caspases

using previously established *Caspase::V5::TurboID* knock-in fly lines (Shinoda et al., 2019).

Expression patterns were analyzed in larval tissues, including the salivary glands, fat bodies,

and brain (Figure 2–figure supplement 1A), with confirmed expression in wing imaginal discs

(Figure 2A) as well as in adult tissues, including the thorax, ovary, and testis (Figure 2–figure

supplement 1B). Drice was ubiquitously expressed in all examined tissues. In contrast, Dcp-1

expression was restricted to the larval brain, imaginal discs and adult ovaries and testes.

Drosophila executioner caspases associate with distinct sets of proximal proteins

Next, we compared the proximal protein profiles of *Drosophila* executioner caspases, based on

our previous observation that Dcp-1 and Drice are associated with distinct sets of proximal

proteins, potentially reflecting differences in their activation mechanisms (Shinoda et al., 2019).

Streptavidin staining, used to visualize proteins biotinylated via TurboID-mediated proximity

labeling, revealed distinct patterns of Dcp-1 and Drice expression across tissues (Figure 2A,

Figure 2–figure supplement 1A, B). To directly compare their proximal landscapes in the same

cellular context, we focused on the imaginal discs. NeutrAvidin-based purification of

biotinylated proteins, followed by mass spectrometry, identified 818 proteins (Figure 2B, C).

As expected, the proximal proteomes of Dcp-1 and Drice were distinct. BubR1 and Fasciclin 3

isoform G, previously identified as Drice-proximal in adult brain and ovary tissues (Muramoto

et al., 2025; Shinoda et al., 2023), were enriched in the Drice-associated proteome (Figure 2C,

Table S1), suggesting that Drice maintains its proximity to specific partners across tissues.

Among the top Dcp-1-enriched proximal proteins was the chaperonin containing TCP1 subunit

8 (CCT8). To validate these findings, we generated an antibody against CCT8 (Figure 2D).

Western blotting confirmed the presence of CCT8 in Dcp-1::V5::TurboID samples but not in

Drice::V5::TurboID samples (Figure 2E). Taken together, these results show that, even when

expressed at comparable levels in the same tissue, Dcp-1 and Drice are associated with distinct

sets of proximal proteins, which is consistent with their differential regulation and functional

specificity.

Screening of Dcp-1 proximal proteins identifies Sirt1 as a Dcp-1-specific activator

To identify potential regulators of Dcp-1 activation, we performed an RNAi screen targeting

eight genes that were selectively proximal to Dcp-1 relative to Drice and wild-type controls

(Dcp-1/Drice fold change > 1.5, Dcp-1/Control fold change > 10 and PSM $\geqq$ 2; Table S1),

using two independent RNAi lines per gene (Figure 2F). Under normal conditions, RNAi-

mediated knockdown of the chaperonin components *CCT2* and *CCT8* resulted in a wingless

phenotype, whereas knockdown of other genes had no effect (Figure 2–figure supplement 2A).

Under *Dcp*-1 overexpression conditions, knockdown of *Fkbp59* or *Sirt1* partially suppressed

Dcp-1-induced cell death (Figure 2G, Figure 2–figure supplement 2B). The requirement for

Sirt1 was further confirmed using a transheteroallelic combination of two previously described

Sirt1 deletion alleles, *Sirt1$^{2A-7-11}$* (Furuyama et al., 2004; Xie and Golic, 2004) and *Sirt1$^{4.5}$*

(Newman et al., 2002), which significantly rescued the wingless phenotype (Figure 2H, I).

These results indicate that the Dcp-1-proximal protein Sirt1 is required for Dcp-1 activation

upon overexpression.

Dcp-1 activation specifically depends on Debcl, Buffy, and autophagy

Sirt1, also known as dSir2, is a lysine deacetylase that promotes the expression of autophagy-

related genes via Atg8a deacetylation under starvation (Jacomin et al., 2020). Multiple studies

have linked Dcp-1 to the regulation of autophagy. Dcp-1, together with the Bcl-2 family

proteins Debcl and Buffy, but not Drice, has been shown to be required for starvation-induced

autophagy in *Drosophila* l(2)mbn cells (Hou et al., 2008), and Dcp-1 is also required for

autophagy in the ovary (Choutka et al., 2017; DeVorkin et al., 2014; Hou et al., 2008). More

recently, the BH3-only protein Sayonara (Synr) was reported to induce Dcp-1-dependent cell

death, which is regulated by Debcl, Buffy, and autophagy (Ikegawa et al., 2023). In contrast, a

gain-of-function modifier screen revealed that autophagy antagonizes Dcp-1-induced cell death

(Kim et al., 2010) and that Dcp-1 suppresses autophagy through cleavage-mediated inactivation

of Acinus (Haberman et al., 2010; Nandi et al., 2014).

Based on these reports, we examined whether Synr, Debcl, Buffy, and core autophagy

components regulate Dcp-1 activation. Knockdown of *Synr* did not suppress Dcp-1-induced

cell death, whereas knockdown of *Debcl* or *Buffy* significantly rescued this wingless phenotype

(Figure 3A, B). Similarly, knockdown of the autophagy-related genes *Atg2* and *Atg8a*

suppressed Dcp-1-induced cell death (Figure 3A, B), indicating that Debcl, Buffy, and

autophagy are required for Dcp-1 activation. To assess the specificity of these regulators, we

analyzed the eye ablation phenotype caused by overexpression of *Hid*, which triggers apoptosis

via canonical apoptotic signaling. As expected, this phenotype was rescued by *p35* expression

or knockdown of *RHG* genes and partially suppressed by inhibition of the Dark-, Dronc-, and

Drice-dependent apoptosome pathways (Figure 3C), confirming its dependence on canonical

apoptosis (Figure 1C). In contrast, *Hid*-induced apoptosis was not rescued by RNAi against

Dcp-1 (Figure 3C), nor by knockdown of *Sirt1* or *Fkbp59* (Figure 3D). Furthermore,

knockdown of *Debcl*, *Buffy*, and autophagy genes failed to rescue *Hid*-induced apoptosis

(Figure 3E). Taken together, these results indicate that Sirt1, Debcl, Buffy, and autophagy

components are not general regulators of apoptosis but function as specific activators of Dcp-

1.

Cleaved Dcp-1 specifically interacts with a giant-IAP Bruce

Next, we examined the molecular link between autophagy and Dcp-1 activation. Caspase

activation is negatively regulated by IAPs. In *Drosophila*, DIAP1 suppresses caspase activity

through ubiquitination (Ditzel et al., 2008) (Figure 1D). In addition to DIAP1, *Drosophila* has

another giant IAP, *Bruce*, which is degraded via autophagy during oogenesis (Nezis et al., 2010).

Bruce inhibits autophagy under nutrient-rich conditions (Hou et al., 2008), suggesting mutual

negative regulation of Bruce and autophagy. Bruce selectively inhibits *reaper*-induced cell

death, but not *hid*-induced apoptosis, suggesting a degree of pathway specificity (Domingues

and Ryoo, 2012). These observations led us to focus on Bruce as a potential selective regulator

of Dcp-1 activation.

The structure of human Bruce/BIRC6 has recently been resolved using cryo-electron

microscopy (Dietz et al., 2023; Ehrmann et al., 2023; Hunkeler et al., 2023; Liu et al., 2024),

revealing several conserved domains: linker domain (LD), baculoviral IAP repeat (BIR), WD40,

β-X1, β-X2, carbohydrate-binding module (CBM), ubiquitin-like (UBL), and ubiquitin-

conjugating (UBC) (Figure 4A). Human Bruce/BIRC6 ubiquitinates the activated executioner

caspases caspase-3 and caspase-7, but not their non-processed proforms *in vitro* (Dietz et al.,

2023). The BIR domain mediates interactions with ubiquitination substrates by recognizing N-

terminal IAP-binding motifs (IBMs) (Berthelet and Dubrez, 2013). The consensus IBM

sequence is defined as A(K/T/V/I)(P/A/E)(F/E/I/S/Y) (Vaux and Silke, 2005). We identified

IBM-like sequences at the N-termini of the P20 domains of both Dcp-1 and Drice (Figure 4B,

C). To assess the potential interaction between Bruce and these caspases, we used AlphaFold3

to predict the structure of the complexes between Bruce and the IBM-like motifs of Dcp-1 and

Drice (Abramson et al., 2024). Consistent with the known architecture of BIRC6, AlphaFold3

predicted that the BIR domain of Bruce interacts with LD, WD40, β-X1, and β-X2 domains

(Figure 4–figure supplement 1A, B). For structural prediction, we used a Bruce fragment

spanning residues 1–1960, including the BIR domain and its associated Zn^{2+}-binding site

(Figure 4A), and 15-residue peptides from the Dcp-1 and Drice cleavage sites encompassing

their IBM-like motifs (Figure 4B, C). The predicted structures revealed a high-confidence

interaction between Dcp-1 and the Bruce BIR domain. All five AlphaFold3 models showed

confident pLDDT scores (70–90) for the N-terminal residues of Dcp-1, corresponding to the

IBM-like motif (Figure 4D, F). The predicted binding mode resembled the canonical BIR-IBM

interaction observed in XIAP BIR3-Smac (Wu et al., 2000) (Figure 4–figure supplement 1C).

In contrast, although Drice appeared to bind to Bruce in the predicted models, the pLDDT

scores were consistently low (<70) across the interaction interface, including the four N-

terminal residues, indicating low confidence in the predicted interaction (Figure 4E, G).

To validate these predictions, we performed co-immunoprecipitation assays on

Drosophila S2 cells. We found that only the cleaved form of Dcp-1, which forms a head-to-tail

dimer composed of the P20 and P10 domains (Julien and Wells, 2017), and not the full-length

protein, interacted with the N-terminal BIR domain of Bruce following apoptosis induction

(Figure 4H), consistent with the observation that human Bruce/BIRC6 only targets cleaved

executioner caspases for ubiquitination *in vitro* (Dietz et al., 2023). Conversely, neither the

cleaved nor full-length forms of Drice showed detectable interactions with the Bruce BIR

domain (Figure 4H), corroborating the AlphaFold3 predictions. Taken together, these findings

suggest that Bruce specifically suppresses activated Dcp-1 but not Drice and serves as a

branching point for a Dcp-1-specific, non-canonical regulatory pathway.

Bruce is ubiquitously expressed in wing imaginal discs

Bruce has been reported to function in oogenesis and spermatogenesis in the ovaries and testes,

respectively (Arama et al., 2007, 2003; Hou et al., 2008; Kaplan et al., 2010; Nezis et al., 2010).

However, its expression in other tissues has not been thoroughly characterized. To examine the

tissue distribution of Bruce, we generated CRISPR/Cas9-mediated knock-in flies in which an

mStayGold::C4 linker::V5-tag (Ando et al., 2024) was inserted into the N-terminus of Bruce

(Figure 5–figure supplement 1A). Using the knocked-in line, we confirmed that Bruce was

expressed in the ovaries (Figure 5–figure supplement 1B) and testes (Figure 5–figure

supplement 1C), consistent with previous reports (Arama et al., 2007, 2003; Hou et al., 2008;

Kaplan et al., 2010; Nezis et al., 2010). Western blot analysis revealed Bruce expression in both

adult males and females (Figure 5–figure supplement 1D). We also detected Bruce expression

in wing imaginal discs (Figure 5–figure supplement 1E). This signal was abolished by the

RNAi-mediated knockdown of *Bruce*, confirming the specificity of the signal (Figure 5–figure

supplement 1F). These findings indicate that Bruce not only functions in the gonads but can

also exert Dcp-1 regulatory functions in wing imaginal discs.

Bruce specifically suppresses Dcp-1 overexpression-induced cell death

Next, we investigated whether Bruce regulated Dcp-1 activity. Upon overexpression of *Bruce*

(Figure 5A), we observed marked suppression of Dcp-1-induced cell death (Figure 5B, C).

Although *DIAP1* overexpression also inhibited Dcp-1 activation, this suppressive effect was

stronger following *Bruce* overexpression (Figure 5B, C). We examined the specificity of Bruce

in the regulation of other forms of cell death. As mentioned earlier, the BH3-only protein Synr

induces Debcl/Buffy, autophagy and Dcp-1-dependent cell death (Ikegawa et al., 2023). To

investigate whether Bruce was involved in Debcl/Buffy-autophagy-Dcp-1-mediated cell death,

we examined the role of Bruce in Synr-induced cell death. Overexpression of *Bruce*, but not

DIAP1, specifically suppressed Synr-induced cell death (Figure 5D). In contrast, *Bruce*

overexpression failed to rescue the eye ablation phenotype independent of autophagy caused

by *Hid* overexpression, whereas DIAP1 fully rescued this phenotype (Figure 5E). Taken

together, these results suggest that Bruce and DIAP1 regulate distinct cell death pathways, with

Bruce acting as a selective inhibitor of autophagy-regulated Dcp-1- and Synr-dependent cell

death.

Bruce suppresses wing tissue growth

We have previously reported that Dcp-1 activity can promote imaginal tissue growth

independent of the core apoptosis signaling pathway, including Dronc (Shinoda et al., 2019).

To investigate the role of Bruce in imaginal tissue growth, we analyzed its effect on wing

imaginal discs using *WP-Gal4*, which drives expression throughout the wing pouch.

Overexpression of *p35* suppressed wing tissue growth (Figure 5F, G), whereas inhibition of

canonical apoptotic signaling by *Dronc^{DN}* and *DIAP1* had no effect (Figure 5F, G). In contrast,

Bruce overexpression suppressed wing growth to an extent similar to *p35* overexpression,

whereas *Bruce* knockdown tended to increase wing size, although the difference was not

statistically significant (Figure 5F, G). We confirmed these results using *Engrailed-Gal4*, which

is expressed in the posterior compartment of the wing imaginal disc. *p35* overexpression

suppressed wing tissue growth (Figure 5H, I). In this context, the inhibition of canonical

apoptosis signaling by *Dronc^{DN}* and *DIAP1* had a minor but negative effect on wing size (Figure

5H, I). *Bruce* overexpression suppressed wing growth, whereas *Bruce* knockdown increased

wing size (Figure 5H, I), indicating that Bruce negatively regulates wing tissue growth. Finally,

to enhance the sensitivity of the assay, we overexpressed the dominant-negative form of the

insulin receptor (*InR^{DN}*) using *WP-Gal4*, which significantly reduced wing size (Figure 5J, K).

Under this sensitized condition, simultaneous *Bruce* knockdown significantly enhanced wing

tissue growth (Figure 5J, K). Overall, these results indicate that Bruce suppresses wing tissue

growth.

Discussion

Autophagy-facilitated alternative caspase activation pathway

In this study, we identified a distinct regulatory mechanism governing the activation of the *Drosophila* executioner caspase, Dcp-1, in sharp contrast to its paralog Drice. Although both caspases share structural similarities and are expressed in overlapping tissues, Dcp-1, but not Drice, is activated upon overexpression in an apoptosome-independent, autophagy-regulated manner. Previous studies have implicated Dcp-1 in the induction of autophagy (Choutka et al., 2017; DeVorkin et al., 2014; Hou et al., 2008). Our findings extend this view by showing that autophagy is required for Dcp-1 activation, suggesting a positive feedback loop between Dcp-1 and autophagy during apoptosis. Although the precise downstream targets of the Bcl-2 family proteins, Debcl and Buffy, remain unknown, their roles in autophagy regulation (Hou et al., 2008) suggest that they influence Dcp-1 activity through autophagic pathways. Additionally, we identified Bruce, a giant inhibitor of apoptotic proteins, as a Dcp-1-specific suppressor. Functionally, Bruce limits wing tissue growth, suggesting that Dcp-1 activation contributes to growth regulation in a non-lethal manner (Shinoda et al., 2019). Taken together, our results demonstrate that the regulatory mechanisms of Dcp-1 and Drice diverge at the level of IAPs and that Dcp-1 activation is regulated by autophagy, enabling specific non-lethal functions that promote tissue growth independent of canonical apoptotic signaling (Figure 5L). It is important to note that Dcp-1 can also suppress autophagy through cleavage of Acinus (Haberman et al., 2010; Nandi et al., 2014). Although this may seem contradictory, the ability of caspases to function in a non-lethal manner requires strict prevention of excessive activation. Such suppression may represent a key mechanism that enables non-lethal caspase activity.

It is also important to understand how cells determine the usage of canonical versus alternative caspase activation pathways depending on physiological or environmental conditions. In mammals, two major apoptotic pathways are activated by distinct initiator

caspases, caspase-8 and caspase-9, in response to different upstream signals (Newton et al.,

2024). Similarly, in *Drosophila*, while most developmental cell death is driven by canonical

apoptotic signaling involving Dronc (Xu et al., 2005), alternative caspase activation pathways

may be preferentially engaged in response to environmental stimuli, such as starvation or heat

shock.

Initiator caspase-independent activation mechanism for Dcp-1

The mechanism by which Dcp-1 is activated by itself remains an open and important question.

In contrast to Drice, Dcp-1 is activated upon overexpression in an apoptosome-independent but

autophagy-dependent manner. Unlike initiator caspases, executioner caspases such as Dcp-1

generally lack long N-terminal prodomains and require proteolytic cleavage to become

catalytically active. Proximity-labeling analysis revealed that Dcp-1 was predominantly

associated with the chaperonin subunit CCT8 (Figure 2E). Given that the overexpression of

Dcp-1 increases its local concentration for activation, we propose that its accumulation in the

chaperonin complex may serve as a local scaffold for caspase activation. This model implies a

spatially restricted mode of regulation in which caspase activity is compartmentalized within

the cell. In this model, we speculated that Dcp-1 activation occurs sporadically in subcellular

regions enriched in activators or scaffolding molecules such as CCT8. Under these conditions,

Dcp-1 activation may need to be tightly regulated to prevent its unintended proteolytic activity.

Supporting this notion, we proposed that Bruce functions as a basal suppressor of Dcp-1 in non-

lethal contexts, restraining caspase activity when full apoptotic signaling is not engaged.

Distinct roles of evolutionarily multiplicated executioner caspases

Caspases are essential cysteine proteases that execute apoptotic cell death and are conserved

from *C. elegans* to mammals (Shinoda and Miura, 2024). Multiple executioner caspases exist,

including caspase-3 and caspase-7 in mammals and Dcp-1 and Drice in *Drosophila*. Although

these enzymes are often functionally redundant in apoptosis, growing evidence indicates that

each executioner caspase harbors distinct substrate specificities and fulfills non-overlapping

roles (Boucher et al., 2012; Desroches and Denault, 2019; Shinoda et al., 2023, 2019; Walsh et

al., 2008). In mammals, caspase-3 and caspase-7 exhibit high sequence similarity but differ in

their substrate preferences. Caspase-7, for instance, contains polybasic residues in its exosite

that enhance its ability to cleave RNA-binding proteins, a function not observed in caspase-3

(Boucher et al., 2012). Mechanistically, this exosite enables caspase-7 to interact directly with

RNA, thereby promoting selective proteolysis of RNA-associated substrates (Desroches and

Denault, 2019). Furthermore, we have recently demonstrated that caspase-7 is essential for

early-peaking caspase activity at the plasma membrane during apoptosis, a process mediated

by its N-terminal intrinsically disordered region, which facilitates rapid efferocytosis of

apoptotic cells (Taira et al., 2025). In this study, we demonstrated that *Drosophila* executioner

caspases Dcp-1 and Drice are also subject to distinct activation mechanisms for the specific

non-lethal function of tissue growth. These findings underscore that the evolutionary expansion

of executioner caspases does not merely provide functional redundancy for apoptotic execution

but rather confers unique biochemical properties and regulatory roles under distinct cellular

contexts. This diversification allows executioner caspases to participate in both lethal and non-

lethal cellular processes, highlighting their broader functional repertoire beyond canonical

apoptosis.

 Methods

 Fly strains and rearing conditions

 We raised the flies in standard *Drosophila* medium at 25°C. For in vivo biotin labeling

 experiments, flies were raised in standard *Drosophila* medium supplemented with 100 μM (+)-

 biotin (#029-08713, WAKO). The fly strains used in this study are listed in Table S2.

 Genotypes corresponding to each figure are shown in Table S3.

 Intrinsically disordered region prediction

 Amino acid sequences were obtained from the FlyBase database [https://flybase.org/].

 Disordered regions were predicted using the PSIPRED protein sequence analysis workbench in

 DISOPRED3 [http://bioinf.cs.ucl.ac.uk/psipred/].

 Adult wings and eyes phenotypes scoring

 Adult female flies were anesthetized using CO_2. The wing and eye images were acquired using

 a Leica MZ16F fluorescent stereomicroscope (Leica Microsystems) equipped with a Leica

 DFC450 C CCD camera. Each wing was manually classified into one of four categories: *Intact*

 (no defect), *Mild* (≧50% of wing area intact), *Severe* (<50% but ≧5% of wing area intact), and

 Wingless (<5% of wing area intact).

 Molecular cloning

 For *pAc5-Caspases-myc-mNeonGreen*, the coding sequences of *Dcp-1* and *Drice* were

 amplified using PCR from the corresponding *pUASz-Caspase::V5::TurboID* plasmids

 (Muramoto et al., 2025). The coding sequence of myc-tagged mNeonGreen was PCR-amplified

 from *pUASz-Drice::myc::mNeonGreen plasmid* (Muramoto et al., 2025). The two fragments

were ligated into EcoRI/XhoI-digested *pAc5_STABLE2_neo* (González et al., 2011) using In-Fusion (TaKaRa, Z9648N).

For *pCold I-EcoRI-CCT8-EcoRI*, the coding sequence of *CCT8* was PCR-amplified from *Drosophila* cDNA, and the fragment was ligated into the *Eco*RI-digested *pCold I DNA* vector (#3361, TaKaRa) using In-Fusion.

For *pAc5-Caspases^{CG}-myc*, the coding sequences of *Dcp-1* and *Drice* were PCR amplified from the corresponding *pAc5-Caspases-myc-mNeonGreen* plasmids. The fragments were ligated into EcoRI/BamHI-digested *pAc5_STABLE2_neo* (González et al., 2011) using In-Fusion. For pAct-Bruce^{1-900a}a-3xFLAG, the first 900 aa of the coding sequences of *Bruce* were PCR-amplified from *Drosophila* cDNA, and the fragment was ligated into the *Eco*RI-digested *pAc5-Fasciclin 3 isoform A-3xFLAG* plasmids (Muramoto et al., 2025) using In-Fusion. For *pAc5-3xFLAG-mRuby3*, *Drosophila* codon-optimized mRuby3 was synthesized (Integrated DNA Technologies, Coralville, IA, USA) and ligated into the BamHI-digested *pAc5-3xFLAG-Bub3* plasmids (Shinoda et al., 2023) using In-Fusion.

For mStayGold::C4 linker::V5-tag knock-in for Bruce, the guide RNA was identified using the CRISPR Optimal Target Finder tool available on flyCRISPR [http://flycrispr.molbio.wisc.edu/]. The DNA fragments for the guide RNA were subcloned into BbsI-digested pDCC6 vectors (#59985, Addgene) (Gokcezade et al., 2014) to establish the *pDCC6-dBruce-sgRNA* plasmid. The following primers were annealed to generate the DNA fragments for guide RNAs: 5′-CTTCGCTTGCTTGAGGACGCGCTGC-3′ and 5′-AAACGCAGCGCGTCCTCAAGCAAGC-3′. For the generation of the homology-directed repair template (*pBac-mStayGold-c4-V5-dBruce-donor-3xP3-dsRed-polyA*), DNA fragments of the 5′ homology arm were assembled into the BsaI site, and the DNA fragment of the 3′ homology arm with mStayGold::C4 linker::V5-tag was assembled into the SapI site of pBac[3xP3-DsRed_polyA_Scarless_TK] (Shinoda et al., 2019) using In-Fusion. The PAM

sequence (TGG) of the gRNA-binding sites in the donor template was mutated (TGA) to

prevent Cas9-directed cleavage following homology-directed repair. The mStayGold::C4

linker::V5-tag fragment was amplified from pRSETB/mStayGold(c4) (#212018; Addgene)

(Ando et al., 2024) with V5-tag harboring primers. DNA fragments were amplified using PCR

with the following primers: dBruce 5′ homology arm (forward primer, 5′-

TGAAGGTCTCCTTAAAAGCCCGATTGGCGGCGCAGGAAGT-3′, and reverse primer,

5′-TCTTTCTAGGGTTAATCGAGTGCCCAGGGGGCGCGCGAGT-3′), dBruce 3′

homology arm-1 (forward primer, 5′-

TCTTTCTAGGGTTAATTTTTGGCATTCACAGGCGATTGCT-3′, and reverse primer, 5′-

TCCGGAGGTGCGCCGCGGCAATTTG-3′), mStayGold-C4 linker-V5-tag (forward primer,

5′-CGGCGCACCTCCGGAATGGTGTCTACAGGCGAGGAGCTGT-3′, and reverse primer,

5′-

GGTGCTGTCCAGGCCCAGCAGGGGGTTGGGGATGGGCTTGCCCACGGCAGAAGC

AGAAGGCTCGTGC-3′), 3′ homology arm-2 with mutated PAM (forward primer, 5′-

GGCCTGGACAGCACCGCCACGGAGCAGCATCATCAGCAGCGCGTCCTCAA-3′, and

reverse primer, 5′-GACGGCTCTTCATTAACTCTTAAATCAAATTTAAATTGAA-3′). The

established plasmids were injected into *yw; nos-Cas9(II-attP40)* for CRISPR/Cas9-mediated

transgenesis (BestGene Inc.). Each DsRed-positive transformant was isogenized and confirmed

using genomic PCR and sequencing.

All established plasmids were sequenced by Eurofins, Inc. The detailed plasmid DNA

sequences are available upon request.

Cell culture

Drosophila S2 cells were grown at 25°C in Schneider's *Drosophila* medium (GIBCO,

21720001) supplemented with 10% (v/v) heat-inactivated fetal bovine serum, 100 U/mL

penicillin, and 100 μg/mL streptomycin (WAKO, 168-23191).

mNeonGreen signal quantification of S2 cells

Drosophila S2 cells (2.5×10^5) seeded in 24-well plates were transfected with 200 ng of the

desired plasmid using Effectene Transfection Reagent (QIAGEN, 301427) following the

manufacturer's protocol. Sixteen hours after transfection, cells were transferred to four-

compartment glass-bottomed dishes (Greiner #627870). After another 24 hours, images were

captured using a Leica TCS SP8 confocal microscope (Leica Microsystems).

The mNeon Green intensity was quantified using Fiji (ImageJ) software (NIH Image).

Images were analyzed automatically using the ImageJ macro, as previously described (Fujisawa

et al., 2020) with modifications. Briefly, the mNeonGreen channel was filtered by the

"Gaussian Blur (sigma = 1)", processed with "Auto Threshold ("Mean" method)," processed

with "Watershed," and processed with "Analyze particles (size = 40–200 μm^2, circularity =

0.2–1.0)" to define the region of interest. For a given region of interest, the average mNeon

Green intensity was measured.

Protein preparation of Drosophila tissues

Whole larvae without digestive tract and fatobodies, larval salivary glands, fatbodies, brains,

wing discs and whole adults, adult thoraxes, ovaries, and testises of desired genotype were

dissected and lysed with RIPAi buffer (50 mM Tris-HCl, pH 8.0, 150 mM sodium chloride, 0.5

wt/vol% sodium deoxycholate, 0.1% w/v sodium dodecyl sulfate, and 1.0% w/v NP-40)

supplemented with cOmplete ULTRA EDTA-free protease inhibitor cocktail (#05892953001,

Roche). Samples were homogenized and centrifuged at 20,000 × *g*, 4°C for 10 min. The

supernatants were collected and snap-frozen in liquid nitrogen. Protein concentrations were

determined using the BCA assay (#297-73101, Wako) following the manufacturer's protocol.

The samples were mixed with Laemmli buffer, boiled at 95°C for 5 min, and then subjected to

western blotting analysis.

Western blotting

Each sample was separated using SDS-PAGE. The proteins were then transferred onto

Immobilon-P polyvinylidene fluoride membranes (#IPVH00010; Millipore) for

immunoblotting. Membranes were blocked with 4% skim milk diluted in 1× TBST.

Immunoblotting was performed using the antibodies mentioned below, which were diluted with

4% skim milk. Signals were visualized using Immobilon Western Chemiluminescent HRP

Substrate (#WBKLS0500; Millipore) and FUSION SOLO. 7S. EDGE (Vilber-Lourmat).

Contrast and brightness adjustments were applied equally using Fiji (ImageJ) software (NIH

Image).

The primary antibodies used included mouse anti-V5 monoclonal antibody (1:1,000,

#46-0705, Invitrogen), mouse anti-FLAG M2 monoclonal antibody (1:5,000, Sigma), rabbit

anti-CCT8 polyclonal antibody (1:5,000, this study), rabbit anti-Myc-Tag (71D10) monoclonal

antibody (1:5,000, #2278, CST), rabbit anti-3xDYKDDDDK Tag (3xFLAG Tag) (E7C5T)

monoclonal antibody (1:5,000, #87537, CST), mouse anti-alpha tubulin (DM1A) monoclonal

antibody (1:10,000, #T9026, Sigma), mouse anti-Histone H3 (1B1B2) monoclonal antibody

(1:10,000, #14269, CST), and mouse anti-Actin monoclonal antibody (1:5,000; #A4700,

Sigma). The secondary antibodies used were goat anti-rabbit IgG horseradish peroxidase

(HRP)-conjugated antibody (1:5,000, #7074S, CST) and goat/rabbit/donkey anti-mouse IgG

HRP Conjugate (1:5,000, #W402B, Promega). The membranes for streptavidin blotting were

blocked with 3% bovine serum albumin. Streptavidin blotting was performed using

streptavidin-HRP (1:10,000, #SA10001; Invitrogen) diluted in 3% bovine serum albumin.

Experimental design for liquid chromatography–mass spectrometry (LC-MS/MS) analyses

Two biological replicates were analyzed for each experimental condition to determine the

abundance of Drice/Dcp-1::TurboID relative to the control.

Purification of biotinylated proteins

Samples were prepared as previously described (Shinoda et al., 2023). Briefly, 100 µg of

biotinylated protein-containing lysate was subjected to FG-NeutrAvidin bead (#TAS8848

N1171, Tamagawa Seiki) purification. FG-NeutrAvidin beads (25 µL, approximately 500 µg)

were washed three times with RIPA buffer. Benzonase-treated biotinylated protein samples

suspended in 1 mL of RIPA buffer were incubated overnight at 4°C. Conjugated beads were

magnetically isolated and washed with 500 µL of an ice-cold RIPA buffer solution, 1 M KCl

solution, 0.1 M Na_2CO_3 solution, and 4 M urea solution. For western blot analyses, the purified

samples were mixed with Laemmli buffer, boiled at 95°C for 5 min, and then subjected to

western blot analysis. For LC-MS/MS analysis, the purified samples were washed with 500 µL

ultrapure water (#214-01301, Wako) and 500 µL of 50 mM ammonium bicarbonate. The

samples were then mixed with 50 µL of 0.1% Rapigest diluted in 50 mM ammonium

bicarbonate, and 5 µL of 50 mM TCEP was subsequently added. Samples were incubated at

60°C for 5 min, and then 2.5 µL of 200 mM methyl methanethiosulfonate was added. One

microgram sequence-grade modified trypsin (#V5111, Promega) was then added for on-bead

trypsin digestion at 37°C, leaving the treatment for 16 h. The beads were then magnetically

isolated, and 60 µL of the supernatant was collected. Then, 3 µL of 10% trifluoroacetic acid

was added to the supernatants, incubating them at 37°C for 60 min with gentle agitation. The

samples were then centrifuged at 20,000 × *g*, 4°C for 10 min. The peptides were then

desalinated and purified using a GL-tip SDB (#7820-11200, GL Sciences) following the

manufacturer's instructions. The samples were speed-backed at 45°C for 30 min and dissolved

in 25 µL of 0.1% formic acid. The samples were then centrifuged at 20,000 × *g*, 4°C for 10

min, and the supernatants were collected. The peptide concentrations were determined using

the BCA assay. Finally, 500 ng of the purified protein was subjected to LC-MS/MS analysis.

LC-MS/MS analyses

Samples were loaded onto an Acclaim PepMap 100 C18 column (75 µm x 2 cm, 3 µm particle

size and 100 Å pore size; #164946, Thermo Fisher Scientific) and separated on a nano-capillary

C18 column, (75 µm x 12.5 cm, 3 µm particle size, #NTCC-360/75-3-125, Nikkyo Technos)

using EASY-nLC 1200 system (Thermo Fisher Scientific). Elution conditions are listed in

Table S4. The separated peptides were analyzed using QExactive (Thermo Fisher Scientific) in

data-dependent MS/MS mode. The parameters for the MS/MS analysis are listed in Table S5.

The collected data were analyzed using Proteome Discoverer (PD) 2.2 software with the

Sequest HT search engine. The parameters for the PD 2.2 analysis are described in Table S6.

Peptides were filtered at a false discovery rate of 0.01 using the Percolator node. Label-free

quantification was performed based on the intensities of the precursor ions using a precursor-

ion quantifier node. Normalization was performed using the total amount of peptides in all

average scaling modes. MS proteomic data were deposited in the ProteomeXchange

Consortium via the jPOST partner repository with the dataset identifier PXD067425 (Okuda et

al., 2017).

Anti-CCT8 antibody generation

CCT8 was expressed in *Escherichia coli BL21* transformed with *pCold I-EcoRI-CCT8-EcoRI*.

Cells were grown at 37 °C overnight, diluted to OD_{600} 0.2 and cultured until OD_{600} 0.5 at 37°C.

Protein expression was induced by the addition of 1 mM IPTG at 15°C for 24 h. Cells were

centrifuged at 4000 × *g*, 4 °C for 20 min and resuspended in phosphate-buffered saline (PBS).

Cells were centrifuged at 4000 × *g*, 4°C for 10 min. Purification of 6x His-tagged CCT8 was

performed using the QIAexpress Ni-NTA Fast Start Kit (#30600, Qiagen) according to the

manufacturer's protocol: purification of 6xHis-tagged proteins under denaturing conditions.

Briefly, the cell pellets were lysed in Lysis Buffer (Denaturing Buffer adjusted to pH 8.0) with

gentle agitation for 60 min. The sample was centrifuged at 9,200 × *g*, 20°C for 30 min. The

supernatant was applied to Fast Start column and washed twice with Wash Buffer (Denaturing

Buffer adjusted to pH 6.3). Proteins were eluted using Elution Buffer (Denaturing Buffer

adjusted to pH 4.5). Polyclonal antibodies were raised in rabbits using purified 6xHis-tagged

CCT8 as an immunogen (Eurofins Inc.).

Protein complexes prediction with AlphaFold3

Amino acid sequences were obtained from the FlyBase database [https://flybase.org/]. The

protein complex structures were predicted using AlphaFold3 (Abramson et al., 2024) on the

AlphaFold server [https://alphafoldserver.com/].

Co-immunoprecipitation

Drosophila S2 cells ($2.0 × 10^6$) seeded in 6-well plates were transfected with the desired

plasmid using Effectene Transfection Reagent. For live cell lysate, *Drosophila* S2 cells were

lysed on ice using 550 µL of IP lysis buffer (50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.5%

NP-40) supplemented with cOmplete ULTRA EDTA-free protease inhibitor cocktail. For

apoptotic cell lysate, the cells were treated with 10 µg/mL cycloheximide (#C7698, Sigma) for

apoptosis induction. After 6 h of incubation, cells were harvested and centrifuged at $800 \times g$,

4 °C for 5 min and washed twice. Then, cells were lysed by using 550 μL of IP lysis buffer (50

mM Tris-HCl (pH 7.5), 150 mM NaCl, and 0.5% NP-40) supplemented with cOmplete

ULTRA EDTA-free protease inhibitor cocktail. The cell lysates were centrifuged at $20,000 \times g$,

4 °C for 5 min. Supernatants (50 μL) were collected as inputs. The remainder of the supernatant

was incubated for 2 h at 4°C with 20 μL of anti-DDDDK-tag mAb-Magnetic Agarose (#M185-

10R, MBL) equilibrated with the IP lysis buffer. The beads were washed thrice in IP lysis buffer

and boiled for 5 min at 95°C with 50 μL 1× Laemmli buffer. The samples were magnetically

separated, and the supernatants were subjected to SDS-PAGE.

Immunohistochemistry

Wing imaginal discs of the late third-instar larvae and testes and ovaries of the adults were fixed

with 4% paraformaldehyde in PBS. The wing imaginal discs and testes samples were washed

in PBS containing 0.1% Triton X-100 (PBST), and the ovary samples were washed in 1.0%

PBST, blocked in PBST with 5% normal donkey serum (PBSTn) for 30 min, and incubated

with primary antibodies in PBSTn overnight at 4°C. The samples were washed with PBST,

incubated for 2 h with secondary antibodies suspended in PBSTn, and washed again with PBST.

Images were captured using a TCS SP8 microscope (Leica Microsystems).

The primary antibodies used were rabbit anti-HA (Y11) polyclonal antibody (1:100,

#sc-805, Santa Cruz), mouse anti-Myc-tag monoclonal antibody (1:500, #R950-25, Invitrogen)

and mouse anti-V5 tag monoclonal antibody (1:100, #960-25, Invitrogen). The secondary

antibodies used were Cy3-conjugated donkey anti-rabbit IgG (1:250, #711-165-152, Jackson

Immuno Research), Alexa488-conjugated donkey anti-mouse IgG (1:500 (Myc) or 1:100 (V5),

#A-21202, Thermo Fisher Scientific). Hoechst 33342 (8 μM; #H3570, Invitrogen) for nuclear

staining and Phalloidin-iFluor633 (1 : 1000, #ab176758, Abcam) for F-actin staining were used

during the secondary antibody incubation. Images were analyzed and edited using ImageJ

software.

Wing size measurement

Flies were raised, and their wings were mounted and digitized as previously described (Shinoda

et al., 2019). Briefly, flies were placed on an acetic acid agar plate (2% agar, 1% sucrose, and

0.36% acetic acid) to lay eggs overnight. The plates were collected, and the eggs were incubated

at 25°C for 24 h. The 1st instar larvae were collected from the agar plate, and 30 larvae were

placed in one vial to precisely control the larval density. These flies were raised at 25°C. After

more than two days of hatching, the adult female flies were collected and submerged in 99.5%

ethanol to dehydrate for 24 hours. Dehydrated flies were dissected to mount their wings using

Euparal (Waldeck) with the dorsal side upward. Images were digitized using an upright

microscope (Leica DM5000B; Leica Microsystems) equipped with a CCD camera. The wing

size was measured using a Drosophila wing digital twin

[https://datamarkin.com/models/drosophila-wings-beta] developed by Datamarkin. Areas 0, 1,

and 2 were quantified as the anterior parts, whereas areas 4, 6, 7, and 8 were quantified as the

posterior parts, respectively.

Statistical analysis

The data are presented as the means ± standard error of measurement (SEM). *p*-values were

calculated using a one-way analysis of variance (ANOVA), followed by Sidak's multiple

comparison test for selected pairs (Figure 1–figure supplement 1B, C, D; Figure 5K), or

Dunnett's multiple comparison test using GraphPad Prism 8. NS: $p \geqq 0.05$, †: $p < 0.05$.

Acknowledgments

We would like to thank G. W. Davis, R. Carthew, B. Hay, C-H. Cheng, H. Richardson, S. K. Yoo, H. Steller, Y. Hiromi, H. D. Ryoo, the FlyORF Zurich ORFeome Project, Bloomington *Drosophila* Stock Center, Vienna Drosophila Resource Center, Kyoto Stock Center, and NIG-FLY for providing the fly strains. We thank Datamarkin for providing the Drosophila wing digital twin tool. We thank Dr. S. Hirayama, Dr. S. Murata, and the One-Stop Sharing Faculty Center for Future Drug Discovery at the Graduate School of Pharmaceutical Sciences, University of Tokyo, for the LC-MS/MS analysis. We thank Miura's lab members for their technical assistance and discussions; in particular, K. Takenaga for preparing the fly food and R. Takamoto for experimental support. We thank *Editage* [www.editage.com] for the English language editing of this manuscript. This work was supported by grants from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (KAKENHI Grant Numbers 19K16137, 21K15080, 22H05586, and 23K05747 to N.S. and 21H04774, 23H04766, 24H00567, and 25H01842 to M.M.), the Japan Agency for Medical Research and Development (AMED; Grant number JP21gm5010001 to M.M.), and grants from the Takeda Science Foundation and Sumitomo Foundation to N.S.

Author Contributions

N.S. and M.M. designed the study; N.S. conducted most of the experiments and analyzed most of the data; Y.H conducted protein structural predictions; N.H. analyzed the expression patterns of Drosophila caspases; and N.S. and M.M. wrote the manuscript.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M, Reiman D, Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**:493–500.

Ando R, Shimozono S, Ago H, Takagi M, Sugiyama M, Kurokawa H, Hirano M, Niino Y, Ueno G, Ishidate F, Fujiwara T, Okada Y, Yamamoto M, Miyawaki A. 2024. StayGold variants for molecular fusion and membrane-targeting applications. *Nat Methods* **21**:648–656.

Aram L, Yacobi-Sharon K, Arama E. 2017. CDPs: caspase-dependent non-lethal cellular processes. *Cell Death Differ* **24**:1307–1310.

Arama E, Agapite J, Steller H. 2003. Caspase activity and a specific cytochrome C are required for sperm differentiation in Drosophila. *Dev Cell* **4**:687–697.

Arama E, Bader M, Rieckhof GE, Steller H. 2007. A ubiquitin ligase complex regulates caspase activation during sperm differentiation in Drosophila. *PLoS Biol* **5**:e251.

Berthelet J, Dubrez L. 2013. Regulation of apoptosis by inhibitors of apoptosis (IAPs). *Cells* **2**:163–187.

Boucher D, Blais V, Denault J-B. 2012. Caspase-7 uses an exosite to promote poly(ADP ribose) polymerase 1 proteolysis. *Proc Natl Acad Sci U S A* **109**:5669–5674.

Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N, Ting AY. 2018. Efficient proximity labeling in living cells and organisms with TurboID. *Nat Biotechnol* **36**:880–887.

Choutka C, DeVorkin L, Go NE, Hou Y-CC, Moradian A, Morin GB, Gorski SM. 2017. Hsp83 loss suppresses proteasomal activity resulting in an upregulation of caspase-dependent compensatory autophagy. *Autophagy* **13**:1573–1589.

Desroches A, Denault J-B. 2019. Caspase-7 uses RNA to enhance proteolysis of poly(ADP-ribose) polymerase 1 and other RNA-binding proteins. *Proc Natl Acad Sci U S A* **116**:21521–21528.

DeVorkin L, Go NE, Hou Y-CC, Moradian A, Morin GB, Gorski SM. 2014. The Drosophila effector caspase Dcp-1 regulates mitochondrial dynamics and autophagic flux via SesB. *J Cell Biol* **205**:477–492.

Dietz L, Ellison CJ, Riechmann C, Cassidy CK, Felfoldi FD, Pinto-Fernández A, Kessler BM, Elliott PR. 2023. Structural basis for SMAC-mediated antagonism of caspase inhibition by the giant ubiquitin ligase BIRC6. *Science* **379**:1112–1117.

Ditzel M, Broemer M, Tenev T, Bolduc C, Lee TV, Rigbolt KTG, Elliott R, Zvelebil M, Blagoev B, Bergmann A, Meier P. 2008. Inactivation of effector caspases through nondegradative polyubiquitylation. *Mol Cell* **32**:540–553.

Domingues C, Ryoo HD. 2012. Drosophila BRUCE inhibits apoptosis through non-lysine ubiquitination of the IAP-antagonist REAPER. *Cell Death Differ* **19**:470–477.

Ehrmann JF, Grabarczyk DB, Heinke M, Deszcz L, Kurzbauer R, Hudecz O, Shulkina A, Gogova R, Meinhart A, Versteeg GA, Clausen T. 2023. Structural basis for regulation of apoptosis and autophagy by the BIRC6/SMAC complex. *Science* **379**:1117–1123.

Florentin A, Arama E. 2012. Caspase levels and execution efficiencies determine the apoptotic potential of the cell. *J Cell Biol* **196**:513–527.

Fujisawa Y, Shinoda N, Chihara T, Miura M. 2020. ROS Regulate Caspase-Dependent Cell
Delamination without Apoptosis in the Drosophila Pupal Notum. *iScience* **23**:101413.

Furuyama T, Banerjee R, Breen TR, Harte PJ. 2004. SIR2 is required for polycomb silencing
and is associated with an E(Z) histone methyltransferase complex. *Curr Biol* **14**:1812–
1821.

Gokcezade J, Sienski G, Duchek P. 2014. Efficient CRISPR/Cas9 plasmids for rapid and
versatile genome editing in Drosophila. *G3 (Bethesda)* **4**:2279–2282.

González M, Martín-Ruíz I, Jiménez S, Pirone L, Barrio R, Sutherland JD. 2011. Generation
of stable Drosophila cell lines using multicistronic vectors. *Sci Rep* **1**:75.

Haberman AS, Akbar MA, Ray S, Krämer H. 2010. Drosophila acinus encodes a novel
regulator of endocytic and autophagic trafficking. *Development* **137**:2157–2166.

Hou Y-CC, Chittaranjan S, Barbosa SG, McCall K, Gorski SM. 2008. Effector caspase Dcp-1
and IAP protein Bruce regulate starvation-induced autophagy during Drosophila
melanogaster oogenesis. *J Cell Biol* **182**:1127–1139.

Hunkeler M, Jin CY, Fischer ES. 2023. Structures of BIRC6-client complexes provide a
mechanism of SMAC-mediated release of caspases. *Science* **379**:1105–1111.

Ikegawa Y, Combet C, Groussin M, Navratil V, Safar-Remali S, Shiota T, Aouacheria A, Yoo
SK. 2023. Evidence for existence of an apoptosis-inducing BH3-only protein, sayonara,
in Drosophila. *EMBO J* **42**:e110454.

Jacomin A-C, Petridi S, Di Monaco M, Bhujabal Z, Jain A, Mulakkal NC, Palara A, Powell EL,
Chung B, Zampronio C, Jones A, Cameron A, Johansen T, Nezis IP. 2020. Regulation
of Expression of Autophagy Genes by Atg8a-Interacting Partners Sequoia, YL-1, and
Sir2 in Drosophila. *Cell Rep* **31**:107695.

Julien O, Wells JA. 2017. Caspases and their substrates. *Cell Death Differ* **24**:1380–1389.

Kaplan Y, Gibbs-Bar L, Kalifa Y, Feinstein-Rotkopf Y, Arama E. 2010. Gradients of a
ubiquitin E3 ligase inhibitor and a caspase inhibitor determine differentiation or death
in spermatids. *Dev Cell* **19**:160–173.

Kim Y-I, Ryu T, Lee J, Heo Y-S, Ahnn J, Lee S-J, Yoo O. 2010. A genetic screen for modifiers
of Drosophila caspase Dcp-1 reveals caspase involvement in autophagy and novel
caspase-related genes. *BMC Cell Biol* **11**:9.

Liu S-S, Jiang T-X, Bu F, Zhao J-L, Wang G-F, Yang G-H, Kong J-Y, Qie Y-F, Wen P, Fan
L-B, Li N-N, Gao N, Qiu X-B. 2024. Molecular mechanisms underlying the BIRC6-
mediated regulation of apoptosis and autophagy. *Nat Commun* **15**:891.

Muramoto M, Hanawa N, Okumura M, Chihara T, Miura M, Shinoda N. 2025. Executioner
caspase is proximal to Fasciclin 3 which facilitates non-lethal activation in Drosophila
olfactory receptor neurons. *Elife* **13**:RP99650.

Nakajima Y-I, Kuranaga E. 2017. Caspase-dependent non-apoptotic processes in development.
Cell Death Differ **24**:1422–1430.

Nandi N, Tyra LK, Stenesen D, Krämer H. 2014. Acinus integrates AKT1 and subapoptotic
caspase activities to regulate basal autophagy. *J Cell Biol* **207**:253–268.

Newman BL, Lundblad JR, Chen Y, Smolik SM. 2002. A Drosophila homologue of Sir2
modifies position-effect variegation but does not affect life span. *Genetics* **162**:1675–
1685.

Newton K, Strasser A, Kayagaki N, Dixit VM. 2024. Cell death. *Cell* **187**:235–256.

Nezis IP, Shravage BV, Sagona AP, Lamark T, Bjørkøy G, Johansen T, Rusten TE, Brech A,
Baehrecke EH, Stenmark H. 2010. Autophagic degradation of dBruce controls DNA
fragmentation in nurse cells during late Drosophila melanogaster oogenesis. *J Cell Biol*
190:523–531.

Obata F, Kuranaga E, Tomioka K, Ming M, Takeishi A, Chen C-H, Soga T, Miura M. 2014.
Necrosis-driven systemic immune response alters SAM metabolism through the FOXO-
GNMT axis. *Cell Rep* **7**:821–833.
Okuda S, Watanabe Y, Moriya Y, Kawano S, Yamamoto T, Matsumoto M, Takami T,
Kobayashi D, Araki N, Yoshizawa AC, Tabata T, Sugiyama N, Goto S, Ishihama Y.
2017. jPOSTrepo: an international standard data repository for proteomes. *Nucleic*
Acids Res **45**:D1107–D1111.
Qin W, Cho KF, Cavanagh PE, Ting AY. 2021. Deciphering molecular interactions by
proximity labeling. *Nat Methods* **18**:133–143.
Shinoda N, Hanawa N, Chihara T, Koto A, Miura M. 2019. Dronc-independent basal
executioner caspase activity sustains Drosophila imaginal tissue growth. *Proc Natl*
Acad Sci U S A **116**:20539–20544.
Shinoda N, Horikoshi M, Taira Y, Muramoto M, Hirayama S, Murata S, Miura M. 2023.
Caspase cleaves Drosophila BubR1 to modulate spindle assembly checkpoint function
and lifespan of the organism. *FEBS J* **290**:4200–4223.
Shinoda N, Miura M. 2024. Exploring caspase-dependent non-lethal cellular processes using
Drosophila. *Front Cell Death* **3**:1472108.
Taira Y, Yang Z, Miyata Y, Shirasaki Y, Segawa K, Miura M, Shinoda N. 2025. Early-peaking
caspase-7 activity at the plasma membrane drives apoptotic phosphatidylserine
exposure. *bioRxiv*. doi:10.1101/2025.03.17.643603
Tyra LK, Nandi N, Tracy C, Krämer H. 2020. Yorkie growth-promoting activity is limited by
Atg1-mediated phosphorylation. *Dev Cell* **52**:605-616.e7.
Vaux DL, Silke J. 2005. IAPs, RINGs and ubiquitylation. *Nat Rev Mol Cell Biol* **6**:287–297.
Walsh JG, Cullen SP, Sheridan C, Lüthi AU, Gerner C, Martin SJ. 2008. Executioner caspase-
3 and caspase-7 are functionally distinct proteases. *Proc Natl Acad Sci U S A*
105:12815–12819.
Wu G, Chai J, Suber TL, Wu JW, Du C, Wang X, Shi Y. 2000. Structural basis of IAP
recognition by Smac/DIABLO. *Nature* **408**:1008–1012.
Xie HB, Golic KG. 2004. Gene deletions by ends-in targeting in Drosophila melanogaster.
Genetics **168**:1477–1489.
Xu D, Li Y, Arcaro M, Lackey M, Bergmann A. 2005. The CARD-carrying caspase Dronc is
essential for most, but not all,developmental cell death in *Drosophila*. *Development*
132:2125–2134.
774

 Figures and figure legends

 Figure 1

777
Figure 1. *Dcp-1* overexpression induces apoptosis independent of the apoptosome.
(A) Schematic structures and intrinsically disordered region prediction of Dcp-1 (dark cyan)
and Drice (red). The predicted disordered score plot is shown. Regions wherein the score is
more than 0.5 are predicted to be disordered.
(B) Representative images of female wing phenotypes upon caspase overexpression using *WP-*
Gal4.
(C) Representative images of larval wing imaginal disc upon *3xHA-tagged executioner caspase*
(magenta) overexpression using *WP-Gal4*. Nuclei are visualized by Hoechst 33342 (green).
Magnified images are shown in right below. Arrowheads indicate condensed nuclei. Scale bar:
50 μm (top) and 10 μm (bottom).
(D) Schematic diagram of canonical apoptosis signaling in *Drosophila*.

(E, F) Representative images of female wing phenotypes upon *Dcp-1::VENUS* overexpression
using *WP-Gal4* with the genetic perturbation of canonical apoptosis signaling.
(G) Quantification of (E) and (F). Each wing is manually classified into four (wingless, severe,
mild, and intact) categories. Schematic images of the four categories are shown on the right.
Sample sizes are shown in the figure.
794

Figure 1–figure supplement 1

Figure 1–figure supplement 1. Dcp-1 overexpression induces apoptosis in *Drosophila* S2 cells.

(A) Representative images of *Drosophila* S2 cells transfected with mNeonGreen (mNG)-tagged wild-type (WT) or catalytically inactive (catalytic cysteine mutated to glycine; CG) Dcp-1 and Drice. Scale bars: 50 μm (left) and 5 μm (right).

(B) Quantification of mNG intensity for each cell, shown as a violin plot. Quartiles are indicated by black broken lines, and the median is indicated by a red bar. (B′) Magnified view of the data in (B), with the Y-axis limited to 0–40. *p*-values were calculated using one-way analysis of variance (ANOVA) followed by Sidak's multiple-comparison test for selected pairs. NS, $p > 0.05$; †, $p < 0.05$. Sample sizes: mNG (n = 712 cells), Dcp-1WT::mNG (n = 212 cells), Dcp-1CG::mNG (n = 380 cells), DriceWT::mNG (n = 434 cells), and DriceCG::mNG (n = 442 cells).

(C) Quantification of the average mNG intensity in each field of view. Data are presented as mean ± SEM. *p*-values were calculated using one-way ANOVA followed by Sidak's multiple-comparison test for selected pairs. NS, $p \geqq 0.05$; †, $p < 0.05$. Sample size for each condition is 4.

812   (D) Quantification of the mNG-positive cell number in each field of view. Data are presented
813   as mean ± SEM. p-values were calculated using one-way ANOVA followed by Sidak's
814   multiple-comparison test for selected pairs. NS, $p \geqq 0.05$; †, $p < 0.05$. Sample size for each
815   condition is 4.
816

Figure 2

818
Figure 2. Dcp-1-enriched proximal proteins are required for Dcp-1 activation.
(A) Western blotting of expression of C-terminally V5::TurboID knocked-in tagged caspases
and biotinylation patterns of their proximal proteins detected by streptavidin (SA) in larval wing
imaginal discs.
(B) A schematic diagram of the TurboID-mediated identification of proximal proteins. Dcp-1-
and Drice-proximal proteins (Protein C, D, E, F, G) are promiscuously labeled in vivo with the
administration of biotin. Then, biotinylated proteins are purified using NeutrAvidin magnetic
beads and are subsequently analyzed by mass spectrometry.
(C) A summary of mass spectrometry analysis of proteins in the larval imaginal discs. Among
818 proteins identified, 230 proteins were detected as highly enriched proteins in
Drice::V5::TurboID flies (FC [Drice::V5::TurboID/Dcp-1::V5::TurboID] > 2) and 16 proteins

were detected as highly enriched proteins in Dcp-1::V5::TurboID flies flies (FC [Dcp-
1::V5::TurboID/Drice::V5::TurboID] > 2).

(D) Western blotting validation of the anti-CCT8 antibody in whole larval lysates without the
digestive tract and fat body. The anti-CCT8 antibody detects both endogenous CCT8 (MW:
59.4 kDa) and 3xFLAG::StrepII::VENUS::StrepII (FSVS)-tagged CCT8 (MW: 91.1 kDa).

(E) Western blotting of endogenous CCT8 expressed in larval wing imaginal disc. Dcp-1-
proximal and Drice-proximal proteins biotinylated by C-terminally knocked-in tagged
V5::TurboID are purified using NeutrAvidin. The red asterisk represents a non-specific band
in the marker lane.

(F) Schematic diagram of the RNAi screen targeting eight Dcp-1-enriched proximal proteins.
The *WP > Dcp-1::VENUS* line was crossed with each *UAS-RNAi* line, and wing phenotypes of
the F1 progeny were scored.

(G) Representative images of female wing phenotypes from *Dcp-1::VENUS* overexpression
driven by *WP-Gal4* with knockdown of Dcp-1-enriched proximal proteins. "XX" denotes the
knocked-down gene, and "#1" and "#2" indicate independent RNAi lines.

(H) Representative images of female wing phenotypes upon *Dcp-1::VENUS* overexpression
driven by *WP-Gal4* in the *Sirt1* null mutant background.

(I) Quantification of (H). Each wing is manually classified into four (wingless, severe, mild,
and intact) categories. Sample sizes are shown in the figure.

Figure 2–figure supplement 1

Figure 2–figure supplement 1. Expression patterns of *Drosophila* caspases in larval and

adult tissues.

(A) Western blotting of expression of C-terminally V5::TurboID knocked-in tagged caspases
and biotinylation patterns of their proximal proteins detected by streptavidin (SA) in larval
salivary gland, fat body, and brain.
(B) Western blotting of expression of C-terminally V5::TurboID knocked-in tagged caspases
and biotinylation patterns of their proximal proteins detected by SA in adult thorax, ovary, and
testis.
860

 Figure 2–figure supplement 2

Figure 2–figure supplement 2. RNAi screening of Dcp-1-enriched proximal proteins for

regulators of Dcp-1 activation.

(A) Quantification of female wing phenotypes upon knockdown of Dcp-1-enriched proximal

proteins driven by *WP-Gal4*. "XX" denotes the knocked-down gene, and "#1" and "#2" indicate

independent RNAi lines. Each wing is manually classified into four (wingless, severe, mild,

and intact) categories. Sample sizes are shown in the figure.

(B) Quantification of female wing phenotypes upon *Dcp-1::VENUS* overexpression driven by

WP-Gal4 with simultaneous knockdown of Dcp-1-enriched proximal proteins. "XX" denotes

the knocked-down gene, and "#1" and "#2" indicate independent RNAi lines. Each wing is

manually classified into four (wingless, severe, mild, and intact) categories. Sample sizes are

shown in the figure.

Figure 3

Figure 3. Debcl, Buffy, and autophagy are specifically required for *Dcp-1* overexpression-induced apoptosis.

(A) Representative images of female wing phenotypes from *Dcp-1::VENUS* overexpression driven by *WP-Gal4* with the genetic perturbation of *Synr*, *Debcl*, *Buffy,* and autophagy. "XX" denotes the knocked-down gene, and "#1" and "#2" indicate independent RNAi lines.

(B) Quantification of (A). "XX" denotes the knocked-down gene, and "#1" and "#2" indicate independent RNAi lines. Each wing is manually classified into four (wingless, severe, mild, and intact) categories. Sample sizes are shown in the figure.

(C) Representative images of female eye phenotypes upon *Hid* overexpression with the genetic perturbation of canonical apoptosis signaling.

(D) Representative images of female eye phenotypes upon *Hid* overexpression with the genetic perturbation of *Sirt1* or *Fkbp59*. "#1" and "#2" indicate independent RNAi lines.

(E) Representative images of female eye phenotypes upon *Hid* overexpression with the genetic perturbation of *Debcl*, *Buffy*, and autophagy.

Figure 4

Figure 4. Cleaved Dcp-1, but not Drice, interacts with Bruce.

(A) Domain structure of Bruce. LD denotes the linker domain, WD40 a WD40 domain, BIR the Baculoviral IAP repeat, β-X a β-sandwich-like fold, CBM a domain resembling the carbohydrate-binding module family 32 (CBM), UBL the ubiquitin-like domain, and UBC the E2-E3 hybrid ubiquitin-conjugating (UBC) domain. The region used for structural prediction using AlphaFold3 is indicated in red.

(B) IBM-like sequence in Dcp-1. The propeptide cleavage site of Dcp-1 is indicated by a black
dashed line; the caspase recognition motif is highlighted in black; the IBM-like sequence is
highlighted in pink; and the region used for AlphaFold3 prediction is shown in red.
(C) IBM-like sequence in Drice. The propeptide cleavage site of Drice is indicated by a black
dashed line; the caspase recognition motif is highlighted in black; the IBM-like sequence is
highlighted in pink; and the region used for AlphaFold3 prediction is shown in red.
(D) Predicted complex structure of Bruce BIR and Dcp-1 IBM-like sequence. The AlphaFold3
prediction [model 0] is shown. The Bruce β-X2 and BIR domains are displayed as surfaces in
green and pink, respectively. Dcp-1 is shown in stick representation and color-coded according
to pLDDT scores, as indicated in the figure. See also panel (F).
(E) Predicted complex structure of Bruce BIR and Drice IBM-like sequence. The AlphaFold3
prediction [model 0] is shown. The Bruce β-X2 and BIR domains are displayed as surfaces in
green and pink, respectively. Drice is shown in stick representation and color-coded according
to pLDDT scores, as indicated in the figure. See also panel (G).
(F) pLDDT scores of Dcp-1 in the five predicted complex structures with Bruce. For all five
AlphaFold3 models, per-residue pLDDT scores of Dcp-1 are plotted. Prediction confidence is
color-coded as follows: very high (pLDDT > 90) in dark blue, confident (90 > pLDDT > 70)
in light blue, low (70 > pLDDT > 50) in yellow, and very low (pLDDT < 50) in orange.
(G) pLDDT scores of Drice in the five predicted complex structures with dBruce. For all five
AlphaFold3 models, per-residue pLDDT scores of Drice are plotted. Prediction confidence is
color-coded as follows: very high (pLDDT > 90) in dark blue, confident (90 > pLDDT > 70)
in light blue, low (70 > pLDDT > 50) in yellow, and very low (pLDDT < 50) in orange.
(H) Co-immunoprecipitation of N-terminally 3xFLAG-tagged N-terminal region of Bruce (Br,
1–900 aa; containing LD, BIR, and WD40 domains) with C-terminally myc-tagged,
catalytically inactive Dcp-1 or Drice (catalytic cysteine mutated to glycine; CG) expressed in
Drosophila S2 cells. N-terminally 3xFLAG-tagged mRuby3 (mRb3) was used as a control.
Apoptosis was induced by cycloheximide (CHX) treatment. 3xFLAG-tagged proteins were
immunoprecipitated using an anti-3xFLAG antibody.
928

929   **Figure 4–figure supplement 1**

930
Figure 4–figure supplement 1. Predicted structures of Bruce complexed with Dcp-1 and

Drice IBM-like sequences.

(A) Overall view of the predicted complex structure of Bruce and Dcp-1. Model 0 is shown
with the same color scheme as in Figure 4A and B. The predicted position error is shown in
(A').
(B) Overall view of the predicted complex structure of Bruce and Drice. Model 0 is shown with
the same color scheme as in Fig. 4A and C. The predicted position error is shown in (B').
(C) Structural comparison between the experimentally determined complex of XIAP BIR3 and
Smac IBM (C), and the predicted complex of dBruce BIR and Dcp-1 IBM (C'). XIAP BIR3 is
shown as a pink cartoon and the Smac IBM as green sticks (PDB ID: 1G73, right panel). dBruce
BIR domain is shown as a pink cartoon and the Dcp-1 IBM as dark green sticks (model 0, left
panel). Zn^{2+} ions are shown as gray spheres. Salt bridges are indicated by black dashed lines,
along with the corresponding interacting residues.

Figure 5

Figure 5. Bruce selectively suppresses Dcp-1 activity and wing tissue growth.

(A) Expression pattern of overexpressed *myc::Bruce* (green) by using *WP-Gal4* in the wing

imaginal discs. Nuclei are visualized using Hoechst 33342 (magenta). Scale bar: 50 µm.

(B) Representative images of female wing phenotypes from *Dcp-1::VENUS* overexpression
driven by *WP-Gal4* with *DIAP1* or *Bruce* overexpression.
(C) Quantification of (B). Each wing is manually classified into four (wingless, severe, mild,
and intact) categories. Sample sizes are shown in the figure.
(D) Representative images of female wing phenotypes upon *Synr* overexpression driven by
Nub-Gal4 with *DIAP1* or *Bruce* overexpression.
(E) Representative images of female eye phenotypes upon *Hid* overexpression with *DIAP1* or
Bruce overexpression.
(F) Representative images of female wing upon *Bruce* or *Bruce-RNAi* (*Bruce-i*) overexpression
using *WP-Gal4*. Scale bar: 1 mm.
(G) Quantification of relative wing size normalized to the corresponding *No-Gal4* control. Data
are presented as mean ± SEM. *p*-values were calculated using one-way analysis of variance
(ANOVA) followed by Dunnett's multiple-comparison test against no UAS control. NS, $p \geqq$
0.05; †, $p < 0.05$. Sample sizes are shown in the figure.
(H) Representative images of female wing upon *Bruce* or *Bruce-RNAi* (*Bruce-i*) overexpression
using *En-Gal4*. The grey area (labelled as A) represents the anterior part, and the green area
(labelled as P) represents the posterior part of the wing. Scale bar: 1 mm.
(I) Quantification of relative posterior part of wing size normalized to corresponding anterior
part of wing size. Data are presented as mean ± SEM. *p*-values were calculated using one-way
ANOVA followed by Dunnett's multiple comparison test against no UAS control. †, $p < 0.05$.
Sample sizes are shown in the figure.
(J) Representative images of female wing upon InR^{DN} and *Bruce-RNAi* (*Bruce-i*)
overexpression using *WP-Gal4*. Scale bar: 1 mm.
(K) Quantification of relative wing size normalized to control (*WP > +*). Data are presented as
mean ± SEM. p-values were calculated using one-way ANOVA followed by Sidak's multiple
comparison test for selected pairs.†, $p < 0.05$. Sample sizes are shown in the figure.
(L) Schematic diagram of Dcp-1 activity-regulating alternative apoptosis signaling pathway in
Drosophila mediated by autophagy-Bruce axis.
977

Figure 5–figure supplement 1

Figure 5–figure supplement 1. Expression patterns of mStayGold::V5-tag knocked-in Bruce.

(A) Schematic diagram of CRISPR/Cas9-mediated knock-in of mStayGold::C4 linker (C4)::V5 tag at the N-terminus of Bruce. A 3xP3-DsRed cassette was integrated at the endogenous TTAA site as a transformation marker. Removal of the cassette restores the genomic TTAA site, enabling scarless genome engineering.

(B) Expression pattern of mStayGold (mSG)::C4::V5-tag knocked-in Bruce (green) in the ovariole. Nuclei are visualized using Hoechst 33342 (magenta). Scale bar: 50 μm.

988 (C) Expression pattern of mSG::C4::V5-tag knocked-in Bruce (green) in the testis. Nuclei are
989 visualized using Hoechst 33342 (magenta). Scale bar: 50 µm.
990 (D) Western blotting of expression of mSG::C4::V5-tag knocked-in tagged Bruce in the whole
991 adult male and female.
992 (E) Expression pattern of mSG::C4::V5-tag knocked-in tagged Bruce (green) in the wing
993 imaginal discs. Nuclei are visualized using Hoechst 33342 (magenta). Scale bar: 50 µm.
994 (F) Expression pattern of mSG::C4::V5-tag knocked-in tagged Bruce (green) in the wing
995 imaginal discs upon *Bruce-RNAi* using *WP-Gal4*. Nuclei are visualized using Hoechst 33342
996 (magenta). Scale bar: 50 µm.
997

998   **Legends for Supplemental Tables**
999

Table S1. Dcp-1 and Drice proximal protein lists.
Lists of the identified Dcp-1 proximal proteins (abundance ratio (Dcp-
1::V5::TurboID/Drice::V5::TurboID) > 2), Drice proximal proteins (abundance ratio
(Drice::V5::TurboID/Dcp-1::V5::TurboID) > 2), and candidates for RNAi screening
(abundance ratio (Dcp-1::V5::TurboID/Drice::V5::TurboID) > 1.5, abundance ratio (Dcp-
1::V5::TurboID/Control) > 10, PSM $\geqq$ 2). The UniProt accessions, gene description,
abundance ratio, and # PSMs are listed.
1007

1008   **Table S2. Fly stock list.**
1009   List of fly strains. Genotypes, maps, source or references, and stock numbers are listed.
1010

1011   **Table S3. Detailed genotypes.**
1012   Detailed genotypes and their related figure # are presented.
1013

1014   **Table S4. Liquid chromatography (LC) settings.**
1015   Gradient settings for LC analysis are presented.
1016

1017   **Table S5. Tandem mass spectrometry (MS/MS) settings.**
1018   Settings used for the MS/MS analysis are presented.
1019

1020   **Table S6. Proteome Discoverer 2.2 settings.**
1021   Proteome Discoverer 2.2 settings for proteomics analysis are presented.
1022