1 # Decoding emergent properties of microbial community functions through
2 # sub-community observations and interpretable machine learning
3

4 Hidehiro Ishizawa[1*], Sunao Noguchi[1], Miku Kito[1], Yui Nomura[2], Kodai Kimura[2],
5 Masahiro Takeo[1]
6
7 Short title: Dissecting emergent microbiome functions
8
9 [1]Department of Applied Chemistry, Graduate School of Engineering, University of
10 Hyogo, Himeji, Hyogo, 671-2280, Japan
11 [2]Department of Applied Chemistry and Chemical Engineering, School of Engineering,
12 University of Hyogo, Himeji, Hyogo, 671-2280, Japan
13
14 *Corresponding author
15 Hidehiro Ishizawa
16 Address: Shosha 2167, Himeji, Hyogo, 671-2280 Japan
17 Phone: (+81)79-267-4969
18 Mail to: ishizawahidehiro@gmail.com
19

**ABSTRACT**

The functions of microbial communities, including substrate conversion and pathogen suppression, arise not as a simple sum of individual species' capabilities but through complex interspecies interactions. Understanding how such functions arise from individual species and their interactions remains a major challenge, limiting efforts to rationally understand microbial roles in both natural and engineered ecosystems. Because current holistic (meta-omics) and reductionist (isolation- or single-cell-based) approaches struggle to capture these emergent microbial community functions, this study explores an intermediate strategy: analyzing simple sub-community combinations to enable a bottom-up understanding of community-level functions. To examine the validity of this approach, we used a nine-member synthetic microbial community capable of degrading the environmental pollutant aniline, and systematically generated a dataset of 256 sub-community combinations and their associated functions. Analyses using random forest models revealed that the sub-community combinations of just three to four species enabled the quantitative prediction of functions in larger communities (5–9-member; Pearson's $r = 0.78$–$0.80$). Prediction performance remained robust even with limited sub-community data, suggesting applicability to more diverse microbial communities where exhaustive sub-community observation is infeasible. Moreover, interpreting models trained on these simple sub-community combinations enabled the identification of key species and interspecies interactions that strongly influence the overall community function. These findings provide a methodological framework for mechanistically dissecting complex microbial community functions through sub-community-based analysis.

## INTRODUCTION

Microbial communities constitute a significant proportion of Earth's total biomass (approximately 90 Gt-C) [1], encompassing diverse, interacting species. Far from being mere background organisms, they are key drivers of climate regulation, biogeochemical cycling, and phenotypic expression in multicellular hosts [2-4]. Their diverse metabolic capabilities are increasingly being applied in agriculture, bioproduction, and environmental remediation [5-6]. However, despite decades of research, our ability to rationally manipulate microbial community functions remains limited.

Difficulties in microbial community management largely stem from the emergent nature of microbial community functions, which are shaped by complex interspecies interactions rather than the additive effects of individual species' capabilities [7]. For instance, in pollutant-degrading microbial communities, species lacking genetic capacity for direct degradation still influence overall system performance by supporting downstream metabolism, modulating population dynamics, or providing essential cofactors [8-10]. Existing analytical frameworks often struggle to capture such indirect contributions. Holistic meta-omics approaches provide comprehensive data but cannot easily resolve causal relationships between specific species and functional outcomes [11-12]. Reductionist approaches using isolates or single-cell data often miss the interspecies interactions that drive emergent properties. Consequently, understanding interspecies interactions and the true functional contributions of individual species remains challenging, hindering our ability to predict, design, and control microbial community functions.

A promising approach to dissect complex microbial interactions is to focus on sub-community combinations as an intermediate unit of observation. This approach involves sampling multiple combinations of a few species, typically three to five, from a larger community and evaluating their functional outputs. Unlike statistical inference from whole-community data, this approach facilitates the examination of species' functional roles in simpler and more interpretable contexts, with still capturing emergent properties not detectable in isolates or single-cell assays. The conceptual basis for this approach derives from the theoretical food web ecology, where ecosystem-level properties have been modeled by decomposing complex systems into small ecological

modules of three to four species [13-15]. Empirical studies have also shown that

interactions among as few as three species can recapitulate emergent behaviors relevant

to community assembly in both plant and microbial systems [16-18].

Existing studies have primarily focused on predicting species abundance or

community composition, leaving open the question of how sub-community data can be

leveraged to dissect functional outputs, such as substrate conversion or host phenotypic

modulation. To address this gap, this study aimed to establish an analytical framework

that uses sub-community information to achieve bottom-up, mechanistic understanding

of microbial community functions. Specifically, a machine learning approach was

applied to explain community-level functional performance based on measurements

from sub-community combinations. Although previous studies have trained machine

learning models on whole community data to predict functional outputs [19-20], we

expected that models trained on sub-community data would further highlight influential

species and interspecies interactions because of their greater simplicity and

interpretability.

To evaluate the sub-community-based analytical framework in a controlled

setting, we constructed a nine-member synthetic microbial community capable of

degrading the environmental pollutant aniline. Aniline, a representative aromatic amine

widely used to produce pesticides and dyes, is considered a high-priority bioremediation

target due to its toxicity and environmental persistence [21]. Using this model system,

we systematically assessed all possible sub-community combinations to test two

hypotheses: (1) functional outputs of more complex communities can be predicted from

simpler sub-community data, and (2) sub-community observation allows identification

of key species and interspecies interactions driving emergent community functions.

Although derived from a controlled synthetic system, our findings provide a

methodological blueprint for leveraging sub-community data to gain mechanistic insight

into complex microbial community functions.

**MATERIALS AND METHODS**

**Establishment of a reproducible aniline-degrading community**

To establish a model aniline-degrading microbial community, 2.5 g of soil collected from a household vegetable garden in Kakogawa City, Hyogo, Japan (34.77 N 134.85 E), was suspended in 30 mL of MSM medium [22], mixed vigorously, and centrifuged (1,000 × $g$, 1 min). Then, 5 mL of the supernatant containing microbial cells was transferred to 50 mL of MSM medium supplemented with 0.5 mM aniline (hereafter referred to as MSM aniline medium) and incubated at 30°C with shaking at 150 rpm for 48 h. Subsequently, 1 mL of this culture was inoculated into 50 mL of fresh MSM aniline medium and incubated for four days. During cultivation, 1 mL samples were periodically collected to measure residual aniline concentrations using HPLC (Table S1). The resulting enriched culture was preserved at −80°C in cryotubes containing 16.7% glycerol.

For community restoration, 1.2 mL of thawed glycerol stock was added to 10 mL of MSM medium and microbial cells were collected via centrifugation (10,000 × $g$, 4°C, 10 min), washed twice, and resuspended in 5 mL of MSM. Then, 1 mL of the cell suspension was inoculated into 50 mL of MSM aniline medium and incubated at 30°C with shaking at 150 rpm for 48 h. The resulting culture was used for subsequent analyses. To assess the community structure and aniline degradation ability, 1 mL of the culture was inoculated into 50 mL of MSM aniline medium and incubated in four replicate flasks at 30°C with shaking at 150 rpm for 48 h. Reproducibility was assessed by performing the restoration procedure twice on separate days, followed by the evaluation of residual aniline concentration using HPLC and community structure using 16S rRNA gene amplicon sequencing (methods described in Supplementary Note 1).

**Metagenome analysis**

DNA was extracted from the restored aniline-degrading community using NucleoSpin Microbial DNA (Macherey-Nagel, Düren, Germany). For shotgun metagenome sequencing, libraries were prepared with the MGIEasyFS DNA Library Prep Set (MGI Tech, Shenzhen, China), and 150 bp paired-end sequencing was performed on a DNBSEQ-G400RS platform (MGI Tech) at Genome-Lead Inc. (Takamatsu, Japan). Sequencing reads were quality-filtered and adapter-trimmed with fastp v0.23.2 and assemblies were generated using metaSPAdes v3.15.5 [23]. Subsequently, reads were

mapped to assembled scaffolds using BWA v0.7.17 [24], and metagenome binning was performed using MetaBATt2 v2.15 [25]. The resulting bins were evaluated using checkM v1.2.2 [26], and those with completeness >50% and contamination <5% were regarded as valid metagenome-assembled genomes (MAGs). MAGs were annotated using DFAST v1.6.0 [27] and BlastKOALA v3.0 [28]. The relative abundances of the MAGs were estimated by mapping the metagenome reads using BBMap v36.01.1. Homology search and gene feature visualizations were performed using the BLAST and CAGECAT [29].

**Isolation, identification, and characterization of synthetic community members**

The restored model aniline-degrading community was diluted, spread on R2A and PCAT agar [30], and incubated at 30°C for up to one week. Morphologically distinct colonies were isolated and subjected to PCR amplification of the 16S rRNA gene using the 27F–1392R primer set [31]. Amplicons were purified using the NucleoSpin Gel and PCR Purification Kit (Macherey-Nagel) and sequenced via Sanger sequencing on a 3730xl DNA analyzer (Applied Biosystems, Foster City, CA, USA) at Macrogen Japan Corp. (Tokyo, Japan). The taxonomic positions of the isolates were identified using similarity searches against the EzBioCloud 16S database. Based on these results, nine bacterial strains were selected as members of the synthetic community: *Achromobacter veterisilvae* MA031 (Ach031), *Pseudomonas monteilii* MA033 (Pse033), *Sphingomonas pseudosanguinis* MA034 (Sph034), *Microbacterium laevaniformans* MA039 (Mic039), *Bosea spartocytisi* (Bos046), *Bacillus mexicanus* MA050 (Bac050), *Pseudomonas* sp. MA056 (Pse056), *Caballeronia cordobensis* MA074 (Cab074), and *Sphingomonas* sp. MA075 (Sph075) (Table S2). Given the absence of rRNA gene sequences in our MAG sequences, we predicted the closest MAG–isolate correspondence by comparing species-level classifications obtained independently for MAGs and isolates. The nine strains were cultivated overnight in 2 mL R2A medium in a 24-well plate (30°C, 150 rpm), washed twice with MSM medium, and used for subsequent experiments.

For one isolate (Cab074), identified as the primary aniline degrader, a fluorescence marker (eGFP) was introduced using the pMRE152 plasmid as previously

described [32]. The obtained fluorescent transformants (hereafter Cab074F) were

purified and used as the member of synthetic communities in place of the wildtype.

To evaluate the ability of each synthetic community member to utilize aniline,

individual strains were inoculated into 500 µL MSM aniline medium in a 24-well plate

at an initial $OD_{600}$ of 0.01. After 48 h of incubation, the residual aniline content was

quantified using HPLC.

**Full factorial evaluation of the sub-community combinations**

A 384-well plate-based assay was conducted to evaluate all possible sub-community

combinations of the synthetic community that included Cab074F (i.e., $2\square = 256$

patterns; $n = 4$). Each well of the black 384-well plate contained $20\,\square\mu L$ of MSM

aniline medium supplemented with $2\,\square\mu g/mL$ of resazurin, a metabolic marker reduced

to fluorescent resorufin in proportion to microbial aerobic metabolism [33]. Because

aniline was the sole carbon and energy source, fluorescence intensity was considered a

proxy for aniline utilization. Strains were inoculated into wells at an initial $OD\,\square\square\square$ of

0.001 each. To minimize pipetting errors, a robotic pipettor (Assist Plus, Integra

Biosciences, Zizers, Switzerland) was used for culture preparation. The plates were

incubated in a microplate reader (Synergy HTX, Agilent Technologies, Santa Clara, CA,

USA) at 30°C with orbital shaking at 237 rpm. During the 72 h cultivation period, GFP

(excitation/emission: 485/528 nm) and resorufin (528/590 nm) fluorescence were

measured every 10 min. Three plates were analyzed to cover all sub-community

combinations in four replicates each. Every plate included at least 16 Cab074F-only and

16 blank wells without microbial inoculation.

The "aniline-utilization activity" and the "Cab074F growth" in each well were

defined as the area under the curve (AUC) calculated from the fluorescence time

courses of resorufin and GFP during cultivations, respectively. The detailed calculation

method is described in Supplementary Note 2. We also checked the correlation between

aniline degradation rate and resorufin fluorescence, as well as between colony-forming

unit of Cab074F and GFP fluorescence, using selected combinations (Supplementary

Note 3).

The functional effect of each non-degrader strain was calculated as the change

in aniline-utilization activity between sub-communities that differed only by the

presence or absence of the focal strain. The Shapley value, representing the overall

contribution of each species to community function, was calculated as the average

functional effect across all possible assembly pathways [34]. The functional interactions

between all pairs of non-degrader strains were estimated as the difference in the mean

functional effect of a focal strain in the presence versus absence of a partner strain.

Statistical significance of differences in functional effect was assessed using two-sided

*t*-test in R v4.4.2.

**Random forest model**

Random forest models were employed to examine the bottom-up predictability of

aniline-utilization activity. Models were constructed using the "randomForest" R

package, with the presence–absence data of eight non-degrader strains as explanatory

variables and the measured aniline-utilization activity as the response variable. We

chose species presence/absence rather than their abundances, because species

abundances in microbial communities are highly dynamic, and relying on single

snapshot can introduce sampling bias that does not reflect overall system performance.

215            To determine which levels of sub-community diversity (species richness) carry

important information for predicting community-level functions, we first trained models

with observed sub-community combinations at fixed richness levels: two (n = 8), three

(n = 28), four (n = 56), five (n = 70), six (n = 56), and seven-member combinations (n =

28). Prediction performance was then evaluated as Pearson's correlation coefficient

using data from 5–9-member combinations (n = 163) as the test set. To prevent data

leakage, when a test combination is also present in the training pool, we rebuilt the

model after removing only that single overlapping data point from the training data.

Furthermore, to assess model performance under limited training data, models were also

trained on using randomly sampled subsets comprising 50% or 25% of the original

training set.

226            To identify important species and interspecies interactions contributing to

prediction accuracy, permutation importance and Friedman's H statistic were extracted

from the random forest models using the "vivid" R package [35]. Models trained on all

sub-community combinations at each richness level (two-to six-members) and a model

trained on all 2–4-member sub-communities were used for this analysis. The obtained

values were compared with empirically calculated Shapley value and functional

interaction strength to evaluate their ability to identify key species and interactions.

To account for variability in training and sampling, the entire procedure

(training and evaluation) was repeated 100 times with different random seeds; we report

the mean and standard deviation across runs. Figures and parameter inference were

generated from the model with median performance (51st of 100 trials).

**Data availability**

The 16S rRNA gene sequences of the bacterial isolates were deposited in the

DDBJ/EMBL/GenBank database under accession numbers LC878797-LC878805.

Metagenome sequence reads were deposited under the accession number DRR707044.

All experimental data and analytical codes have been deposited in Zenodo [36] .

**RESULTS**

**Reproducibility of the model aniline-degrading microbial community**

A major challenge in systematic analysis of microbial communities is the lack of

reproducibility (i.e., the ability to regenerate the same community across experiments).

To address this, we first established a reproducible community using a

whole-community glycerol stock derived from an aniline-degrading community.

Restoring the preserved stock through a standardized procedure confirmed consistent

community degradation of 0.5 mM aniline within 48 h and a stable community

structure dominated by specific genera including *Achromobacter*, *Burkholderia*, and

*Pseudomonas*, with only minor variations in relative abundances (Fig. S1). The

dominant bacteria in the restored communities were consistent at the ASV level,

suggesting that the preserved community stock provides a reliable source of

reproducible model aniline-degrading communities.

**Sequence-based prediction of aniline- and catechol-degrading bacteria**

Shotgun metagenome analysis was performed to identify species in the

aniline-degrading community harboring genes responsible for aniline degradation. The assembly of approximately 107 million reads reconstructed 13 MAGs, encompassing all dominant taxa detected via the 16S rDNA amplicon sequencing (Fig. 1A; Table S3).

Reportedly, aniline degradation is initiated by a unique reaction in which a glutamate synthase-like protein (AtdA1) attaches a glutamate molecule to aniline, forming γ-glutamylanilide. Subsequently, dioxygenase proteins (AtdA3–5) oxidize the aromatic ring, yielding catechol, glutamate, and ammonium [37-38] (Fig. 1B). Genes encoding these enzymes are often clustered with *atdA2*, encoding the reverse reaction of AtdA1. Catechol, a common intermediate in aromatic compound biodegradation is typically introduced into central anabolic metabolism via either the ortho- or meta-cleavage pathways [39].

Within the obtained MAGs, only MAG01, assigned to the genus *Caballeronia*, contained a gene with significant similarity to known *atdA1* genes (43.6% amino acid sequence similarity to AtdA1 of *Acinetobacter* sp. YAA) [37]. Adjacent genes encoded components of an aromatic ring-hydroxylating dioxygenase (a functional analog of AtdA3–5) and a glutamyl hydrolase (a functional analog of AtdA2) (Fig. 1C). Although these surrounding genes exhibited <20% similarity to previously reported aniline-degrading gene clusters, we considered this gene cluster responsible for aniline degradation in our community.

For catechol degradation, several MAGs dominant in the community—including those affiliated with *Caballeronia*, *Achromobacter*, *Pseudomonas*, and *Pseudorhodopherax*—possessed most genes involved in the ortho-cleavage pathway (Fig. 1D). In contrast, no MAG carried the candidate genes for 2-keto-4-pentenoate hydratase (K02554), a key enzyme in the meta-cleavage pathway, indicating that catechol degradation in this community proceeds primarily through the ortho-cleavage pathway.

In summary, the metagenomic analysis suggests a potential metabolic relay within the model aniline-degrading community, in which *Caballeronia* first converts aniline into catechol, whereas several other strains contribute to its subsequent metabolism. Bacteria lacking the ability to degrade neither aniline nor catechol may persist through metabolic cross-feeding or scavenging of cell debris.

**Fig. 1** Metagenomic characterization of the model aniline-degrading community served

as the model system in this study. A) Community composition of the aniline-degrading

community. Relative abundances of metagenome-assembled genomes (MAGs) and their

closest genera are shown. B) Schematic representation of the known aniline degradation

pathway. AtdA1-5 catalyze the initial oxidation of aniline, and the intermediate catechol

is degraded via the ortho- or meta-cleavage pathway. C) The putative aniline oxidation

gene cluster identified in MAG01. Annotation of the six gene products and their

sequence or functional similarity to known aniline oxidation enzymes are shown. D)

Distribution of catechol-degradation genes among the eight most abundant MAGs in the

community. Presence/absence of KEGG entries associated with the catechol ortho- and

meta-cleavage pathway is indicated.

**Characterization of synthetic community members**

To construct a tractable synthetic community reflecting the diversity, functionality, and

ecological characteristics of the model aniline-degrading community, nine bacterial

strains representing a broad range of phylogenetic groups present in the original

community were isolated and selected as members of the synthetic community (Table

S2). Species-level taxonomic comparisons indicated that Cab074, Ach031, Pse033,

Sph075, Pse056, and Mic039 corresponded most closely to the six dominant MAGs (MAG01–MAG06). In contrast, Sph034, Bos046, and Bac050 were considered minor community members without corresponding MAGs. To facilitate detection of the primary aniline degrader, a GFP-labeled strain of Cab074 (Cab074F) was constructed and used in place of the wildtype in subsequent experiments. Degradation assays for individual synthetic community members confirmed that only Cab074F degraded aniline (Fig. S2), consistent with the metagenome characterization of the original community. Overall, the synthetic community recapitulates key characteristics of the original community: a single *Caballeronia* genotype predominantly performs initial aniline oxidation and other strains would participate in downstream metabolism.

**Composition–function landscape**

Using the synthetic community, we investigated how community-level functions emerge from interactions between the aniline degrader and non-degrader strains, using a microplate-based assay (Fig. 2A). To increase throughput, we employed a fluorescence-based assay of aniline-utilization activity using resazurin. Preliminary tests showed a positive correlation between RFU and aniline degradation, as evaluated using HPLC (Pearson's $r = 0.77$) (Fig. S3). Although the correlation was moderate, deviations likely arose because resazurin reduction reflects not only aniline degradation but also the metabolism of downstream intermediates. Thus, the resazurin-based assay would capture overall aniline-utilization activity, encompassing both aniline degradation and downstream metabolic processes. Simultaneously, Cab074F abundance was monitored by using GFP fluorescence, which showed strong linear correlation between GFP RFU and MA074F abundance as determined via colony counting (Pearson's $r = 0.89$) (Fig. S3).

By quantifying the aniline-utilization activity of all possible sub-community combinations containing Cab074F (i.e., $2^\square = 256$ patterns), we generated a composition–function landscape in which each data point represents a sub-community combination and each line denotes the functional effect of a single non-degrader (Fig. 2B). In this study, "functional effect" was defined as the change in aniline-utilization activity between two sub-community combinations differing only in the presence or

absence of the focal strain. If the functional effect of each strain were constant

regardless of co-existing members, the line angles would remain consistent. Indeed,

Ach031 and Pse033 consistently exhibited negative and positive functional effects,

respectively, with few exceptions (Fig. 2C). Sph034 and Mic039 also showed relatively

consistent effects, whereas the remaining strains exhibited highly variable effects,

ranging from strongly positive to strongly negative depending on the composition of the

surrounding community.

To systematically quantify the contribution of each species to community-level

function, we applied Shapley values to the composition–function landscape, calculating

the average functional effect of each strain across all possible assembly pathways from

the Cab074F-only community to the nine-member community ($8! = 40,320$ pathways).

As the results, Pse033 contributed most positively, followed by Bac050 and Sph075,

whereas Ach031 and Sph034 contributed negatively (Fig. 2D).

We evaluated how the presence or absence of a given non-degrader strain

altered the functional effect of another. For example, Sph034 had a near-neutral effect in

the absence of Pse033, but became negative when Pse033 was present (Fig. 2E). We

quantified such "functional interaction" as changes in the mean functional effect of a

focal strain due to a partner strain's presence or absence. We identified ten significant

negative and three positive functional interactions ($t$-test, $P < 0.05$), indicating

combinations that lead to either diminished or enhanced community-level functions,

respectively, compared with simple additive assumptions (Fig. 2F). Despite its high

individual contribution to aniline degradation, Pse033 exhibited negative interactions

with six other strains, suggesting diminished contributions in co-culture. In contrast,

Mic039 showed no significant interactions, indicating that its functional effect was

largely independent of other species.

Fig. 2 Full factorial evaluation of the synthetic aniline-degrading community. A)
Schematic overview of the experimental workflow. Sub-community combinations
consisting of the primary aniline degrader (Cab074F) and all possible combinations of
eight non-degrader strains were cultivated for 72 h in medium containing aniline as the
sole carbon source. Aniline-utilization activity and Cab074F growth were quantified as
the area under the fluorescence-time curve of resorufin and GFP, respectively. B)
Composition–function landscape of aniline-utilization activity for all 256 combinations.
Each data point represents aniline-utilization activity of a sub-community combination
at a given species richness. Line angles reflect the functional effect of each strain on
aniline-utilization activity, and colors indicate the identity of the strain present in the
right-side node but absent in the left-hand node (color-coded as in panel C). C)
Strain-resolved composition–function landscapes for eight non-degrader strains. D)
Overall contributions of the eight non-degrader strains to community-level function,
quantified as Shapley values. E) Example of a functional interaction: Sph034 exhibited
a near-neutral effect in the absence of Pse033 but a negative effect when Pse033 was
present. The change in the mean functional effect was defined as the functional
interaction between two species. F) Pairwise functional interactions among the eight

non-degrader strains. Positive values indicate the synergistic contributions to

community-level functions; negative values indicate sub-additive contributions.

Statistically significant interactions (t-test, $P < 0.05$) are indicated.

**Abundance of primary aniline degrader**

Given that Cab074F was the sole strain capable of converting aniline to catechol, we

hypothesized that variations in its population success could explain differences in

community-level aniline-utilization activity. To test this hypothesis, we generated a

composition–abundance landscape for Cab074F using GFP fluorescence data (Fig. S4)

and assessed the correlations between the functional effect on aniline-utilization activity

and the corresponding effects on Cab074F growth within the same community contexts

(Fig. 3). The results varied significantly across non-degrader strains: Ach031, Pse033,

Sph034, and Mic039 showed significant positive correlations, whereas the others

showed weak or no correlations. These positive correlations suggest that the influence

of these non-degrader strains on community-level aniline-utilization activity can be

largely explained by their facilitation or inhibition of Cab074F growth. In contrast, the

absence of correlation implies the involvement of additional mechanisms, such as

effects on downstream metabolism or interactions with coexisting non-degraders. It was

found that Ach031 and Pse033, which exhibited consistent functional effects regardless

of community composition (Fig. 2C), also showed exceptionally strong positive

correlations. In contrast, strains that exhibited highly variable functional effects (Bos046,

Bac050, Pse056, and Sph075) showed weak or no correlations. These findings suggest

that the functional effects mediated through Cab074F growth tend to be robust to

community composition, whereas effects involving other mechanisms are more

sensitive to the community context.

**Fig. 3** Correlation between changes in Cab074F's growth and aniline-utilization activity.

Each point represents differences in Cab074F growth (measured using GFP) and

aniline-utilization activity (measured using resazurin) between sub-community

combinations differing only by the presence or absence of a focal non-degrader strain

(corresponding to a line in Fig. 2C). A positive correlation suggests that each species'

functional effect is related to promoting or inhibiting the growth of the primary aniline

degrader, Cab074F, whereas the absence of a correlation implies that other mechanisms

predominantly contribute. Asterisks denote significance levels in Pearson's correlation

analysis: *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$.

**Bottom-up predictability of the composition-abundance landscape**

Using the obtained dataset, we tested whether the functions of complex microbial

communities can be predicted based on observations of simpler sub-community

combinations. To assess how species richness in the training data influences predictive

performance, we trained random forest models using sub-communities of defined

richness levels and applied them to predict aniline utilization activity in communities

containing 5–9 species. Models trained solely on three-member combinations (n = 28)

achieved high predictive accuracy for more complex communities ($r = 0.78$) (Fig. 4A).

Incorporating more diverse combinations into the training set did not substantially

improve the prediction performance (Fig. 4B). However, models trained solely on

two-member combinations (i.e., Cab074F plus one other strain, n = 8) showed markedly

lower accuracy ($r = 0.49$). These results indicate that sub-community combinations of

just three to four species can support robust bottom-up prediction of microbial

community functions.

Because exhaustively sampling all sub-community combinations is infeasible

in real microbial communities, we tested performance with reduced training data (50%

and 25%). Reducing the dataset caused only modest decreases in predictive

performance. For example, models trained on 25% of the three-member combinations

(n = 7) exhibited lower performance ($r = 0.51$), whereas those trained on 25% of the

four-member combinations (n = 14) still achieved high performance ($r = 0.71$) (Fig. 4B).

Across conditions, prediction performance was tightly associated with the number of

unique sub-community combinations used for training (Fig. 4C), explaining the

apparent dip in performance at species richness of 7, where relatively few combinations

are possible (n = 28). Moreover, when comparing models at similar training set sizes,

those trained on 3–4-member combinations tended to outperform models trained on

more diverse (6–7-member) combinations. Overall, random forest models trained with

simple sub-community combinations can reliably predict functions of diverse

communities without exhaustive data, provided that the training set included the

sufficient number of unique combinations.

**Fig. 4.** Bottom-up prediction of aniline-utilization activity using random forest models.

A) Schematic of the bottom-up prediction framework. To understand which levels of

sub-community diversity (species richness) are informative for predicting

community-level functions, random forest models were trained on observed

sub-community combinations at fixed richness levels (example shown for species

richness = 3). Model performance in predicting aniline-utilization activity of

5–9-member combinations was evaluated as Pearson's correlation coefficient. B)

Prediction performance of random forest models trained on sub-community

combinations of varying species richness. To assess model performance under limited

training data, models trained on reduced datasets (50% or 25% random sampling) were

also evaluated. Error bars indicate the standard deviations across 100 runs. C)

Relationship between prediction performance and the number of sub-community

combinations used for model training. Each data point represents a model trained at the

indicated species richness and saturation level.

**Interpretation of the random forest model to identify key species and interactions**

We tested whether interpretating the trained random forest models could reveal key

species and interspecies interactions that shape community-level functions. To this end,

we evaluated two model-derived metrics: permutation importance, which quantifies

each feature's contribution by measuring changes in prediction accuracy after random

shuffling of focal features [40] and Friedman's H statistic, which estimates the strength

of interactions between features by assessing how the effect of one feature on aniline

utilization activity depends on another [41].

We applied these interpretability techniques to the random forest model trained

on three-, four-, and five-member sub-community combinations (Fig. 5A–C). The

permutation importance scores consistently identified Ach031 and Pse033 as the most

influential species, followed by Sph034, Bac050, and Sph075. This ranking closely

matched species contributions estimated from the full dataset using Shapley values (Fig.

2D), indicating that random forest models trained on simple sub-community data can

reliably identify the dominant contributors to community-level functions.

Additionally, Friedman's H statistics derived from models trained on three- and

four-member sub-communities correlated significantly with empirical functional

interaction strengths (Fig. 2E) ($r = 0.40$–$0.45$, $P < 0.05$), whereas those trained on

two-member sub-communities showed no correlation ($r = 0.14$, $P = 0.457$). The three-

and four-member models highlighted different sets of pairwise relationships, and

predictive accuracy improved markedly when models were trained on combined dataset of all two- to four-member combinations ($r = 0.71$, $P < 0.0001$) (Fig. 5D). In contrast, Friedman's H statistics derived from models trained on five-member sub-communities were no longer significantly correlated with the observed functional interactions ($r = 0.20$, $P > 0.05$), and a similar loss of correspondence occurred in six-member models ($r = 0.14$, $P > 0.05$). These results suggest that increasing the species richness of training data reduces the ability to resolve specific interspecies interactions, likely due to statistical difficulties in pinpointing influential feature pairs within high-dimensional observational units.

Together, these findings show that machine-learning models trained on simple sub-community combinations can identify key species and interspecies interactions underlying emergent community functions. In particular, sub-community combinations of three to four species appear to represent the most informative unit for dissecting the emergent properties of microbial community functions.

**Fig. 5.** Key species and interspecies interactions inferred from the random forest models. Permutation importance and Friedman's H statistic were calculated from random forest models trained on sub-community combinations comprising three, four, and five members, as well as from models trained on two- to four-member combinations. Permutation importance was used to quantify the relative contributions of individual species to community-level aniline utilization activity, whereas Friedman's H statistic was used to assess the strength of pairwise functional interactions. To evaluate the

reliability of inferred interactions, Friedman's H values were compared with the

empirically derived functional interaction strengths (Fig. 2F). Permutation importance

values were consistent with empirically calculated Shapley values (Fig. 2D).

**DISCUSSIONS**

Using a synthetic microbial community capable of degrading aniline, this study showed

that analyzing simple sub-community combinations enables the identification of key

species and interactions shaping community-level functions (Fig. 5). Traditional

analytical approaches to microbial community analysis have tended to follow either

holistic (meta-omics-based) or reductionist (isolated strain/single-cell-based) strategies.

In contrast, this study proposes an intermediate, sub-community-based framework that

effectively captures the emergent properties of microbial community functions. Our

proposed framework involves three steps: (1) generating multiple sub-community

combinations from a target microbial community, (2) evaluating their functions and

compositions, and (3) analyzing the resulting dataset to infer the species contributions

and interactions, thereby identifying rational targets for follow-up analyses or microbial

community engineering. Combining with meta-omics approaches (e.g.,

metatranscriptomics), this framework would further strengthen mechanistic inference.

Recent advances in high-throughput microbial cultivation platforms [42-43] have made

it increasingly feasible to generate and characterize thousands of sub-community

combinations without isolation. Thus, the findings and analytical framework presented

here provide a foundation for applying this approach to more complex natural microbial

communities in the near future.

534           Consistent with recent studies investigating composition–function landscapes

of synthetic microbial communities [44-45], our results demonstrate that the functional

effects of individual species are highly context-dependent (Fig. 2). To explain such

context dependency, a "global epistasis" has been proposed in which a species'

functional effect can be approximated by simple linear regression against the functional

level of the background community—tending to be positive when the baseline function

is low, and negative when it is high [7]. This pattern was evident in our dataset (Fig. S5)

and appears to reflect both functional redundancy among species and complex

interaction networks that obscure individual contributions. Beyond this statistical

explanation, our results suggest that mechanistic differences in species' modes of action

also contribute: strains that affected aniline degradation through direct promotion of the

degrader Cab074F's growth showed relatively stable functional effects, whereas those

acting through other mechanisms exhibited greater context sensitivity (Fig. 3). Although

detailed interaction mechanisms are beyond the scope of this study, linking statistical

patterns in composition–function landscapes with mechanistic insights constitutes a

future avenue for effectively controlling microbial community functions.

Another key finding is that microbial community functions can be predicted in

a bottom-up manner by combining simple sub-community observations with machine

learning. Similar bottom-up approaches have recently gained traction in studies using

synthetic microbial communities, with several strategies showing promise. For example,

genome-scale metabolic models have been applied to predict the metabolic output of

complex microbial consortia [46-48]. In parallel, mathematical models that infer

community function from the geometry of composition–function landscapes have also

been developed [7,45,49], enabling prediction without detailed knowledge of metabolic

pathways. Although many of these modeling studies relied on full datasets in

composition–function landscapes and leave-one-out validation strategies, our

framework demonstrated that reasonable prediction can be achieved using only a

limited set of simple sub-community combinations. This flexibility makes our approach

suitable not only for the rational design of synthetic microbial consortia but also for

investigating natural communities, where exhaustive sub-community observation is

infeasible.

This study also introduced a method for identifying key species and

interspecies interactions by interpreting variable-importance measures derived from

machine-learning models. Whereas models trained on 3–4-member communities

yielded reasonable predictions, increasing species richness in observed sub-community

combinations did not necessarily improve analytical insights (Fig. 5), likely due to

statistical challenges in resolving interspecies interactions from more complex

combinations. These findings suggest that sub-community combinations consisting of

three to four species may represent an analytical sweet spot, balancing predictive power and interpretability. This observation aligns with previous reports where three- to four-member sub-communities provided robust models for predicting species abundance or survival [17-18], highlighting their promise as a unit for dissecting emergent properties in microbial communities.

In summary, this study demonstrates that simple sub-community combinations contain valuable information for bottom-up, mechanistic understanding of microbial community functions. The machine learning–based framework proved robust to data sparsity, making it a promising tool for analyzing diverse and noisy microbial systems. Although further validation is needed to assess its applicability to highly diverse and dynamic microbial ecosystems, integrating this sub-community-based strategy with existing meta-omics and cultivation-based methods offer a promising path towards rationally predicting and controlling emergent properties of microbial community functions.

**CONFLICTS OF INTEREST**

The authors declare no conflicts of interest.

**ACKNOWLEDGEMENTS**

This work was supported by JST, ACT-X (JPMJAX22B1), Research Grants by Kurita Water and Environment Foundation (22E053, 23K008), and Gene-grant 2023 by Nippon Genetics Co., Ltd. to Hidehiro Ishizawa. The manuscript's English was polished using generative AI for grammar and style and further edited by Enago (www.enago.jp). The authors reviewed and edited all output.

**AUTHOR CONTRIBUTIONS**

Hidehiro Ishizawa: Conceptualization, Formal analysis, Funding acquisition, Project administration, Software, Visualization, Writing – Original Draft. Sunao Noguchi: Formal analysis, Investigation, Visualization, Writing – Review & Editing. Miku Kito: Investigation, Writing – Review & Editing. Yui Nomura: Investigation, Writing – Review & Editing. Kodai Kimura: Investigation, Writing – Review & Editing. Masahiro

Takeo: Project Administration, Writing – Review & Editing, Supervision.

**References**

1. Bar-On YM, Phillips R, Milo R. The biomass distribution on Earth. *Proc Natl Acad Sci USA* 2018;**115**:6506–11. https://doi.org/10.1073/pnas.1711842115

2. Bardgett RD, Freeman C, Ostle NJ. Microbial contributions to climate change through carbon cycle feedbacks. *ISME J* 2008;**2**:805–14. https://doi.org/10.1038/ismej.2008.58

3. Nicholson JK, Holmes E, Kinross J et al. Host-gut microbiota metabolic interactions. *Science* 2012;**336**:1262–7. https://doi.org/10.1126/science.1223813

4. Hassani MA, Durán P, Hacquard S. Microbial interactions within the plant holobiont. *Microbiome* 2018;**6**:58. https://doi.org/10.1186/s40168-018-0445-0

5. McCarty NS, Ledesma-Amaro R. Synthetic biology tools to engineer microbial communities for biotechnology. *Trends Biotechnol* 2019;**37**:181–97. https://doi.org/10.1016/j.tibtech.2018.11.002

6. Compant S, Cassan F, Kostić T et al. Harnessing the plant microbiome for sustainable crop production. *Nat Rev Microbiol* 2025;**23**:9–23. https://doi.org/10.1038/s41579-024-01079-1

7. Diaz-Colunga J, Skwara A, Vila JCC et al. Global epistasis and the emergence of function in microbial consortia. *Cell* 2024;**187**:3108–19. https://doi.org/10.1016/j.cell.2024.04.016

8. Hu B, Wang M, Geng S et al. Metabolic exchange with non-alkane-consuming *Pseudomonas stutzeri* SLG510A3-8 improves n-alkane biodegradation by the alkane degrader *Dietzia* sp. Strain DQ12-45-1b. *Appl Environ Microbiol* 2020;**86**:e02931-19. https://doi.org/10.1128/AEM.02931-19

9. D'Souza G, Schwartzman J, Keegstra J et al. Interspecies interactions determine growth dynamics of biopolymer-degrading populations in microbial communities. *Proc Natl Acad Sci USA* 2023;**120**:e2305198120. https://doi.org/10.1073/pnas.2305198120

10. Peng Q, Zhao C, Wang X et al. Modeling bacterial interactions uncovers the importance of outliers in the coastal lignin-degrading consortium. *Nat Commun*

2025;**16**:639. https://doi.org/10.1038/s41467-025-56012-8

11. Noecker C, Chiu HC, McNally CP et al. Defining and evaluating microbial contributions to metabolite variation in microbiome-metabolome association studies. *mSystems* 2019;**4**:e00579-19. https://doi.org/10.1128/msystems.00579-19

12. Pinto S, Benincà E, Nes EH et al. Species abundance correlations carry limited information about microbial network interactions. *PLoS Comput Biol* 2022;**18**:e1010491. https://doi.org/10.1371/journal.pcbi.1010491

13. Holt RD. Predation, apparent competition, and the structure of prey communities. *Theor Popul Biol* 1977;**12**:197–229. https://doi.org/10.1016/0040-5809(77)90042-9

14. Kondoh M. Building trophic modules into a persistent food web. *Proc Natl Acad Sci USA* 2008;**105**:16631–5. https://doi.org/10.1073/pnas.0805870105

15. Stouffer DB, Bascompte J. Understanding food□web persistence from local to global scales. *Ecol Lett* 2010;**13**:154–61. https://doi.org/10.1111/j.1461-0248.2009.01407.x

16. Mayfield MM, Stouffer DB. Higher-order interactions capture unexplained complexity in diverse communities. *Nat Ecol Evol* 2017;**1**:0062. https://doi.org/10.1038/s41559-016-0062

17. Friedman J, Higgins LM, Gore J. Community structure follows simple assembly rules in microbial microcosms. *Nat Ecol Evol* 2017;**1**:0109. https://doi.org/10.1038/s41559-017-0109

18. Ishizawa H, Tashiro Y, Inoue D et al. Learning beyond-pairwise interactions enables the bottom–up prediction of microbial community structure. *Proc Natl Acad Sci USA* 2024;**121**:e2312396121. https://doi.org/10.1073/pnas.2312396121

19. Thompson J, Johansen R, Dunbar J et al. Machine learning to predict microbial community functions: An analysis of dissolved organic carbon from litter decomposition. *PLoS One* 2019;**14**:e0215502. https://doi.org/10.1371/journal.pone.0215502

20. Dutta A, Goldman T, Keating J et al. Machine learning predictions biogeochemistry from microbial community structure in a complex model system. *Microbiol Spectr* 2022;**10**:e01909-21. https://doi.org/10.1128/spectrum.01909-21

21. Kulkarni M, Chaudhari A. Microbial remediation of nitro-aromatic compounds: An

overview. *J Environ Manag* 2007;**85**:496–512.

https://doi.org/10.1016/j.jenvman.2007.06.009

22. Coleman NV, Mattes TE, Gossett JM et al. Biodegradation of cis-dichloroethene as

the sole carbon source by a β-Proteobacterium. *Appl Environ Microbiol*

2002;**68**:2726–30. https://doi.org/10.1128/AEM.68.6.2726-2730.2002

23. Nurk S, Meleshko D, Korobeynikov A et al. metaSPAdes: a new versatile

metagenomic assembler. *Genome Res* 2017;**27**:824–34.

https://doi.org/10.1101/gr.213959.116

24. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* 2009;**25**:1754–60.

https://doi.org/10.1093/bioinformatics/btp324

25. Kang DD, Li F, Kirton E et al. MetaBAT 2: an adaptive binning algorithm for

robust and efficient genome reconstruction from metagenome assemblies. *PeerJ*

2019;**7**:e7359. https://doi.org/10.7717/peerj.7359

26. Parks DH, Imelfort M, Skennerton CT et al. CheckM: assessing the quality of

microbial genomes recovered from isolates, single cells, and metagenomes.

*Genome Res* 2015;**25**:1043–1055. https://doi.org/10.1101/gr.186072.114

27. Tanizawa Y, Fujisawa T, Nakamura Y. DFAST: a flexible prokaryotic genome

annotation pipeline for faster genome publication. *Bioinformatics* 2018;**34**:1037–9.

https://doi.org/10.1093/bioinformatics/btx713

28. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools

for functional characterization of genome and metagenome sequences. *J Mol Biol*

2016;**428**:726–31. https://doi.org/10.1016/j.jmb.2015.11.006

29. van den Belt M, Gilchrist C, Booth TJ et al. CAGECAT: the comparative gene

cluster analysis toolbox for rapid search and visualization of homologous gene

clusters. *BMC Bioinform* 2023;**24**:181. https://doi.org/10.1186/s12859-023-05311-2

30. Draghi WO, Peeters C, Cnockaert M et al. *Burkholderia cordobensis* sp. nov., from

agricultural soils. *Int J Syst Evol Microbiol* 2014;**64**:2003–8.

https://doi.org/10.1099/ijs.0.059667-0

31. Amann RI, Ludwig W, Schleifer KH. Phylogenetic identification and in situ

detection of individual microbial cells without cultivation. *Microbiol Rev*

1995;**59**:143–69. https://doi.org/10.1128/mr.59.1.143-169.1995.

32. Schlechter RO, Jun H, Bernach M et al. Chromatic bacteria – a broad host-range

plasmid and chromosomal insertion toolbox for fluorescent protein expression in

bacteria. *Front Microbiol* 2018;**9**:3052. https://doi.org/10.3389/fmicb.2018.03052

33. Sarker SD, Nahar L, Kumarasamy Y. Microtitre plate-based antibacterial assay

incorporating resazurin as an indicator of cell growth, and its application in the in

vitro antibacterial screening of phytochemicals. *Methods* 2007;**42**:321–4.

https://doi.org/10.1016/j.ymeth.2007.01.006

34. Shapley L. A value for n-person games. In: Kuhn H, Tucker A (ed.), Contributions

to the Theory of Games II. Princeton: Princeton University Press, 1953, 307–317.

https://doi.org/10.1515/9781400881970-018

35. Inglis A, Parnell A, Hurley CB. Visualizing variable importance and variable

interaction effects in machine learning models. *J Comput Graph Stat*

2022;**31**:766–78. https://doi.org/10.1080/10618600.2021.2007935

36. Ishizawa H. Data and analytical codes for: Ishizawa et al. (2025) Decoding

emergent properties of microbial community functions through sub-community

observations and interpretable machine learning. Zenodo. 2025.

https://doi.org/10.5281/zenodo.17364288

37. Takeo M, Fujii T, Maeda Y. Sequence analysis of the genes encoding a

multicomponent dioxygenase involved in oxidation of aniline and o-toluidine in

*Acinetobacter* sp. strain YAA. *J Ferment Bioeng* 1998;**85**:17–24.

https://doi.org/10.1016/S0922-338X(97)80347-9

38. Takeo M, Ohara A, Sakae S et al. Function of a glutamine synthetase-like protein in

bacterial aniline oxidation via γ-glutamylanilide. *J Bacteriol* 2013;**195**:4406–14.

https://doi.org/10.1128/JB.00397-13

39. Barnsley EA. Role and regulation of the ortho and meta pathways of catechol

metabolism in pseudomonads metabolizing naphthalene and salicylate. *J Bacteriol*

1976;**125**:404–8. https://doi.org/10.1128/jb.125.2.404-408.1976

40. Altmann A, Toloşi L, Sander O. Permutation importance: a corrected feature

importance measure. *Bioinformatics* 2010;**26**:1340–7.

https://doi.org/10.1093/bioinformatics/btq134

41. Friedman JH, Popescu BE. Predictive learning via rule ensembles. *Ann Appl Stat*
2008;**2**:916–54. https://doi.org/10.1214/07-AOAS148

42. Hsu RH, Clark RL, Tan JW et al. Microbial interaction network inference in
microfluidic droplets. *Cell Syst* 2019;**9**:229–42.
https://doi.org/10.1016/j.cels.2019.06.008

43. Jackman CM, Deans KW, Forney LJ et al. Microdroplet co-cultivation and
interaction characterization of human vaginal bacteria. *Integr Biol* 2019;**11**:69–78.
https://doi.org/10.1093/intbio/zyz006

44. Sanchez-Gorostiaga A, Bajić D, Osborne ML et al. High-order interactions distort
the functional landscape of microbial consortia. *PLoS Biol* 2019;**17**:e3000550.
https://doi.org/10.1371/journal.pbio.3000550

45. Ruiz J, De Celis M, Diaz□Colunga J et al. Predictability of the
community□function landscape in wine yeast ecosystems. *Mol Syst Biol*
2023;**19**:e11613. https://doi.org/10.15252/msb.202311613

46. Clark RL, Connors BM, Stevenson DM et al. Design of synthetic human gut
microbiome assembly and butyrate production. *Nat Commun* 2021;**12**:3254.
https://doi.org/10.1038/s41467-021-22938-y

47. Gowda K, Ping D, Mani M et al. Genomic structure predicts metabolite dynamics
in microbial communities. *Cell* 2022;**185**:530–46.
https://doi.org/10.1016/j.cell.2021.12.036

48. Ruan Z, Chen K, Cao W et al. Engineering natural microbiomes toward enhanced
bioremediation by microbiome modeling. *Nat Commun* 2024;**15**:4694.
https://doi.org/10.1038/s41467-024-49098-z

49. Skwara A, Gowda K, Yousef M et al. Statistically learning the functional landscape
of microbial communities. *Nat Ecol Evol* 2023;**7**:1823–33.
https://doi.org/10.1038/s41559-023-02197-4