Correspondence to

Pascal Baelo

Centre de Surveillance de la Biodiversité, University of Kisangani, Kisangani City,

2012, Democratic Republic of Congo

Léa Fourchault

Vertebrate group, Directorate Taxonomy and Phylogeny, Royal Belgian Institute of

Natural Sciences, 1000, Brussels, Belgium

Running head: Distribution of African rope squirrels

REASSESSING THE DIVERSITY AND DISTRIBUTION OF AFRICAN ROPE

SQUIRRELS (*FUNISCIURUS* TROUESSART, 1880)

Pascal Baelo[1, *], Léa Fourchault[2, 3] *, Nicolas Laurent[2, 4], Claude Mande[5], Anne

Laudisoit[7], Nicaise Amundala[5], Ganiyat Temidayo Saliu[2], Guy-Crispin Gembu[1, 5],

Joachim Mariën[6, 7], Sophie Gryseels[2, 7], Jan Hulselmans[7] and Erik Verheyen[2, 7]

Full e-mail addresses, contributions and author statements and data availability see end

of paper.

ORCID

PB, https://orcid.org/0000-0003-3134-9881

LF, https://orcid.org/0000-0003-4026-7319

CM, https://orcid.org/0000-0002-9851-3677

AL, https://orcid.org/0000-0001-7626-9426

NA,

NL, https://orcid.org/0000-0001-9347-6507

GTS, https://orcid.org/0009-0003-5553-9070

JN,

JM, https://orcid.org/0000-0001-7023-445X

SG, https://orcid.org/0000-0002-4851-9384

GG,

JH,

EV, https://orcid.org/0000-0001-7157-1474

Affiliations

¹Centre de Surveillance de la Biodiversité, University of Kisangani, Kisangani City,

Democratic Republic of Congo

²Vertebrate group, Directorate Taxonomy and Phylogeny, Royal Belgian Institute of

Natural Sciences, Brussels, Belgium

³Systems Ecology and Resource Management, Université Libre de Bruxelles, Brussels,

Belgium

⁴Department of Environmental Sciences, Open Universiteit, Heerlen, The Netherlands

⁵Faculty of Science, University of Kisangani, Kisangani, Kisangani City, Democratic

Republic of the Congo

⁶Department of Biomedical Sciences - Virus Ecology, Institute of Tropical Medicine,

Antwerp, Belgium

⁷Evolutionary Ecology Group, Department of Biology, University of Antwerp, Antwerp,

Belgium

Teaser text

We studied the diversity and distribution of rope squirrels, genus *Funisciurus* Trouessart

1880, in the Congo Basin. By gathering the most comprehensive genetic and

morphometric data to date on a significant part of the geographical and taxonomic range

of these squirrels (specimens from seven provinces of the Democratic Republic of

Congo), we identified six main taxa: one previously undescribed species, three allopatric

taxa, and two genetically distinct sympatric taxa. We then mapped their distribution,

showing that the Congo and Kasai rivers represent significant biogeographical barriers

limiting the distribution of *Funisciurus* species. These results support research into the

role of rope squirrel species in the transmission of pathogens of public health

importance.

Image for the online table of contents: African rope squirrel

Photo credit: Bushnell photo trap set by Pascal Baelo

ABSTRACT

The diversity, taxonomy, distribution and ecology of sub-Saharan tree squirrels remain

under-researched. This study aims to elucidate the diversity and distribution of rope

squirrels, genus *Funisciurus* Trouessart 1880, in the Congo Basin. We assembled the

most comprehensive genetic and morphometric data set to date, on a significant portion

of the geographical and taxonomic range of these squirrels (470 specimens from seven

provinces of the Democratic Republic of Congo). We identified six main taxa: a

previously undescribed *Funisciurus* species (here, Fx), three allopatric taxa of

Funisciurus anerythrus (Fc, FaRB and FaLB)*,* and two sympatric genetically distinct

taxa (FbLB1, FbLB2), potentially belonging to a *Funisciurus bayonii* species complex.

We then mapped their distribution, showing that the Congo and Kasaï Rivers represent

significant biogeographical barriers limiting the distribution of *Funisciurus* species. In

addition to providing a baseline for research on the impact of climate change and habitat

degradation on the abundance and distribution of rope squirrels, our findings support

research on the role of rope squirrel species in the transmission of pathogens of public

health importance.

Keywords: Biodiversity, Congo Basin, DNA-barcoding, Sciuridae, rope squirrels

INTRODUCTION

The Sciuridae family, which includes approximately 50 genera and 302 species (**Zachos**

et al., 2021, Steiner & Huettmann, 2023), is distributed globally, with the exception of

Antarctica. While some squirrel species are valued as a food source in certain regions

(**Fa et al., 2009; Baelo et al., 2018**), they are increasingly recognized as important

reservoirs and incidental hosts for zoonotic pathogens (**Han et al., 2015; e.g.,**

Sklenovska et al., 2018; Schulze et al., 2020). However, the phylogenetic, ecological,

and life history characteristics of many squirrels, particularly those in sub-Saharan

Africa, remain poorly understood. In Central Africa, the Sciuridae family is represented

by eight genera, including the African pygmy squirrel (*Myosciurus pumilio*) and several

species within the rope (*Funisciurus*), sun (*Heliosciurus*), ground (*Xerus*), bush

(*Paraxerus*), palm (*Epixerus*), and giant (*Protoxerus*) genera (**Kingdon & Hoffmann,**

2013; Thorington et al., 2012). Four of these genera—*Funisciurus*, *Paraxerus*,

Heliosciurus, and *Protoxerus*—are common across diverse habitats in the Democratic

Republic of Congo (DRC), spanning dense rainforest to savannah ecosystems.

Unlike many large mammals that are highly susceptible to habitat disturbances, a

few Sciuridae species have shown a remarkable capacity to adapt to human-altered

environments. For instance, *Funisciurus anerythrus* has been observed at high

population densities in palm oil plantations, reaching up to 500 individuals per square

kilometer—far exceeding densities in primary lowland rainforests (**Tiee et al., 2018**).

This adaptability heightens the frequency of human-squirrel interactions and raises

concerns about potential zoonotic spillover risks, particularly in agricultural and peri-

domestic settings (**Rimoin et al., 2017; Kumakamba et al., 2021**).

Recent studies underscore the role of squirrels in zoonotic disease transmission,

for example bornavirus (**Hoffmann et al., 2015; Schulze et al., 2020**). Sub-Saharan

squirrels, particularly those within the *Funisciurus* and *Heliosciurus* genera, have also

been associated with the mpox virus (MPXV), of which a variant led to a global

outbreak infecting over 80,000 people in 2022, emphasizing the importance of

understanding the status of these squirrels as zoonotic reservoirs (**Chen et al., 2012;**

Curaudeau et al., 2023; Kinganda-Lusamaki et al., 2025).

Studying the distribution and phylogenetic relationships of *Funisciurus* in sub-

Saharan Africa could illuminate patterns of pathogen-host co-evolution within this

genus. MPXV, for instance, comprises two main clades: the more virulent clade I and

the less virulent clade II (**Nakazawa et al., 2015; Kinganda-Lusamaki et al., 2025**).

Clade I MPXV occurs in the Congo Basin, and clade II in West Africa - though both

clades have been exported via human infections to other regions in Africa and the rest of

the world (**Otieno et al., 2025**). The distinct clades found to be infecting humans could

be a reflection of genetically and geographically distinct natural reservoir(s). These

geographically distinct clades may correlate with specific squirrel populations,

suggesting possible co-evolutionary and biogeographical relationships affecting disease

transmission dynamics.

Geographical barriers, particularly rivers, can influence the distribution of rodents

and the spread of pathogens, limiting the ranges of arboreal mammals in West and

Central Africa. While natural barriers such as the Volta, Sanaga and Congo rivers are

known to limit the ranges of several small mammal species (**Kennis et al., 2011**), the

influence of the Congo River on the distribution and evolutionary history of

Sciuromorph rodents remains underexplored, leaving significant gaps in our

understanding of their evolutionary history across the Congo Basin.

This study addresses these gaps by clarifying the diversity of *Funisciurus* taxa within the

Congo Basin and assessing whether the Congo River serves as a significant geographic

barrier affecting their distribution. Along with advancing biodiversity, evolutionary and

biogeographic insights, our research provides critical data on the distribution of

Funisciurus taxa across the Congo Basin, to serve as a baseline for comparison with the

phylogeographies of zoonotic pathogens relevant for public health.

2. MATERIALS AND METHODS

2.1. SAMPLE COLLECTION

2.1.1. SQUIRREL CAPTURE AND SAMPLING; we trapped and sampled squirrels

from 2014 to 2022 in 17 localities across four provinces (Tshopo, Bas-Uele, Ituri and

Tshuapa) in the Democratic Republic of Congo (DRC). Further, we used specimens

from three other localities in three provinces (Kwilu, Kinshasa and Kongo Central),

where squirrels were sampled in 1995 and preserved at the Royal Museum for Central

Africa-Tervuren (Belgium) (**Supplementary Figure S1**).

We captured squirrels using traditional traps (**Supplementary Figure S2**), baited

with a ripe palm nut and placed in different habitats (primary forests, secondary forests

and fallow land), at heights ranging from 0.00 m (on the ground) to 7.37 m (tree

branches) (average height =1.58m±1.29m). The various standard traps and techniques

used to capture rodents (pitfall traps, Havahart traps, flap traps, Sherman traps) proved

unsuitable or ineffective when applied to squirrels. Consequently, traditional traps were

used, given their effectiveness in capturing tree squirrels.

The traditional traps were constructed from stems, flexible vines and rattan, which

were wrapped around a branch to facilitate the animals' movement towards the bait

(**Supplementary Figure S2**). A wooden wedge was then attached to the wire to act as a

trigger for the bait. The trap was baited to attract the animal and catch it by the neck.

Traditional traps had one or two sliding knots, allowing squirrels to be caught in both

directions of approach. Other traps were constructed from flexible rods and wires with a

radius of 0.5 mm and rattan. These traps had a single noose and functioned in the same

way as traps with two nooses.

We euthanized live captured squirrels with an overdose of isoflurane, adhering to

the guidelines of the American Society of Mammalogists for euthanasia (**Sikes et al.,**

2016), before collecting blood samples on Whatman filter paper and collecting tissue

samples (tongue, kidney, liver, lung, spleen, and colon) and feces. These tissues were

preserved either in 70-94% ethanol or in a DNA/RNA preservation solution (e.g.

RNAlater or DNA/RNA Shield, Zymo Research). The preserved samples were sent to

the Royal Belgian Institute of Natural Sciences (RBINS) and the University of Antwerp

(Belgium) for genetic analysis. The carcasses were preserved in a 10% formalin solution

and stored partly at the Centre de Surveillance de la Biodiversité in Kisangani (DRC)

and partly at the University of Antwerp (Belgium). A list of all specimens is provided in

Supplementary Table S1.

Field research permits were granted by the Centre de Surveillance de la

Biodiversité in Kisangani of University of Kisangani at the University of Kisangani,

with provincial and   local authorities approving all mission orders. Samples exported

to the Royal Belgian Institute of Natural Sciences and the University of Antwerp

(Belgium) prior to the implementation of the Nagoya Protocol were covered by Centre

de Surveillance de la Biodiversité material transfer agreements, while those exported

after 2018 were authorized by a Nagoya export permit issued by the Congolese Ministry

of the Environment and Sustainable Development (MEDD) (Permit numbers

003/ANCCB-RDC/SG-EDD/BTB/08/2019, 018/ANCCB-RDC/SG-EDD/BTB/07/2022,

021/SG-EDD/BTB/ANCCB-RDC/08/2023, 011/ANCCB-RDC/SG-EDD/BTB/2023).

No CITES permits were required. Permission to import the samples into Belgium was

granted by the Federal Agency for the Safety of the Food Chain (FASFC), permit

numbers 2022DBP504, 2023DBP370, 2023DBP371, 2024DBP424.

2.1.2. EXTERNAL MORPHOMETRY AND CRANIOMETRY; Squirrels were first

identified *in situ* to genus or species level using **Kingdon & Hoffmann (2013)**. We

examined captured squirrels on the basis of external morphological criteria, *i.e.* coat,

body length, tail length, hind foot length, and ear length. Sex and reproductive stages

were determined by external observation, including the presence or absence of nipples

(small or swollen) and the appearance of the vagina (closed or open) in females or the

presence of testes (abdominal or scrotal) in males. Length was measured to the nearest

millimeter on fresh specimens using a Mitutoyo caliper to an accuracy of 0.05 mm.

Weight was measured using a spring balance (Pesola) and expressed in grams.

Only the skulls of adult squirrels were extracted and measured (**Supplementary**

Table S1). A total of 23 measurements were taken on each skull (**Verheyen & Bracke,**

1966; Van Der Straeten & Dieterlen, 1987; Verheyen et al., 1996). The used

measurements are listed with its corresponding number, abbreviation, and

morphological character (**Supplementary Table S2).**

We also measured 29 skulls from the collections of the Royal Museum for Central

Africa-Tervuren to link a craniometrically defined Operational Taxonomic Unit (OTU)

to cytochrome b sequences previously published in GenBank (**Falendysz at al., 2017**).

2.1.3. ANALYSIS OF MORPHOMETRIC AND CRANIOMETRIC DATA; The

approach consisted of several steps. First, OTUs were designated based on preliminary

Molecular Taxonomic Units (MOTUs) identified using cytochrome b data. Next, we

tested the hypothesis that variation in craniometrics and external morphological data

aligns with the MOTUs through multivariate analysis using R version 4.2.3 and

Statistica version 12 (StatSoft Inc).

To evaluate sexual dimorphism within OTUs, all measurements were analyzed.

However, as no significant sexual dimorphism was detected (**Supplementary Tables S3**

and S4), subsequent analyses were conducted using specimens of both sexes. Basic

statistics, including the number of observations, mean, minimum, maximum, standard

deviation, and coefficient of variation, were calculated for all OTUs (**Supplementary**

Tables S5 and S6)

A forward discriminant analysis was then performed using all 23 variables from

the complete craniometric dataset, clearly highlighting the sequenced specimens. A

second discriminant analysis focused on the OTUs *Funisciurus anerythrus* Left Bank

(FaLB), *Funisciurus anerythrus* Right Bank (FaRB), and *Funisciurus cf congicus* (Fc)

near Kikwit, using a partial craniometric dataset. For the Kikwit specimens, where some

measurements were unavailable due to skull damage, a mean substitution method was

applied to estimate missing values (Statistica) (**Supplementary Figure S4).** Finally, a

third discriminant analysis incorporated all available external measurements for all

OTUs, again clearly highlighting the sequenced specimens.

Not all specimens were intact, the tail and/or ear having been cut off and the skull

bones fractured, the skull not extracted or not recovered, which explains the variations in

the number of individuals (N) used in each type of analysis. The subset of tissues

bearing a DNA barcode was selected at random on the basis of the geographical origin

of the capture and the availability of morphometric measurements. We also measured 41

squirrel skulls collected in Kikwit, Kinshasa and Kongo Central (DRC) and stored at the

Royal Museum for Central Africa, and analyzed their tissue samples stored at the Royal

Belgian Institute of Natural Sciences.

2.2. MOLECULAR DATA ACQUISITION

We extracted DNA of squirrel tissue samples (tongue or liver or kidney or lung) using

the Nucleospin Tissue mini kit according to the manufacturer's instructions (Machery-

Nagel, Düren, Germany), with an overnight pre-lysis incubation instead of 1-3h, and a

one-minute elution at 70°C instead of RT. We then amplified a mitochondrial gene

fragment using universal cytochrome b primers L14723 and H15915 (**Montgelard et**

al., 2002) with Promega MasterMix PCR following the manufacturer's instructions

(Promega, Madison, USA). The initial denaturation was at 94°C for 3 min, followed by

35 cycles at 94°C for 1 min, 52°C for 1 min and 72°C for 1 min. Final extension was

carried out at 72°C for 7 minutes.

We then purified the amplified products using ThermoFisher Exo-SAP IT PCR

product clean-up reagent (ThermoFisher Scientific, Santa Clara, USA) and sent them to

Macrogen (Amsterdam, The Netherlands) for sequencing. Finally, we manually cleaned

the sequences based on the chromatograms and aligned them using the MAFFT

algorithm within Geneious 2019.2.3. This alignment included in-house generated

Funisciurus sequences, as well as two *Paraxerus* sequences serving as an outgroup. This

alignment also included *Funisciurus sp* sequences downloaded from GenBank, on the

condition that they did not contain stop-codons and clustered with our *Funisciurus*

sequences rather than with the outgroup (accession numbers: MF694315, MF694313,

MF694317, MF694320, MN597594 and MN597668), yielding a total of 143 sequences.

2.3. SPECIES DELIMITATION USING GENETIC DATA

We first inferred an approximately-maximum-likelihood phylogenetic tree using

MAFFT-aligned mitochondrial sequences and the FastTree plugin in Geneious 2019.2.3,

using default settings and rooting on the *Paraxerus* outgroup (**Price et al., 2009**).

Briefly, the FastTree plugin worked as follows: a rough tree topology was first obtained

through a neighbour-joining approach; tree length was then reduced using nearest-

neighbour interchanges and subtree-prune-regraft-moves. Tree topology and branch

length were improved using maximum-likelihood rearrangements, where each site was

assigned to one of 20 geometrically spaced categories (0.05-20) using a Bayesian

approach (gamma prior), representing each site's most likely evolutionary category.

Finally, to estimate the reliability of each split in the tree, the Shimodaira-Hasegawa test

was used on three alternate topologies (NNIs) around the focal split. The resampling size

was 1,000 (**Price et al., 2009**).

To further explore *Funisciurus* taxon delineation within our dataset, we used

species delimitation algorithms that generate species partitions based either on distance

(ASAP, Assemble Species by Autonomous Partitioning) or phylograms (bPTP,

Bayesian Poisson Tree Processes), to offer complementary perspectives (**DeSalle, 2024**).

First, we employed the ASAP method on our MAFFT alignment, with the following

adjustments to meet ASAP requirements: five GenBank sequences (MF694315,

MF694313, MF694317, MF694318, and MF694320) were removed to ensure overlap

across sequences. ASAP is a frugal, computationally efficient method, which is not

based on the reconstruction of phylogenies (**Puillandre et al., 2021**). Rather, this

distance-based approach identifies species partitions by detecting barcode gaps—the

difference between small intraspecific and large interspecific sequence distances

(**Puillandre et al., 2021**).

ASAP uses an ascending hierarchical clustering algorithm, which sequentially

merges sequences, based on pairwise distances, until all sequences are grouped within

the same partition (**Additional file 1**). For each partition, a *p*-value (likelihood of

panmixia, *i.e.*, a low *p*-value signifies that merging the proposed subsets would lead to

intra-subset genetic diversity that is higher than expected within a single panmictic

species), and W-score (barcode gap width metric, calculated using the threshold

distance, *i.e.*, the midpoint between the distance that triggered the merging of subsets

and the distance before merging) are computed (**Additional file 1**). A larger W-score

reflects a larger barcoding gap, *i.e.*, species are well separated. After clustering, the *p*-

value and W-score of each partition are sorted by ascending and descending order,

respectively, to compute the ASAP-score (averaged ranks of *p*-value and W-score for

each partition). Consequently, smaller ASAP-scores reflect better partitions. Its main

advantage over other automated barcode gap discovery approaches is that the maximal

intraspecific genetic distance does not need to be specified by the user *a priori*, which

makes it suitable for under-researched taxa such as African rope squirrels. We used the

most recent ASAP tool, (**Puillandre et al., 2021**), and compared simple p-distances,

Jukes-Cantor (JC69) and Kimura 2-parameter (K80) substitution models to investigate

whether substitution models impacted the proposed species delimitation. Only results

from JC69 are presented here.

For additional robustness, we applied the bPTP (**Zhang et al., 2013**) method. This

method uses phylogenetic tree branching patterns to propose species partitions (*i.e.*,

evaluating whether the number of substitutions follow a Poisson distribution within

versus between species) and provides Bayesian support values for these partitions.To

meet bPTP requirements (**Zhang et al., 2013**), we ran an IQ-TREE analysis (using

default settings from the IQ-TREE tool, (**Nguyen et al., 2015; Hoang, et al., 2017;**

Kalyaanamoorthy et al., 2017; Minh et al., 2020) on the MAFFT alignment used in

the ASAP method, with the following adjustments: only *Funisciurus* sequences, non-

redundant haplotypes, and fully overlapping sequences were included, for a total of 47

sequences. The resulting phylogenetic tree was used as an input file for the bPTP

analysis (**Additional file 2**).

Finally, we compared the species partitions proposed by ASAP and bPTP with the

topology of the FastTree and the results of discriminant analyses based on morphometric

data.

3. RESULTS

3.1. TRAPPING EFFORT

During a total of 99,035 trapping nights, the trapping effort resulted in the capture of

429 squirrels belonging to the genus *Funisciurus*, (success rate of 0.053%). Of these

squirrels, 301 were caught on the left bank of the Congo River and 128 on the right

bank. No reliable classification could be established based on external morphometric

characteristics alone.

3.2. SPECIES PARTITIONING

Overall, approaches based on mitochondrial sequences yielded between four and six

MOTUs, when excluding singletons and outgroups. The following MOTUs could be

identified based on ASAP and bPTP results: FaLB, FaRB, Fc, FbLB and Fx. Based on

ASAP results only, FbLB was further split into FbLB1 and FbLB2. The general species

partitioning obtained using mitochondrial data was supported by craniometric data, with

the exception of Fx, for which an insufficient number of cranial measures could be

taken.

3.2.1. *SPECIES PARTITIONING USING ASAP;* Using the best available species

partition, with the lowest ASAP score (ASAP score = 1.50, **Table 1**, hereafter ASAP1),

seven MOTUs of *Funisciurus* were distinguished, in addition to two outgroup MOTUs.

However, one MOTU is represented by a single sequence (YK156), giving a total of six

well-supported MOTUs consisting of two or more specimens each (**Figure 1,**

Additional file 1). Using the second-best species partition, with the second-lowest

ASAP score, (ASAP score = 3, **Table 1**), eight MOTUs of *Funisciurus* were

distinguished, in addition to two outgroup MOTUs. However, two MOTUs are

represented by a single sequence each (YK156 and NDU803), resulting in the same six

well-supported MOTUs identified in the first partition (**Additional file 1**). Using the

smallest *p*-value to determine the best available species partition (ASAP score = 5.50, p

= 0.00022, **Table 1,** hereafter ASAP2), five *Funisciurus* MOTUs can be distinguished,

in addition to two outgroup taxa. These include, again, a MOTU represented by a single

sequence (YK156), yielding a total of four well supported MOTUs consisting of two or

more specimens each. In contrast to previous partitions, FaLB, FaRB and Fc are merged

within a single MOTU (**Figure 1, Additional file 1**).

3.2.2. *SPECIES PARTITIONING USING bPTP;* Using the bPTP method applied to

our IQ-TREE, six MOTUs of *Funisciurus* were identified with support values ranging

from 0.697-0.984. However, one of them consists of a single sequence (YK110, support

value: 1.000). Thus, bPTP identifies five well supported MOTUs represented by two or

more specimens: Fx (0.984), Fc (0.900), FaLB (0.820), FaRB (0.817), and FbLB (0.697)

(**Figure 1, Additional file 2**).

3.2.3. *SPECIES PARTITIONING BASED ON EXTERNAL MORPHOMETRIC*

CHARACTERISTICS AND SEXUAL DIMORPHISM; Comparing Student's t-tests

with the craniometric (**Supplementary Table S3**) and external (**Supplementary Table**

S4) measurements of males and females of each species did not reveal any sexual

dimorphism. In all subsequent analyses, sexes were therefore lumped.

The discriminant analysis carried out on the external measurements (Weight,

Length of hind foot, Length of ear, Length of tail, Length of body) revealed low

discriminating power (Wilk's Lambda = 0.229) (**Supplementary Figure S3**). There is

no distinction between *F. anerythrus* from the right bank and from the left bank. *F.cf*

bayonii (Fb) is clearly separated from *F. anerythrus* along root 1. No external

measurements are available for *F.cf congicus*.

3.2.4. *SPECIES PARTITIONING BASED ON CRANIAL MORPHOMETRIC*

CHARACTERISTICS; Unlike external measurements, craniometric measurements of

adult rodents offer greater stability and are among the most effective methods for

identifying rodent species. For this reason, 220 adult squirrel skulls were measured and

23 measurements were taken on each skull (**Supplementary Table S2**).

A forward discriminant analysis, (**Supplementary Figure S5**) omitting measures

M8 (width of the interorbital narrowing), M9 (width of the zygomatic arch) and M19

(maximum width of the cranial cavity) to maximize the number of observations for *F. cf*

congicus (Fc), measures (M1: maximum length of the skull; M2: condilo-basal length;

M3: henselion-basion length; M4: henselion-palation length; M5: length of the cleft

palate; M6: length of the diastema; M7: distance between the anterior edge of the

alveolus of M1 and the cutting edge of the upper incisor; M10: minimum width of the

palate; M11: length of the upper molar row; M12 : external width of upper molar row at

M1; M13: maximum molar width; M14: minimum width of zygomatic plate; M15:

maximum width of nasals; M16: maximum length of nasals; M17: length of lower molar

row; M18: length of the tympanic bulla; M20: depth of the upper incisors; M21: height

of the rostrum at the anterior edge of the alveolus of M1; M22: maximum width of the

rostrum; M23: distance between the extreme points of the coronoid and angular

processes) reveals a good discrimination. *F. anerythrus* (FaRB and FaLB) and *F cf*

bayonii (FbLb) are clearly separated along root 1.

F.cf congicus (only six sequenced specimens with mesured skulls, see **figure 1**) is

clearly different from both other species. *F. anerythrus* Right Bank (FaRB) and Left

Bank (FaLB) are well differentiated along root 2. The number of sequenced specimens

of the two genotypes of *F.cf bayonii* (FbLB1 and FbLB2) are too small to separate them

in this analysis (**Supplementary Figure S5**).

A discriminant analysis, performed on the OTUs of *F. anerythrus* from the right

bank and left bank, as well as FbLB1 and FbLB2 using a craniometric dataset limited to

the skulls of specimens with matching genetic sequences, while omitting the often

missing M6, M7 and M20 measurements (**Supplementary Figure S3**), reveals subtle

differences between *F. cf bayonii* sp1 (FbLB1) and *F. cf bayonii* sp2 (FbLB2). However,

the limited availability of specimens with both sequence data and craniometric

measurements has hampered the differentiation of these potentially distinct OTUs. The

same figure also reveals clear differences between *F. anerythrus* from the two banks of

the Congo River (FaRB and FaLB) and no difference between *F. anerythrus* from the

right bank (FaRB and Fx in **Supplementary Figure S6**).

3.2.5 *OVERALL SPECIES PARTITIONING;* Our most conservative results, found

using a barcoding gap analysis with partition scoring where we retained the partition

with the lowest *p*-value (ASAP2), suggest the presence of four MOTUs consisting of at

least two specimens each. Here, *F. anerythrus* and *F. cf congicus* form one MOTU,

whereas *F. cf bayonii*, and a previously undescribed *Funisciurus* species (here, Fx), each

form another MOTU.

Our second most conservative result, using a tree-based algorithm inferring

species delimitation based on the distribution of substitutions between and within

species (bPTP), suggests the presence of five MOTUs. These are *F. cf bayonii* (FbLB),

the undescribed *Funisciurus* species (Fx), *F. anerythrus* individuals found on the left

bank of the Congo River (FaLB), *F. anerythrus* individuals found on the right bank of

the Congo River (FaRB), and *F. congicus* (Fc).

The discriminant analysis conducted on craniometric data suggests the presence of

five OTUs, however differently distributed: *F.cf bayonii* individuals form two weakly

supported OTUs (FbLB1 and FbLB2), while Fc, FaLB and FaRB each form a well-

supported OTU. While Fx does not appear to be different from FaRB, the limited

number of measurable skulls for Fx prevents any conclusions about craniometric

differences between Fx and FaRB.

Last, our least conservative results, based on a barcoding gap analysis with

partition scoring where we retained the partition with the lowest ASAP score (ASAP1),

suggest the presence of six well-supported MOTUs of *Funisciurus* represented by at

least two individuals each. These MOTUs are Fc, FaLB, FaRB, Fx, FbLB1, and FbLB2,

and are in line with the FastTree topology (**Figure 1**).

3.3. DISTRIBUTION AND HYDROLOGICAL BARRIERS

The six main well-supported taxa occupy distribution areas separated by hydrological

barriers (**Figure 2**). The previously undescribed *Funisciurus* species (Fx) originates

exclusively from the right bank of the Congo River (Figure 2a) and the *F. cf bayonii*

species (FbLB1 and FbLB2) (Figure 2a) are sympatric and distributed exclusively on the

left bank of the Congo River. In addition, FaRB is distributed exclusively on the right

bank of the Congo River, and FaLB is distributed exclusively on the left bank of the

Congo River. *F. cf congicus* (Fc) is distributed exclusively on the left bank of the Kasaï

River (Figure 2b).

4. DISCUSSION

This study had two main objectives: first, to ascertain the number of *Funisciurus* OTUs

present in the Congo Basin; and second, to investigate the role of the Congo River as a

geographical barrier between *Funisciurus* taxa. For this purpose, we integrated
morphometric, molecular and zoogeographic data from the most extensive *Funisciurus*

species dataset to date. We first identified a discrepancy between the previously

described species of *Funisciurus* in our study area, and the individuals present in our

dataset (**Table 2, Figure 1**). We then found that both the Congo River and the Kasaï

River delineate the distribution of *Funisciurus* OTUs (**Figure 2**). This highlights the

importance of hydrological barriers in the speciation processes of small arboreal

mammals in the Congo Basin and calls for an update in the taxonomy and IUCN

distribution maps of rope squirrels.

Previously, five rope squirrel species had been described in our study area: *F.*

anerythrus (Thomas, 1890), *F. bayonii* (Bocage, 1890), *F. congicus* (Kuhl, 1820), *F.*

pyropus (F. Cuvier, 1833), and the mountain-dwelling *F. carruthersi* Thomas, 1906

(**Kingdon & Hoffmann, 2013; Thorington et al., 2012; Gerrie et al., 2016**).

To minimise the risk of OTU over-segregation or fusion in our analysis (**Sloan et**

al., 2017; Nesi et al., 2011), our final OTU partitioning is based on both craniometric

and molecular OTU delineation, as well as a zoogeographic analysis of their

distribution. To do so, we have integrated multiple OTU delineation methods (**Demos et**

al., 2013), *i.e.,* a discriminant analysis of craniometric data, two barcoding gap analyses

with partition scoring (ASAP1 and ASAP2), and a tree-based algorithm inferring species

delimitation based on the distribution of substitutions between and within species

(bPTP), with OTU distribution maps, where well-supported OTUs distributed in the

same hydrological basin indicate the presence of non-interbreeding, sympatric species;

while allopatric OTUs could theoretically interbreed, but are prevented to do so by

hydrological barriers. Thus, using our most conservative results (ASAP2) and our

updated OTU distribution, we suggest that there exist four *Funisciurus* species in the

Congo Basin, namely *F. anerythrus* (composed of three allopatric subspecies: Fc, FaLB,

and FaRB); two sympatric, but genetically and craniometrically different, thus non-

interbreeding *F. cf bayonii* species (FbLB1, FbLB2); and the previously undescribed

Funisciurus species (Fx).

Several key points regarding the taxonomy of *F. anerythrus*, *F. bayonii*, and an

undescribed *Funisciurus* species warrant further discussion. According to previous

studies, *F. anerythrus*—originally described from Buguera, Uganda—has a distribution

spanning from Uganda to Angola and Togo (**Thorington & Hoffmann, 2005**), covering

all central Congolese forest blocks (**Colin, 1991**). **Kingdon & Hoffmann (2013)**

identified four subspecies based on geographical and morphological variations. The

southernmost, *F. a. anerythrus*, occurs from western Uganda to southern DRC, reaching

Mount Kabobo and northern Angola. It has a reddish-brown dorsal coat and a yellow to

red ventral coat. The northernmost, *F. a. bandarum*, ranges from the Central African

Republic to southern Chad, featuring a buff-tinged dorsal coat and a pale grey to orange

ventral side. Either of these may correspond to our OTU FaRB, but limited skull and

DNA data prevent verification. *F. a. mystax*, found in southern Cameroon and Gabon,

has a reddish-brown dorsal and orange ventral coat, while *F. a. raptorum*, endemic to

Benin and Nigeria, displays a grey dorsal and whitish ventral surface. Due to

morphological similarities and limited overlap in our dataset, these findings reinforce the

need for a thorough taxonomic reassessment of the *F. anerythrus* complex.

Conversely, previous studies indicate that *F. bayonii* is found mostly in western

DRC, with no described subspecies (**Thorington et al., 2012; Kingdon & Hoffmann,**

2013; Gerrie et al., 2016). This species has only been documented outside the Central

African forest block and was previously considered a species of wooded savannah

mosaics (**Thorington et al. 2012**). However, our results show that two species

morphologically similar to *F. bayonii* are present in Yoko and Yasikia, located near the

left bank of the Congo River, in evergreen tropical forests. This result corroborates the

need for further taxonomic revision to clarify the status of these species, including their

ecological niches.

Last, we report the existence of a species that does not cluster with either *F.*

anerythrus nor *F.* cf *bayonii*, and which is distributed north of the Congo River (e.g.,

UMA30). Its range does not overlap with the known range of the mountain-restricted *F.*

carruthersi Thomas, 1906, and it does not display the typical red-leggedness of *F.*

pyrropus (F. Cuvier, 1833). Although its appearance resembles that of individuals of *F.*

anerythrus found on the right bank of the Congo River (FaRB), all OTU delineation

techniques based on genetic data suggest that it is a yet undescribed species of rope

squirrel.

Beyond taxonomic considerations, we highlight that the distribution ranges of

these squirrels are different, and much smaller than previously thought, rendering their

populations more vulnerable to environmental change and habitat degradation (**Steiner**

M. & Huettmann F., 2023). Current IUCN distribution maps indicate that *F. congicus*'

northeastern expansion is limited by the Congo River, while *F. bayonii* is confined

southwest of the Kasaï River. (data retrieved from IUCN Red List

485 (https://www.iucnredlist.org)**; Gerrie et al., 2016**). However, our findings contradict

486 this. We observed that *F. congicus* (Fc) is restricted southwest of the Kasaï River, with

487 no specimens found between the Kasaï and Congo Rivers. Additionally, unlike the

488 IUCN map for *F. bayonii*, we found that the sympatric FbLB1 and FbLB2 OTUs are

489 exclusively limited to this interfluvial region, where the Kasaï River restricts

490 southwestern expansion, and the Congo River blocks northeastern expansion. Similarly,

491 contrary to IUCN data for *F. anerythrus*, the Congo River separates FaLB (southwest)

492 and FaRB (northeast). The undescribed Fx is found only northeast of the Congo River

493 (**Figure 2**). These findings highlight the role of hydrological barriers in driving

494 allopatric speciation in arboreal Sciuridae, consistent with research on primates

495 (**Anthony et al., 2007; Harcourt & Wood, 2012; Fonteyn et al., 2023**) and small

496 mammals rodentia (**Katuala et al., 2008; Kennis et al., 2011**). They further highlight

497 that the distribution range of each *Funisciurus* OTU is smaller than previously thought

498 (**Supplementary Figure S7**). Consequently, habitat degradation or environmental

499 change may have unexpectedly severe impacts on *Funisciuru*s populations

500 (**Supplementary Figure S8**), of which regular monitoring is necessary. We therefore

501 advocate for updated IUCN distribution maps and regular monitoring of the *Funisciurus*

502 OTUs identified above, to ensure that conservation measures can be taken where

503 appropriate.Regular monitoring will also support research on the potential

504 epidemiological importance of *Funisciurus* species.

505 **CONCLUSION**

506 Our morphometric and molecular analyses enabled us to identify six main rope squirrel

507 OTUs in the Congo Basin, namely *F. cf congicus*, *F. anerythrus*_LB, *F. anerythrus*_RB,

508 *Funisciurus sp*, *F. cf bayonii*_LB1, and *F. cf bayonii*_LB2. Further integration of these

509 approaches with zoogeographic analyses suggests that Fc (*F. cf congicus* (Kuhl, 1820)), *F.*

*anerythrus*_LB and *F. anerythrus*_RB are subspecies of *F. anerythrus* (Thomas, 1890),

while the two sympatric and non-interbreeding *F. cf bayonii*_LB1, and *F. cf bayonii*_LB2

each are a separate species, possibly forming a complex with *F. bayonii* (Bocage, 1890).

Last, we report on a previously undescribed species of *Funisciurus* (Fx). We further

highlight the role of the Congo and Kasaï as hydrological barriers, delineating the

distribution of our OTUs, with *F. cf congicus* found exclusively southwest of the Kasaï, *F.*

*anerythrus*_LB, *F. cf bayonii*_LB1 and *F. cf bayonii*_LB2 found between the Kasaï and

Congo, and *F. anerythrus*_RB and Fx found northeast of the Congo River. These results

call for an update of IUCN maps, and for regular monitoring of *Funisciurus* populations,

which may be more vulnerable to environmental change and habitat degradation than

previously thought. Last, we highlight that further taxonomic and ecological studies are

needed to describe each OTU and ascertain its ecological niche, which will support

research on zoonotic pathogen spillover risks associated with each OTU. In this

perspective, we also recommend a complete review of the taxonomic status of other tree

squirrels, in particular the genera *Paraxerus, Heliosciurus* and *Protoxerus.*

ACKNOWLEDGEMENTS

This research was funded by the Flemish Interuniversity Council South-Initiative (VLIR-

SI) (AL, EV), the Flemish Interuniversity Council (VLIR-CUI) (GG, EV), the Research

Foundation—Flanders (G051322N), and a FED-tWIN scholarship funded by the Belgian

federal government (Prf-2019-prf004_OMEga) (SG). The CEBioS program (Royal

Belgian Institute of Natural Sciences) provided financial support for the CSB team's

fieldwork (PB, CM, GG) and research stays in Belgium (PB).

We thank the Faculty of Science and the Centre de Surveillance de la Biodiversité

of University of Kisangani for their supervision. We also thank Emmanuel Gilissen and

Marthys Rotonda for allowing us access to Squirrel specimens of the collection of

MRAC-Tervuren.

REFERENCES

Anthony NM, Johnson-Bawe M, Jeffery K, Clifford SL, Abernethy KA, Tutin CE, Lahm

SA, White LJT, Utley JF, Wickings EJ, et al. 2007. The role of Pleistocene refugia

and rivers in shaping gorilla genetic diversity in central Africa. Proceedings of the

National Academy of Sciences, 104(51), 2043220436.

https://doi.org/10.1073/pnas.0704816105

Baelo P, Asimonyio J, Amundala N, Akaibe D, Gambalemoke S, Erik V, Anne L, Bongo

G, Koto-te-Nyiwa Ngbolua K. 2018. Preliminary Inventory of Squirrels (Sciuridae,

Rodentia) of Kisangani Region in the Democratic Republic of the Congo. American

Journal of Zoology . Vol. 1, No. 1, pp. 15-19.

https://doi.org/10.11648/j.ajz.20180101.13

Chen NLiG, Liszewski MK, Atkinson JP, Jahrling PB, Feng Z. 2012.Virulence

differences between monkeypox virus isolates from West Africa and the Congo

basin. Virology Journal. 340:46–63. https://doi.org/10.1016/j.virol.2005.05.030

Colyn M. 1991. L'importance zoogéographique du bassin du fleuve Zaïre pour la

spéciation: le cas des primates simiens. Annales-Musée Royal de l'Afrique

Centrale-Sciences Zoologiques (Belgium). Annalen-Koninklijk Museum voor

Midden-Afrika-Zoologische Wetenschappen. no. 264. http://agris.fao.org

Curaudeau M, Besombes C, Nakouné E, Fontanet A, Gessain A, and Hassanin A. 2023.

Identifying the most probable mammal reservoir hosts for monkeypox virus based

on ecologicalniche comparisons. Viruses, 15(3),

727. https://doi.org/10.3390/v15030727

Demos TC, Agwanda B, Hickerson MJ. 2013. Integrative taxonomy within the

Hylomyscus denniae complex (Rodentia: Muridae) and a new species from Kenya.

Journal of Mammalogy 2013, 95(1):E1-E15. https://doi.org/10.1644/13-MAMM-

A-268

DeSalle R. 2024. DNA Barcoding: Methods and Protocols ISBN978-1-0716-3580-

3ISBN978-1-0716-3581-0(eBook) https://doi.org/10.1007/978-1-0716-3581-0

Fa J E, Peres CA, & Meeuwig J. 2002. Bush meat exploitation in tropical forests: an

intercontinental comparison. Conservation biology, 16(1), 232-237.

https://doi.org/10.1046/j.1523-1739.2002.00275.x

Falendysz E A, Lopera JG, Doty JB, Nakazawa Y, Crill C, Lorenzsonn F, Kalemba, F.,

Ronderos, L.N, Mejia, M.D., Malekani, J.M, Karem K., Carroll D.S., Osorio J.O.,

and Rocke, T. E. (2017). Characterization of Monkeypox virus infection in

African rope squirrels (*Funisciurus sp*.). PLoS neglected tropical diseases, 11(8),

e0005809. https://doi.org/10.1371/journal.pntd.0005809

Fonteyn D, Vermeulen C, Gorel AP, Silva de Miranda PL, Lhoest S, and Fayolle A.

2023. Biogeography of central African forests: Determinants, ongoing threats and

conservation priorities of mammal assemblages. Diversity and

Distributions, 29(6), 698-712. DOI: 10.1111/ddi.13677

Gerrie R, Kennerley and Koprowski RJ. 2016. *Funisciurus bayonii*. The IUCN Red List

of Threatened Species; http://dx.doi.org/10.2305/IUCN.UK.2016-

3.RLTS.T8757A22270331.

Han BA, Schmidt JP, Bowden SE et Drake JM. 2015. Réservoirs de rongeurs de futures

zoonoses. Actes de l'Académie nationale des sciences, 112(22), 7039-7044.

https://doi.org/10.1073/pnas.1501598112

Harcourt AH & Wood MA. 2012. Rivers as Barriers to Primate Distributions in

Africa. International Jornal of Primatolology **33**, 168–183

https://doi.org/10.1007/s10764-011-9558-z

Hoang DT, Chernomor O, Haeseler A, Minh BQ, and Vinh LS. 2017. UFBoot2:

Improving the ultrafast bootstrapapproximation. Molecular Biology and Evolution,

in press. https://doi.org/10.1093/molbev/msx281

Bernd Hoffmann DVM., Dennis Tappe MD, Dirk Höper MSc, Christiane Herden DVM,

Annemarie Boldt MD, Christian Mawrin MD, Olaf Niederstraßer MD, Tobias

Müller MD, Maria Jenckel MSc, Elisabeth van der Grinten DVM, et al. 2015. A

variegated squirrel bornavirus associated with fatal human encephalitis. New

England Journal of Medicine, 373(2), 154-162.

https://doi.org/10.1056/NEJMoa1415627

Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A, and Jermiin LS. 2017.

ModelFinder: Fast model selection foraccurate phylogenetic estimates. Nature

Methods, 14:587–589.https://doi.org/10.1038/nmeth.4285

Katuala P, Kennis J, Nicolas V, Wendelen W, Hulselmans J, Verheyen E, Van Houtte

N, Dierckx T, Dudu A andLeirs H. 2008. The presence of *Praomys*, *Lophuromys*,

and *Deomys* species (Muridae, Mammalia) in the forest blocks separated by the

Congo River and its tributaries (Kisangani region, Democratic Republic of

Congo). Mammalia, 72(3), 223-228. https://doi.org/10.1515/MAMM.2008.044

Kennis J, Nicolas V, Hulsemans J, Katuala GBP, Wendlen W, Verheyen E, Dudu A,

Leirs H, 2011. The impact of the Congo River and its tributaries on the rodent

genus Praomys: speciation origin or range expansion limit?, Zoological Journal of

the Linnean Society, Volume 163, Issue 3, November 2011, Page 983-1002,

https://doi.org/10.1111/j.1096-3642.2011.00733.x

Kinganda LE, Amuri AA, Nuñez FN, Makangara JC, Pratt C, Vakaniaki EH, Hoff NA,

Luakanda GN, Sabiti SN, Mulopo NM et al. 2024. Clade I mpox virus genomic

diversity in the Democratic Republic of the Congo, 2018–2024: predominance of

zoonotic transmission. Cell journal. DOI: 10.1016/j.cell.2024.10.017

Kingdon J & Hoffmann M. 2013. 'Mammals of Africa. Volume V: Carnivores,

Pangolins, Equids and Rhinoceroses' 2nd ed. Ed J Kingdon. (Bloomsbury

Publishing Plc: London) https://doi.org/10.5860/CHOICE.50-4188

Kumakamba C, Niama FR, Muyembe F, Mombouli JV, Kingebeni PM, Nina RA, Ngay

LI, Bounga G, N'Kawa F, Goma NC et al. 2021. Coronavirus surveillance in

wildlife from two Congo basin countries detects RNA of multiple species

circulating in bats and rodents. PLoS One, 16(6), e0236971.

https://doi.org/10.1371/journal.pone.0236971

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Haeseler A,

Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era, Molecular Biology and Evolution,

Volume 37, Issue 5, May 2020, Pages 1530–

1534, https://doi.org/10.1093/molbev/msaa015

Montgelard C, Sophie B, Claire T, Olivier V, François M. 2002. Systématique

moléculaire de Sciurognathi (Rodentia) : les gènes mitochondriaux du

cytochrome *b* et de l'ARNr 12S soutiennent les Anomaluroidea (Pedetidae et

Anomaluridae) Volume 22, Numéro 2, février 2002, pages 220-233

https://doi.org/10.1006/mpev.2001.1056

Nakazawa Y, Mauldin MR, Emerson GL, Reynolds MG, Lash RR, Gao J, Zhao H, Yu

Li, Muyembe JJ et al. 2015. A phylogeographic investigation of African

monkeypox. *Viruses*, *7*(4), 2168-2184. https://doi.org/10.3390/v7042168

Nesi N, Nakouné E, Cruaud C, and Hassanin A. 2011. DNA barcoding of African fruit

bats (Mammalia, Pteropodidae). The mitochondrial genome does not provide a

reliable discrimination between *Epomophorus gambianus* and *Micropteropus*

pusillus. Comptes Rendus. Biologies, Volume 3334 (2011) no. 7, pp. 544-

554.https://comptes-rendus.academie

sciences.fr/biologies/articles/10.1016/j.crvi.2011.05.003/

Nguyen LT, Schmidt HA, Haeseler A, and Minh BQ. 2015. IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum likelihood phylogenies. Molecular

Biology and Evolution, 32:268-274. https://doi.org/10.1093/molbev/msu300

Otieno JR, Ruis C, Onoja AB, Kuppalli K, Hoxha A, Nitsche A, Brinkmann A, Michel

J, Kingebeni PM, Bamuleka DM, et al. 2025. Global genomic surveillance of

monkeypox virus. Natural Medicines **31**, 342–350 (2025).

https://doi.org/10.1038/s41591-024-03370-3

Price MN., Dehal PS, & Arkin AP. 2009. FastTree: computing large minimum evolution

trees with profiles instead of a distance matrix. Molecular biology and

evolution, 26(7), 1641-1650. https://doi.org/10.1093/molbev/msp077

Puillandre N, Brouillet S, Achez G. 2021. ASAP: assemble species by automatic

partitioning. Molecular Ecology Resources 21:609–620.

https://doi.org/10.1111/1755-0998.13281

Rimoin AW, Alfonso VH, Hoff NA, Doshi RH, Mulembakani P, Kisalu NK, Wright

LL. 2017. Human exposure to wild animals in the Sankuru Province of the

Democratic Republic of the Congo. Ecohealth, 14, 552-

563.https://doi.org/10.1007/s10393-017-1262-9

Schulze V, Lurz PW, Ferrari N, Romeo C, Steele MA, Marino S, Vittoria MM,

Calvignac SS, Schlottau K, Beer M, Ulrich RG, and Ehlers B. 2020. Search for

polyoma-, herpes-, and bornaviruses in squirrels of the family Sciuridae. Virology

journal, 17, 1-21. https://doi.org/10.1186/s12985-020-01310-4

Sikes RS, & Animal Care and Use Committee of the American Society of

Mammalogists. 2016. 2016 Guidelines of the American Society of Mammalogists

for the use of wild mammals in research and education. Journal of

mammalogy, 97(3), 663-688. https://doi.org/10.1093/jmammal/gyw078.

Sklenovská N, & Van Ranst M. 2018. Emergence of monkeypox as the most important

orthopoxvirus infection in humans. *Frontiers in public health*, *6*, 383729.

https://doi.org/10.3389/fpubh.2018.00241

Sloan DB, Havird JC, & Sharbrough J. 2017. The on-again, off-again relationship

between mitochondrial genomes and species boundaries. Molecular

ecology, 26(8), 2212-2236. https://doi.org/10.1111/mec.13959

StatSoft, Inc. 2014. STATISTICA (data analysis software system), version 12. Available

at: https://www.statsoft.com.

Steiner M. & Huettmann F. 2023. Chapter 1. Taxonomy for the Squirrels of the World:

Hairy Classifications and Conservation Category Games. in: Sustainable Squirrel

Conservation, https://doi.org/10.1007/978-3-031-23547-4_1

Thorington JRW, Koprowski JL, Steele MA, & Whatton JF. 2012. Squirrels of the

world. JHU Press.

Thorington RW & Hoffmann RS. Family Sciuridae. Pp 754-818 in Wilson D.E. &

Reeder, D.M. (Eds), (2005). Mammals species of the World: A taxonomic and

Geographic reference. The Johns Hopkins University Press, Baltimore; Vol. 2, 21-

42 pp.

Tiee MS, Harrigan RJ, Thomassen HA, & Smith TB. 2018. Ghosts of infections past:

using archival samples to understand a century of monkeypox virus prevalence

among host communities across space and time. Royal Society Open

Science, 5(1), 171089.

Van der Straeten E, & Dieterlen F. 1987. *Praomys misonnei*, a new species of Muridae

from Eastern Zaïre (Mammalia). *Stuttg. Beitr. Naturk*., 402: 1-11.

Verheyen W, & Bracke E. 1966. The influence of aging on the craniometrical characters

of *Praomys jacksoni* (De Winton 1897). Annales du Musée Royal d'Afrique

Centrale de Tervuren, 144, 91-101.

Verheyen WN, Colyn M, & Hulselmans J. 1996. Re-evaluation of the *Lophuromys*

nudicaudus HELLER, 1911 species-complex with a description of a new species

from Zaire (Muridae-Rodentia). Bulletin de l'Institut Royal des Sciences

Naturelles de Belgique Biologie, 66, 241-273.

Zachos FE, Burgin CJ, Wilson DE, Mittermeier RA, Rylands AB, Lacher TE, Sechrest

W. 2021. Illustrated Checklist of the Mammals of the World. Boxed set of two

volumes, with a foreword by Razan Al Mubarak. Mammalian Biology 101, 125–

126 (2021). https://doi.org/10.1007/s42991-020-00088-w

Zhang J, Kapli P, Pavlidis P, Stamatakis P. 2013. A general species delimitation method

With applications to phylogenetic placements. Bioinformatics 29:2869–2876.

https://doi.org/10.1093/bioinformatics/btt499

List of figures

Fig. 1. Proposed rope squirrel Operational Taxonomic Units using a combination of

methods. Above the branches, methods used to propose species partitions.

Fig. 2. Maps showing the distribution of the species and subspecies identified in

relation to the Congo River and its tributaries.

List of table

Table 1. Species partitions ranked in ascending order of ASAP scores. The *p*-value reflects

the likelihood that Partition is different from Partition$_{n-1}$ and is ranked in ascending order.

The threshold distance is used to compute the W-score and represents the mid-point

between the distance that triggered the merging of subsets into Partition and the previous

distance in the list. The W-score is the relative barcode gap width associated with

Partition$_{n-1}$ and is ranked in descending order. The ASAP-score is the average of the ranks

of the *p*-value and of the W-score. A smaller ASAP-score thus signifies a better partition

(**Puillandre et al., 2021**). Read: Considering an ASAP-score of 2.00 and a p-value of

0.0177, 9 subsets are identifiable in the dataset, including two outgroups.

Number of subsets	ASAP-score	P-val (rank)	W (rank)	Threshold dist.
*9	2.00	1.77e-02 (3)	7.83e-04 (1)	0.020430
5	3.50	1.96e-01	2.03e-04 (3)	0.118381
*8	4.50	3.46e-03 (02)	4.68e-05 (7)	0.029208
6	5.00	3.80e-04 (1)	2.56e-05 (9)	0.075144
*12	7.00	5.45e-01 (8)	4.72e-05 (6)	0.0008821
8	11.50	6.87e-01 (15)	3.11e-05 (8)	0.052757
*10	12.00	8.40e-01 (22)	5.20e-04 (2)	0.015671
59	14.00	5.73e-01 (10)	6.22e-06 (18)	0.000899
60	15.50	6.79e-01 (14)	6.22e-06 (17)	0.000450
2	15.50	8.14e-01 (20)	1.15e-05 (11)	0.180112

Figures

Figure 1. Proposed rope squirrel Operational Taxonomic Units using a combination of

methods. Above the branches, methods used to propose species partitions. Cranio:

Craniometry; ASAP1: Assemble Species by Autonomous Partitioning (ASAP), lowest

ASAP score; ASAP2: *idem*, lowest p-value; bPTP: bayesian Poisson Tree Processes;

RB: right bank of the Congo River; LB: left bank of the Congo River. Fx: previously

undescribed *Funisciurus* species; Fb: *Funisciurus cf bayonii*; Fc: *Funisciurus cf*

congicus; Fa: *Funisciurus anerythrus*. The tree was made using FastTree on MAFFT-

assembled cytochrome b sequences.

736   **Figure 2.** Maps showing the distribution of the species and subspecies identified in

737   relation to the Congo River and its tributaries, (**a**) *Funisciurus sp* from the right bank of

738   the Congo River and the sympatric *F. cf bayonii sp1* and *F. cf bayonii sp2* on the left

739   bank of the Congo River; (**b**) *F. anerythrus_RB* on the right bank, *F. anerythrus_LB* on

740   the left bank of the Congo River, and *F. cf congicus* on the left bank of the Kasaï River.

Full e-mail addresses of authors

743   **Pascal Baelo,** pascal.baelo@unikis.ac.cd **ou** baelopascal@gmail.com

744   **Léa Fourchault,** lfourchault@naturalsciences.be

745   **Claude Mande,** claude.mande@unikis.ac.cd

746   **Anne Laudisoit,** anne@mwambao.or.tz **ou** anne@gmail.com

747   **Nicaise Amundala**, nicaise.amundala@gmail.com

748   **Nicolas Laurent**, nicolas.laurent@ou.nl

749   **Ganiyat Temidayo Saliu,** ganiyat.temidayo.saliu@vub.be

750   **Joachim Mariën**, Joachim.Marien@uantwerpen.be

751   **Sophie Gryseels**, sophie.gryseels@uantwerpen.be

752   **Guy-crispin Gembu**, gembuguycrispin@gmail.com

753   **Jan Hulselmans**, jan.hulselmans@uantwerpen.be

754   **Erik Verheyen**, everheyen@naturalsciences.be

Contributions of the authors

PB, EV and JH conceptualized the study and designed the methodology. PB collected

the data and identified the species in the field. LF, PB, GTS and NL generated molecular

data. PB, JH, and EV analyzed the morphometric data. PB produced the map. LF, SG

and EV analysed the molecular data. PB, LF, JH and EV wrote the original version. PB,

LF, CM, AL, NA, NL, GTS, JN, JM, SG, GG, JH and EV reviewed and edited the

document. All authors approved the final manuscript.

Data availability

765   The generated and/or analyzed datasets can be made available for research purposes upon

766   request to the corresponding authors. The sequences from this study will be submitted to

767   GenBank upon acceptance.

769   **Conflict of interest declaration.**

770   The authors declare no conflict of interest.

772