***Streptococcus mitis* bacteriocins drive contact-dependent lysis of**

***S. pneumoniae* facilitating transformation in multispecies environments**

João Borralho[1], João Lança,[1], Joana Bryton[1], Wilson Antunes[2], Raquel Sá-Leão[1]*

[1] Laboratory of Molecular Microbiology of Human Pathogens, Instituto de Tecnologia Química e

Biológica António Xavier, Universidade Nova de Lisboa, Oeiras, Portugal

[2] Centro de Investigação da Academia Militar (CINAMIL), Unidade Militar Laboratorial de Defesa

Biológica e Química (UMLDBQ), Av. Dr. Alfredo Bensaúde, 1849-012, Lisboa, Portugal.

* Corresponding author: rsaleao@itqb.unl.pt

Keywords: Horizontal gene transfer, competence, interspecies interactions, predation, biofilms

Abstract

Natural competence allows bacterial species like *Streptococcus pneumoniae* and *S. mitis* to acquire environmental DNA, driving horizontal gene transfer (HGT) and adaptation. In *S. pneumoniae*, a human pathogen, competence-induced predation is well characterized and involves the release of bacteriocins and a murein hydrolase to lyse noncompetent siblings and liberate DNA. In contrast, in the human commensal *S. mitis*, mechanisms mediating DNA acquisition remain poorly understood. Here, we identify a diverse set of competence-associated bacteriocins (*cab*) that are produced by *S. mitis* during the late phase of competence. We focus on one bacteriocin pair, CabAB, that triggers contact-dependent growth inhibition and lysis of *S. pneumoniae* through activation of the major pneumococcal autolysin LytA. We demonstrate that CabAB compromises *S. pneumoniae* membrane integrity, leading to formation of intracellular membrane aggregates and the release of cytoplasmatic content, thereby increasing available DNA, which enhances HGT from *S. pneumoniae* to *S. mitis* in biofilms. These findings uncover a mechanism of interspecies predation and gene acquisition, revealing a critical role for competence-associated bacteriocins in shaping evolutionary dynamics of streptococci.

Importance

Many streptococci are naturally competent, acquiring environmental DNA through transformation. This includes pathogens like *S. pneumoniae* and commensals like *S. mitis,* which can exchange genetic material through horizontal gene transfer (HGT). For example, *S. mitis* can acquire pneumococcal capsules, leading to its misidentification in polymicrobial samples such as those obtained from the upper respiratory tract. Understanding the drivers of HGT between these species is therefore critical. Here, we characterize a competence-induced bacteriocin cluster in *S. mitis*. These bacteriocins lyse pneumococci, promoting DNA release and enhancing gene transfer in dual-species biofilms. Our findings uncover a mechanism by which competence-associated predation promotes interspecies HGT, shaping the evolution and epidemiology of streptococcal populations.

Introduction

Natural competence for genetic transformation is a developmental program found in certain bacterial species that enables the uptake of naked DNA from the environment. Once internalized, this DNA can be incorporated into the genome through homologous recombination, making natural transformation one of the main modes of horizontal gene transfer (HGT). This phenomenon was first described in *Streptococcus pneumoniae* (or pneumococcus), a species capable of exchanging pathogenic traits such as the capsule by taking up DNA from other pneumococci (1).

In *S. pneumoniae*, competence is a tightly regulated transient physiological state which is governed by a cell-cell signaling system. The competence-stimulating peptide (CSP), encoded by *comC*, is processed and exported via the ATP-binding cassette transporter ComAB. Extracellular CSP is sensed by the membrane-bound histidine kinase ComD (2-4), which undergoes autophosphorylation triggering subsequent phosphorylation of the response regulator ComE (5). Phosphorylated ComE increases expression of the early competence cascade including *comAB* and *comCDE*, creating a positive feedback loop. It also induces expression of the two copies of the alternative sigma factor (*sigX1* and *sigX2*), marking the onset of the late phase of competence (6).

In the late phase of competence, expression of all genes required for DNA uptake, processing and recombination – the transformasome – is induced (7). Here, pneumococci produce the competence-induced bacteriocins A and B (CibAB), along with the immunity protein CibC, which provides self-immunity to CibAB (8). Because competence may not be synchronized across the whole population (9-11), CibAB release on solid media may lead to lysis of genetically identical (kin) noncompetent cells that have not yet expressed *cibC*. Additionally, expression of the murein hydrolase CbpD, against which competent cells are protected via ComM expression, is essential for kin predation in liquid cultures. Notably, CbpD has also been implicated in lysis of non-kin, closely related streptococci, thereby expanding the gene pool from which *S. pneumoniae* may acquire DNA (12). CibAB and CbpD have also been associated to competitive interactions *in vivo*, where their expression by colonizing pneumococci provides a competitive advantage over other *S. pneumoniae* invaders (13).

A close relative of the pneumococcus, *S. mitis*, is also naturally transformable. *S. mitis* is a

common commensal of the human nasopharynx and oral cavity, where it typically establishes long-

term colonization without causing disease (14). Despite close phylogenetic relationship with *S.*

pneumoniae, and a strikingly similar competence regulatory network, *S. mitis* exhibits key differences

in terms of bacteriocin content. While *S. mitis* typically encodes *cbpD*, most strains lack *cibAB*,

suggesting that *S. mitis* may rely on an alternative mechanism for predation during competence (15-

17).

Although the frequency at which *S. mitis* acquires DNA from *S. pneumoniae* is not well

established, such events have been documented. These include, for example, the acquisition of

pneumococcal *pbp2x* mosaic blocks (18-20). Some *S. mitis* have also been found to express some

pneumococcal or pneumococcal-like capsules, which can cross-react with antisera against serotypes

1, 5, 17F, 18A, 19C, 21, 39 and 40 (21-24).

A competence-associated bacteriocin (*cab*) cluster, which is absent in *S. pneumoniae*, has

been identified in *S. mitis*. This cluster is located upstream of the competence transporter *comAB* and

can encode multiple *cab* bacteriocins distinct from CibAB (15, 16, 25-27). However, the role of *cab* in

S. mitis predation of kin and non-kin, and its contribution to DNA acquisition, remains unexplored.

Here, we characterize the arsenal of *cab* bacteriocins carried by *S. mitis* and provide evidence

that some specifically mediate lysis of *S. pneumoniae*. We also show that pneumococcal DNA can

be efficiently taken up and recombined by *S. mitis* and that Cab-mediated lysis accelerates gene

transfer from *S. pneumoniae* to *S. mitis*. Together, these findings shed light on previous observations

of pneumococcal DNA integration in *S. mitis* genomes.

Results

***S. mitis* encodes a diverse arsenal of competence-associated bacteriocins (*cab*)**

We previously identified and characterized a limited number of *S. mitis* strains encoding multiple

competence-associated bacteriocins (*cab*), most of which lacked the pneumococcal bacteriocins

cibAB and associated immunity protein *cibC* (27). To investigate the broader distribution of *cab* and

cibAB in *S. mitis*, we surveyed publicly available genomes, leveraging the conserved genomic context of *cab* clusters – located upstream of *comAB* and downstream of *plsX-acpP*. This region was identified and extracted from 253 of 332 *S. mitis* genomes (Fig. 1A and Table S1). The remaining genomes had contig breaks and were excluded from further analysis.

Among the 253 genomes analyzed, 26.9% harbored *cibABC*, a striking contrast with *S. pneumoniae*, where these genes are universally present. An additional 13.0% of strains carried *cibC* but not *cibAB* – often referred to as "cheater" phenotype (28), in which strains are immune to CibAB-mediated killing but do not produce the bacteriocins themselves. In contrast, 47.4% of *S. mitis* encoded between one to eleven unique *cab* genes that contain a double glycine leader sequence ((M/L/V)XXXXGG), which is recognized by ComAB (29). We defined *cab* bacteriocins as all non-*cib* bacteriocin genes that were found in this particular location. In total, 95 distinct Cab variants were found, which grouped into 28 clusters based on sequence similarity (Fig. 1B and Fig. S1). Overall, 61.3% of *S. mitis* genomes carried at least one *cab* gene or the *cibABC* operon (Table S1).

Cab variants were present at low frequencies (<10%) in the *S. mitis* population, with 32 of 95 bacteriocin variants only being found in a single strain (Fig. 1B). Despite this extensive diversity, rarefaction analysis suggested that Cab diversity is not yet saturated (Fig. 1C); sequencing more *S. mitis* will likely reveal new Cab sequences.

These findings highlight *S. mitis* as a reservoir of highly diverse and largely unexplored competence-associated bacteriocins.

Cab bacteriocins are expressed in the late phase of competence

S. mitis G22, previously described by us, encodes a *cab* operon consisting of two putative bacteriocins and an associated immunity protein (hereafter referred as *cabAB* and *cabC*, respectively; *cabA* belongs to cluster 7 and *cabB* belongs to cluster 21) (Fig. 2A and Fig. S1) (27). This strain was selected to study *cab* regulation in *S. mitis* and its role in competitive behaviors. Both bacteriocins and the immunity protein it encodes are rare (30) and thus unlikely to encounter pre-existing immunity in other strains. Notably, its *cab* promoter contains five putative binding sites for the alternative sigma

factors (SigX boxes) (27), suggesting that *cabABC* may be expressed during the late phase of competence.

To investigate the dynamics of *cab* expression, the native *cabABC* promoter was ectopically inserted into the *S. mitis* G22 genome, upstream of a firefly luciferase reporter (*luc*). Expression of *cabABC* was then inferred by expression of *luc* and compared to known early (*comCDE*) and late (*sigX1* and *sigX2*) competence regulators. As expected, we observed that competence was not induced under acidic pH (Fig. S2), unless exogenous cognate competence-stimulating peptide (CSP$_{G22}$) was supplied (Fig. S3A). Upon CSP$_{G22}$ treatment, *cabABC* expression followed a similar temporal pattern to *comCDE*, *sigX1* and *sigX2*, peaking at approximately 70 min post-induction (Fig. S3B). Thus, based on expression kinetics alone, early and late competence genes could not be distinguished in this assay. This response was specific to CSP, as exogenous addition of cognate bacteriocin-like peptide pheromones (BlpC$_1$ and BlpC$_4$), encoded elsewhere in the *S. mitis* G22 genome (30), failed to induce *cabABC* expression (Fig. S4).

Deletion of the early regulator *comE* abolished *cabABC* expression, a result expected for both early- and late-regulated genes (Fig. 2B). Individual deletion of *sigX1* or *sigX2* partially reduced expression of *cabABC*, while deletion of both resulted in complete loss of *cabABC* expression, consistent with *cab* being expressed during the late phase.

Although multiple SigX boxes have been observed in *cab* promoters previously (16, 27), their functional relevance has not been tested. To assess this, we compared expression driven by the full-length *cabABC* promoter (containing all five SigX boxes) to a truncated version containing only the most proximal SigX box. Deletion of four out of five SigX boxes reduced *cabABC* expression (Fig. 2C), indicating that tandem SigX binding sites enhance *cab* expression. In the *S. mitis* collection, all *cab* loci harbored at least two SigX boxes, with one strain containing up to 38 boxes (Fig. 2D and Table S1). These boxes were often spread throughout *cab* loci, and larger loci were associated with higher number of SigX boxes, suggesting accumulation of boxes throughout these sequences.

Together, these results demonstrate that *cab* bacteriocins are specifically expressed during the late phase of competence, and that the presence of multiple SigX binding sites likely amplifies their expression.

Expression of *S. mitis cabABC* is induced by pneumococcal CSP-1

Mitis group streptococci naturally produce and recognize different CSP variants, known as pherotypes (31). To investigate the specificity of *cab* expression, synthetic CSP variants from closely related strains were tested for their ability to activate *cabABC*. CSP variants from three other *S. mitis*, one *S. oralis*, and one *S. pneumoniae* strain failed to induce *cab* expression in *S. mitis* G22 (Fig. 3A). By contrast, the most prevalent *S. pneumoniae* pheromone, CSP-1 (32), efficiently induced *cab* to the same levels as native CSP$_{G22}$.

Comparison of the amino acid sequences of CSP$_{G22}$ and the other CSP variants revealed a single amino acid residue, Q14 (glutamine), that may explain the differences in induction. While both CSP$_{G22}$ and CSP-1 contain Q14, a polar residue, the non-inducing variants contain either a L14 (leucine) or a P14 (proline), both nonpolar. To investigate the importance of Q14 and polarity in competence induction, CSP$_{G22}$ variants with single amino acid substitutions at position 14 were tested. Substituting Q14 with L significantly reduced *cab* expression, whereas replacing it with another polar residue (N, asparagine) restored expression (Fig. 3B). This suggests that polarity of amino acid 14 is important for CSP recognition by *S. mitis* G22. However, introducing Q14 into a non-inducing peptide (CSP-2; L14Q) was not sufficient to convert it into a functional inducer, indicating that additional residues beyond Q14 are required for efficient recognition by the membrane-bound histidine kinase ComD of *S. mitis* G22.

To assess whether *S. pneumoniae* can also respond to CSP$_{G22}$, the *comCDE* promoter from *S. pneumoniae* D39, a strain that produces and recognizes CSP-1, was transcriptionally fused to a *luc* reporter and integrated in its chromosome. Competence in D39 was efficiently induced by both CSP-1 and CSP$_{G22}$, confirming that *S. mitis* G22 and *S. pneumoniae* D39 can cross-communicate through distinct CSP peptides (Fig. 3C). Similar to G22, *S. pneumoniae* D39 was also affected by mutations at position 14: Q14L (polar to nonpolar) reduced induction. In contrast to *S. mitis* G22, the substitution Q14N (polar with another polar amino acid), failed to restore it. Notably, the substitution L14Q into CSP-2 (the second most frequent CSP in pneumococci) enabled competence induction.

These findings highligh the importance of Q14 in CSP recognition of pneumococci of the CSP-1

pherotype.

In the 253 *S. mitis* genomes analyzed, 2.8% encode CSP_{G22} (Table S1) indicating that more

S. mitis strains can induce competence in pneumococci of pherotype CSP-1.

These results reveal that cross-communications via CSP is possible between *S. mitis* and *S.*

pneumoniae, and that residue 14 plays a critical, but context-dependent, role in pheromone

recognition.

CabAB mediates inhibition of *S. pneumoniae* in a contact-dependent manner

Given that *S. mitis* G22 and *S. pneumoniae* D39 can cross-communicate through CSP, we

hypothesized both species would engage in competence-induced predation. Supporting this, *S. mitis*

G22 harbors the orphan immunity gene *cibC* (but not *cibAB*), suggesting it may be resistant to

pneumococcal CibAB attack (Table S1).

To investigate the specific role of the competence associated bacteriocins CabAB in *S.*

pneumoniae inhibition, we first deleted two bacteriocin clusters (*blp4* and *blp1*) with known anti-

pneumococcal activity (27), from the genome of *S. mitis* G22, as their activity could obscure CabAB-

mediated effects. All experiments in this section were, thus, performed using *S. mitis* G22 Δ*blp4*Δ*blp1*

and its derivatives as described below.

We grew dual-species biofilms of *S. mitis* G22 Δ*blp4*Δ*blp1* and *S. pneumoniae* D39 under

competence-permissive conditions. After 24 h, both species persisted in the biofilm (Fig. 4A and Fig.

S5). Deletion of *cab* from *S. mitis* led to increased pneumococcal loads, while constitutive expression

of *cab* via plasmid p(*cab*+) (overexpressor mutant contains the native *cab* copy and carries an

additional constitutively expressed copy in plasmid p(*cab*+)) reduced them, indicating that the

presence and expression levels of *cab* modulate competitive outcomes.

Bacteriocins are typically secreted to the extracellular environment (33). To test whether

CabAB exerts its activity as a secreted factor, we harvested cell-free supernatants (CFS) from *S. mitis*

strains and added them to pneumococcal cultures. CFS of *S. mitis* natively or overexpressing *cab* did

not inhibit *S. pneumoniae* growth (Fig. 4B), suggesting CabAB is not released into the medium. This

aligns with recent reports of bacteriocins, such as Listeriolysin S and contact-dependent inhibition by

glycine zipper proteins CdzCD (34, 35), that remain attached to the producer cell surface. In these

cases, inhibitory activity occurs via direct contact between producer and target cells.

To test whether CabAB could remain cell-associated, we heat-killed *S. mitis* strains (as

bacteriocins are typically resistant to high temperature treatments) and exposed them to *S.*

pneumoniae. Heat-killed *cab*[+] and p(*cab*[+]) *S. mitis* extracts significantly inhibited pneumococcal

growth, while extracts from Δ*cab S. mitis* did not (Fig. 4B), supporting the idea that CabAB is retained

in producer cells.

We then tested whether direct contact between *S. mitis* and *S. pneumoniae* was required for

inhibition. Dual-species biofilms were grown either with or without a 0.4 μm transwell membrane

separating both species. When the strains were in direct contact with each other, *cab* expression was

associated with pneumococcal inhibition, whereas no inhibition was observed when both species

were separated (Fig. 4C). These effects were not due to differences in *S. mitis* growth, as bacterial

loads were consistent across conditions and genotypes (Fig. 4D).

We then sought to detect CabA and CabB on the surface of competent *S. mitis* cells. A mutant

lacking the native *cab* locus was used, into which we ectopically integrated cassettes containing

haemagglutinin-tagged (HA) CabA (*cabA-HA cabBC*) or CabB (*cabA cabB-HA cabC*), under native

regulation. Immunofluorescence microscopy of live *S. mitis* expressing *cabA-HA* using anti-HA

antibodies revealed discrete clusters of CabA on the cell surface (Fig. 4E). Of note, CabA was not

detectable in all cells. This may be due to several factors, such as CabA being occluded by the cell

wall or competence development being asynchronous, with different subpopulations of *S. mitis*

becoming competent at slightly different times. No signal was detected in strains lacking the HA

epitope, confirming specificity. Interestingly, no signal was observed in the *cabB-HA* strain,

suggesting that CabB may be less surface-exposed, that the N-terminal HA tag was not accessible

to antibody binding, or that tagging interfered with localization.

Together, these results demonstrate that *S. mitis* inhibits *S. pneumoniae* via CabAB in a

contact-dependent manner.

CabAB is a two-peptide bacteriocin with LytA-dependent bactericidal activity

CibAB has previously been shown to be a two-peptide bacteriocin, meaning its inhibitory activity occurs only when both peptide components are present (8). To test if CabAB also functions as a two-peptide bacteriocin, synthetic CabA and CabB were tested individually and in combination against *S. pneumoniae* D39. No effect on pneumococcal growth was observed upon addition of DMSO (the solvent used to resuspend the peptides), CabA alone, or CabB alone (Fig. 5A). In contrast, a single treatment with equimolar amounts of CabA and CabB delayed *S. pneumoniae* growth for ~12 h, indicating that both peptides are required for activity. This is consistent with the presence of multiple GxxxG or GxxxG-like motifs in both bacteriocins (Fig. 5B), known to facilitate and stabilize two-component bacteriocins (36-38). We also confirmed that CabAB can cross the membrane used in the transwell assays, leading to a decrease in *S. pneumoniae* loads when added to opposite compartments (Fig. S6).

We next investigated whether CabAB exerts bactericidal activity. Exponentially growing *S. pneumoniae* D39 cultures were treated with CabAB or DMSO. CabAB-treated cultures showed evident decrease in optical density, indicating lysis, while DMSO-treated controls did not (Fig. 5C). Lysis induced by the related bacteriocin CibAB has been shown to depend on the action of the pneumococcal autolysins LytA and LytC (8). Here, deletion of *lytC* from D39 had no effect on CabAB-mediated lysis, whereas a Δ*lytA* mutant exhibited delayed growth but no decrease in optical density, indicating that LytA plays a key role in CabAB-induced lysis. This does not exclude, however, the possibility that limited lysis occurs concomitantly with growth.

Together, these findings show that *S. mitis* CabAB is a two-peptide bacteriocin with bactericidal activity against *S. pneumoniae*, and that its lytic activity depends on the target cell LytA autolysin.

CabAB affects membrane homeostasis in *S. pneumoniae*

We sought to shed light on how CabAB affects pneumococcal physiology. To this end, *S. pneumoniae* D39 cells were treated with CabAB or DMSO and imaged by scanning electron microscopy (SEM).

While DMSO-treated cultures displayed normal morphology, CabAB-treated cells appeared to be

encased in an extracellular matrix (Fig. 6A), suggesting leakage of intracellular material.

To test if membrane permeability was compromised, cells were stained with propidium iodide

(PI), a DNA-binding dye that only enters cells with compromised membranes. In DMSO-treated

cultures, only 3% were PI-positive, while CabAB-treated cultures contained 85% PI-positive cells (Fig.

6B). Doubling the concentration of CabAB further increased this proportion to 97%, indicating this

phenotype is concentration dependent. Deletion of *lytA* had no effect on these phenotypes,

demonstrating that membrane permeabilization occurs independently of autolysin-mediated lysis (Fig.

S7).

Time series measurements showed that PI signal intensity peaked at 30 min post CabAB

exposure, still being detected after 120 min (Fig. 6C). We also assessed pneumococcal membrane

depolarization using the fluorescent dye $DiBAC_4(3)$, which enters cells only when their membranes

are depolarized. Compared to DMSO, CabAB treatment led to a rapid and sustained increase in

$DiBAC_4(3)$ signal (Fig. 6D), indicating membrane depolarization.

Since some characterized two-peptide bacteriocins act by forming pores (39-42), we tested

whether CabAB treatment causes efflux of intracellular components from target cells. In wild-type *S.*

pneumoniae treated with CabAB, 80% of total ATP was detected in the supernatant (Fig. 6E), in

agreement with our previous observation of significant cell lysis (Fig. 5C). Interestingly, in CabAB-

treated Δ*lytA S. pneumoniae* cells, no decrease in optical density was observed over the course of

the experiment, yet 30% of ATP was still detected in the supernatant, suggesting that CabAB induces

leakage independently of cell lysis.

We next visualized CabAB-induced damages using transmission electron microscopy (TEM).

DMSO-treated cells appeared intact, whereas 17% of CabAB-treated cells contained large

intracellular aggregates (Fig. 6F). Based on our prior observations, we hypothesized these

aggregates would be membranous. Consistently, staining with the membrane dye FM4-64 revealed

intracellular foci in CabAB-treated cells, but not in control cells (Fig. 6G), confirming the formation of

membrane aggregates by CabAB.

Finally, lower-magnification fluorescent microscopy of CabAB-treated cultures stained with

DAPI and FM4-64 revealed frequent clumps of cells encased in nucleic acid and membrane material,

mirroring the SEM observations of surviving cells remaining embedded in an extracellular matrix (Fig.

S8)

Altogether, these findings show that *S. mitis* CabAB disrupts membrane homeostasis in *S.*

pneumoniae by permeabilizing and depolarizing membranes, leading to release of its intracellular

content.

CabAB facilitates the acquisition of pneumococcal DNA by *S. mitis* in biofilms

As previously shown, CabAB promotes release of DNA from *S. pneumoniae* cultures. Since *cab*

expression is synchronized with the competence period, we hypothesized that CabAB could enhance

DNA acquisition and HGT within biofilms. To test this, we grew dual-species biofilms under conditions

designed to study HGT (Fig. 7A), where: (i) DNA transfer is unidirectional, since the DNA donor strain

(*S. pneumoniae* D39) carries a deletion of the major competence pilus subunit gene, *comGC,*

rendering it nontransformable; (ii) the *S. pneumoniae* DNA donor harbors chloramphenicol (Chl^R) and

tetracycline (Tet^R) resistance cassettes integrated at opposite regions of the chromosome with high

sequence homology (89-98%) between DNA donor and recipient (Fig. S9); (iii) the DNA recipients (*S.*

mitis G22, wt and variants) are kanamycin resistant, allowing for discrimination between non-

transformed (Kan^R) and transformed *S. mitis* (Kan^R Chl^R, Kan^R Tet^R, and Kan^R Chl^R Tet^R); (iv) biofilms

are pre-grown under acidic pH for 24 h to promote biofilm establishment without inducing competence;

and (v) CSP$_{G22}$ is added after 24 h to transiently induce competence. A noncompetent *S. mitis* Δ*comE*

mutant was used as a negative control, as it is unable to acquire Chl^R and/or Tet^R by transformation.

These experiments were conducted using the *S. mitis* G22 Δ*blp4* Δ*blp1* background.

To rule out the possibility that CabAB affects transformability rather than DNA availability, we

first replaced the donor strain with purified gDNA. Under these conditions, both *cab*$^+$ and Δ*cab S. mitis*

acquired resistance at similar frequencies (Fig. 7B), indicating that native *cab* expression does not

impact transformability. In contrast, overexpression of *cab* reduced transformation efficiency for one

antibiotic resistance, possibly due to the fitness burden associated with carrying p(*cab*$^+$).

In dual-species biofilms, *cab*$^+$ *S. mitis* efficiently acquired Chl^R and Tet^R from the DNA donor,

at a significantly higher frequency than the Δ*cab* mutant (Fig. 7C), indicating that CabAB contributes

to DNA acquisition. In contrast, overexpression of CabAB did not enhance transformation efficiency,

suggesting that a precise *cab* regulation might be necessary for optimal transformation. We extended

these findings to an additional *S. mitis* strain, C22, which harbors a distinct *cab* bacteriocin repertoire

(27). *S. mitis* C22 contains a *cab* cluster encoding two copies of the bacteriocin *cabD* (cluster 10), two

copies of the bacteriocin *cabE* (cluster 26), and two putative immunity proteins, *cabF* and *cabG* (Fig.

S10A). Wild-type C22 was able to acquire Chl^R and Tet^R from the *S. pneumoniae* DNA donor and

transformation frequencies were also reduced in the Δ*cab* mutant (Fig. S10B). Consistently, the

proportion of *S. pneumoniae* in dual-species biofilms with *S. mitis* C22 was lower when co-cultured

with C22 *cab*$^+$ than with Δ*cab* cells (Fig. S10C), showing that the cab locus of a different *S. mitis* strain

also mediates inhibition of *S. pneumoniae* and facilitates genetic exchange.

We then sought to investigate the specificity of CabAB-induced lysis. Synthetic CabAB was

tested against one representative strain of 26 bacterial species, including 18 streptococcal species,

four additional non-streptococcal Gram-positive and four Gram-negative species. All selected species

are human commensals or pathogens that can inhabit the oral and/or respiratory tract and could

theoretically encounter *S. mitis* and be exposed to Cab bacteriocins in natural environments. Addition

of 0.975 µM of CabAB to exponential cultures of these bacteria stalled the growth of some species

(Fig. 7D and Fig. S11 and S12). The most sensitive to CabAB were the representative strains of

Pyogenic Group streptococci (*S. pyogenes* and *S. agalactiae*). Gram-negative bacteria (*Haemophilus*

influenzae, *Klebsiella pneumoniae*, *Moraxella catarrhalis*, and *Neisseria lactamica*) were largely

unaffected. Despite growth inhibition in some species, lysis was only observed in two - *S. pneumoniae*

and *S. parasanguinis* - suggesting that CabAB-mediated lysis has limited specificity.

Finally, we investigated whether membrane aggregate formation was specifically associated

with species that undergo CabAB-mediated lysis. Aggregate formation was observed exclusively in

S. pneumoniae and not in *S. parasanguinis*, *S. pyogenes* and *S. agalactiae* (Fig. S13). Notably,

membrane aggregates were also absent in Δ*lytA S. pneumoniae* cells, indicating that aggregate

formation is specific to *S. pneumoniae* and that it requires LytA.

These results suggest that CabAB provides a competitive advantage by inhibiting the growth

of some bacteria, but that it also enhances DNA release in biofilms by selectively lysing a few closely

related species during competence, notably *S. pneumoniae*, thereby promoting gene exchange.

Discussion

During competence, streptococci scavenge extracellular DNA. In *S. pneumoniae*, where competence-

induced predation is best characterized, the coordinated action of CibAB, LytA, LytC, and the murein

hydrolase CbpD leads to lysis of noncompetent siblings (8). This process promotes DNA release and

facilitates HGT (12, 43, 44).

Although extensive genomic data suggests that *S. mitis*, a close relative of *S. pneumoniae*,

frequently acquires *S. pneumoniae* DNA (18, 19, 21-24, 45), the molecular mechanisms underlying

this exchange remained poorly understood. Understanding these mechanisms is relevant because

capsule acquisition by *S. mitis* can complicate DNA-based pneumococcal serotyping. In addition, *S.*

mitis has been associated with cases of infective endocarditis (23), highlighting the value of studying

how it acquires potentially pathogenic traits.

Here, we investigated the role of *S. mitis* competence-associated bacteriocins, CabAB, in

mediating interspecies interactions (Fig. 8). We show that CabAB is a two-peptide bacteriocin

produced during the late phase of competence, when cells are primed for DNA uptake and

recombination, capable of lysing *S. pneumoniae* in a LytA-dependent manner. Although LytA is

constitutively expressed throughout growth, its concentration increases during competence, owing to

a genomic architecture comprising three constitutive promoters and one competence-induced

promoter (7, 46-48). In this context, it is tempting to speculate that by inducing *S. pneumoniae*

competence via CSP crosstalk, and therefore increasing LytA concentration, some *S. mitis* could

further sensitize *S. pneumoniae* to Cab bacteriocins. Nonetheless, CabAB-induced lysis appears to

be independent of pherotype of target cells, as both *S. pneumoniae* D39 (CSP-1; used in most

experiments) and TIGR4 (CSP-2; included in the panel of 26 bacterial species) were lysed. These

pherotypes account for the vast majority (>95%) of pneumococcal strains (49-51).

Of the initial 332 *S. mitis* genomes analyzed, we were not able to retrieve the region upstream

of *comAB* from a substantial fraction of strains (23.8%), as their assemblies showed contig breaks

precisely at this locus. We propose this difficulty arises from the repetitive nature of the region, as

indicated by the presence of numerous SigX boxes. The number of boxes scaled linearly with the size

of *cab* loci, suggesting that SigX boxes have accumulated throughout these sequences. Repeats are

known to both promote and result from genomic rearrangements (52, 53). They are often remnants

of previous recombination events, with no known biological function (54). In this case, however,

accumulation of SigX boxes may provide a selective advantage by enhancing bacteriocin expression,

which could explain why they are enriched in this region. Consistent with this, the extensive genetic

rearrangements observed across multiple streptococcal species at this locus also suggests this is a

highly recombinogenic region (25, 26, 55). To the best of our knowledge, the *cab* locus appears to be

the only competence-induced locus in *S. mitis* whose promoters contain multiple SigX boxes; other

competence-induced genes typically harbor a single box (16).

Microscopic analysis revealed that CabAB-treated pneumococci often contain large

membrane-derived intracellular aggregates. Whether these structures are part of CabAB's

mechanism of action or represent a cellular stress response remains unclear. Intracellular vesicle

formation has been shown to be essential for the lytic activity of bacteriocin A37 (56). Consistently,

we have observed that deletion of LytA, required for CabAB-mediated lysis in *S. pneumoniae*, blocks

formation of intracellular aggregates.

We further demonstrate that CabAB-mediated inhibition of *S. pneumoniae* requires direct

contact between producer and target cells. Contact-dependent bacteriocin delivery has been

described in *Caulobacter crescentus* and *Listeria monocytogenes* (34, 35). In *L. monocytogenes*,

bacteriocins were only detected on the surface of producer cells following cell wall digestion. In

contrast, we detected CabAB on live *S. mitis* with intact cell walls, suggesting a distinct delivery

mechanism. Although CabAB activity strictly required both peptides, only CabA could be detected on

the surface of competent *S. mitis*. CabB was not detectable by immunofluorescence, which could reflect limited surface exposure, inaccessibility of the HA tag, or altered localization caused by C-terminal tagging. A similar pattern of differential detectability has been reported in the CdzC/D contact-dependent bacteriocin system in *Caulobacter crescentus*: both CdzC and CdzD are cell-associated and required for activity, but only CdzC is clearly visualized in surface aggregates, whereas CdzD is not direcly observed on the surface (35). These observations suggest that, in two-peptide contact-dependent killing systems, one peptide may be more readily surface-exposed. Contact-dependent killing during competence is not unique to streptococci, despite differences in killing factors. *Vibrio cholerae* produces a competence-induced type VI secretion system that kills nonimmune cells, releasing DNA that becomes accessible for HGT (57). Contact dependence may be particularly advantageous in biofilm contexts, a niche where extracellular DNA could be subject to rapid degradation nucleases (58). Proximity within multispecies biofilms could enhance transformation efficiency by increasing the local DNA concentration, given that DNA abundance and size influence recombination success (59). Notably, biofilm conditions and cell-to-cell contact have been shown to promote both the frequency and size of recombination events in pneumococci, respectively (60, 61). Notably, the frequency of double transformants was higher under cocultivation conditions (*S. mitis* and *S. pneumoniae*) than when genomic DNA was added to *S. mitis* biofilms. In coculture, DNA is likely released continuously over the course of incubation, increasing its local concentration and availability, which may favor sequencial uptake and recombination events. In contrast, purified gDNA was added only once and could be subject to degradation by extracellular nucleases, thereby reducing the probability of independent recombination events. Strikingly, among 26 tested species, only *S. pneumoniae* and *S. parasanguinis* were lysed by CabAB, suggesting a narrow-spectrum bactericidal activity. Whether this specificity is unique to CabAB or shared among other Cab bacteriocins remains to be determined. Nonetheless, the ability to selectively target phylogenetically related but distinct species suggests a potential trade-off between maximizing recombination efficiency and accessing a broader gene pool. While high sequence identity between donor and recipient promotes recombination (62), restricting lysis to closely related species could expand the range of accessible alleles beyond conspecifics (*i.e.* beyond members of the same species). Thus,

the large diversity of *cab* sequences in *S. mitis*, in contrast to *S. pneumoniae*, which typically harbors

only *cibAB*, may represent one of several evolutionary forces contributing to the exceptional genetic

diversity observed in *S. mitis* (23, 63).

Importantly, CabAB also strongly inhibited the growth of other streptococcal pathogens,

including *S. pyogenes* and *S. agalactiae*, both of which are listed as priority pathogens by the World

Health Organization (30). This potent inhibitory activity suggests that the Cab arsenal carried by *S.*

mitis could be a promising source of antimicrobial agents against important clinical pathogens as

recently suggested by us (27).

In summary, our findings uncover a contact-dependent mechanism of bacteriocin delivery in

S. mitis that selectively targets closely related streptococci including *S. pneumoniae*. This process

promotes HGT and likely contributed to competitive fitness and evolutionary plasticity of streptococcal

populations.

Materials and Methods

Bacterial strains and culture conditions

Bacterial strains, oligonucleotides, plasmids and peptides used in the study are listed in Table S2.

Streptococci were routinely grown in C+Y$_{YB}$ (64) (pH 7.4, unless otherwise specified) and plated on

tryptic soy agar with 5% defibrinated sheep blood (blood agar, BA). When appropriate, media was

supplemented with antibiotics at the following concentrations: 150 µg.mL^{-1} kanamycin, 200 µg.mL^{-1}

spectinomycin, 4 µg.mL^{-1} chloramphenicol, or 4 µg.mL^{-1} tetracycline.

Staphylococcus aureus, Staphylococcus epidermidis and *Moraxella catarrhalis* were grown in

tryptic soy broth (TSB); *Corynebacterium accolens* was grown in TSB with 0.5% Tween 80; *Klebsiella*

pneumoniae and *Escherichia coli* were grown in lysogeny broth; and *Neisseria lactamica*,

Haemophilus influenzae and *Dolosigranulum pigrum* were grown in C+Y$_{YB}$ with 20 µg.mL^{-1} hemin and

20 µg.mL^{-1} NAD. For propagation of plasmids in *E. coli,* media was supplemented with 100 µg.mL^{-1}

kanamycin or 100 µg.mL^{-1} spectinomycin, as required.

Identification of *cab* bacteriocins and *cibABC* in *S. mitis*

Publicly available *S. mitis* genomes (n = 322) were downloaded from PubMLST. An additional 10 genomes from reference or commonly used *S. mitis* strains were included, resulting in a total of 332 genomes (Table S1). To detect *cab* clusters, nucleotide sequences of *plsX-acpP* and *comAB* (from *S. mitis* B6; GenBank Accession GCA_000027165.1), which flank the *cab* locus, were aligned against all genomes using pairwise sequence alignment (Biopython v1.81; match = 2; mismatch = -1; gap open = -5; gap extension = -0.5; minimum overall sequence identity $\geq$ 90%). Genomes in which *plsX-acpP* and *comAB* were absent or located on separate contigs were excluded from further analysis. This region was identified in 253 genomes.

To identify mature Cab bacteriocin sequences, extracted sequences were scanned for open reading frames (ORFs) of 100-300 nucleotides, beginning with an ATG start codon and ending with a TAA, TAG or TGA stop codon. Only ORFs whose translated sequences contained a (M/L/V)XXXXGG double-glycine cleavage motif were retained. Mature Cab bacteriocins were obtained by removing the N-terminal signal sequence upstream of the cleavage site. In this work, we defined *cab* bacteriocins as all non-*cib* bacteriocin genes that were found in this particular location. To explore diversity of Cab bacteriocins, sequences were converted into amino acid frequency vectors. Pairwise dissimilarities between sequences were calculated using the Bray-Curtis metric (ScyPy), followed by agglomerative hierarchical clustering (scikit-learn). To determine the optimal number of clusters, silhouette scores (scikit-learn) were computed for cluster numbers ranging from 2 to 50. The number of clusters yielding the highest score was selected as optimal.

To assess the prevalence of *cibABC* in *S. mitis*, the nucleotide sequence of *cibABC* (from *S. pneumoniae* D39, GenBank Accession NC_008533.2) was aligned against all *S. mitis* genomes using NCBI BLASTn. Genomes with no detectable hit were considered *cibABC*-negative. For genomes with partial matches (<100% query coverage), the locus was manually inspected to verify the presence or absence of *cibA*, *cibB*, and *cibC* individually – a common occurrence in strains harboring *cibC* but not *cibAB*.

To find SigX boxes, the consensus sequence (65) was searched within *cab* loci using Find Individual Motif Occurrences (FIMO) v5.5.8 (66). Hits with *p*-value <0.001 were considered true SigX

boxes. The detection of Cab bacteriocins and putative SigX boxes was benchmarked against six *S.*

mitis strains with complete genomes and manually curated *cab* loci (27, 67).

Phylogeny of *S. mitis*

Core genome single-nucleotide polymorphisms (SNPs) were called using Snippy v4.6.0

(https://github.com/tseemann/snippy) by aligning *S. mitis* genomes to the *S. mitis* B6 reference

genome. The resulting alignment was used to construct a maximum-likelihood phylogenetic tree with

FastTree v2.2 (68). The phylogeny was visualized and annotated using the Interactive Tree of Life

(iTOL) platform (69).

Synthetic peptides

Peptides were produced by solid-phase peptide synthesis with >95% purity (NZYTech) (Table S2)

(27, 49-51). CSP and $BlpC_1$ pheromones were resuspended in sterile ddH_2O; $BlpC_4$ was resuspended

in 10% acetic acid; and CabA and CabB were resuspended in DMSO. Peptides were serially diluted

in sterile ddH_2O and stored at -20ºC.

Transformation

For transformation of *S. mitis* and *S. pneumoniae*, cells were grown to an OD_{600} of 0.5, diluted 1:100

in pre-warmed $C+Y_{YB}$ and grown until an OD_{600} of 0.04-0.1. Cells were treated with 1 µM of cognate

CSP and transformed with 0.4 µg linear DNA, 1 µg plasmid DNA, or 10 µg chromosomal DNA.

Transformants were recovered for 4 h and plated in selective medium. Linear, plasmid and genomic

DNA were purified using the Zymoclean Gel DNA Recovery Kit (Zymo Research), ZymoPURE

Plasmid Miniprep Kit (Zymo Research) and EZ1 DSP Virus Kit (Qiagen), respectively.

For heat-shock transformation of *E. coli* DH10B or DH5α, $CaCl_2$-competent cells were mixed

with 10% (v/v) of ligation reactions and incubated on ice for 15 min. Cells were then heat-shocked at

42ºC for 40 s, immediately chilled on ice, and recovered with 900 µL of SOC (2% tryptone, 0.5% yeast

extract, 10 mM NaCl, 2.5 mM KCl, 10 mM $MgCl_2$, 10 mM $MgSO_4$ and 20 mM glucose; pH 7) for 1 h

before plating in selective medium.

Construction of a fluorescence-luminescence transcriptional reporter

A DNA cassette containing a *hlpA-mNeonGreen* fusion and plasmid pBIR, which integrates in the genome of *S. mitis* G22 and *S. pneumoniae* D39, were synthesized (Twist Bioscience). The reporter cassette and plasmid were amplified and blunt end ligated to generate plasmid pJB10. The firefly luciferase *luc* (from pPEP-LGZ) and plasmid pJB10 were amplified and blunt-end ligated to create plasmid pJB11 encoding a dual fluorescence-luminescence reporter. Transcriptional fusions were created by digesting both pJB11 and promoter fragments with BsaI, followed by ligation. Variants of pJB11 were linearized with BamHI and transformed into *S. mitis* G22 or *S. pneumoniae* D39. Transformants were selected in BA with chloramphenicol.

Expression of *cabABC* and other competence genes

S. mitis and *S. pneumoniae* strains harboring transcriptional fusions of competence promoters were grown in C+Y$_{YB}$ at pH 6.6 (to prevent natural competence induction) until an OD$_{600}$ of 0.5. To evaluate natural competence dynamics, acidic precultures were diluted 1:100 in C+Y$_{YB}$ (pH 6.6, 7.4 or 8.2) containing 340 µg/mL D-luciferin (Abcam). To assess the effect of different pheromones on competence induction, acidic precultures were diluted to OD$_{600}$ = 0.1 in C+Y$_{YB}$ (pH 6.6) with 340 µg/mL D-luciferin. Then, CSP or BlpC variants (up to 1 µM), were added. Cultures were transferred to white 96-well plates with clear bottoms (Nunc). The OD$_{595}$ and luminescence was recorded every 10 min for 24 h in a BioTek Neo2 Multimode Reader with a gain of 185. Luminescence values were normalized to the OD$_{595}$. D-luciferin was resuspended in sterile ddH$_2$O to 15 mg.mL^{-1} and kept at -80 ºC in single-use aliquots.

Construction of ComE, SigX1 and SigX2 deletion mutants

The upstream and downstream flanking regions of *comE, sigX1, sigX2* were amplified from *S. mitis* G22 gDNA. Kanamycin and spectinomycin resistance genes were amplified from pKan and pPEP-LGZ (70), respectively. DNAs were ligated using Gibson Assembly (New England Biolabs) and

534 amplified with nested primers. Transformants were selected in BA with kanamycin (for *comE* and

535 *sigX1*) or spectinomycin (for *sigX2*).

Biofilms and transwells

S. mitis and *S. pneumoniae* were grown to exponential phase and mixed at a 1:1 ratio (10^5 CFU of

each strain) in C+Y$_{YB}$ with 1600 U.mL^{-1} catalase. Two milliliters of the mixtures were added to tissue-

culture treated 12- or 24-well plates (Corning) and incubated 24 h at 34ºC with 5% CO_2. Non-adherent

cells were removed by aspiration, and biofilms were thoroughly resuspended in PBS. Initial biofilms

were performed in 24-well plates, while the remaining were performed in 12-well plates (since these

were the ones where we could fit a transwell insert).

For transwell biofilms, 1.5 mL of C+Y$_{YB}$ containing catalase and 10^5 CFU of *S. pneumoniae*

was added to wells of tissue-culture treated 12-well plates. Transwell inserts containing a 0.4 µm pore

polycarbonate membrane (Corning) were placed in the wells, and 0.5 mL of C+Y$_{YB}$ with catalase and

10^5 CFU of *S. mitis* were added to the upper insert. Biofilms were recovered from both compartments.

In dual-species biofilms, *S. pneumoniae* and *S. mitis* cells were enumerated by plating on BA

with chloramphenicol or kanamycin, respectively. In the transwell setup, where *S. pneumoniae* and

S. mitis were physically separated, biofilm suspensions from both compartments were also plated on

selective media to confirm that neither strain had crossed the membrane.

Construction of an *S. mitis cab* overexpressor mutant

554 The plasmid backbone was amplified from pDL278_P23-DsRed-Express2 (Addgene plasmid

555 #121505), and *cabABC* was amplified from *S. mitis* G22 gDNA. The two fragments were blunt-end

556 ligated, generating plasmid pJB09-*cabABC*, in which *cabABC* is placed under the control of the strong

557 constitutive P23 promoter. *S. mitis* transformants were selected in BA with spectinomycin.

Inhibitory activity of cell-free supernatants and heat-killed cells

S. mitis was grown to early stationary phase and centrifuged at 10,000 g for 30 min at 4ºC. Cell-free

supernatants (CFS) were obtained by filtration (0.22 µm). CFS were lyophilized by vacuum

evaporation using a SpeedVac concentrator (LabConco) and resuspended in ddH$_2$O to a

concentration of 10x before testing. C+Y$_{YB}$ medium was concentrated in parallel and used as a

negative control for inhibitory activity. Cell pellets were washed once in PBS and heat-killed for 15

min at 95 ºC. Complete lysis of pellet extracts was confirmed by plating in BA.

To test for inhibitory activity of CFS and heat-killed cells, *S. pneumoniae* D39 was grown to

an OD$_{600}$ of 0.5, diluted 1:100 in fresh medium, and 180 μL of the dilution was added to clear 96-well

plates. Twenty microliters of treatments or water was added to each well and growth was monitored

by measuring the OD$_{595}$ every 30 min for 24 h using a Tecan Infinite 200 Pro.

Sensitivity of bacteria to CabAB

Exponential-phase cultures of bacterial species (18 *Streptococcus* spp., four additional non-

streptococcal Gram-positive species, and four Gram-negative species) were diluted to OD$_{600}$ 0.1 and

195 μL was added to wells of clear 96-well plates. CabAB (0.975 μM) or DMSO (0.00975%) was

added to wells and growth was recorded by measuring the OD$_{595}$ every 10 min for 24 h using the

Tecan Infinite 200. Growth arrest was quantified by comparing the area under the curve (AUC)

between CabAB- and DMSO-treated cultures. All further assays using CabAB and DMSO treatments

were performed in the same conditions, unless specified otherwise.

Construction of autolysis-deficient *S. pneumoniae* mutants

The upstream and downstream regions of *lytA* and *lytC* were amplified from *S. pneumoniae* D39

gDNA. The PhunSweet cassette containing *pheS-sacB*-aph*A-3* which confers susceptibility to

chlorinated phenylalanine and sucrose, and resistance to kanamycin, was amplified from *S.

pneumoniae* D39 Δ*cps*::*PhunSweet* gDNA (71). DNAs were assembled using Gibson Assembly and

amplified with nested primers. *S. pneumoniae* D39 transformants were selected on BA with

kanamycin.

To remove the antibiotic resistance cassette, the flanking regions of *lytA* and *lytC* were

amplified, assembled with Gibson Assembly, and amplified with nested primers. Transformants were

recovered by counterselection on BA with 15 mM chlorinated phenylalanine and 10% sucrose.

Scanning electron microscopy (SEM)

S. pneumoniae D39 cultures (30 mL) were treated with DMSO or CabAB and incubated for 15 min at

37ºC. Cells were harvested by centrifugation (all centrifugations were carried out at 800 g for 10 min

at room temperature) and washed three times with PBS. Fixation was performed in 500 μL of 2.5%

glutaraldehyde and 1% paraformaldehyde in 0.1 M sodium cacodylate buffer for 30 min at room

temperature. Cells were washed three times with 0.1 M sodium cacodylate buffer and dehydrated in

increasingly higher ethanol concentrations (50%, 70%, 90%, 100%), by incubating 10 min in 500 μL

of each ethanol concentration. Ethanol was replaced with 500 μL tert-butyl alcohol and samples were

incubated at 30ºC for 1 h. Then, tert-butyl was removed and replaced with 500 μL of fresh tert-butyl

alcohol. Samples were kept at -20ºC overnight. Samples were then lyophilized under vacuum, gold-

sputtered using a Cressington 108 sputter coater, and imaged with a Hitachi SU-8010 scanning

electron microscope at 1.5kV.

Transmission electron microscopy (TEM)

S. pneumoniae D39 cultures (30 mL) were treated with DMSO or CabAB, incubated for 15 min at

37ºC, and recovered by centrifugation at 800 g. Cells were fixed with 2% formaldehyde, 2.5%

glutaraldehyde, 0.075% ruthenium red and 1.55 % lysine acetate in distilled water for 20 min at 4ºC.

Cells were washed twice in 0.075% ruthenium red, resuspended in 2% formaldehyde, 2.5%

glutaraldehyde and 0.075% ruthenium red, and incubated overnight at 4ºC. Cells were washed three

times with 0.075% ruthenium red and further processed at Instituto Gulbenkian de Ciência. Cells were

embedded in 2% low-melting point agarose and fixed with 1% osmium tetroxide and 0.06% ruthenium

red for 1 h at room temperature. Cells were washed once with 0.1 M sodium cacodylate buffer and

twice with distilled water. Dehydration was performed on ice using increasing concentrations of

ethanol (10%, 30%, 50%; 10 min each). Samples were stained overnight at 4ºC with 2% uranyl

acetate in 70% ethanol, followed by further dehydration with 90% ethanol and three times with 100%

ethanol. Infiltration was carried out with increasing concentrations of epoxy resin in ethanol (3:1, 1:1,

1:3), and 100% resin overnight, followed by two exchanges of 100% resin. Ultrathin sections were cut

and imaged on a Tecnai G2 Spirit BioTWIN Transmission Electron Microscope (FEI) equipped with

an Olympus-SIS Veleta CCD camera.

Large intracellular aggregates were observed in some CabAB-treated cells. To quantify these

structures, micrographs of CabAB- and DMSO-treated cells were blindly evaluated by two

researchers. Each researcher was first shown representative cells with and without aggregates, and

then independently counted the number of cells containing these structures. The final percentage of

cells with aggregates was calculated as the average between both researcher counts.

Permeabilization and depolarization of *S. pneumoniae* membranes

S. pneumoniae D39 cultures were treated with DMSO or CabAB. One-milliliter aliquots were taken at

0, 2, 5, 15, 30, 60, and 120 min post treatment and immediately chilled on ice. Cells were washed

twice with PBS and kept on ice. Cells were stained with 6 μM propidium iodide (Invitrogen) and 3 μM

DiBAC$_4$(3) (Enzo Life Sciences) for 5 min at room temperature, protected from light. Cells were

centrifuged, resuspended in PBS, and added to black 96-well plates with clear bottoms (Nunc).

Fluorescence intensity was measured using a BioTek Neo2 Multimode Reader under the following

settings: shake for 5 s; propidium iodide – top reading, 535 ± 12 nm excitation, 617 ± 12 nm emission

and 120 gain; DiBAC$_4$(3) – top reading, 492 ± 9 nm excitation, 518 ± 9 emission, 120 gain).

Fluorescent values were normalized to the OD$_{595}$.

Efflux of ATP

Cultures of *S. pneumoniae* were treated with DMSO or CabAB. Aliquots were taken at 0, 15 and 120

min post-treatment and kept on ice. Cells were centrifuged, and supernatants were filtered (0.22 μm)

and maintained on ice. Cells were permeabilized in DMSO for 10 min at room temperature. Both cells

and supernatants were diluted 1:10 in ddH$_2$O and kept on ice.

Extracellular and intracellular ATP was measured with the ATP Luminescent Cell Viability

Assay (Millipore) according to instructions. Luminescence was measured in a BioTek Neo2 Multimode

Reader using a gain of 150 and integration time of 10 s. An ATP standard curve confirmed linear

detection within the range of 0.2 μM to 2 mM ATP. Total ATP was calculated by summing

luminescence values from both the cell and supernatant fractions under each condition.

Construction of *S. mitis* harboring HA-tagged bacteriocins

DNAs containing HA (YPYDVPDYA)-tagged *cabA* and *cabB* were synthesized (Twist Bioscience).

The regions upstream (including the native promoter) and downstream (including a chloramphenicol

resistance cassette) of the *cab* operon were amplified from pJB11-*cabABC*. DNAs were assembled

by Gibson Assembly and amplified with nested primers. The resulting constructs were transformed

into the *S. mitis* Δ*cabABC* background, generating two mutant strains: *cabA-HA cabBC* and *cabA*

cab-HA cabC. Transformants were selected in BA with chloramphenicol.

Confocal microscopy and immunofluorescence

To quantify the percentage of cells that become membrane-permeable, *S. pneumoniae* was treated

with DMSO or CabAB. Cells were recovered by centrifugation (9600 g, 4ºC, 3 min) and washed three

times with PBS. Cells were stained with 30 µM propidium iodide and 5 µM SYTO 9 for 15 min at room

temperature, protected from light.

To determine whether large intracellular structures observed in CabAB-treated *S. pneumoniae*

cells corresponded to membranes, treated cells were washed three times with PBS and stained with

20 µg.mL^{-1} FM4-64 and 15 µg.mL^{-1} DAPI for 5 min at room temperature, protected from light.

For immunofluorescence of live competent bacteria, *S. mitis* harboring HA-tagged bacteriocins

were grown in acidic C+Y$_{YB}$ and diluted to OD$_{600}$ = 0.1 in pre-warmed C+Y$_{YB}$. CSP$_{G22}$ was added at 1

μM, and cells were incubated for 70 min at 37ºC and then chilled on ice. To retain the competent

status of cells, all subsequent steps were performed on ice using ice-cold solutions. Cells were

centrifuged (9600 g, 4ºC, 3 min) and washed twice in PBS. Blocking was performed in PBS with 5%

bovine serum albumin (BSA) for 1 h. Cells were pelleted and resuspended in 100 µL of primary rabbit

monoclonal anti-HA antibody (clone C29F4; Cell Signaling Technology) diluted 1:100 in PBS with 1%

BSA, and incubated for 1 h. Cells were washed twice in PBS with 0.1% Tween 20, resuspended in 100 µL of Alexa Fluor 488-conjugated goat anti-rabbit secondary antibody (Invitrogen) diluted 1:500 in PBS with 1% BSA, and incubated 1 h protected from light. Cells were washed twice with PBS containing 0.1% Tween 20 and once with PBS. Cells were then stained with 20 µg.mL^{-1} FM4-64 and 15 µg.mL^{-1} DAPI for 5 at room temperature, covered from light.

Immediately before imaging, cells were mounted on thin 1.7% agarose pads. Fluorescence imaging was performed using a Zeiss LSM 880 point scanning confocal microscope equipped with an Airyscan detector. Samples were imaged with a Plan-Apochromat 63x/1.4 Oil DIC M27 objective (Zeiss), using laser lines 405 nm (DAPI), 488 nm (SYTO 9 and Alexa Fluor 488), and 561 nm (FM4-64 and propidium iodide). Imaging was performed at laser intensities of 2-3% and a gain of 700 for DAPI, 700 for SYTO 9, 1000 for Alexa Fluor 488, 900 for FM4-64, and 700 for propidium iodide. The Zeiss Zen v3.0 was used to control the microscope and for Airyscan processing. Image analysis was conducted using ImageJ/FIJI v2.14 (72).

Construction of a transformation-deficient *S. pneumoniae* mutant

The upstream and downstream regions of *comGC* were amplified from *S. pneumoniae* D39 gDNA, and *tetM* was amplified from *S. mitis* F22 gDNA. DNAs were ligated by Gibson Assembly and amplified using nested primers. *S. pneumoniae* transformants were selected in BA with tetracycline.

Horizontal gene transfer in biofilms

Dual-species biofilms of *S. pneumoniae* D39-GFP Δ*comGC* (chloramphenicol and tetracycline resistant; Chl^R Tet^R) and *S. mitis* (kanamycin resistant; Kan^R) were grown as described above, except that C+Y$_{YB}$ was adjusted to pH 6.6 to prevent natural competence induction. After 24h, non-adherent cells were removed by aspiration, and 2 mL of fresh C+Y$_{YB}$ (pH 7.4) containing catalase and 1 µM of CSP$_{G22}$ were added to biofilms. After 6 h of incubation at 34ºC with 5% CO$_2$, biofilm cells were thoroughly resuspended in PBS and recovered by centrifugation.

Cells were plated on the following selective media: (i) BA with kanamycin to quantify total *S. mitis* (Kan^R); (ii) BA with kanamycin and chloramphenicol to detect *S. mitis* transformants that have

acquired chloramphenicol resistance (Kan^R Chl^R), either with or without tetracycline resistance; (iii)

BA with kanamycin and tetracycline to detect *S. mitis* transformants that acquired tetracycline

resistance (Kan^R Tet^R), either with or without chloramphenicol resistance; (iv) BA with kanamycin,

chloramphenicol and tetracycline to identify *S. mitis* transformants that acquired both resistances

(Kan^R Chl^R Tet^R); (v) and BA with chloramphenicol and tetracycline (without kanamycin) to detect *S.*

pneumoniae loads.

Transformants were counted after 48 h of incubation. *S. mitis* transformation frequencies were

calculated as the ratio between the number of transformants (Kan^R colonies also resistant to

chloramphenicol and/or tetracycline) and the total number of Kan^R colonies. *S. mitis* G22 Δ*comE* was

used as a non-transformable control, in which acquisition of chloramphenicol and tetracycline

resistance should not occur via transformation.

Statistical analyses

The two-tailed unpaired *Mann-Whitney U* test with Benjamini and Hochberg correction for FDR was

used to compare bacterial loads. Student's *t*-test with Benjamini and Hochberg correction for FDR

was used to compare gene expression. The ratio paired *t*-test was used for the statistical analysis of

bacterial proportions in multi-species biofilms. All statistical analysis were computed using GraphPad

Prism 9.0 and considered significant at *P*-value ≤0.05.

Data availability

All data is available from the corresponding author upon reasonable request.

Acknowledgements

We are grateful to Jan-Willem Veening (University of Lausanne) and Morten Kjos (Norwegian

University of Life Sciences) for providing plasmids pPEP-LGZ and Jason Rosch (St. Jude Children's

Research Hospital) for providing *S. pneumoniae* strain harboring a PhunSweet cassette. We also

acknowledge Luisa N. Hiller (Carnegie Mellon University) and Adriano O. Henriques (ITQB NOVA)

for fruitful discussions.

Funding

This work was supported by FCT – Fundação para a Ciência e Tecnologia, I.P., through project

STOPneumo   (PTDC/BIA-MIC/30703/2017),   MOSTMICRO-ITQB   R&D   Unit

(doi.org/10.54499/UID/04612/2025, UID/PRR/4612/2025), LS4FUTURE Associated Laboratory (DOI

10.54499/LA/P/0087/2020), and PPBI - Portuguese Platform of BioImaging (PPBI-POCI-01-0145-

FEDER-022122) co-funded by national funds from OE - "Orçamento de Estado" and by european

funds from FEDER - "Fundo Europeu de Desenvolvimento Regional". J. Borralho and J. Lança were

supported   by   PhD   fellowships   (2021.07866.BD   -   doi.org/10.54499/2021.07866.BD   and

UI/BD/153385/2022 - doi.org/10.54499/UI/BD/153385/2022).

Author Contributions: R. Sá-Leão and J. Borralho contributed to the concept and design of the study. R. Sá-Leão contributed with reagents and materials. Data acquisition was performed by J. Borralho, J. Lança, J. Bryton and W. Antunes. Data was interpreted by J. Borralho, J. Lança, J. Bryton, W. Antunes and R. Sá-Leão. The manuscript was drafted by J. Borralho and R. Sá-Leão. All authors critically revised and approved the final version of the manuscript.

Competing Interest Statement: Universidade NOVA de Lisboa has filed a provisional patent application that covers pharmaceutical compositions comprising strain G22 and/or bacteriocin molecules, and derivatives thereof, which can inhibit the growth and persistence of the pathogen *S. pneumoniae* (PT119647).

References

1.   Griffith F. 1928. The significance of pneumococcal types. J Hyg (Lond) 27:113-59.

2.   Håvarstein LS, Coomaraswamy G, Morrison DA. 1995. An unmodified heptadecapeptide
pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*.
Proc Natl Acad Sci U S A 92:11140-4.

3.   Håvarstein LS, Gaustad P, Nes IF, Morrison DA. 1996. Identification of the streptococcal
competence-pheromone receptor. Mol Microbiol 21:863-9.

4.   Hui FM, Morrison DA. 1991. Genetic transformation in *Streptococcus pneumoniae*: nucleotide
sequence analysis shows *comA*, a gene required for competence induction, to be a member
of the bacterial ATP-dependent transport protein family. J Bacteriol 173:372-81.

5.   Pestova EV, Håvarstein LS, Morrison DA. 1996. Regulation of competence for genetic
transformation in *Streptococcus pneumoniae* by an auto-induced peptide pheromone and a
two-component regulatory system. Mol Microbiol 21:853-62.

6.   Lee MS, Morrison DA. 1999. Identification of a new regulator in *Streptococcus pneumoniae*
linking quorum sensing to competence for genetic transformation. J Bacteriol 181:5004-16.

7.   Peterson SN, Sung CK, Cline R, Desai BV, Snesrud EC, Luo P, Walling J, Li H, Mintz M,
Tsegaye G, Burr PC, Do Y, Ahn S, Gilbert J, Fleischmann RD, Morrison DA. 2004.
Identification of competence pheromone responsive genes in *Streptococcus pneumoniae* by
use of DNA microarrays. Mol Microbiol 51:1051-70.

8.   Guiral S, Mitchell TJ, Martin B, Claverys JP. 2005. Competence-programmed predation of
noncompetent cells in the human pathogen *Streptococcus pneumoniae*: genetic
requirements. Proc Natl Acad Sci U S A 102:8710-5.

9.   Moreno-Gámez S, Sorg RA, Domenech A, Kjos M, Weissing FJ, van Doorn GS, Veening JW.
2017. Quorum sensing integrates environmental cues, cell density and cell history to control
bacterial competence. Nat Commun 8:854.

10.   Prudhomme M, Berge M, Martin B, Polard P. 2016. Pneumococcal competence coordination
relies on a cell-contact sensing mechanism. PLoS Genet 12:e1006113.

11.   Prudhomme M, Johnston CHG, Soulet AL, Boyeldieu A, De Lemos D, Campo N, Polard P.
2024. Pneumococcal competence is a populational health sensor driving multilevel
heterogeneity in response to antibiotics. Nat Commun 15:5625.

12.   Johnsborg O, Eldholm V, Bjørnstad ML, Håvarstein LS. 2008. A predatory mechanism

780   dramatically increases the efficiency of lateral gene transfer in *Streptococcus pneumoniae* and

781   related commensal species. Mol Microbiol 69:245-53.

782   13.   Shen P, Lees JA, Bee GCW, Brown SP, Weiser JN. 2019. Pneumococcal quorum sensing

783   drives an asymmetric owner-intruder competitive strategy during carriage via the competence

784   regulon. Nat Microbiol 4:198-208.

785   14.   Hohwy J, Reinholdt J, Kilian M. 2001. Population dynamics of *Streptococcus mitis* in its natural

786   habitat. Infect Immun 69:6055-63.

787   15.   Salvadori G, Junges R, Morrison DA, Petersen FC. 2019. Competence in *Streptococcus*

788   *pneumoniae* and close commensal relatives: mechanisms and implications. Front Cell Infect

789   Microbiol 9:94.

790   16.   Salvadori G, Junges R, Åmdal HA, Chen T, Morrison DA, Petersen FC. 2018. High-resolution

791   profiles of the *Streptococcus mitis* CSP signaling pathway reveal core and strain-specific

792   regulated genes. BMC Genomics 19:453.

793   17.   Berg KH, Biørnstad TJ, Johnsborg O, Håvarstein LS. 2012. Properties and biological role of

794   streptococcal fratricins. Appl Environ Microbiol 78:3515-22.

795   18.   Potgieter E, Chalkley LJ. 1995. Relatedness among penicillin-binding protein 2b genes of

796   *Streptococcus mitis*, *Streptococcus oralis*, and *Streptococcus pneumoniae*. Microb Drug

797   Resist 1:35-42.

798   19.   Hakenbeck R. 2000. Transformation in *Streptococcus pneumoniae*: mosaic genes and the

799   regulation of competence. Res Microbiol 151:453-6.

800   20.   Rieger M, Mauch H, Hakenbeck R. 2017. Long persistence of a *Streptococcus pneumoniae*

801   23F clone in a cystic fibrosis patient. mSphere 2.

802   21.   Lessa FC, Milucky J, Rouphael NG, Bennett NM, Talbot HK, Harrison LH, Farley MM, Walston

803   J, Pimenta F, Gertz RE, Rajam G, Carvalho MDG, Beall B, Whitney CG. 2018. *Streptococcus*

804   *mitis* expressing pneumococcal serotype 1 capsule. Sci Rep 8:17959.

805   22.   Pimenta F, Gertz RE, Park SH, Kim E, Moura I, Milucky J, Rouphael N, Farley MM, Harrison

806   LH, Bennett NM, Bigogo G, Feikin DR, Breiman R, Lessa FC, Whitney CG, Rajam G, Schiffer

J, da Gloria Carvalho M, Beall B. 2018. *Streptococcus infantis*, *Streptococcus mitis*, and

Streptococcus oralis strains with highly similar *cps5* loci and antigenic relatedness to serotype

5 pneumococci. Front Microbiol 9:3199.

23.   Kalizang'oma A, Richard D, Kwambana-Adams B, Coelho J, Broughton K, Pichon B, Hopkins

KL, Chalker V, Beleza S, Bentley SD, Chaguza C, Heyderman RS. 2024. Population genomics

of *Streptococcus mitis* in UK and Ireland bloodstream infection and infective endocarditis

cases. Nat Commun 15:7812.

24.   Skov Sørensen UB, Yao K, Yang Y, Tettelin H, Kilian M. 2016. Capsular polysaccharide

expression in commensal *Streptococcus* species: genetic and antigenic similarities to

Streptococcus pneumoniae. mBio 7.

25.   Denapaite D, Brückner R, Nuhn M, Reichmann P, Henrich B, Maurer P, Schähle Y, Selbmann

P, Zimmermann W, Wambutt R, Hakenbeck R. 2010. The genome of *Streptococcus mitis* B6

- what is a commensal? PLoS One 5:e9426.

26.   Ikryannikova LN, Malakhova MV, Lominadze GG, Karpova IY, Kostryukova ES, Mayansky

NA, Kruglov AN, Klimova EA, Lisitsina ES, Ilina EN, Govorun VM. 2017. Inhibitory effect of

streptococci on the growth of *M. catarrhalis* strains and the diversity of putative bacteriocin-

like gene loci in the genomes of *S. pneumoniae* and its relatives. AMB Express 7:218.

27.   Borralho J, Handem S, Lança J, Ferreira B, Candeias C, Henriques AO, Hiller NL, Valente C,

Sá-Leão R. 2025. Inhibition of pneumococcal growth and biofilm formation by human isolates

of *Streptococcus mitis* and *Streptococcus oralis*. Appl Environ Microbiol 91:e0133624.

28.   Lehtinen S, Croucher NJ, Blanquart F, Fraser C. 2022. Epidemiological dynamics of

bacteriocin competition and antibiotic resistance. Proc Biol Sci 289:20221197.

29.   Håvarstein LS, Diep DB, Nes IF. 1995. A family of bacteriocin ABC transporters carry out

proteolytic processing of their substrates concomitant with export. Mol Microbiol 16:229-40.

30.   Sati H, Carrara E, Savoldi A, Hansen P, Garlasco J, Campagnaro E, Boccia S, Castillo-Polo

JA, Magrini E, Garcia-Vello P, Wool E, Gigante V, Duffy E, Cassini A, Huttner B, Pardo PR,

Naghavi M, Mirzayev F, Zignol M, Cameron A, Tacconelli E, Group WBPPLA. 2025. The WHO

Bacterial Priority Pathogens List 2024: a prioritisation study to guide research, development,

and public health strategies against antimicrobial resistance. Lancet Infect Dis.

31.   Kilian M, Poulsen K, Blomqvist T, Håvarstein LS, Bek-Thomsen M, Tettelin H, Sørensen UB.

2008. Evolution of *Streptococcus pneumoniae* and its close commensal relatives. PLoS One

3:e2683.

32.   Pozzi G, Masala L, Iannelli F, Manganelli R, Havarstein LS, Piccoli L, Simon D, Morrison DA.

1996. Competence for genetic transformation in encapsulated strains of *Streptococcus*

pneumoniae: two allelic variants of the peptide pheromone. J Bacteriol 178:6087-90.

33.   Nes IF, Diep DB, Holo H. 2007. Bacteriocin diversity in *Streptococcus* and *Enterococcus*. J

Bacteriol 189:1189-98.

34.   Meza-Torres J, Lelek M, Quereda JJ, Sachse M, Manina G, Ershov D, Tinevez JY,

Radoshevich L, Maudet C, Chaze T, Giai Gianetto Q, Matondo M, Lecuit M, Martin-Verstraete

I, Zimmer C, Bierne H, Dussurget O, Cossart P, Pizarro-Cerdá J. 2021. Listeriolysin S: A

bacteriocin from *Listeria monocytogenes* that induces membrane permeabilization in a

contact-dependent manner. Proc Natl Acad Sci U S A 118.

35.   García-Bayona L, Guo MS, Laub MT. 2017. Contact-dependent killing by

Caulobacter crescentus via cell surface-associated, glycine zipper proteins. Elife 6.

36.   Oppegård C, Schmidt J, Kristiansen PE, Nissen-Meyer J. 2008. Mutational analysis of putative

helix-helix interacting GxxxG-motifs and tryptophan residues in the two-peptide bacteriocin

lactococcin G. Biochemistry 47:5242-9.

37.   Ekblad B, Kyriakou PK, Oppegård C, Nissen-Meyer J, Kaznessis YN, Kristiansen PE. 2016.

Structure-function analysis of the two-peptide bacteriocin plantaricin EF. Biochemistry

55:5106-16.

38.   Ekblad B, Kristiansen PE. 2019. NMR structures and mutational analysis of the two peptides

constituting the bacteriocin plantaricin S. Sci Rep 9:2333.

39.   Moll G, Ubbink-Kok T, Hildeng-Hauge H, Nissen-Meyer J, Nes IF, Konings WN, Driessen AJ.

1996. Lactococcin G is a potassium ion-conducting, two-component bacteriocin. J Bacteriol

178:600-5.

40. McAuliffe O, Ryan MP, Ross RP, Hill C, Breeuwer P, Abee T. 1998. Lacticin 3147, a broad-
spectrum bacteriocin which selectively dissipates the membrane potential. Appl Environ
Microbiol 64:439-45.

41. Moll GN, van den Akker E, Hauge HH, Nissen-Meyer J, Nes IF, Konings WN, Driessen AJ.
1999. Complementary and overlapping selectivity of the two-peptide bacteriocins plantaricin
EF and JK. J Bacteriol 181:4848-52.

42. Zhang X, Wang Y, Liu L, Wei Y, Shang N, Li P. 2016. Two-peptide bacteriocin PlnEF causes
cell membrane damage to Lactobacillus plantarum. Biochim Biophys Acta 1858:274-80.

43. Wei H, Håvarstein LS. 2012. Fratricide is essential for efficient gene transfer between
pneumococci in biofilms. Appl Environ Microbiol 78:5897-905.

44. Steinmoen H, Knutsen E, Håvarstein LS. 2002. Induction of natural competence in
Streptococcus pneumoniae triggers lysis and DNA release from a subfraction of the cell
population. Proc Natl Acad Sci U S A 99:7681-6.

45. Ganaie F, Branche AR, Peasley M, Rosch JW, Nahm MH. 2022. Effect of oral streptococci
expressing pneumococcus-like cross-reactive capsule types on World Health Organization
recommended pneumococcal carriage detection procedure. Clin Infect Dis 75:647-656.

46. Peterson S, Cline RT, Tettelin H, Sharov V, Morrison DA. 2000. Gene expression analysis of
the *Streptococcus pneumoniae* competence regulons by use of DNA microarrays. J Bacteriol
182:6192-202.

47. Rimini R, Jansson B, Feger G, Roberts TC, de Francesco M, Gozzi A, Faggioni F, Domenici
E, Wallace DM, Frandsen N, Polissi A. 2000. Global analysis of transcription kinetics during
competence development in *Streptococcus pneumoniae* using high density DNA arrays. Mol
Microbiol 36:1279-92.

48. García E. 2025. Structure, function, and regulation of LytA: the N-acetylmuramoyl-L-alanine
amidase driving the "suicidal tendencies" of *Streptococcus pneumoniae-a review.*
Microorganisms 13.

49. Whatmore AM, Barcus VA, Dowson CG. 1999. Genetic diversity of the streptococcal
competence (*com*) gene locus. J Bacteriol 181:3144-54.

50.   Carrolo M, Pinto FR, Melo-Cristino J, Ramirez M. 2009. Pherotypes are driving genetic
differentiation within *Streptococcus pneumoniae.* BMC Microbiol 9:191.

51.   Vestrheim DF, Gaustad P, Aaberge IS, Caugant DA. 2011. Pherotypes of pneumococcal
strains co-existing in healthy children. Infect Genet Evol 11:1703-8.

52.   Aras RA, Kang J, Tschumi AI, Harasaki Y, Blaser MJ. 2003. Extensive repetitive DNA
facilitates prokaryotic genome plasticity. Proc Natl Acad Sci U S A 100:13579-84.

53.   Wang D, Li S, Guo F, Ning K, Wang L. 2017. Core-genome scaffold comparison reveals the
prevalence that inversion events are associated with pairs of inverted repeats. BMC Genomics
18:268.

54.   Treangen TJ, Abraham AL, Touchon M, Rocha EP. 2009. Genesis, effects and fates of
repeats in prokaryotic genomes. FEMS Microbiol Rev 33:539-71.

55.   Bogaardt C, van Tonder AJ, Brueggemann AB. 2015. Genomic analyses of pneumococci
reveal a wide diversity of bacteriocins - including pneumocyclicin, a novel circular bacteriocin.
BMC Genomics 16:554.

56.   Puls JS, Winnerling B, Power JJ, Krüger AM, Brajtenbach D, Johnson M, Bilici K, Camus L,
Fließwasser T, Schneider T, Sahl HG, Ghosal D, Kubitscheck U, Heilbronner S, Grein F. 2024.
Staphylococcus epidermidis bacteriocin A37 kills natural competitors with a unique
mechanism of action. ISME J 18.

57.   Borgeaud S, Metzger LC, Scrignari T, Blokesch M. 2015. The type VI secretion system of
Vibrio cholerae fosters horizontal gene transfer. Science 347:63-7.

58.   Zhu L, Kuang Z, Wilson BA, Lau GW. 2013. Competence-independent activity of
pneumococcal EndA mediates degradation of extracellular DNA and nets and is important for
virulence. PLoS One 8:e70363.

59.   Salvadori G, Junges R, Morrison DA, Petersen FC. 2016. Overcoming the barrier of low
efficiency during genetic transformation of *Streptococcus mitis.* Front Microbiol 7:1009.

60.   Cowley LA, Petersen FC, Junges R, Jimson D Jimenez M, Morrison DA, Hanage WP. 2018.
Evolution via recombination: cell-to-cell contact facilitates larger recombination events in
Streptococcus pneumoniae. PLoS Genet 14:e1007410.

61.   Marks LR, Reddinger RM, Hakansson AP. 2012. High levels of genetic recombination during
nasopharyngeal carriage and biofilm formation in *Streptococcus pneumoniae*. mBio 3.

62.   Majewski J, Zawadzki P, Pickerill P, Cohan FM, Dowson CG. 2000. Barriers to genetic
exchange between bacterial species: *Streptococcus pneumoniae* transformation. J Bacteriol
182:1016-23.

63.   Davison C, Tallman S, de Ste-Croix M, Antonio M, Oggioni MR, Kwambana-Adams B, Freund
F, Beleza S. 2024. Long-term evolution of *Streptococcus mitis* and *Streptococcus pneumoniae*
leads to higher genetic diversity within rather than between human populations. PLoS Genet
20:e1011317.

64.   Stevens KE, Chang D, Zwack EE, Sebert ME. 2011. Competence in *Streptococcus*
pneumoniae is regulated by the rate of ribosomal decoding errors. mBio 2.

65.   Campbell EA, Choi SY, Masure HR. 1998. A competence regulon in *Streptococcus*
pneumoniae revealed by genomic analysis. Mol Microbiol 27:929-39.

66.   Grant CE, Bailey TL, Noble WS. 2011. FIMO: scanning for occurrences of a given motif.
Bioinformatics 27:1017-8.

67.   Handem S, Ferreira B, Valente C, Sá-Leão R. 2025. Genomic and functional insights into
commensal streptococci with anti-pneumococcal activity. BMC Genomics 26:577.

68.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for
large alignments. PLoS One 5:e9490.

69.   Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. Nucleic Acids Res 52:W78-W82.

70.   Kjos M, Miller E, Slager J, Lake FB, Gericke O, Roberts IS, Rozen DE, Veening JW. 2016.
Expression of *Streptococcus pneumoniae* bacteriocins is induced by antibiotics via regulatory
interplay with the competence system. PLoS Pathog 12:e1005422.

71.   Echlin H, Rosch JW. 2020. Advancing genetic tools in *Streptococcus pneumoniae*. Genes
(Basel) 11.

72.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak

P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. Nat Methods

9:676-82.

Figure Legends

Fig. 1. *Streptococcus mitis* genomes encode a plethora of competence-associated bacteriocins (*cab*). (A) Core-SNP phylogeny of 253 *S. mitis* genomes and distribution of competence-induced bacteriocins A and B (*cibAB*), dedicated immunity (*cibC*), and competence-associated bacteriocin (*cab*) genes. (B) Prevalence of CibAB and 95 Cab variants in the *S. mitis* population. (C) Rarefaction analysis of mature Cab amino acid sequences found in the *S. mitis* population.

Fig. 2. Cab bacteriocins are expressed at the late phase of competence. (A) Genetic organization of *cab* cluster of strain *S. mitis* G22, containing two putative bacteriocin (*cabAB*) and an immunity protein (*cabC*) upstream of the ABC-transporter *comAB* genes. The *cabABC* operon promoter (P*cabABC*) contains five tandem copies of SigX-binding sites (SigX box; SigX is the alternative sigma factor that controls gene expression during the late phase of competence). Expression of *comAB* is driven from a different promoter and induced by ComE (represented as an orange circle). (B) Expression of *cabABC* in *S. mitis* G22 containing a P*cabABC*-luciferase (*luc*) reporter, ectopically inserted in its genome. Expression of *cabABC* was inferred by expression of *luc*. *S. mitis* was pre-grown in acidic medium to avoid natural competence development and synthetic cognate competence-stimulating peptide (CSP$_{G22}$) was added to wild-type *S. mitis* and mutants lacking competence regulators. (C) Dose-response of *cabABC* expression across CSP$_{G22}$ concentrations for the *S. mitis luc* reporter containing either the full-length P*cabABC* promoter (which contains five SigX boxes), or a truncated version containing only the most proximal SigX box. (D) Number of SigX boxes within *cab* clusters of varied sizes. Data in panels B and C represent three to four independent experiments. Unpaired *t*-tests were performed to compare *cab* expression. ** $P < 0.05$, *** $P < 0.001$, **** $P < 0.0001$.

Fig. 3. *S. mitis* and *S. pneumoniae* crosstalk through production of distinct CSP pheromones. (A) Effect of CSP pheromones of close streptococci on *cabABC* expression in *S. mitis* G22 and alignment of tested CSP variants (purple – Q14 amino acid residue, present only in inducing peptides). *S. mitis* G22 P*cabABC-luc* was pre-grown in acidic medium, to which exogenous CSP was added. (B) Effect

1001 of CSP with substitutions in residue 14 in the activation of *cabABC* expression in *S. mitis* G22. (C)

1002 Effect of CSP with substitutions in residue 14 in the activation of competence in *S. pneumoniae* of

1003 pherotype CSP-1 (P*comCDE-luc*). Dotted lines represent average luminescence value for nontreated

1004 controls. Experiments were repeated three times, independently. Unpaired *t*-tests were performed to

1005 compare *cab* expression. **** $P < 0.0001$. ns not significant.

1006

Fig. 4. *S. mitis* bacteriocins CabAB mediate contact-dependent growth inhibition of *S. pneumoniae*.

Experiments were done using *S. mitis* G22 Δ*blp4*Δ*blp1* and *S. pneumoniae* D39. The bacteriocin

clusters *blp4* and *blp1* were first deleted from the genome of *S. mitis* G22 as they have anti-

pneumococcal activity (27), which could obscure CabAB-mediated effects. (A) Proportion of *S. mitis*

and *S. pneumoniae* in dual-species biofilm grown for 24 h in competence-permissive conditions in

24-well plates. *S. mitis* mutant p(*cab*⁺) contains the native *cabABC* copy, and an additional copy under

constitutive regulation (promoter *p23*) in a multicopy plasmid. (B) *S. mitis* was grown in competence-

permissive conditions. Cell-free supernatants (CFS) were recovered and cell pellets were boiled. CFS

and heat-killed cells were added to *S. pneumoniae* cultures and growth was monitored. (C) Bacterial

loads of *S. pneumoniae* grown in the presence of *S. mitis* in direct contact (left) or separated with a

0.4 µm pore transwell membrane (right). Biofilms were grown in 12-well plates. As the surface area

available for cell attachment is higher in 12-well plates than in 24-well plates, we hypothesize that this

contributed to the observed differences in maximum cell counts for D39. (D) Corresponding bacterial

loads of *S. mitis* G22 in experiment (C). (E) Representative images of CabA-HA and CabB-HA

immunodetection on the surface of live competent *S. mitis* cells containing the native *cabABC*

(negative control), *cabA-HA cabBC* or *cabA cabB-HA cabC* genotypes. *S. mitis* cultures were induced

with CSP$_{G22}$, incubated with a primary anti-HA rabbit antibody, and then with an Alexa Fluor 488-

conjugated goat anti-rabbit secondary antibody. Cells were stained with FM4-64 (membrane dye) and

DAPI (nucleic acid dye). At least ten independent fields were detected per experiment/condition.

Experiments were repeated at least three times, independently. The ratio paired *t*-test was used to

compare bacterial proportions in dual-species biofilms. The two-tailed unpaired Mann-Whitney U test

was used to compare bacterial loads. ** $P < 0.01$. *** $P < 0.001$. ns not significant.

1029

Fig. 5. CabAB is a two-peptide bacteriocin with LytA-dependent bactericidal activity against *S.*

pneumoniae. (A) *S. pneumoniae* was treated with synthetic CabA, CabB, an equimolar mixture of

CabA and CabB (0.975 µM), or with DMSO (solvent where peptides were resuspended). Growth was

recorded for 24 h. (B) Presence of GxxxG and GxxxG-like amino acid motifs in CabA and CabB

sequences. (C) Exponential cultures of wild-type and autolysin-deficient *S. pneumoniae* were treated

with DMSO (grey) or synthetic CabAB (purple). Growth was recorded for 24 h. For easier visualization

of lysis (decrease in optical density), the first 12 h of growth is represented. Growth curves were

repeated three times independently, and one representative experiment with two technical replicates

per condition is shown.

1039

Fig. 6. CabAB leads to permeabilization and depolarization of *S. pneumoniae* membranes, leading

to metabolite leakage. (A) *S. pneumoniae* treated with DMSO or CabAB and imaged by scanning

electron microscopy (SEM). (B) Quantification of membrane-permeable *S. pneumoniae* (stained with

propidium iodide (PI)) upon treatment with DMSO (560 cells) or increasing concentrations of CabAB

(0.975 µM – 481 cells; or 1.95 µM – 397 cells). (C-D) Quantification of populational membrane

permeability (C) and depolarization (D) upon DMSO or CabAB treatment at several timepoints. (E)

Proportion of extracellular *vs.* total ATP in wild-type and Δ*lytA S. pneumoniae* cultures upon DMSO

or CabAB treatment. (F) Representative transmission electron microscopy (TEM) image and

quantification of intracellular aggregates in DMSO- (n = 118 cells) or CabAB- treated *S. pneumoniae*

(n = 138 cells). (G) Labeling of large intracellular aggregates by membrane dye FM4-64 in CabAB-

treated *S. pneumoniae*. SEM and TEM were done once. Other data was taken from at least three

independent experiments.

1052

Fig. 7. Expression of *cabAB* facilitates horizontal gene transfer in dual-species biofilms. (A)

Experimental setup to study gene transfer in dual-species biofilms. The *S. pneumoniae* DNA donor is

nontransformable due to a *comGC* deletion, and contains chloramphenicol and tetracycline resistance

cassettes at opposite ends of the chromosome (Chl^R and Tet^R); the *S. mitis* DNA receiver is labeled

with a kanamycin resistance cassette (Kan^R); biofilms were pre-grown at a starting 1:1 ratio for 24 h

in acidic medium to prevent natural competence induction; cells were aspirated and fresh medium

containing CSP$_{G22}$ was added to biofilms to transiently induce competence; after 6 h of incubation,

cells were plated; *S. mitis* transformants were recovered and counted by selection in plates with

kanamycin and combinations of chloramphenicol and/or tetracycline. (B-C) Acquisition of Chl^R and/or

Tet^R by *S. mitis* in biofilms containing purified donor gDNA (B) or the strain itself (C). (D) Growth

inhibition and lysis of 26 bacterial species when treated with 0.975 µM CabAB *vs.* DMSO during their

exponential phase. Growth was recorded for 24 h. Lysis is shown when a decrease from the initial

culture optical density was observed in all replicates. Experiments were repeated three times,

independently. Transformation efficiencies were compared using unpaired *t*-tests. LOD limit of

detection. * $P < 0.05$. ** $P < 0.01$.

1068

Fig. 8. Model for the role of Cab bacteriocins in the acquisition of heterologous DNA. *S. mitis* produces

the CSP precursor (preCSP) intracellularly, which is exported and processed by ComAB. Extracellular

CSP(CSP$_{G22}$), as well as other CSP variants such as pneumococcal CSP-1, may be recognized by

the membrane-bound histidine kinase ComD, leading to phosphorylation of the response regulator

ComE. Phosphorylated ComE (ComE~P) activates expression of the early competence genes,

establishing a positive feedback loop. This includes *sigX1 and sigX2*, which encode for the alternative

sigma factor that drives expression of late competence genes. Among the late genes are the *cab*

bacteriocins. Cab localize to the surface of producer cells. Surface-associated Cab mediate lysis of

neighbouring *S. pneumoniae* cells in a cell contact- and LytA-dependent manner, resulting in the

release of DNA. During the late competence phase, *S. mitis* cells also assemble a competence pilus

responsible for the uptake of extracellular DNA, which can subsequently be integrated into the

genome by homologous recombination.

Figure 1

Figure 2

Figure 3

CSP type	Amino acid sequence	Identity
CSP_{G22}	**E S R L S R L L R D F I F Q I K Q**	-
CSP-1	E M R L S K F F R D F I L Q R K K	10/17
CSP-2	E M R I S R I I L D F L F L R K K	8/17
CSP_{A22}	D W R I S E T I R N L I F P R R K	5/17
CSP_{B22}	E S R V S R I I L D F L F L R K K	9/17
CSP_{D22}	D S R I S K F L R D F I F P R K K	10/17
CSP_{E22}	E M R L P K I L R D F I F P R K K	10/17

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplemental material

Streptococcus mitis bacteriocins drive contact-dependent lysis of

S. pneumoniae facilitating transformation in multispecies environments

João Borralho[1], João Lança,[1], Joana Bryton[1], Wilson Antunes[2], Raquel Sá-Leão[1]*

[1] Laboratory of Molecular Microbiology of Human Pathogens, Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Oeiras, Portugal

[2] Centro de Investigação da Academia Militar (CINAMIL), Unidade Militar Laboratorial de Defesa Biológica e Química (UMLDBQ), Av. Dr. Alfredo Bensaúde, 1849-012, Lisboa, Portugal.

* Corresponding author: rsaleao@itqb.unl.pt

Keywords: Horizontal gene transfer, competence, interspecies interactions, predation, biofilms

Fig. S1. Alignment-based clustering of Cab variants. The 95 Cab variants found *S. mitis* populations were clustered based on pairwise sequence alignments (see Materials and Methods). Alignments for Cab sequences within each cluster is shown, with associated Alphafold2 (ColabFold v1.5.5) structural model of the most prevalent variant. Amino acids are colored according to biochemical traits (RasMol coloring). Structures are colored according to pIDDT (predicted local distance difference test) confidence.

(continues next page)

Fig. S1. (cont.)

Cluster 15
RVSWRGNINIDVHKVIDSTGRFFGGIYNAGRAEGRNVYNAFY
RVSWRGNINIDVHKVIDSTGRFFGGIYNAGRSFGRNVYNAFY
RVSWRGNININIDVHKVIDSTGRFFGGIYNAGRXFGRNVYNAFY

Cluster 16
FVPSPSSVSDSMTSSNRVPS

Cluster 17
LIITTSTAIAWGVLGVVTYMYGRHLGSRR
LIITTSTAIAWGILGVVTYMYGRHLGSRR
LIITTSTAIAWGXLGVVTYMYGRHLGSRR

Cluster 18
GCNWGDFAKAGVGGGAARGLQLGIKTGTWQGAATGAAGGAILGGVAYAATCWW

Cluster 19
IAPIIIGGVVIGWKLIGGVATLATVSAGAGIAAGYYANRP
IAPIIIGGVVIGWKLIGGVAALATVSAGAGIAAGYYANRP
IAPIIIGGVVISWKLIGGVAALATASAGAGIAAGYYANRS
IAPIIIGGVVISWKLIGGVAALATASAGAGIAAGYYANRP
IAPIIIGGVVIXWKLIGGVAALATXSAGAGIAAGYYANRP

Cluster 20
SVVGALALAWGVTKTVGIIGGTAVSVYVAGKSVKAIGDWGLSRLR
SVVGALALAWGVTKAVGIIGGTAVSVYVAGKSVKAIGDWGLSRLR
SVVGA-ALAWGVTKALGIIGGTAVSVYVAGKSVKAIGDWGLSRLR
SVVGALALAWGVTKAVGIIIGGTAVSVYVAGKSVKAIGDWGLSRLR

Cluster 21
IAPLVVAFGVAAFKGFSVTGGAIILGAGLYQVGKNLK

Cluster 22
VGLLIVIGGSLLAAGGAGVAGYWLSRTFG
VGLLIVIGGSLLAAGGAGVVGYWLSRTFG
VGLLIVIGGSLLAAGGAGVXGYWLSRTFG

Cluster 23
IIPIAVAASIIAGSAFAGLSAGVAIGISRNKR

Cluster 24
FFGPVLFTIW

Cluster 25
I-PISVIFGLSAIAFGAFGAGYTFGSDLARRGR
IEPISIIFGLSAIAFGAFGAGYTFGSDLARRGR
IEPISXIFGLSAIAFGAFGAGYTFGSDLARRGR

Cluster 26
ALPLLIPIAAGFIKGFIYTGGAILTTRALLSGN
ALPLLIPIAAGFVKGFVYTGGAILTTRALLSGN
A-PLLIPIAAGFVKGFVYTGGAILTTRALLSGK
ALPLLIPIAAGFVKGFVYTGGAILTTRALLSGN

Cluster 27
LLFGLVVVGGVAIAGTFLLGAVDGASGVHAKRRGR
LLFGLVVVGGVAIAGTFLLGAVDGASGAHAKRRGR
L-FGLVVVGGVAIAGTFLLGAVDGASGAHSKRRGR
L-FGLAVVGGVAIAGTFLLGAVDGASGAHAKRRGR
LLFGLVVVGGVAIAGTFLLGAVDGASGAHAKRRGR

Cluster 28
LLFGLVVVGGVAIAGTFYWVR

Acidic - Asp (D), Glu (E)
Small/flexible – Gly (G)
Sulfur-containing – Cys (C), Met (M)
Aliphatic hydrophobic – Leu (L), Val (V), Ile (I)
Basic – Lys (K), Arg (R)
Small hydrophobic – Ala (A)
Polar uncharged – Ser (S), Thr (T)
Aromatic bulky   Trp (W)
Aromatic – Phe (F), Tyr (Y)
Weakly basic – His (H)
Polar uncharged (amide) – Asn (N), Gln (Q)
Cyclic/ring – Pro (P)

pIDDT: Very low (<50)   Low (60)   OK (70)   Confident (80)   Very high (>90)

Fig. S2. *S. mitis* competence is inhibited in medium with acidic pH. *S. mitis* G22 harboring an early-competence reporter (P*comCDE-luc*) was pre-grown in medium at pH 6.6, before being diluted 1:100 in fresh medium at pH 6.6, 7.4, or 8.2. Optical density (closed circles) and luminescence (open circles) values were measured every 10 min for 24 h. Growth curves were repeated three times independently, and one representative experiment with two technical replicates per condition is shown.

Fig. S3. Expression of *cabABC* follows the same dynamics of other competence genes. (A) Expression of *comCDE*, *sigX1*, *sigX2* and *cabABC* evaluated by luminescence produced by luminescent reporter *S. mitis*. Cells were pre-grown in acidic pH and diluted in fresh medium containing synthetic CSP$_{G22}$. Values represent the highest expression value (RLU.OD$_{595}^{-1}$) obtained during the 3 h post CSP$_{G22}$ treatment. (B) Dynamics of *comCDE*, *sigX1*, *sigX2* and *cabABC* expression in *S. mitis*. Optical density – closed circles. Luminescence – open circles. Growth curves were repeated four times independently, and one representative experiment with two technical replicates per condition is shown. For clarity only the first 8 h of 24 h are shown, since there is no visible expression of the four reporters after this period.

Fig. S4. Expression of *cabABC* is not induced by bacteriocin-like peptide pheromones (BlpC). *S. mitis* G22 harboring a P*cabABC-luc* reporter was pre-grown in acidic pH and diluted in fresh medium containing 1 μM of synthetic CSP_{G22} (grey), 1 μM of $BlpC_1$ or $BlpC_4$ pheromones (blue) or left untreated (dashed line). $BlpC_1$ or $BlpC_4$ are encoded in two *blp*-type bacteriocin clusters elsewhere in the genome of G22. Experiment were repeated four times independently.

■ *S. mitis* loads   ■ *S. pneumoniae* loads   ▧ Total loads (*S. mitis* + *S. pneumoniae*)

Fig. S5. Expression of *cab* by *S. mitis* reduces *S. pneumoniae* loads in dual-species biofilms. Individual and total bacterial loads of biofilm proportions presented in Figure 4A of the main text. *S. pneumoniae* and *S. mitis* of different *cab* genotypes were grown in 24 h biofilms in competence-permissive conditions. *S. mitis* and *S. pneumoniae* cell loads were enumerated by plating in BA with kanamycin or chloramphenicol, respectively. The experiment was repeated three times, independently.

Fig. S6. Synthetic CabAB peptides can cross the transwell membrane. *S. pneumoniae* D39 cells were diluted to ~10^5 cfu/mL in C+Y$_{YB}$ and 1.5 mL was added to 12-well plates. An insert containing a 0.4 µm pore polycarbonate membrane was placed on top and 0.5 mL of C+Y$_{YB}$ with 0.975 µM CabAB or 0.00975% DMSO was added. Aliquots were taken from the bottom compartment at indicated timepoints and plated do assess *S. pneumoniae* loads. The experiments was repeated three times, independently.

Fig. S7. Permeabilization of *S. pneumoniae* membranes does not depend on autolysin LytA. Confocal imaging of wild-type and Δ*lytA S. pneumoniae* treated with 0.00975% DMSO or 0.975 µM CabAB peptides. Membrane permeabilization is shown in red, by entry of propidium iodide. The total number and percentage of permeable cells is shown next to panels. Microscopy images are representative of three independent experiments.

Fig. S8. CabAB treatment induces formation of large clumps of *S. pneumoniae* cells encased in a matrix of DNA and membrane material. *S. pneumoniae* D39 was treated with 0.00975% DMSO or 0.975 µM CabAB and stained with FM4-64 (membrane dye) and DAPI (nucleic acid dye), before being imaged in a confocal microscope. Experiment was repeated three times, independently.

Fig. S9. Nucleotide sequence homology between *S. pneumoniae* D39 and *S. mitis* G22 flanking the sites of antibiotic resistance gene integration. Upstream and downstream regions of *cat* and *tet(M)* in the *S. pneumoniae* DNA donor were divided in five 1,000 bp blocks each and compared to the *S. mitis* G22 genome using BLASTn. Query coverage was consistently 100%, and the percentage of sequence identity is shown for each 1,000 bp block.

A

Competence-associated
bacteriocin (Cab) locus

SigX boxes   D E F D E G   ComAB

CabD mature sequence (Cluster 10):

AIPLVAWLIANPVTAAKIAGGVAAVGIAGYHYMTGKW

CabE mature sequence (Cluster 26):

ALPLLIPIAAGFVKGFVYTGGAILTTRALLSGN

B

Transformation efficiency ($\log_{10}$)

*** ** ns

TetR   ChlR   TetR + ChlR

LOD

Antibiotic resistance acquired

C

****

Proportion in biofilm

cab^+ Δcab

S. mitis C22 genotype

S. mitis C22 genotype: ● cab^+ ● Δcab ■ *S. mitis* C22   ■ *S. pneumoniae* D39

Fig. S10. The Cab bacteriocins of *S. mitis* C22 enhance HGT and mediate inhibition of *S. pneumoniae* in dual-species biofilms. (A) Genomic arrangement of the *S. mitis* C22 *cab* locus, which contains two copies of the bacteriocin *cabD*, two copies of the bacteriocin *cabE*, and the putative immunity proteins *cabF and cabG.* (B) Acquisition of TetR and ChlR by *S. mitis* C22 when in 12-well biofilms with *S. pneumoniae* DNA donor (D39 *gfp* Δ*comGC*). (C) Proportion of *S. mitis* C22 and *S. pneumoniae* D39 when grown in dual-species biofilm. Experiments were repeated three times, independently. Results in panel C represent the proportion of each species in the dual-species biofilms of panel B. Transformation efficiencies were compared using unpaired t-tests. LOD limit of detection. ** *P* < 0.01. *** *P* < 0.001. **** *P* < 0.0001.

Fig. S11. Individual growth curves of 18 streptococcal species and their sensitivity to CabAB. Exponential cultures of streptococcal species were diluted to OD_{600} 0.1 in appropriate medium and treated with either DMSO or CabAB. Optical density was measured every 10 min for 24 h. For clarity, the first 12 h are shown. Species are colored according to streptococcal group. The experiment was repeated three times, independently. Summary data of the growth curves can be found in the main manuscript.

Fig. S12. Individual growth curves of 4 non-streptococcal Gram-positive and 4 Gram-negative bacteria and their sensitivity to CabAB. Exponential cultures of species were diluted to OD_{600} in appropriate medium and treated with either DMSO or CabAB. Optical density was measured every 10 min for 24 h. For clarity, the first 12 h are shown. Species are colored according to Gram staining. The experiment was repeated three times, independently. Summary data of the growth curves can be found in the main manuscript.

Fig. S13. CabAB treatment induces formation of membrane aggregates exclusively in *S. pneumoniae*. Cells were treated with 0.975 μM CabAB or 0.00975% DMSO for 15 min and stained with FM4-64 (membrane dye) and DAPI (nucleic acid dye), before being imaged in a confocal microscope. The experiment was repeated three times, independently. Scale – 5 μm.

Table S1. *Streptococcus mitis* genomes used in the study.

Strain	Region flanked by *plsX-acpP* and comAB recovered from genome (Y/N)	Presence of *cab* (Y/N)	*cab* locus length (bp)	Number of unique Cabs	Number of SigX boxes in *cab* locus	Presence of *cibAB* (Y/N)	Presence of *cibC* (Y/N)	Mature CSP sequence	4th last amino acid in CSP	Genbank accession	Location of region flanked by *plsX-acpP* and *comAB*
103U_S92	Y	Y	9862	4	20	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946186075.1	23700 to 33569 (contig 5)
125U_S4	Y	N	-	-	-	N	N	EMRRSNNNFFNFLRRI	L	GCA_946184935.1	121366 to 123329 (contig 6)
128U_S6	N	-	-	-	-	-	-		-	GCA_946186995.1	-
12P_S5	Y	Y	6869	5	15	N	N	EIRQTHNIFFNFFKRR	F	GCA_946186085.1	27830 to 34698 (contig 6)
13P_S6	Y	Y	9282	5	16	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946187105.1	25159 to 34440 (contig 14)
14P_S7	Y	Y	2287	2	3	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946186685.1	19479 to 21765 (contig 2)
161U_S19	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946188375.1	109310 to 116321 (contig 4)
1P_S1	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946183265.1	10854 to 17772 (contig 11)
22P_S11	N	-	-	-	-	-	-		-	GCA_946183395.1	-
27P_S14	Y	Y	7997	5	17	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946184735.1	22263 to 30259 (contig 10)
32P_S19	Y	Y	6869	5	15	N	N	EIRQTHNIFFNFFKRR	F	GCA_946223905.1	28620 to 35488 (contig 11)
38P_S23	Y	Y	6883	3	17	N	N	ESRLSKLLRDFIFKIKQ	K	GCA_946183455.1	26015 to 32897 (contig 9)
3P_S2	N	-	-	-	-	-	-		-	GCA_946183495.1	-
44P_S26	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946186695.1	36560 to 43673 (contig 10)
50P_S29	Y	Y	7547	4	15	Y	Y	ESRLSRLLRDFIFQIKQ	Q	GCA_946188075.1	25384 to 32930 (contig 8)
52P_S30	N	-	-	-	-	-	-		-	GCA_946186135.1	-
61P_S37	N	-	-	-	-	-	-		-	GCA_946183505.1	-
62P_S38	Y	Y	7457	5	15	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946183485.1	24444 to 24837 (contig 5)
68U_S84	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_946183445.1	13397 to 18312 (contig 11)
7P_S3	N	-	-	-	-	-	-		-	GCA_946185985.1	-
B1_S23	Y	Y	3685	6	4	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946189985.1	18059 to 21743 (contig 6)
B10_S32	N	-	-	-	-	-	-		-	GCA_946183015.1	-
B100_S21	N	-	-	-	-	-	-		-	GCA_946186585.1	-
B1015B24C5	Y	Y	8698	4	17	N	Y	EMRKSNSTFFHFLRRI	L	GCA_905220945.1	27478 to 36175 (contig 5)
B1015G24C3	Y	Y	3544	6	7	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_905221115.1	3577 to 7120 (contig 16)
B1015H24C1	Y	Y	9990	4	18	Y	Y	ESRVSDLILDFLFQRKK	Q	GCA_905221235.1	24516 to 34505 (contig 8)
B1015H24C3	Y	N	-	-	-	N	N	ESRLPKILLDFLFLRKK	L	GCA_905221375.1	45744 to 49408 (contig 11)
B1015H24C4	Y	N	-	-	-	N	N	EMRRIGSVLLNFFKRR	F	GCA_905221455.1	110170 to 117841 (contig 8)
B102_S23	N	-	-	-	-	-	-		-	GCA_946183435.1	-
B105_S26	Y	Y	3557	4	9	N	Y	EMRKMNEKSFNFFNIFRR	I	GCA_946185425.1	22400 to 25956 (contig 13)
B106_S27	Y	Y	6472	5	16	N	N	ESRLSRLLRDFILQRKK	Q	GCA_946184175.1	27052 to 33524 (contig 3)
B107_S28	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946184385.1	73855 to 74301 (contig 3)
B110_S31	Y	N	-	-	-	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946185245.1	58142 to 67806 (contig 5)
B111_S32	Y	N	-	-	-	Y	Y	EIRIADKIFFNLFKRR	F	GCA_946185595.1	35645 to 42337 (contig 8)
B112_S33	Y	Y	8000	5	17	N	Y	EMRLPKILRDFIFPRKK	P	GCA_946185795.1	16961 to 24960 (contig 6)
B113_S34	Y	Y	7352	5	14	N	N	EMRKPDGALFNLFRRR	F	GCA_946186035.1	28045 to 35396 (contig 4)
B114_S35	Y	Y	7477	1	16	Y	Y	ESRLPKILLDFIFPRKK	P	GCA_946184755.1	130119 to 137595 (contig 5)
B116_S37	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946186655.1	65112 to 74936 (contig 6)
B117_S38	Y	N	-	-	-	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946184245.1	112994 to 121923 (contig 7)
B119_S40	Y	N	-	-	-	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946184705.1	31961 to 19319 (contig 13)
B12_S34	Y	Y	6838	5	15	N	N	EMRKPDGALFNLFRRR	F	GCA_946187535.1	27777 to 34614 (contig 5)
B121_S42	Y	N	-	-	-	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_946184465.1	3398 to 5219 (contig 19)
B122_S43	Y	Y	9952	5	14	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946183295.1	27547 to 37498 (contig 7)
B124_S44	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946184765.1	33241 to 41252 (contig 13)
B125_S45	Y	Y	6733	3	14	Y	Y	EIRIADKIFFNLFKRR	F	GCA_946186825.1	10414 to 17146 (contig 9)
B126_S46	Y	Y	7012	5	14	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946183535.1	26548 to 33559 (contig 6)
B128_S47	N	-	-	-	-	-	-		-	GCA_946187305.1	-
B129_S48	N	-	-	-	-	-	-		-	GCA_946183555.1	-
B13_S35	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946184335.1	124084 to 131222 (contig 4)
B132_S51	Y	Y	5685	5	11	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946188115.1	26091 to 31775 (contig 7)
B133-2_S52	N	-	-	-	-	-	-		-	GCA_946187925.1	-

(continues next page)

ID								Sequence		Accession	Position
B134_S53	N	-	-	-	-	-	-	-	-	GCA_946183695.1	-
B137_S55	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_946183685.1	37817 to 47227 (contig 10)
B139_S57	Y	N	-	-	-	Y	Y	ESRMPKILRDFIFPRKK	P	GCA_946183965.1	129975 to 133398 (contig 6)
B14_S36	Y	N	-	-	-	Y	Y	DMRISESIRNLIFPRKKK	R	GCA_946188365.1	132557 to 140438 (contig 5)
B140_S58	Y	Y	7981	7	17	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946188035.1	21526 to 21918 (contig 7)
B146_S64	N	-	-	-	-	-	-	-	-	GCA_946184845.1	
B147_S65	Y	N	-	-	-	Y	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946188285.1	33777 to 37954 (contig 129
B148_S66	N	-	-	-	-	-	-	-	-	GCA_946185855.1	-
B15_S37	Y	N	-	-	-	N	N	EMRKMNDKSFNLFNFFRRR	F	GCA_946188735.1	108533 to 116391 (contig 7)
B151_S69	N	-	-	-	-	-	-	-	-	GCA_946185905.1	-
B153_S71	Y	Y	10165	3	13	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188725.1	12287 to 22451 (contig 7)
B155_S73	N	-	-	-	-	-	-	-	-	GCA_946185295.1	-
B156_S74	N	-	-	-	-	-	-	-	-	GCA_946189305.1	-
B157_S75	Y	N	-	-	-	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_946188715.1	47861 to 48593 (contig 11)
B158_S76	N	-	-	-	-	-	-	-	-	GCA_946188825.1	-
B159_S77	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188815.1	126333 to 135213 (contig 1)
B161_S79	Y	N	-	-	-	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946188505.1	121745 to 133038 (contig 7)
B162_S80	N	-	-	-	-	-	-	-	-	GCA_946188645.1	-
B163_S81	N	-	-	-	-	-	-	-	-	GCA_946188665.1	-
B165_S83	Y	Y	7454	5	12	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946188525.1	9840 to 17293 (contig 6)
B166_S84	N	-	-	-	-	-	-	-	-	GCA_946188555.1	-
B167_S85	Y	N	-	-	-	N	Y	ESRLPKILLDFLFLRKK	L	GCA_946188435.1	51984 to 55661 (contig 7)
B168_S86	Y	N	-	-	-	N	N	ESRLPKILLDFLFLRKK	L	GCA_946188425.1	129601 to 136812 (contig 5)
B169_S87	Y	N	-	-	-	N	Y	EMRLPKILRDFIFPRKK	P	GCA_946189595.1	100635 to 105207 (contig 7)
B170_S88	Y	Y	2522	2	7	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946189625.1	16848 to 19369 (contig 119
B172_S90	N	-	-	-	-	-	-	-	-	GCA_946189565.1	-
B173_S91	N	-	-	-	-	-	-	-	-	GCA_946188785.1	-
B174_S92	Y	N	-	-	-	N	Y	EMRKPDGALFNLFRRR	F	GCA_946189255.1	44630 to 50769 (contig 13)
B176_S94	N	-	-	-	-	-	-	-	-	GCA_946188865.1	-
B19_S40	Y	Y	6430	2	12	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_946190005.1	24129 to 30558 (contig 12)
B20_S41	N	-	-	-	-	-	-	-	-	GCA_946190035.1	-
B21_S42	N	-	-	-	-	-	-	-	-	GCA_946189125.1	-
B22_S43	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946189035.1	45816 to 51979 (contig 12)
B26_S45	Y	Y	7324	1	14	N	N	EMRRIDKIFINFLKRR	L	GCA_946189055.1	36560 to 43883 (contig 12)
B3_S25	Y	Y	6276	6	13	Y	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946189635.1	27034 to 33309 (contig 11)
B30_S48	Y	N	-	-	-	Y	Y	ESRVSDILLDFLFRRKK	R	GCA_946189895.1	13875 to 25318 (contig 11)
B31_S49	Y	Y	7447	5	12	Y	Y	EMRKMNDKSFNLFNFFRRR	F	GCA_946188485.1	24297 to 31743 (contig 13)
B33_S51	Y	Y	11446	5	20	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188605.1	9714 to 21159 (contig 14)
B34_S52	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFNIFRR	F	GCA_946189235.1	27396 to 35035 (contig 17)
B37_S55	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946188755.1	8105 to 15978 (contig 6)
B38_S56	N	-	-	-	-	-	-	-	-	GCA_946188795.1	-
B4_S26	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946189945.1	13981 to 20965 (contig 29)
B42_S60	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188675.1	147283 to 158055 (contig 5)
B43_S61	Y	Y	7325	4	16	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946188595.1	23294 to 30618 (contig 15)
B44_S62	Y	Y	6767	5	14	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188765.1	24640 to 31406 (contig 5)
B45_S63	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946189355.1	8513 to 9245 (contig 6)
B46_S64	N	-	-	-	-	-	-	-	-	GCA_946188655.1	-
B47_S65	Y	N	-	-	-	N	Y	EMRKSNNNFFHFLRRI	L	GCA_946188745.1	43466 to 50055 (contig 7)
B49_S67	N	-	-	-	-	-	-	-	-	GCA_946188965.1	-
B5_S27	N	-	-	-	-	-	-	-	-	GCA_946189025.1	-
B50_S68	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188955.1	126680 to 135327 (contig 7)

Table S1. (cont.)

B52_S70	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946189865.1	13665 to 14681 (contig 16)
B54_S72	N	-	-	-	-	-	-		-	GCA_946188925.1	-
B55_S73	Y	N	-	-	-	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_946188895.1	49166 to 58656 (contig 9)
B56_S74	N	-	-	-	-	-	-		-	GCA_946188995.1	-
B57_S75	Y	N	-	-	-	N	Y	EMRRIDKMFINFLKRR	L	GCA_946188985.1	117917 to 125697 (contig 4)
B59_S77	N	-	-	-	-	-	-		-	GCA_946188875.1	-
B60_S78	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_946189965.1	124476 to 131218 (contig 4)
B61_S79	Y	N	-	-	-	Y	Y	EIRKSNSALVNFFKRR	F	GCA_946188855.1	55573 to 66461 (contig 10)
B63_S81	Y	Y	7221	5	14	N	Y	ESRVSDILDFLFQRKK	Q	GCA_946189015.1	10888 to 18108 (contig 15)
B65_S83	Y	N	-	-	-	N	N	ESRVSDILLDFLFRRKK	R	GCA_946189955.1	118327 to 122899 (contig 3)
B67_S85	Y	Y	3156	4	10	Y	Y	EMRKSNNTFFNFLRRI	L	GCA_946188885.1	7767 to 10922 (contig 13)
B69_S87	N	-	-	-	-	-	-		-	GCA_946188945.1	-
B73_S91	Y	Y	8931	3	19	N	N	ESRVSDILLDFLFRRKK	R	GCA_946188935.1	13555 to 22485 (contig 9)
B74_S92	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_946188975.1	10138 to 10870 (contig 11)
B76_S94	Y	Y	7437	4	12	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_946189935.1	30822 to 38258 (contig 8)
B77_S95	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946189755.1	118938 to 126810 (contig 5)
B78_S96	Y	Y	8714	4	21	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946189875.1	25049 to 33762 (contig 5)
B79_S1	Y	Y	3832	6	10	Y	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946188405.1	8778 to 12609 (contig 18)
B82_S4	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946188465.1	68030 to 72644 (contig 9)
B85_S7	Y	N	-	-	-	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946189515.1	124429 to 125974 (contig 6)
B88_S10	Y	N	-	-	-	N	Y	EMRLPKILRDFIFPRKK	P	GCA_946189785.1	67460 to 77667 (contig 6)
B89L_S11	Y	Y	8758	4	19	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946188455.1	23760 to 32517 (contig 5)
B9_S31	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946189075.1	32842 to 41229 (contig 13)
B91_S12	Y	N	-	-	-	Y	Y	EMRLSKFLRDFILQRKN	Q	GCA_946189535.1	19728 to 28296 (contig 10)
B93_S14	N	-	-	-	-	-	-		-	GCA_946189095.1	-
B95_S16	Y	N	-	-	-	N	N	MKNTVKLEQFVALKEKTCRR FEVGKLEKQLIVYLISSKEDK	K	GCA_946188475.1	57945 to 58714 (contig 10)
B96_S17	Y	Y	7329	1	16	N	N	EIRIADKIFFNLFKRR	F	GCA_946189115.1	35203 to 42531 (contig 12)
B97_S18	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_946189825.1	102775 to 103025 (contig 4)
B99_S20	N	-	-	-	-	-	-		-	GCA_946189005.1	-
GCF_000027165.1_ASM2716v1	Y	Y	9068	5	18	Y	Y	EMRKPDGALFNLFRRR	F	GCA_000027165.1 ASM2716v1	53854 to 62921 (contig 1)
GCF_000148505.1_ASM14850v1	N	-	-	-	-	-	-		-	GCA_000148505.1 ASM14850v1	-
GCF_000148525.1_ASM14852v1	Y	Y	4573	7	7	N	N	ESRVSDILLDFLFRRKK	R	GCA_000148525.1 ASM14852v1	23154 to 27726 (contig 22)
GCF_000148545.1_ASM14854v1	N	-	-	-	-	-	-		-	GCA_000148545.1 ASM14854v1	-
GCF_000148585.2_ASM14858v3	Y	Y	8195	5	17	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_000148585.2 ASM14858v3	46590 to 54784 (contig 1)
GCF_000220045.1_ASM22004v2	Y	N	-	-	-	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_000220045.1 ASM22004v2	113804 to 121511 (contig 10)
GCF_000220085.1_ASM22008v2	Y	Y	7601	3	18	N	N	EMRRIDKIFNFLKRR	L	GCA_000220085.1 ASM22008v2	33312 to 40912 (contig 6)
GCF_000222785.1_ASM22278v2	Y	N	-	-	-	Y	Y	EMRLPKILRDFIFPRKK	P	GCA_000222785.1 ASM22278v2	80829 to 88913 (contig 17)
GCF_000257805.1_ASM25780v1	Y	N	-	-	-	N	N	EMRRIDKIFINFLKRR	L	GCA_000257805.1 ASM25780v1	30349 to 36794 (contig 6)
GCF_000257865.1_ASM25786v1	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_000257865.1 ASM25786v1	108462 to 111048 (contig 6)
SK579v1.0	Y	N	-	-	-	N	N	EIRKSNSALVNFFKRR	F	GCA_000257905.1 SmitisSK579v1.0	51648 to 53976 (contig 13)
GCF_000385815.1_ASM38581v1	N	-	-	-	-	-	-		-	GCA_000385815.1 ASM38581v1	-
Smit.str17_34_v1.0	Y	Y	8147	3	16	N	N	ESRISDLLLDFLFQRKK	Q	GCA_000430305.1 Smit.str17_34_v1.	24690 to 32836 (contig 5)
Smit.str18_56_v1.0	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_000430345.1 Smit.str18_56_v1.	6585 to 7319 (contig 12)
Smit.str29_42_v1.0	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_000430365.1 Smit.str29_42_v1.	28257 to 34755 (contig 8)
Sm.str21_39_v1.0	Y	N	-	-	-	N	N	EMRLPKILRDFIFPRKK	P	GCA_000506725.1 Sm.str21_39_v1.0	107853 to 111953 (contig 6)
Sm.str27_7_v1.0	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_000506765.1 Sm.str27_7_v1.0	48091 to 54587 (contig 8)
GCF_000722685.1_ASM72268v1	Y	Y	6593	3	15	N	N	EIRKSNSALVNFFKRR	F	GCA_000722685.1 ASM72268v1	33237 to 39817 (contig 3)
GCF_000722695.1_ASM72269v1	Y	Y	8106	4	20	N	Y	EMRLPKILRDFIFPRKK	P	GCA_000722695.1 ASM72269v1	28640 to 36745 (contig 2)
GCF_000722705.1_ASM72270v1	Y	Y	7521	3	19	N	Y	EMRRIDKMFINFLKRR	L	GCA_000722705.1 ASM72270v1	26310 to 33830 (contig 1)
GCF_000722755.1_ASM72275v1	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_000722755.1 ASM72275v1	19662 to 20232 (contig 8)

(continues next page)

Table S1. (cont.)

GCF_000722765.2_ASM72276v2	Y	Y	7760	4	18	N	Y	EMRKSNNNFFHFLRRI	L	GCA_000722765.2_ASM72276v2	49608 to 57367 (contig 1)
GCF_000722775.1_ASM72277v1	Y	Y	8930	3	21	N	N	EIRKTSNSLLNFFKRR	F	GCA_000722775.1_ASM72277v1	27763 to 36692 (contig 6)
GCF_000722815.1_ASM72281v1	Y	Y	1789	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_000722815.1_ASM72281v1	20061 to 21849 (contig 1)
GCF_000722835.1_ASM72283v1	N	-	-	-	-	-	-	-	-	GCA_000722835.1_ASM72283v1	-
GCF_000724165.1_ASM72416v1	Y	N	-	-	-	N	N	Not found	-	GCA_000724165.1_ASM72416v1	438770 to 446249 (contig 12)
GCF_000960005.1_ASM96000v1	Y	Y	8611	2	18	Y	Y	EMRKPDGALFNLFRRR	F	GCA_000960005.1_ASM96000v1	29135 to 37745 (contig 1)
GCF_000960025.1_ASM96002v1	Y	Y	12265	4	24	N	N	ESRLPKIRFDFIFPRKK	P	GCA_000960025.1_ASM96002v1	32833 to 45097 (contig 3)
GCF_000960065.1_ASM96006v1	Y	N	-	-	-	N	N	DMRISESIRNLIFPRKKK	R	GCA_000960065.1_ASM96006v1	33075 to 43002 (contig 5)
GCF_001070035.1_ASM107003v1	N	-	-	-	-	-	-	-	-	GCA_001070035.1_ASM107003v1	-
GCF_001070205.1_ASM107020v1	N	-	-	-	-	-	-	-	-	GCA_001070205.1_ASM107020v1	-
GCF_001070375.1_ASM107037v1	N	-	-	-	-	-	-	-	-	GCA_001070375.1_ASM107037v1	-
GCF_001070705.1_ASM107070v1	Y	N	-	-	-	N	N	ESRLSKLLRDFIFKIKQ	K	GCA_001070705.1_ASM107070v1	91104 to 97864 (contig 1)
GCF_001070985.1_ASM107098v1	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_001070985.1_ASM107098v1	125336 to 135287 (contig 33)
GCF_001074635.1_ASM107463v1	N	-	-	-	-	-	-	-	-	GCA_001074635.1_ASM107463v1	-
GCF_001281025.1_ASM128102v1	Y	N	-	-	-	Y	Y	EMRKSNNNFFNFLRRI	L	GCA_001281025.1_ASM128102v1	326587 to 336055 (contig 1)
GCF_001546795.1_ASM154679v1	N	-	-	-	-	-	-	-	-	GCA_001546795.1_ASM154679v1	-
GCF_001560895.1_ASM156089v1	Y	Y	2200	2	5	N	N	EMRKSNNTFFNFLRRI	-	GCA_001560895.1_ASM156089v1	73674 to 75873 (contig 1)
GCF_001579045.1_ASM157904v1	Y	N	-	-	-	Y	Y	EMRLPKILRDFIFPRKK	P	GCA_001579045.1_ASM157904v1	416341 to 423519 (contig 63)
GCF_001588595.1_ASM158859v1	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_001588595.1_ASM158859v1	14693 to 15462 (contig 4)
GCF_001588605.1_ASM158860v1	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_001588605.1_ASM158860v1	34154 to 34922 (contig 15)
GCF_001588625.1_ASM158862v1	Y	Y	9205	5	23	N	N	EMRKPDGALFNLFRRR	F	GCA_001588625.1_ASM158862v1	9739 to 18943 (contig 27)
GCF_002005545.1_ASM200554v1	Y	Y	3980	6	9	N	N	EMRLPKILRDFIFPRKK	P	GCA_002005545.1_ASM200554v1	495219 to 499198 (contig 1)
GCF_002014755.1_ASM201475v1	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_002014755.1_ASM201475v1	115535 to 123729 (contig 6)
GCF_002014795.1_ASM201479v1	Y	Y	8075	4	21	N	Y	EMRKPDGALFNLFRRR	F	GCA_002014795.1_ASM201479v1	26394 to 34468 (contig 6)
GCF_002014875.1_ASM201487v1	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_002014875.1_ASM201487v1	63548 to 69821 (contig 7)
GCF_002096735.1_ASM209673v1	Y	Y	9752	2	14	N	N	EMRKSNNNFFNFLRRI	L	GCA_002096735.1_ASM209673v1	14476 to 24227 (contig 13)
GCF_002096755.1_ASM209675v1	N	-	-	-	-	-	-	-	-	GCA_002096755.1_ASM209675v1	-
GCF_002096765.1_ASM209676v1	Y	Y	6682	6	16	Y	Y	EMRKPDGALFNLFRRR	F	GCA_002096765.1_ASM209676v1	26701 to 33382 (contig 48)
GCF_002096775.1_ASM209677v1	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_002096775.1_ASM209677v1	23754 to 24138 (contig 24)
GCF_002096815.1_ASM209681v1	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_002096815.1_ASM209681v1	14082 to 19251 (contig 18)
GCF_002096835.1_ASM209683v1	Y	N	-	-	-	Y	Y	ESRVSDILLDFLFRRKK	R	GCA_002096835.1_ASM209683v1	4746 to 5469 (contig 19)
GCF_002096845.1_ASM209684v1	N	-	-	-	-	-	-	-	-	GCA_002096845.1_ASM209684v1	-
GCF_002096895.1_ASM209689v1	Y	N	-	-	-	N	N	DSRISKYLRDFIFPRKK	P	GCA_002096895.1_ASM209689v1	125698 to 126269 (contig 19)
GCF_002096925.1_ASM209692v1	N	-	-	-	-	-	-	-	-	GCA_002096925.1_ASM209692v1	-
GCF_002096935.1_ASM209693v1	Y	Y	6733	3	14	N	N	EIRIADKIFFNLFKRR	F	GCA_002096935.1_ASM209693v1	24923 to 31655 (contig 25)
GCF_002860825.1_ASM286082v1	Y	Y	3524	6	5	N	N	EMRKSNNTFFNFLRRI	L	GCA_002860825.1_ASM286082v1	18587 to 22110 (contig 13)
GCF_002860865.1_ASM286086v1	Y	Y	2525	2	5	N	N	ESRVSDLILDFLFQRKK	Q	GCA_002860865.1_ASM286086v1	58524 to 61048 (contig 3)
GCF_002860905.1_ASM286090v1	Y	Y	3156	4	10	Y	Y	EMRRIDKIFINFLKRR	L	GCA_002860905.1_ASM286090v1	18653 to 21808 (contig 8)
GCF_002871315.1_ASM287131v1	Y	Y	3524	6	5	N	N	EMRKSNNTFFNFLRRI	L	GCA_002871315.1_ASM287131v1	18509 to 22032 (contig 13)
GCF_003942775.1_ASM394277v1	N	-	-	-	-	-	-	-	-	GCA_003942775.1_ASM394277v1	-
GCF_003942885.1_ASM394288v1	N	-	-	-	-	-	-	-	-	GCA_003942885.1_ASM394288v1	
GCF_003942965.1_ASM394296v1	Y	N	-	-	-	N	N	Not found	-	GCA_003942965.1_ASM394296v1	11262 to 11512 (contig 3)
GCF_003942995.1_ASM394299v1	Y	N	-	-	-	Y	Y	ESRVSDILLDFLFRRKK	R	GCA_003942995.1_ASM394299v1	30869 to 40944 (contig 14)
GCF_003943915.1_ASM394391v1	N	-	-	-	-	-	-	-	-	GCA_003943915.1_ASM394391v1	-
GCF_003943965.1_ASM394396v1	Y	N	-	-	-	N	Y	EMRRSNNNFFNFLRRI	L	GCA_003943965.1_ASM394396v1	40655 to 42425 (contig 16)
GCF_005144795.1_ASM514479v1	Y	Y	1780	2	4	Y	Y	ESRLSKIFLDFIFPRKK	P	GCA_005144795.1_ASM514479v1	13402 to 15181 (contig 16)
GCF_005844455.1_ASM584445v1	N	-	-	-	-	-	-	-	-	GCA_005844455.1_ASM584445v1	-
GCF_008763935.1_FS623	Y	Y	6780	5	16	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_008763935.1_FS623	107171 to 113950 (contig 5)
GCF_009495695.1_ASM949569v1	N	-	-	-	-	-	-	-	-	GCA_009495695.1_ASM949569v1	-
GCF_009495705.1_ASM949570v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009495705.1_ASM949570v1	23517 to 30981 (contig 17)

(continues next page)

GCF_009495735.1_ASM949573v1	Y	N	-	-	-	N	N	ESRVSDLILDFLFQRKK	Q	GCA_009495735.1 ASM949573v1	88623 to 96087 (contig 8)
GCF_009495895.1_ASM949589v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009495895.1 ASM949589v1	23636 to 31100 (contig 8)
GCF_009495955.1_ASM949595v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009495955.1 ASM949595v1	58616 to 66080 (contig 10)
GCF_009496035.1_ASM949603v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496035.1 ASM949603v1	88623 to 96087 (contig 9)
GCF_009496045.1_ASM949604v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496045.1 ASM949604v1	88615 to 96079 (contig 9)
GCF_009496065.1_ASM949606v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496065.1 ASM949606v1	23577 to 31041 (contig 10)
GCF_009496075.1_ASM949607v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496075.1 ASM949607v1	23517 to 30981 (contig 11)
GCF_009496165.1_ASM949616v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496165.1 ASM949616v1	23517 to 30981 (contig 8)
GCF_009496185.1_ASM949618v1	N	-	-	-	-	-	-	-	-	GCA_009496185.1 ASM949618v1	-
GCF_009496205.1_ASM949620v1	N	-	-	-	-	-	-	-	-	GCA_009496205.1 ASM949620v1	-
GCF_009496295.1_ASM949629v1	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_009496295.1 ASM949629v1	124953 to 131821 (contig 6)
GCF_009496335.1_ASM949633v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496335.1 ASM949633v1	88622 to 96086 (contig 8)
GCF_009496355.1_ASM949635v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496355.1 ASM949635v1	88623 to 96087 (contig 7)
GCF_009496385.1_ASM949638v1	Y	Y	2273	2	6	N	N	EMRKSNNTFFNFLRRI	L	GCA_009496385.1 ASM949638v1	111359 to 113631 (contig 5)
GCF_009496395.1_ASM949639v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496395.1 ASM949639v1	23577 to 31041 (contig 8)
GCF_009496435.1_ASM949643v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496435.1 ASM949643v1	88623 to 96087 (contig 8)
GCF_009496505.1_ASM949650v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496505.1 ASM949650v1	23517 to 30981 (contig 6)
GCF_009496725.1_ASM949672v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496725.1 ASM949672v1	23577 to 31041 (contig 7)
GCF_009496735.1_ASM949673v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496735.1 ASM949673v1	23577 to 31041 (contig 7)
GCF_009496755.1_ASM949675v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496755.1 ASM949675v1	18419 to 25883 (contig 15)
GCF_009496795.1_ASM949679v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496795.1 ASM949679v1	23577 to 31041 (contig 8)
GCF_009496805.1_ASM949680v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496805.1 ASM949680v1	23577 to 31041 (contig 8)
GCF_009496815.1_ASM949681v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496815.1 ASM949681v1	23577 to 31041 (contig 9)
GCF_009496825.1_ASM949682v1	N	-	-	-	-	-	-	-	-	GCA_009496825.1 ASM949682v1	-
GCF_009496835.1_ASM949683v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496835.1 ASM949683v1	23577 to 31041 (contig 9)
GCF_009496895.1_ASM949689v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496895.1 ASM949689v1	23517 to 30981 (contig 15)
GCF_009496905.1_ASM949690v1	Y	N	-	-	-	N	N	EIRKTINGLLNFFKRR	F	GCA_009496905.1 ASM949690v1	51468 to 56727 (contig 9)
GCF_009730515.1_ASM973051v1	Y	Y	8195	5	17	N	N	EIRQTHNIFFNFFKRR	-	GCA_009730515.1 ASM973051v1	31493 to 39687 (contig 1)
GCF_900459425.1_42042_B02	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_900459425.1 42042_B02	116797 to 124991 (contig 2)
MSCx	Y	Y	8886	4	18	Y	Y	DSRISKYLRDFIFPRKK	P	GCA_905187535.1	25487 to 34372 (contig 11)
NM2_1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_905187515.1	32077 to 39698 (contig 13)
NM4_2	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_905187545.1	32687 to 40308 (contig 13)
NM5_2	N	-	-	-	-	-	-	-	-	GCA_905187495.1	-
P11_S4	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_946189525.1	42441 to 48870 (contig 6)
P164_S40	N	-	-	-	-	-	-	-	-	GCA_946189105.1	-
P21_S10	Y	Y	7902	6	14	N	N	EMRKPDGALFNLFRRR	F	GCA_946189045.1	26201 to 34102 (contig 7)
P28_S15	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188915.1	120578 to 128595 (contig 6)
P42_S25	N	-	-	-	-	-	-	-	-	GCA_946189085.1	-
P55_S33	Y	Y	7359	5	16	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946189995.1	28034 to 35392 (contig 7)
P59_S35	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946189495.1	151368 to 161798 (contig 4)
P60_S36	N	-	-	-	-	-	-	-	-	GCA_946190025.1	-
S1071B24C1	Y	Y	2587	2	9	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221505.1	3951 to 6537 (contig 12)
S1071B24C2	Y	Y	2461	2	8	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221525.1	3630 to 6090 (contig 13)
S1071G24C1	Y	N	-	-	-	N	N	EMRKIDDALFNLFNFFRRR	F	GCA_905221535.1	47372 to 58002 (contig 11)
S1071G24C2	Y	N	-	-	-	N	N	EMRKIDDALFNLFNFFRRR	F	GCA_905221545.1	47372 to 58002 (contig 11)
S1071G24C3	Y	Y	2461	2	8	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221005.1	3630 to 6090 (contig 12)
S1071G24C4	N	-	-	-	-	-	-	-	-	GCA_905221035.1	-
S1071G24C5	N	-	-	-	-	-	-	-	-	GCA_905221045.1	-
S1072B24C4	Y	Y	2873	4	7	N	N	EMRMLKSPFSFLFPRKK	P	GCA_905221065.1	23060 to 25952 (contig 4)
S1072B24C5	Y	Y	2873	4	7	N	N	EMRMLKSPFSFLFPRKK	P	GCA_905221055.1	23060 to 25952 (contig 4)

(continues next page)

S1072G24C4	Y	N	-	-	-	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221105.1	26882 to 35329 (contig 7)
S1075M24C2	Y	N	-	-	-	Y	Y	ESRVADLLLDFLFQRKK	Q	GCA_905221085.1	103323 to 113219 (contig 6)
S1075M24C3	Y	N	-	-	-	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221095.1	31726 to 42537 (contig 11)
S1075M24C5	Y	Y	10812	2	26	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221075.1	27291 to 37102 (contig 8)
S1086G24C1	N	-	-	-	-	-	-		-	GCA_905221125.1	-
S1088B24C2	Y	Y	6473	2	18	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_905221135.1	26672 to 33144 (contig 6)
S1088M24C1	Y	N	-	-	-	N	N	EMRIADKIFFNLFKRR	F	GCA_905221145.1	54156 to 65018 (contig 7)
S1091B24C4	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_905221155.1	125261 to 134199 (contig 4)
S1092G24C1	Y	Y	1740	2	2	N	N	DMRISGSIRNLIFPRKKK	R	GCA_905221165.1	12912 to 14651 (contig 9)
S1092G24C3	Y	Y	1768	2	2	N	N	EIRKKSNSLLNFFKRR	F	GCA_905221175.1	13104 to 14871 (contig 5)
S1092G24C4	N	-	-	-	-	-	-		-	GCA_905221185.1	-
S1093G24C1	Y	Y	8698	4	17	N	Y	EMRKSNSTFFHFLRRI	L	GCA_905221195.1	27478 to 36175 (contig 5)
S1093G24C2	Y	N	-	-	-	N	N	ESRLPKILRDFIFPRKK	P	GCA_905221205.1	66159 to 73603 (contig 4)
S1093G24C3	Y	N	-	-	-	N	N	EMRKTSYSLLNLFKRR	F	GCA_905221225.1	104898 to 109878 (contig 7)
S1096G24C5	N	-	-	-	-	-	-		-	GCA_905221215.1	-
S1096M24C3	Y	Y	9580	5	22	N	N	EIRKTSNSLFNFFKRR	F	GCA_905221245.1	11901 to 21480 (contig 6)
S1096M24C5	N	-	-	-	-	-	-		-	GCA_905221285.1	-
U10_S50	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946188615.1	136100 to 146454 (contig 6)
U104_S93	Y	Y	9859	4	20	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946188685.1	23700 to 33558 (contig 6)
U105_S94	N	-	-	-	-	-	-		-	GCA_946188575.1	-
U109_S95	Y	N	-	-	-	N	N	EIRKTSNSLFNFFKRR	F	GCA_946188695.1	67998 to 77255 (contig 5)
U114_S96	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946188415.1	84542 to 94928 (contig 4)
U117_S1	Y	N	-	-	-	N	N	DSRISKYLRDFIFPRKK	P	GCA_946188585.1	15202 to 15765 (contig 7)
U12_S51	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946224775.1	52874 to 63087 (contig 6)
U122_S2	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946189145.1	149195 to 157067 (contig 2)
U123_S3	Y	N	-	-	-	N	N	ESRVSDILLNFLFRRKK	R	GCA_946190065.1	49685 to 50417 (contig 16)
U126_S5	Y	Y	8396	4	19	Y	Y	ESRVSDLILDFLFQRKK	Q	GCA_946190085.1	25538 to 25928 (contig 3)
U13_S52	Y	N	-	-	-	N	N	ESRLSKLLRDFIFKIKQ	K	GCA_946190075.1	21460 to 34396 (contig 12)
U131_S7	Y	N	-	-	-	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946190095.1	127239 to 129511 (contig 5)
U134_S8	Y	Y	1964	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_946190055.1	12955 to 14918 (contig 6)
U135_S9	Y	Y	1964	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_946190045.1	12955 to 14918 (contig 6)
U14_S53	N	-	-	-	-	-	-		-	GCA_946190385.1	-
U143_S13	Y	N	-	-	-	N	N	EIRKKSNSLLNFFKRR	F	GCA_946190395.1	13391 to 14114 (contig 3)
U144_S14	Y	N	-	-	-	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946190245.1	31975 to 35539 (contig 15)
U148_S15	Y	Y	1964	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_946190215.1	12955 to 14918 (contig 6)
U15_S54	Y	Y	3970	1	7	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946190375.1	44742 to 48711 (contig 14)
U16_S55	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946190205.1	52874 to 63087 (contig 9)
U164_S21	Y	N	-	-	-	N	N	DSRISKYLRDFIFPRKK	P	GCA_946190365.1	56909 to 62429 (contig 9)
U165-2_S22	N	-	-	-	-	-	-		-	GCA_946186005.1	-
U23_S57	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190355.1	34140 to 39308 (contig 18)
U24_S58	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190345.1	13215 to 18383 (contig 16)
U25_S59	N	-	-	-	-	-	-		-	GCA_946224995.1	-
U28_S60	N	-	-	-	-	-	-		-	GCA_946225005.1	-
U32_S62	Y	Y	6537	3	14	N	N	EMRRIDKIFINFLKRR	L	GCA_946224725.1	25517 to 32053 (contig 4)
U35_S65	Y	N	-	-	-	N	N	EMRRIDKIFINFLKRR	L	GCA_946224685.1	125817 to 132353 (contig 5)
U36_S66	N	-	-	-	-	-	-		-	GCA_946224705.1	-
U39_S69	N	-	-	-	-	-	-		-	GCA_946224645.1	-
U40_S70	Y	Y	1815	3	7	N	N	EIRKTINGLLNFFKRR	F	GCA_946190305.1	49454 to 51267 (contig 3)
U48_S72	N	-	-	-	-	-	-		-	GCA_946224865.1	-
U52_S74	N	-	-	-	-	-	-		-	GCA_946190225.1	-

(continues next page)

U54_S75	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190185.1	18384 to 28422 (contig 12)
U56_S76	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190325.1	71434 to 73220 (contig 5)
U57_S77	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190235.1	37034 to 38822 (contig 12)
U61_S78	N	-	-	-	-	-	-	-	-	GCA_946190115.1	-
U63_S79	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190175.1	37034 to 38822 (contig 15)
U65_S81	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190125.1	10058 to 15226 (contig 20)
U67_S83	Y	Y	7327	1	14	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946190165.1	35610 to 42936 (contig 11)
U8_S49	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946190315.1	52874 to 63087 (contig 6)
U92_S86	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190145.1	58367 to 59135 (contig 6)
U98_S90	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946190155.1	14745 to 15455 (contig 11)
U99_S91	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946190155.1	14745 to 15455 (contig 12)
WTCHG_774611_72895361	N	-	-	-	-	-	-	-	-	GCA_946190255.1	-
WTCHG_774611_72965368	Y	N	-	-	-	N	N	ESRVSDILLDFLFRRKK	R	GCA_946190195.1	114512 to 120897 (contig 4)
WTCHG_774611_73005372	Y	Y	12164	2	23	N	N	EMRKMNDKSFNLFNFFRRR	F	GCA_946190135.1	24172 to 36335 (contig 6)
WTCHG_774611_73015373	N	-	-	-	-	-	-	-	-	GCA_946190275.1	-
WTCHG_774611_73075379	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946190295.1	119791 to 129649 (contig 6)
WTCHG_774611_73095381	Y	N	-	-	-	N	N	EMRKMNDKSFNLFNFFRRR	F	GCA_946190335.1	118632 to 130795 (contig 6)
WTCHG_774611_73775353	Y	Y	11984	5	25	N	N	EIRLSKFLRDFILQRKK	Q	GCA_946190285.1	28199 to 40182 (contig 5)
WTCHG_774611_73825358	Y	Y	11984	5	25	N	N	EIRLSKFLRDFILQRKK	Q	GCA_946190255.1	28199 to 40182 (contig 5)
WTCHG_774611_73845360	Y	N	-	-	-	N	N	EMRKMNDKSFNLFNFFRRR	F	GCA_946185975.1	118632 to 130795 (contig 6)
B22	Y	Y	15566	11	38	N	Y	ESRVSRIILDFLFLRKK	L	GCA_965111745.1	61443 to 77008 (contig 1)
C22	Y	Y	7274	3	15	N	Y	ESRLSRLLRDFIFQIKQ	Q	GCA_965111725.1	50761 to 58034 (contig 1)
D22	Y	Y	2127	2	2	N	N	DSRISKFLRDFIFPRKK	P	GCA_965111735.1	34443 to 36569 (contig 1)
E22	Y	Y	14434	1	32	Y	Y	EMRLPKILRDFIFPRKK	P	GCA_965111705.1	263852 to 278285 (contig 1)
F22	Y	Y	7848	5	13	Y	Y	ESRVSRIILDFLFLRKK	L	GCA_965111715.1	42896 to 50743 (contig 1)
G22	Y	Y	4404	2	12	N	Y	ESRLSRLLRDFIFQIKQ	Q	GCA_965111695.1	57304 to 61707 (contig 1)
B6	Y	Y	9068	5	18	Y	Y	EMRKPDGALFNLFRRR	F	GCA_000027165.1	53854 to 62921 (contig 1)
FDA684	Y	Y	8195	5	17	N	N	EIRQTHNIFFNFFKRR	F	GCA_009730515.1	31493 to 39687 (contig 1)
SK637	Y	Y	7760	4	18	N	Y	EMRKSNNNFFHFLRRI	L	GCA_000722765.2	49608 to 57367 (contig 1)
SVGS061	Y	Y	2200	2	5	N	N	EMRKSNNTFFNFLRRI	L	GCA_001560895.1	73674 to 75873 (contig 1)

The genomic region between *plsX-acpP* and *comAB* was extracted from publicly available *S. mitis* genomes. Of the 332 genomes analyzed, this region was successfully retrieved from 253; the remaining genomes contained contig breaks and were excluded from further analyses. The 253 genomes were screened for the presence of competence-associated bacteriocin (*cab*), competence-induced bacteriocins A and B (*cibAB*), and the cognate immunity protein (*cibC*) genes. The number of SigX boxes in *cab* loci was determined. Mature CSP sequences of each strain was predicted by identifying small ORFs upstream of *comDE* that contained a double-glycine cleavage motif.

Table S2. Bacterial strains, plasmids, peptides and oligonucleotides used in the study.

Bacterial strain, plasmid, oligonucleotide or peptide	Relevant genotype or sequence	Note	Source
Bacterial strains			
Corynebacterium accolens LMG 19042	Wild type	-	BCCM/LMG
Dolosigranulum pigrum LMG 15126	Wild type	-	BCCM/LMG
Escherichia coli DH5α	Wild type	-	Thermo Fisher
Escherichia coli DH10B	Wild type	-	Thermo Fisher
Haemophilus influenzae FP7132	Wild type		P. Lavado
Klebsiella pneumoniae OND482	Wild type	-	Simões *et al*., 2022
Moraxella catarrhalis ATCC 25238	Wild type	-	ATCC
Neisseria lactamica ATCC 23970	Wild type	-	ATCC
Staphylococcus aureus ATCC 25293	Wild type	-	ATCC
Staphylococcus epidermidis ATCC 14990	Wild type	-	ATCC
Streptococcus agalactiae ATCC 13813	Wild type	-	ATCC
Streptococcus anginosus ATCC 12395	Wild type	-	ATCC
Streptococcus constellatus ATCC 27823	Wild type	-	ATCC
Streptococcus gordonii ATCC 33399	Wild type	-	ATCC
Streptococcus infantis ATCC 700779	Wild type	-	ATCC
Streptococcus mitis SK142	Wild type	-	ATCC
Streptococcus mitis C22	Wild type	-	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis C22 Δcab	Δ*cab* ::*aphA-3*	kan^R	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis C22 kan^R	*ZIP::aphA-3*	kan^R	This study
Streptococcus mitis F22	Wild type	-	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis G22	Wild type	-	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis G22 P*comCDE*-pJB11	*CIL* ::P*comCDE* -mNeonGreen-luciferase *cat*	chl^R	This study
Streptococcus mitis G22 P*sigX1*-pJB11	*CIL::PsigX1-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 P*sigX2*-pJB11	*CIL::PsigX2-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 PcabABC-pJB11	*CIL::PcabABC-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 PcabABC (truncated)-pJB11	*CIL::P cabABC(truncated)-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 Δ*comE* PcabABC-pJB11	Δ*comE::aphA-3 CIL::PcabABC-mNeonGreen-luciferase cat*	kan^R chl^R	This study
Streptococcus mitis G22 Δ*sigX1* PcabABC-pJB11	Δ*sigX1::aphA-3 CIL::PcabABC-mNeonGreen-luciferase cat*	kan^R chl^R	This study
Streptococcus mitis G22 Δ*sigX2* PcabABC-pJB11	Δ*sigX2::ant(9) CIL::PcabABC-mNeonGreen-luciferase cat*	spc^R chl^R	This study
Streptococcus mitis G22 Δ*sigX1* Δ*sigX2* PcabABC-	Δ*sigX1* ::*aphA-3* Δ*sigX2* ::*ant(9) CIL* ::PcabABC-mNeonGreen-luciferase cat*	kan^R spc^R chl^R	This study
Streptococcus mitis G22 Δ*blp4* Δ*blp1*	Δ*blp4* ::*lox72* Δ*blp1::aphA-3*	kan^R	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis G22 Δ*blp4* Δ*blp1* (markerless)	Δ*blp4* ::*lox72* Δ*blp1::lox72*	-	This study
Streptococcus mitis Δ*blp4* Δ*blp1* Δ*cabABC*	Δ*blp4* ::*lox72* Δ*blp1::lox72* Δ*cabABC* ::*aphA-3*	kan^R	This study
Streptococcus mitis G22 p(*cab* $^+$)	Δ*blp4* ::*lox72* Δ*blp1::lox72* Δ*cabABC* ::*aphA-3* pJB09-P23-*cabABC*	spc^R kan^R	This study

(continues next page)

Streptococcus mitis G22 *cabA-HA*	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*cabABC*::*aphA-3* CIL::*cabA-HA cabBC cat*	kan^R chl^R	This study
Streptococcus mitis G22 *cabB-HA*	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*cabABC*::*aphA-3* CIL::*cabA cabB-HA cabC cat*	kan^R chl^R	This study
Streptococcus mitis G22 Δ*blp4* Δ*blp1* Δ*comE*	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*comE*::*aphA-3*	kan^R	This study
Streptococcus mutans ATCC 25175	Wild type	-	ATCC
Streptococcus oralis DSM 20066	Wild type	-	DSMZ
Streptococcus parasanguinis ATCC 15912	Wild type	-	ATCC
Streptococcus peroris ATCC 700780	Wild type	-	ATCC
Streptococcus pneumoniae D39	Wild type	-	Avery *et al.*, 1944
Streptococcus pneumoniae D39 PhunSweet	Δ*cps*::*pheS-sacB-aphA-3*	kan^R	Echlin and Rosch, 2020
Streptococcus pneumoniae D39 P*comCDE*-pJB11	CIL::P*comCDE*-mNeonGreen-luciferase *cat*	chl^R	This study
Streptococcus pneumoniae D39 Δ*lytA*	Δ*lytA*::*pheS-sacB-aphA-3*	kan^R	This study
Streptococcus pneumoniae D39 Δ*lytA* (markerless)	Δ*lytA*	-	This study
Streptococcus pneumoniae D39 Δ*lytC*	Δ*lytC*::*pheS-sacB-aphA-3*	kan^R	This study
Streptococcus pneumoniae D39 Δ*lytC* (markerless)	Δ*lytC*	-	This study
Streptococcus pneumoniae D39-GFP	P*hlpA*-*hlpA-gfp cat*	chl^R	Valente *et al.*, 2022
Streptococcus pneumoniae D39-GFP ΔcomGC	P*hlpA*-*hlpA-gfp cat* Δ*comGC*::*tet(M)*	tet^R	This study
Streptococcus pneumoniae TIGR4	Wild type	-	ATCC
Streptococcus pseudopneumoniae BAA-960	Wild type	-	ATCC
Streptococcus pyogenes ATCC 12344	Wild type	-	ATCC
Streptococcus salivarius ATCC 7073	Wild type	-	ATCC
Streptococcus sanguinis ATCC 10556	Wild type	-	ATCC
Streptococcus sinensis HKU4	Wild type	-	DSMZ
Streptococcus sobrinus ATCC 27351	Wild type	-	ATCC
Streptococcus vestibularis ATCC MM1	Wild type	-	ATCC
Plasmids			
pDL278-P23-DsRed-Express2	*P23-dsRed-Express2 pMB1 pVA380-1 ant(9)*	*E. coli*-streptococcal	Plasmid #121505 (Addgene)
pJB09-cabABC	pDL278-P23-DsRed-Express2 but Δ*dsRed-Express2*::*cabABC*	*E. coli*-streptococcal	This study
pBIR	*p15A aphA-3 tdTomato cat*	Integrates in *CIL* locus	This study (Twist Bioscience)
pPEP-LGZ	*luc sfGFP lacZ cat p15A ant(9)*	Integrates in *CEP* locus	Kjos *et al*., 2016
pJB10	pBIR but Δ*tdTomato*::*hlpA-mNeonGreen*	Integrates in *CIL* locus	This study
pJB11	pJB10 but *luc*	Integrates in *CIL* locus	This study
pJB11-PcomCDE(G22)	pJB11 but P*comCDE* reporter (*S. mitis* G22)	Integrates in *CIL* locus	This study
pJB11-PcomCDE(D39)	pJB11 but P*comCDE* reporter (*S. pneumoniae* D39)	Integrates in *CIL* locus	This study
pJB11-PsigX1	pJB11 but P*sigX1* reporter	Integrates in *CIL* locus	This study
pJB11-PsigX2	pJB11 but P*sigX2* reporter	Integrates in *CIL* locus	This study

(continues next page)

Table S2. (cont.)

pJB11-PcabABC	pJB11 but P*cabABC* reporter	Integrates in *CIL* locus	This study
pJB11-PcabABC(truncated)	pJB11 but P*cabABC* (truncated) reporter	Integrates in *CIL* locus	This study
pKan	*pMB1 lox66-p3-aphA-3-lox71*	Integrates in *cab* locus	Borralho, Handem and Lança *et al.*, 2025
Peptides			
BlpC$_1$	GLWEELYETITTPLKLRTTFKP	-	NZYTech
BlpC$_4$	GWWEELLHETILSKFKITKALELPVQL	-	NZYTech
CSP-1	EMRLSKFFRDFILQRKK	-	NZYTech
CSP-2	EMRISRIILDFLFLRKK	-	NZYTech
CSP-2(L14Q)	EMRISRIILDFLFQRKK	-	NZYTech
CSP$_{A22}$	DWRISETIRNLIFPRRK	-	NZYTech
CSP$_{B22}$	ESRVSRIILDFLFLRKK	-	NZYTech
CSP$_{D22}$	DSRISKFLRDFIFPRKK	-	NZYTech
CSP$_{E22}$	EMRLPKILRDFIFPRKK	-	NZYTech
CSP$_{G22}$	ESRLSRLLRDFIFQIKQ	-	NZYTech
CSP$_{G22}$(Q14L)	ESRLSRLLRDFIFLIKQ	-	NZYTech
CSP$_{G22}$(Q14N)	ESRLSRLLRDFIFNIKQ	-	NZYTech
Oligonucleotides			
P1F_cab7_OE	ATGACAAATTTTGACAAAATGGAAC	Construction of plasmid pJB09-cabABC	This study
P1R_cab7_OE	CCCTATAGAACGTCTACTACCTC		
P1F_pDL278_OE	CTCGGTACCAAATTCCAGAAAAG		
P1F_pDL278_OE	GTTTGTCCTCCTTAACTAGTAATTAATAC		
P1F_hlpA_mNG_twist	AAAGCACTCAAAAGTTTACCTTATGG	Construction of plasmid pJB10	This study
P1R_hlpA_mNG_twist	CGCAAAAAACCCCGCTTCGGCGGGGTTTTTTCGCTTATTTATAC		
P1F_pBIR_backbone	ATCGGAGAAATTGGACACACC		
P1R_pBIR_backbone	CCCTATCAAAAGAATCTTGCTTGGC		
PF_luc_pPEP	GCTGAAGGAGGAATAATGAGATCC	Construction of plasmid pJB11	This study
PR_luc_pPEP	TTACAATTTGGGCTTTCCGCC		
PF_pJB10_backbone	GAATGGACGAGTTGTATAAG		
PR_pJB10_backbone	TTATTTATACAATTCATCCATACCC		
PF_reporter_PcomCDE(G22)	AGCTGGTCTCTGACTATATAGTAGCAAAGCTGGG	Construction of plasmid pJB11-PcomCDE (G22)	This study
PR_reporter_PcomCDE(G22)	CGATGGTCTCAAGTCAATAAAATCTCCTAAAATGTTTTTTTC		
PF_reporter_PcomCDE(D39)	AGCTGGTCTCTGACTATATAATAGCAAAGCTGGG	Construction of plasmid pJB11-PcomCDE (D39)	This study
PR_reporter_PcomCDE(D39)	CGATGGTCTCAAGTCAATAAAATCTCCTAAAATGTTTTTTC		
PF_reporter_PsigX1	AGCTGGTCTCTGACTCCCAAAGAGAGGCAAGACC	Construction of plasmid pJB11-PsigX1	This study
PR_reporter_PsigX1	CGATGGTCTCAAGTCTCTTTTCCCCTTTCTGTGTAC		

Table S2. (cont.)

PF_reporter_PsigX2	AGCTGGTCTCTGACTCCCTGAGAGAGGCCCGTC	Construction of plasmid pJB11-PsigX2	This study
PR_reporter_PsigX2	CGATGGTCTCAAGTCAGTCTTCCCCTTTCTGTGTAC		
PF_reporter_PcabABC	AGCTGGTCTCTGACTAAATAACATCTTCACAACTAATCG	Construction of plasmid pJB11-PcabABC	This study
PR_reporter_PcabABC	CGATGGTCTCAAGTCCTTTTTTCTCCTTTATTTCTATTTTTTAG		
PF_reporter_PcabABC_trunc	AGCTGGTCTCTGACTTTTAGTTGTGTATTTTCGAATTG	Construction of plasmid pJB11-PcabABC (truncated)	This study
PR_reporter_PcabABC_trunc	CGATGGTCTCAAGTCCTTTTTTCTCCTTTATTTCTATTTTTTAG		
P1F_comE_G22	GGATTAAACAGTATGAGAGG	Deletion of *comE* in *S. mitis* G22	This study
P1F_comE_G22	GTCAATACCGTTCGTATAGCATACATTATACGAAGTTATTCATTCAAATTCCCTCTTAAATC		
P2F_comE_G22_pKan	CAAATTATTAGATTTAAGAGGGAATTTGAATGAATAACTTCGTATAATGTATGC		
P2R_comE_G22_pKan	CATTATATCAATTTCTTGCTAATTGTCAATCACTATAACTTCGTATAGCATAC		
P3F_comE_G22	GTTTTAGTACCGTTCGTATAATGTATGCTATACGAAGTTATAGTGATTGACAATTAGCAAGAAATTG		
P3R_comE_G22	TATTACCATTATAGGAAGCC		
P4F_comE_G22	TATAACAGGAGATAATTTTG		
P4R_comE_G22	ACAGAGCTGGAATAAGAGAAATCC		
P1F_sigX1_G22	ATTGTCATCGCTGCGACAAAC	Deletion of *sigX1* in *S. mitis* G22	This study
P1R_sigX1_G22	GTCAATACCGTTCGTATAGCATACATTATACGAAGTTATCATTCTTTTCCCCTTTCTGTGTACAG		
P2F_sigX1_G22_pKan	GACTGTACACAGAAAGGGGAAAAGAATGATAACTTCGTATAATGTATGC		
P2R_sigX1_G22_pKan	CTTTTTTGCATGAATTCCTAATGGGTACGATAACTTCGTATAGCATAC		
P3F_sigX1_G22	GTTTTAGTACCGTTCGTATAATGTATGCTATACGAAGTTATCGTACCCATTAGGAATTCATGC		
P3R_sigX1_G22	CGCTACACATGGAATTCCAC		
P4F_sigX1_G22	AGCAATCTTGAAAGTTCATGCTAAG		
P4R_sigX1_G22	ATCCGGACAACGCTCGGGACCTAC		
P1F_sigX2_G22	TGCTTGTCAAAGCCTGGCTTTCTTG	Deletion of *sigX2* in *S. mitis* G22	This study
P1F_sigX2_G22	CATGCAAAACAATGAATAGGCATAGTCTTCCCCTTTCTGTGTAC		
P2F_sigX2_G22_pPEP	GACTGTACACAGAAAGGGGAAGACTATGCATGCAAAACAATGAATAGG		
P2R_sigX2_G22_pPEP	CTTTTTTGCATGAATTCCTAATGAGTACGCTAATTGAGAGAAGTTTCTATAG		
P3F_sigX2_G22	CTAATTGAGAGAAGTTTCTATAGCGTACTCATTAGGAATTCATGCAAA		
P3R_sigX2_G22	CCGCTACACATGGAATTCCACTC		
P4F_sigX2_G22	CTATTCAATCAAGATTAGTGGAAAC		
P4R_sigX2_G22	TAGTTAGCCGTCCCTTTCTGGTAAG		

(continues next page)

Table S2. (cont.)

P1F_PS_lytA_D39	GGTTTATCAGCGGTTCTCTGC		
P1R_PS_lytA_D39	CGGATCGATCCTTAATAGTGGGTTCCATATTCTACTCCTTATCAA		
P2F_PS_lytA_D39	GTTTTAATTGATAAGGAGTAGAATATGGAACCCACTATTAAGGATCGATC		
P2R_PS_lytA_D39	CTGTTCTGATTTGAAAGACATTCCATTATTACTAAAACAATTCATCCAGTAAAATATAAT	Deletion of *lytA* in *S. pneumoniae D39*	This study
P3F_PS_lytA_D39	CTGGATGAATTGTTTTATAATAATGGAATGTCTTTCAAATCAG		
P3R_PS_lytA_D39	TTTTACTTTCTGACAAAGGACG		
P4F_PS_lytA_D39	TTCAATTGTATCTATCGGCAG		
P4R_PS_lytA_D39	TTCCTCAATCTATATAACATAGCTTTATG		
P5F_PS_lytA_D39	GTTTTAATTGATAAGGAGTAGAATATGGAATAATAATGGAATGTCTTTCAAATCAG		
P5R_PS_lytA_D39	CTGTTCTGATTTGAAAGACATTCCATTATTATTCCATATTCTACTCCTTATCAA		
P1F_PS_lytC_D39	ACTAAAGGAAGATGGAACCAG		
P1R_PS_lytC_D39	CGGATCGATCCTTAATAGTGGGAGACATCTGAATGCTGTTCC		
P2F_PS_lytC_D39	CTAGGTGGAACAGCATTCAGATGTCTCCCACTATTAAGGATCGATC		
P2R_PS_lytC_D39	CACATCCCTCTTTCAAATCATCGCTTACTAAAACAATTCATCCAGTAAAATATAAT	Deletion of *lytC* in *S. pneumoniae D39*	This study
P3F_PS_lytC_D39	CTGGATGAATTGTTTTATAAGCGATGATTTGAAAGAGGG		
P3R_PS_lytC_D39	TTCTTAGTCATGCTTCCATCC		
P4F_PS_lytC_D39	CTCAAATTGAGGCCAAGAGAG		
P4R_PS_lytC_D39	GTCCAAGTTCGGCTGCAAG		
P5F_PS_lytC_D39	CTAGGTGGAACAGCATTCAGATGTCTTAAGCGATGATTTGAAAGAGGG		
P5R_PS_lytC_D39	CACATCCCTCTTTCAAATCATCGCTTAAGACATCTGAATGCTGTTCC		
P1F_comGC_del	AATCGTTGACTTTGCAAGCAG		
P1R_comGC_del	TAATTTTCCTCCTATTTAGATCTTGCATGTATAGTTATTATACCAGGGGGACAGTGCAATGTCAATTTAAAAATTTACCTCCATATTTGATACATG		
P2F_comGC_del	TTGACATTGCACTGTCCCCCTGGTATAATAACTATACATGCAAGATCTAAATAGGAGGAAAATTAATGAAAATTATTAATATTGGAGTTTTAGC		
P2R_comGC_del	CATTGACTTTACGATTTGCTCCTCCATTCTAAGTTATTTTATTGAACATATATCG	Deletion of *comGC* in *S. pneumoniae D39*	This study
P3F_comGC_del	GTACGATATATGTTCAATAAAATAACTTAGAATGGAGGAGCAAATCGTAAAGTC		
P3R_comGC_del	GTGACTGCGTAGAGGAGAAC		
P4F_comGC_del	TTCAGGCTGAGACGGAAAAAATTAG		
P4R_comGC_del	CAGATTTGAGTCCATAAACC		
P1F_cabAB_HA	CAAGCGCCAGGGTTTTCC		
P1R_cabAB_HA	GTTCTGTTCCATTTTGTCAAAATTTGTCATCTTTTTTCTCCTTTATTTCTATTTTTTAG		
P2F_cabAB_HA	GTAAACTAAAAAATAGAAATAAAGGAGAAAAAAGATGACAAATTTTGACAAAATGGAAC		
P2F_cabAB_HA	CCAGGGGGACAGTGCAATGTCAATTATCCATTCTTTTTTTCTTTACTTC	Construction of *S. mitis* harboring HA-tagged bacteriocins	This study
P3F_cabAB_HA	CATTCTTTTATATAGAAGTAAAGAAAAAAAGAATGGATAATTGACATTGCACTGTCCCC		
P3F_cabAB_HA	GCCCTCTGAGCGGATAACAATTTC		
P4F_cabAB_HA	ATCGGAGAAATTGGACACACC		
P4R_cabAB_HA	TTCGCAAATAGCGAAACCAATG		

(continues next page)

Table S2. (cont.)

P1F_C22_kan_ZIP	GCTGAACAACAGTGTGTCATA		
P1R_C22_kan_ZIP	CAGGGGGACAGTGCAATGTCAATAGCTCAATTTGACACTTTTCTTC		
P2F_C22_kan_ZIP	CGAAGAAGAAAAGTGTCAAATTGAGCTATTGACATTGCACTGTCC		
P2R_C22_kan_ZIP	CTATGCAGGTCATTTCTACTCTTATAGCTAAAACAATTCATCCAGTAAAATA	Construction of *S. mitis* C22 mutant with kanamycin resistance	This study
P3F_C22_kan_ZIP	TATTATATTTTACTGGATGAATTGTTTTAGCTATAAGAGTAGAAATGACCTGC		
P3R_C22_kan_ZIP	TCCCTTCCTATCAATTAGGTC		
P4F_C22_kan_ZIP	AGATAGATCAGCTATTGGAGAC		
P4R_C22_kan_ZIP	TATTTAAGTCCATCATAGCTTGG		

chl^R - chloramphenicol resistant; kan^R - kanamycin resistant; tet^R - tetracycline resistant; spc^R - spectinomycin resistant.

Supplementary references

Simões AS, Touret T, Faria NA, Peres Ladeiro S, Costa J, Bispo S, Serrano M, Palos C, Miragaia M, Bastos Leite R, Sá-Leão R. Using whole genome sequencing to investigate a mock-outbreak of carbapenem-resistant *Klebsiella pneumoniae* in real-time. *Acta Med Port*. 2022 Jan 3;35(1):36–41. doi: 10.20344/amp.15174. Epub 2021 Nov 10. PMID: 34755594.

Borralho J, Handem S, Lança J, Ferreira B, Candeias C, Henriques AO, Hiller NL, Valente C, Sá-Leão R. Inhibition of pneumococcal growth and biofilm formation by human isolates of *Streptococcus mitis* and *Streptococcus oralis*. *Appl Environ Microbiol*. 2025 Mar 19;91(3):e0133624. doi: 10.1128/aem.01336-24. Epub 2025 Feb 26. PMID: 40008876; PMCID: PMC11921387.

Avery OT, Macleod CM, McCarty M. Studies on the chemical nature of the substance inducing transformation of pneumococcal types: induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III. *J Exp Med*. 1944;79:137–158.

Echlin H, Rosch JW. Advancing genetic tools in *Streptococcus pneumoniae*. *Genes (Basel)*. 2020 Aug 20;11(9):965. doi: 10.3390/genes11090965. PMID: 32825523; PMCID: PMC7563404.

Valente C, Cruz AR, Henriques AO, Sá-Leão R. Intra-species interactions in *Streptococcus pneumoniae* biofilms. *Front Cell Infect Microbiol*. 2022 Jan 5;11:803286. doi: 10.3389/fcimb.2021.803286. PMID: 35071049; PMCID: PMC8767070.

Kjos M, Miller E, Slager J, Lake FB, Gericke O, Roberts IS, Rozen DE, Veening JW. Expression of *Streptococcus pneumoniae* bacteriocins is induced by antibiotics via regulatory interplay with the competence system. *PLoS Pathog*. 2016 Feb 3;12(2):e1005422. doi: 10.1371/journal.ppat.1005422. PMID: 26840404; PMCID: PMC4739728.