Streptococcus mitis* bacteriocins drive contact-dependent lysis of *S.

***pneumoniae* facilitating transformation in multispecies environments**

João Borralho[1], João Lança,[1], Joana Bryton[1], Wilson Antunes[2], Raquel Sá-Leão[1]*

[1] Laboratory of Molecular Microbiology of Human Pathogens, Instituto de Tecnologia Química e

Biológica António Xavier, Universidade Nova de Lisboa, Oeiras, Portugal

[2] Centro de Investigação da Academia Militar (CINAMIL), Unidade Militar Laboratorial de Defesa

Biológica e Química (UMLDBQ), Av. Dr. Alfredo Bensaúde, 1849-012, Lisboa, Portugal.

* Corresponding author: rsaleao@itqb.unl.pt

Author Contributions: R. Sá-Leão and J. Borralho contributed to the concept and design of the

study. R. Sá-Leão contributed with reagents and materials. Data acquisition was performed by J.

Borralho, J. Lança, J. Bryton and W. Antunes. Data was interpreted by J. Borralho, J. Lança, J. Bryton,

W. Antunes and R. Sá-Leão. The manuscript was drafted by J. Borralho and R. Sá-Leão. All authors

critically revised and approved the final version of the manuscript.

Competing Interest Statement: Universidade NOVA de Lisboa has filed a provisional patent

application that covers pharmaceutical compositions comprising strain G22 and/or bacteriocin

molecules, and derivatives thereof, which can inhibit the growth and persistence of the pathogen *S.*

pneumoniae (PT119647).

26 **Keywords:** Horizontal gene transfer, competence, interspecies interactions, predation, biofilms

Abstract

Natural competence allows bacterial species like *Streptococcus pneumoniae* and *S. mitis* to acquire environmental DNA, driving horizontal gene transfer (HGT) and adaptation. In *S. pneumoniae*, a human pathogen, competence-induced predation is well characterized and involves the release of bacteriocins and a murein hydrolase to lyse noncompetent siblings and liberate DNA. In contrast, in the human commensal *S. mitis*, mechanisms mediating DNA acquisition remain poorly understood. Here, we identify a diverse set of competence-associated bacteriocins (*cab*) that are produced by *S. mitis* during the late phase of competence. We focus on one bacteriocin pair, CabAB, that triggers contact-dependent growth inhibition and lysis of *S. pneumoniae* through activation of the major pneumococcal autolysin LytA. We demonstrate that CabAB compromises *S. pneumoniae* membrane integrity, leading to formation of intracellular membrane aggregates and the release of cytoplasmatic content, thereby increasing available DNA, which enhances HGT from *S. pneumoniae* to *S. mitis* in biofilms. These findings uncover a mechanism of interspecies predation and gene acquisition, revealing a critical role for competence-associated bacteriocins in shaping evolutionary dynamics of streptococci.

Significance Statement

Many streptococci are naturally competent, acquiring environmental DNA through transformation. This includes pathogens like *S. pneumoniae* and commensals like *S. mitis,* which can exchange genetic material through horizontal gene transfer (HGT). For example, *S. mitis* can acquire pneumococcal capsules, leading to its misidentification in polymicrobial samples such as those obtained from the upper respiratory tract. Understanding the drivers of HGT between these species is therefore critical. Here, we characterize a competence-induced bacteriocin cluster in *S. mitis*. These bacteriocins lyse pneumococci, promoting DNA release and enhancing gene transfer in dual-species biofilms. Our findings uncover a mechanism by which competence-associated predation promotes interspecies HGT, shaping the evolution and epidemiology of streptococcal populations.

Main Text

Introduction

Natural competence for genetic transformation is a developmental program found in certain bacterial species that enables the uptake of naked DNA from the environment. Once internalized, this DNA can be incorporated into the genome through homologous recombination, making natural transformation one of the main modes of horizontal gene transfer (HGT). This phenomenon was first described in *Streptococcus pneumoniae* (or pneumococcus), a species capable of exchanging pathogenic traits such as the capsule by taking up DNA from other pneumococci (1).

In *S. pneumoniae*, competence is a tightly regulated transient physiological state which is governed by a cell-cell signaling system. The competence-stimulating peptide (CSP), encoded by *comC*, is processed and exported via the ATP-binding cassette transporter ComAB. Extracellular CSP is sensed by the membrane-bound histidine kinase ComD (2-4), which undergoes autophosphorylation triggering subsequent phosphorylation of the response regulator ComE (5). Phosphorylated ComE increases expression of the early competence cascade including *comAB* and *comCDE*, creating a positive feedback loop. It also induces expression of the two copies of the alternative sigma factor (*sigX1* and *sigX2*), marking the onset of the late phase of competence (6).

In the late phase of competence, expression of all genes required for DNA uptake, processing and recombination – the transformasome – is induced (7). Here, pneumococci produce the competence-induced bacteriocins A and B (CibAB), along with the immunity protein CibC, which provides self-immunity to CibAB (8). Because competence is a bistable process, where only a subset of the population becomes competent at any given time, CibAB release on solid media leads to lysis of genetically identical (kin) noncompetent cells that have not yet expressed *cibC*. Additionally, expression of the murein hydrolase CbpD, against which competent cells are protected via ComM expression, is essential for kin predation in liquid cultures. Notably, CbpD has also been implicated in lysis of non-kin, closely related streptococci, thereby expanding the gene pool from which *S. pneumoniae* may acquire DNA (9). CibAB and CbpD have also been associated to competitive

interactions *in vivo*, where their expression by colonizing pneumococci provides a competitive

advantage over other *S. pneumoniae* invaders (10).

A close relative of the pneumococcus, *S. mitis*, is also naturally transformable. *S. mitis* is a

common commensal of the human nasopharynx and oral cavity, where it typically establishes long-

term colonization without causing disease (11). Despite close phylogenetic relationship with *S.*

pneumoniae, and a strikingly similar competence regulatory network, *S. mitis* exhibits key differences

in terms of bacteriocin content. While *cibAB* is highly conserved in *S. pneumoniae*, most *S. mitis* lack

cibAB, suggesting that *S. mitis* may rely on an alternative mechanism for predation during

competence (12, 13).

Although the frequency at which *S. mitis* acquires DNA from *S. pneumoniae* is not well

established, such events have been documented. These includes the acquisition of pneumococcal

pbp mosaic blocks (14, 15), as well as the expression of pneumococcal or pneumococcal-like

capsules by certain *S. mitis* strains (16-20).

A competence-associated bacteriocin (*cab*) cluster, which is absent in *S. pneumoniae*, has

been identified in *S. mitis*. This cluster is located upstream of the competence transporter *comAB* and

can encode multiple *cab* bacteriocins distinct from CibAB (12, 13, 21-23). However, the role of *cab* in

S. mitis predation of kin and non-kin, and its contribution to DNA acquisition, remains unexplored.

Here, we characterize the arsenal of *cab* bacteriocins carried by *S. mitis* and provide evidence

that some specifically mediate lysis of *S. pneumoniae*. We also show that pneumococcal DNA can

be efficiently taken up and recombined by *S. mitis* and that Cab-mediated lysis accelerates gene

transfer from *S. pneumoniae* to *S. mitis*. Together, these findings shed light on previous observations

of pneumococcal DNA integration in *S. mitis* genomes.

Results

***S. mitis* encodes a diverse arsenal of competence-associated bacteriocins (*cab*)**

We previously identified and characterized a limited number of *S. mitis* strains encoding multiple

competence-associated bacteriocins (*cab*), most of which lacked the pneumococcal bacteriocins

cibAB and associated immunity protein *cibC* (23). To investigate the broader distribution of *cab* and *cibAB* in *S. mitis*, we surveyed publicly available genomes, leveraging the conserved genomic context of *cab* clusters – located upstream of *comAB* and downstream of *plsX-acpP*. This region was identified and extracted from 253 of 332 *S. mitis* genomes (Fig. 1A and *SI Appendix*, Table S1). The remaining genomes had contig breaks and were excluded from further analysis.

Among the 253 genomes analyzed, 26.9% harbored *cibABC*, a striking contrast with *S. pneumoniae*, where these genes are universally present. An additional 13.0% of strains carried *cibC* but not *cibAB* – often referred to as "cheater" phenotype (24), in which strains are immune to CibAB-mediated killing but do not produce the bacteriocins themselves. In contrast, 47.4% of *S. mitis* encoded between one to eleven unique *cab* genes. In total, 95 distinct Cab variants were found, which grouped into 28 clusters based on sequence similarity (Fig. 1B and *SI Appendix*, Fig. S1). Overall, 61.3% of *S. mitis* genomes carried at least one *cab* gene or the *cibABC* operon (*SI Appendix*, Table S1).

Cab variants were present at low frequencies (<10%) in the *S. mitis* population, with 32 of 95 bacteriocin variants only being found in a single strain (Fig. 1B). Despite this extensive diversity, rarefaction analysis suggested that Cab diversity is not yet saturated (Fig. 1C); sequencing more *S. mitis* will likely reveal new Cab sequences.

These findings highlight *S. mitis* as a reservoir of highly diverse and largely unexplored competence-associated bacteriocins.

Cab bacteriocins are expressed in the late phase of competence

S. mitis G22, previously described by us, encodes a *cab* operon consisting of two putative bacteriocins and an associated immunity protein (hereafter referred as *cabAB* and *cabC*, respectively) (Fig. 2A) (23). This strain was selected to study *cab* regulation in *S. mitis* and its role in competitive behaviors. Both bacteriocins and the immunity protein it encodes are rare (25) and thus unlikely to encounter pre-existing immunity in other strains. Notably, its *cab* promoter contains five putative binding sites for the alternative sigma factors (SigX boxes) (23), suggesting that *cabABC* may be expressed during the late phase of competence.

To investigate the dynamics of *cab* expression, the native *cabABC* promoter was ectopically

inserted into the *S. mitis* G22 genome, upstream of a firefly luciferase reporter (*luc*). Expression of

cabABC was then inferred by expression of *luc* and compared to known early (*comCDE*) and late

(*sigX1* and *sigX2*) competence regulators. As expected, we observed that competence was not

induced under acidic pH (*SI Appendix*, Fig. S2), unless exogenous cognate competence-stimulating

peptide (CSP$_{G22}$) was supplied (*SI Appendix*, Fig. S3A). Upon CSP$_{G22}$ treatment, *cabABC* expression

followed a similar temporal pattern to *comCDE*, *sigX1* and *sigX2*, peaking at approximately 70 min

post-induction (*SI Appendix*, Fig. S3B). This response was specific to CSP, as exogenous addition of

cognate bacteriocin-like peptide pheromones (BlpC$_1$ and BlpC$_4$), encoded elsewhere in the *S. mitis*

G22 genome (25), failed to induce *cabABC* expression (*SI Appendix*, Fig. S4).

Deletion of the early regulator *comE* abolished *cabABC* expression, a result expected for both

early- and late-regulated genes (Fig. 2B). Individual deletion of *sigX1* or *sigX2* partially reduced

expression of *cabABC*, while deletion of both resulted in complete loss of *cabABC* expression,

consistent with *cab* being expressed during the late phase.

Although multiple SigX boxes have been observed in *cab* promoters previously (13, 23), their

functional relevance has not been tested. To assess this, we compared expression driven by the full-

length *cabABC* promoter (containing all five SigX boxes) to a truncated version containing only the

most proximal SigX box. Deletion of four out of five SigX boxes reduced *cabABC* expression (Fig.

2C), indicating that tandem SigX binding sites enhance *cab* expression. In the *S. mitis* collection, all

cab loci harbored at least two SigX boxes, with one strain containing up to 38 boxes (Fig. 2D and *SI*

Appendix, Table S1). Larger *cab* loci were associated with higher number of SigX boxes, suggesting

accumulation of boxes throughout these sequences.

Together, these results demonstrate that *cab* bacteriocins are specifically expressed during

the late phase of competence, and that the presence of multiple SigX binding sites amplifies their

expression.

Expression of *S. mitis cabABC* is induced by pneumococcal CSP-1

Mitis group streptococci naturally produce and recognize different CSP variants, known as pherotypes

(26). To investigate the specificity of *cab* expression, synthetic CSP variants from closely related

strains were tested for their ability to activate *cabABC*. CSP variants from three other *S. mitis*, one *S.*

oralis, and one *S. pneumoniae* strain failed to induce *cab* expression in *S. mitis* G22 (Fig. 3A). By

contrast, the most prevalent *S. pneumoniae* pheromone, CSP-1 (27), efficiently induced *cab* to the

same levels as native CSP$_{G22}$.

Comparison of the amino acid sequences CSP$_{G22}$ and the other CSP variants revealed a

single amino acid residue, Q14 (glutamine), that may explain the differences in induction. While both

CSP$_{G22}$ and CSP-1 contain Q14, a polar residue, the non-inducing variants contain either a L14

(leucine) or a P14 (proline), both nonpolar. To investigate the importance of Q14 and polarity in

competence induction, CSP$_{G22}$ variants with single amino acid substitutions at position 14 were tested.

Substituting Q14 with L significantly reduced *cab* expression, whereas replacing it with another polar

residue (N, asparagine) restored expression (Fig. 3B). This suggests that polarity of amino acid 14 is

important for CSP recognition by *S. mitis* G22. However, introducing Q14 into a non-inducing peptide

(CSP-2; L14Q) was not sufficient to convert it into a functional inducer, indicating that additional

residues beyond Q14 are required for efficient recognition by the membrane-bound histidine kinase

ComD of *S. mitis* G22.

To assess whether *S. pneumoniae* can also respond to CSP$_{G22}$, the *comCDE* promoter from

S. pneumoniae D39, a strain that produces and recognizes CSP-1, was transcriptionally fused to a

luc reporter and integrated in its chromosome. Competence in D39 was efficiently induced by both

CSP-1 and CSP$_{G22}$, confirming that *S. mitis* G22 and *S. pneumoniae* D39 can cross-communicate

through distinct CSP peptides (Fig. 3C). Similar to G22, *S. pneumoniae* D39 was also affected by

mutations at position 14: Q14L (polar to nonpolar) reduced induction. In contrast to *S. mitis* G22, the

substitution Q14N (polar with another polar aminoacid), failed to restore it. Notably, the substitution

L14Q into CSP-2 (the second most frequent CSP in pneumococci) enabled competence induction.

These findings highligh the importance of Q14 in CSP recognition of pneumococci of the CSP-1

pherotype.

In the 253 *S. mitis* genomes analyzed, 2.8% encode CSP$_{G22}$, and 13.0% encode a CSP

containing Q14 (*SI Appendix*, Table S1), suggesting that more *S. mitis* strains could induce

competence in pneumococci of pherotype CSP-1.

These results reveal that cross-communications via CSP is possible between *S. mitis* and *S.*

pneumoniae, and that residue 14 plays a critical, but context-dependent, role in pheromone

recognition.

CabAB mediates inhibition of *S. pneumoniae* in a contact-dependent manner

Given that *S. mitis* G22 and *S. pneumoniae* D39 can cross-communicate through CSP, we

hypothesized both species would engage in competence-induced predation. Supporting this, *S. mitis*

G22 harbors the orphan immunity gene *cibC* (but not *cibAB*), suggesting it may be resistant to

pneumococcal CibAB attack (*SI Appendix*, Table S1).

To investigate the specific role of the competence associated bacteriocins CabAB in *S.*

pneumoniae inhibition, we first deleted two bacteriocin clusters (*blp4* and *blp1*) with known anti-

pneumococcal activity (23), from the genome of *S. mitis* G22, as their activity could obscure CabAB-

mediated effects. All experiments in this section were, thus, performed using *S. mitis* G22 Δ*blp4*Δ*blp1*

and its derivatives as described below.

We grew dual-species biofilms of *S. mitis* G22 Δ*blp4*Δ*blp1* and *S. pneumoniae* D39 under

competence-permissive conditions. After 24 h, both species persisted in the biofilm (Fig. 4A and *SI*

Appendix, Fig. 5). Deletion of *cab* from *S. mitis* led to increased pneumococcal loads, while

constitutive expression of *cab* via plasmid p(*cab*$^+$) (overexpressor mutant contains the native *cab* copy

and carries an additional constitutively expressed copy in plasmid p(*cab*$^+$)) reduced them, indicating

that the presence and expression levels of *cab* modulate competitive outcomes.

Bacteriocins are typically secreted to the extracellular environment (28). To test whether

CabAB exerts its activity as a secreted factor, we harvested cell-free supernatants (CFS) from *S. mitis*

strains and added them to pneumococcal cultures. CFS of *S. mitis* natively or overexpressing *cab* did

not inhibit *S. pneumoniae* growth (Fig. 4B), suggesting CabAB is not released into the medium. This

aligns with recent reports of bacteriocins, such as Listeriolysin S and contact-dependent inhibition by glycine zipper proteins CdzCD (29, 30), that remain attached to the producer cell surface. In these cases, inhibitory activity occurs via direct contact between producer and target cells.

To test whether CabAB could remain cell-associated, we heat-killed *S. mitis* strains (as bacteriocins are typically resistant to high temperature treatments) and exposed them to *S. pneumoniae*. Heat-killed *cab*+ and p(*cab*+) *S. mitis* extracts significantly inhibited pneumococcal growth, while extracts from Δ*cab S. mitis* did not (Fig. 4B), supporting the idea that CabAB is retained in producer cells.

We then tested whether direct contact between *S. mitis* and *S. pneumoniae* was required for inhibition. Dual-species biofilms were grown either with or without a 0.4 μm transwell membrane separating both species. When the strains were in direct contact with each other, *cab* expression was associated with pneumococcal inhibition, whereas no inhibition was observed when both species were separated (Fig. 4C). These effects were not due to differences in *S. mitis* growth, as bacterial loads were consistent across conditions and genotypes (Fig. 4D).

We then sought to detect CabA and CabB on the surface of competent *S. mitis* cells. A mutant lacking the native *cab* locus was used, into which we ectopically integrated cassettes containing haemagglutinin-tagged (HA) CabA (*cabA-HA cabBC*) or CabB (*cabA cabB-HA cabC*), under native regulation. Immunofluorescence microscopy of live *S. mitis* expressing *cabA-HA* using anti-HA antibodies revealed discrete clusters of CabA on the cell surface (Fig. 4E). No signal was detected in strains lacking the HA epitope, confirming specificity. Interestingly, no signal was observed in the *cabB-HA* strain, suggesting that CabB may be less surface-exposed, that the N-terminal HA tag was not accessible to antibody binding, or that tagging interfered with localization.

Together, these results demonstrate that *S. mitis* inhibits *S. pneumoniae* via CabAB in a contact-dependent manner.

CabAB is a two-peptide bacteriocin with LytA-dependent bactericidal activity

CibAB has previously been shown to be a two-peptide bacteriocin, meaning its inhibitory activity occurs only when both peptide components are present (8). To test if CabAB also functions as a two-

peptide bacteriocin, synthetic CabA and CabB were tested individually and in combination against *S.*

pneumoniae D39. No effect on pneumococcal growth was observed upon addition of DMSO (the

solvent used to resuspend the peptides), CabA alone, or CabB alone (Fig. 5A). In contrast, a single

treatment with equimolar amounts of CabA and CabB delayed *S. pneumoniae* growth for ~12 h,

indicating that both peptides are required for activity. This is consistent with the presence of multiple

GxxxG or GxxxG-like motifs in both bacteriocins (Fig. 5B), known to facilitate and stabilize two-

component bacteriocins (31-33).

We next investigated whether CabAB exerts bactericidal activity. Exponentially growing *S.*

pneumoniae D39 cultures were treated with CabAB or DMSO. CabAB-treated cultures showed

evident decrease in optical density, indicating lysis, while DMSO-treated controls did not (Fig. 5C).

Lysis induced by the related bacteriocin CibAB has been shown to depend on the action of the

pneumococcal autolysins LytA and LytC (8). Here, deletion of *lytC* from D39 had no effect on CabAB-

mediated lysis, whereas a Δ*lytA* mutant exhibited delayed growth but no lysis, indicating that LytA is

required for this phenotype.

Together, these findings show that *S. mitis* CabAB is a two-peptide bacteriocin with

bactericidal activity against *S. pneumoniae*, and that its lytic activity depends on the target cell LytA

autolysin.

CabAB affects membrane homeostasis in *S. pneumoniae*

We sought to shed light on how CabAB affects pneumococcal physiology. To this end, *S. pneumoniae*

D39 cells were treated with CabAB or DMSO and imaged by scanning electron microscopy (SEM).

While DMSO-treated cultures displayed normal morphology, CabAB-treated cells appeared to be

encased in an extracellular matrix (Fig. 6A), suggesting leakage of intracellular material.

To test if membrane permeability was compromised, cells were stained with propidium iodine

(PI), a DNA-binding dye that only enters cells with compromised membranes. In DMSO-treated

cultures, only 3% were PI-positive, while CabAB-treated cultures contained 85% PI-positive cells (Fig.

6B). Doubling the concentration of CabAB further increased this proportion to 97%, indicating this

phenotype is concentration dependent. Deletion of *lytA* had no effect on these phenotypes,

demonstrating that membrane permeabilization occurs independently of autolysin-mediated lysis (*SI Appendix*, Fig. S6).

Time series measurements showed that PI signal intensity peaked at 30 min post CabAB exposure, still being detected after 120 min (Fig. 6C). We also assessed pneumococcal membrane depolarization using the fluorescent dye $DiBAC_4(3)$, which enters cells only when their membranes are depolarized. Compared to DMSO, CabAB treatment led to a rapid and sustained increase in $DiBAC_4(3)$ signal (Fig. 6D), indicating membrane depolarization.

Since some characterized two-peptide bacteriocins act by forming pores (34-37), we tested whether CabAB treatment causes efflux of intracellular components from target cells. In wild-type *S. pneumoniae* treated with CabAB, 80% of total ATP was detected in the supernatant (Fig. 6E), in agreement with our previous observation of cell lysis (Fig. 5C). Interestingly, in treated Δ*lytA S. pneumoniae* cells, which are not lysed by CabAB (Fig. 5C), 30% of ATP was still released, suggesting that CabAB induces leakage independently of cell lysis.

We next visualized CabAB-induced damages using transmission electron microscopy (TEM). DMSO-treated cells appeared intact, whereas 17% of CabAB-treated cells contained large intracellular aggregates (Fig. 6F). Based on our prior observations, we hypothesized these aggregates would be membranous. Consistently, staining with the membrane dye FM4-64 revealed intracellular foci in CabAB-treated cells, but not in control cells (Fig. 6G), confirming the formation of membrane aggregates by CabAB.

Finally, lower-magnification fluorescent microscopy of CabAB-treated cultures stained with DAPI and FM4-64 revealed frequent clumps of cells encased in nucleic acid and membrane material, mirroring the SEM observations of surviving cells remaining embedded in an extracellular matrix (*SI Appendix*, Fig. 7)

Altogether, these findings show that *S. mitis* CabAB disrupts membrane homeostasis in *S. pneumoniae* by permeabilizing and depolarizing membranes, leading to release of its intracellular content.

CabAB facilitates the acquisition of pneumococcal DNA by *S. mitis* in biofilms

As previously shown, CabAB promotes release of DNA from *S. pneumoniae* cultures. Since *cab* expression is synchronized with the competence period, we hypothesized that CabAB could enhance DNA acquisition and HGT within biofilms. To test this, we grew dual-species biofilms under conditions designed to study HGT (Fig. 7A), where: (i) DNA transfer is unidirectional, since the DNA donor strain (*S. pneumoniae* D39) carries a deletion of the major competence pilus subunit gene, *comGC*, rendering it nontransformable; (ii) the *S. pneumoniae* DNA donor harbors chloramphenicol (Chl^R) and tetracycline (Tet^R) resistance cassettes integrated at opposite ends of the chromosome; (iii) the DNA recipients (*S. mitis* G22, wt and variants) are kanamycin resistant, allowing for discrimination between non-transformed (Kan^R) and transformed *S. mitis* (Kan^R Chl^R, Kan^R Tet^R, and Kan^R Chl^R Tet^R); (iv) biofilms are pre-grown under acidic pH for 24 h to promote biofilm establishment without inducing competence; and (v) CSP$_{G22}$ is added after 24 h to transiently induce competence. A noncompetent *S. mitis* ΔcomE mutant was used as a negative control, as it is unable to acquire Chl^R and/or Tet^R by transformation.

To rule out the possibility that CabAB affects transformability rather than DNA availability, we first replaced the donor strain with purified gDNA. Under these conditions, both *cab*$^+$ and Δ*cab S. mitis* acquired resistance at similar frequencies (Fig. 7B), indicating that native *cab* expression does not impact transformability. In contrast, overexpression of *cab* reduced transformation efficiency for one antibiotic resistance, possibly due to the fitness burden associated with carrying p(*cab*$^+$).

In dual-species biofilms, *cab*$^+$ *S. mitis* efficiently acquired Chl^R and Tet^R from the DNA donor, at a significantly higher frequency than the Δ*cab* mutant (Fig. 7C), indicating that CabAB contributes to DNA acquisition. In contrast, overexpression of CabAB did not enhance transformation efficiency, suggesting that a precise *cab* regulation might be necessary for optimal transformation.

Finally, we sought to investigate the specificity of CabAB-induced lysis. Synthetic CabAB was tested against 26 bacterial species, including 18 streptococcal species, four additional non-streptococcal Gram-positive and four Gram-negative species. All selected species are human commensals or pathogens that can inhabit the oral and/or respiratory tract and could theoretically encounter *S. mitis* and be exposed to Cab bacteriocins in natural environments. Addition of CabAB

to exponential cultures of these bacteria stalled the growth of some species (Fig. 7D and *SI Appendix*,

Fig. S8 and S9). The most sensitive to CabAB were Pyogenic Group streptococci (*S. pyogenes* and

S. agalactiae). Gram-negative bacteria (*Haemophilus influenzae*, *Klebsiella pneumoniae*, *Moraxella*

catarrhalis, and *Neisseria lactamica*) were largely unaffected. Despite growth inhibition in some

species, lysis was only observed in two - *S. pneumoniae* and *S. parasanguinis* - highlighting the

specificity of CabAB-mediated lysis.

These results suggest that CabAB provides a competitive advantage by inhibiting the growth

of some bacteria, but that it also enhances DNA release in biofilms by selectively lysing a few closely

related species during competence, notably *S. pneumoniae*, thereby promoting gene exchange.

Discussion

During competence, streptococci scavenge extracellular DNA. In *S. pneumoniae*, where competence-

induced predation is best characterized, the coordinated action of CibAB, LytA, LytC, and the murein

hydrolase CbpD leads to lysis of noncompetent siblings (8). This process promotes DNA release and

facilitates HGT (9, 38, 39).

Although extensive genomic data suggests that *S. mitis*, a close relative of *S. pneumoniae*,

frequently acquires *S. pneumoniae* DNA (14-20), the molecular mechanisms underlying this

exchange remained poorly understood. Understanding these mechanisms is relevant because

capsule acquisition by *S. mitis* can complicate DNA-based pneumococcal serotyping. In addition, *S.*

mitis has been associated with cases of infective endocarditis (19), highlighting the value of studying

how it acquires potentially pathogenic traits.

Here, we investigated the role of *S. mitis* competence-associated bacteriocins, CabAB, in

mediating interspecies interactions. We show that CabAB is a two-peptide bacteriocin produced

during the late phase of competence, when cells are primed for DNA uptake and recombination,

capable of lysing *S. pneumoniae* in a LytA-dependent manner. Although LytA is constitutively

expressed throughout growth, its concentration increases during competence, owing to a genomic

architecture comprising three constitutive promoters and one competence-induced promoter (7, 40-

42). In this context, it is tempting to speculate that by inducing *S. pneumoniae* competence via CSP

crosstalk, and therefore increasing LytA concentration, some *S. mitis* could further sensitize *S.*

pneumoniae to Cab bacteriocins. Nonetheless, CabAB-induced lysis appears to be independent of

pherotype of target cells, as both *S. pneumoniae* D39 (CSP-1; used in most experiments) and TIGR4

(CSP-2; included in the panel of 26 bacterial species) were lysed. These pherotypes account for the

vast majority (>95%) of pneumococcal strains (43-45).

Of the initial 332 *S. mitis* genomes analyzed, we were not able to retrieve the region upstream

of *comAB* from a substantial fraction of strains (23.8%), as their assemblies showed contig breaks

precisely at this locus. We propose this difficulty arises from the repetitive nature of the region, as

indicated by the presence of numerous SigX boxes. The number of boxes scaled linearly with the size

of *cab* loci, suggesting that SigX boxes have accumulated throughout these sequences. Repeats are

known to both promote and result from genomic rearrangements (46, 47). They are often remnants

of previous recombination events, with no known biological function (48). In this case, however,

accumulation of SigX boxes may provide a selective advantage by enhancing bacteriocin expression,

which could explain why they are enriched in this region. Consistent with this, the extensive genetic

rearrangements observed across multiple streptococcal species at this locus also suggests this is a

highly recombinogenic region (21, 22, 49).

Microscopic analysis revealed that CabAB-treated pneumococci often contain large

membrane-derived intracellular aggregates. Whether these structures are part of CabAB's

mechanism of action or represent a cellular stress response remains unclear. Intracellular vesicle

formation has been shown to be essential for the lytic activity of bacteriocin A37 (50). However, the

disordered morphology of CabAB-induced structures suggests an alternative process. We

hypothesize that aggregate formation may reflect a protective response, where membrane

invaginations at CabAB-enriched sites reduce metabolite leakage and trap affected membrane

material intracellularly.

We further demonstrate that CabAB-mediated inhibition of *S. pneumoniae* requires direct

contact between producer and target cells. Contact-dependent bacteriocin delivery has been

described in *Caulobacter crescentus* and *Listeria monocytogenes* (29, 30). In *L. monocytogenes*,

bacteriocins were only detected on the surface of producer cells following cell wall digestion. In
contrast, we detected CabAB on live *S. mitis* with intact cell walls, suggesting a distinct delivery
mechanism. Contact-dependent killing during competence is not unique to streptococci, despite
differences in killing factors. *Vibrio cholerae* produces a competence-induced type VI secretion
system that kills nonimmune cells, releasing DNA that becomes accessible for HGT (51). Contact
dependence may be particularly advantageous in biofilm contexts, a niche where extracellular DNA
could be subject to rapid degradation nucleases (52). Proximity within multispecies biofilms could
enhance transformation efficiency by increasing the local DNA concentration, given that DNA
abundance and size influence recombination success (53). Notably, biofilm conditions and cell-to-cell
contact have been shown to promote both the frequency and size of recombination events in
pneumococci, respectively (54, 55).

Strikingly, among 26 tested species, only *S. pneumoniae* and *S. parasanguinis* were lysed by
CabAB, suggesting a narrow-spectrum bactericidal activity. Whether this specificity is unique to
CabAB or shared among other Cab bacteriocins remains to be determined. Nonetheless, the ability
to selectively target phylogenetically related but distinct species suggests a potential trade-off
between maximizing recombination efficiency and accessing a broader gene pool. While high
sequence identity between donor and recipient promotes recombination (56), restricting lysis to
closely related species could expand the range of accessible alleles beyond conspecifics.

Importantly, CabAB also strongly inhibited the growth of other streptococcal pathogens,
including *S. pyogenes* and *S. agalactiae*, both of which are listed as priority pathogens by the World
Health Organization (25). This potent inhibitory activity suggests that the Cab arsenal carried by *S.*
mitis could be a promising source of antimicrobial agents against important clinical pathogens as
recently suggested by us (25).

In summary, our findings uncover a contact-dependent mechanism of bacteriocin delivery in
S. mitis that selectively targets closely related streptococci including *S. pneumoniae*. This process
promotes HGT and likely contributed to competitive fitness and evolutionary plasticity of streptococcal
populations.

Materials and Methods

Bacterial strains and culture conditions

Bacterial strains, oligonucleotides, plasmids and peptides used in the study are listed in *SI Appendix*, Table S2. Streptococci were routinely grown in C+Y$_{YB}$ (57) (pH 7.4, unless otherwise specified) and plated on tryptic soy agar with 5% defibrinated sheep blood (blood agar, BA). When appropriate, media was supplemented with antibiotics at the following concentrations: 150 µg.mL^{-1} kanamycin, 200 µg.mL^{-1} spectinomycin, 4 µg.mL^{-1} chloramphenicol, or 4 µg.mL^{-1} tetracycline.

Staphylococcus aureus, Staphylococcus epidermidis and *Moraxella catarrhalis* were grown in tryptic soy broth (TSB); *Corynebacterium accolens* was grown in TSB with 0.5% Tween 80; *Klebsiella pneumoniae* and *Escherichia coli* were grown in lysogeny broth; and *Neisseria lactamica*, *Haemophilus influenzae* and *Dolosigranulum pigrum* were grown in C+Y$_{YB}$ with 20 µg.mL^{-1} hemin and 20 µg.mL^{-1} NAD. For propagation of plasmids in *E. coli*, media was supplemented with 100 µg.mL^{-1} kanamycin or 100 µg.mL^{-1} spectinomycin, as required.

Identification of *cab* bacteriocins and *cibABC* in *S. mitis*

Publicly available *S. mitis* genomes (n = 322) were downloaded from PubMLST. An additional 10 genomes from reference or commonly used *S. mitis* strains were included, resulting in a total of 332 genomes (*SI Appendix*, Table S1). To detect *cab* clusters, nucleotide sequences of *plsX-acpP* and *comAB* (from *S. mitis* B6; GenBank Accession GCA_000027165.1), which flank the *cab* locus, were aligned against all genomes using pairwise sequence alignment (Biopython v1.81; match = 2; mismatch = -1; gap open = -5; gap extension = -0.5; minimum overall sequence identity $\geq$ 90%). Genomes in which *plsX-acpP* and *comAB* were absent or located on separate contigs were excluded from further analysis. This region was identified in 253 genomes.

To identify mature Cab bacteriocin sequences, extracted sequences were scanned for open reading frames (ORFs) of 100-300 nucleotides, beginning with an ATG start codon and ending with a TAA, TAG or TGA stop codon. Only ORFs whose translated sequences contained a (M/L/V)XXXXGG double-glycine cleavage motif were retained. Mature Cab bacteriocins were

obtained by removing the N-terminal signal sequence upstream of the cleavage site. To explore

diversity of Cab bacteriocins, sequences were converted into amino acid frequency vectors. Pairwise

dissimilarities between sequences were calculated using the Bray-Curtis metric (ScyPy), followed by

agglomerative hierarchical clustering (scikit-learn). To determine the optimal number of clusters,

silhouette scores (scikit-learn) were computed for cluster numbers ranging from 2 to 50. The number

of clusters yielding the highest score was selected as optimal.

To assess the prevalence of *cibABC* in *S. mitis*, the nucleotide sequence of *cibABC* (from *S.*

pneumoniae D39, GenBank Accession NC_008533.2) was aligned against all *S. mitis* genomes using

NCBI BLASTn. Genomes with no detectable hit were considered *cibABC*-negative. For genomes with

partial matches (<100% query coverage), the locus was manually inspected to verify the presence or

absence of *cibA*, *cibB*, and *cibC* individually – a common occurrence in strains harboring *cibC* but not

cibAB.

To find SigX boxes, the consensus sequence (58) was searched within *cab* loci using Find

Individual Motif Occurrences (FIMO) v5.5.8 (59). Hits with *p*-value <0.001 were considered true SigX

boxes. The detection of Cab bacteriocins and putative SigX boxes was benchmarked against six *S.*

mitis strains with complete genomes and manually curated *cab* loci (23, 60).

Phylogeny of *S. mitis*

456 Core genome single-nucleotide polymorphisms (SNPs) were called using Snippy v4.6.0

457 (https://github.com/tseemann/snippy) by aligning *S. mitis* genomes to the *S. mitis* B6 reference

458 genome. The resulting alignment was used to construct a maximum-likelihood phylogenetic tree with

459 FastTree v2.2 (61). The phylogeny was visualized and annotated using the Interactive Tree of Life

460 (iTOL) platform (62).

Synthetic peptides

Peptides were produced by solid-phase peptide synthesis with >95% purity (NZYTech) (*SI Appendix*,

Table S2) (23, 43-45). CSP and BlpC$_1$ pheromones were resuspended in sterile ddH$_2$O; BlpC$_4$ was

resuspended in 10% acetic acid; and CabA and CabB were resuspended in DMSO. Peptides were

serially diluted in sterile ddH_2O and stored at -20ºC.

Transformation

For transformation of *S. mitis* and *S. pneumoniae*, cells were grown to an OD_{600} of 0.5, diluted 1:100

in pre-warmed C+Y_{YB} and grown until an OD_{600} of 0.04-0.1. Cells were treated with 1 µM of cognate

CSP and transformed with 0.4 µg linear DNA, 1 µg plasmid DNA, or 10 µg chromosomal DNA.

Transformants were recovered for 4 h and plated in selective medium. Linear, plasmid and genomic

DNA were purified using the Zymoclean Gel DNA Recovery Kit (Zymo Research), ZymoPURE

Plasmid Miniprep Kit (Zymo Research) and EZ1 DSP Virus Kit (Qiagen), respectively.

For heat-shock transformation of *E. coli* DH10B or DH5α, $CaCl_2$-competent cells were mixed

with 10% (v/v) of ligation reactions and incubated on ice for 15 min. Cells were then heat-shocked at

42ºC for 40 s, immediately chilled on ice, and recovered with 900 µL of SOC (2% tryptone, 0.5% yeast

extract, 10 mM NaCl, 2.5 mM KCl, 10 mM $MgCl_2$, 10 mM $MgSO_4$ and 20 mM glucose; pH 7) for 1 h

before plating in selective medium.

Construction of a fluorescence-luminescence transcriptional reporter

A DNA cassette containing a *hlpA-mNeonGreen* fusion and plasmid pBIR, which integrates in the

genome of *S. mitis* G22 and *S. pneumoniae* D39, were synthesized (Twist Bioscience). The reporter

cassette and plasmid were amplified and blunt end ligated to generate plasmid pJB10. The firefly

luciferase *luc* (from pPEP-LGZ) and plasmid pJB10 were amplified and blunt-end ligated to create

plasmid pJB11 encoding a dual fluorescence-luminescence reporter. Transcriptional fusions were

created by digesting both pJB11 and promoter fragments with BsaI, followed by ligation. Variants of

pJB11 were linearized with BamHI and transformed into *S. mitis* G22 or *S. pneumoniae* D39.

Transformants were selected in BA with chloramphenicol.

Expression of *cabABC* and other competence genes

S. mitis and *S. pneumoniae* strains harboring transcriptional fusions of competence promoters were

grown in C+Y_{YB} at pH 6.6 (to prevent natural competence induction) until an OD_{600} of 0.5. To evaluate

natural competence dynamics, acidic precultures were diluted 1:100 in C+Y_{YB} (pH 6.6, 7.4 or 8.2)

containing 340 µg/mL D-luciferin (Abcam). To assess the effect of different pheromones on

competence induction, acidic precultures were diluted to OD_{600} = 0.1 in C+Y_{YB} (pH 6.6) with 340 µg/mL

D-luciferin. Then, CSP or BlpC variants (up to 1 µM), were added. Cultures were transferred to white

96-well plates with clear bottoms (Nunc). The OD_{595} and luminescence was recorded every 10 min

for 24 h in a BioTek Neo2 Multimode Reader with a gain of 185. Luminescence values were

normalized to the OD_{595}. D-luciferin was resuspended in sterile ddH_2O to 15 mg.mL^{-1} and kept at -80

ºC in single-use aliquots.

Construction of ComE, SigX1 and SigX2 deletion mutants

The upstream and downstream flanking regions of *comE, sigX1, sigX2* were amplified from *S. mitis*

G22 gDNA. Kanamycin and spectinomycin resistance genes were amplified from pKan and pPEP-

LGZ (63), respectively. DNAs were ligated using Gibson Assembly (New England Biolabs) and

amplified with nested primers. Transformants were selected in BA with kanamycin (for *comE* and

sigX1) or spectinomycin (for *sigX2*).

Biofilms and transwells

S. mitis and *S. pneumoniae* were grown to exponential phase and mixed at a 1:1 ratio (10^5 CFU of

each strain) in C+Y_{YB} with 1600 U.mL^{-1} catalase. Two milliliters of the mixtures were added to tissue-

culture treated 12-well plates (Corning) and incubated 24 h at 34ºC with 5% CO_2. Non-adherent cells

were removed by aspiration, and biofilms were thoroughly resuspended in PBS.

For transwell biofilms, 1.5 mL of C+Y_{YB} containing catalase and 10^5 CFU of *S. pneumoniae*

was added to wells of tissue-culture treated 12-well plates. Transwell inserts containing a 0.4 µm pore

polycarbonate membrane (Corning) were placed in the wells, and 0.5 mL of C+Y_{YB} with catalase and

10^5 CFU of *S. mitis* were added to the upper insert. Biofilms were recovered from both compartments.

In dual-species biofilms, *S. pneumoniae* and *S. mitis* cells were enumerated by plating on BA

with chloramphenicol or kanamycin, respectively. In the transwell setup, where *S. pneumoniae* and

S. mitis were physically separated, biofilm suspensions from both compartments were also plated on

selective media to confirm that neither strain had crossed the membrane.

Construction of an *S. mitis cab* overexpressor mutant

The plasmid backbone was amplified from pDL278_P23-DsRed-Express2 (Addgene plasmid

#121505), and *cabABC* was amplified from *S. mitis* G22 gDNA. The two fragments were blunt-end

ligated, generating plasmid pJB09-*cabABC*, in which *cabABC* is placed under the control of the strong

constitutive P23 promoter. *S. mitis* transformants were selected in BA with spectinomycin.

Inhibitory activity of cell-free supernatants and heat-killed cells

S. mitis was grown to early stationary phase and centrifuged at 10,000 g for 30 min at 4ºC. Cell-free

supernatants (CFS) were obtained by filtration (0.22 µm). CFS were lyophilized by vacuum

evaporation using a SpeedVac concentrator (LabConco) and resuspended in ddH$_2$O to a

concentration of 10x before testing. C+Y$_{YB}$ medium was concentrated in parallel and used as a

negative control for inhibitory activity. Cell pellets were washed once in PBS and heat-killed for 15

min at 95 ºC. Complete lysis of pellet extracts was confirmed by plating in BA.

To test for inhibitory activity of CFS and heat-killed cells, *S. pneumoniae* D39 was grown to

an OD$_{600}$ of 0.5, diluted 1:100 in fresh medium, and 180 µL of the dilution was added to clear 96-well

plates. Twenty microliters of treatments or water was added to each well and growth was monitored

by measuring the OD$_{595}$ every 30 min for 24 h using a Tecan Infinite 200 Pro.

Sensitivity of bacteria to CabAB

Exponential-phase cultures of bacterial species (18 *Streptococcus* spp., four additional non-

streptococcal Gram-positive species, and four Gram-negative species) were diluted to OD$_{600}$ 0.1 and

195 µL was added to wells of clear 96-well plates. CabAB (0.975 µM) or DMSO (0.00975%) was

added to wells and growth was recorded by measuring the OD$_{595}$ every 10 min for 24 h using the

Tecan Infinite 200. Growth arrest was quantified by comparing the area under the curve (AUC)

between CabAB- and DMSO-treated cultures. All further assays using CabAB and DMSO treatments

were performed in the same conditions, unless specified otherwise.

Construction of autolysis-deficient *S. pneumoniae* mutants

The upstream and downstream regions of *lytA* and *lytC* were amplified from *S. pneumoniae* D39

gDNA. The PhunSweet cassette containing *pheS*-*sacB*-aph*A-3* which confers susceptibility to

chlorinated phenylalanine and sucrose, and resistance to kanamycin, was amplified from *S.*

pneumoniae D39 Δ*cps*::*PhunSweet* gDNA (64). DNAs were assembled using Gibson Assembly and

amplified with nested primers. *S. pneumoniae* D39 transformants were selected on BA with

kanamycin.

To remove the antibiotic resistance cassette, the flanking regions of *lytA* and *lytC* were

amplified, assembled with Gibson Assembly, and amplified with nested primers. Transformants were

recovered by counterselection on BA with 15 mM chlorinated phenylalanine and 10% sucrose.

Scanning electron microscopy (SEM)

S. pneumoniae D39 cultures (30 mL) were treated with DMSO or CabAB and incubated for 15 min at

37ºC. Cells were harvested by centrifugation (all centrifugations were carried out at 800 g for 10 min

at room temperature) and washed three times with PBS. Fixation was performed in 500 μL of 2.5%

glutaraldehyde and 1% paraformaldehyde in 0.1 M sodium cacodylate buffer for 30 min at room

temperature. Cells were washed three times with 0.1 M sodium cacodylate buffer and dehydrated in

increasingly higher ethanol concentrations (50%, 70%, 90%, 100%), by incubating 10 min in 500 μL

of each ethanol concentration. Ethanol was replaced with 500 μL tert-butyl alcohol and samples were

incubated at 30ºC for 1 h. Then, tert-butyl was removed and replaced with 500 μL of fresh tert-butyl

alcohol. Samples were kept at -20ºC overnight. Samples were then lyophilized under vacuum, gold-

sputtered using a Cressington 108 sputter coater, and imaged with a Hitachi SU-8010 scanning

electron microscope at 1.5kV.

Transmission electron microscopy (TEM)

S. pneumoniae D39 cultures (30 mL) were treated with DMSO or CabAB, incubated for 15 min at 37ºC, and recovered by centrifugation at 800 g. Cells were fixed with 2% formaldehyde, 2.5% glutaraldehyde, 0.075% ruthenium red and 1.55 % lysine acetate in distilled water for 20 min at 4ºC. Cells were washed twice in 0.075% ruthenium red, resuspended in 2% formaldehyde, 2.5% glutaraldehyde and 0.075% ruthenium red, and incubated overnight at 4ºC. Cells were washed three times with 0.075% ruthenium red and further processed at Instituto Gulbenkian de Ciência. Cells were embedded in 2% low-melting point agarose and fixed with 1% osmium tetroxide and 0.06% ruthenium red for 1 h at room temperature. Cells were washed once with 0.1 M sodium cacodylate buffer and twice with distilled water. Dehydration was performed on ice using increasing concentrations of ethanol (10%, 30%, 50%; 10 min each). Samples were stained overnight at 4ºC with 2% uranyl acetate in 70% ethanol, followed by further dehydration with 90% ethanol and three times with 100% ethanol. Infiltration was carried out with increasing concentrations of epoxy resin in ethanol (3:1, 1:1, 1:3), and 100% resin overnight, followed by two exchanges of 100% resin. Ultrathin sections were cut and imaged on a Tecnai G2 Spirit BioTWIN Transmission Electron Microscope (FEI) equipped with an Olympus-SIS Veleta CCD camera.

Large intracellular aggregates were observed in some CabAB-treated cells. To quantify these structures, micrographs of CabAB- and DMSO-treated cells were blindly evaluated by two researchers. Each researcher was first shown representative cells with and without aggregates, and then independently counted the number of cells containing these structures. The final percentage of cells with aggregates was calculated as the average between both researcher counts.

Permeabilization and depolarization of *S. pneumoniae* membranes

S. pneumoniae D39 cultures were treated with DMSO or CabAB. One-milliliter aliquots were taken at 0, 2, 5, 15, 30, 60, and 120 min post treatment and immediately chilled on ice. Cells were washed twice with PBS and kept on ice. Cells were stained with 6 µM propidium iodine (Invitrogen) and 3 µM DiBAC$_4$(3) (Enzo Life Sciences) for 5 min at room temperature, protected from light. Cells were

centrifuged, resuspended in PBS, and added to black 96-well plates with clear bottoms (Nunc).

Fluorescence intensity was measured using a BioTek Neo2 Multimode Reader under the following

settings: shake for 5 s; propidium iodine – top reading, 535 ± 12 nm excitation, 617 ± 12 nm emission

and 120 gain; $DiBAC_4(3)$ – top reading, 492 ± 9 nm excitation, 518 ± 9 emission, 120 gain).

Fluorescent values were normalized to the OD_{595}.

Efflux of ATP

Cultures of *S. pneumoniae* were treated with DMSO or CabAB. Aliquots were taken at 0, 15 and 120

min post-treatment and kept on ice. Cells were centrifuged, and supernatants were filtered (0.22 µm)

and maintained on ice. Cells were permeabilized in DMSO for 10 min at room temperature. Both cells

and supernatants were diluted 1:10 in ddH_2O and kept on ice.

Extracellular and intracellular ATP was measured with the ATP Luminescent Cell Viability

Assay (Millipore) according to instructions. Luminescence was measured in a BioTek Neo2 Multimode

Reader using a gain of 150 and integration time of 10 s. An ATP standard curve confirmed linear

detection within the range of 0.2 μM to 2 mM ATP. Total ATP was calculated by summing

luminescence values from both the cell and supernatant fractions under each condition.

Construction of *S. mitis* harboring HA-tagged bacteriocins

DNAs containing HA (YPYDVPDYA)-tagged *cabA* and *cabB* were synthesized (Twist Bioscience).

The regions upstream (including the native promoter) and downstream (including a chloramphenicol

resistance cassette) of the *cab* operon were amplified from pJB11-*cabABC*. DNAs were assembled

by Gibson Assembly and amplified with nested primers. The resulting constructs were transformed

into the *S. mitis* Δ*cabABC* background, generating two mutant strains: *cabA-HA cabBC* and *cabA*

cab-HA cabC. Transformants were selected in BA with chloramphenicol.

Confocal microscopy and immunofluorescence

To quantify the percentage of cells that become membrane-permeable, *S. pneumoniae* was treated

with DMSO or CabAB. Cells were recovered by centrifugation (9600 g, 4ºC, 3 min) and washed three

times with PBS. Cells were stained with 30 µM propidium iodine and 5 µM SYTO 9 for 15 min at room

temperature, protected from light.

To determine whether large intracellular structures observed in CabAB-treated *S. pneumoniae*

cells corresponded to membranes, treated cells were washed three times with PBS and stained with

20 µg.mL^{-1} FM4-64 and 15 µg.mL^{-1} DAPI for 5 min at room temperature, protected from light.

For immunofluorescence of live competent bacteria, *S. mitis* harboring HA-tagged bacteriocins

were grown in acidic C+Y$_{YB}$ and diluted to OD$_{600}$ = 0.1 in pre-warmed C+Y$_{YB}$. CSP$_{G22}$ was added at 1

µM, and cells were incubated for 70 min at 37ºC and then chilled on ice. To retain the competent

status of cells, all subsequent steps were performed on ice using ice-cold solutions. Cells were

centrifuged (9600 g, 4ºC, 3 min) and washed twice in PBS. Blocking was performed in PBS with 5%

bovine serum albumin (BSA) for 1 h. Cells were pelleted and resuspended in 100 µL of primary rabbit

monoclonal anti-HA antibody (clone C29F4; Cell Signaling Technology) diluted 1:100 in PBS with 1%

BSA, and incubated for 1 h. Cells were washed twice in PBS with 0.1% Tween 20, resuspended in

100 µL of Alexa Fluor 488-conjugated goat anti-rabbit secondary antibody (Invitrogen) diluted 1:500

in PBS with 1% BSA, and incubated 1 h protected from light. Cells were washed twice with PBS

containing 0.1% Tween 20 and once with PBS. Cells were then stained with 20 µg.mL^{-1} FM4-64 and

15 µg.mL^{-1} DAPI for 5 at room temperature, covered from light.

Immediately before imaging, cells were mounted on thin 1.7% agarose pads. Fluorescence

imaging was performed using a Zeiss LSM 880 point scanning confocal microscope equipped with

an Airyscan detector. Samples were imaged with a Plan-Apochromat 63x/1.4 Oil DIC M27 objective

(Zeiss), using laser lines 405 nm (DAPI), 488 nm (SYTO 9 and Alexa Fluor 488), and 561 nm (FM4-

64 and propidium iodine). Imaging was performed at laser intensities of 2-3% and a gain of 700 for

DAPI, 700 for SYTO 9, 1000 for Alexa Fluor 488, 900 for FM4-64, and 700 for propidium iodine. The

Zeiss Zen v3.0 was used to control the microscope and for Airyscan processing. Image analysis was

conducted using ImageJ/FIJI v2.14 (65).

Construction of a transformation-deficient *S. pneumoniae* mutant

The upstream and downstream regions of *comGC* were amplified from *S. pneumoniae* D39 gDNA, and *tetM* was amplified from *S. mitis* F22 gDNA. DNAs were ligated by Gibson Assembly and amplified using nested primers. *S. pneumoniae* transformants were selected in BA with tetracycline.

Horizontal gene transfer in biofilms

Dual-species biofilms of *S. pneumoniae* D39-GFP Δ*comGC* (chloramphenicol and tetracycline resistant; Chl[R] Tet[R]) and *S. mitis* (kanamycin resistant; Kan[R]) were grown as described above, except that C+Y$_{YB}$ was adjusted to pH 6.6 to prevent natural competence induction. After 24h, non-adherent cells were removed by aspiration, and 2 mL of fresh C+Y$_{YB}$ (pH 7.4) containing catalase and 1 µM of CSP$_{G22}$ were added to biofilms. After 6 h of incubation at 34ºC with 5% CO_2, biofilm cells were thoroughly resuspended in PBS and recovered by centrifugation.

Cells were plated on the following selective media: (i) BA with kanamycin to quantify total *S. mitis* (Kan[R]); (ii) BA with kanamycin and chloramphenicol to detect *S. mitis* transformants that have acquired chloramphenicol resistance (Kan[R] Chl[R]), either with or without tetracycline resistance; (iii) BA with kanamycin and tetracycline to detect *S. mitis* transformants that acquired tetracycline resistance (Kan[R] Tet[R]), either with or without chloramphenicol resistance; (iv) BA with kanamycin, chloramphenicol and tetracycline to identify *S. mitis* transformants that acquired both resistances (Kan[R] Chl[R] Tet[R]); (v) and BA with chloramphenicol and tetracycline (without kanamycin) to detect *S. pneumoniae* loads.

Transformants were counted after 48 h of incubation. *S. mitis* transformation frequencies were calculated as the ratio between the number of transformants (Kan[R] colonies also resistant to chloramphenicol and/or tetracycline) and the total number of Kan[R] colonies. *S. mitis* G22 Δ*comE* was used as a non-transformable control, in which acquisition of chloramphenicol and tetracycline resistance should not occur via transformation.

Statistical analyses

The two-tailed unpaired *Mann-Whitney U* test with Benjamini and Hochberg correction for FDR was

used to compare bacterial loads. Student's *t*-test with Benjamini and Hochberg correction for FDR

was used to compare gene expression. The ratio paired *t*-test was used for the statistical analysis of

bacterial proportions in multi-species biofilms. All statistical analysis were computed using GraphPad

Prism 9.0 and considered significant at *P*-value ≤0.05.

Data availability

All data is available from the corresponding author upon reasonable request.

Acknowledgements

We are grateful to Jan-Willem Veening (University of Lausanne) and Morten Kjos (Norwegian

University of Life Sciences) for providing plasmids pPEP-LGZ and Jason Rosch (St. Jude Children's

Research Hospital) for providing *S. pneumoniae* strain harboring a PhunSweet cassette. We also

acknowledge Luisa N. Hiller (Carnegie Mellon University) and Adriano O. Henriques (ITQB NOVA)

for fruitful discussions.

Funding

This work was supported by FCT – Fundação para a Ciência e Tecnologia, I.P., through project

STOPneumo (PTDC/BIA-MIC/30703/2017), MOSTMICRO-ITQB R&D Unit (UIDB/04612/2020,

UIDP/04612/2020), LS4FUTURE Associated Laboratory (LA/P/0087/2020), and PPBI - Portuguese

Platform of BioImaging (PPBI-POCI-01-0145-FEDER-022122) co-funded by national funds from OE

- "Orçamento de Estado" and by european funds from FEDER - "Fundo Europeu de Desenvolvimento

Regional". J. Borralho and J. Lança were supported by PhD fellowships (2021.07866.BD -

doi.org/10.54499/2021.07866.BD and UI/BD/153385/2022 - doi.org/10.54499/UI/BD/153385/2022).

References

1.   F. Griffith, The significance of pneumococcal types. *J Hyg (Lond)* **27**, 113-159 (1928).

2.   L. S. Håvarstein, G. Coomaraswamy, D. A. Morrison, An unmodified heptadecapeptide
pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*.
Proc Natl Acad Sci U S A **92**, 11140-11144 (1995).

3.   L. S. Håvarstein, P. Gaustad, I. F. Nes, D. A. Morrison, Identification of the streptococcal
competence-pheromone receptor. *Mol Microbiol* **21**, 863-869 (1996).

4.   F. M. Hui, D. A. Morrison, Genetic transformation in *Streptococcus pneumoniae*: nucleotide
sequence analysis shows *comA*, a gene required for competence induction, to be a member
of the bacterial ATP-dependent transport protein family. *J Bacteriol* **173**, 372-381 (1991).

5.   E. V. Pestova, L. S. Håvarstein, D. A. Morrison, Regulation of competence for genetic
transformation in *Streptococcus pneumoniae* by an auto-induced peptide pheromone and a
two-component regulatory system. *Mol Microbiol* **21**, 853-862 (1996).

6.   M. S. Lee, D. A. Morrison, Identification of a new regulator in *Streptococcus pneumoniae*
linking quorum sensing to competence for genetic transformation. *J Bacteriol* **181**, 5004-5016
(1999).

7.   S. N. Peterson *et al.*, Identification of competence pheromone responsive genes in
Streptococcus pneumoniae by use of DNA microarrays. *Mol Microbiol* **51**, 1051-1070 (2004).

8.   S. Guiral, T. J. Mitchell, B. Martin, J. P. Claverys, Competence-programmed predation of
noncompetent cells in the human pathogen *Streptococcus pneumoniae*: genetic
requirements. *Proc Natl Acad Sci U S A* **102**, 8710-8715 (2005).

9.   O. Johnsborg, V. Eldholm, M. L. Bjørnstad, L. S. Håvarstein, A predatory mechanism
dramatically increases the efficiency of lateral gene transfer in *Streptococcus pneumoniae* and
related commensal species. *Mol Microbiol* **69**, 245-253 (2008).

10. P. Shen, J. A. Lees, G. C. W. Bee, S. P. Brown, J. N. Weiser, Pneumococcal quorum sensing
drives an asymmetric owner-intruder competitive strategy during carriage via the competence
regulon. *Nat Microbiol* **4**, 198-208 (2019).

11. J. Hohwy, J. Reinholdt, M. Kilian, Population dynamics of *Streptococcus mitis* in its natural
habitat. *Infect Immun* **69**, 6055-6063 (2001).

12.   G. Salvadori, R. Junges, D. A. Morrison, F. C. Petersen, Competence in *Streptococcus*

pneumoniae and close commensal relatives: mechanisms and implications. *Front Cell Infect*

Microbiol **9**, 94 (2019).

13.   G. Salvadori *et al.*, High-resolution profiles of the *Streptococcus mitis* CSP signaling pathway

reveal core and strain-specific regulated genes. *BMC Genomics* **19**, 453 (2018).

14.   E. Potgieter, L. J. Chalkley, Relatedness among penicillin-binding protein 2b genes of

Streptococcus mitis, *Streptococcus oralis*, and *Streptococcus pneumoniae*. *Microb Drug*

Resist **1**, 35-42 (1995).

15.   R. Hakenbeck, Transformation in *Streptococcus pneumoniae*: mosaic genes and the

regulation of competence. *Res Microbiol* **151**, 453-456 (2000).

16.   F. C. Lessa *et al.*, *Streptococcus mitis* expressing pneumococcal serotype 1 capsule. *Sci Rep*

8, 17959 (2018).

17.   F. Pimenta *et al.*, *Streptococcus infantis*, *Streptococcus mitis*, and *Streptococcus oralis* strains

with highly similar *cps5* loci and antigenic relatedness to serotype 5 pneumococci. *Front*

Microbiol **9**, 3199 (2018).

18.   F. Ganaie, A. R. Branche, M. Peasley, J. W. Rosch, M. H. Nahm, Effect of oral streptococci

expressing pneumococcus-like cross-reactive capsule types on World Health Organization

recommended pneumococcal carriage detection procedure. *Clin Infect Dis* **75**, 647-656

(2022).

19.   A. Kalizang'oma *et al.*, Population genomics of *Streptococcus mitis* in UK and Ireland

bloodstream infection and infective endocarditis cases. *Nat Commun* **15**, 7812 (2024).

20.   U. B. Skov Sørensen, K. Yao, Y. Yang, H. Tettelin, M. Kilian, Capsular polysaccharide

expression in commensal *Streptococcus* species: genetic and antigenic similarities to

Streptococcus pneumoniae. mBio **7** (2016).

21.   D. Denapaite *et al.*, The genome of *Streptococcus mitis* B6 - what is a commensal? *PLoS One*

5, e9426 (2010).

22.   L. N. Ikryannikova *et al.*, Inhibitory effect of streptococci on the growth of *M. catarrhalis* strains
and the diversity of putative bacteriocin-like gene loci in the genomes of *S. pneumoniae* and
its relatives. *AMB Express* **7**, 218 (2017).

23.   J. Borralho *et al.*, Inhibition of pneumococcal growth and biofilm formation by human isolates
of *Streptococcus mitis* and *Streptococcus oralis*. *Appl Environ Microbiol* **91**, e0133624 (2025).

24.   S. Lehtinen, N. J. Croucher, F. Blanquart, C. Fraser, Epidemiological dynamics of bacteriocin
competition and antibiotic resistance. *Proc Biol Sci* **289**, 20221197 (2022).

25.   H. Sati *et al.*, The WHO Bacterial Priority Pathogens List 2024: a prioritisation study to guide
research, development, and public health strategies against antimicrobial resistance. *Lancet*
Infect Dis **25**, 1033-1043 (2025).

26.   M. Kilian *et al.*, Evolution of *Streptococcus pneumoniae* and its close commensal relatives.
PLoS One **3**, e2683 (2008).

27.   G. Pozzi *et al.*, Competence for genetic transformation in encapsulated strains of
Streptococcus pneumoniae: two allelic variants of the peptide pheromone. *J Bacteriol* **178**,
6087-6090 (1996).

28.   I. F. Nes, D. B. Diep, H. Holo, Bacteriocin diversity in *Streptococcus* and *Enterococcus*. *J*
Bacteriol **189**, 1189-1198 (2007).

29.   J. Meza-Torres *et al.*, Listeriolysin S: A bacteriocin from *Listeria monocytogenes* that induces
membrane permeabilization in a contact-dependent manner. *Proc Natl Acad Sci U S A* **118**,
e2108155118 (2021).

30.   L. García-Bayona, M. S. Guo, M. T. Laub, Contact-dependent killing by
Caulobacter crescentus via cell surface-associated, glycine zipper proteins. *Elife* **6** (2017).

31.   C. Oppegård, J. Schmidt, P. E. Kristiansen, J. Nissen-Meyer, Mutational analysis of putative
helix-helix interacting GxxxG-motifs and tryptophan residues in the two-peptide bacteriocin
lactococcin G. *Biochemistry* **47**, 5242-5249 (2008).

32.   B. Ekblad *et al.*, Structure-function analysis of the two-peptide bacteriocin plantaricin EF.
Biochemistry **55**, 5106-5116 (2016).

33. B. Ekblad, P. E. Kristiansen, NMR structures and mutational analysis of the two peptides
constituting the bacteriocin plantaricin S. *Sci Rep* **9**, 2333 (2019).

34. G. Moll *et al.*, Lactococcin G is a potassium ion-conducting, two-component bacteriocin. *J
Bacteriol* **178**, 600-605 (1996).

35. O. McAuliffe *et al.*, Lacticin 3147, a broad-spectrum bacteriocin which selectively dissipates
the membrane potential. *Appl Environ Microbiol* **64**, 439-445 (1998).

36. G. N. Moll *et al.*, Complementary and overlapping selectivity of the two-peptide bacteriocins
plantaricin EF and JK. *J Bacteriol* **181**, 4848-4852 (1999).

37. X. Zhang *et al.*, Two-peptide bacteriocin PlnEF causes cell membrane damage to
Lactobacillus plantarum. *Biochim Biophys Acta* **1858**, 274-280 (2016).

38. H. Wei, L. S. Håvarstein, Fratricide is essential for efficient gene transfer between
pneumococci in biofilms. *Appl Environ Microbiol* **78**, 5897-5905 (2012).

39. H. Steinmoen, E. Knutsen, L. S. Håvarstein, Induction of natural competence in *Streptococcus
pneumoniae* triggers lysis and DNA release from a subfraction of the cell population. *Proc Natl
Acad Sci U S A* **99**, 7681-7686 (2002).

40. S. Peterson, R. T. Cline, H. Tettelin, V. Sharov, D. A. Morrison, Gene expression analysis of
the *Streptococcus pneumoniae* competence regulons by use of DNA microarrays. *J Bacteriol*
182, 6192-6202 (2000).

41. R. Rimini *et al.*, Global analysis of transcription kinetics during competence development in
Streptococcus pneumoniae using high density DNA arrays. *Mol Microbiol* **36**, 1279-1292
(2000).

42. E. García, Structure, function, and regulation of LytA: the N-acetylmuramoyl-L-alanine
amidase driving the "suicidal tendencies" of *Streptococcus pneumoniae-a review.
Microorganisms* **13**, 827 (2025).

43. A. M. Whatmore, V. A. Barcus, C. G. Dowson, Genetic diversity of the streptococcal
competence (*com*) gene locus. *J Bacteriol* **181**, 3144-3154 (1999).

44. M. Carrolo, F. R. Pinto, J. Melo-Cristino, M. Ramirez, Pherotypes are driving genetic
differentiation within *Streptococcus pneumoniae. BMC Microbiol* **9**, 191 (2009).

45.   D. F. Vestrheim, P. Gaustad, I. S. Aaberge, D. A. Caugant, Pherotypes of pneumococcal
strains co-existing in healthy children. *Infect Genet Evol* **11**, 1703-1708 (2011).

46.   R. A. Aras, J. Kang, A. I. Tschumi, Y. Harasaki, M. J. Blaser, Extensive repetitive DNA
facilitates prokaryotic genome plasticity. *Proc Natl Acad Sci U S A* **100**, 13579-13584 (2003).

47.   D. Wang, S. Li, F. Guo, K. Ning, L. Wang, Core-genome scaffold comparison reveals the
prevalence that inversion events are associated with pairs of inverted repeats. *BMC Genomics*
18, 268 (2017).

48.   T. J. Treangen, A. L. Abraham, M. Touchon, E. P. Rocha, Genesis, effects and fates of repeats
in prokaryotic genomes. *FEMS Microbiol Rev* **33**, 539-571 (2009).

49.   C. Bogaardt, A. J. van Tonder, A. B. Brueggemann, Genomic analyses of pneumococci reveal
a wide diversity of bacteriocins - including pneumocyclicin, a novel circular bacteriocin. *BMC*
Genomics **16**, 554 (2015).

50.   J. S. Puls *et al.*, *Staphylococcus epidermidis* bacteriocin A37 kills natural competitors with a
unique mechanism of action. *ISME J* **18**, wrae044 (2024).

51.   S. Borgeaud, L. C. Metzger, T. Scrignari, M. Blokesch, The type VI secretion system of *Vibrio*
cholerae fosters horizontal gene transfer. *Science* **347**, 63-67 (2015).

52.   L. Zhu, Z. Kuang, B. A. Wilson, G. W. Lau, Competence-independent activity of pneumococcal
EndA mediates degradation of extracellular DNA and nets and is important for virulence. *PLoS*
One **8**, e70363 (2013).

53.   G. Salvadori, R. Junges, D. A. Morrison, F. C. Petersen, Overcoming the barrier of low
efficiency during genetic transformation of *Streptococcus mitis*. *Front Microbiol* **7**, 1009 (2016).

54.   L. A. Cowley *et al.*, Evolution via recombination: cell-to-cell contact facilitates larger
recombination events in *Streptococcus pneumoniae*. *PLoS Genet* **14**, e1007410 (2018).

55.   L. R. Marks, R. M. Reddinger, A. P. Hakansson, High levels of genetic recombination during
nasopharyngeal carriage and biofilm formation in *Streptococcus pneumoniae*. *mBio* **3**,
e00200-12 (2012).

56.   J. Majewski, P. Zawadzki, P. Pickerill, F. M. Cohan, C. G. Dowson, Barriers to genetic exchange between bacterial species: *Streptococcus pneumoniae* transformation. *J Bacteriol* **182**, 1016-1023 (2000).

57.   K. E. Stevens, D. Chang, E. E. Zwack, M. E. Sebert, Competence in *Streptococcus pneumoniae* is regulated by the rate of ribosomal decoding errors. *mBio* **2**, e00071-11 (2011).

58.   E. A. Campbell, S. Y. Choi, H. R. Masure, A competence regulon in *Streptococcus pneumoniae* revealed by genomic analysis. *Mol Microbiol* **27**, 929-939 (1998).

59.   C. E. Grant, T. L. Bailey, W. S. Noble, FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**, 1017-1018 (2011).

60.   S. Handem, B. Ferreira, C. Valente, R. Sá-Leão, Genomic and functional insights into commensal streptococci with anti-pneumococcal activity. *BMC Genomics* **26**, 577 (2025).

61.   M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2--approximately maximum-likelihood trees for large alignments. *PLoS One* **5**, e9490 (2010).

62.   I. Letunic, P. Bork, Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. *Nucleic Acids Res* **52**, W78-W82 (2024).

63.   M. Kjos *et al.*, Expression of *Streptococcus pneumoniae* bacteriocins is induced by antibiotics via regulatory interplay with the competence system. *PLoS Pathog* **12**, e1005422 (2016).

64.   H. Echlin, J. W. Rosch, Advancing genetic tools in *Streptococcus pneumoniae*. *Genes (Basel)* **11**, 965 (2020).

65.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012).

Figure Legends

Fig. 1. *Streptococcus mitis* genomes encode a plethora of competence-associated bacteriocins (*cab*).

(A) Core-SNP phylogeny of 253 *S. mitis* genomes and distribution of competence-induced

bacteriocins A and B (*cibAB*), dedicated immunity (*cibC*), and competence-associated bacteriocin

(*cab*) genes. (B) Prevalence of CibAB and 95 Cab variants in the *S. mitis* population. (C) Rarefaction

analysis of mature Cab amino acid sequences found in the *S. mitis* population.

Fig. 2. Cab bacteriocins are expressed at the late phase of competence. (A) Genetic organization of

cab cluster of strain *S. mitis* G22, containing two putative bacteriocin (*cabAB*) and an immunity protein

(*cabC*) upstream of the ABC-transporter *comAB* genes. The *cabABC* operon promoter (P*cabABC*)

contains five tandem copies of SigX-binding sites (SigX box; SigX is the alternative sigma factor that

controls gene expression during the late phase of competence). (B) Expression of *cabABC* in *S. mitis*

G22 containing a P*cabABC*-luciferase (*luc*) reporter, ectopically inserted in its genome. Expression of

cabABC was inferred by expression of *luc*. *S. mitis* was pre-grown in acidic medium to avoid natural

competence development and synthetic cognate competence-stimulating peptide (CSP$_{G22}$) was

added to wild-type *S. mitis* and mutants lacking competence regulators. (C) Dose-response of

cabABC expression across CSP$_{G22}$ concentrations for the *S. mitis luc* reporter containing either the

full-length P*cabABC* promoter (which contains five SigX boxes), or a truncated version containing

only the most proximal SigX box. (D) Number of SigX boxes within *cab* clusters of varied sizes. Data

in panels B and C represent three to four independent experiments. Unpaired *t*-tests were performed

to compare *cab* expression. **** $P < 0.0001$.

Fig. 3. *S. mitis* and *S. pneumoniae* crosstalk through production of distinct CSP pheromones. (A)

Effect of CSP pheromones of close streptococci on *cabABC* expression in *S. mitis* G22 and alignment

of tested CSP variants (purple – Q14 amino acid residue, present only in inducing peptides). *S. mitis*

G22 P*cabABC-luc* was pre-grown in acidic medium, to which exogenous CSP was added. (B) Effect

of CSP with substitutions in residue 14 in the activation of *cabABC* expression in *S. mitis* G22. (C)

Effect of CSP with substitutions in residue 14 in the activation of competence in *S. pneumoniae* of

pherotype CSP-1 (P*comCDE-luc*). Dotted lines represent average luminescence value for nontreated

controls. Experiments were repeated three times, independently. Unpaired *t*-tests were performed to

compare *cab* expression. **** $P < 0.0001$. ns not significant.

Fig. 4. *S. mitis* bacteriocins CabAB mediate contact-dependent growth inhibition of *S. pneumoniae.* Experiments were done using *S. mitis* G22 Δ*blp4*Δ*blp1* and *S. pneumoniae* D39. The bacteriocin clusters *blp4* and *blp1* were first deleted from the genome of *S. mitis* G22 as they have anti-pneumococcal activity (23), which could obscure CabAB-mediated effects. (A) Proportion of *S. mitis* and *S. pneumoniae* in dual-species biofilm grown for 24 h in competence-permissive conditions. *S. mitis* mutant p(*cab*⁺) contains the native *cabABC* copy, and an additional copy under constitutive regulation (promoter *p23*) in a multicopy plasmid. (B) *S. mitis* was grown in competence-permissive conditions. Cell-free supernatants (CFS) were recovered and cell pellets were boiled. CFS and heat-killed cells were added to *S. pneumoniae* cultures and growth was monitored. (C) Bacterial loads of *S. pneumoniae* grown in the presence of *S. mitis* in direct contact (left) or separated with a 0.4 µm pore transwell membrane (right). (D) Corresponding bacterial loads of *S. mitis* G22 in experiment (C). (E) Representative images of CabA-HA and CabB-HA immunodetection on the surface of live competent *S. mitis* cells containing the native *cabABC* (negative control), *cabA-HA cabBC* or *cabA cabB-HA cabC* genotypes. *S. mitis* cultures were induced with CSP$_{G22}$, incubated with a primary anti-HA rabbit antibody, and then with an Alexa Fluor 488-conjugated goat anti-rabbit secondary antibody. At least ten independent fields were detected per experiment/condition. Experiments were repeated at least three times, independently. The ratio paired *t*-test was used to compare bacterial proportions in dual-species biofilms. The two-tailed unpaired Mann-Whitney U test was used to compare bacterial loads. ** *P* < 0.01. *** *P* < 0.001. ns not significant.

Fig. 5. CabAB is a two-peptide bacteriocin with LytA-dependent bactericidal activity against *S. pneumoniae.* (A) *S. pneumoniae* was treated with synthetic CabA, CabB, an equimolar mixture of CabA and CabB, or with DMSO (solvent where peptides were resuspended). Growth was recorded for 24 h. (B) Presence of GxxxG and GxxxG-like amino acid motifs in CabA and CabB sequences. (C) Exponential cultures of wild-type and autolysin-deficient *S. pneumoniae* were treated with DMSO (grey) or synthetic CabAB (purple). Growth was recorded for 24 h. For easier visualization of lysis

(decrease in optical density), the first 12 h of growth is represented. Experiments were repeated three

times independently.

Fig. 6. CabAB leads to permeabilization and depolarization of *S. pneumoniae* membranes, leading

to metabolite leakage. (A) *S. pneumoniae* treated with DMSO or CabAB and imaged by scanning

electron microscopy (SEM). (B) Quantification of membrane-permeable *S. pneumoniae* (stained with

propidium iodine (PI)) upon treatment with DMSO or increasing concentrations of CabAB. (C-D)

Quantification of populational membrane permeability (C) and depolarization (D) upon DMSO or

CabAB treatment at several timepoints. (E) Proportion of extracellular *vs.* total ATP in wild-type and

Δ*lytA S. pneumoniae* cultures upon DMSO or CabAB treatment. (F) Representative transmission

electron microscopy (TEM) image and quantification of intracellular aggregates in DMSO- (n = 118

cells) or CabAB- treated *S. pneumoniae* (n = 138 cells). (G) Labeling of large intracellular aggregates

by membrane dye FM4-64 in CabAB-treated *S. pneumoniae*. SEM and TEM were done once. Other

data was taken from at least three independent experiments.

Fig. 7. Expression of *cabAB* facilitates horizontal gene transfer in dual-species biofilms. (A)

Experimental setup to study gene transfer in dual-species biofilms. The *S. pneumoniae* DNA donor is

nontransformable due to a *comGC* deletion, and contains chloramphenicol and tetracycline resistance

cassettes at opposite ends of the chromosome (Chl[R] and Tet[R]); the *S. mitis* DNA receiver is labeled

with a kanamycin resistance cassette (Kan[R]); biofilms were pre-grown at a starting 1:1 ratio for 24 h

in acidic medium to prevent natural competence induction; cells were aspirated and fresh medium

containing CSP$_{G22}$ was added to biofilms to transiently induce competence; after 6 h of incubation,

cells were plated; *S. mitis* transformants were recovered and counted by selection in plates with

kanamycin and combinations of chloramphenicol and/or tetracycline. (B-C) Acquisition of Chl[R] and/or

Tet[R] by *S. mitis* in biofilms containing purified donor gDNA (B) or the strain itself (C). (D) Growth

inhibition and lysis of 26 bacterial species when treated with CabAB *vs.* DMSO during their

exponential phase. Growth was recorded for 24 h. Lysis is shown when a decrease from the initial

culture optical density was observed in all replicates. Experiments were repeated three times,

independently. Transformation efficiencies were compared using unpaired *t*-tests. LOD limit of

detection. * *P* < 0.05. ** *P* < 0.01.

Figure 1

Figure 2

Figure 3

A

Luminescence (RLU.OD$_{595}^{-1}$.10^4)

Addition of CSP from:
- *S. pneumoniae* (red)
- *S. oralis* (blue)
- *S. mitis* (green)

**** / **** / ns

Nontreated, CSP$_{G22}$, CSP-1, CSP-2, CSP$_{A22}$, CSP$_{B22}$, CSP$_{D22}$, CSP$_{E22}$

CSP type	Amino acid sequence	Identity
CSP$_{G22}$	E S R L S R L L R D F I F Q I K Q	-
CSP-1	E M R L S K F F R D F I L Q R K K	10/17
CSP-2	E M R I S R I I L D F L F L R K K	8/17
CSP$_{A22}$	D W R I S E T I R N L I F P R R K	5/17
CSP$_{B22}$	E S R V S R I I L D F L F L R K K	9/17
CSP$_{D22}$	D S R I S K F L R D F I F P R K K	10/17
CSP$_{E22}$	E M R L P K I L R D F I F P R K K	10/17

B

Luminescence (RLU.OD$_{595}^{-1}$.10^4)

S. mitis G22

ns / **** / **** / ns / ns

WT	WT	Q14L	Q14N	WT	L14Q
CSP-1	CSP$_{G22}$			CSP-2	

C

Luminescence (RLU.OD$_{595}^{-1}$.10^6)

S. pneumoniae D39
(CSP-1 pherotype)

ns / **** / **** / ns / ****

WT	WT	Q14L	Q14N	WT	L14Q
CSP-1	CSP$_{G22}$			CSP-2	

Figure 4

Figure 5

A

S. pneumoniae D39 treated with:

DMSO · CabA · CabB · CabA + CabB

B

CabA

IAPLVV**A**FGV**A**AFK**G**F**S**VTG**GA**IIL**G**AGLYQVGKNLK

CabB

VLP**A**AVV**A**VPV**G**IKI**A****G**YVV**G**GLS**A**VGL**A**AW**G**YFR

Gxxx**G** motif   **G**xxx**G**-like motif

C

WT

ΔlytA

ΔlytC

Figure 6

Figure 7

Fig. S1. Alignment-based clustering of Cab variants. The 95 Cab variants found *S. mitis* populations were clustered based on pairwise sequence alignments (see Materials and Methods). Alignments for Cab sequences within each cluster is shown, with associated Alphafold2 (ColabFold v1.5.5) structural model of the most prevalent variant. Amino acids are colored according to biochemical traits (RasMol coloring). Structures are colored according to plDDT (predicted local distance difference test) confidence.

Fig. S1. (cont.)

Fig. S2. *S. mitis* competence is inhibited in medium with acidic pH. *S. mitis* G22 harboring an early-competence reporter (P*comCDE-luc*) was pre-grown in medium at pH 6.6, before being diluted 1:100 in fresh medium at pH 6.6, 7.4, or 8.2. Optical density (closed circles) and luminescence (open circles) values were measured every 10 min for 24 h. Experiment was repeated three times independently.

Fig. S3. Expression of *cabABC* follows the same dynamics of other competence genes. (A) Expression of *comCDE*, *sigX1*, *sigX2* and *cabABC* evaluated by luminescence produced by luminescent reporter *S. mitis*. Cells were pre-grown in acidic pH and diluted in fresh medium containing synthetic CSP_{G22}. Values represent the highest expression value ($RLU.OD_{595}^{-1}$) obtained during the 3 h post CSP_{G22} treatment. (B) Dynamics of *comCDE*, *sigX1*, *sigX2* and *cabABC* expression in *S. mitis*. Optical density – closed circles. Luminescence – open circles. Experiments were repeated 3 times independently for 24 h. For clarity only the first 8 h are shown, since there is no visible expression of the four reporters after this period.

Fig. S4. Expression of *cabABC* is not induced by bacteriocin-like peptide pheromones (BlpC). *S. mitis* G22 harboring a P*cabABC-luc* reporter was pre-grown in acidic pH and diluted in fresh medium containing 1 μM of synthetic CSP_{G22} (grey), 1 μM of $BlpC_1$ or $BlpC_4$ pheromones (blue) or left untreated (dashed line). $BlpC_1$ or $BlpC_4$ are encoded in two *blp*-type bacteriocin clusters elsewhere in the genome of G22.

Fig. S5. Expression of *cab* by *S. mitis* reduces *S. pneumoniae* loads in dual-species biofilms. Figure individual and total bacterial loads of biofilm proportions presented in Figure 4A of the main text. *S. pneumoniae* and *S. mitis* of different *cab* genotypes were grown in 24 h biofilms in competence-permissive conditions. *S. mitis* and *S. pneumoniae* cell loads were enumerated by plating in BA with kanamycin or chloramphenicol, respectively. The experiment was repeated three times, independently.

Fig. S6. Permeabilization of *S. pneumoniae* membranes does not depend on autolysin LytA. Confocal imaging of wild-type and Δ*lytA S. pneumoniae* treated with DMSO or synthetic CabAB peptides. Membrane permeabilization is shown in red, by entry of propidium iodine. The total number and percentage of permeable cells is shown next to panels. Microscopy images are representative of three independent experiments.

Fig. S7. CabAB treatment induces formation of large clumps of *S. pneumoniae* cells encased in a matrix of DNA and membrane material. *S. pneumoniae* D39 was treated with DMSO or CabAB and stained with FM4-64 (membrane dye) and DAPI (nucleic acid dye), before being imaged in a confocal microscope. Experiment was repeated three times, independently.

Fig. S8. Individual growth curves of 18 streptococcal species and their sensitivity to CabAB. Exponential cultures of streptococcal species were diluted to OD_{600} 0.1 in appropriate medium and treated with either DMSO or CabAB. Optical density was measured every 10 min for 24 h. For clarity, the first 12 h are shown. Species are colored according to streptococcal group. The experiment was repeated three times, independently. Summary data of the growth curves can be found in the main manuscript.

Fig. S9. Individual growth curves of 4 non-streptococcal Gram-positive and 4 Gram-negative bacteria and their sensitivity to CabAB. Exponential cultures of species were diluted to OD_{600} in appropriate medium and treated with either DMSO or CabAB. Optical density was measured every 10 min for 24 h. For clarity, the first 12 h are shown. Species are colored according to Gram staining. The experiment was repeated three times, independently. Summary data of the growth curves can be found in the main manuscript.

Table S1. *Streptococcus mitis* genomes used in the study.

Strain	Region flanked by *plsX-acpP* and comAB recovered from genome (Y/N)	Presence of *cab* (Y/N)	*cab* locus length (bp)	Number of unique Cabs	Number of SigX boxes in *cab* locus	Presence of *cibAB* (Y/N)	Presence of *cibC* (Y/N)	Mature CSP sequence	4th last amino acid in CSP	Genbank accession	Location of region flanked by *plsX-acpP* and *comAB*
103U_S92	Y	Y	9862	4	20	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946186075.1	23700 to 33569 (contig 5)
125U_S4	Y	N	-	-	-	N	N	EMRRSNNNFFNFLRRI	L	GCA_946184935.1	121366 to 123329 (contig 6)
128U_S6	N	-	-	-	-	-	-		-	GCA_946186995.1	-
12P_S5	Y	Y	6869	5	15	N	N	EIRQTHNIFFNFFKRR	F	GCA_946186085.1	27830 to 34698 (contig 6)
13P_S6	Y	Y	9282	5	16	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946187105.1	25159 to 34440 (contig 14)
14P_S7	Y	Y	2287	2	3	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946186685.1	19479 to 21765 (contig 2)
161U_S19	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946188375.1	109310 to 116321 (contig 4)
1P_S1	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946183265.1	10854 to 17772 (contig 11)
22P_S11	N	-	-	-	-	-	-		-	GCA_946183395.1	-
27P_S14	Y	Y	7997	5	17	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946184735.1	22263 to 30259 (contig 10)
32P_S19	Y	Y	6869	5	15	N	N	EIRQTHNIFFNFFKRR	F	GCA_946223905.1	28620 to 35488 (contig 11)
38P_S23	Y	Y	6883	3	17	N	N	ESRLSKLLRDFIFKIKQ	K	GCA_946183455.1	26015 to 32897 (contig 9)
3P_S2	N	-	-	-	-	-	-		-	GCA_946183495.1	-
44P_S26	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946186695.1	36560 to 43673 (contig 10)
50P_S29	Y	Y	7547	4	15	Y	Y	ESRLSRLLRDFIFQIKQ	Q	GCA_946188075.1	25384 to 32930 (contig 8)
52P_S30	N	-	-	-	-	-	-		-	GCA_946186135.1	-
61P_S37	N	-	-	-	-	-	-		-	GCA_946183505.1	-
62P_S38	Y	Y	7457	5	15	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946183485.1	24444 to 24837 (contig 5)
68U_S84	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_946183445.1	13397 to 18312 (contig 11)
7P_S3	N	-	-	-	-	-	-		-	GCA_946185985.1	-
B1_S23	Y	Y	3685	6	4	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946189985.1	18059 to 21743 (contig 6)
B10_S32	N	-	-	-	-	-	-		-	GCA_946183015.1	-
B100_S21	N	-	-	-	-	-	-		-	GCA_946186585.1	-
B1015B24C5	Y	Y	8698	4	17	N	Y	EMRKSNSTFFHFLRRI	L	GCA_905220945.1	27478 to 36175 (contig 5)
B1015G24C3	Y	Y	3544	6	7	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_905221115.1	3577 to 7120 (contig 16)
B1015H24C1	Y	Y	9990	4	18	Y	Y	ESRVSDLILDFLFQRKK	Q	GCA_905221235.1	24516 to 34505 (contig 8)
B1015H24C3	Y	N	-	-	-	N	N	ESRLPKILLDFLFLRKK	L	GCA_905221375.1	45744 to 49408 (contig 11)
B1015H24C4	Y	N	-	-	-	N	N	EMRRIGSVLLNFFKRR	F	GCA_905221455.1	110170 to 117841 (contig 8)
B102_S23	N	-	-	-	-	-	-		-	GCA_946183435.1	-
B105_S26	Y	Y	3557	4	9	N	Y	EMRKMNEKSFNFFNIFRRR	I	GCA_946185425.1	22400 to 25956 (contig 13)
B106_S27	Y	Y	6472	5	16	N	N	ESRLSRLLRDFILQRKK	Q	GCA_946184175.1	27052 to 33524 (contig 3)
B107_S28	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946184385.1	73855 to 74301 (contig 3)
B110_S31	Y	N	-	-	-	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946185245.1	58142 to 67806 (contig 5)
B111_S32	Y	N	-	-	-	Y	Y	EIRIADKIFFNLFKRR	F	GCA_946185595.1	35645 to 42337 (contig 8)
B112_S33	Y	Y	8000	5	17	N	Y	EMRLPKILRDFIFPRKK	P	GCA_946185795.1	16961 to 24960 (contig 6)
B113_S34	Y	Y	7352	5	14	N	N	EMRKPDGALFNLFRRR	F	GCA_946186035.1	28045 to 35396 (contig 4)
B114_S35	Y	Y	7477	1	16	Y	Y	ESRLPKILLDFPRKK	P	GCA_946184755.1	130119 to 137595 (contig 5)
B116_S37	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946186655.1	65112 to 74936 (contig 6)
B117_S38	Y	N	-	-	-	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946184245.1	112994 to 121923 (contig 7)
B119_S40	Y	N	-	-	-	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946184705.1	31961 to 19319 (contig 13)
B12_S34	Y	Y	6838	5	15	N	N	EMRKPDGALFNLFRRR	F	GCA_946187535.1	27777 to 34614 (contig 5)
B121_S42	Y	N	-	-	-	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_946184465.1	3398 to 5219 (contig 19)
B122_S43	Y	Y	9952	5	14	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946183295.1	27547 to 37498 (contig 7)
B124_S44	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946184765.1	33241 to 41252 (contig 13)
B125_S45	Y	Y	6733	3	14	Y	Y	EIRIADKIFFNLFKRR	F	GCA_946186825.1	10414 to 17146 (contig 9)
B126_S46	Y	Y	7012	5	14	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946183535.1	26548 to 33559 (contig 6)
B128_S47	N	-	-	-	-	-	-		-	GCA_946187305.1	-
B129_S48	N	-	-	-	-	-	-		-	GCA_946183555.1	-
B13_S35	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946184335.1	124084 to 131222 (contig 4)
B132_S51	Y	Y	5685	5	11	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946188115.1	26091 to 31775 (contig 7)
B133-2_S52	N	-	-	-	-	-	-		-	GCA_946187925.1	-
B134_S53	N	-	-	-	-	-	-		-	GCA_946183695.1	-
B137_S55	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_946183685.1	37817 to 47227 (contig 10)
B139_S57	Y	N	-	-	-	Y	Y	ESRMPKILRDFIFPRKK	P	GCA_946183965.1	129975 to 133398 (contig 6)
B14_S36	Y	N	-	-	-	Y	Y	DMRISESIRNLIFPRKKK	R	GCA_946188365.1	132557 to 140438 (contig 5)

(continues on next page)

Table S1. (cont.)

B140_S58	Y	Y	7981	7	17	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946188035.1	21526 to 21918 (contig 7)
B146_S64	N	-	-	-	-	-	-		-	GCA_946184845.1	-
B147_S65	Y	N	-	-	-	Y	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946188285.1	33777 to 37954 (contig 129
B148_S66	N	-	-	-	-	-	-		-	GCA_946185855.1	-
B15_S37	Y	N	-	-	-	N	N	EMRKMNDKSFNLFNFFRRR	F	GCA_946188735.1	108533 to 116391 (contig 7)
B151_S69	N	-	-	-	-	-	-		-	GCA_946185905.1	-
B153_S71	Y	Y	10165	3	13	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188725.1	12287 to 22451 (contig 7)
B155_S73	N	-	-	-	-	-	-		-	GCA_946185295.1	-
B156_S74	N	-	-	-	-	-	-		-	GCA_946189305.1	-
B157_S75	Y	N	-	-	-	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_946188715.1	47861 to 48593 (contig 11)
B158_S76	N	-	-	-	-	-	-		-	GCA_946188825.1	-
B159_S77	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188815.1	126333 to 135213 (contig 1)
B161_S79	Y	N	-	-	-	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946188505.1	121745 to 133038 (contig 7)
B162_S80	N	-	-	-	-	-	-		-	GCA_946188645.1	-
B163_S81	N	-	-	-	-	-	-		-	GCA_946188665.1	-
B165_S83	Y	Y	7454	5	12	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946188525.1	9840 to 17293 (contig 6)
B166_S84	N	-	-	-	-	-	-		-	GCA_946188555.1	-
B167_S85	Y	N	-	-	-	N	Y	ESRLPKILLDFLFLRKK	L	GCA_946188435.1	51984 to 55661 (contig 7)
B168_S86	Y	N	-	-	-	N	N	ESRLPKILLDFLFLRKK	L	GCA_946188425.1	129601 to 136812 (contig 5)
B169_S87	Y	N	-	-	-	N	Y	EMRLPKILRDFIFPRKK	P	GCA_946189595.1	100635 to 105207 (contig 7)
B170_S88	Y	Y	2522	2	7	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946189625.1	16848 to 19369 (contig 119
B172_S90	N	-	-	-	-	-	-		-	GCA_946189565.1	-
B173_S91	N	-	-	-	-	-	-		-	GCA_946188785.1	-
B174_S92	Y	N	-	-	-	N	Y	EMRKPDGALFNLFRRR	F	GCA_946189255.1	44630 to 50769 (contig 13)
B176_S94	N	-	-	-	-	-	-		-	GCA_946188865.1	-
B19_S40	Y	Y	6430	2	12	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_946190005.1	24129 to 30558 (contig 12)
B20_S41	N	-	-	-	-	-	-		-	GCA_946190035.1	-
B21_S42	N	-	-	-	-	-	-		-	GCA_946189125.1	-
B22_S43	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946189035.1	45816 to 51979 (contig 12)
B26_S45	Y	Y	7324	1	14	N	N	EMRRIDKIFINFLKRR	L	GCA_946189055.1	36560 to 43883 (contig 12)
B3_S25	Y	Y	6276	6	13	Y	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946189635.1	27034 to 33309 (contig 11)
B30_S48	Y	N	-	-	-	Y	Y	ESRVSDILLDFLFRRKK	R	GCA_946189895.1	13875 to 25318 (contig 11)
B31_S49	Y	Y	7447	5	12	Y	Y	EMRKMNDKSFNLFNFFRRR	F	GCA_946188485.1	24297 to 31743 (contig 13)
B33_S51	Y	Y	11446	5	20	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188605.1	9714 to 21159 (contig 14)
B34_S52	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFNIFRRR	F	GCA_946189235.1	27396 to 35035 (contig 17)
B37_S55	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946188755.1	8105 to 15978 (contig 6)
B38_S56	N	-	-	-	-	-	-		-	GCA_946188795.1	-
B4_S26	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946189945.1	13981 to 20965 (contig 29)
B42_S60	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188675.1	147283 to 158055 (contig 5)
B43_S61	Y	Y	7325	4	16	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946188595.1	23294 to 30618 (contig 15)
B44_S62	Y	Y	6767	5	14	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188765.1	24640 to 31406 (contig 5)
B45_S63	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_946189355.1	8513 to 9245 (contig 6)
B46_S64	N	-	-	-	-	-	-		-	GCA_946188655.1	-
B47_S65	Y	N	-	-	-	N	Y	EMRKSNNNFFHFLRRI	L	GCA_946188745.1	43466 to 50055 (contig 7)
B49_S67	N	-	-	-	-	-	-		-	GCA_946188965.1	-
B5_S27	N	-	-	-	-	-	-		-	GCA_946189025.1	-
B50_S68	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188955.1	126680 to 135327 (contig 7)
B52_S70	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946189865.1	13665 to 14681 (contig 16)
B54_S72	N	-	-	-	-	-	-		-	GCA_946188925.1	-
B55_S73	Y	N	-	-	-	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_946188895.1	49166 to 58656 (contig 9)
B56_S74	N	-	-	-	-	-	-		-	GCA_946188995.1	-
B57_S75	Y	N	-	-	-	N	Y	EMRRIDKMFINFLKRR	L	GCA_946188985.1	117917 to 125697 (contig 4)
B59_S77	N	-	-	-	-	-	-		-	GCA_946188875.1	-
B60_S78	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_946189965.1	124476 to 131218 (contig 4)
B61_S79	Y	N	-	-	-	Y	Y	EIRKSNSALVNFFKRR	F	GCA_946188855.1	55573 to 66461 (contig 10)
B63_S81	Y	Y	7221	5	14	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_946189015.1	10888 to 18108 (contig 15)
B65_S83	Y	N	-	-	-	N	N	ESRVSDILLDFLFRRKK	R	GCA_946189955.1	118327 to 122899 (contig 3)
B67_S85	Y	Y	3156	4	10	Y	Y	EMRKSNNTFFNFLRRI	L	GCA_946188885.1	7767 to 10922 (contig 13)
B69_S87	N	-	-	-	-	-	-		-	GCA_946188945.1	-
B73_S91	Y	Y	8931	3	19	N	N	ESRVSDILLDFLFRRKK	R	GCA_946188935.1	13555 to 22485 (contig 9)

(continues on next page)

B74_S92	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_946188975.1	10138 to 10870 (contig 11)
B76_S94	Y	Y	7437	4	12	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_946189935.1	30822 to 38258 (contig 8)
B77_S95	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946189755.1	118938 to 126810 (contig 5)
B78_S96	Y	Y	8714	4	21	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946189875.1	25049 to 33762 (contig 5)
B79_S1	Y	Y	3832	6	10	Y	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946188405.1	8778 to 12609 (contig 18)
B82_S4	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946188465.1	68030 to 72644 (contig 9)
B85_S7	Y	N	-	-	-	N	N	ESRLPKIRFDFIFPRKK	P	GCA_946189515.1	124429 to 125974 (contig 6)
B88_S10	Y	N	-	-	-	N	Y	EMRLPKILRDFIFPRKK	P	GCA_946189785.1	67460 to 77667 (contig 6)
B89L_S11	Y	Y	8758	4	19	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946188455.1	23760 to 32517 (contig 5)
B9_S31	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_946189075.1	32842 to 41229 (contig 13)
B91_S12	Y	N	-	-	-	Y	Y	EMRLSKFLRDFILQRKN	Q	GCA_946189535.1	19728 to 28296 (contig 10)
B93_S14	N	-	-	-	-	-	-			GCA_946189095.1	-
B95_S16	Y	N	-	-	-	N	N	MKNTVKLEQFVALKEKTCRR FEVGKLEKQLIVYLISSKEDK	K	GCA_946188475.1	57945 to 58714 (contig 10)
B96_S17	Y	Y	7329	1	16	N	N	EIRIADKIFFNLKRR	F	GCA_946189115.1	35203 to 42531 (contig 12)
B97_S18	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_946189825.1	102775 to 103025 (contig 4)
B99_S20	N	-	-	-	-	-	-			GCA_946189005.1	-
GCF_000027165.1_ASM2716v1	Y	Y	9068	5	18	Y	Y	EMRKPDGALFNLFRRR	F	GCA_000027165.1 ASM2716v1	53854 to 62921 (contig 1)
GCF_000148505.1_ASM14850v1	N	-	-	-	-	-	-			GCA_000148505.1 ASM14850v1	-
GCF_000148525.1_ASM14852v1	Y	Y	4573	7	7	N	N	ESRVSDILLDFLFRRKK	R	GCA_000148525.1 ASM14852v1	23154 to 27726 (contig 22)
GCF_000148545.1_ASM14854v1	N	-	-	-	-	-	-			GCA_000148545.1 ASM14854v1	-
GCF_000148585.2_ASM14858v3	Y	Y	8195	5	17	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_000148585.2 ASM14858v3	46590 to 54784 (contig 1)
GCF_000220045.1_ASM22004v2	Y	N	-	-	-	N	N	ESRLSRLLRDFIFQIKQ	Q	GCA_000220045.1 ASM22004v2	113804 to 121511 (contig 10)
GCF_000220085.1_ASM22008v2	Y	Y	7601	3	18	N	N	EMRRIDKIFNFLKRR	L	GCA_000220085.1 ASM22008v2	33312 to 40912 (contig 6)
GCF_000222785.1_ASM22278v2	Y	N	-	-	-	Y	Y	EMRLPKILRDFIFPRKK	P	GCA_000222785.1 ASM22278v2	80829 to 88913 (contig 17)
GCF_000257805.1_ASM25780v1	Y	N	-	-	-	N	N	EMRRIDKIFINFLKRR	L	GCA_000257805.1 ASM25780v1	30349 to 36794 (contig 56)
GCF_000257865.1_ASM25786v1	Y	N	-	-	-	N	N	EMRKMNEKSFNFFNIFRRR	F	GCA_000257865.1 ASM25786v1	108462 to 111048 (contig 56)
SK579v1.0	Y	N	-	-	-	N	N	EIRKSNSALVNFFKRR	F	GCA_000257905.1 SmitisSK579v1.0	51648 to 53976 (contig 13)
GCF_000385815.1_ASM38581v1	N	-	-	-	-	-	-			GCA_000385815.1 ASM38581v1	-
Smit.str17_34_v1.0	Y	Y	8147	3	16	N	N	ESRISDLLLDFLFQRKK	Q	GCA_000430305.1 Smit.str17_34_v1.	24690 to 32836 (contig 5)
Smit.str18_56_v1.0	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_000430345.1 Smit.str18_56_v1.	6585 to 7319 (contig 12)
Smit.str29_42_v1.0	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_000430365.1 Smit.str29_42_v1.	28257 to 34755 (contig 8)
Sm.str21_39_v1.0	Y	N	-	-	-	N	N	EMRLPKILRDFIFPRKK	P	GCA_000506725.1 Sm.str21_39_v1.0	107853 to 111953 (contig 6)
Sm.str27_7_v1.0	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_000506765.1 Sm.str27_7_v1.0	48091 to 54587 (contig 8)
GCF_000722685.1_ASM72268v1	Y	Y	6593	3	15	N	N	EIRKSNSALVNFFKRR	F	GCA_000722685.1 ASM72268v1	33237 to 39817 (contig 3)
GCF_000722695.1_ASM72269v1	Y	Y	8106	4	20	N	Y	EMRLPKILRDFIFPRKK	P	GCA_000722695.1 ASM72269v1	28640 to 36745 (contig 2)
GCF_000722705.1_ASM72270v1	Y	Y	7521	3	19	N	Y	EMRRIDKMFINFLKRR	L	GCA_000722705.1 ASM72270v1	26310 to 33830 (contig 1)
GCF_000722755.1_ASM72275v1	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_000722755.1 ASM72275v1	19662 to 20232 (contig 8)
GCF_000722765.2_ASM72276v2	Y	Y	7760	4	18	N	Y	EMRKSNNNFFHFLRRI	L	GCA_000722765.2 ASM72276v2	49608 to 57367 (contig 1)
GCF_000722775.1_ASM72277v1	Y	Y	8930	3	21	N	N	EIRKTSNSLLNFFKRR	F	GCA_000722775.1 ASM72277v1	27763 to 36692 (contig 6)
GCF_000722815.1_ASM72281v1	Y	Y	1789	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_000722815.1 ASM72281v1	20061 to 21849 (contig 1)
GCF_000722835.1_ASM72283v1	N	-	-	-	-	-	-			GCA_000722835.1 ASM72283v1	-
GCF_000724165.1_ASM72416v1	Y	N	-	-	-	N	N	Not found	-	GCA_000724165.1 ASM72416v1	438770 to 446249 (contig 12)
GCF_000960005.1_ASM96000v1	Y	Y	8611	2	18	Y	Y	EMRKPDGALFNLFRRR	F	GCA_000960005.1 ASM96000v1	29135 to 37745 (contig 1)
GCF_000960025.1_ASM96002v1	Y	Y	12265	4	24	N	N	ESRLPKIRFDFIFPRKK	P	GCA_000960025.1 ASM96002v1	32833 to 45097 (contig 3)
GCF_000960065.1_ASM96006v1	Y	N	-	-	-	N	N	DMRISESIRNLIFPRKKK	R	GCA_000960065.1 ASM96006v1	33075 to 43002 (contig 5)
GCF_001070035.1_ASM107003v1	N	-	-	-	-	-	-			GCA_001070035.1 ASM107003v1	-
GCF_001070205.1_ASM107020v1	N	-	-	-	-	-	-			GCA_001070205.1 ASM107020v1	
GCF_001070375.1_ASM107037v1	N	-	-	-	-	-	-			GCA_001070375.1 ASM107037v1	-
GCF_001070705.1_ASM107070v1	Y	N	-	-	-	N	N	ESRLSKLLRDFIFKIKQ	K	GCA_001070705.1 ASM107070v1	91104 to 97864 (contig 13)
GCF_001070985.1_ASM107098v1	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_001070985.1 ASM107098v1	125336 to 135287 (contig 33)
GCF_001074635.1_ASM107463v1	N	-	-	-	-	-	-			GCA_001074635.1 ASM107463v1	-
GCF_001281025.1_ASM128102v1	Y	N	-	-	-	Y	Y	EMRKSNNNFFNFLRRI	L	GCA_001281025.1 ASM128102v1	326587 to 336055 (contig 1)
GCF_001546795.1_ASM154679v1	N	-	-	-	-	-	-			GCA_001546795.1 ASM154679v1	-
GCF_001560895.1_ASM156089v1	Y	Y	2200	2	5	N	N	EMRKSNNTFFNFLRRI	-	GCA_001560895.1 ASM156089v1	73674 to 75873 (contig 1)
GCF_001579045.1_ASM157904v1	Y	N	-	-	-	Y	Y	EMRLPKILRDFIFPRKK	P	GCA_001579045.1 ASM157904v1	416341 to 423519 (contig 63)
GCF_001588595.1_ASM158859v1	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_001588595.1 ASM158859v1	14693 to 15462 (contig 4)
GCF_001588605.1_ASM158860v1	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_001588605.1 ASM158860v1	34154 to 34922 (contig 15)
GCF_001588625.1_ASM158862v1	Y	Y	9205	5	23	N	N	EMRKPDGALFNLFRRR	F	GCA_001588625.1 ASM158862v1	9739 to 18943 (contig 27)
GCF_002005545.1_ASM200554v1	Y	Y	3980	6	9	N	N	EMRLPKILRDFIFPRKK	P	GCA_002005545.1 ASM200554v1	495219 to 499198 (contig 1)

(continues on next page)

GCF_002014755.1_ASM201475v1	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_002014755.1 ASM201475v1	115535 to 123729 (contig 6)
GCF_002014795.1_ASM201479v1	Y	Y	8075	4	21	N	Y	EMRKPDGALFNLFRRR	F	GCA_002014795.1 ASM201479v1	26394 to 34468 (contig 6)
GCF_002014875.1_ASM201487v1	Y	N	-	-	-	Y	Y	EIRQTHNIFFNFFKRR	F	GCA_002014875.1 ASM201487v1	63548 to 69821 (contig 7)
GCF_002096735.1_ASM209673v1	Y	Y	9752	2	14	N	N	EMRKSNNNFFNFLRRI	L	GCA_002096735.1 ASM209673v1	14476 to 24227 (contig 13)
GCF_002096755.1_ASM209675v1	N	-	-	-	-	-	-	-	-	GCA_002096755.1 ASM209675v1	-
GCF_002096765.1_ASM209676v1	Y	Y	6682	6	16	Y	Y	EMRKPDGALFNLFRRR	F	GCA_002096765.1 ASM209676v1	26701 to 33382 (contig 48)
GCF_002096775.1_ASM209677v1	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_002096775.1 ASM209677v1	23754 to 24138 (contig 24)
GCF_002096815.1_ASM209681v1	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_002096815.1 ASM209681v1	14082 to 19251 (contig 18)
GCF_002096835.1_ASM209683v1	Y	N	-	-	-	Y	Y	ESRVSDILLDFLFRRKK	R	GCA_002096835.1 ASM209683v1	4746 to 5469 (contig 19)
GCF_002096845.1_ASM209684v1	N	-	-	-	-	-	-	-	-	GCA_002096845.1 ASM209684v1	-
GCF_002096895.1_ASM209689v1	Y	N	-	-	-	N	N	DSRISKYLRDFIFPRKK	P	GCA_002096895.1 ASM209689v1	125698 to 126269 (contig 19)
GCF_002096925.1_ASM209692v1	N	-	-	-	-	-	-	-	-	GCA_002096925.1 ASM209692v1	-
GCF_002096935.1_ASM209693v1	Y	Y	6733	3	14	N	N	EIRIADKIFFNLFKRR	F	GCA_002096935.1 ASM209693v1	24923 to 31655 (contig 25)
GCF_002860825.1_ASM286082v1	Y	Y	3524	6	5	N	N	EMRKSNNTFFNFLRRI	L	GCA_002860825.1 ASM286082v1	18587 to 22110 (contig 13)
GCF_002860865.1_ASM286086v1	Y	Y	2525	2	5	N	N	ESRVSDILDFLFQRKK	Q	GCA_002860865.1 ASM286086v1	58524 to 61048 (contig 3)
GCF_002860905.1_ASM286090v1	Y	Y	3156	4	10	Y	Y	EMRRIDKIFINFLKRR	L	GCA_002860905.1 ASM286090v1	18653 to 21808 (contig 8)
GCF_002871315.1_ASM287131v1	Y	Y	3524	6	5	N	N	EMRKSNNTFFNFLRRI	L	GCA_002871315.1 ASM287131v1	18509 to 22032 (contig 13)
GCF_003942775.1_ASM394277v1	N	-	-	-	-	-	-	-	-	GCA_003942775.1 ASM394277v1	-
GCF_003942885.1_ASM394288v1	N	-	-	-	-	-	-	-	-	GCA_003942885.1 ASM394288v1	-
GCF_003942965.1_ASM394296v1	Y	N	-	-	-	N	N	Not found	-	GCA_003942965.1 ASM394296v1	11262 to 11512 (contig 3)
GCF_003942995.1_ASM394299v1	Y	N	-	-	-	Y	Y	ESRVSDILLDFLFRRKK	R	GCA_003942995.1 ASM394299v1	30869 to 40944 (contig 14)
GCF_003943915.1_ASM394391v1	N	-	-	-	-	-	-	-	-	GCA_003943915.1 ASM394391v1	-
GCF_003943965.1_ASM394396v1	Y	N	-	-	-	N	Y	EMRRSNNNFFNFLRRI	L	GCA_003943965.1 ASM394396v1	40655 to 42425 (contig 16)
GCF_005144795.1_ASM514479v1	Y	Y	1780	2	4	Y	Y	ESRLSKIFLDFIFPRKK	P	GCA_005144795.1 ASM514479v1	13402 to 15181 (contig 16)
GCF_005844455.1_ASM584445v1	N	-	-	-	-	-	-	-	-	GCA_005844455.1 ASM584445v1	-
GCF_008763935.1_FS623	Y	Y	6780	5	16	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_008763935.1 FS623	107171 to 113950 (contig 5)
GCF_009495695.1_ASM949569v1	N	-	-	-	-	-	-	-	-	GCA_009495695.1 ASM949569v1	-
GCF_009495705.1_ASM949570v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009495705.1 ASM949570v1	23517 to 30981 (contig 17)
GCF_009495735.1_ASM949573v1	Y	N	-	-	-	N	N	ESRVSDILDFLFQRKK	Q	GCA_009495735.1 ASM949573v1	88623 to 96087 (contig 8)
GCF_009495895.1_ASM949589v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009495895.1 ASM949589v1	23636 to 31100 (contig 8)
GCF_009495955.1_ASM949595v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009495955.1 ASM949595v1	58616 to 66080 (contig 10)
GCF_009496035.1_ASM949603v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496035.1 ASM949603v1	88623 to 96087 (contig 9)
GCF_009496045.1_ASM949604v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496045.1 ASM949604v1	88615 to 96079 (contig 9)
GCF_009496065.1_ASM949606v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496065.1 ASM949606v1	23577 to 31041 (contig 10)
GCF_009496075.1_ASM949607v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496075.1 ASM949607v1	23517 to 30981 (contig 11)
GCF_009496165.1_ASM949616v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496165.1 ASM949616v1	23517 to 30981 (contig 8)
GCF_009496185.1_ASM949618v1	N	-	-	-	-	-	-	-	-	GCA_009496185.1 ASM949618v1	-
GCF_009496205.1_ASM949620v1	N	-	-	-	-	-	-	-	-	GCA_009496205.1 ASM949620v1	-
GCF_009496295.1_ASM949629v1	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_009496295.1 ASM949629v1	124953 to 131821 (contig 6)
GCF_009496335.1_ASM949633v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496335.1 ASM949633v1	88622 to 96086 (contig 8)
GCF_009496355.1_ASM949635v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496355.1 ASM949635v1	88623 to 96087 (contig 7)
GCF_009496385.1_ASM949638v1	Y	Y	2273	2	6	N	N	EMRKSNNTFFNFLRRI	L	GCA_009496385.1 ASM949638v1	111359 to 113631 (contig 5)
GCF_009496395.1_ASM949639v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496395.1 ASM949639v1	23577 to 31041 (contig 8)
GCF_009496435.1_ASM949643v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496435.1 ASM949643v1	88623 to 96087 (contig 8)
GCF_009496505.1_ASM949650v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496505.1 ASM949650v1	23517 to 30981 (contig 6)
GCF_009496725.1_ASM949672v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496725.1 ASM949672v1	23577 to 31041 (contig 7)
GCF_009496735.1_ASM949673v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496735.1 ASM949673v1	23577 to 31041 (contig 7)
GCF_009496755.1_ASM949675v1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496755.1 ASM949675v1	18419 to 25883 (contig 15)
GCF_009496795.1_ASM949679v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496795.1 ASM949679v1	23577 to 31041 (contig 8)
GCF_009496805.1_ASM949680v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496805.1 ASM949680v1	23577 to 31041 (contig 8)
GCF_009496815.1_ASM949681v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496815.1 ASM949681v1	23577 to 31041 (contig 9)
GCF_009496825.1_ASM949682v1	N	-	-	-	-	-	-	-	-	GCA_009496825.1 ASM949682v1	-
GCF_009496835.1_ASM949683v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496835.1 ASM949683v1	23577 to 31041 (contig 9)
GCF_009496895.1_ASM949689v1	Y	Y	7465	7	16	N	N	EIRKTSNSLLNFFKRR	F	GCA_009496895.1 ASM949689v1	23517 to 30981 (contig 15)
GCF_009496905.1_ASM949690v1	Y	N	-	-	-	N	N	EIRKTINGLLNFFKRR	F	GCA_009496905.1 ASM949690v1	51468 to 56727 (contig 9)
GCF_009730515.1_ASM973051v1	Y	Y	8195	5	17	N	N	EIRQTHNIFFNFFKRR	-	GCA_009730515.1 ASM973051v1	31493 to 39687 (contig 1)
GCF_900459425.1_42042_B02	Y	N	-	-	-	N	N	EIRQTHNIFFNFFKRR	F	GCA_900459425.1 42042_B02	116797 to 124991 (contig 2)
MSCx	Y	Y	8886	4	18	Y	Y	DSRISKYLRDFIFPRKK	P	GCA_905187535.1	25487 to 34372 (contig 11)
NM2_1	Y	N	-	-	-	N	N	EIRKTSNSLLNFFKRR	F	GCA_905187515.1	32077 to 39698 (contig 13)

Table S1. (cont.)

NM4_2	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_905187545.1	32687 to 40308 (contig 13)
NM5_2	N	-	-	-	-	-	-		-	GCA_905187495.1	-
P11_S4	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_946189525.1	42441 to 48870 (contig 6)
P164_S40	N	-	-	-	-	-	-		-	GCA_946189105.1	-
P21_S10	Y	Y	7902	6	14	N	N	EMRKPDGALFNLFRRR	F	GCA_946189045.1	26201 to 34102 (contig 7)
P28_S15	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946188915.1	120578 to 128595 (contig 6)
P42_S25	N	-	-	-	-	-	-		-	GCA_946189085.1	-
P55_S33	Y	Y	7359	5	16	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946189995.1	28034 to 35392 (contig 7)
P59_S35	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946189495.1	151368 to 161798 (contig 4)
P60_S36	N	-	-	-	-	-	-		-	GCA_946190025.1	-
S1071B24C1	Y	Y	2587	2	9	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221505.1	3951 to 6537 (contig 12)
S1071B24C2	Y	Y	2461	2	8	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221525.1	3630 to 6090 (contig 13)
S1071G24C1	Y	N	-	-	-	N	N	EMRKIDDALFNLFNFFRRR	F	GCA_905221535.1	47372 to 58002 (contig 11)
S1071G24C2	Y	N	-	-	-	N	N	EMRKIDDALFNLFNFFRRR	F	GCA_905221545.1	47372 to 58002 (contig 11)
S1071G24C3	Y	Y	2461	2	8	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221005.1	3630 to 6090 (contig 12)
S1071G24C4	N	-	-	-	-	-	-		-	GCA_905221035.1	-
S1071G24C5	N	-	-	-	-	-	-		-	GCA_905221045.1	-
S1072B24C4	Y	Y	2873	4	7	N	N	EMRMLKSPFSFLFPRKK	P	GCA_905221065.1	23060 to 25952 (contig 4)
S1072B24C5	Y	Y	2873	4	7	N	N	EMRMLKSPFSFLFPRKK	P	GCA_905221055.1	23060 to 25952 (contig 4)
S1072G24C4	Y	N	-	-	-	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221105.1	26882 to 35329 (contig 7)
S1075M24C2	Y	N	-	-	-	Y	Y	ESRVADLLLDFLFQRKK	Q	GCA_905221085.1	103323 to 113219 (contig 6)
S1075M24C3	Y	N	-	-	-	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221095.1	31726 to 42537 (contig 11)
S1075M24C5	Y	Y	10812	2	26	N	N	ESRISDLLLDFLFQRKK	Q	GCA_905221075.1	27291 to 37102 (contig 8)
S1086G24C1	N	-	-	-	-	-	-		-	GCA_905221125.1	-
S1088B24C2	Y	Y	6473	2	18	N	Y	ESRVSDLILDFLFQRKK	Q	GCA_905221135.1	26672 to 33144 (contig 6)
S1088M24C1	Y	N	-	-	-	N	N	EMRIADKIFFNLFKRR	F	GCA_905221145.1	54156 to 65018 (contig 7)
S1091B24C4	Y	N	-	-	-	N	N	EMRKPDGALFNLFRRR	F	GCA_905221155.1	125261 to 134199 (contig 4)
S1092G24C1	Y	Y	1740	2	2	N	N	DMRISGSIRNLIFPRKKK	R	GCA_905221165.1	12912 to 14651 (contig 9)
S1092G24C3	Y	Y	1768	2	2	N	N	EIRKKSNSLLNFFKRR	F	GCA_905221175.1	13104 to 14871 (contig 5)
S1092G24C4	N	-	-	-	-	-	-		-	GCA_905221185.1	-
S1093G24C1	Y	Y	8698	4	17	N	Y	EMRKSNSTFFHFLRRI	L	GCA_905221195.1	27478 to 36175 (contig 5)
S1093G24C2	Y	N	-	-	-	N	N	ESRLPKILRDFIFPRKK	P	GCA_905221205.1	66159 to 73603 (contig 4)
S1093G24C3	Y	N	-	-	-	N	N	EMRKTSYSLLNLFKRR	F	GCA_905221225.1	104898 to 109878 (contig 7)
S1096G24C5	N	-	-	-	-	-	-		-	GCA_905221215.1	-
S1096M24C3	Y	Y	9580	5	22	N	N	EIRKTSNSLNFFKRR	F	GCA_905221245.1	11901 to 21480 (contig 6)
S1096M24C5	N	-	-	-	-	-	-		-	GCA_905221285.1	-
U10_S50	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946188615.1	136100 to 146454 (contig 6)
U104_S93	Y	Y	9859	4	20	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946188685.1	23700 to 33558 (contig 6)
U105_S94	N	-	-	-	-	-	-		-	GCA_946188575.1	-
U109_S95	Y	N	-	-	-	N	N	EIRKTSNSLNFFKRR	F	GCA_946188695.1	67998 to 77255 (contig 5)
U114_S96	Y	N	-	-	-	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946188415.1	84542 to 94928 (contig 4)
U117_S1	Y	N	-	-	-	N	N	DSRISKYLRDFIFPRKK	P	GCA_946188585.1	15202 to 15765 (contig 7)
U12_S51	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946224775.1	52874 to 63087 (contig 6)
U122_S2	Y	N	-	-	-	N	Y	EIRKTSNSLLNFFKRR	F	GCA_946189145.1	149195 to 157067 (contig 2)
U123_S3	Y	N	-	-	-	N	N	ESRVSDILLNFLFRRKK	R	GCA_946190065.1	49685 to 50417 (contig 16)
U126_S5	Y	Y	8396	4	19	Y	Y	ESRVSDLILDFLFQRKK	Q	GCA_946190085.1	25538 to 25928 (contig 3)
U13_S52	Y	N	-	-	-	N	N	ESRLSKLLRDFIFKIKQ	K	GCA_946190075.1	21460 to 34396 (contig 12)
U131_S7	Y	N	-	-	-	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946190095.1	127239 to 129511 (contig 5)
U134_S8	Y	Y	1964	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_946190055.1	12955 to 14918 (contig 6)
U135_S9	Y	Y	1964	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_946190045.1	12955 to 14918 (contig 6)
U14_S53	N	-	-	-	-	-	-		-	GCA_946190385.1	-
U143_S13	Y	N	-	-	-	N	N	EIRKKSNSLLNFFKRR	F	GCA_946190395.1	13391 to 14114 (contig 3)
U144_S14	Y	N	-	-	-	N	Y	EMRKMNEKSFNFFNIFRRR	F	GCA_946190245.1	31975 to 35539 (contig 15)
U148_S15	Y	Y	1964	3	4	N	N	EMRRSNNNFFNFLRRI	L	GCA_946190215.1	12955 to 14918 (contig 6)
U15_S54	Y	Y	3970	1	7	N	N	ESRVSDLILDFLFQRKK	Q	GCA_946190375.1	44742 to 48711 (contig 14)
U16_S55	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946190205.1	52874 to 63087 (contig 9)
U164_S21	Y	N	-	-	-	N	N	DSRISKYLRDFIFPRKK	P	GCA_946190365.1	56909 to 62429 (contig 9)
U165-2_S22	N	-	-	-	-	-	-		-	GCA_946186005.1	-
U23_S57	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190355.1	34140 to 39308 (contig 18)

(continues on next page)

Table S1. (cont.)

U24_S58	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190345.1	13215 to 18383 (contig 16)
U25_S59	N	-	-	-	-	-	-	-	-	GCA_946224995.1	-
U28_S60	N	-	-	-	-	-	-	-	-	GCA_946225005.1	-
U32_S62	Y	Y	6537	3	14	N	N	EMRRIDKIFINFLKRR	L	GCA_946224725.1	25517 to 32053 (contig 4)
U35_S65	Y	N	-	-	-	N	N	EMRRIDKIFINFLKRR	L	GCA_946224685.1	125817 to 132353 (contig 5)
U36_S66	N	-	-	-	-	-	-	-	-	GCA_946224705.1	-
U39_S69	N	-	-	-	-	-	-	-	-	GCA_946224645.1	-
U40_S70	Y	Y	1815	3	7	N	N	EIRKTINGLLNFFKRR	F	GCA_946190305.1	49454 to 51267 (contig 3)
U48_S72	N	-	-	-	-	-	-	-	-	GCA_946224865.1	-
U52_S74	N	-	-	-	-	-	-	-	-	GCA_946190225.1	-
U54_S75	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190185.1	18384 to 28422 (contig 12)
U56_S76	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190325.1	71434 to 73220 (contig 5)
U57_S77	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190235.1	37034 to 38822 (contig 12)
U61_S78	N	-	-	-	-	-	-	-	-	GCA_946190115.1	-
U63_S79	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190175.1	37034 to 38822 (contig 15)
U65_S81	Y	N	-	-	-	N	N	EMRRIDKMFINFLKRR	L	GCA_946190125.1	10058 to 15226 (contig 20)
U67_S83	Y	Y	7327	1	14	Y	Y	EMRRIDKMFINFLKRR	L	GCA_946190165.1	35610 to 42936 (contig 11)
U8_S49	Y	N	-	-	-	Y	Y	ESRLPKIRFDFIFPRKK	P	GCA_946190315.1	52874 to 63087 (contig 6)
U92_S86	Y	N	-	-	-	N	N	ETRKTINSLLNLFKRR	F	GCA_946190145.1	58367 to 59135 (contig 6)
U98_S90	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946190155.1	14745 to 15455 (contig 11)
U99_S91	Y	N	-	-	-	N	N	EIRIADKIFFNLFKRR	F	GCA_946190155.1	14745 to 15455 (contig 12)
WTCHG_774611_72895361	N	-	-	-	-	-	-	-	-	GCA_946190255.1	-
WTCHG_774611_72965368	Y	N	-	-	-	N	N	ESRVSDILLDFLFRRKK	R	GCA_946190195.1	114512 to 120897 (contig 4)
WTCHG_774611_73005372	Y	Y	12164	2	23	N	N	EMRKMNDKSFNLFNFRRR	F	GCA_946190135.1	24172 to 36335 (contig 6)
WTCHG_774611_73015373	N	-	-	-	-	-	-	-	-	GCA_946190275.1	-
WTCHG_774611_73075379	Y	N	-	-	-	Y	Y	EMRKPDGALFNLFRRR	F	GCA_946190295.1	119791 to 129649 (contig 6)
WTCHG_774611_73095381	Y	N	-	-	-	N	N	EMRKMNDKSFNLFNFRRR	F	GCA_946190335.1	118632 to 130795 (contig 6)
WTCHG_774611_73775353	Y	Y	11984	5	25	N	N	EIRLSKFLRDFILQRKK	Q	GCA_946190285.1	28199 to 40182 (contig 5)
WTCHG_774611_73825358	Y	Y	11984	5	25	N	N	EIRLSKFLRDFILQRKK	Q	GCA_946190255.1	28199 to 40182 (contig 5)
WTCHG_774611_73845360	Y	N	-	-	-	N	N	EMRKMNDKSFNLFNFRRR	F	GCA_946185975.1	118632 to 130795 (contig 6)
B22	Y	Y	15566	11	38	N	Y	ESRVSRIILDFLFLRKK	L	GCA_965111745.1	61443 to 77008 (contig 1)
C22	Y	Y	7274	3	15	N	Y	ESRLSRLLRDFIFQIKQ	Q	GCA_965111725.1	50761 to 58034 (contig 1)
D22	Y	Y	2127	2	2	N	N	DSRISKFLRDFIFPRKK	P	GCA_965111735.1	34443 to 36569 (contig 1)
E22	Y	Y	14434	1	32	Y	Y	EMRLPKILRDFIFPRKK	P	GCA_965111705.1	263852 to 278285 (contig 1)
F22	Y	Y	7848	5	13	Y	Y	ESRVSRIILDFLFLRKK	L	GCA_965111715.1	42896 to 50743 (contig 1)
G22	Y	Y	4404	2	12	N	Y	ESRLSRLLRDFIFQIKQ	Q	GCA_965111695.1	57304 to 61707 (contig 1)
B6	Y	Y	9068	5	18	Y	Y	EMRKPDGALFNLFRRR	F	GCA_000027165.1	53854 to 62921 (contig 1)
FDA684	Y	Y	8195	5	17	N	N	EIRQTHNIFFNFFKRR	F	GCA_009730515.1	31493 to 39687 (contig 1)
SK637	Y	Y	7760	4	18	N	Y	EMRKSNNNFFHFLRRI	L	GCA_000722765.2	49608 to 57367 (contig 1)
SVGS061	Y	Y	2200	2	5	N	N	EMRKSNNTFFNFLRRI	L	GCA_001560895.1	73674 to 75873 (contig 1)

The genomic region between *plsX-acpP* and *comAB* was extracted from publicly available *S. mitis* genomes. Of the 332 genomes analyzed, this region was successfully retrieved from 253; the remaining genomes contained contig breaks and were excluded from further analyses. The 253 genomes were screened for the presence of competence-associated bacteriocin (*cab*), competence-induced bacteriocins A and B (*cibAB*), and the cognate immunity protein (*cibC*) genes. The number of SigX boxes in cab loci was determined. Mature CSP sequences of each strain was predicted by identifying small ORFs upstream of *comDE* that contained a double-glycine cleavage motif.

Table S2. Bacterial strains, plasmids, peptides and oligonucleotides used in study.

Bacterial strain, plasmid, oligonucleotide or peptide	Relevant genotype or sequence	Note	Source
Bacterial strains			
Corynebacterium accolens LMG 19042	Wild type	-	BCCM/LMG
Dolosigranulum pigrum LMG 15126	Wild type	-	BCCM/LMG
Escherichia coli DH5α	Wild type	-	Thermo Fisher
Escherichia coli DH10B	Wild type	-	Thermo Fisher
Haemophilus influenzae FP7132	Wild type		P. Lavado
Klebsiella pneumoniae OND482	Wild type	-	Simões *et al.*, 2022
Moraxella catarrhalis ATCC 25238	Wild type	-	ATCC
Neisseria lactamica ATCC 23970	Wild type	-	ATCC
Staphylococcus aureus ATCC 25293	Wild type	-	ATCC
Staphylococcus epidermidis ATCC 14990	Wild type	-	ATCC
Streptococcus agalactiae ATCC 13813	Wild type	-	ATCC
Streptococcus anginosus ATCC 12395	Wild type	-	ATCC
Streptococcus constellatus ATCC 27823	Wild type	-	ATCC
Streptococcus gordonii ATCC 33399	Wild type	-	ATCC
Streptococcus infantis ATCC 700779	Wild type	-	ATCC
Streptococcus mitis SK142	Wild type	-	ATCC
Streptococcus mitis F22	Wild type	-	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis G22	Wild type	-	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis G22 P*comCDE*-pJB11	*CIL*::P*comCDE*-mNeonGreen-luciferase *cat*	chl^R	This study
Streptococcus mitis G22 P*sigX1*-pJB11	*CIL::PsigX1-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 P*sigX2*-pJB11	*CIL::PsigX2-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 P*cabABC*-pJB11	*CIL::PcabABC-mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 P*cabABC* (truncated)-*pJB11*	*CIL::P*cabABC(truncated)-*mNeonGreen-luciferase cat*	chl^R	This study
Streptococcus mitis G22 Δ*comE* P*cabABC*-pJB11	Δ*comE*::*aphA-3 CIL::PcabABC-mNeonGreen-luciferase cat*	kan^R chl^R	This study
Streptococcus mitis G22 Δ*sigX1* P*cabABC*-pJB11	Δ*sigX1*::*aphA-3 CIL::PcabABC-mNeonGreen-luciferase cat*	kan^R chl^R	This study
Streptococcus mitis G22 Δ*sigX2* P*cabABC*-pJB11	Δ*sigX2*::*ant(9) CIL::PcabABC-mNeonGreen-luciferase cat*	spc^R chl^R	This study
Streptococcus mitis G22 Δ*sigX1* Δ*sigX2* P*cabABC*-	Δ*sigX1*::*aphA-3* Δ*sigX2*::*ant(9) CIL*::*PcabABC-mNeonGreen-luciferase cat*	kan^R spc^R chl^R	This study
Streptococcus mitis Δ*blp4* Δ*blp1*	Δ*blp4*::*lox72* Δ*blp1*::*aphA-3*	kan^R	Borralho, Handem and Lança *et al.,* 2025
Streptococcus mitis Δ*blp4* Δ*blp1* (markerless)	Δ*blp4*::*lox72* Δ*blp1*::*lox72*	-	This study
Streptococcus mitis Δ*blp4* Δ*blp1* Δ*cabABC*	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*cabABC*::*aphA-3*	kan^R	This study
Streptococcus mitis p(*cab$^+$*)	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*cabABC*::*aphA-3* pJB09-P23-*cabABC*	spc^R kan^R	This study
Streptococcus mitis G22 *cabA*-HA	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*cabABC*::*aphA-3 CIL*::*cabA-HA cabBC cat*	kan^R chl^R	This study
Streptococcus mitis G22 *cabB*-HA	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*cabABC*::*aphA-3 CIL*::*cabA cabB-HA cabC cat*	kan^R chl^R	This study
Streptococcus mitis Δ*blp4* Δ*blp1* Δ*comE*	Δ*blp4*::*lox72* Δ*blp1*::*lox72* Δ*comE*::*aphA-3*	kan^R	This study
Streptococcus mutans ATCC 25175	Wild type	-	ATCC
Streptococcus oralis DSM 20066	Wild type	-	DSMZ

(continues on next page)

Table S2. (cont.)

Streptococcus parasanguinis ATCC 15912	Wild type	-	ATCC
Streptococcus peroris ATCC 700780	Wild type	-	ATCC
Streptococcus pneumoniae D39	Wild type	-	Avery *et al.,* 1944
Streptococcus pneumoniae D39 PhunSweet	Δ*cps* ::*pheS-sacB-aphA-3*	kan^R	Echlin and Rosch, 2020
Streptococcus pneumoniae D39 P*comCDE*-pJB11	*CIL* ::P*comCDE* -mNeonGreen-luciferase *cat*	chl^R	This study
Streptococcus pneumoniae D39 Δ*lytA*	Δ*lytA* ::*pheS-sacB-aphA-3*	kan^R	This study
Streptococcus pneumoniae D39 Δ*lytA* (markerless)	Δ*lytA*	-	This study
Streptococcuspneumoniae D39 Δ*lytC*	Δ*lytC* ::*pheS-sacB-aphA-3*	kan^R	This study
Streptococcuspneumoniae D39 Δ*lytC* (markerless)	Δ*lytC*	-	This study
Streptococcus pneumoniae D39-GFP	P*hlpA* -*hlpA-gfp cat*	chl^R	Valente *et* al., 2022
Streptococcus pneumoniae D39-GFP ΔcomGC	P*hlpA* -*hlpA-gfp cat* Δ*comGC* ::*tet(M)*	tet^R	This study
Streptococcus pneumoniae TIGR4	Wild type	-	ATCC
Streptococcus pseudopneumoniae BAA-960	Wild type	-	ATCC
Streptococcus pyogenes ATCC 12344	Wild type	-	ATCC
Streptococcus salivarius ATCC 7073	Wild type	-	ATCC
Streptococcus sanguinis ATCC 10556	Wild type	-	ATCC
Streptococcus sinensis HKU4	Wild type	-	DSMZ
Streptococcus sobrinus ATCC 27351	Wild type	-	ATCC
Streptococcus vestibularis ATCC MM1	Wild type	-	ATCC

Plasmids

pDL278-P23-DsRed-Express2	*P23-dsRed-Express2 pMB1 pVA380-1 ant(9)*	*E. coli*-streptococcal	Plasmid #121505 (Addgene)
pJB09-cabABC	pDL278-P23-DsRed-Express2 but Δ*dsRed-Express2* ::*cabABC*	*E. coli*-streptococcal	This study
pBIR	*p15A aphA-3 tdTomato cat*	Integrates in *CIL* locus	This study (Twist Bioscience)
pPEP-LGZ	*luc sfGFP lacZ cat p15A ant(9)*	Integrates in *CEP* locus	Kjos *et al* ., 2016
pJB10	pBIR but Δ*tdTomato* ::*hlpA-mNeonGreen*	Integrates in *CIL* locus	This study
pJB11	pJB10 but *luc*	Integrates in *CIL* locus	This study
pJB11-PcomCDE(G22)	pJB11 but P*comCDE* reporter (*S. mitis* G22)	Integrates in *CIL* locus	This study
pJB11-PcomCDE(D39)	pJB11 but P*comCDE* reporter (S. *pneumoniae* D39)	Integrates in *CIL* locus	This study
pJB11-PsigX1	pJB11 but P*sigX1* reporter	Integrates in *CIL* locus	This study
pJB11-PsigX2	pJB11 but P*sigX2* reporter	Integrates in *CIL* locus	This study
pJB11-PcabABC	pJB11 but P*cabABC* reporter	Integrates in *CIL* locus	This study
pJB11-PcabABC(truncated)	pJB11 but P*cabABC* (truncated) reporter	Integrates in *CIL* locus	This study
pKan	*pMB1 lox66-p3-aphA-3-lox71*	Integrates in *cab* locus	Borralho, Handem and Lança *et al.* , 2025

Peptides

BlpC$_1$	GLWEELYETITTPLKLRTTFKP	-	NZYTech
BlpC$_4$	GWWEELLHETILSKFKITKALELPVQL	-	NZYTech
CSP-1	EMRLSKFFRDFILQRKK	-	NZYTech
CSP-2	EMRISRIILDFLFLRKK	-	NZYTech
CSP-2(L14Q)	EMRISRIILDFLFQRKK	-	NZYTech

Table S2. (cont.)

CSP$_{A22}$	DWRISETIRNLIFPRRK	-	NZYTech
CSP$_{B22}$	ESRVSRIILDFLFLRKK	-	NZYTech
CSP$_{D22}$	DSRISKFLRDFIFPRKK	-	NZYTech
CSP$_{E22}$	EMRLPKILRDFIFPRKK	-	NZYTech
CSP$_{G22}$	ESRLSRLLRDFIFQIKQ	-	NZYTech
CSP$_{G22}$(Q14L)	ESRLSRLLRDFIFLIKQ	-	NZYTech
CSP$_{G22}$(Q14N)	ESRLSRLLRDFIFNIKQ	-	NZYTech
Oligonucleotides			
P1F_cab7_OE	ATGACAAATTTTGACAAAATGGAAC	Construction of plasmid pJB09-cabABC	This study
P1R_cab7_OE	CCCTATAGAACGTCTACTACCTC		
P1F_pDL278_OE	CTCGGTACCAAATTCCAGAAAAG		
P1F_pDL278_OE	GTTTGTCCTCCTTAACTAGTAATTAATAC		
P1F_hlpA_mNG_twist	AAAGCACTCAAAAGTTTACCTTATGG	Construction of plasmid pJB10	This study
P1R_hlpA_mNG_twist	CGCAAAAAACCCCGCTTCGGCGGGGTTTTTTCGCTTATTTATAC		
P1F_pBIR_backbone	ATCGGAGAAATTGGACACACC		
P1R_pBIR_backbone	CCCTATCAAAAGAATCTTGCTTGGC		
PF_luc_pPEP	GCTGAAGGAGGAATAATGAGATCC	Construction of plasmid pJB11	This study
PR_luc_pPEP	TTACAATTTGGGCTTTCCGCC		
PF_pJB10_backbone	GAATGGACGAGTTGTATAAG		
PR_pJB10_backbone	TTATTTATACAATTCATCCATACCC		
PF_reporter_PcomCDE(G22)	AGCTGGTCTCTGACTATATAGTAGCAAAGCTGGG	Construction of plasmid pJB11-PcomCDE (G22)	This study
PR_reporter_PcomCDE(G22)	CGATGGTCTCAAGTCAATAAAATCTCCTAAAATGTTTTTTTC		
PF_reporter_PcomCDE(D39)	AGCTGGTCTCTGACTATATAATAGCAAAGCTGGG	Construction of plasmid pJB11-PcomCDE (D39)	This study
PR_reporter_PcomCDE(D39)	CGATGGTCTCAAGTCAATAAAATCTCCTAAAATGTTTTTTTC		
PF_reporter_PsigX1	AGCTGGTCTCTGACTCCCAAAGAGAGGCAAGACC	Construction of plasmid pJB11-PsigX1	This study
PR_reporter_PsigX1	CGATGGTCTCAAGTCTCTTTTCCCCTTTCTGTGTAC		
PF_reporter_PsigX2	AGCTGGTCTCTGACTCCCTGAGAGAGGCCCGTC	Construction of plasmid pJB11-PsigX2	This study
PR_reporter_PsigX2	CGATGGTCTCAAGTCAGTCTTCCCCTTTCTGTGTAC		
PF_reporter_PcabABC	AGCTGGTCTCTGACTAAATAACATCTTCACAACTAATCG	Construction of plasmid pJB11-PcabABC	This study
PR_reporter_PcabABC	CGATGGTCTCAAGTCCTTTTTTCTCCTTTATTTCTATTTTTTAG		
PF_reporter_PcabABC_trunc	AGCTGGTCTCTGACTTTTAGTTGTGTATTTTCGAATTG	Construction of plasmid pJB11-PcabABC (truncated)	This study
PR_reporter_PcabABC_trunc	CGATGGTCTCAAGTCCTTTTTTCTCCTTTATTTCTATTTTTTAG		
P1F_comE_G22	GGATTAAACAGTATGAGAGG	Deletion of *comE* in *S. mitis* G22	This study
P1F_comE_G22	GTCAATACCGTTCGTATAGCATACATTATACGAAGTTATTCATTCAAATTCCCTCTTAAATC		
P2F_comE_G22_pKan	CAAATTATTAGATTTAAGAGGGAATTTGAATGAATAACTTCGTATAATGTATGC		
P2R_comE_G22_pKan	CATTATATCAATTTCTTGCTAATTGTCAATCACTATAACTTCGTATAGCATAC		
P3F_comE_G22	GTTTTAGTACCGTTCGTATAATGTATGCTATACGAAGTTATAGTGATTGACAATTAGCAAGAAATTG		
P3R_comE_G22	TATTACCATTATAGGAAGCC		
P4F_comE_G22	TATAACAGGAGATAATTTTG		
P4R_comE_G22	ACAGAGCTGGAATAAGAGAAATCC		

(continues on next page)

Table S2. (cont.)

P1F_sigX1_G22	ATTGTCATCGCTGCGACAAAC		
P1R_sigX1_G22	GTCAATACCGTTCGTATAGCATACATTATACGAAGTTATCATTCTTTTCCCCTTTCTGTGTACAG		
P2F_sigX1_G22_pKan	GACTGTACACAGAAAGGGGAAAAGAATGATAACTTCGTATAATGTATGC		
P2R_sigX1_G22_pKan	CTTTTTTGCATGAATTCCTAATGGGTACGATAACTTCGTATAGCATAC	Deletion of *sigX1* in *S. mitis* G22	This study
P3F_sigX1_G22	GTTTTAGTACCGTTCGTATAATGTATGCTATACGAAGTTATCGTACCCATTAGGAATTCATGC		
P3R_sigX1_G22	CGCTACACATGGAATTCCAC		
P4F_sigX1_G22	AGCAATCTTGAAAGTTCATGCTAAG		
P4R_sigX1_G22	ATCCGGACAACGCTCGGGACCTAC		
P1F_sigX2_G22	TGCTTGTCAAAGCCTGGCTTTCTTG		
P1F_sigX2_G22	CATGCAAAACAATGAATAGGCATAGTCTTCCCCTTTCTGTGTAC		
P2F_sigX2_G22_pPEP	GACTGTACACAGAAAGGGGAAGACTATGCATGCAAAACAATGAATAGG		
P2R_sigX2_G22_pPEP	CTTTTTTGCATGAATTCCTAATGAGTACGCTAATTGAGAGAAGTTTCTATAG	Deletion of *sigX2* in *S. mitis* G22	This study
P3F_sigX2_G22	CTAATTGAGAGAAGTTTCTATAGCGTACTCATTAGGAATTCATGCAAA		
P3R_sigX2_G22	CCGCTACACATGGAATTCCACTC		
P4F_sigX2_G22	CTATTCAATCAAGATTAGTGGAAAC		
P4R_sigX2_G22	TAGTTAGCCGTCCCTTTCTGGTAAG		
P1F_PS_lytA_D39	GGTTTATCAGCGGTTCTCTGC		
P1R_PS_lytA_D39	CGGATCGATCCTTAATAGTGGGTTCCATATTCTACTCCTTATCAA		
P2F_PS_lytA_D39	GTTTTAATTGATAAGGAGTAGAATATGGAACCCACTATTAAGGATCGATC		
P2R_PS_lytA_D39	CTGTTCTGATTTGAAAGACATTCCATTATTACTAAAACAATTCATCCAGTAAAATATAAT		
P3F_PS_lytA_D39	CTGGATGAATTGTTTTATAATAATGGAATGTCTTTCAAATCAG	Deletion of *lytA* in *S. pneumoniae* D39	This study
P3R_PS_lytA_D39	TTTTACTTTCTGACAAAGGACG		
P4F_PS_lytA_D39	TTCAATTGTATCTATCGGCAG		
P4R_PS_lytA_D39	TTCCTCAATCTATATAACATAGCTTTATG		
P5F_PS_lytA_D39	GTTTTAATTGATAAGGAGTAGAATATGGAATAATAATGGAATGTCTTTCAAATCAG		
P5R_PS_lytA_D39	CTGTTCTGATTTGAAAGACATTCCATTATTATTCCATATTCTACTCCTTATCAA		
P1F_PS_lytC_D39	ACTAAAGGAAGATGGAACCAG		
P1R_PS_lytC_D39	CGGATCGATCCTTAATAGTGGGAGACATCTGAATGCTGTTCC		
P2F_PS_lytC_D39	CTAGGTGGAACAGCATTCAGATGTCTCCCACTATTAAGGATCGATC		
P2R_PS_lytC_D39	CACATCCCTCTTTCAAATCATCGCTTACTAAAACAATTCATCCAGTAAAATATAAT		
P3F_PS_lytC_D39	CTGGATGAATTGTTTTATAAGCGATGATTTGAAAGAGGG	Deletion of *lytC* in *S. pneumoniae* D39	This study
P3R_PS_lytC_D39	TTCTTAGTCATGCTTCCATCC		
P4F_PS_lytC_D39	CTCAAATTGAGGCCAAGAGAG		
P4R_PS_lytC_D39	GTCCAAGTTCGGCTGCAAG		
P5F_PS_lytC_D39	CTAGGTGGAACAGCATTCAGATGTCTTAAGCGATGATTTGAAAGAGGG		
P5R_PS_lytC_D39	CACATCCCTCTTTCAAATCATCGCTTAAGACATCTGAATGCTGTTCC		

(continues on next page)

Table S2. (cont.)

P1F_comGC_del	AATCGTTGACTTTGCAAGCAG	Deletion of *comGC* in *S. pneumoniae* D39	This study
P1R_comGC_del	TAATTTTCCTCCTATTTAGATCTTGCATGTATAGTTATTATACCAGGGGGACAGTGCAATGTCAATTTAAAAATTTACCTCCATATTTTGATACATG		
P2F_comGC_del	TTGACATTGCACTGTCCCCCTGGTATAATAACTATACATGCAAGATCTAAATAGGAGGAAAATTAATGAAAATTATTAATATTGGAGTTTTAGC		
P2R_comGC_del	CATTGACTTTACGATTTGCTCCTCCATTCTAAGTTATTTTATTGAACATATATCG		
P3F_comGC_del	GTACGATATATGTTCAATAAAATAACTTAGAATGGAGGAGCAAATCGTAAAGTC		
P3R_comGC_del	GTGACTGCGTAGAGGAGAAC		
P4F_comGC_del	TTCAGGCTGAGACGGAAAAAATTAG		
P4R_comGC_del	CAGATTTGAGTCCATAAACC		
P1F_cabAB_HA	CAAGCGCCAGGGTTTTCC	Construction of *S. mitis* harboring HA-tagged bacteriocins	This study
P1R_cabAB_HA	GTTCTGTTCCATTTTGTCAAAATTTGTCATCTTTTTTCTCCTTTATTTCTATTTTTTAG		
P2F_cabAB_HA	GTAAACTAAAAAATAGAAATAAAGGAGAAAAAAGATGACAAATTTTGACAAAATGGAAC		
P2F_cabAB_HA	CCAGGGGGACAGTGCAATGTCAATTATCCATTCTTTTTTTCTTTACTTC		
P3F_cabAB_HA	CATTCTTTTATATAGAAGTAAAGAAAAAAAGAATGGATAATTGACATTGCACTGTCCCC		
P3F_cabAB_HA	GCCCTCTGAGCGGATAACAATTTC		
P4F_cabAB_HA	ATCGGAGAAATTGGACACACC		
P4R_cabAB_HA	TTCGCAAATAGCGAAACCAATG		

chl[R] - chloramphenicol resistant; kan[R] - kanamycin resistant; tet[R] - tetracycline resistant; spc[R] - spectinomycin resistant.

SI References

1.   A. S. Simões *et al*., Using whole genome sequencing to investigate a mock-outbreak of carbapenem-resistant *Klebsiella pneumoniae* in real-time. *Acta Med Port* **35**, 36–41 (2022).

2.   J. Borralho *et al*., Inhibition of pneumococcal growth and biofilm formation by human isolates of *Streptococcus mitis* and *Streptococcus oralis. Appl Environ Microbiol* **91**, e01336-24 (2025).

3.   O. T. Avery, C. M. MacLeod, M. McCarty, Studies on the chemical nature of the substance inducing transformation of pneumococcal types: induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III. *J Exp Med* **79**, 137–158 (1944).

4.   H. Echlin, J. W. Rosch, Advancing genetic tools in *Streptococcus pneumoniae. Genes (Basel)* **11**, 965 (2020).

5.   C. Valente, A. R. Cruz, A. O. Henriques, R. Sá-Leão, Intra-species interactions in *Streptococcus pneumoniae* biofilms. *Front Cell Infect Microbiol* **11**, 803286 (2022).

6.   M. Kjos *et al*., Expression of *Streptococcus pneumoniae* bacteriocins is induced by antibiotics via regulatory interplay with the competence system. *PLoS Pathog* **12**, e1005422 (2016).