Diet and temperature interactively impact brown adipose tissue gene regulation controlled by DNA methylation

Tobias Hagemann[1,*], Anne Hoffmann[1,*], Kerstin Rohde-Zimmermann[1,*], Helen Broghammer[1], Lucas Massier[1,2], Peter Kovacs[3], Michael Stumvoll[1,3], Matthias Blüher[1,3], John T. Heiker[1,3,*, §] & Juliane Weiner[1,3,*, §]

[1]Helmholtz Institute for Metabolic, Obesity and Vascular Research (HI-MAG) of the Helmholtz Zentrum München at the University of Leipzig and University Hospital Leipzig, Leipzig, Germany
[2]Department of Medicine H7, Karolinska Institute, Stockholm, Sweden
[3]Leipzig University Medical Center, Divisions of Endocrinology and Nephrology, University of Leipzig, Leipzig, Germany

*These authors contribute equally

§**Corresponding authors:**
Juliane Weiner
[2]Leipzig University Medical Center, Divisions of Endocrinology and Nephrology, University of Leipzig, Leipzig, Germany,
Liebigstr. 21, D-04103 Leipzig, Germany
Email: Juliane.Weiner@medizin.uni-leipzig.de
Phone: +49-341-97 20919

John T. Heiker
Helmholtz Institute for Metabolic, Obesity and Vascular Research (HI-MAG)
Helmholtz Zentrum München at the University of Leipzig and University Hospital Leipzig
Philipp-Rosenthal-Straße 27, D-04103 Leipzig, Germany
Email: John.Heiker@helmholtz-munich.de
Phone: +49-341-97 22909

Keywords: brown adipose tissue, DNA methylation, high fat diet, cold exposure, epigenetics

Abbreviations: Aza: 5'aza-2'-deoxycytidine (Aza), BAT: Brown adipose tissue, BH: Bonferroni-Holm, DEG: differentially expressed gene, DGE: differential gene expression, DMEG: Differentially methylated and expressed gene, DMP: differentially methylated positions, DMR: differentially methylated regions, eWAT: epididymal white adipose tissue, FC: fold change, FDR: false discovery rate, HFD: high-fat diet, imBA: immortalized brown adipocytes, iWAT: white adipose tissue, PCA: principal component analysis, qPCR: quantitative real-time-PCR, RNAseq: RNA sequencing, RRBS: reduced representation bisulfite sequencing, SAM: S-adenosylmethionine, SVF: stromal vascular fraction, VST: variance stabilization transformation

Abstract

Controlling brown adipose tissue (BAT) plasticity in adulthood holds promising potential for effective new obesity therapies by targeting the mechanisms of adaptive thermogenesis. Recent studies have shown that BAT development and function are under epigenetic control, with DNA methylation linked to the regulation of key thermogenic and metabolic genes. Here we sought to understand how diet and cold exposure interactively shape BAT gene regulation controlled by DNA methylation. Mice (N = 8 per group) were housed under cold exposure (8°C) or thermoneutrality (30°C) and fed either chow or high-fat diet (HFD). BAT was isolated for transcriptome (RNAseq) and methylome (RRBS) analyses. We identified differentially methylated and expressed genes (DMEGs) by comparing the effects of cold exposure under chow and high-fat diet, as well as by analyzing the interaction between temperature and diet. Functional pathway enrichment and EpiFactors Database screening were used to assess epigenetic regulators, and candidate gene expression was validated by modulating DNA-methylation in vitro.

We identified ~1,360 differentially expressed genes (DEGs) uniquely affected by the diet-temperature interaction with most downregulated in HFD-fed mice, indicating that obesity limits the transcriptional response of BAT to cold. 65 DMEGs (4% of DEGs) were largely diet-specific in response to cold exposure, suggesting that DNA methylation contributes to a selective layer of gene regulation during BAT adaptation to distinct metabolic states. In HFD-fed mice, DMEGs were enriched in pathways related to mitochondrial dysfunction, altered lipid metabolism, neuroendocrine signaling, and compensatory stress responses, contrasting the adaptive thermogenic profile observed in chow-fed controls. Differentially expressed genes of epigenetic regulators such as *Tet2*, *Dnmt3a* and *Apobec1* showed diet- and cold-dependent regulation, indicating impaired methylation flexibility under obesogenic conditions. Using cell culture experiments, we confirmed the regulation of gene expression of candidate genes, validating the functional link between DNA methylation and thermogenic gene regulation. This is the first study to demonstrate an epigenetic response to cold exposure that differs by dietary condition, particularly in obesity. Our findings highlight a coordinated transcriptional and epigenetic remodeling of BAT, shaped by both environmental and metabolic signals. These insights may inform targeted epigenetic or nutritional strategies to restore BAT function and would further strengthen our understanding of how these might be used as therapeutic basis to improve metabolic health in obesity.

Graphical abstract

Mice were housed under cold exposure or thermoneutrality and fed either chow or high-fat diet (HFD). RNA sequencing (RNAseq) and reduced representation bisulfite sequencing (RRBS) was performed on brown adipose tissue (BAT) which provide the fundament for the identification of differentially methylated positions (DMP) and regions (DMR) as well as differentially expressed genes (DEG) in three models: $COLD_{chow}$, comparing 8° vs 30°C mice on chow diet; $COLD_{HFD}$, comparing 8° vs 30°C mice on HFD diet and $\Delta COLD_{diet}$, comparing $COLD_{chow}$ vs $COLD_{HFD}$. Differentially methylated and expressed genes (DMEGs; circos plot shows a representative example) were identified for all comparisons based on those DMPs which correlated significantly with differentially expressed genes (DEGs; the heatmap shows a representative example). DMEGs were taken forward for pathways enrichment analyses and to identify differentially regulated epigenetic regulators using the EpiFactors Database. The figure was created with BioRender.

1. Introduction

Brown adipose tissue (BAT) is a specialized fat tissue that is rich in mitochondria and promotes non-shivering thermogenesis by expressing the *uncoupling protein 1 (UCP1).* Unlike white adipose tissue, it burns calories to produce heat and therefore plays a key role in energy regulation and controlling metabolic health. Increasing energy expenditure by BAT activation is an intriguing therapeutic approach to combat the overwhelming obesity pandemic, either alone or to complement the current pharmacotherapy that mainly addresses energy intake based on the incretin-mimetic poly-agonist class of drugs [1]. With this in mind it is not surprising that a lot of research was conducted to understand the molecular underpinnings of BAT regulation specifically addressing environmental cues. Cold exposure is the most powerful inducer of BAT activation leading to the upregulation of thermogenic gene program and adrenergic receptor-mediated activation of lipolysis and metabolism. BAT activation also occurs post-prandially, especially after acute overfeeding, to trigger diet-induced thermogenesis [2,3]. However, this compensatory component of energy-expenditure is impaired during chronic overfeeding, a phenomenon that was termed adaptive thermogenesis, and is believed to further drive weight gain and obesity [2,4,5].

In any case the mechanistical switch enabling the high plasticity of BAT changing from passive thermoneutral to active thermogenic states may be regulated by epigenetic mechanisms that are affected by, e.g. temperature and diet [6]. Indeed, studies have shown changes in DNA methylation for key BAT regulatory genes such as *Ucp1, PR domain containing 16* (*Prdm16*) or *PPARG coactivator 1 alpha* (*Ppargc1a*) , altering their expression level and impacting BAT function [7,8]. Moreover, during the development of precursor cells into major brown adipocytes, DNA methylation patterns change and lineage commitment towards a thermogenic adipocyte is epigenetically controlled [8]. The major relevance of the ambient temperature in this context has just recently been implied, as cold exposure of parents at conception is related to increased brown adipogenesis and thermogenesis along with reduced obesity risk in the offspring [9]. Mechanistically, sperm DNA methylation is altered in response to cold exposure which leads to adaptations in BAT activity of the offspring [10,11]. In addition, DNA methylation in key genes of thermogenesis and fatty acid oxidation, such as *acetyl-CoA acyltransferase 2* (*Acaa2*) and *acyl-CoA synthetase long chain family member 1* (*Acsl1*), is increased in BAT of offsprings from mice fed a high fat diet [12]. These studies strongly imply the relevance of cold, diet and obesity on the transgenerational modulation of BAT function. However, these environmental cues, either alone or in combination, also trigger and control BAT plasticity during adulthood. Rats with a blockade in afferent vagal nerve signaling that triggered by high-fat diet showed a reduced BAT activation after cold exposure [13]. The relevance of diet for regulating DNA methylation in later life comes with nutritional compounds such as folate or methionine as critical components of the one-carbon metabolism, which are classically enriched in vegetable-rich diets but diminished in high fat diets.

However, the interaction of especially cold exposure and high fat diet on BAT's epigenetic machinery is less understood. In this study, we aimed to address DNA methylation alterations in response to temperature and diet, and link this to changes in gene expression of BAT in mice on a

genome-wide level. We explored the interactive impact of diet and cold exposure on the epigenetic
regulation of BAT gene expression. Using a stringent bioinformatic workflow we focused on
differentially methylated and expressed genes (DMEGs) by taking forward only candidates whose
differential methylation signal correlates to the gene expression level in order to provide only top
functional relevant candidates. Providing a profound overview over the impact of DNA
methylation on genes expression affected by cold and diet is of strong relevance to understand adult
BAT plasticity and adaptive thermogenesis and provides the fundament for new obesity treatment
strategies.

2. Material and methods

2.1. Animal studies

Male C57BL/6NTac mice (Taconic Bioscience, Lille Skensved, Denmark) were housed in
pathogen-free facilities at 23°C on a 12h light/dark cycle at the Sächsische Inkubator für Klinische
Translation (SIKT), Leipzig. Starting from five weeks of age, mice were either fed a standard chow
diet (EV153, 3.3% from fat, Ssniff, Soest, Germany) or a high fat diet (HFD, E15742, 60 kJ% from
fat, Ssniff) with ad libitum access to food and water. At 11 (chow) or 22 (HFD) weeks of age, mice
were adapted to single housing in rodent climate chambers (MKKL1200, Flohr Instruments,
Netherlands) for 5 days, before randomization into four groups (each N = 8 mice) and housed either
at thermoneutrality (30°C) or in the cold (8°C) for 7 days. After a 4-hour fasting period, body
weight and rectal temperature were measured. BAT and tail surface temperatures were measured
by thermal imaging (VarioCAM®, InfraTec, Dresden, Germany). Mice were sacrificed and tissue
samples of intrascapular brown adipose tissue (BAT) were collected, snap frozen in liquid nitrogen
and stored at -80°C before further analyses. All animal experiments were approved by the local
authorities of the Free State of Saxony, Germany (Landesdirektion Leipzig: TVV51/20), as
recommended by the responsible local animal ethics review board.

2.2. RNA sequencing of brown adipose tissue

RNA from BAT was isolated using RNeasy Lipid Tissue Mini kit (Qiagen, Hilden, Germany) as
specified by the manufacturer. RNA sequencing (RNAseq) was performed by the core unit DNA-
Technologien, Medical Faculty Leipzig. 50ng of total RNA were depleted of ribosomal RNA using
the NEBNext rRNA Depletion Kit v2 (NEB, USA) according to the manufacturer's instructions.
Depleted RNA was transcribed using SuperScript IV reverse transcriptase (ThermoFisher
Scientific, USA) for 2h at 55°C. After second strand synthesis (TargetAmp kit, EPICENTRE,
USA), the DNA was fragmented using the Illumina Tagment DNA TDE1 Enzyme and Buffer Kits,
which fragments DNA and inserts partial sequencing adapter sequences. Final PCR amplification
of the libraries was done using KAPA HiFi HotStart Library Amplification Kit with unique dual
indexing by IDT for Illumina Nextera DNA Unique Dual Indexes Sets. The barcoded libraries
were purified and quantified using Qubit Fluorometric Quantification (ThermoFisher Scientific,
USA). Size distribution of the library DNA was analyzed using the FragmentAnalyzer (Agilent,

USA). Sequencing of 2x150 bp was performed with NovaSeq sequencer (Illumina, USA)
according to the manufacturer's instructions, with a read depth of 20 million reads per sample.

2.3. Reduced representation bisulfite sequencing in brown adipose tissue

Genomic DNA from BAT samples were isolated using the DNeasy blood and tissue kit according
to the manufacturer's instructions (Qiagen, Hilden, Germany). Reduced representation bisulfite
sequencing (RRBS) library preparation was performed on 100 ng DNA using the Zymo-Seq RRBS
Library Kit (Zymo Research, Freiburg, Germany) by the core unit DNA-Technologien, Medical
Faculty Leipzig. Briefly, DNA was digested with 20U/µl MspI for 4h followed by RRBS adapter
ligation using 400U/µl T4 DNA ligase and gap filling with 2U/µl Taq DNA polymerase and 10mM
5-methylcytosine dNTP Mix. Bisulfite conversion was performed with lightning conversion
reagent, and the final samples were amplified using 5µM of the unique dual index Primer Set
(indexes 1-12) supplied by the manufacturer. Quality control was performed on the Bioanalyzer
(Illumina, USA) prior paired-end sequencing on the NovaSeq6000 platform (Illumina, USA) were
20 million reads per sample were obtained.

2.4. RNAseq data preprocessing

Raw RNAseq reads where quality-filtered and adapter-trimmed using Trimmomatic (v0.39) [14],
discarding reads shorter than 36 bp. Afterwards, reads were aligned to GRCm38.p6 reference
assembly using STAR (v2.7.8a) [15], allowing for a maximum of 50 multi-mapped reads. Gene
counts were quantified with featureCounts from the subread software package (v2.0.1) [16] with
multimappings counted by fractions.

2.5. Differential gene expression analysis

Transcripts without gene annotation and genes with a total count sum <10 were removed resulting
in 18k genes for subsequent gene expression analysis. Variance stabilization transformation (VST)
was applied to the filtered gene count matrix using the R package DESeq2 (v1.42.1) [17]. A
principal component analysis (PCA) was conducted to verify clustering of diet (HFD, chow) and
temperature (8°C, 30°C) groups within the gene expression signals. Three samples were identified
as outliers which did not cluster in their respective group according to the first two principal
components and were subsequently removed from the analysis. This resulted in group sizes of 7-8
mice reflecting all diet-temperature combinations ($N_{total} = 29$; $N_{HFD\_30°C} = 7$; $N_{HFD\_8°C} = 7$; $N_{chow\_30°C}$
$= 7$; $N_{chow\_8°C} = 8$). Afterwards, differential gene expression (DGE) analysis was performed with
DESeq2. Three comparisons were conducted: chow-fed mice at 8°C versus those at 30°C
($COLD_{chow}$), HFD-fed mice at 8°C versus those at 30°C ($COLD_{HFD}$), and an interaction model
comparing $COLD_{HFD}$ versus $COLD_{chow}$ contrasts ($\Delta COLD_{diet}$). DEGs with a false discovery rate
(FDR) < 0.05 were considered statistically significant. DEG lists from each contrast where
subsequently matched with entries in the EpiFactors Database (v2.0,
https://epifactors.autosome.org, accessed on 10.09.2024) to investigate diet-specific methylation
regulators.

2.6. RBBS data processing

Raw RRBS reads from N = 29 mice ($N_{HFD\_30°C}$ = 7; $N_{HFD\_8°C}$ = 7; $N_{chow\_30°C}$ = 7; $N_{chow\_8°C}$ = 8) were
processed with the nextflow (v 23.04.3) [18] pipeline nf-core/methylseq (v2.4.0) [19] using the
parameters *--genome GRCm38 --non_directional --rrbs --skip_deduplication* and *--*
cytosine_report. In brief, quality filtering and adapter trimming of raw RRBS fastq files was
conducted with trimgalore (v0.6.7) [20] before running sequence alignments with Bismark
(v0.24.0) [21]. Afterwards, methylation signals are called with the Bismark methylation extractor
script including the build-in *coverage2cytosine* module for a genome-wide cytosine methylation
report which served as input for downstream analysis. Sex chromosomes were excluded.

2.7. Differential methylation analysis

To identify differentially methylated positions (DMPs) and differentially methylated regions
(DMRs), differential methylation analysis was conducted for *COLD_{chow}*, *COLD_{HFD}*, and
ΔCOLD_{diet}, utilizing beta values and the R packages limma (v3.54.0) [22] and RnBeads (v2.16.0)
[23] with parameters filtering.low.coverage.masking = TRUE, filtering.high.coverage.outliers =
TRUE and filtering.snp = "any". Based on their chromosomal position and respective annotation,
individual methylations can be contextualized within gene bodies or promoter regions (1500 bases
upstream and 500 bases downstream of the transcription start site), both of which were considered
in our analysis.

2.8. Integration of methylation and transcriptome data

After differential analysis of individual omics data, both gene expression and methylation signals
were combined to find biological associative effects between transcriptomics and epigenomics.
The R package biomaRt (v2.58.2) [24] was used to gather gene loci information for differentially
expressed genes (DEGs) and DMRs in gene and promotor context. Beta methylation values were
transformed to M-values using RnBeads. For each DEG, the corresponding DMR was selected
based on the alignment of gene IDs. M-values of all significantly DMPs overlapping with the DMR
were considered using the bedR (v1.0.7) [25] and the GenomicRanges (v1.54.1) [26] R packages.
Pearson correlation was then applied to find associations between DEG expression values and M-
values of each intersecting DMP using the multtest.cor function from RVAideMemoire (v0.9-83-
7) [27]. All DMPs with significant absolute Pearson correlations >0.5 between expression and
methylation signal (p <0.05 without adjustment for multiple testing) were defined as differentially
methylated and expressed genes (DMEGs). If a DMEG is associated to both promoter and gene
body region, it was assigned to the DMR with the highest significance. This step was parallelized
with the foreach package (v1.5.2) [28]. The final list of DMEGs is comprised of genes with
significantly regulated expression signal (absolute $\log_2$ fold change (FC) >0.5) and methylation
signal (absolute $\log_2$ FC >1). Since several DMRs or DMPs can be differentially methylated within
a gene, several DMEGs were generated for such genes.

Subsequently, functional analysis of DMEGs was done with the use of QIAGEN Ingenuity
Pathway Analysis software (QIAGEN Inc., https://digitalinsights.qiagen.com/IPA, Winter Release
Q4 2024) [29] covering the catalog for disease and bio functions and canonical pathways. The
database query was limited within the IPA software to relationships with adipose tissue or
adipocytes in mice that are experimentally observed. Enrichments are reported with p after
Bonferroni-Holm (BH) adjustment and activation z-scores. Enrichments with BH corrected p <0.05
are considered significant.

2.9. In vitro validation experiments

2.9.1. Immortalized and primary brown adipocytes

Immortalized (imBA [30]) and primary brown adipocytes were cultured and differentiated as
previously described [31]. At day 10, cells were harvested and subjected to further analyses.

2.9.2. In vitro manipulation of cellular DNA methylation levels

To analyze the effect of DNA methylation status on gene expression in imBA and primary brown
adipocytes, pre-adipocytes were treated with indicated concentrations of 5'aza-2'-deoxycytidine
(Aza) for 48h and subsequently differentiated [31]. Differentiated adipocytes at day 6 were treated
with indicated concentrations of S-adenosylmethionine (SAM, New England Biolabs) for 48h.
Differentiated adipocytes were harvested and subjected to further analysis.

2.9.3. Quantitative real-time-PCR (qPCR)

RNA from imBA and human primary SAT cells was isolated using RNeasy Lipid Tissue Mini kit.
qPCR was performed using the LightCycler System LC480 and LightCycler-DNA Master SYBR
Green I Kit (Roche, Mannheim, Germany). Adipocyte gene expression of candidate genes was
calculated by ΔΔCT method and normalized to *Nono* or *36b4* levels as indicated. Primer sequences
are listed in (Suppl. Table 1).

2.10. Statistical analysis

Analyses were performed under R version 4.3.2, except for mouse phenotyping data that was
analyzed using GraphPad Prism 10 (GraphPad, San Diego, CA, USA). Statistical tests are stated
in the figure legends. If not stated otherwise, adj. p <0.05 were considered statistically significant.

3. Results

3.1. Phenotyping of chow and high fat diet-fed mice held at thermoneutrality or exposed to cold

Groups of 8 mice per diet were held at thermoneutrality (30°C) or in the cold (8°C) for 7 days (Figure 1A). Before the temperature challenge, mice were held at 23°C with HFD-fed mice presenting significantly higher body weights than chow fed mice (48.2 g vs 28.3 g, Figure 1B). In the cold, mice on chow diet maintained a steady body weight, while mice on HFD lost about 10% body weight (Figure 1B, C). All mice in the cold showed a significant drop in body temperature during the first days, but this loss in temperature was significantly higher in mice on HFD (-0.5°C vs -2°C after d1, Figure 1D). After 7 days, mice on chow diet recovered their body temperature to normal levels, while mice on HFD still had significantly lower temperatures (-1°C) (Figure 1D, E). Under thermoneutrality, mice on HFD had significantly higher BAT surface temperatures, consistent with increased diet-induced thermogenesis, but BAT surface temperatures were not different after cold exposure (Figure 1F, G). Extended phenotyping data is presented in Suppl. Figure 1.

3.2. Obesity impairs core transcriptome remodeling of BAT in response to cold

Gene expression data was used from all mice samples that also had methylation signal data available. After removal of outliers, N = 29 samples ($N_{HFD\_30°C}$ = 7; $N_{HFD\_8°C}$ = 7; $N_{chow\_30°C}$ = 7; $N_{chow\_8°C}$ = 8) remained for subsequent differential gene expression and methylation analysis. Analysis of VST normalized RNAseq expression counts shows clear clustering of all diet (chow, HFD)- and temperature (8°C, 30°C)-specific subgroups in the principal component analysis (Figure 2A), providing a solid basis for subsequent differential gene expression analysis. Moreover, temperature groups are separated by principal component 1 describing 33% of variance while diet groups are rather distinguished by principal component 2 covering 16,8% of total variance demonstrating an overall stronger impact of temperature compared to diet on overall gene expression patterns. Mice on HFD are older compared to chow fed mice. The age gap of 11 weeks could potentially interfere and explain a variance in the principal component analyses which was not addressed here.

To dissect BAT's physiological response to cold normal-weight control mice, we compared chow-fed mice held in the cold (8°C) with mice held at thermoneutrality (30°C) for one week. We will refer to this condition pair as $COLD_{chow}$. Additionally, to assess the effects of obesity on cold-induced BAT activation, we compared HFD-induced obese mice held in the cold (8°C) with mice held at thermoneutrality (30°C) for one week. This comparison will be referred to as $COLD_{HFD}$. Differential gene expression analysis (FDR <0.05) resulted in 7,089 DEGs for the $COLD_{chow}$ (Suppl. Figure 2A, B; Suppl. Table 2) and 5,249 DEGs for the $COLD_{HFD}$ (Suppl. Figure 2C, D; Suppl. Table 3) condition pairs. Reflecting the dominating effect of temperature on BAT activity, the majority of the significant DEGs overlap between the $COLD_{chow}$ and $COLD_{HFD}$ conditions are regulated similarly (N = 3,786; $COLD_{chow}$: 53%; $COLD_{HFD}$: 72%). Furthermore, 45% and 26% are

specifically regulated in $COLD_{chow}$ and $COLD_{HFD}$, respectively, while a total of 92 DEGs are
contrarily regulated (Suppl. Figure 4A).

To benchmark our $COLD_{chow}$ results with previously published data, we overlapped our DEGs with
those reported by Taylor et al. (2024) [32], who performed a comparable study in chow-fed male
mice exposed to severe cold (8 °C) versus thermoneutrality (28 °C) (Suppl. Figure 3; Suppl. Table
2). We found that 80% of our upregulated and 52% of our downregulated genes overlapped with
their DEGs, demonstrating a strong concordance across both directions in these independent
analyses.

Additionally, to investigate how diet affects BAT activation in response to cold exposure, we
employed an interaction model ($\Delta COLD_{diet}$) that compares the difference between both $COLD_{HFD}$
and $COLD_{chow}$ contrasts, rather than relying solely on simple group comparisons. This approach
enabled us to assess the combined and potentially synergistic or antagonistic effects of diet and
environment on BAT function [33]. Although an age gap of 11 weeks is indicated, mice on chow
or HFD are in a comparable age-range regarding BAT function and diet-independent body weight
development [34–37]. In total, we observed 1,364 genes (FDR <0.05) which are differentially
expressed between $COLD_{HFD}$ and $COLD_{chow}$ (Figure 2B, C; Suppl. Table 4). Since the most
significant DEGs are downregulated in the $COLD_{HFD}$ condition compared to $COLD_{chow}$, this
highlights a potential mechanism by which obesity may impair thermogenic capacity and energy
expenditure, suggesting that a high-fat diet disrupts the normal activation of BAT in response to
cold exposure.

3.3. Associations of gene expression and methylation signals underscore the influence of diet on
gene regulatory mechanisms

To explore the associative effects between transcriptomics and epigenomics, we correlated all
expression signals of DEGs with significant DMPs found within the DMRs of the corresponding
genes. This approach allowed us to define a set of differentially methylated and expressed genes
(DMEGs), using specific criteria: a Pearson correlation coefficient greater than ± 0.5, DEGs with
a $\log_2$ FC greater than ± 0.5 (adj. P <0.05), and DMPs with a $\log_2$ FC greater than ± 1 (p <0.05).
Importantly, we set a higher threshold for DMPs and focused on DMEGs located in both the
promoter and gene body regions, ensuring that our analysis captures the most relevant regulatory
elements influencing gene expression. To verify our findings, we analyzed the effect of DNA-
demethylation (using Aza) or hypermethylation (using SAM) on the expression of selected genes
in mouse adipocytes *in vitro* (Suppl. Figure 5). The observed effects of DNA methylation and the
manipulation on gene expression were in line with the results of our analysis correlating gene
expression and methylation and confirmed epigenetic control of gene expression in adipocytes.

In the $COLD_{chow}$ control comparison, we identified a total of 3,142 DMEGs among 21% (N =
1,524) of all DEGs. The data showed a balanced distribution of DMEGs, with 55% located within
gene bodies and 45% found in promoter regions (Figure 3A; Suppl. Table 5), indicating that
methylation can affect gene regulation through various mechanisms, such as promoter accessibility

or splice site manipulation. As illustrated in Figure 3B, DMEGs within the top 100 DMRs (ranked by absolute $\log_2$ FC) are predominantly found in promoter regions and show distinctly higher prevalence on chromosome 11.

To investigate which cell types might drive DNA methylation changes in BAT, we overlapped DMEGs from the $COLD_{chow}$ condition with cell-type-specific marker genes derived from single-cell RNAseq (scRNAseq) of BAT from male mice exposed to cold (5 °C) versus thermoneutrality (30 °C) [38]. This analysis revealed that DMEGs were distributed across multiple cell types, with the highest proportions in vascular smooth muscle cells (VSM, 43%), adipocytes (35%), endothelial cells (EC, 31.8%), and stromal populations, including $Pdgfra^+$ adipocyte progenitor cells (APCs, 28.1%) and pericytes (20.9%). In contrast, immune cells contributed more modestly, with macrophages at 18.2%, NK cells at 9.3%, and T cell subsets (CD4$^+$, CD8$^+$, Treg) ranging from 5–9%. Across nearly all cell types, most DMEGs belonged to the hypo up category - genes exhibiting hypomethylation accompanied by increased expression. This pattern was particularly pronounced in adipocytes (26.4%) and vascular populations, including VSM (21.6%) and EC (11.9%). Hyper up DMEGs were most notable in adipocytes (10%), whereas hypo down DMEGs were enriched in vascular and stromal populations, including VSM (15%), EC (14.9%), $Pdgfra^+$ APCs (13.6%), and to a lesser extent in adipocytes (2.8%). Conversely, the fractions of hyper down DMEGs were relatively small, ranging from 1-6% depending on the cell type (Figure 3C; Suppl. Table 5).

Additionally, we detected 1,886 DMEGs from 19% (N = 1001) of all DGEs in the $COLD_{HFD}$ condition. Similar to the $COLD_{chow}$ condition, 60% of the DMEGs are located within gene bodies and 40% in promoter regions (Figure 4A; Suppl. Table 6); however, the most highly regulated DMEGs do not exhibit significant enrichment in the promoter regions (Figure 4B).

Although we previously observed that the majority of significant DEGs overlapping between the $COLD_{chow}$ and $COLD_{HFD}$ conditions are regulated in a similar direction (Suppl. Figure 5), we identified a contrasting pattern among the DMEGs. The gene expression of DMEGs shows consistent regulatory patterns, with only 245 ($COLD_{chow}$: 16%; $COLD_{HFD}$: 24%) genes being upregulated and 159 ($COLD_{chow}$: 10%; $COLD_{HFD}$: 16%) genes being downregulated in both conditions (Suppl. Figure 4B; Suppl. Figure 6). In contrast, 73% and 60% are specifically regulated in the $COLD_{chow}$ and $COLD_{HFD}$ conditions, respectively, while only one gene (*guanylyl cyclase domain containing 1* (*Gucd1*)) is regulated in the opposite direction (Suppl. Figure 4B; Suppl. Figure 7). This suggests that the response to temperature conditions is strongly dependent on the specific environment and that methylation changes may be purposefully employed to adapt to different metabolic or physiological demands.

Our previous findings revealed pronounced differences in the regulatory patterns of DMEGs between the $COLD_{chow}$ and $COLD_{HFD}$ conditions by emphasizing the direction of regulation. To further elucidate the nuances of diet-specific regulation, we utilized the $\Delta COLD_{diet}$ contrast to detect more subtle differences in regulatory effects including regulation strength, directly

identifying diet-related differences in DMEGs associated with BAT activation induced by cold exposure (Figure 5; Suppl. Table 7). For the $\Delta COLD_{diet}$ contrast, we detected 65 DMEGs from only 4% (N = 56) of all DEGs. Interestingly, these are primarily enriched in the gene bodies (69%), suggesting that the regulation of these genes may be more influenced by post-transcriptional mechanisms in response to cold exposure.

3.4. Adaptive divergence in thermogenic signaling reflects metabolic stress in obese BAT

For further characterization of the reported DMEGs, pathway enrichment analysis was performed using the IPA software. The DMEGs were categorized into four groups: those exhibiting similar regulation in both $COLD_{chow}$ and $COLD_{HFD}$ condition pairs, and those that are uniquely regulated in each individual condition ($\Delta COLD_{diet,}$ $COLD_{chow}$, and $COLD_{HFD}$). Querying the catalog for significant (BH-adjusted p <0.05) canonical pathways (Figure 6; Suppl. Table 8) shows similar activation of pathways with respect to lipid metabolism except for ketolysis which is only significantly activated in the $COLD_{HFD}$ condition. Cell signaling and energy expenditure seem to be activated or inhibited more differentially between $COLD_{chow}$ and $COLD_{HFD}$ condition pairs. For example, the process of white adipose tissue browning is significantly inhibited in $COLD_{HFD}$ but shows (unsignificant) activation in $COLD_{chow}$. Additionally, type II diabetes mellitus signaling is significantly activated under $COLD_{HFD}$ considerations but tends to be (insignificantly) inhibited under $COLD_{chow}$.

3.5. Diet and cold exposure shape epigenetic regulation in BAT

To gain deeper insights into the epigenetic mechanisms that regulate BAT activation and thermogenesis under different dietary conditions, we matched DEGs from each condition pair with the EpiFactors database (v2.0, https://epifactors.autosome.org, accessed on 10.09.2024), with the analysis being restricted to DNA-methylation-related factors, while histone-modifying enzymes, chromatin remodelers and RNA-based epigenetic regulators were omitted. As summarized in Figure 7, we identified eleven differentially expressed methylation regulators for the $COLD_{chow}$, seven for $COLD_{HFD}$, and four for $\Delta COLD_{diet}$ condition pairs. Notably, five regulators were found to be common to both the $COLD_{chow}$ and $COLD_{HFD}$ conditions including *Tet2* (*ten-eleven translocation methylase 2*), *Dnmt3* (*DNA methyltransferase 3*), *Cenpc1* (*centromere protein C*), *Apobec1* (*apolipoprotein B mRNA editing enzyme catalytic subunit 1*) which are all downregulated in $COLD_{chow}$ and $COLD_{HFD}$. This overlap indicates a conserved epigenetic mechanism that is essential for the physiological response to cold exposure, regardless of diet. However, the majority of methylation enzymes exhibited unique regulation within each condition pair, highlighting the nuanced regulation of epigenetic mechanisms based on dietary composition. For instance, *Tet3* and *Dnmt1* were differently regulated within the $\Delta COLD_{diet}$ and $COLD_{chow}$ conditions, while *Mettl4* (*methyltransferase-like protein 4*) and *Apobec3b* showed differential regulation within the $\Delta COLD_{diet}$ contrast.

4. Discussion

We aimed to explore how environmental factors and DNA methylation may influence BAT function. To investigate potential links between DNA methylation and gene expression, we integrated RNA-seq data with RRBS methylation profiles to identify genes with coordinated changes in expression and methylation (DMEGs) under different dietary and thermal conditions. This approach provides insights into BAT plasticity and adaptive thermogenesis, contributing to a foundational understanding that could inform future obesity-related research and interventions.

Consistent with BAT's physiological adaptation to cold, our data show that cold exposure robustly increases BAT activity regardless of diet, evidenced by strong transcriptomic responses with 7,089 DEGs in chow-fed and 5,249 DEGs in HFD-fed mice. A substantial overlap (~3,780 genes) between both dietary groups suggests a conserved "core" thermogenic program that remains inducible under obesogenic conditions. Notably, *Ucp1*, a central mediator of non-shivering thermogenesis, and *Letmd1* (*leucine zipper-EF-hand-containing transmembrane protein domain containing 1*), which is essential for mitochondrial integrity and cold response, were among the top upregulated genes across both diets. Their consistent induction under cold exposure, aligns with previous reports of diet-insensitive thermogenic regulators [39,40]. Beside this shared response, ~1,360 genes were differentially expressed between the cold-exposed groups ($\Delta COLD_{diet}$), with most downregulated in diet-induced obese mice, consistent with a high-fat diet impairing thermoregulatory capacity driven by mitochondrial dysfunction, inflammation, and insulin resistance [41,42]. This functional attenuation under high-fat diet conditions is also reflected among the most strongly regulated DGEs. Downregulation of *Hk2* (*hexokinase 2*), a key glycolytic enzyme, has been linked to reduced glucose uptake [43]. Similarly, reduced *Gpd1* (*glycerol-3-phosphate dehydrogenase 1*) expression, part of the glycerol-3-phosphate shuttle, may reflect compromised lipid metabolism. *Ucp1* stayed cold-inducible in all diets but was lower in HFD-fed mice, showing that thermogenic response is maintained but reduced under metabolic stress. This aligns with reports of reduced BAT activity in obesity despite retained gene expression [39]. Our results of $COLD_{chow}$ are in line with those of Taylor et al. (2024) [32], who performed a comparable study in chow-fed mice exposed to severe cold (8°C) versus thermoneutrality (28°C). The strong overlap of DEGs highlights the reproducibility of cold-induced BAT transcriptional responses across independent studies and supports the generalizability of our findings.

To identify epigenetically regulated genes, we correlated methylation changes (M-values of DMPs overlapping DMRs) with gene expression levels to define DMEGs. While these correlations suggest potential regulatory links, they do not establish causality, as gene expression is also influenced by other epigenetic modifications and transcription factors [32,44,45]. More comprehensive approaches integrating histone modifications, proteomics, or single-cell epigenomics could further elucidate the complex regulation within heterogeneous BAT cell populations [46]. Compared to generalized linear models with distributional assumptions [47,48], our site-specific correlation approach offers higher sensitivity for detecting local methylation-expression relationships, though it may miss broader regulatory patterns and is more sensitive to

noise. Our thresholds for defining significant DMEGs were set at |log FC| >0.5 for expression
changes and |log FC| >1 for methylation differences; correlations were considered significant if |r|
>0.5 and p <0.05. These cutoffs balance sensitivity and specificity but also limit detection power
given our sample size. When DMEGs overlapped promoter and gene body regions, assignment
was based on the strongest correlation, which enhances interpretability but may introduce bias due
to regional variability or technical factors.

By analyzing DMEGs, ~3,100 DMEGs (21% of DGEs) were identified in $COLD_{chow}$, with a
roughly equal distribution between promoter and gene body methylation. In contrast, HFD-fed
mice exhibited only ~1,900 DMEGs (19% of DEGs), with a higher proportion of gene body
methylation. As gene body methylation is often linked to transcript fine-tuning rather than
activation, this may reflect constrained thermogenic adaptability under obesogenic conditions [49].
Notably, DMEGs were largely condition-specific despite substantial overlap in DEGs, highlighting
that diet and cold exposure interact to shape BAT-specific regulatory states, rather than acting
independently. These condition-specific methylation-expression relationships suggest that DNA
methylation may play an active role in modulating transcriptional plasticity and energy
metabolism, enabling BAT to adapt epigenetically to distinct environmental challenges.

To gain further insight into the cellular context of these DNA methylation changes, we mapped
DMEGs of the $COLD_{chow}$ condition onto cell-type-specific markers derived from male mice
scRNAseq data of cold-exposed BAT [38]. This analysis revealed that the majority of DMEGs
belonged to the hypo up category, particularly in adipocytes, vascular cells, and $Pdgfra^+$ adipocyte
progenitors, suggesting coordinated activation of thermogenic programs and vascular remodeling.
Hypo down DMEGs were enriched in vascular and stromal populations, indicating selective
repression of pathways supporting tissue structure and function. Hyper up DMEGs were primarily
observed in adipocytes, whereas hyper down DMEGs were rare (1–6%), implying that DNA
methylation mediated silencing plays a limited role in the cold response. The observed cell type-
specific DMEG distributions are consistent with prior evidence of increased intercellular signaling
among adipocytes, vascular, stromal, and immune cells under cold conditions in chow-fed mice,
suggesting an epigenetic basis for the transcriptional remodeling in BAT [38].

Pathway enrichment analysis further supports this adaptive divergence of DMEGs and highlighting
the diet-dependent rewiring of thermogenic signaling. Cold exposure activated distinct lipid
metabolic pathways (triacylglycerol, phosphatidylglycerol, and CDP−diacylglycerol biosynthesis)
in both dietary groups, however, ketolysis was selectively enriched in HFD-fed mice. In line with
previous findings that HFD suppresses ketone body utilization for lipogenesis in BAT via
downregulation of *acetoacetyl-CoA synthetase* (*Aacs*) [50], our DGE analysis show that *Aacs* was
significantly upregulated by cold in both dietary groups, but this response was markedly stronger
in chow-fed mice ($\log_2$ FC: 1.77) than in HFD-fed mice ($\log_2$ FC: 0.74), with a negative interaction
effect ($\Delta COLD_{diet}$ l $\log_2$ FC: –0.81). These results indicate that cold-induced ketone body recycling
remains inducible but is dampened under HFD, potentially limiting metabolic flexibility and
compensatory lipid synthesis in insulin-resistant BAT. Consistently, only $COLD_{HFD}$ showed

activation of the Type II Diabetes Mellitus signaling pathway under cold stress, reflecting a
diabetogenic transcriptional environment marked by insulin resistance and impaired glucose
homeostasis [51]. This suggests that cold exposure does not fully overcome metabolic dysfunction
in BAT during obesity, potentially limiting its systemic metabolic benefits.

Further, cold-exposed HFD mice uniquely activated Sirtuin signaling, potentially a compensatory
response to mitochondrial stress, while suppressing pathways related to mitochondrial dysfunction
and white adipose tissue browning, indicating a shift from efficient thermogenesis to cellular
maintenance [52]. Further, a selective activation of neuroendocrine pathways (e.g., corticotropin-
releasing hormone and orexin signaling) may reflect rewiring of central and peripheral regulatory
mechanisms to compensate for reduced intrinsic thermogenic capacity [53]. Concurrent
suppression of nitric oxide and neuronal nitric oxide synthase (nNOS) signaling, crucial for
vascular perfusion and mitochondrial function, points to impaired BAT structural and vascular
adaptation under HFD.

We identified several epigenetic regulators differentially expressed in response to cold and diet.
Core factors such as *Tet2*, *Dnmt3*, *Cenpc1*, and *Apobec1* responded similarly under both diets,
suggesting a foundational epigenetic program supporting cold-induced chromatin remodeling and
transcriptomic flexibility. In contrast to white adipose tissue, where *Mettl4* is upregulated under
obesogenic conditions [54], we observed its downregulation in BAT only in $COLD_{HFD}$. Given that
Mettl4-mediated N6-adenine methylation has been implicated in adipocyte differentiation and lipid
production [55], it is conceivable that high-fat diets constrain the differentiation and functional
capacity of BAT through epigenetic mechanisms. Conversely, cold exposure decreased *Tet3* and
Dnmt1 expression under chow but not HFD. As Tet3 and Dnmt1 regulate adipocyte expansion and
mitochondrial function as well as chromatin architecture, respectively, their cold-induced
downregulation may favor thermogenesis by limiting hypertrophy and enhancing mitochondrial
and epigenetic adaptability [56–58]. *Dnmt1* deletion in BAT has no metabolic impact under
standard diets [59], but may gain relevance under thermogenic or obesogenic stress. A previous
study demonstrated the downregulation of *Dnmt1*, *Dnmt3a,* and *Dnmt3b* in BAT upon cold
exposure under chow diet conditions, linking reduced DNA methylation to enhanced thermogenic
activation [60]. Our findings confirm this pattern under chow conditions but reveal a lack of
regulation under HFD, supporting the notion that epigenetic flexibility is impaired in obesogenic
environments. *Apobec3*, a cytidine deaminase implicated in RNA and potentially DNA editing
[61], was downregulated by cold under both diets, though significantly more under chow. This
pattern may reflect its role in facilitating transcriptomic remodeling during thermogenic activation
that may be blunted under HFD. Therefore, BAT activation appears to be governed by a conserved
core of methylation regulators responsive to cold, while additional, context-specific pathways are
modulated by the dietary environment. This dual mechanism supports a model in which a stable
epigenetic framework is dynamically adjusted to accommodate metabolic demands, thereby fine-
tuning the thermogenic capacity of BAT.

By conducting in vitro methylation analyses, we confirmed the epigenetic transcription control of certain DMEGs and important BAT regulatory genes such as *Bmp8b* or *Cpne5* pointing towards a valid impact of the DNA methylation on BAT function. By using Aza and SAM, we cannot rule out the possibility that other epigenetic mechanisms than DNA methylation led to these observations as especially SAM serves as methyldonor for various processes [62]. In any case, our results strongly support the highly dynamic control of BAT by epigenetic alterations.

4.1. Limitations

A limitation arises from age differences between diet groups (11 weeks chow vs. 22 weeks HFD). First, the difference in bodyweight between chow and HFD mice may have partially arisen from the age gap, as there is still a slow increase in body weight beyond adulthood at 12 weeks of age [37]. However, male chow fed mice at 22 weeks of age would reach approximately 30g bodyweight which still is significantly different from mice on HFD at the same age. Second, although BAT function declines with age, mice used in this study are of a comparable age-range related to BAT function. Studies show that mice aged between 12 weeks and six months present a stable metabolic active BAT, with stable mitochondrial content and thermogenic potential before these factors start to decline and lipid accumulation increases [34–36]. Although we cannot bioinformatically address this issue in our comparative analyses, we believe that age is not a main driving factor of the here observed effects as we addressed a comparable age range regarding bodyweight and cold adaptation of BAT. Anyway, regardless of BAT function, global DNA methylation reduces with age while specifically promoter regions tend to get hypermethylated with age [63]. While we cannot address the age factor in our analyses due to group specific age differences, we expect a limited impact of age on our results based on comparing mice in a suitable age range meaning only "young adult mice" [64]. A further limitation is the use of only male mice, which may limit generalizability to females. However, given the known sex-specific differences in adipose biology, BAT activity, and epigenetic regulation, addressing only males is beneficial regarding the addressed methodology. Future studies should include female mice to assess potential sex-by-diet or sex-by-temperature interactions. Small sample sizes (7-8 per group) limit statistical power to detect subtle effects. In addition, RRBS data provides only partial genome coverage, primarily focusing on CpG-rich regions, potentially overlooking important regulatory areas outside these regions. In contrast, whole-genome bisulfite sequencing could provide more comprehensive methylation profiling.

5. Conclusions

In conclusion, this study provides the first evidence that cold-induced DNA methylation remodeling in BAT is strongly shaped by dietary context, particularly under obesogenic conditions. Our integrative analysis reveals that, while cold exposure activates a conserved thermogenic transcriptional program, high-fat diet attenuates epigenetic flexibility and restricts full BAT activation. These findings underscore that environmental temperature and dietary intake interact to regulate BAT function via interconnected transcriptional and epigenetic mechanisms. Importantly,

we identify both core and context-specific regulators that mediate this response, offering potential molecular targets for future therapies. By demonstrating that obesity alters the epigenetic landscape of cold adaptation, our study opens new avenues for nutritional or epigenetic interventions aimed at restoring BAT plasticity and enhancing thermogenic capacity in metabolic disease.

Data Availability

Raw RNAseq and RRBS data have been deposited in the Sequence Read Archive (SRA, https://www.ncbi.nlm.nih.gov/sra/) [65] under the BioProject number PRJNA1294891.

Acknowledgements

We acknowledge the technical support of the Core Unit DNA Technologies of the Faculty of Medicine, University of Leipzig, and thank Dr. Knut Krohn, Stefanie Ziesche and Kathleen Schön. We thank Jenny Schuster and Lisa Gärtner from the animal facility, and Ines Müller, Claudia Gebhardt and Olivia Paetow for excellent technical assistance.

Author contributions

Conceptualization: TH, AH, KRZ, JTH, and JW; Methodology: AH, TH, KRZ, and JW; Investigation: KRZ, HB, and JW; Validation: KRZ, HB, and JW; Formal analysis: TH; Data curation: TH and AH; Writing – original draft: TH, AH, KRZ, JW, and JTH; Writing – review & editing: TH, AH, KRZ, HB, LM, PK, MS, MB, JTH, and JW; Visualization: TH; Supervision: AH and JTH; Funding acquisition: MB, JW, and JTH.

Funding

This work was funded or supported by grants of the Deutsche Forschungsgemeinschaft, project number 209933838 (SFB1052 "Obesity Mechanisms": B1 to MB, C7 to JTH) and by the Formel1 Nachwuchsförderung of the Medical Faculty of the University of Leipzig (JW). HB is supported by a doctoral scholarship of the Studienstiftung des Deutschen Volkes. LM is supported by the Swedish Research Council, the European for the Study of Diabetes and the German Diabetes Association.

Conflict of interest

MB received honoraria as a consultant and speaker from Amgen, AstraZeneca, Bayer, Boehringer-Ingelheim, Lilly, Novo Nordisk, Novartis, and Sanofi. All other authors declare no conflicts of interest. The funders had no role in the design of the study, the collection, analyses, or interpretation of data, in the decision to publish the results and the writing of the manuscript.

References

[1]   Kusminski CM, Bickel PE, Scherer PE. Targeting adipose tissue in the treatment of obesity-associated diabetes. Nat Rev Drug Discov 2016;15:639–60. https://doi.org/10.1038/nrd.2016.75.

[2]   Saito M, Matsushita M, Yoneshiro T, Okamatsu-Ogura Y. Brown Adipose Tissue, Diet-Induced Thermogenesis, and Thermogenic Food Ingredients: From Mice to Men. Front Endocrinol 2020;11. https://doi.org/10.3389/fendo.2020.00222.

[3]   U Din M, Saari T, Raiko J, Kudomi N, Maurer SF, Lahesmaa M, et al. Postprandial Oxidative Metabolism of Human Brown Fat Indicates Thermogenesis. Cell Metab 2018;28:207-216.e3. https://doi.org/10.1016/j.cmet.2018.05.020.

[4]   Chouchani ET, Kazak L, Spiegelman BM. New Advances in Adaptive Thermogenesis: UCP1 and Beyond. Cell Metab 2019;29:27–37. https://doi.org/10.1016/j.cmet.2018.11.002.

[5]   Rosenbaum M, Leibel RL. Adaptive thermogenesis in humans. Int J Obes 2010;34:S47–55. https://doi.org/10.1038/ijo.2010.184.

[6]   Scheele C, Wolfrum C. Brown Adipose Crosstalk in Tissue Plasticity and Human Metabolism. Endocr Rev 2020;41:53–65. https://doi.org/10.1210/endrev/bnz007.

[7]   Xiao H, Kang S. The role of DNA methylation in thermogenic adipose biology. Epigenetics 2019;14:837–43. https://doi.org/10.1080/15592294.2019.1625670.

[8]   Wang S, Cao Q, Cui X, Jing J, Li F, Shi H, et al. Dnmt3b Deficiency in Myf5+-Brown Fat Precursor Cells Promotes Obesity in Female Mice. Biomolecules 2021;11:1087. https://doi.org/10.3390/biom11081087.

[9]   Teperino R. Intergenerational metabolic effects of cold exposure. Nat Metab 2025;7:635–6. https://doi.org/10.1038/s42255-025-01222-z.

[10]   Sun S, Hood M, Scott L, Peng Q, Mukherjee S, Tung J, et al. Differential expression analysis for RNAseq using Poisson mixed models. Nucleic Acids Res 2017;45:e106–e106. https://doi.org/10.1093/nar/gkx204.

[11]   Skinner MK. Preconception cold–induced epigenetic inheritance. Nat Med 2018;24:1308–9. https://doi.org/10.1038/s41591-018-0187-3.

[12]   Zhang Q, Xiao X, Zheng J, Li M, Yu M, Ping F, et al. Maternal High-Fat Diet Disturbs the DNA Methylation Profile in the Brown Adipose Tissue of Offspring Mice. Front Endocrinol 2021;12. https://doi.org/10.3389/fendo.2021.705827.

[13]   Madden CJ, Morrison SF. A high-fat diet impairs cooling-evoked brown adipose tissue activation via a vagal afferent mechanism. Am J Physiol-Endocrinol Metab 2016;311:E287–92. https://doi.org/10.1152/ajpendo.00081.2016.

[14]   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinforma Oxf Engl 2014;30:2114–20. https://doi.org/10.1093/bioinformatics/btu170.

[15]   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinforma Oxf Engl 2013;29:15–21. https://doi.org/10.1093/bioinformatics/bts635.

[16]   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinforma Oxf Engl 2014;30:923–30. https://doi.org/10.1093/bioinformatics/btt656.

[17]   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 2014;15:1–21. https://doi.org/10.1186/s13059-014-0550-8.

[18]   Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow enables reproducible computational workflows. Nat Biotechnol 2017;35:316–9.

https://doi.org/10.1038/nbt.3820.

[19]   Ewels P, Hammarén R, Peltzer A, phue, F S, Tommaso PD, et al. nf-core/methylseq 2019.

https://doi.org/10.5281/zenodo.2555454.

[20]   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnetJournal 2011;17:10–2. https://doi.org/10.14806/ej.17.1.200.

[21]   Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-

Seq applications. Bioinformatics 2011;27:1571–2. https://doi.org/10.1093/bioinformatics/btr167.

[22]   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res

2015;43:e47. https://doi.org/10.1093/nar/gkv007.

[23]   Müller F, Scherer M, Assenov Y, Lutsik P, Walter J, Lengauer T, et al. RnBeads 2.0:

comprehensive analysis of DNA methylation data. Genome Biol 2019;20:55.

https://doi.org/10.1186/s13059-019-1664-9.

[24]   Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, et al. BioMart and

Bioconductor: a powerful link between biological databases and microarray data analysis.

Bioinformatics 2005;21:3439–40. https://doi.org/10.1093/bioinformatics/bti525.

[25]   Haider S, Waggott D, Boutros P. bedr: Genomic Region Processing using Tools Such as

'BEDTools','BEDOPS'and 'Tabix' 2019.

[26]   Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software

for Computing and Annotating Genomic Ranges. PLoS Comput Biol 2013;9.

https://doi.org/10.1371/journal.pcbi.1003118.

[27]   HERVE M. RVAideMemoire: Testing and Plotting Procedures for Biostatistics. 2023.

[28]   Microsoft, Weston S. foreach: Provides Foreach Looping Construct. 2022.

[29]   Krämer A, Green J, Pollard J, Tugendreich S. Causal analysis approaches in Ingenuity

Pathway Analysis. Bioinformatics 2014;30:523–30.

https://doi.org/10.1093/bioinformatics/btt703.

[30]   Klein J, Fasshauer M, Klein HH, Benito M, Kahn CR. Novel adipocyte lines from brown

fat: A model system for the study of differentiation, energy metabolism, and insulin action.

BioEssays 2002;24:382–8. https://doi.org/10.1002/bies.10058.

[31]   Weiner J, Rohde K, Krause K, Zieger K, Klöting N, Kralisch S, et al. Brown adipose

tissue (BAT) specific vaspin expression is increased after obesogenic diets and cold exposure and

linked to acute changes in DNA-methylation. Mol Metab 2017;6:482–93.

https://doi.org/10.1016/j.molmet.2017.03.004.

[32]   Taylor BC, Steinthal LH, Dias M, Yalamanchili HK, Ochsner SA, Zapata GE, et al.

Histone proteoform analysis reveals epigenetic changes in adult mouse brown adipose tissue in

response to cold stress. Epigenetics Chromatin 2024;17. https://doi.org/10.1186/s13072-024-

00536-8.

[33]   Nedergaard J, Cannon B. The browning of white adipose tissue: some burning issues. Cell

Metab 2014;20:396–407. https://doi.org/10.1016/j.cmet.2014.07.005.

[34]   Cannon B, De Jong JMA, Fischer AW, Nedergaard J, Petrovic N. Human brown adipose

tissue: Classical brown rather than brite/beige? Exp Physiol 2020;105:1191–200.

https://doi.org/10.1113/EP087875.

[35]   Silva GDN, Amato AA. Thermogenic adipose tissue aging: Mechanisms and

implications. Front Cell Dev Biol 2022;10:955612. https://doi.org/10.3389/fcell.2022.955612.

[36]   Zoico E, Rubele S, De Caro A, Nori N, Mazzali G, Fantin F, et al. Brown and Beige

Adipose Tissue and Aging. Front Endocrinol 2019;10:368.

https://doi.org/10.3389/fendo.2019.00368.

[37]   The Jackson Laboratory. Body Weight Information for B6J (000664) 2025.

https://www.jax.org/jax-mice-and-services/strain-data-sheet-pages/body-weight-chart-000664.

[38]   Shamsi F, Zheng R, Ho L-L, Chen K, Tseng Y-H. Comprehensive analysis of
intercellular communication in the thermogenic adipose niche. Commun Biol 2023;6:761.
https://doi.org/10.1038/s42003-023-05140-2.

[39]   Fromme T, Klingenspor M. Uncoupling protein 1 expression and high-fat diets. Am J
Physiol-Regul Integr Comp Physiol 2011;300:R1–8. https://doi.org/10.1152/ajpregu.00411.2010.

[40]   Choi K-M, Kim JH, Kong X, Isik M, Zhang J, Lim H-W, et al. Defective brown adipose
tissue thermogenesis and impaired glucose metabolism in mice lacking Letmd1. Cell Rep
2021;37:110104. https://doi.org/10.1016/j.celrep.2021.110104.

[41]   Bournat JC, Brown CW. Mitochondrial dysfunction in obesity. Curr Opin Endocrinol
Diabetes Obes 2010;17:446–52. https://doi.org/10.1097/MED.0b013e32833c3026.

[42]   Maliszewska K, Kretowski A. Brown Adipose Tissue and Its Role in Insulin and Glucose
Homeostasis. Int J Mol Sci 2021;22:1530. https://doi.org/10.3390/ijms22041530.

[43]   Shimobayashi M, Thomas A, Shetty S, Frei IC, Wölnerhanssen BK, Weissenberger D, et
al. Diet-induced loss of adipose hexokinase 2 correlates with hyperglycemia. eLife
2023;12:e85103. https://doi.org/10.7554/eLife.85103.

[44]   Kouzarides T. Chromatin Modifications and Their Function. Cell 2007;128:693–705.
https://doi.org/10.1016/j.cell.2007.02.005.

[45]   Lee TI, Young RA. Transcriptional regulation and its misregulation in disease. Cell
2013;152:1237–51. https://doi.org/10.1016/j.cell.2013.02.014.

[46]   Smallwood SA, Lee HJ, Angermueller C, Krueger F, Saadeh H, Peat J, et al. Single-cell
genome-wide bisulfite sequencing for assessing epigenetic heterogeneity. Nat Methods
2014;11:817–20. https://doi.org/10.1038/nmeth.3035.

[47]   Gevaert O, Tibshirani R, Plevritis SK. Pancancer analysis of DNA methylation-driven
genes using MethylMix. Genome Biol 2015;16:17. https://doi.org/10.1186/s13059-014-0579-8.

[48]   Mishra NK, Guda C. Genome-wide DNA methylation analysis reveals molecular
subtypes of pancreatic cancer. Oncotarget 2017;8:28990–9012.
https://doi.org/10.18632/oncotarget.15993.

[49]   Moore LD, Le T, Fan G. DNA Methylation and Its Basic Function.
Neuropsychopharmacology 2013;38:23–38. https://doi.org/10.1038/npp.2012.112.

[50]   Yamasaki M, Hasegawa S, Ozaki S, Imai M, Saito D, Takahashi N. High-Fat-Diet
Suppressed Ketone Body Utilization for Lipogenic Pathway in Brown Adipose Tissues.
Metabolites 2023;13:519. https://doi.org/10.3390/metabo13040519.

[51]   Chondronikola M, Volpi E, Børsheim E, Porter C, Saraf MK, Annamalai P, et al. Brown
Adipose Tissue Activation Is Linked to Distinct Systemic Effects on Lipid Metabolism in
Humans. Cell Metab 2016;23:1200–6. https://doi.org/10.1016/j.cmet.2016.04.029.

[52]   Escalona-Garrido C, Vázquez P, Mera P, Zagmutt S, García-Casarrubios E, Montero-
Pedrazuela A, et al. Moderate SIRT1 overexpression protects against brown adipose tissue
inflammation. Mol Metab 2020;42:101097. https://doi.org/10.1016/j.molmet.2020.101097.

[53]   Madden CJ, Tupone D, Morrison SF. Orexin modulates brown adipose tissue
thermogenesis. Biomol Concepts 2012;3:381–6. https://doi.org/10.1515/bmc-2011-0066.

[54]   Kang Q, Zhu X, Ren D, Ky A, MacDougald OA, O'Rourke RW, et al. Adipose
METTL14-Elicited N^6-Methyladenosine Promotes Obesity, Insulin Resistance, and NAFLD
Through Suppressing β Adrenergic Signaling and Lipolysis. Adv Sci 2023;10.
https://doi.org/10.1002/advs.202301645.

[55]   Zhang Z, Hou Y, Wang Y, Gao T, Ma Z, Yang Y, et al. Regulation of Adipocyte
Differentiation by METTL4, a 6 mA Methylase. Sci Rep 2020;10.
https://doi.org/10.1038/s41598-020-64873-w.

[56]   Jung BC, You D, Lee I, Li D, Schill RL, Ma K, et al. TET3 plays a critical role in white
adipose development and diet-induced remodeling. Cell Rep 2023;42:113196.
https://doi.org/10.1016/j.celrep.2023.113196.
[57]   Park J, Lee DH, Ham S, Oh J, Noh J-R, Lee YK, et al. Targeted erasure of DNA
methylation by TET3 drives adipogenic reprogramming and differentiation. Nat Metab
2022;4:918–31. https://doi.org/10.1038/s42255-022-00597-7.
[58]   Kajimura S. The epigenetic regulation of adipose tissue plasticity. Proc Natl Acad Sci
2021;118. https://doi.org/10.1073/pnas.2102944118.
[59]   Park YJ, Lee S, Lim S, Nahmgoong H, Ji Y, Huh JY, et al. DNMT1 maintains metabolic
fitness of adipocytes through acting as an epigenetic safeguard of mitochondrial dynamics. Proc
Natl Acad Sci 2021;118. https://doi.org/10.1073/pnas.2021073118.
[60]   Taylor BC, Steinthal LH, Dias M, Yalamanchili HK, Ochsner SA, Zapata GE, et al.
Histone proteoform analysis reveals epigenetic changes in adult mouse brown adipose tissue in
response to cold stress. Epigenetics Chromatin 2024;17:12. https://doi.org/10.1186/s13072-024-
00536-8.
[61]   Silvas TV, Schiffer CA. APOBEC3s: DNA-editing human cytidine deaminases. Protein
Sci Publ Protein Soc 2019;28:1552–66. https://doi.org/10.1002/pro.3670.
[62]   Schmidt T. S-Adenosylmethionine affects ERK1/2 and STAT3 pathway in androgen-
independent prostate cancer cells. Mol Biol Rep 2022;49:4805–17.
https://doi.org/10.1007/s11033-022-07331-2.
[63]   Perez-Correa J-F, Tharmapalan V, Geiger H, Wagner W. Epigenetic Clocks for Mice
Based on Age-Associated Regions That are Conserved Between Mouse Strains and Human.
Front Cell Dev Biol 2022;10:902857. https://doi.org/10.3389/fcell.2022.902857.
[64]   Berry DC, Stenesen D, Zeve D, Graff JM. The developmental origins of adipose tissue.
Development 2013;140:3939–49. https://doi.org/10.1242/dev.080549.
[65]   Katz K, Shutov O, Lapoint R, Kimelman M, Brister JR, O'Sullivan C. The Sequence
Read Archive: a decade more of explosive growth. Nucleic Acids Res 2022;50:D387–90.
https://doi.org/10.1093/nar/gkab1053.
780

    Figures

Figure 1: Phenotyping of chow and HFD-fed mice housed at 30°C or 8°C. (A) Schematic
overview and corresponding color code of the cohort mouse groups stratified by dietary (chow,
HFD) and environmental (thermoneutrality vs cold) conditions. (B) Body weight development and
(C) body weight change in chow and HFD-fed mice housed at thermoneutrality (30°C) or in the
cold (8°C) for 7 days. Body temperature (D) and temperature change (E) in chow and HFD-fed
mice housed at thermoneutrality (30°C) or in the cold (8°C) for 7 days. (F) Thermal images from
brown adipose tissue (BAT) and (G) BAT surface temperature in chow and HFD-fed mice housed
at thermoneutrality (30°C) or in the cold (8°C) for 7 days. N = 7-8 per condition. (B, D) Statistical
significance was evaluated by two-way ANOVA with uncorrected Fischer's LSD, (E) Šídák's post-
hoc test, (C, G) or one-way ANOVA with Tukey's post-hoc test. $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$
(***). Scale bar: 100 µm.

**Figure 2: RNA sequencing-based gene expression data revealse distinct clusters of treatment
groups.** (A) Principal component analysis separates temperature and diet. All considered
subgroups reveal distinct expression profiles and cluster according to the first two principal
components. (B) Volcano plot highlights most significantly (false discovery rate (FDR) <0.05)
regulated genes under *ΔCOLD*$_{diet}$ condition. (C) Heatmap of expression values shows that the
subgroups can be clearly separated by the top significant regulated genes. Abbreviations: HFD:
high fat diet; PC: principal component.

Figure 3: Multi-layered insights into DMEGs under *COLD$_{chow}$* condition. (A) The scatter plot
displays the log$_2$ fold changes (FC) in gene expression and DNA methylation signals for all
differentially methylated and expressed genes (DMEGs) in both gene body and promoter contexts
under the *COLD$_{chow}$* condition. Colors indicate the direction of regulation. DMEGs with absolute
log$_2$ FC >0.5 in expression and >1 in methylation signals are highlighted as significant, while
DMEGs in the 5% and 95% quantile of both differentially methylated position (DMP) and
differentially expressed gene (DEG) log$_2$ FC signals are labeled. (B) The circos visualization
illustrates the DMEGs covering the top 100 significant differentially methylated regions (DMRs;
ranked by log$_2$ FC). This plot integrates multiple layers of information from outer to inner ring: Dot
colors indicate the direction of regulation, while dot positions denote whether the DMR of the
DMEG is located in the gene body or promoter context (1); a stylized ideogram of chromosomes
provides a genomic reference frame; and a heatmap displays scaled log2 FC for both gene
expression and DNA methylation signals in gene body and promoter contexts for each DMEG (2).
(C) Overview of assigned cell types to the DMEGs, categorized as hypo up, hypo down, hyper up,
and hyper down. The color scale reflects the percentage of DMEGs, and the size of the dots reflects
the absolute number of DMEGs.

Figure 4: Multi-layered insights into DMEGs under _COLD_$_{HFD}$ condition. (A) The scatter plot
displays the log$_2$ FC in gene expression and DNA methylation signals for all DMEGs in both gene
body and promoter contexts under the _COLD_$_{HFD}$ condition. Colors indicate the direction of
regulation. DMEGs with absolute log$_2$ FC >0.5 in expression and >1 in methylation signal are
highlighted as significant, while DMEGs in the 5% and 95% quantile of both DMP and DEG log$_2$
FC signals are labeled. (B) The circos visualization illustrates the DMEGs covering the top 100
most significant DMRs (ranked by log$_2$ FC). This plot integrates multiple layers of information
from outer to inner ring: Dot colors indicate the direction of regulation, while dot positions denote
whether the DMR of the DMEG is located in the gene body or promoter context (1); a stylized
ideogram of chromosomes provides a genomic reference frame; and a heatmap displays scaled log$_2$
FC for both gene expression and DNA methylation signals in gene body and promoter contexts for
each DMEG (2).

Figure 5: Multi-layered insights into DMEGs under $\varDelta COLD_{diet}$ condition. (A) The scatter plot
displays the $\log_2$ FC in gene expression and DNA methylation signals for all DMEGs in both gene
body and promoter contexts under the $\varDelta COLD_{diet}$ condition. Colors indicate the direction of
regulation. DMEGs with absolute $\log_2$ FC >0.5 in expression and >1 in methylation signals are
highlighted as significant, while DMEGs in the 5% and 95% quantile of both DMP and DEG $\log_2$
FC signals are labeled. (B) The circos visualization illustrates all detected DMEGs in their
respective DMRs. This plot integrates multiple layers of information from outer to inner ring: Dot
colors indicate the direction of regulation, while dot positions denote whether the DMR of the
DMEG is located in the gene body or promoter context (1); a stylized ideogram of chromosomes
provides a genomic reference frame; and a heatmap displays scaled $\log_2$ FC for both gene
expression and DNA methylation signals in gene body and promoter contexts for each DMEG (2).

Figure 6: Pathway enrichment analysis of DMEGs for canonical pathways, diseases, and biological functions. Ingenuity Pathway Analysis (IPA) with enrichments of DMEGs were categorized into four groups: those exhibiting similar regulation in both $COLD_{chow}$ and $COLD_{HFD}$ condition pairs, and those that are uniquely regulated in each individual condition ($\Delta COLD_{diet,}$ $COLD_{chow}$, and $COLD_{HFD}$). Top ten statistically significant enrichments are shown for canonical pathways. More then ten enrichments are listed in case of p ties. Circle sizes reflect the number of genes in each enrichment while the color highlights the activation z-score. Significance levels for p after adjustment for multiple testing with Bonferroni-Holm are indicated as p <0.05 (*), p <0.01 (**).

Figure 7: Overlap of methylation regulators in cold-induced BAT activation across dietary conditions. Upset plot displays the overlap of differentially expressed methylation regulators detected for the $COLD_{chow}$, $COLD_{HFD}$, and $\Delta COLD_{diet}$ condition pairs. The scatterplot shows log2 FC values for each differentially expressed methylation regulator of the respective intersect in each condition highlighted by coloring.

Supplementary Figures

Suppl. Figure 1: Extended phenotyping of chow and HFD-fed mice housed at 30°C or 8°C.
Overview thermal images from brown adipose tissue (BAT) (A), tail surface temperature (B) and
heat loss (Ttail – Trectal; (C) in chow and HFD-fed mice housed at thermoneutrality (30°C) or in
the cold (8°C) for 7 days. Average daily calory intake (D), final fasted blood glucose (E) and
absolute organ weights (F) of inguinal white adipose tissue (iWAT), epididymal white adipose
tissue (eWAT), BAT and liver of chow and HFD-fed mice housed at thermoneutrality (30°C) or in
the cold (8°C) for 7 days. N = 7-8 per condition. Statistical significance was evaluated one-way
ANOVA with Tukey's post-hoc test (B-D) or uncorrected Fischer's LDS (E) or two-way ANOVA
with uncorrected Fischer's LDS (F). $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***).

Suppl. Figure 2: Differential gene expression in response to cold exposure in chow diet and HFD-fed mice. Summary of differential gene expression signals for cold-induced BAT activation in (A, B) normal-weight control mice ($COLD_{chow}$) and (C, D) high fat diet (HFD)-induced obese mice ($COLD_{HFD}$). Volcano plots (A, C) and heatmaps (B, D) display differentially expressed genes (DGEs) based on $\log_2$ fold change (FC) and false discovery rate (FDR) signals and show clustering of the top 25 DGEs, respectively.

Suppl. Figure 3: Overlap of DEGs in the *COLD_chow* condition with published data. Upset plots show the intersection of DEGs from the *COLD_chow* condition and DEGs reported by Tayler et al. (2024) [60] for chow-fed male mice (8°C vs 28°C), categorized by positive (up) and negative (down) regulation. DEGs with adj. P >0.05 were compared. Set size indicates the total number of DEGs.

Suppl. Figure 4: Overlap of DEGs and DMEGs between the *COLD_chow* and *COLD_HFD* conditions. Upset plots display the overlap of (A) DEGs and (B) differentially methylated and expressed genes (DMEGs) for the *COLD_chow* and *COLD_HFD* condition pairs, categorized by positive (up) and negative (down) regulation. Significant DEGs or DMEGs that show the same and contrary regulatory direction in both conditions are highlighted in red and black respectively. Those significant only in the *COLD_chow* condition are shown in blue, while those significant only in the *COLD_HFD* condition are marked in yellow. Genes are counted uniquely, even though multiple DMEGs may be associated with the same gene. Set size indicates the total number of DEGs or DMEGs.

Suppl. Figure 5: In vitro validation of DNA methylation affecting expression of candidate genes. (A) Inhibition of DNA methylation using indicated concentrations of Aza in mouse immortalized (imBA, left) and primary brown adipocytes (right). Gene expression was analyzed by qPCR, N = 6-10 per condition. (B) Induction of DNA hypermethylation using indicated concentrations of SAM in mouse imBA. Gene expression was analyzed by qPCR, n = 3-5 per condition. Statistical significance was evaluated by two-way ANOVA with uncorrected Fischer's LSD. $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***).

Suppl. Figure 6: Overlapping DMEGs between *COLD_chow* and *COLD_HFD*. The scatter plot
displays the log₂ FC in gene expression and DNA methylation signals for all DMEGs which
overlap in both *COLD_chow* and *COLD_HFD* condition. Colors correspond to highlights in Suppl.
Figure 3. DMEGs with absolute log₂ FC >0.5 in expression and >1 in methylation signals are
shown, while DMEGs in the 5% and 95% quantile of both DMP and DEG log₂ FC signals are
labeled.

Suppl. Figure 7: DMEGs that are specifically regulated within each condition. The scatter plot
displays the log$_2$ FC in gene expression and DNA methylation signals for all DMEGs which are
specifically regulated in both *COLD$_{chow}$* and *COLD$_{HFD}$* conditions. Colors correspond to highlights
in Suppl. Figure 3. DMEGs with absolute log$_2$ FC >0.5 in expression and >1 in methylation signals
are shown, while DMEGs in the 5% and 95% quantile of both DMP and DEG log$_2$ FC signals are
labeled.

Supplementary Table legends

Suppl. Table 1: Overview of primers used for PCR.

Suppl. Table 2: Overview of significant (FDR <0.05) DEGs for the *COLD$_{chow}$* contrast. DEGs that
overlap with Tayler et al. (2024) [60] (chow-fed mice, 8°C vs 28°C) are listed.

Suppl. Table 3: Overview of significant (FDR <0.05) DEGs for the *COLD$_{HFD}$* contrast.

Suppl. Table 4: Overview of significant (FDR <0.05) DEGs for the *ΔCOLD$_{diet}$* condition.

Suppl. Table 5: Overview of DMEGs in the *COLD$_{chow}$* contrast with absolute expression log$_2$ FC
>0.5 and absolute methylation log$_2$ FC >1, including the cell type assignment derived from single-
cell RNAseq data obtained from [38].

Suppl. Table 6: Overview of DMEGs in the *COLD$_{HFD}$* contrast with absolute expression log$_2$ FC
>0.5 and absolute methylation log$_2$ FC >1.

Suppl. Table 7: Overview of DMEGs in the $\Delta COLD_{diet}$ contrast with absolute expression $\log_2$ FC
>0.5 and absolute methylation $\log_2$ FC >1.

Suppl. Table 8: Significant IPA enrichment (before p adjustment) for canonical pathways
categorized by four groups: those exhibiting similar regulation in both $COLD_{HFD}$ and $COLD_{chow}$
condition pairs, and those that are uniquely regulated in each individual condition ($\Delta COLD_{diet}$,
$COLD_{chow}$, and $COLD_{HFD}$.)