CDK4/6 inhibition induces a DNA damage-independent senescence-associated
secretory phenotype driven by delayed activation of NF-κB

Joanna Lan-Hing Yeung,[1,*] Justin Rendleman,[1,*] Lauren Anderson,[1] Matthew Pressler,[1]
Nicole Pagane,[1] Irene Duba,[1] Ria Hosuru,[1] Bat-Ider Tumenbayar,[1] Andrew Koff,[2]
Viviana I. Risca[1]

[1]Laboratory of Genome Architecture and Dynamics, The Rockefeller University, New
York, NY 10065, USA.
[2]Molecular Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY
10065, USA.
*These authors contributed equally
Correspondence: vrisca@rockefeller.edu

Running title: CDK4/6 inhibition induces NF-κB-driven SASP in cancer
13

COMPETING INTERESTS
A. K. reports grants from NIH, NCI, Jennifer Linn Fund Cycle for Survival, The Maloris
Foundation, and Sarcoma Foundation of America and other support from Eli Lilly during
the conduct of the study as well as personal fees from Eli Lilly, Concarlo Therapeutics,
and PTT Global Chemical; grants from Eli Lilly and Novartis; and other support from
Atropos Therapeutics outside the submitted work. In addition, A. K. has a patent for
US9889135 issued. V. I. R. has submitted a patent application (PCT/US2024/059554)
for a chromatin conformation capture method.

ABSTRACT

Cellular senescence protects against aberrant cancer proliferation by enforcing stable cell cycle arrest. The best-studied triggers of senescence involve DNA damage signaling, while DNA damage-independent senescence is less understood. In the cancer context, senescence has recently been shown to be induced by CDK4/6 inhibitors (CDK4/6i), a class of targeted therapies currently applied to breast cancer and sarcoma, among other tumors. The senescence-associated secretory phenotype (SASP) can exert either pro- or anti-tumorigenic effects depending on its composition, making it a critical determinant of therapy outcome. The dependence of the SASP on the senescence trigger has been a controversial question, which we investigate here, particularly the involvement of NF-κB signaling. Using a multi-omics approach, we profiled the temporal dynamics of transcriptional and epigenetic state rewiring during therapy-induced senescence in CDK4/6i-treated liposarcoma cells over 28 days; we compared this to the effects of the DNA damaging chemotherapeutic, doxorubicin. Doxorubicin triggers an early and robust activation of p53 and NF-κB, as expected. Surprisingly, a delayed NF-κB-driven SASP signature also emerges in CDK4/6i-induced senescence after a 10-day exposure, in the absence of additional DNA damage. We replicated this effect in breast cancer cells over similar time courses. Inhibition of NF-κB blunts pro-inflammatory SASP expression without reversing cell cycle arrest in both treatment contexts, showing that NF-κB is a conserved SASP regulator, regardless of trigger and cancer type. Instead, the nature of the trigger regulates differential dynamics of the NF-κB response and additional SASP gene signatures, including a CDK4/6i-specific gene module driven by TEAD factors. Overall, our 28-day multi-omics time course reveals previously unknown dynamics of the SASP and demonstrates that both common and trigger-specific components of therapy-induced SASP can be separately tuned and uncoupled from cell cycle arrest, with implications for the rational design of adjuvant therapies.

INTRODUCTION

Cellular senescence protects against aberrant proliferation through the enforcement of stable cell cycle arrest in response to aging, oncogene activation, and other stressors [1]. Although senescence has historically been linked to activation of DNA damage[2, 3], it is now understood that cells can also undergo geroconversion—the transition from reversible to stable cell cycle arrest[4]—without incurring DNA damage[5, 6]. Unlike traditional chemotherapeutics, CDK4/6 inhibitors (CDK4/6i) are a class of targeted cancer therapies that halt cell cycle progression at the G1/S transition and induce senescence through a mechanism that does not appear to involve DNA damage[7, 8, 9, 10]. Whether senescence triggered by distinct mechanisms converges on a common cell state, or engages fundamentally different regulatory programs, continues to be debated. This remains challenging to resolve, in part because no universal biomarker defines the

senescent state[5, 11, 12]. While features such as increased lysosomal content and the
formation of senescence-associated heterochromatic foci (SAHFs) are common
hallmarks used to identify senescent cells[12, 13, 14], it is still unclear whether the presence
or absence of DNA damage drives fundamentally distinct senescent states.

A key regulatory feature of senescence in cancer cells is the secretion of pro-
inflammatory molecules, collectively known as the senescence-associated secretory
phenotype (SASP)[13, 15, 16, 17]. Although NF-κB is a known master transcriptional regulator
of the SASP in DNA damage-induced senescence[18, 19], its activation in CDK4/6i induced
senescence remains highly debated[8, 20, 21, 22]. While this can be attributed to differences
in dosages and model systems used, studies on CDK4/6i thus far remain focused on its
acute effects, not the late-stage senescence response. A recent study extended the
duration of CDK4/6i treatment to investigate the transition to stable arrest, identifying
the angiogenic factor, ANGPTL4 as a key regulator of this process in the liposarcoma
context[9]. However, the full breadth of transcriptional networks that regulate the late-
stage senescence response beyond this transition remain to be investigated.

The context-dependent effects of the SASP have important implications in the benefits
of therapy-induced senescence[16, 17]. The SASP's composition can vary depending on
the trigger, cell type, and treatment duration, and this heterogeneity influences its pro- or
anti-tumorigenic effects on the tumor microenvironment through modulation of immune
responses, inflammation, cellular plasticity, and epigenome remodeling[17, 23]. Hence, an
area of active interest has been to manipulate the SASP composition to achieve
therapeutically beneficial outcomes through selective inhibition of its gene modules[5, 11,
17]. Given the ongoing evaluation of CDK4/6i in for cancers such as breast cancer,
liposarcoma, non-small cell lung carcinoma, glioblastoma and colorectal cancer[24, 25, 26,
27, 28], we investigated this as a model of cancer therapy-induced senescence to
interrogate the epigenomic regulation of the SASP.

In this study, we addressed two key unresolved questions in the senescence field: (1)
How does the epigenome evolve as arrested cells progress into late-stage therapy-
induced senescence, and does this differ from DNA damage induced senescence? (2)
Does NF-κB play a role in driving the SASP when triggered by non-DNA damage
mechanisms? We hypothesized that senescence caused by different triggers would
exhibit distinct regulatory features, as blockage of CDK4/6 signaling at the G1/S
boundary should give rise to a distinct cell state when compared to arrest caused by
DNA damage (Fig. 1A), which can occur in either G1, S, or G2 and lead to the activation
of p53[3, 29]. Moreover, since CDK4/6i's main mechanism of action does not engage in
DNA damage signaling, we hypothesized that NF-κB would not be involved in shaping
the resulting SASP, consistent with prior studies that did not detect NF-κB activation in
CDK4/6i-induced senescence [21, 22].

We chose to address these questions in a liposarcoma model, a type of malignant
mesenchymal tumor characterized by the co-amplification of *CDK4* and *MDM2* on the
chromosome segment 12q13-15 in >90% of cases[30]. Traditionally, treatment is focused
on surgical resection, but recurrence rates are high for patients with dedifferentiated
histology and when tumor resides in the retroperitoneum[31, 32]. In these cases, the
disease is managed nonoperatively, however, radiation and chemotherapy exhibit
minimal efficacy for local disease[26, 30]. As such, the 10-year survival rate for these
patients is only 40.1%[33]. Over the past decade, the development of CDK4/6i has
emerged as a promising therapeutic strategy for patients with this challenging
malignancy, as a targeted means of countering the CDK4 amplification[9, 25, 34]. We
systematically compared the remodeling of both the transcriptome and chromatin
accessibility over time in response to the CDK4/6i, palbociclib, and the DNA-damaging
chemotherapeutic, doxorubicin, in a patient-derived dedifferentiated liposarcoma cell
line (Fig. 1A). Importantly, these analyses were conducted with the goal of capturing not
only the initial geroconversion entry phase, but also the cell state that arises in late-
stage senescence, specifically up to 28 days of CDK4/6i exposure.

Our results unexpectedly show that NF-κB is activated in DNA damage-independent
CDK4/6i-induced senescence, but it is not evident until cells transition beyond
quiescence into stable arrest. Although there are immediate differences upon cell cycle
exit which are maintained throughout all stages of senescence, we observe a core
SASP module driven by NF-κB signaling that is conserved across senescence triggers,
as well as a trigger-specific module for which TEAD factors are key regulators. Through
pharmacological inhibition experiments, we show that targeting NF-κB and TEAD
transcriptional networks can alter SASP profiles, without reversing cell cycle arrest,
highlighting that SASP regulation consists of potentially separable and tunable modules
apart from the impact on proliferative capacity.

RESULTS

CDK4/6i and doxorubicin both induce senescence features with distinct DNA
damage levels

To determine whether cellular hallmarks of senescence were similar in response to
CDK4/6i and doxorubicin, we profiled a variety of senescence assays over a 28-day
time course of the CDK4/6i, palbociclib (we will refer to this treatment course as Palbo),
and a 21-day time course of doxorubicin (referred to as Doxo) (Fig. 1B); we used a
slightly shorter period for the Doxo experiment as doxorubicin-induced geroconversion
occurs more rapidly. We observed an increasing number of cells with heightened
lysosomal activity after treatment with either palbociclib or doxorubicin, as measured by
positive staining for senescence-associated beta galactosidase activity (SABgal+) (Fig.

1C). In clonogenic outgrowth assays, it was apparent that less time was needed for
doxorubicin to cause a drastic reduction in the number of cells capable of forming
colonies after drug wash-out, indicating doxorubicin induces stable cell cycle arrest
earlier than palbociclib (Fig. 1D). We also observed senescence-related chromatin
changes by immunofluorescence with both treatments, consisting of increased ATRX
foci and senescence-associated heterochromatin foci (SAHF) marked by macroH2A
(Fig. 1E).[35, 36] Due to differences in the mechanisms of action for each drug (Fig. 1A),[7,
37, 38, 39] we anticipated the amount of DNA damage would be a distinctive feature
between treatments. Indeed, DNA damage in cells following palbociclib treatment was
comparable to basal levels observed in untreated cells at all timepoints, whereas the
number of γH2AX DNA damage foci and 53BP1 DNA repair foci increased rapidly upon
doxorubicin treatment and remained high over 21 days (Fig. 1F). Thus, our results
demonstrated that CDK4/6i treatment can induce senescence without the need for
excess DNA damage beyond what was detected in untreated cycling cells.

Global changes in the chromatin and transcriptional landscape define drug-
induced senescence trajectories

We next wanted to compare how transcriptional regulation changes over time between
these two distinct modes of therapy-induced senescence. To that end, we performed
RNA-seq and ATAC-seq in liposarcoma cells treated with palbociclib or doxorubicin and
sampled over early and late geroconversion, capturing the same timepoints as before
(Fig. 1B). We analyzed these two data types and mapped drug responses to determine
how the global transcriptome changed over time, relative to changes in peak
accessibility, after assessing for standard data quality metrics (Supplementary Table 1).
In both RNA-seq & ATAC-seq, sample distance measures and principal component
analysis showed that, as expected, samples grouped by drug treatment, which were
both distinct from untreated controls (Cycling) (Fig. 2A, B). Global analyses of both
transcriptome (RNA-seq) and DNA accessibility landscape (ATAC-seq) dynamics
showed subclusters representing early-, mid-, and late-phases of senescence driven by
each drug (Fig. 2A). The first two principal components (PCs), accounting for ~60-70%
of variation, were able to largely separate samples by drug and timepoint (Fig. 2B).
These overall changes corresponded to the timing of progressive geroconversion as
observed by SABgal+ cells, SAHF formation, and reduced clonogenic outgrowth (Fig. 1).

We asked how differing levels of DNA damage influences molecular changes between
these distinct types of therapy-induced senescence and identified 3,744 differentially
expressed genes across drug treatments and time points in our RNA-seq data (Fig. 2C;
Supplementary Table 2). Genes could be grouped into four expression-based clusters
that distinguish the effects of each drug, which was further broken down depending on
the dynamics of response. Assessing global patterns, we noticed that most

downregulated genes were common across the two treatments, annotated as "Shared-
down," and this cluster was enriched for genes involved in cell cycle progression. On
the other hand, SASP genes were largely split across clusters that were upregulated,
either by both drugs ("Shared-up"), or preferentially by one drug ("Palbo-upregulated"
and "Doxo-upregulated") (Fig. 2C). Taken together this suggests that, while stable arrest
is a common feature between these two forms of therapy-induced senescence, their
transcriptomic compositions have distinct outputs. We also noted that while most
differentially regulated genes were of a protein coding nature, approximately 20% were
annotated as lncRNA (Supplementary Fig. 1A).

Next, we performed differential peak analysis on our ATAC-seq data, comparing
genome-wide chromatin accessibility at each timepoint relative to Cycling and day 3-
treated samples, within each drug time course (summarized in Supplementary Fig. 1B).
Out of the 215,671 unique ATAC-seq signal peaks in our non-redundant master peak
set, 88,735 peaks (41%) underwent significant changes: a total of 49,107 unique peaks
increased in accessibility in at least one drug, while 42,164 unique peaks decreased
(Fig. 2D, E). We asked whether accessibility changes due to drug treatment were
largely specific or shared, and we found that while palbociclib and doxorubicin did have
unique sets of peaks changing in either direction, reflecting drug-distinct responses,
they also have common regions with altered accessibility, denoting a shared chromatin
component of therapy-induced senescence (Fig. 2D, E). Of note, while most ATAC-seq
peaks with increasing accessibility were localized to enhancer-like regions (i.e., distal
intergenic or intronic), those with decreasing accessibility had a stronger preference for
promoter proximal regions (Supplementary Fig. 1C, D). We observed that the contrast
between accessibility at transcription start sites (TSS) and surrounding regions dropped
toward later stages of senescence; however, our libraries maintained a TSS-to-
background accessibility ratio that passed our data quality standards (Supplementary
Fig. 1E). Interestingly, we also observed that nucleosome peaks in late-stage
geroconversion were left-shifted, consistent with a reduction in inter-nucleosome linker
lengths (Supplementary Fig. 1F). Given previous work, we expected largescale
chromatin reorganization and potential decompaction upon geroconversion[40, 41, 42]; we
found that globally, doxorubicin induced a comparable number of ATAC-seq peaks with
increasing and decreasing accessibility, while palbociclib treatment resulted in more
peaks with increased accessibility overall (Fig. 1G, H).

The shutdown of E2F target genes is more efficient with direct CDK4/6 inhibition

Given the nature of stable growth arrest in therapy-induced senescence, and the
targeted effects of CDK4/6i on Rb (Fig. 1A), we asked whether E2F transcriptional
targets were being shut down similarly between palbociclib and doxorubicin treatments.
Indeed, E2F targets were enriched amongst "Shared-down" genes (Supplementary Fig.

2A B; Supplementary Table 3) corresponding to cell cycle exit. Although both drugs led
to an eventual attenuation of E2F target genes (Supplementary Fig. 2C, D), this was
achieved at the earliest timepoints with palbociclib while it was a gradual shutdown with
doxorubicin (RNA Cluster 1 in Fig. 2C; Supplementary Fig. 2E), underscoring the direct
inhibitory effect on G1/S blockade imposed by CDK4/6i. We noticed the smaller
subcluster of genes decreased expression (RNA Cluster 2; Fig. 2C) behaved very
differently, by shutting down earlier in the Doxo time course than in Palbo. This cluster
was not enriched for cell cycle progression or E2F targets (Supplementary Fig. 2A, B),
but instead contained several genes related to cell adhesion and laminin interactions
(e.g., *NID1*, *ITGB4*, *COL4A2*). As for the downregulation of cell cycle progression genes,
the E2F shutdown was also observed at the chromatin level in our ATAC-seq data, with
E2F motif-containing peaks decreasing in accessibility in response to both palbociclib
and doxorubicin (Supplementary Fig. 2F, G). Although there was a shared set of
downregulated E2F motif-containing loci, there was also many drug-specific E2F
sequences that became less accessible (Supplementary Fig. 2F), suggesting that while
both drugs converge on inhibiting the same set of E2F target genes, they may do so
through distinct changes to the E2F binding landscape.

Signatures of enriched transcriptional regulators in therapy-induced senescence

We next focused our attention on characterizing features of upregulated peaks and
genes, as these would likely include drivers of geroconversion and not solely reflect cell
cycle exit. To understand what transcription factors may be binding in regions of newly
open chromatin, we determined enriched motifs under increasingly accessible peaks at
each timepoint in the Palbo and Doxo experiments (Supplementary Fig. 3A;
Supplementary Table 4). Since static ATAC-seq peaks often contain many of the same
motifs found in dynamic peaks, due to their roles in cell identity and maintenance, we
focused on identifying motifs with the greatest dynamic contrast. To this end, we
performed motif enrichment analysis comparing regions with increasing accessibility
against regions with decreasing accessibility at each timepoint (see Methods). In line
with what others have reported previously in senescence,[43, 44] the AP-1 family of
transcription factors was strongly enriched in response to DNA damage and increased
in enrichment over time with CDK4/6 inhibition (Supplementary Fig. 3A). Additional
motifs with similar patterns of increased accessibility were from the CEBP, STAT, TEAD,
GATA, SOX and FOX families of transcription factors, which play roles in cell fate
determination, metabolism, and inflammatory responses.[45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55] We
asked whether these transcription factor effects could also be observed in our RNA-seq
data, and to test this we looked at activation scores of upstream transcriptional
regulators predicted by Ingenuity Pathway Analysis (IPA) (Supplementary Fig. 3B;
Supplementary Table 5). In these networks, we found *JUN*, a member of the AP-1 family,
was among the top predicted transcriptional regulators across four out of five

upregulated RNA clusters, consistent with a role in general chromatin remodeling. On
the other hand, we found *STAT3* and *CEBPB* displayed stronger activation scores
among Palbo-upregulated genes, whereas *TP53* had the highest score for the Doxo
cluster with early induction after treatment. Additionally, *NFKB1* (encoding a subunit of
NF-κB) appeared in the network that was predicted to control shared upregulated genes,
as well as clusters specific to each drug (Fig. 3B), which also aligned with ATAC-based
motif enrichment results (Fig. 3A).

p53 activity in senescence is dependent on DNA damage

Performing pathway enrichment and Gene Set Enrichment Analysis (GSEA) on our
RNA-seq data, we found the p53 pathway overrepresented among genes that were
specifically induced by doxorubicin treatment (Fig. 3A; Supplementary FIg.S4A;
Supplementary Table 3). Examining genes under the "P53 PATHWAY" term that met our
significance criteria (fold change > 4 and *Padj* < 0.05), confirmed many were more
strongly upregulated in Doxo samples compared to Palbo (Fig. 3B). This preferential
doxorubicin-induced upregulation was also observed for well-established p53 targets,
such as p21 (*CDKN1A*) and *MDM2* (Fig. 3C). Moreover, the number of p53 targets in
Doxo-upregulated clusters was much higher than in Palbo-upregulated clusters (Fig. 3D,
E).

To further compare the effects of each condition on p53 target expression, we
overlapped our list of significantly changing genes with previously reported p53 targets
curated by Fischer,[56] as well as genes associated with the "P53 PATHWAY" Hallmark
term (Fig. 3F). Overall, doxorubicin treatment resulted in a stronger induction of target
expression compared to palbociclib treatment (Fig. 3F). Taken together, doxorubicin
clearly displayed a stronger activation of p53 signaling, however, a subset of p53 targets
also increased in response to palbociclib; we found that many of these genes were also
annotated as SASP or NF-κB targets, suggesting their upregulation in palbociclib-
treated cells may not be exclusively or primarily regulated by p53 (Fig. 3E).

To investigate whether a differential p53 chromatin binding signature existed between
palbociclib- and doxorubicin-induced cell cycle arrest, we analyzed Tn5 insertion site
coverage within 1-kb of p53 motifs located in ATAC-seq peaks with increased
accessibility. Among Doxo-upregulated ATAC-seq peaks, both the total number of peaks
containing p53 motifs as well as the accessibility centered on the motif was higher than
in Palbo (Fig. 3G). This trend was also observed in shared upregulated peaks, though
the difference was less pronounced; in contrast, in Palbo-upregulated peaks,
accessibility was not preferentially concentrated at p53 motifs (Fig. 3G). These findings
suggest that doxorubicin-induced cell cycle arrest is associated with a stronger p53
signature at both the gene expression and chromatin accessibility levels.

Differential regulatory pathways distinguish doxorubicin- and palbociclib-driven therapy-induced senescence

We next focused on drug-specific responses, based on our finding of Hallmark gene sets and Reactome pathways with strong preferential enrichment scores for Palbo (Supplementary Fig. 4B) or Doxo (Supplementary Fig. 4C) in the GSEA results. From the drug-specific clusters in our RNA-seq results (Fig. 2C), we used log2FC values to better visualize the distinct differences between these subclusters (Fig. 4A; Supplementary Table 2). We again performed an IPA upstream analysis, only this time directly looking for the most differential predicted regulators between drugs (Fig. 4B; Supplementary Table 5). Approaching the data in this way reiterated a p53-centered network specific to doxorubicin, but it also revealed a palbociclib-specific network involving HIF-1 and TGFβ regulation (Fig. 4A, B). Genes in this cluster included many related to the extracellular matrix (ECM) (Supplementary Fig. 4A), including *COL3A1, COL5A2, SERPINE1*, and *TIMP3*; notably these were also classified as SASP factors under our criteria, suggesting this may represent a CDK4/6i-specific SASP signature.

Given these distinct transcriptional programs, we next sought to identify differentially enriched motifs between drugs at increased accessibility regions that could be involved in drug-distinct gene expression changes. To do so, we performed motif enrichment analysis on upregulated ATAC-seq peaks that were significantly different between drugs (Fig. 4C; Supplementary Table 6). This analysis identified several Palbo-enriched motifs, including CEBPβ, TEAD4, MEF2A/B/C, SOX5 and GATA3 (Fig. 4C, left)—transcription factors that can modulate binding to mesenchymal enhancer elements and are involved in the epithelial-mesenchymal transition process,[57, 58, 59, 60, 61, 62, 63, 64] which was also reflected in our RNA-seq data (Supplementary Fig. 4A, B).

Consistent with our broader motif enrichment results (Supplementary Fig. 3A), this targeted differential analysis found the AP-1 family enriched in regions with increased accessibility specific to doxorubicin treatment (Fig. 4C, right), reinforcing AP-1's preferential role in a DNA damage-induced senescence transcriptional response. We also observed preferential enrichment of ZIC1 in Doxo-upregulated accessible regions, a factor overexpressed in liposarcomas and implicated in tumor invasiveness.[65] Taken together, these findings describe a chromatin landscape where, although many of the same transcriptional motifs were enriched across regions of increased accessibility shared between the responses to both drugs (Supplementary Fig. 3A), when we specifically look at regions that became more accessible in one condition (Fig. 4C), these regions contained binding sites for alternative transcription factors that may activate distinct modules of the senescence program and differentiate RNA outputs (Fig. 4A, B).

A shared senescence transcriptomic signature is driven by NF-κB signaling across cancer cell types

Although multiple features of therapy-induced senescence varied between doxorubicin and palbociclib treatments (e.g., DNA damage activation, p53 signaling, and a large fraction of RNA and chromatin accessibility changes), we were particularly interested in what aspects these two very different drivers of geroconversion had in common, with respect to regulatory networks and transcriptional outputs. To investigate this, we focused on shared upregulated genes (RNA Cluster 7 in Fig. 2C), which were enriched in inflammation-associated pathways (Fig. 5A). We previously noted NF-κB appeared as a mutual signature of therapy-induced senescence in both drugs (Supplementary Fig. 3A, B). Given the nature of NF-κB as a pivotal mediator of inflammatory responses,[66] we asked whether the shared transcriptional response may be NF-κB-driven. Indeed, when we looked at predicted upstream regulators of genes comprising the shared RNA response, the NF-κB complex and RELA, the p65 subunit of NF-κB, emerged as the top candidates (Fig. 5B). GSEA further supported this, showing the Hallmark pathway "TNFA SIGNALING VIA NFKB" was similarly enriched at day 14 of both treatments, relative to untreated cycling controls and day 3 within each drug time course, representing early response (Fig. 5C). Furthermore, we identified 40 NF-κB target genes within the shared cluster, half of which were annotated as SASP genes (Fig. 5D), and analyzed their RNA levels at days 3 and 14 of treatment (Fig. 5E). Both drugs showed a significant increase in expression over time, consistent with a progressive NF-κB response. However, this response was delayed in palbociclib-treated cells, with the median fold change at day 14 resembling that of doxorubicin at day 3. To assess whether delayed NF-κB activation also occurs in other cancer types treated with CDK4/6i, we performed RNA-seq on MCF7 breast cancer cells treated with 100 nM palbociclib and 10 nM fulvestrant—the standard-of-care regimen for ER+ breast cancer. A previous study did not report NF-κB activation with 5 days of CDK4/6i treatment using the MCF7 cell line[21]. Therefore, we wondered whether later timepoints would show NF-kB activation regulating the SASP in this system as well. We confirmed that CDK4/6i induces senescence in MCF7 cells without triggering excess DNA damage beyond what is observed in cycling cells, unlike doxorubicin (Fig. 5F, G). Transcriptomic analysis revealed that the 7,510 significantly altered genes based on a *Padj* threshold of < 0.05, which are part of NF-κB-related pathways, are enriched in cluster 3; a group of genes that are modestly upregulated at early time points (days 3-5) but show strong induction after day 10 (Fig. 5H, I). Thus, even in a model where NF-κB activity was previously undetected in short-term studies,[21, 22] we observed the eventual activation of NF-κB as a late treatment response. This supports the notion that NF-κB-SASP network activation is a shared feature of therapy-induced senescence across cancer types (Fig. 5; Supplementary Fig. 5).

Notably, while both drugs shared a core set of upregulated NF-κB target genes,
additional drug-specific NF-κB target genes, many of which are additional SASP factors,
were identified within drug-distinct RNA clusters (Supplementary Fig. 5B). However,
when we compiled a list of SASP genes that were induced in our context by either drug,
performing an upstream analysis with IPA predicted the Palbo-specific SASP was driven
by AGT and TGFβ, the Doxo-specific SASP was driven by IFN-γ and p53, while only the
shared SASP was likely driven by TNF and NF-κB (Supplementary Fig. 5C).

Increased chromatin accessibility at putative NF-κB binding sites occurs in both
types of therapy-induced senescence

371
372

Observing both drugs give rise to distinct effects on the chromatin landscape (Fig. 2D;
Supplementary Fig. 1G), we asked whether the shared and drug-specific NF-κB target
gene expression patterns could be explained by a model where there is a common
activation of NF-κB, however differential effects on transcription are determined by motif
accessibility. Therefore, we next investigated how the NF-κB signature was reflected in
chromatin accessibility changes between the two treatments. Motif enrichment analysis
of shared peaks with increased accessibility revealed significant enrichment of NF-κB
family motifs in both drugs (Fig. 6A). We found that among all ATAC-seq peaks that
displayed increasing accessibility in either condition (N = 49,107), there were 9,112
regions containing at least one NF-κB motif, however, in line with our model, only 1,718
of these were in mutual regions of increasing accessibility between drugs, while a
majority of NF-κB motif-containing ATAC-seq peaks with increasing accessibility were
instead drug-specific (Fig. 6B).

To determine whether changes in accessibility at putative NF-κB binding sites contribute
to the shared upregulation of NF-κB target genes, we looked specifically at regions
within 50 kb of these genes (see Methods). Tn5 insertion site coverage around the
NFKB1 motif in these nearby ATAC-seq peaks showed increased accessibility in both
treatments relative to cycling or day 3 controls (Fig. 6C). Specifically, we observed
shared increases in accessibility within the gene bodies of NF-κB target genes and
particularly around their transcription termination sites, including *IL1A, IL1B, CXCL1,*
and CXCL8 (Fig. 6D)*,* which are also part of the shared SASP. This is consistent with
their transcriptional upregulation in both drug treatments. Interestingly, we also detected
increased accessibility in distal intergenic regions near these genes (Fig. 6D); these
regions correlate with previously characterized RELA binding sites and H3K27ac-
depleted regions in untreated cells (Fig. 6D), providing evidence that NF-κB may only
gain access to these regulatory elements upon drug treatment.

Further motif enrichment analysis of genomic regions with increasing chromatin
accessibility near transcriptionally induced genes reaffirmed the consistent enrichment

of NF-κB family motifs across both drugs (Supplementary Fig. 6A; Supplementary Table
7). Notably, these motifs were highly enriched in drug-upregulated RNA clusters but
were less enriched—or absent—near shared downregulated genes (Supplementary Fig.
6B, C; Supplementary Table 7). These findings suggest that, while a proinflammatory
NF-κB-driven SASP arises early in doxorubicin treatment due to excess and persistent
DNA damage, it is gradually activated in palbociclib-treated cells following the
establishment of stable cell cycle arrest; a generalizable response despite the distinct
mechanisms through which the two drugs induce senescence.

Distinct components of the SASP, including NF-κB, can be uncoupled from stable
cell cycle arrest

To verify the shared NF-κB signatures found in the multi-omics analysis, we performed
immunofluorescence on p65 to directly measure NF-κB activation; phosphorylation of
IkB☐ by the IKK complex leads to proteasomal degradation and subsequently the
release of the p50/p65 canonical subunits that form NF-κB.[67] This results in the rapid
and transient nuclear translocation of NF-κB where it can bind to chromatin to regulate
transcription of its target genes. In both treatments, we observed p65 localization to the
nucleus with timing that correlated approximately with the timing of stable cell cycle
arrest (Fig. 7A; Supplementary Fig. 7A), i.e., steadily increasing up to day 14 in
response to palbociclib and much earlier by day 3 in doxorubicin.

After verifying that NF-κB was indeed being activated in both drugs, we attempted to
alter the NF-κB-driven shared SASP response through pharmacological inhibition with
BAY11-7082 (BAY)[68] during both treatments, after senescence was established
(Supplementary Fig. 7B). Principal component analysis shows NF-κB inhibition had
minimal impact on the overall transcriptome, as BAY- and DMSO-treated samples
clustered together within each drug group (Fig. 7B). However, BAY suppressed
expression of key shared NF-κB target SASP genes, *CXCL1, CXCL8, IL1A, IL1B,*
GMCSF (CSF2), IL6 and SOD2 (Fig. 7C; Supplementary Fig. 7C, D). Beyond these,
NF-κB inhibition downregulated 57 shared genes as well as additional drug-specific
targets (Supplementary Fig. 7E), suggesting that while NF-κB is a common feature of
senescence, its regulated may differ across treatments. Like previous studies of NF-κB
in other types of senescence[18, 69], inhibiting its activity was not enough to reverse
senescence, as indicated by no differences in clonogenic outgrowth assays (Fig. 7D).
Hence, although NF-κB activation does not appear to be necessary for maintaining
stable arrest, it is responsible for driving part of the inflammatory SASP response.

Given the growing interest in targeting specific components of the SASP, [69, 70, 71] we
asked whether additional trigger-specific branches of the SASP could be uncoupled
from cell cycle arrest, in addition to the shared NF-κB-driven program. We focused on

the TEAD, a family of transcription factors involved in the HIPPO pathway, whose motifs
were preferentially enriched near genes and in accessible regions upregulated by
palbociclib compared to doxorubicin (Fig. 4C; Supplementary Fig. 6A). Pharmacological
inhibition of TEAD with MYF-01-37[72] suppressed key palbociclib-upregulated SASP
factors, most notably *ANGPTL4* and *IGFBP3*; without disrupting stable cell cycle arrest
(Fig. 7E, F). These results identify TEAD as a previously unrecognized regulator of a
CDK4/6i-induced SASP program and demonstrate that both shared and drug-distinct
branches of the SASP can be selectively modulated.

DISCUSSION

Therapeutic response in various cancers is in part determined by how effectively tumors
are driven into stable arrest. Understanding the evolution of the corresponding SASP
that arises in different therapies will be essential in identifying new targeted approaches
that can drive more favorable outcomes.

Here we have mapped the temporal dynamics of epigenomic and transcriptomic states
that develop during therapy-induced senescence, which appear dramatically different
when triggered by a DNA-damaging agent (e.g., doxorubicin) or a DNA damage-
independent mechanism (e.g., the CDK4/6 inhibitor palbociclib). Unlike replicative
senescence that arises during aging[73], therapy-induced senescence is an acute
response to cancer treatment, exhibiting many features of canonical senescence
although mediated through distinct molecular triggers[10, 71, 74]. Our findings indicate that
most changes in both chromatin accessibility and the transcriptional landscape that
occur during prolonged cell cycle arrest are in fact drug-distinct, however there exists a
set of shared molecular responses across both drugs, which emerge at trigger-specific
rates. Our results bring new insights that help explain conflicting reports on various
features of therapy-induced senescence, including the involvement of NF-κB in
therapeutics that do not rely on DNA damage as a mechanism of action.

Furthermore, while the field has long viewed stable arrest and the SASP as tightly
linked, prior studies have challenged this assumption by showing that these two
hallmarks of senescence can be uncoupled[69, 70, 71]. Our findings contribute to this
growing, but still under-explored body of work. We were able to suppress both shared
and CDK4/6i-distinct components of the SASP while preserving cell cycle arrest, by
inhibiting NF-κB and TEAD, two key transcriptional regulators identified from our multi-
omics analysis. This suggests that selective inhibition of targeted components of the
SASP could modulate the long-term effects of treatment across different drug classes.

Although several studies have pointed to CDK4/6i-induced senescence occurring
without DNA damage [7, 8, 9, 21], some recent reports do show DNA damage is triggered
following CDK4/6i washout, amongst cells that re-enter the cell cycle[75, 76]. In our study,

we did not detect significant DNA damage following prolonged CDK4/6i treatment, even
by day 28, which we consider a late-stage senescence timepoint. Our experimental
design did not include washout, and we focused on stably arrested populations. The
lack of persistent DNA damage supports the notion that CDK4/6i triggers senescence
through distinct mechanisms from genotoxic agents like doxorubicin. This is consistent
with other observations of DNA damage-independent senescence, such as that
triggered by over expression of the endogenous CDK4/6 inhibitor, p16[5, 77]. This
difference is reflected in their divergent transcriptomic and chromatin accessibility
profiles that remain distinct over time. One of the most distinct differences between
treatments was their p53 signature, for which doxorubicin elicited a much stronger
response relative to CDK4/6i, in contrast with a recent study in CDK4/6i-treated
fibroblasts where they reported a p53-driven SASP [22].

In addition to the significant trigger-specific transcriptional responses, we unexpectedly
observed a SASP module shared by CDK4/6i- and doxorubicin-induced senescence,
albeit with distinct temporal dynamics. Both treatments induce dramatic chromatin
remodeling: reduced accessibility at promoters coinciding with decreased expression of
E2F target genes, with increased accessibility predominantly at putative enhancer
elements. NF-κB motifs stood out as being similarly enriched in both treatments, in
contrast to most other motifs that showed differential levels of enrichment. Notably, we
observed increased accessibility at putative NF-κB binding sites around shared
upregulated SASP genes, coinciding with nuclear accumulation of NF-κB and stable
arrest.

Although NF-κB's role in oncogene-induced senescence is well-established[18, 19], its
activation in CDK4/6i-induced senescence was surprising, as prior studies which
focused on early responses failed to detect NF-κB activity[21, 22]. This discrepancy likely
reflects differences in the duration of treatment. We did not observe increased NF-κB
activity in MCF7 breast cancer cells at day 5 of treatment, a similar observation to that
made by Lee *et al.* [21]. However, by increasing the duration of CDK4/6i treatment (i.e., 14,
21, 28 days), we observed delayed but robust NF-κB activation across both liposarcoma
and breast cancer cells. This reveals a previously underappreciated inflammatory
component of CDK4/6i-induced senescence, with potential implications for long-term
therapeutic outcomes.

These findings raise important questions about the temporal regulation of the SASP
across different senescence triggers. A prominent example of this is increased
accessibility at AP-1 motifs, which was shown to activate enhancers in response to
CDK4/6i in breast cancer[43]. Here, AP-1 activation emerged as a global response, with
dynamics that differed between treatments: while immediately enriched upon
doxorubicin exposure, accessible AP-1 sites gradually accumulated with palbociclib,

ultimately reaching comparable enrichment levels after a month of treatment. AP-1 is
known to act synergistically with NF-κB to drive the expression of its target genes[78, 79, 80].
Our results are consistent with cooperation between AP-1 and NF-κB to drive genome-
wide chromatin rewiring, with both factors exhibiting delayed activation coinciding with
palbociclib-induced geroconversion, offering an explanation for the timing and
magnitude differences of NF-κB-driven SASP activation across treatments.

Our results underscore two important considerations, which likely contribute to apparent
discrepancies in recent studies: 1) the importance of the underlying chromatin
landscape that can vary intrinsically by cell type and extrinsically by trigger response,
and 2) the dynamic nature of the SASP. In conclusion, we propose a model in which
both doxorubicin and CDK4/6i induce senescence through distinct molecular
mechanisms that converge over time, leading to NF-κB activation (Fig. 8A). This
promotes a shared senescence response that manifests as a common set of cytokines
and chemokines, which are secreted as part of the SASP (Fig. 8B). However, at the
same time differential chromatin accessibility, possibly through differential chromatin
remodeler activity (e.g., targeted by AP-1), allows NF-κB to act on alternative genomic
regions to give rise to drug-specific expression and temporal patterns.

Here, we show that a subset of shared SASP factors *(CXCL1, CXCL8, IL6, IL1A,* and
IL1B) can be effectively suppressed by NF-κB inhibition; many of these have known
tumor-promoting functions, including immunosuppression, metastasis, and
chemoresistance[17, 81, 82, 83, 84, 85, 86]. Beyond shared targets, CDK4/6i not only
upregulates additional NF-κB-target SASP genes compared to doxorubicin, but also
induces a distinct ECM-enriched SASP program regulated by TEAD, including
ANGPTL4, a recently identified regulator of geroconversion in liposarcoma[9]. Future
work is needed to understand the interplay and regulation between CDK4/6i-distinct and
shared components of the SASP. Nonetheless, our data suggests a paradigm shift in
the way we think about the clinical implications of CDK4/6i treatment. While it may offer
initial benefits to patients over traditional chemotherapy, over time it leads to the
emergence of a pro-inflammatory SASP driven by NF-κB that could promote tumor
progression. The deeper understanding of transcriptional and epigenetic dynamics
made possible by our extended comparative time course provides a potential solution to
this problem. We showed that the expression of both NF-κB- and TEAD-driven SASP
can be separately reduced by small molecule inhibition of these transcription factors,
without compromising cell cycle arrest. The senescence response and SASP may
therefore be tunable by adjuvant therapies to diminish pro-tumorigenic side effects.

By addressing gaps in our understanding of long-term CDK4/6i-induced senescence,
these results provide insights into its molecular regulation and effects on cell fate

decisions over time. This will be useful in furthering our understanding of patient
outcomes, especially as new CDK4 and CDK2 inhibitors enter the clinic.

DATA AND MATERIALS AVAILABILITY

All sequencing datasets have been deposited in the Gene Expression Omnibus (GEO) database. Code supporting analysis of sequencing datasets can be found at https://github.com/yeungj234/Manuscript_Figures.

Further information and request for resources and reagents should be directed to and will be fulfilled by the lead contact, Dr. Viviana Risca (vrisca@rockefeller.edu).

ATACseq: GSE295258 (Reviewer token: ojihqayqnhethgj)

RNA-seq: GSE295260 (Reviewer token: qbsxqqouzxufzub)

CUT&Tag: GSE295261 (Reviewer token: shapgsuwfnqthcp)

ACKNOWLEDGMENTS

J. L. Y. acknowledges support from a Natural Sciences and Engineering Research Council of Canada Postgraduate Scholarships – Doctoral scholarship. J. R. was supported by the American Cancer Society – Fairfield Research Council– Postdoctoral Fellowship, PF-23-1034949-01-CCB, https://doi.org/10.53354/ACS.PF-23-1034949-01-CCB.pc.gr.168157. L. J. A acknowledges support from the NSF Graduate Research Fellowship Program (1946429). This work was supported in part by a V Scholar Award (V2019-011) from the V Foundation for Cancer Research, by a Starr Cancer Consortium Grant (I16-0058), a Scholar Award from The Rita Allen Foundation (S-2108-02147), and a Career Scientist Award from the Irma T. Hirschl/Monique Weill-Caulier Trust.

We acknowledge support from the Rockefeller University Genomics Resource Center (RRID:SCR_020986), and the High-Performance Computing Resource Center (RRID:SCR_025889). We thank Andrew Scortea for administrative support and his work in setting up the Risca laboratory. We thank Wei Zhou for technical assistance. We acknowledge Novogene for providing RNA and DNA sequencing services. We thank all members of the Risca and Koff laboratories, Hironori Funabiki, Lydia Finley and Junyue Cao for insightful discussions and advice.

AUTHOR CONTRIBUTIONS

J. L. Y.: Conceptualization of project, performed experiments, data analysis, visualization and curation, manuscript writing and editing. J. R.: Performed experiments, data analysis, manuscript writing and editing, and provided mentorship. N. P.: Bioinformatics analysis. L. J. A.: Performed experiments, image analysis and manuscript editing. M. P.: Performed experiments and manuscript editing. I. D.: Bioinformatics analysis. R. H.: Bioinformatics analysis. B. T.: Performed experiments and image analysis. A. K.: Mentorship, resources, and manuscript editing. V. I. R.: Conceptualization of project, resources, project supervision, mentorship, funding acquisition, manuscript writing and editing.

MATERIALS AND METHODS:

Cell lines and cell culture

Human LS8817[TetON FLAG-MDM2] liposarcoma cell line was a gift from Dr. Andrew Koff's laboratory at Memorial Sloan Kettering Cancer Center and has been extensively characterized in the literature.[65, 87, 88, 89, 90, 91] LS8817 have previously been described using the nomenclature DDLS8817 (RRID:CVCL_M814). Cells were cultured in DMEM high glucose no glutamine (Cat# 11960044, Gibco™, Waltham, Massachusetts), supplemented with 9% Fetal Bovine Serum (Cat# 10082147, Gibco™), 1% of 100x GlutaMAX™ Supplement (Cat# 35050061, Gibco™) and 1% of 10,000U/mL Penicillin/Streptomycin (Cat# 15140122, Gibco™) and maintained at 37°C and 5% CO_2. Cells were authenticated by performing STR profiling to another LS8817 isogenic cell line.

Human MCF-7 cells (RRID:CVCL_0031) were cultured in in 75 cm2 flasks DMEM/F-12 (#Cat 10565018, Thermo Fisher Scientific, Waltham, Massachusetts) supplemented with GlutaMAX, 10% Fetal Bovine Serum (Cat# F4135, Sigma-Aldrich) and 1% Penicillin-Streptomycin at 37 °C and 5% CO2. Prior to their use, MCF7 cell identity was verified by STR profiling.

Cells were passaged at a 1:3 ratio every 3 days and regularly tested for mycoplasma.

RNA-sequencing

RNA extraction was done using the Quick-RNA Microprep Kit (Cat# R1050, Zymo Research, Irvine, California). Total RNA from at least 2 biological replicates per condition was sent to Novogene for stranded library preparation and sequencing. cDNA libraries were made with NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina (Cat# E7765, New England Biolabs, Ipswich, Massachusetts) and sequenced on Illumina NovaSeq 6000 Sequencing System as 150-base pair-end reads.

ATAC-sequencing

ATAC libraries were generated following Corces et al. protocol for OMNI-ATAC.[92] In brief, cells were treated in culture medium with DNase (Cat# LS002007, Worthington, Lakewood, New Jersey) at a final concentration of 200 U/mL at 37 °C for 30 minutes, then washed twice with PBS at 500 RCF, room temperature (RT). Approximately ~50,000-60,000 cells were used for each transposition reaction. Following permeabilization, 2.5 µL of Illumina TDE1 enzyme (Cat# 20034198, Illumina, San Diego, California) was used for transposition at 37 °C for 30 minutes. Libraries were PCR amplified for 8-13 cycles with Illumina i5 or i7 barcoded primers and were cleaned up pre- and post-amplification following the manufacturer's instructions using Zymo DNA Clean and Concentrator-5 Kit (Cat# D4014, Zymo Research). At least 2 technical replicates and 2 biological replicates per condition were sequenced. Fragment length distribution and DNA amount was quantified via TapeStation and Qubit to assess for

630 quality prior to sequencing. Libraries were sequenced on the Illumina NovaSeq 6000 or
631 on the NextSeq 1000 Sequencing System.
632

CUT&Tag sequencing
CUT&Tag on untreated cycling LS8817 cells was performed following the protocol from
Kaya-Okur et al.[93] and was previously described in Soroczynski et al.[94] Antibodies used
include: Tri-Methyl-Histone H3 (Lys27) (C36B11) Rabbit mAb (9733T, Cell Signaling
Technology, Danvers, Massachusetts), 1:50; Anti-Histone H3 (acetyl K27) antibody -
ChIP Grade (ab4729, Abcam, Cambridge, United Kingdom), 1:100; Guinea Pig anti-
Rabbit IgG (Heavy & Light Chain) antibody (ABIN101961, Antibodies-online, Aachen,
Germany), 1:100.
641

Drug treatments
Cells were passaged at a 1:3 ratio one day before drug treatment. For samples sent for
sequencing, cells were treated with either 1 µM palbociclib (Cat# S1116, Selleck
Chemicals, Houston, Texas) or 100 nM doxorubicin (Cat# S1208, Selleck Chemicals),
with media changes every 2–3 days until harvest. For NF-κB inhibition experiments, 10
µM BAY11-7082 (Cat# S2913, Selleck Chemicals) was co-administered with 1 µM
palbociclib in cells that had been treated with Palbo 3 days before harvest. For the
corresponding timepoints in Doxo, cells were treated with 100 nM doxorubicin for 24
hours, after which the drug was washed out. 10 µM BAY11-7082 was then added for 2–
3 days prior to harvesting the Doxo-treated cells. For TEAD inhibition experiments, cells
were treated with 1 µM palbociclib for 3 days before co-administration with 10 µM MYF-
01-37 (Cat# S8950, Selleck Chemicals) until harvest. Stock solutions were prepared as
followed: palbociclib (10 mM in ultrapure dH_2O, stored at -80°C), doxorubicin (1 mM in
ultrapure dH_2O, stored at -20°C), BAY11-7082 and MYF-01-37 (50 mM in DMSO, stored
at -80°C).
657

Immunofluorescence
50,000-60,000 cells were plated on acid washed glass coverslips and allowed to adhere
overnight. Cells were fixed for 15 minutes with 4% formaldehyde followed by
permeabilization with 0.1% Triton-X. Blocking was done with 0.5% Tween-20 and 1%
normal goat serum for 30 minutes-1 hour. Antibody dilutions are as follows: ATRX (Cat#
A301-045A, Bethyl, Montgomery, Texas) 1:2000, macroH2A (Cat# sc-377452, Santa
Cruz, Dallas, Texas) 1:500, γH2A.X (Cat# 05-636, Millipore, Burlington, Massachusetts)
1:1000, 53BP1 (Cat# ab172580, Abcam) 1:1000, NF-κB p65 (Cat #8242, Cell Signaling
Technology) 1:800. Primary antibody incubations were done at 4°C overnight. The next
day, secondary Alexa Fluor™ antibodies (Cat# A-11004 and A-11008, Invitrogen,
Waltham, Massachusetts) were diluted 1:500 with 1:2000 Hoechst (Cat# H3570, Fisher
Scientific, Waltham, Massachusetts) and incubated for 1 hour at RT. All reagents,

including antibodies, were diluted in 1X PBS. All steps were followed by 3 washes of 1X
PBS. Coverslips were mounted on homemade mounting medium (1 mL 10X PBS, 9 mL
Ultrapure glycerol and 50mg of N-propyl gallate (Cat# A10877-06, Alfa Aesar, Ward Hill,
Massachusetts)).
674
675 **SA-Bgal staining**
676 Senescence-associated beta galactosidase (SABgal) staining was conducted following
677 the manufacturer's protocol (Cat# 9860S, Cell Signaling Technology), with all reagents
678 diluted in 1X PBS instead of dH2O.
679
Colony formation assays
Clonogenic outgrowth was performed in a 6-well plate by plating 1600, 800, and 400
cells per well using a 2× serial dilution method. First, 6400 cells were suspended in 16
mL of media, and 4 mL was plated into each of two wells (1600 cells/well). The
remaining 8 mL of suspension was then diluted with an additional 8 mL of media (2X
dilution), and 4 mL was plated into two more wells (800 cells/well). This dilution step
was repeated once more to obtain 400 cells/well. Cells were allowed to grow for 10–12
days in drug-free media, followed by staining with crystal violet (1.25 g in 100 mL
methanol and 400 mL distilled water) for 10 minutes at room temperature. Plates were
washed with cold water, dried overnight, and scanned using an Epson Perfection V600
photo scanner.
691
RT-qPCR
Total RNA was extracted using the Quick-RNA Microprep Kit (Cat# R1050, Zymo
Research) from at least two biological replicates per condition. cDNA was synthesized
with the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit (Cat#
4368814, Thermo Fisher Scientific) and SUPERase• In™ RNase Inhibitor (Cat#
AM2696, Thermo Fisher Scientific). Prior to qPCR, cDNA was diluted to 5 ng/µL in
dH2O.
qPCR was performed in triplicate in a 96-well microplate using a 15 µL reaction
containing 7.5 µL PowerUp SYBR Green Master Mix (Cat# A25742, Thermo Fisher
Scientific), 4 µL dH2O, 0.75 µL each of forward and reverse primers, and 1.5 µL of 5
ng/µL cDNA. Reactions were run on a QuantStudio™ 5 Real-Time PCR System with
the following untreated cycling conditions: 98°C for 30 seconds, followed by 40–45
cycles of 98°C for 10 seconds, 63°C for 30 seconds, and 72°C for 1 minute. Cq values
were normalized to the housekeeping gene *PUM1*, which was selected based on its
stable expression across conditions in RNA-seq datasets.
707
<u>**QUANTIFICATION AND STATISTICAL ANALYSIS**</u>
Image quantification

All images were acquired using the Nikon Ti2E Research Microscope System where
about 100 nuclei or more were used for downstream quantification. To quantify ATRX,
macroH2A, γH2AX, and 53BP1 foci in immunofluorescent images, we used a custom
CellProfiler pipeline.[95] Foci and nuclei were identified using the **IdentifyPrimaryObjects**
module, with the **EnhanceOrSuppressFeatures** module applied beforehand to improve
foci detection. The **RelateObjects** module was then used to associate foci with their
respective nuclei.
The NF-κB nuclear-to-perinuclear intensity ratio was calculated using a CellProfiler
pipeline based on
https://www.youtube.com/watch?v=DlQSgDIGuds.
Briefly, **IdentifyPrimaryObjects** detected nuclei, followed by **ExpandOrShrinkObjects**
to expand nuclear boundaries, and **IdentifyTertiaryObjects** to define the perinuclear
region. The **MeasureObjectIntensity** module was then used to quantify intensity in
nuclear and perinuclear regions. SABgal+ associated nuclei were manually counted
using Fiji software.[96] For all quantifications, Dunnett's test was used to test for
significance between pairwise comparisons of conditions.
726
727 **RNA-sequencing pre-processing**
728 Reads containing adapters, N > 10% (N represents the base cannot be determined) and
729 low quality (Qscore<= 5) reads were filtered out. Pseudoalignment and quantification of
730 transcripts were performed with kallisto.[97]
731
Differential gene expression analysis
LS8817: Downstream analysis was performed in RStudio v3.6.3. Kallisto counts were
imported into R using tximport v1.14.2, and differential gene expression analysis was
conducted with DESeq2 1.26.0.[98] Genes with fewer than 50 total counts across all
samples were filtered out. Significantly differentially expressed genes were identified
using a likelihood ratio test ($Padj$ < 0.05) and a Wald test (absolute log2FC ≥ 2) relative
to Cycling or day 3 treatment controls.A total of 3,744 differentially expressed genes
were identified and clustered into seven RNA expression groups using k-means
clustering. To characterize expression patterns across treatments, genes were further
categorized as "Shared-down," "Palbo-upregulated," "Doxo-upregulated," or "Shared-
up." rlog-normalized gene expression values were visualized using the pheatmap R
package, with values scaled by row to enhance comparability.
MCF7: Quantification of transcripts and read alignment were performed with STAR
2.7.10a Genes with at least 1 sample with > 5 counts were kept. Differential gene expression
analysis was conducted with DESeq2[98] and significantly differentially expressed genes
were identified using a likelihood ratio test ($Padj$ < 0.05)
Identification of SASP and transcription factor target genes

SASP genes: We used the manually curated databases provided in Wang et al.,[22]
Gleason et al.,[9] and the SASPAtlas[99] (511 genes). Filtering for significantly changing
genes present in all three databases, we identified 277 SASP genes that were
associated with either therapy-induced senescence.
Cell cycle genes: Significantly changing genes were annotated as Cell Cycle if they
were found in one of the following databases (892 genes). Hallmark: "E2F TARGETS,"
"G2M_CHECKPOINT;" Reactome: "CELL_CYCLE," "CELL_CYCLE_CHECKPOINTS,"
"CELL_CYCLE_MITOTIC;" KEGG: "CELL_CYCLE."
P53 target genes: Significantly changing genes were cross-referenced with the curated
p53 target list from Fischer and the "P53 PATHWAY" Hallmark term (473 genes),
identifying 102 upregulated p53 target genes across RNA clusters.
NF-κB target genes: We used the Gilmore Lab NF-κB target gene database
(https://www.bu.edu/nf-kb/gene-resources/target-genes/), the "TNFA SIGNALING VIA
NFKB" Hallmark term, and Ingenuity Pathway Analysis (IPA) upstream regulators
containing the pattern, "NFKB" (687 genes). Filtering for significantly changing genes
present in all three sources, we identified 163 upregulated NF-κB target genes across
RNA clusters.
E2F target genes: We filtered significantly changing genes against the "E2F TARGETS"
Hallmark term and IPA upstream regulators containing the pattern, "E2F" (291 genes),
identifying 164 downregulated E2F target genes.
769
GO enrichment analysis
GO enrichment was performed using enricher function from clusterProfiler package,[100]
using gene IDs from each RNA cluster examining MSigDB databases.[101, 102] Default
settings of *P*-value cutoff = 0.05 and *P*-adjust method = "BH" were used. Background
gene set for GO enrichment was set as all genes expressed. -log10(*Padj*) values for the
top 5-10 enriched terms across all databases for each RNA cluster were plotted as a
heatmap using pheatmap R package.
777
778 **GSEA analysis**
779 Genes were ranked based on the product of log2FC*-log10(*P*-value), from DESeq2
780 results. GSEA was performed on the ranked list of genes using the GSEA function from
781 clusterProfiler package v3.14.3 using the default settings.[103] Running enrichment score
782 plots were generated using the gseaplot function with the setting "by = runningScore."
783
Ingenuity Pathway Analysis
To determine predicted upstream regulators, data were analyzed with the use of
QIAGEN IPA (QIAGEN Inc., https://digitalinsights.qiagen.com/IPA) where the core
analysis was performed on gene IDs from each RNA cluster.[104] To determine the
activation scores and visualize predicted relationships between upstream regulators,

IPA core analysis was performed on log2FC values of gene IDs across timepoints
relative to Cycling or day 3 treatment controls with a threshold set to an absolute value
of log2FC = 1.
To identify differentially activated upstream regulators and compare between drugs, an
IPA comparison was performed on the individual core analyses results, generating a
matrix of activation scores. To filter for the top 10-12 most differentially activated
upstream regulators, the sum of activation scores across timepoints within each drug
was calculated. Upstream regulators with activation scores significantly higher in one
treatment condition relative to the other were selected based on greatest magnitude
differences between drugs. Conversely, upstream regulators significantly activated in
Doxo-treated conditions but repressed in Palbo-treated conditions were selected based
on an analogous threshold.
To identify the top activated or inhibited upstream regulators for each drug treatment
overall, the sum of activation scores across multiple conditions within drug was ranked
from highest to lowest.
For all analyses, filtered upstream regulators were cross-referenced with a list of
expressed genes in this cell line to ensure biological relevance. -log10(*Padj*) values for
the top 5-10 enriched upstream regulators or activation scores for each condition were
plotted as a heatmap using pheatmap R package.
808
809 **CUT&Tag preprocessing**
810 The preprocessing of CUT&Tag data, including alignment of FASTQ into BAM files, and
811 the generation of BigWig files, was performed as previously described.[94]
812
ATAC-sequencing preprocessing
Adapters were trimmed using cutadapt and reads were aligned with bowtie2 to the hg38
genome. Following this, duplicate reads and reads mapping to the mitochondrial
genome were removed. BAM files were converted to bedpe files, which were then
converted to tagAlign files for MACS2 to call peaks on, areas of the genome enriched
for aligned sequencing reads, indicating hyper-accessibility. MACS2 parameters were `-`
`g 2.7e9 -p 0.2 --shift -75 --extsize 150 --nomodel -B --SPMR --`
`keep-dup all --call-summits`. Reproducible peaks across technical replicates
per condition were kept using Irreproducible Discovery Rate.[105] A master nonredundant
peak set was generated using a custom iterative peak filtering script which selects the
peak with the lowest q-value generated by MACS2 from a set of overlapping peaks.
Peaks were fixed to 500 bp in length, with 250 bp extending from the summit identified
by MACS2.[106] In total we identified 215,671 non-redundant peaks. Fragment counts
under peaks were counted using chromVAR's `getCounts` function.[107]
827
ATAC-seq fragment length distribution:

Biological replicates per condition were merged into a single BAM file using samtools
merge function.[108] The following line of code was used to extract the fragment length
and count the number of unique occurrences for each fragment length: `samtools`
`view $INPUT| awk -F'\t' 'function abs(x)(return ((x < 0.0) ? -x :`
`x)) (print abs($9))' | sort -n | uniq -c | awk -v OFS="\t"`
`'(print $2, $1/2)' >$OUTPUT\_insertdata.txt` where $INPUT is the merged
BAM file and $OUTPUT is the directory path and file name. The output text file
containing fragment length information was imported into R. The overall fragment length
distribution was normalized by dividing the number of each fragment length by the total
number of fragments in the merged BAM file. The resulting fragment length distribution
was plotted using ggplot2 v3.5.1 package in R as a line plot.
840

Differential ATAC-seq peak analysis
Differential peak analysis was performed using DESeq2 with the Wald test, based on
pairwise comparisons of each timepoint relative to Cycling or day 3 treatment controls.
Among the 215,671 unique peaks in our non-redundant master peak set, we identified
49,107 upregulated and 42,164 downregulated peaks across both drug treatments.
Peaks were considered significant based on a Bonferroni-corrected *Padj* threshold
(0.01/215,671 total peaks) and an absolute log2FC > 1 at any timepoint compared to
Cycling or day 3 treatment controls. Upregulated peaks were categorized as "Doxo-
upregulated" if they were significant in Doxo but not Palbo, "Palbo-upregulated" if
significant in Palbo but not Doxo, and "Shared-upregulated" if significant in both
treatments. The same criteria were applied for categorizing downregulated peaks.
To visualize read counts under significantly upregulated peaks, reads were variance-
stabilizing transformed (VST) and ordered based on their classification as Palbo-
upregulated, Shared-upregulated, or Doxo-upregulated. Hierarchical clustering was
applied within each drug category to improve the coherence of the heatmap structure.
VST-normalized counts were visualized using the pheatmap package, with values
scaled across rows to enhance comparability. The same was performed for
downregulated peaks.
859

Peak Annotation
Significantly changing peaks at each timepoint relative to Cycling or day 3 treatment
controls within each drug were merged into a single peak set. Peaks were annotated
using the `annotatePeak` function from ChIPseeker with the UCSC hg38 knownGene
TxDb.[109] For easier visualization of heatmap of reads under peaks, categories "Exon",
"Intron", "3' UTR" and "5' UTR" were merged into a single "Gene Body category" and
categories "Distal intergenic" and "Downstream" were merged into a single "Distal"
category.
868

Nearby peaks identification

Genomic coordinates of genes were obtained from the comprehensive gene annotation file in GENCODE Release 41 (GRCh38.p13).[110] For each RNA cluster of significantly changing genes, gene boundaries were extended by 50 kb upstream and downstream. Significantly changing peaks were then filtered to retain those overlapping with the extended genomic coordinates of each RNA cluster. Additionally, nearby static peaks (peaks that did not meet the significance criteria) were identified using the same approach.

Motif enrichment analysis

Motif enrichment analysis was performed using Analysis of Motif Enrichment (AME) from the MEME SUITE v5.0.2.[111] Enriched motifs were identified under peaks using the HOCOMOCO v11 database with the following command: `ame --verbose 1 --scoring avg --method fisher --hit-lo-fraction 0.25 --evalue-report-threshold 10.0 --control $CONTROL --kmer 2 $INPUT HOCOMOCOv11_core_HUMAN_mono_meme_format.meme.`
Here $CONTROL represents the FASTA file containing sequences from background peaks, and $INPUT contains sequences from foreground peaks.

For motif enrichment analysis in upregulated peaks, FASTA sequences were extracted from upregulated peaks at each time point relative to either Cycling (untreated) or day 3-treated controls within each drug treatment; these were used as the foreground peak set, with the corresponding downregulated peaks serving as the background set. The same approach while switching the foreground and background sets was used to identify motifs enriched in downregulated peaks.

To determine differentially enriched motifs between Palbo- and Doxo-upregulated peaks, Palbo-upregulated peaks were used as the foreground set, while Doxo-upregulated peaks served as the background set, and vice versa.

For motif enrichment in nearby peaks of RNA clusters, up- or downregulated nearby peaks for each drug treatment were used as the foreground set, while static nearby peaks served as the background set.

Motif enrichment results were visualized as heatmaps of -log10($Padj$) values using the pheatmap R package or as dot plots displaying %True Positive – %False Positive and -log10($Padj$) values using ggplot2 v3.5.1.

ATAC-seq coverage plots

Tn5 insertion sites were identified as fragment ends from merged BAM files of biological replicates for each condition and saved as bigWig files. This was performed in R using a custom script. In brief, the `resize` and `shift` functions in GenomicRanges were used to retain and shift the ends of reads to +4, -5 bp to account for Tn5 insertion bias. The `coverage` function was used to convert the fragment ends into a RleList object. To

normalize bigWig files, reads were counted in 500 bp bins tiled across the genome, and
the resulting count matrix was used in DESeq to calculate size factors with the
`estimateSizeFactors` function. Normalization was performed by multiplying the
coverage by a scale factor (1/DESeq size factor) before exporting the resulting object as
a bigWig.
For accessibility signal plots around motifs, motif genomic coordinates were identified
using position weight matrices from HOCOMOCO v11 and scanned across the hg38
genome with motifmatchr 1.8.0.[112] Motifs overlapping up- or downregulated peaks at
any time point for each drug category (Doxo, Shared, Palbo) were saved as BED files
and used for coverage plotting in deepTools.[113] The same approach was applied to
nearby peaks of RNA cluster peak sets. Tn5 insertion site coverage was plotted within 1
kb of motifs using deepTools with the `computeMatrix` function (`--`
`referencePoint center`) and `plotProfile` function.
For transcription start site (TSS) accessibility, the same method was used with a BED
file of genes from GENCODE Release 41 (GRCh38.p13),[110] applying `computeMatrix`
(`--referencePoint TSS`) and `plotProfile` in deepTools.
925
Genome browser visualization
Normalized bigwigs were generated from merged BAM files of biological replicates for
each condition using the `bamCoverage` function from deepTools with the following
command: `bamCoverage -b $INPUT -o $OUTPUT --binSize 5 --`
`scaleFactor $SCALE --extendReads` where $INPUT is the merged BAM file and
$OUTPUT is the resulting bigwig file. $SCALE is the same scale factor used to
generate bigWigs for fragment ends. BigWigs were then visualized on Integrative
Genomics Viewer (IGV).
Visualization of geneHancer data of putative connected enhancers linked to NF-κB
target genes was achieved by converting geneHancer v5.20 gff file to a bedpe file and
filtering for geneHancers that had at least one NF-κB target gene as the connected
gene ID.[114] Putative RelA binding sites were derived from the cis-regulatory modules
(CRMs) ReMap 2022 database.[115]
939
Sample distance and PCA
Distance measures between samples were calculated by using the `dist` function on
the transposed VST-normalized counts matrix for ATAC-seq or rlog-normalized counts
matrix for RNA-seq in R. Prior to this, technical replicates were merged into biological
replicates for ATAC-seq. The resulting distance values were plotted using pheatmap.
Principal component analysis (PCA) plots were generated using DESeq2's `plotPCA`
function.
947
948 **COMPETING INTERESTS**

REFERENCES

1. Di Micco R, Krizhanovsky V, Baker D, d'Adda di Fagagna F. Cellular senescence in
ageing: from mechanisms to therapeutic opportunities. *Nat Rev Mol Cell Biol* 2021,
22(2)**:** 75–95.

2. d'Adda di Fagagna F. Living on a break: cellular senescence as a DNA-damage
response. *Nat Rev Cancer* 2008, **8**(7)**:** 512–522.

3. Campisi J. Senescent cells, tumor suppression, and organismal aging: good citizens,
bad neighbors. *Cell* 2005, **120**(4)**:** 513–522.

4. Blagosklonny MV. Geroconversion: irreversible step to cellular senescence. *Cell*
Cycle 2014, **13**(23)**:** 3628–3635.

5. Sheekey E, Narita M. p53 in senescence - it's a marathon, not a sprint. *FEBS J* 2023,
290(5)**:** 1212–1220.

6. Torres G, Salladay-Perez IA, Dhingra A, Covarrubias AJ. Genetic origins, regulators,
and biomarkers of cellular senescence. *Trends Genet* 2024, **40**(12)**:** 1018–1031.

7. Klein ME, Kovatcheva M, Davis LE, Tap WD, Koff A. CDK4/6 Inhibitors: The
Mechanism of Action May Not Be as Simple as Once Thought. *Cancer Cell* 2018,
34(1)**:** 9–20.

8. Guan X, LaPak KM, Hennessey RC, Yu CY, Shakya R, Zhang J*, et al.* Stromal
Senescence By Prolonged CDK4/6 Inhibition Potentiates Tumor Growth. *Mol Cancer*
Res 2017, **15**(3)**:** 237–249.

9. Gleason CE, Dickson MA, Klein Dooley ME, Antonescu CR, Gularte-Merida R,
Benitez M, *et al.* Therapy-Induced Senescence Contributes to the Efficacy of
Abemaciclib in Patients with Dedifferentiated Liposarcoma. *Clin Cancer Res* 2024,
30(4)**:** 703–718.

10. Wagner V, Gil J. Senescence as a therapeutically relevant response to CDK4/6
inhibitors. *Oncogene* 2020, **39**(29)**:** 5165–5176.

11. Malaquin N, Martinez A, Rodier F. Keeping the senescence secretome under
control: Molecular reins on the senescence-associated secretory phenotype. *Exp*
Gerontol 2016, **82:** 39–49.

12. Hernandez-Segura A, Nehme J, Demaria M. Hallmarks of Cellular Senescence.
Trends Cell Biol 2018, **28**(6)**:** 436–453.

13. Wang B, Han J, Elisseeff JH, Demaria M. The senescence-associated secretory
phenotype and its physiological and pathological implications. *Nat Rev Mol Cell Biol*
2024, **25**(12)**:** 958–978.

1002

1003 14. Suryadevara V, Hudgins AD, Rajesh A, Pappalardo A, Karpova A, Dey AK, *et al.*
1004 SenNet recommendations for detecting senescent cells in different tissues. *Nat Rev*
1005 *Mol Cell Biol* 2024, **25**(12)**:** 1001–1023.

1006

1007 15. Chambers CR, Ritchie S, Pereira BA, Timpson P. Overcoming the senescence-
1008 associated secretory phenotype (SASP): a complex mechanism of resistance in the
1009 treatment of cancer. *Mol Oncol* 2021, **15**(12)**:** 3242–3255.

1010

1011 16. Faget DV, Ren Q, Stewart SA. Unmasking senescence: context-dependent
1012 effects of SASP in cancer. *Nat Rev Cancer* 2019, **19**(8)**:** 439–453.

1013

1014 17. Lau L, David G. Pro- and anti-tumorigenic functions of the senescence-
1015 associated secretory phenotype. *Expert Opin Ther Targets* 2019, **23**(12)**:** 1041–1051.

1016

1017 18. Chien Y, Scuoppo C, Wang X, Fang X, Balgley B, Bolden JE, *et al.* Control of the
1018 senescence-associated secretory phenotype by NF-kappaB promotes senescence
1019 and enhances chemosensitivity. *Genes Dev* 2011, **25**(20)**:** 2125–2136.

19.   Salminen A, Kauppinen A, Kaarniranta K. Emerging role of NF-kappaB signaling in the induction of senescence-associated secretory phenotype (SASP). *Cell Signal* 2012, **24**(4): 835–845.

20.   Goel S, DeCristo MJ, Watt AC, BrinJones H, Sceneay J, Li BB*, et al.* CDK4/6 inhibition triggers anti-tumour immunity. *Nature* 2017, **548**(7668): 471–475.

21.   Lee DH, Imran M, Choi JH, Park YJ, Kim YH, Min S*, et al.* CDK4/6 inhibitors induce breast cancer senescence with enhanced anti-tumor immunogenic properties compared with DNA-damaging agents. *Mol Oncol* 2024, **18**(1): 216–232.

22.   Wang B, Varela-Eirin M, Brandenburg SM, Hernandez-Segura A, van Vliet T, Jongbloed EM*, et al.* Pharmacological CDK4/6 inhibition reveals a p53-dependent senescent state with restricted toxicity. *EMBO J* 2022, **41**(6): e108946.

23.   Lee S, Schmitt CA. The dynamic nature of senescence in cancer. *Nat Cell Biol* 2019, **21**(1): 94–101.

24.   Patnaik A, Rosen LS, Tolaney SM, Tolcher AW, Goldman JW, Gandhi L*, et al.* Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors. *Cancer Discov* 2016, **6**(7): 740–753.

25.   Dickson MA, Schwartz GK, Keohan ML, D'Angelo SP, Gounder MM, Chi P*, et al.* Progression-Free Survival Among Patients With Well-Differentiated or Dedifferentiated Liposarcoma Treated With CDK4 Inhibitor Palbociclib: A Phase 2 Clinical Trial. *JAMA Oncol* 2016, **2**(7): 937–940.

26.   Haddox CL, Hornick JL, Roland CL, Baldini EH, Keedy VL, Riedel RF. Diagnosis and management of dedifferentiated liposarcoma: A multidisciplinary position statement. *Cancer Treat Rev* 2024, **131:** 102846.

27.   Reddy PM, Martin JM, Montero AJ. CDK 4/6 Inhibitors: Evolution and Revolution in the Management of ER+ Metastatic Breast Cancer. *JCO Oncol Pract* 2022, **18**(5): 329–330.

1056

1057   28.   Wang X, Zhao S, Xin Q, Zhang Y, Wang K, Li M. Recent progress of CDK4/6
1058   inhibitors' current practice in breast cancer. *Cancer Gene Ther* 2024, **31**(9)**:** 1283–
1059   1291.

1060

1061   29.   Toettcher J. Cell Cycle Arrest After DNA Damage. In: Dubitzky W, Wolkenhauer O,
1062   Cho K-H, Yokota H (eds). *Encyclopedia of Systems Biology*. Springer New York: New
1063   York, NY, 2013, pp 249–254.

1064

1065   30.   Crago AM, Singer S. Clinical and molecular approaches to well differentiated and
1066   dedifferentiated liposarcoma. *Curr Opin Oncol* 2011, **23**(4)**:** 373–378.

1067

1068   31.   Tan MC, Brennan MF, Kuk D, Agaram NP, Antonescu CR, Qin LX*, et al.*
1069   Histology-based Classification Predicts Pattern of Recurrence and Improves Risk
1070   Stratification in Primary Retroperitoneal Sarcoma. *Ann Surg* 2016, **263**(3)**:** 593–600.

1071

1072   32.   Jolissaint JS, Raut CP, Fairweather M. Management of Recurrent Retroperitoneal
1073   Sarcoma. *Curr Oncol* 2023, **30**(3)**:** 2761–2769.

1074

1075   33.   Knebel C, Lenze U, Pohlig F, Lenze F, Harrasser N, Suren C*, et al.* Prognostic
1076   factors and outcome of Liposarcoma patients: a retrospective evaluation over 15
1077   years. *BMC Cancer* 2017, **17**(1)**:** 410.

1078

1079   34.   Hsu JY, Seligson ND, Hays JL, Miles WO, Chen JL. Clinical Utility of CDK4/6
1080   Inhibitors in Sarcoma: Successes and Future Challenges. *JCO Precis Oncol* 2022, **6:**
1081   e2100211.

1082

1083   35.   Kovatcheva M, Liao W, Klein ME, Robine N, Geiger H, Crago AM*, et al.* ATRX is
1084   a regulator of therapy induced senescence in human cells. *Nat Commun* 2017, **8**(1)**:**
1085   386.

1086

1087   36.   Zhang R, Poustovoitov MV, Ye X, Santos HA, Chen W, Daganzo SM*, et al.*
1088   Formation of MacroH2A-containing senescence-associated heterochromatin foci and
1089   senescence driven by ASF1a and HIRA. *Dev Cell* 2005, **8**(1)**:** 19–30.

1090

1091 37. Kciuk M, Gielecińska A, Mujwar S, Kołat D, Kałuzińska-Kołat Ż, Celik I, *et al.*
1092 Doxorubicin-An Agent with Multiple Mechanisms of Anticancer Activity. *Cells* 2023,
1093 **12**(4).

1094

1095 38. Watt AC, Goel S. Cellular mechanisms underlying response and resistance to
1096 CDK4/6 inhibitors in the treatment of hormone receptor-positive breast cancer. *Breast*
1097 *Cancer Res* 2022, **24**(1)**:** 17.

1098

1099 39. Linders AN, Dias IB, López Fernández T, Tocchetti CG, Bomer N, Van der Meer P.
1100 A review of the pathophysiological mechanisms of doxorubicin-induced cardiotoxicity
1101 and aging. *NPJ Aging* 2024, **10**(1)**:** 9.

1102

1103 40. Chandra T, Ewels PA, Schoenfelder S, Furlan-Magaril M, Wingett SW, Kirschner
1104 K*, et al.* Global reorganization of the nuclear landscape in senescent cells. *Cell Rep*
1105 2015, **10**(4)**:** 471–483.

1106

1107 41. Dasgupta N, Arnold R, Equey A, Gandhi A, Adams PD. The role of the dynamic
1108 epigenetic landscape in senescence: orchestrating SASP expression. *NPJ Aging*
1109 2024, **10**(1)**:** 48.

1110

1111 42. Swanson EC, Manning B, Zhang H, Lawrence JB. Higher-order unfolding of
1112 satellite heterochromatin is a consistent and early event in cell senescence. *J Cell*
1113 *Biol* 2013, **203**(6)**:** 929–942.

1114

1115 43. Watt AC, Cejas P, DeCristo MJ, Metzger-Filho O, Lam EYN, Qiu X*, et al.* CDK4/6
1116 inhibition reprograms the breast cancer enhancer landscape by stimulating AP-1
1117 transcriptional activity. *Nat Cancer* 2021, **2**(1)**:** 34–48.

1118

1119 44. Martínez-Zamudio RI, Roux PF, de Freitas JANL, Robinson L, Doré G, Sun B*, et*
1120 *al.* AP-1 imprints a reversible transcriptional programme of senescent cells. *Nat Cell*
1121 *Biol* 2020, **22**(7)**:** 842–855.

1122

1123 45. Ren Q, Liu Z, Wu L, Yin G, Xie X, Kong W*, et al.* C/EBPβ: The structure,
1124 regulation, and its roles in inflammation-related diseases. *Biomed Pharmacother*
1125 2023, **169:** 115938.
1126

1127   46.   Awasthi N, Liongue C, Ward AC. STAT proteins: a kaleidoscope of canonical and
1128   non-canonical functions in immunity and cancer. *J Hematol Oncol* 2021, **14**(1)**:** 198.

1129
1130   47.   Sarkar A, Hochedlinger K. The sox family of transcription factors: versatile
1131   regulators of stem and progenitor cell fate. *Cell Stem Cell* 2013, **12**(1)**:** 15–30.

1132
1133   48.   Currey L, Thor S, Piper M. TEAD family transcription factors in development and
1134   disease. *Development* 2021, **148**(12).

1135
1136   49.   Huh HD, Kim DH, Jeong HS, Park HW. Regulation of TEAD Transcription Factors
1137   in Cancer Biology. *Cells* 2019, **8**(6).

1138
1139   50.   Romano O, Miccio A. GATA factor transcriptional activity: Insights from genome-
1140   wide binding profiles. *IUBMB Life* 2020, **72**(1)**:** 10–26.

1141
1142   51.   Golson ML, Kaestner KH. Fox transcription factors: from development to disease.
1143   *Development* 2016, **143**(24)**:** 4558–4570.

1144
1145   52.   Balamurugan K, Sterneck E. The many faces of C/EBPδ and their relevance for
1146   inflammation and cancer. *Int J Biol Sci* 2013, **9**(9)**:** 917–933.

1147
1148   53.   Kamachi Y, Kondoh H. Sox proteins: regulators of cell fate specification and
1149   differentiation. *Development* 2013, **140**(20)**:** 4129–4144.

1150
1151   54.   Tremblay M, Sanchez-Ferras O, Bouchard M. GATA transcription factors in
1152   development and disease. *Development* 2018, **145**(20).

1153
1154   55.   Li YJ, Zhang C, Martincuks A, Herrmann A, Yu H. STAT proteins in cancer:
1155   orchestration of metabolism. *Nat Rev Cancer* 2023, **23**(3)**:** 115–134.

1156
1157   56.   Fischer M. Census and evaluation of p53 target genes. *Oncogene* 2017, **36**(28)**:**
1158   3943–3956.

1159

1160   57.   Blokzijl A, ten Dijke P, Ibáñez CF. Physical and functional interaction between
1161   GATA-3 and Smad3 allows TGF-beta regulation of GATA target genes. *Curr Biol* 2002,
1162   **12**(1)**:** 35–45.

1163

1164   58.   Gomis RR, Alarcón C, Nadal C, Van Poznak C, Massagué J. C/EBPbeta at the
1165   core of the TGFbeta cytostatic response and its evasion in metastatic breast cancer
1166   cells. *Cancer Cell* 2006, **10**(3)**:** 203–214.

1167

1168   59.   Liu D, Kang JS, Derynck R. TGF-beta-activated Smad3 represses MEF2-
1169   dependent transcription in myogenic differentiation. *EMBO J* 2004, **23**(7)**:** 1557–1566.

1170

1171   60.   Qiao Y, Wang Z, Tan F, Chen J, Lin J, Yang J*, et al.* Enhancer Reprogramming
1172   within Pre-existing Topologically Associated Domains Promotes TGF-β-Induced EMT
1173   and Cancer Metastasis. *Mol Ther* 2020, **28**(9)**:** 2083–2095.

1174

1175   61.   Yu W, Huang C, Wang Q, Huang T, Ding Y, Ma C*, et al.* MEF2 transcription
1176   factors promotes EMT and invasiveness of hepatocellular carcinoma through TGF-β1
1177   autoregulation circuitry. *Tumour Biol* 2014, **35**(11)**:** 10943–10951.

1178

1179   62.   Kuwahara M, Yamashita M, Shinoda K, Tofukuji S, Onodera A, Shinnakasu R*, et*
1180   *al.* The transcription factor Sox4 is a downstream target of signaling by the cytokine
1181   TGF-β and suppresses T(H)2 differentiation. *Nat Immunol* 2012, **13**(8)**:** 778–786.

1182

1183   63.   Hu J, Tian J, Zhu S, Sun L, Yu J, Tian H*, et al.* Sox5 contributes to prostate
1184   cancer metastasis and is a master regulator of TGF-β-induced epithelial
1185   mesenchymal transition through controlling Twist1 expression. *Br J Cancer* 2018,
1186   **118**(1)**:** 88–97.

1187

1188   64.   Han Y, Lefebvre V. L-Sox5 and Sox6 drive expression of the aggrecan gene in
1189   cartilage by securing binding of Sox9 to a far-upstream enhancer. *Mol Cell Biol* 2008,
1190   **28**(16)**:** 4999–5013.

1191

1192   65.   Brill E, Gobble R, Angeles C, Lagos-Quintana M, Crago A, Laxa B*, et al.* ZIC1
1193   overexpression is oncogenic in liposarcoma. *Cancer Res* 2010, **70**(17)**:** 6891–6901.

1194

1195 66.   Liu T, Zhang L, Joo D, Sun SC. NF-κB signaling in inflammation. *Signal*
1196   *Transduct Target Ther* 2017, **2:** 17023–.

1197

1198 67.   Viatour P, Merville MP, Bours V, Chariot A. Phosphorylation of NF-kappaB and
1199   IkappaB proteins: implications in cancer and inflammation. *Trends Biochem Sci* 2005,
1200   **30**(1)**:** 43–52.

1201

1202 68.   Pierce JW, Schoenleber R, Jesmok G, Best J, Moore SA, Collins T*, et al.* Novel
1203   inhibitors of cytokine-induced IkappaBalpha phosphorylation and endothelial cell
1204   adhesion molecule expression show anti-inflammatory effects in vivo. *J Biol Chem*
1205   1997, **272**(34)**:** 21096–21103.

1206

1207 69.   Lau L, Porciuncula A, Yu A, Iwakura Y, David G. Uncoupling the Senescence-
1208   Associated Secretory Phenotype from Cell Cycle Exit via Interleukin-1 Inactivation
1209   Unveils Its Protumorigenic Role. *Mol Cell Biol* 2019, **39**(12).

1210

1211 70.   Birch J, Gil J. Senescence and the SASP: many therapeutic avenues. *Genes*
1212   *Dev* 2020, **34**(23-24)**:** 1565–1576.

1213

1214 71.   Schmitt CA, Wang B, Demaria M. Senescence and cancer - role and therapeutic
1215   opportunities. *Nat Rev Clin Oncol* 2022, **19**(10)**:** 619–636.

1216

1217 72.   Kurppa KJ, Liu Y, To C, Zhang T, Fan M, Vajdi A*, et al.* Treatment-Induced Tumor
1218   Dormancy through YAP-Mediated Transcriptional Reprogramming of the Apoptotic
1219   Pathway. *Cancer Cell* 2020, **37**(1)**:** 104–122.e112.

1220

1221 73.   McHugh D, Gil J. Senescence and aging: Causes, consequences, and
1222   therapeutic avenues. *J Cell Biol* 2018, **217**(1)**:** 65–77.

1223

1224 74.   Wang B, Kohli J, Demaria M. Senescent Cells in Cancer Therapy: Friends or
1225   Foes? *Trends Cancer* 2020, **6**(10)**:** 838–857.

1226

1227 75.   Crozier L, Foy R, Adib R, Kar A, Holt JA, Pareri AU*, et al.* CDK4/6 inhibitor-
1228   mediated cell overgrowth triggers osmotic and replication stress to promote
1229   senescence. *Mol Cell* 2023, **83**(22)**:** 4062–4077 e4065.

76. Crozier L, Foy R, Mouery BL, Whitaker RH, Corno A, Spanos C*, et al.* CDK4/6 inhibitors induce replication stress to cause long-term cell cycle withdrawal. *EMBO J* 2022, **41**(6)**:** e108599.

77. Rodier F, Coppe JP, Patil CK, Hoeijmakers WA, Munoz DP, Raza SR*, et al.* Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nat Cell Biol* 2009, **11**(8)**:** 973–979.

78. Fujioka S, Niu J, Schmidt C, Sclabas GM, Peng B, Uwagawa T*, et al.* NF-kappaB and AP-1 connection: mechanism of NF-kappaB-dependent regulation of AP-1 activity. *Mol Cell Biol* 2004, **24**(17)**:** 7806–7819.

79. Ji Z, He L, Regev A, Struhl K. Inflammatory regulatory network mediated by the joint action of NF-kB, STAT3, and AP-1 factors is involved in many human cancers. *Proc Natl Acad Sci U S A* 2019, **116**(19)**:** 9453–9462.

80. Oeckinghaus A, Hayden MS, Ghosh S. Crosstalk in NF-kappaB signaling pathways. *Nat Immunol* 2011, **12**(8)**:** 695–708.

81. Ancrile B, Lim KH, Counter CM. Oncogenic Ras-induced secretion of IL6 is required for tumorigenesis. *Genes Dev* 2007, **21**(14)**:** 1714–1719.

82. Acosta JC, Gil J. A role for CXCR2 in senescence, but what about in cancer? *Cancer Res* 2009, **69**(6)**:** 2167–2170.

83. Jones SA, Jenkins BJ. Recent insights into targeting the IL-6 cytokine family in inflammatory diseases and cancer. *Nat Rev Immunol* 2018, **18**(12)**:** 773–789.

84. Rebe C, Ghiringhelli F. Interleukin-1beta and Cancer. *Cancers (Basel)* 2020, **12**(7).

85. Garlanda C, Mantovani A. Interleukin-1 in tumor progression, therapy, and prevention. *Cancer Cell* 2021, **39**(8)**:** 1023–1027.

1265   86.   Wu T, Yang W, Sun A, Wei Z, Lin Q. The Role of CXC Chemokines in Cancer
1266   Progression. *Cancers (Basel)* 2022, **15**(1).

1267

1268   87.   Kovatcheva M, Liu DD, Dickson MA, Klein ME, O'Connor R, Wilder FO*, et al.*
1269   MDM2 turnover and expression of ATRX determine the choice between quiescence
1270   and senescence in response to CDK4 inhibition. *Oncotarget* 2015, **6**(10)**:** 8226–8243.

1271

1272   88.   Gobble RM, Qin LX, Brill ER, Angeles CV, Ugras S, O'Connor RB*, et al.*
1273   Expression profiling of liposarcoma yields a multigene predictor of patient outcome
1274   and identifies genes that contribute to liposarcomagenesis. *Cancer Res* 2011, **71**(7)**:**
1275   2697–2705.

1276

1277   89.   Miller ML, Molinelli EJ, Nair JS, Sheikh T, Samy R, Jing X*, et al.* Drug synergy
1278   screen and network modeling in dedifferentiated liposarcoma identifies CDK4 and
1279   IGF1R as synergistic drug targets. *Sci Signal* 2013, **6**(294)**:** ra85.

1280

1281   90.   Barretina J, Taylor BS, Banerji S, Ramos AH, Lagos-Quintana M, Decarolis PL*, et*
1282   *al.* Subtype-specific genomic alterations define new targets for soft-tissue sarcoma
1283   therapy. *Nat Genet* 2010, **42**(8)**:** 715–721.

1284

1285   91.   Singer S, Socci ND, Ambrosini G, Sambol E, Decarolis P, Wu Y*, et al.* Gene
1286   expression profiling of liposarcoma identifies distinct biological types/subtypes and
1287   potential therapeutic targets in well-differentiated and dedifferentiated liposarcoma.
1288   *Cancer Res* 2007, **67**(14)**:** 6626–6636.

1289

1290   92.   Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA,
1291   Vesuna S*, et al.* An improved ATAC-seq protocol reduces background and enables
1292   interrogation of frozen tissues. *Nat Methods* 2017, **14**(10)**:** 959–962.

1293

1294   93.   Kaya-Okur HS, Janssens DH, Henikoff JG, Ahmad K, Henikoff S. Efficient low-
1295   cost chromatin profiling with CUT&Tag. *Nat Protoc* 2020, **15**(10)**:** 3264–3283.

1296

1297   94.   Soroczynski J, Anderson LJ, Yeung JL, Rendleman JM, Oren DA, Konishi HA*, et*
1298   *al.* OpenTn5: Open-Source Resource for Robust and Scalable Tn5 Transposase
1299   Purification and Characterization. *bioRxiv* 2024.

1300

1301   95.   Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.
1302   CellProfiler 4: improvements in speed, utility and usability. *BMC Bioinformatics* 2021,
1303   **22**(1)**:** 433.

1304

1305   96.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T*, et al.*
1306   Fiji: an open-source platform for biological-image analysis. *Nat Methods* 2012, **9**(7)**:**
1307   676–682.

1308

1309   97.   Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
1310   quantification. *Nat Biotechnol* 2016, **34**(5)**:** 525–527.

1311

1312   98.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
1313   dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014, **15**(12)**:** 550.

1314

1315   99.   Basisty N, Kale A, Jeon OH, Kuehnemann C, Payne T, Rao C*, et al.* A proteomic
1316   atlas of senescence-associated secretomes for aging biomarker development. *PLoS*
1317   *Biol* 2020, **18**(1)**:** e3000599.

1318

1319   100.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing
1320   biological themes among gene clusters. *OMICS* 2012, **16**(5)**:** 284–287.

1321

1322   101.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov
1323   JP. Molecular signatures database (MSigDB) 3.0. *Bioinformatics* 2011, **27**(12)**:** 1739–
1324   1740.

1325

1326   102.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The
1327   Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst*
1328   2015, **1**(6)**:** 417–425.

1329

1330   103.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA*, et al.*
1331   Gene set enrichment analysis: a knowledge-based approach for interpreting genome-
1332   wide expression profiles. *Proc Natl Acad Sci U S A* 2005, **102**(43)**:** 15545–15550.

1333

1334   104.   Krämer A, Green J, Pollard J, Tugendreich S. Causal analysis approaches in
1335   Ingenuity Pathway Analysis. *Bioinformatics* 2014, **30**(4)**:** 523–530.

1336

1337 105. Li Q, Brown JB, Huang H, Bickel PJ. Measuring reproducibility of high-throughput
1338   experiments. *The Annals of Applied Statistics* 2011, **5**(3)**:** 1752–1779, 1728.

1339

1340 106. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE*, et al.* Model-
1341   based analysis of ChIP-Seq (MACS). *Genome Biol* 2008, **9**(9)**:** R137.

1342

1343 107. Schep AN, Wu B, Buenrostro JD, Greenleaf WJ. chromVAR: inferring
1344   transcription-factor-associated accessibility from single-cell epigenomic data. *Nat
1345   Methods* 2017, **14**(10)**:** 975–978.

1346

1347 108. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N*, et al.* The Sequence
1348   Alignment/Map format and SAMtools. *Bioinformatics* 2009, **25**(16)**:** 2078–2079.

1349

1350 109. Yu G, Wang LG, He QY. ChIPseeker: an R/Bioconductor package for ChIP peak
1351   annotation, comparison and visualization. *Bioinformatics* 2015, **31**(14)**:** 2382–2383.

1352

1353 110. Frankish A, Diekhans M, Ferreira AM, Johnson R, Jungreis I, Loveland J*, et al.*
1354   GENCODE reference annotation for the human and mouse genomes. *Nucleic Acids
1355   Res* 2019, **47**(D1)**:** D766–D773.

1356

1357 111. McLeay RC, Bailey TL. Motif Enrichment Analysis: a unified framework and an
1358   evaluation on ChIP data. *BMC Bioinformatics* 2010, **11:** 165.

1359

1360 112. Kulakovskiy IV, Vorontsov IE, Yevshin IS, Sharipov RN, Fedorova AD, Rumynskiy
1361   EI*, et al.* HOCOMOCO: towards a complete collection of transcription factor binding
1362   models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res*
1363   2018, **46**(D1)**:** D252–D259.

1364

1365 113. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS*, et al.*
1366   deepTools2: a next generation web server for deep-sequencing data analysis.
1367   *Nucleic Acids Res* 2016, **44**(W1)**:** W160–165.

1368

1369 114. Fishilevich S, Nudel R, Rappaport N, Hadar R, Plaschkes I, Iny Stein T*, et al.*
1370   GeneHancer: genome-wide integration of enhancers and target genes in GeneCards.
1371   *Database (Oxford)* 2017, **2017**.

1372
1373 115. Hammal F, de Langen P, Bergon A, Lopez F, Ballester B. ReMap 2022: a
1374 database of Human, Mouse, Drosophila and Arabidopsis regulatory regions from an
1375 integrative analysis of DNA-binding sequencing experiments. *Nucleic Acids Res* 2022,
1376 **50**(D1)**:** D316–D325.

1377

FIGURE LEGENDS

Figure 1: CDK4/6i and doxorubicin both induce senescence features with distinct
DNA damage levels

A. Diagram summarizing CDK4/6i (Palbo) vs. doxorubicin (Doxo) mechanisms of
cell cycle arrest.
B. Experimental design of drug comparison time course.
C. Left: Percent nuclei counted that had cellular SABgal+ staining at different
timepoints. Number of nuclei counted was > 300 per condition at 10x
magnification. Each point represents a field of view, with different shapes
indicating biological replicates. Error bars denote standard deviation. Statistical
significance was calculated using Dunnett's test. Right: Representative SABgal
staining images at day 3 and day 14 shown at 10x magnification with scale bar =
50 µm.
D. Crystal violet staining after clonogenic outgrowth for 10-12 days. 800 cells were
seeded per well in a 6-well plate in drug-free media after completion of drug
treatment at each timepoint.
E. Number of ATRX foci (left) and macroH2A foci (indicative of SAHFs; middle) per
nucleus was counted for ~100 nuclei per timepoint. Representative images are
shown at day 14 of each drug treatment at 60x magnification (right). Significance
was calculated using Dunnett's test. The scale bar is 15 µm.
F. Number of γH2AX foci (indicative of DNA damage; left) & 53BP1 foci (indicative
of DNA repair; middle) per nucleus was counted for ~100 nuclei per timepoint &
categorized in 0-4, 5-10 or >10 foci/nuclei. Representative images are shown at
day 14 of each drug treatment at 60x magnification (right). The scale bar is 15
µm.

Figure 2: Distinct and shared transcriptional and chromatin responses define
drug-specific senescence trajectories

A. Distance measures between samples ordered by treatment and timepoint.
B. Principal component analysis on RNA-seq and ATAC-seq samples.
C. K-means clustered heatmap of z-scores from rlog normalized counts for
significantly changing genes across treatment and timepoints (days of treatment).

| 1409 | SASP and cell cycle genes annotated in dark green and blue, respectively. |
| 1410 | Number assigned to each cluster is shown on the right side under RNA cluster. |

1411 D. Heatmap of z-scores from VST-normalized read counts under ATAC-seq peaks
1412   with significantly increased accessibility over time (relative to Cycling or day 3).
1413   Peaks are categorized as: (1) Palbo-upregulated, (2) Shared-up (upregulated in
1414   both drugs), or (3) Doxo-upregulated. Annotations indicate significance in Palbo
1415   (pink) or Doxo (green) and their genomic location (Peak annotation: distal, gene
1416   body, or promoter).

1417 E. Heatmap of z-scores from VST-normalized read counts under ATAC-seq peaks
1418   with significantly decreasing accessibility over time (relative to Cycling or day 3).
1419   Peaks are categorized as: (1) Palbo-downregulated, (2) Shared-down
1420   (downregulated in both drugs), or (3) Doxo-downregulated. Counts matrix and
1421   annotation of peaks were generated the same way as in Figure 2D.

1422 **Figure 3: p53 activity in senescence is dependent on DNA damage**

1423 A. GSEA running enrichment score for the Hallmark term "P53 PATHWAY" in day 14
1424   Doxo vs. Cycling and day 14 Palbo vs. Cycling comparisons. "P53 PATHWAY" is
1425   significantly enriched in day 14 Doxo but not day 14 Palbo.

1426 B. Log2 fold change (log2FC) values of significantly changing genes under enriched
1427   Hallmark term, "P53 PATHWAY."

1428 C. Normalized RNA-seq counts of *CDKN1A* (p21) & *MDM2*, well-known p53 targets,
1429   over time.

1430 D. Venn diagram showing the number of p53 target genes that overlap with SASP
1431   and NF-κB target genes relative to all Doxo-upregulated genes (RNA cluster 5 &
1432   6 from Figure 2C).

1433 E. Venn diagram showing the number of p53 target genes that overlap with SASP
1434   and NF-κB target genes relative to all Palbo-upregulated genes (RNA cluster 3 &
1435   4 from Figure 2C).

1436 F. Log2 fold change (log2FC) values across all significantly changing p53 target
1437   genes in day 14 Palbo and day 14 Doxo, relative to Cycling. Paired student t-test
1438   shows difference between Palbo and Doxo is significant, with *P*-value = 4.47×10^{-8}.

1439 G. Tn5 insertion sites around aggregated p53 motifs overlapping Doxo- (left),
1440   Shared- (middle), or Palbo-upregulated (right) ATAC-seq peaks relative to
1441   Cycling (peaks from Figure 2D). The number of overlapping peaks with p53
1442   motifs per condition is shown.

1443

1444 **Figure 4: Differential regulatory pathways distinguish doxorubicin and palbociclib**
1445 **driven therapy-induced senescence**

A. Log2 fold change (log2FC) values of differentially upregulated genes from RNA-seq clusters in Figure 2C, across different timepoints (days of treatment) and drug conditions, relative to Cycling or day 3 controls within each drug treatment.

B. Activation scores of the most differentially activated predicted upstream regulators between Doxo and Palbo samples, corresponding to RNA clusters in Figure 4A. Differentially activated regulators were identified by summing activation scores across timepoints within each treatment and ranking them from highest to lowest (see STAR Methods). A black outline marks the timepoint used to generate networks from IPA core analysis. Predicted relationships are shown (right) for each RNA cluster with legend at the bottom-right indicating the confidence and type of predicted relationships.

C. MEME-AME analysis of the top 7 differentially enriched motifs in upregulated ATAC-seq peaks at each timepoint, relative to Cycling or day 3 controls within each treatment. First, peaks increasing in accessibility were identified at each timepoint. Then, for each timepoint, Palbo-upregulated peaks were directly compared to Doxo-upregulated peaks (MEME-AME) to identify motifs more enriched in one condition relative to the other. Left: Motifs more enriched in Palbo-upregulated ATAC-seq peaks (foreground) compared to Doxo-upregulated peaks (background). Right: Motifs more enriched in Doxo-upregulated ATAC-seq peaks (foreground) compared to Palbo-upregulated peaks (background). Dot intensity represents -log10($Padj$) values, while dot size reflects the difference between true positive and false positive peaks (%TP - %FP).

Figure 5: A shared senescence transcriptomic signature driven by NF-κB signaling is delayed in CDK4/6i treatment across cancer cell types

A. -Log10(Padj) values of GO enrichment results for Shared-upregulated genes relative to other RNA-seq clusters in LS8817 cells.

B. -Log10(Padj) values of top 5 IPA predicted upstream regulator complexes and transcriptional regulators of Shared-upregulated genes relative to other RNA-seq clusters in LS8817 cells.

C. GSEA running enrichment score for enriched Hallmark term "TNFA SIGNALING VIA NFKB." Left: day 14 Doxo and day 14 Palbo vs. Cycling. Right: day 14 Doxo and day 14 Palbo vs. day 3 within each drug treatment in LS8817 cells. Padj values from GSEA are shown.

D. Number of NF-κB target genes overlapping with p53 and SASP target genes relative to all Shared-upregulated genes in LS8817 cells.

E. Log2 fold change (log2FC) values of NF-κB target genes that are part of the Shared-upregulated genes (RNA Cluster 7) in day 3 and day 14 Palbo and Doxo relative to Cycling in LS8817 cells. Median log2FC values for each condition are shown as a bold white line. The dotted gray line at log2FC = 0 indicates no

change from Cycling. A paired Student's t-test was used to assess significance
between log2FC distributions.

F. Representative images of SABgal staining (top) and quantification of percentage
of SABgal+ MCF7 cells over time (bottom). The scale bar is 50 µm.

G. Representative images showing different levels of DNA damage (γH2AX foci &
53BP1 foci) in doxorubicin and palbociclib treated MCF7's over time. Cells were
treated with 100nM doxorubicin or 100nM palbociclib + 10nM fulvestrant. The
scale bar is 10 µm.

H. K-means clustered heatmap of z-scores from rlog normalized counts for
significantly changing genes across treatment and timepoints (days of treatment)
based on RNA-seq performed in MCF-7 cells. SASP and cell cycle genes are
annotated in dark green and blue, respectively. NF-kB target genes found in each
upregulated gene cluster are shown left of the heatmap. Genes in dark green are
also annotated as SASP. GO enrichment results of NF-kB related terms from
each RNA-seq cluster in MCF7's.

Figure 6: Increased chromatin accessibility at putative NF-κB binding sites
occurs in both types of therapy-induced senescence

A. Left: Schematic illustrating the selection of upregulated ATAC-seq peak sets for
MEME-AME motif enrichment analysis. Right: -Log10(*Padj*) values from MEME-
AME of NF-κB family of motifs in upregulated ATAC-seq peaks relative to
downregulated peaks (based on differential accessibility relative to Cycling)
within each drug. The timepoints indicated represent the peak sets chosen as
input for MEME-AME. Dot intensity represents -log10(*Padj*) values, while dot size
reflects the difference between true positive and false positive peaks (%TP -
%FP).

B. Venn diagram showing the number of upregulated ATAC-seq peaks in each drug
category from Figure 2D that contain NF-κB family of motifs relative to all
upregulated peaks.

C. Tn5 insertion site profiles around aggregated NFKB1 motifs near shared
upregulated genes (RNA Cluster 7). ATAC-seq peaks were filtered to include only
those with increased accessibility at any timepoint in Doxo or Palbo samples
compared to Cycling or day 3 controls within each drug. The number of
overlapping peaks with NFKB1 motifs near Shared-upregulated genes per
condition is shown.

D. Genome browser visualization of ATAC-seq signal around shared upregulated
SASP and NF-κB target genes, *IL1A* & *IL1B* (top) and *CXCL1* & *CXCL8* (bottom).
Regions of increased accessibility are highlighted in shaded boxes. Putative RelA
(p65) binding sites are shown based on the ReMap database. Putative regulatory
enhancers linked to *IL1A*, *IL1B*, *CXCL1*, and *CXCL8* are represented as arc plots

using geneHancer data. CUT&Tag signal for H3K27ac (gold) and H3K27me3
(olive green) in LS8817 untreated cycling cells is also displayed.

Figure 7: Distinct components of the SASP, including NF-κB, can be uncoupled
from stable cell cycle arrest

A. Left: Representative images of nuclear localization of p65 subunit of NF-κB over
time at 60x magnification. The scale bar is 15 µm. Right: Quantification of log10
ratio of mean intensity of p65 in the nucleus to perinuclear region. Statistical
significance was calculated using Dunnett's test.
B. Schematic of pharmacological inhibition of NF-κB with BAY11-7082 (BAY) (see
STAR Methods). 10 µM BAY or equivalent volume of DMSO (negative control)
was added to cells for 2-3 days prior to harvest.
C. RT-qPCR results of palbociclib or doxorubicin treated cells with or without BAY.
Relative fold change values to Cycling are shown. Dotted grey line indicates no
change. Diamonds inside boxplot represent the mean fold change value. Circles,
triangles, and rectangles represent the fold change of individual biological
replicate. Dark purple and dark green shades indicate Palbo and Doxo samples
treated with BAY, while lighter purple and green shades indicate samples that
had DMSO added at the same time as BAY samples. Statistical significance was
calculated using a two-tailed t-test.
D. Crystal violet staining after clonogenic outgrowth after palbociclib or doxorubicin
treatment +/- BAY. 800 cells were seeded per well in a 6-well plate in drug-free
media for 12 days after harvest.
E. RT-qPCR results of palbociclib treated cells with or without TEAD inhibitor, MYF-
01-37 (MYF). Relative fold change values to Cycling are shown. Dotted grey line
indicates no change. Diamonds inside boxplot represent the mean fold change
value. Circles, triangles, and rectangles represent the fold change of individual
biological replicate. Statistical significance was calculated using a two-tailed t-test.
F. Crystal violet staining after clonogenic outgrowth after palbociclib treatment +/-
MYF. 800 cells were seeded per well in a 6-well plate and grown in drug-free
media for 12 days after harvest.

Figure 8: Summary model showing differences in how CDK4/6i and doxorubicin
may activate NF-κB and subsequently the shared SASP

A. NF-κB activation leads to the shared senescence response across treatments.
The shared SASP driven by NF-κB can be suppressed with BAY11-7082. A
component of the SASP more strongly upregulated in CDK4/6i can also be
suppressed by inhibiting TEAD activity. Differences across drugs in enriched
transcription factor (TF) motifs at upregulated ATAC-seq peaks, such as AP-1,

CEBPB, GATA3, MEF2A and TEAD4 may contribute to differences in NF-κB

regulation. DSB=double-stranded break.

B. Z-scores from rlog normalized counts of induced SASP genes common across

treatments and cancer cell types. SASP genes in red are also annotated as NF-

κB target genes.

1566

Figure 1: CDK4/6i and doxorubicin both induce senescence features with distinct DNA damage levels

A. Diagram summarizing CDK4/6i (Palbo) vs. doxorubicin (Doxo) mechanisms of cell cycle arrest.

B. Experimental design of drug comparison time course.

C. Left: Percent nuclei counted that had cellular SABgal+ staining at different timepoints. Number of nuclei counted was > 300 per condition at 10x magnification. Each point represents a field of view, with different shapes indicating biological replicates. Error bars denote standard deviation. Statistical significance was calculated using Dunnett's test. Right: Representative SABgal staining images at day 3 and day 14 shown at 10x magnification with scale bar = 50 μm.

D. Crystal violet staining after clonogenic outgrowth for 10-12 days. 800 cells were seeded per well in a 6-well plate in drug-free media after completion of drug treatment at each timepoint.

E. Number of ATRX foci (left) and macroH2A foci (indicative of SAHFs; middle) per nucleus was counted for ~100 nuclei per timepoint. Representative images are shown at day 14 of each drug treatment at 60x magnification (right). Significance was calculated using Dunnett's test. The scale bar is 15 μm.

F. Number of γH2AX foci (indicative of DNA damage; left) & 53BP1 foci (indicative of DNA repair; middle) per nucleus was counted for ~100 nuclei per timepoint & categorized in 0-4, 5-10 or >10 foci/nuclei. Representative images are shown at day 14 of each drug treatment at 60x magnification (right). The scale bar is 15 μm.

Figure 2: Distinct and shared transcriptional and chromatin responses define drug-specific senescence trajectories

A. Distance measures between samples ordered by treatment and timepoint.
B. Principal component analysis on RNA-seq and ATAC-seq samples.
C. K-means clustered heatmap of z-scores from rlog normalized counts for significantly changing genes across treatment and timepoints (days of treatment). SASP and cell cycle genes annotated in dark green and blue, respectively. Number assigned to RNA cluster is shown on the right.
D. Heatmap of z-scores from VST-normalized read counts under ATAC-seq peaks with significantly increased accessibility over time (relative to Cycling or day 3). Peaks are categorized as: (1) Palbo-upregulated, (2) Shared-up (upregulated in both drugs), or (3) Doxo-upregulated. Annotations indicate significance in Palbo (pink) or Doxo (green) and their genomic location (Peak annotation: distal, gene body, or promoter).
E. Heatmap of z-scores from VST-normalized read counts under ATAC-seq peaks with significantly decreasing accessibility over time (relative to Cycling or day 3). Peaks are categorized as: (1) Palbo-downregulated, (2) Shared-down (downregulated in both drugs), or (3) Doxo-downregulated. Counts matrix and annotation of peaks were generated the same way as upregulated ATAC-seq peaks in Figure 2D.

Figure 3: p53 activity in senescence is dependent on DNA damage

A. GSEA running enrichment score for the Hallmark term "P53 PATHWAY" in day 14 Doxo vs. Cycling and day 14 Palbo vs. Cycling comparison. "P53 PATHWAY" is significantly enriched in day 14 Doxo but not day 14 Palbo.

B. Log2 fold change (log2FC) values of significantly changing genes under enriched Hallmark term, "P53 PATHWAY."

C. Normalized RNA-seq counts of *CDKN1A* (p21) & *MDM2*, well-known p53 targets, over time.

D. Venn diagram showing the number of p53 target genes that overlap with SASP and NF-κB target genes relative to all Doxo-upregulated genes (RNA cluster 5 & 6 from Figure 2C).

E. Venn diagram showing the number of p53 target genes that overlap with SASP and NF-κB target genes relative to all Palbo-upregulated genes (RNA cluster 3 & 4 from Figure 2C).

F. Log2 fold change (log2FC) values across all significantly changing p53 target genes in day 14 Palbo and day 14 Doxo, relative to Cycling. Paired student t-test shows difference between Palbo and Doxo is significant, with *P*-value = 4.468×10^{-8}.

G. Tn5 insertion sites around aggregated p53 motifs overlapping Doxo- (left), Shared- (middle) or Palbo-upregulated (right) ATAC-seq peaks relative to Cycling (peaks from Figure 2D). The number of overlapping peaks with p53 motifs per condition is shown.

Figure 4: Differential regulatory pathways distinguish doxorubicin- and palbociclib-driven therapy-induced senescence

A. Log2 fold change (log2FC) values of differentially upregulated genes from RNA-seq clusters in Figure 2C, across different timepoints (days of treatment) and drug conditions, relative to Cycling or day 3 controls for each drug treatment.

B. Activation scores of the most differentially activated upstream regulators between Doxo and Palbo, corresponding to RNA clusters in Figure 4A. Differentially activated regulators were identified by summing activation scores across timepoints within each treatment and ranking them from highest to lowest (see Methods). A black outline marks the timepoint used to generate networks from IPA core analysis. Predicted relationships are shown (right) for each RNA cluster with legend at the bottom-right indicating the confidence and type of predicted relationships.

C. MEME-AME analysis of the top 7 differentially enriched motifs in upregulated ATAC-seq peaks at each timepoint, relative to Cycling or day 3 controls within each treatment. First, peaks increasing in accessibility were identified at each timepoint. Then, for each timepoint, Palbo-upregulated peaks were directly compared to Doxo-upregulated peaks (MEME-AME) to identify motifs more enriched in one condition relative to the other. Left: Motifs more enriched in Palbo-upregulated ATAC-seq peaks (foreground) compared to Doxo-upregulated peaks (background). Right: Motifs more enriched in Doxo-upregulated ATAC-seq peaks (foreground) compared to Palbo-upregulated peaks (background). Dot intensity represents -log10(*Padj*) values, while dot size reflects the difference between true positive and false positive peaks (%TP - %FP).

Figure 5: A shared senescence transcriptomic signature driven by NF-κB signaling is delayed in CDK4/6i treatment across cancer cell types

A. -Log10(*Padj*) values of GO enrichment results for Shared-upregulated genes relative to other RNA-seq clusters in LS8817 cells.
B. -Log10(*Padj*) values of top 5 IPA predicted upstream regulator complexes and transcriptional regulators of Shared-upregulated genes relative to other RNA-seq clusters in LS8817 cells.
C. GSEA running enrichment score for enriched Hallmark term "TNFA SIGNALING VIA NFKB." Left: day 14 Doxo and day 14 Palbo vs. Cycling. Right: day 14 Doxo and day 14 Palbo vs. day 3 within each drug treatment in LS8817 cells. *Padj* values from GSEA are shown.
D. Number of NF-κB target genes overlapping with p53 and SASP target genes relative to all Shared-upregulated genes in LS8817 cells.
E. Log2 fold change (log2FC) values of NF-κB target genes that are part of the Shared-upregulated genes (RNA Cluster 7) in day 3 and day 14 Palbo and Doxo relative to Cycling in LS8817 cells. Median log2FC values for each condition are shown as a bold white line. The dotted gray line at log2FC = 0 indicates no change from Cycling. A paired Student's t-test was used to assess significance between log2FC distributions.
F. Representative images of SABgal staining (top) and quantification of percentage of SABgal+ MCF7 cells over time (bottom). The scale bar is 50 μm.
G. Representative images showing different levels of DNA damage (γH2AX foci & 53BP1 foci) in doxorubicin and palbociclib treated MCF7's over time. Cells were treated with 100nM doxorubicin or 100nM palbociclib + 10nM fulvestrant. The scale bar is 10 μm.
H. K-means clustered heatmap of z-scores from rlog normalized counts for significantly changing genes across treatment and timepoints (days of treatment) based on RNA-seq performed in MCF-7 cells. SASP and cell cycle genes are annotated in dark green and blue, respectively. NF-kB target genes found in each upregulated gene cluster are shown left of the heatmap. Genes in dark green are also annotated as SASP.
I. GO enrichment results of NF-kB related terms from each RNA-seq cluster in MCF7's.

Figure 6: Increased chromatin accessibility at putative NF-kB binding sites occur in both types of therapy induced senescence

A. Left: Schematic illustrating the selection of upregulated ATAC-seq peak sets for MEME-AME motif enrichment analysis. Right: -Log10(Padj) values from MEME-AME of NF-κB family of motifs in up-regulated ATAC-seq peaks relative to down-regulated peaks (based on differential accessibility relative to Cycling) within each drug. The timepoints indicated represent the peak sets chosen as input for MEME-AME. Dot intensity represents -log10(p-adj) values, while dot size reflects the difference between true positive and false positive peaks (%TP - %FP).

B. Venn diagram showing the number of upregulated ATAC-seq peaks in each drug category from Fig 2D that contain NF-κB family of motifs relative to all upregulated peaks.

C. Tn5 insertion site profiles around aggregated NFKB1 motifs near shared upregulated genes (RNA Cluster 7). ATAC-seq peaks were filtered to include only those with increased accessibility at any timepoint in response to Doxo or Palbo compared to Cycling or day 3 controls within each drug. The number of overlapping peaks with NFKB1 motifs near Shared-upregulated genes per condition is shown.

D. Genome browser visualization of ATAC-seq signal around shared upregulated SASP and NF-κB target genes, IL1A & IL1B (top) and CXCL1 & CXCL8 (bottom). Regions of increased accessibility are highlighted in shaded boxes. Putative RelA (p65) binding sites are shown based on the ReMap database. Putative regulatory enhancers linked to IL1A and IL1B are represented as arc plots using geneHancer data. CUT&Tag signal for H3K27ac (gold) and H3K27me3 (olive green) in LS8817 untreated cycling cells is also displayed.

Figure 7: Distinct components of the SASP, including NF-κB, can be uncoupled from stable cell cycle arrest

A. Representative images of nuclear localization of p65 subunit of NF-κB over time at 60x magnification. The scale bar is 15 μm.
B. Principal component analysis on RNA-seq of day14 palbociclib or doxorubicin treated cells with or without NF-κB inhibitor, BAY11-7082 (BAY).
C. RT-qPCR results of palbociclib or doxorubicin treated cells with or without BAY. Relative fold change values to Cycling are shown. Dotted grey line indicates no change. Diamonds inside boxplot represent the mean fold change value. Circles, triangles, and rectangles represent the fold change of individual biological replicate. Dark purple and dark green shades indicate Palbo and Doxo samples treated with BAY while lighter purple and green shades indicate samples that had DMSO added at the same time as BAY samples. Statistical significance was calculated using a two-tailed t-test.
D. Crystal violet staining after clonogenic outgrowth after palbociclib or doxorubicin treatment +/- BAY. 800 cells were seeded per well in a 6-well plate and grown in drug-free media for 12 days after harvest.
E. RT-qPCR results of palbociclib treated cells with or without TEAD inhibitor, MYF-01-37 (MYF). Relative fold change values to Cycling are shown. Dotted grey line indicates no change. Diamonds inside boxplot represent the mean fold change value. Circles, triangles, and rectangles represent the fold change of individual biological replicate. Statistical significance was calculated using a two-tailed t-test.
F. Crystal violet staining after clonogenic outgrowth after palbociclib treatment +/- MYF. 800 cells were seeded per well in a 6-well plate and grown in drug-free media for 12 days after harvest.

A

B

Figure 8: Summary model showing differences in how CDK4/6i and doxorubicin may activate NF-κB and subsequently the shared SASP.

A. NF-κB activation leads to the shared senescence response across treatments. The shared SASP driven by NF-κB can be suppressed with BAY11-7082. A component of the SASP more strongly upregulated in CDK4/6i can also be suppressed by inhibiting TEAD activity. Differences across drugs in enriched transcription factor (TF) motifs at upregulated ATAC-seq peaks, such as AP-1, CEBPB, GATA3, MEF2A and TEAD4 may contribute to differences in NF-κB regulation. DSB=double-stranded break.

B. Z-scores from rlog normalized counts of induced SASP genes common across treatments and cancer cell types. SASP genes in red are also annotated as NF-κB target genes.