Direct contact between iPSC-derived macrophages and hepatocytes drives reciprocal acquisition of Kupffer cell identity and hepatocyte maturation

Christopher Zhe Wei Lee[1,2,3 #], Farah Tasnim[1,4 #], Xiaozhong Huang[1,4,5], Raman Sethi[6], Yoohyun Song[4], Tatsuya Kozaki[2], Sebastiaan De Schepper[2], Nicholas Ang[2], Ivy Low[2], You Yi Hwang[2], Jinmiao Chen[6, 7], Hanry Yu[1,4,5,8,9,10 *], and Florent Ginhoux[1,2,3,11,12 *]

1. Innovations in Food & Chemical Safety Programme (IFCS), A*STAR

2. Singapore Immunology Network, 8A Biomedical Grove, Singapore 138648, Singapore

3. School of Biological Sciences, Nanyang Technological University, Singapore 637551, Singapore

4. Institute of Bioengineering and Bioimaging, #05-01, 31 Biopolis Way, The Nanos, Singapore 138669, Singapore

5. Department of Physiology, Yong Loo Lin School of Medicine, National University of Singapore, MD9-04-07, 2 Medical Drive, Singapore 117593, Singapore

6. Bioinformatics Institute (BII), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore

7. Institute of Molecular and Cell Biology (IMCB), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore

8. NUS Graduate School – Integrative Sciences and Engineering Programme (ISEP), University Hall, Tan Chin Tuan Wing Level 5, #05-03, 21 Lower Kent Ridge Road, Singapore 119077, Singapore

9. Mechanobiology Institute, T-Labs, #05-01, 5A Engineering Drive 1, Singapore 117411, Singapore

10. CAMP IRG, Singapore-MIT Alliance for Research and Technology, 1 CREATE Way, Enterprise Wing, Level 4, Singapore, 138602, Singapore

11. Shanghai Institute of Immunology, Shanghai JiaoTong University School of Medicine, Shanghai, China

12. Translational Immunology Institute, SingHealth Duke-NUS Academic Medical Centre,

Singapore, Singapore

# These authors contributed equally

* Corresponding authors

Hanry Yu: hyu@a-star.edu.sg Tel: +6568247103, Fax: +68749526

Florent Ginhoux: Florent_Ginhoux@a-star.edu.sg Tel: +6564070410, Fax: +6564642056

Key words: immune-mediated hepatotoxicity, *in vitro* liver models, co-culture,
macrophages, Kupffer cells, induced pluripotent stem cells, iPSC, iMac

39   Abstract

As the resident tissue macrophage of the liver, Kupffer Cells (KC) play an important role in
homeostasis and tissue support. However, current in-vitro liver models often ignore the
contribution of these KCs toward the proper response and function of the tissue. This is
especially relevant when we consider the implications of immune-mediated drug injuries. To
address this issue, we developed an isogenic co-culture system utilizing iPSC-derived
macrophages (iMacs) and hepatocytes (iHeps). Directly co-culturing iHeps with iMacs
improved the differentiation and maturation of the iHeps, with significant downregulation of
fetal hepatocyte markers as well as upregulation of cytochrome genes. Furthermore, the co-
culture also imparted stronger KC identity to the iMacs in a contact-dependent manner, with
iMacs cultured in iHep conditioned media alone showing weaker expression of key KC
markers. Finally, challenging the iHep-iMac co-culture system with 7 paradigm hepatotoxic
compounds showed dose-dependent cytokine response in the 5 compounds associated with
immune-mediated liver injuries, while no significant changes were observed in the 2
compounds with no reported immune-dependent complications. This effect was also not
recapitulated when the co-culture was instead performed with human peripheral blood
monocyte-derived macrophages, suggesting that iMacs are essential for liver toxicity
response. Taken together, our study shows not only the importance of macrophages in tissue
systems, but also that the source of macrophages is critical to the development of accurate in-
vitro human models.

60   1. Introduction

Hepatic macrophages play key roles in immune-mediated hepatotoxicity, liver injury and
repair [1, 2]. They are impacted by autocrine and paracrine signals from hepatic cells,
especially hepatocytes. They also release soluble stress signals in response to external stimuli
from foreign particles, leading to macrophage activation and production of cell signalling and
stress pathway modulators, including reactive oxygen species and cytokines [3, 4]. In return,
hepatic macrophage activation also regulates parenchymal and non-parenchymal cell death
and the metabolic activity of hepatocytes [5]. Such crosstalk between liver parenchymal cells

(hepatocytes) and non-parenchymal modulators (hepatic macrophages) is critical in
modelling liver injury, especially immune-mediated drug-induced responses [6].

Two major populations of hepatic macrophages have been reported: liver-resident Kupffer
cells (KCs) and monocyte-derived macrophages (MoMφs) that infiltrate the liver upon insult
[3]. While MoMφs have been well studied *in vivo* and applied in *in vitro* models [6], in more
recent years it has been revealed that KCs are derived from embryonic macrophages that seed
the liver early on during development [7][8], and are only minimally replaced by circulating
monocytes under steady state conditions [9]. This also implies that human peripheral blood
monocytes, the conventional source of primary human macrophages, might not be able to
fully recapitulate KC biology. However, it has been demonstrated that murine iPSC-derived
macrophages (iMacs) more closely resembled embryonic macrophages than monocyte-
derived macrophages [10], suggesting that human iMacs might also serve as a suitable
analogue for human embryonic macrophages. Based on these new insights, in our previous
work, we differentiated iMacs into iPSC-derived KCs (iKCs) using primary hepatocyte
conditioned media (PHCM) to drive iMacs towards iKC identity [11]. Although these iKCs
partially resembled primary human KCs (PhKCs) based on the expression of KC specific
markers and drug response, PHCM is not readily available and might exhibit differences
depending on donor variability of primary hepatocytes. Most importantly, the direct cellular
interactions between iMacs and hepatocytes and their potential to impart better reciprocal
identity and functionality to each other has not been studied before.

We recently generated macrophage-sufficient brain organoids by coculturing brain organoids
with iMacs generated from the same human iPSC line [12]. In such cocultures, iMac
differentiated into cells with microglia-like phenotypes and functions. Most importantly,
iMac modulated neuronal progenitor cell (NPC) differentiation, limiting NPC proliferation
and promoting axonogenesis, profoundly remodelling organoid physiology. Thus, in this
study, we hypothesized that iMacs could be directly cultured with iPSC-derived hepatocytes
(iHeps) from the same IPSC source to both improve iHep development and maturation as
well as impart KC-like identify to the iMacs. Upon development of such a model, we tested
its application in detecting immune-mediated responses of seven paradigm compounds.
Concurrently, we also cultured iMacs in PHCM for 7 days in a similar way to our previous
study [11] , comparing the transcriptomic effect of direct contact against soluble factors in the
context of hepatic macrophage specialisation. Our results showed improved *in vivo*
correlation based on the effects on cytokine production when iMac-derived KCs were used.
Altogether, this study highlights the effect of direct crosstalk between hepatocytes and
macrophages *in vitro* using iPSCs and provides a new human model to test drug-induced
cytokine responses.

2. Materials and Methods

106
107
2.1 Maintenance of iPSCs

Human IMR90-iPSCs were obtained from WiCell Research Institute (Madison, WI). The
cells were maintained under antibiotic-free conditions in mTESR media (StemCell
Technologies, 85850) on Matrigel (Corning, 354234)-coated plates. Cells were passaged

using ReLESR (StemCell Technologies, 05872) or Dispase (StemCell Technologies, 07913)
as per manufacturer's protocol whenever they reached 80% confluency.

2.2 Differentiation of iPSCs to iHeps

IMR90-iPSCs were differentiated into iPSC-hepatocytes (iHeps) as previously described
[13]. Briefly, the iPSCs were cultured to 80% confluency, then made into single cells using
Cell Dissociation Buffer (Gibco, 13151014) and seeded at 1.5 x 10^5 cells/cm^2 on the desired
well plate format (i.e. 1.5 x 10^6 cells for a 6 well plate) and placed under hypoxic conditions
(5% O_2 and 5% CO_2). Growth factors and media used starting from the next day to day 20
are detailed in [11]. Media was changed every day and hypoxic conditions were used during
days 5 to15 of differentiation. After 20 days of differentiation, iPSC-derived macrophages
(iMacs) were either directly added to the iHeps as described in section 2.5 or iHeps were
harvested for seeding into smaller well formats for drug testing. The harvesting protocol has
been previously optimized by us and has been detailed in [14].

2.3 Differentiation of iPSCs to iMacs

IMR90-iPSCs were differentiated into iMacs as previously described [10]. Briefly, the iPSCs
were cultured to 80% confluency, then passaged using ReLESR. As ReLESR releases the
cells as small cell clumps, the passage conditions were optimised to result in roughly 1 x
10^5 cells in each well of a 6 well plate (Day -1). Starting from the next day (Day 0) to day
16, the cells were cultured in Stempro Media, consisting of Stempro-34 SFM (GIBCO,
10639-011), supplemented with 200 ug/mL Human Transferrin (Roche, 10-652-202-001), 1x
L-Glut, 1x Pen/Strep, 0.5 mM Ascorbic Acid (Sigma, A4544) and 0.45mM MTG (Sigma,
M6145). A full media change was done every other day for the next 16 days, supplemented
with the following cytokines: Differentiation Day 0 (5 ng/mL BMP4, 50 ng/mL VEGF, and 2
uM CHIR99021), Differentiation Day 2 (5 ng/mL BMP4, 50 ng/mL VEGF, and 20 ng/mL
FGF2), Differentiation Day 4 (15 ng/mL VEGF and 5 ng/mL FGF2), Differentiation Day 6 to
10 (10 ng/mL VEGF, 10 ng/mL FGF2, 50 ng/mL SCF, 30 ng/mL DKK-1 (RnD, 5439-DK),
10 ng/mL IL-6 (RnD, 206-IL), and 20 ng/mL IL-3), Differentiation Day 12 and 14 (10 ng/mL
FGF2, 50 ng/mL SCF, 10 ng/mL IL-6, and 20 ng/mL IL-3). From day 16, the cells were
cultured in 75% IMDM with Glutamax (GIBCO, 31980-030), 25% F12 (GIBCO, 11765-
047), 1x N2 supplement (GIBCO, 17502-048), 1x B27 Supplement (GIBCO, 17504-001),
0.05% BSA (GE Healthcare, SH30574) and 1x Pen/Strep, supplemented with 50ng/ml of
CSF-1 (RnD, 216-MC), with a full media change every 3 days. In addition, for the first 8
days of differentiation, the cells were kept in a hypoxic incubator (5% O_2 and 5% CO_2),
before being transferred to a standard cell culture incubator for the next 18 days. After
transfer to a standard cell culture incubator, floating cells were collected every media change
and resuspended back into the culture to retain the hematopoietic progenitors. After 26 days,
the floating cells were collected, and the purity of the iMacs was determined by flow
cytometry.

2.4 Differentiation of human peripheral blood monocytes (PBMCs) into monocyte-derived macrophages

Human PBMCs were purified from blood apheresis cones by ficoll gradient (Cytiva, 17-1440-02), before being magnetically sorted with CD14+ microbeads (Miltenyi, 130-050-021). 1 x 10^6 CD14+ cells were then seeded per well of a 6 well plate in RPMI 1640 (Hyclone, SH30027.01) with 10% FBS (Biowest, S1810-500), 1x Pen/Strep, 1x Sodium Pyruvate, 1x NEAA and 1x L-Glut with 50ng/ml CSF-1 and cultured for 7 days. The purity of the monocyte-derived macrophages was then determined by flow cytometry.

2.5 Co-culture of iMacs and monocyte-derived macrophages with iHeps

2 x 10^5 iMacs or monocyte-derived macrophages were added to 5 x 10^5 iHeps during a full media change (4ml) per well of a 6 well plate on Day 0. This media was comprised of Williams' Medium E containing the following supplements: Penicillin/Streptomycin (10,000 U/mL / (10,000 μg/mL) – final conc = 1%, ITS+ - final conc= 2%, 4mM glutamax, 30mM HEPES buffer. The concentrations of the supplements were double that of standard conditions used due to the infrequent media change (please see section 3.1 for more details). In addition, 50ng/ml of CSF-1 was added to each well. The cells were cultured for 7 days, with a half-media change (2ml) on Day 4 consisting of the same media as described above with 100ng/ml of CSF-1, resulting in a final concentration of 50ng/ml CSF-1 in the well. The cells were collected by digesting with Accutase (StemCell, 07920) for flow cytometry, RNAseq analysis and qPCR. For pharmacological testing, the number of iHeps and iMacs were downscaled to 96-well plates while maintaining the same iHep:iMac ratio (2.5:1).

2.6 Flow cytometry

Standard staining procedures were used to prepare the cells for flow cytometry analysis. Briefly, cells were dissociated into single cells using the different methods described above for the various tissues, before being incubated with 100ul of antibody containing FACS buffer (1% BSA and 4mM EDTA in PBS) per 5 million cells for 20 minutes at 4°C. The cell suspension was then washed with 5ml of FACS buffer, centrifuged at 400g, and the supernatant was removed. The cells were finally resuspended in PBS containing 3uM DAPI (Invitrogen, D1306). Data was acquired by LSRII (BD Bioscience) and analyzed by Flow Jo (Tree Star, Inc.). For cell sorting, cells were sorted using FACS Aria II (BD Bioscience) or FACS Aria III (BD Bioscience). The following antibodies were used: FITC-conjugated anti-human CD14 (Biolegend, 325604), APC/Cy7-conjugated anti-human CD45 (Biolegend, 368516), PE-conjugated anti-human CD163 (R&D Systems, FAB1607P-100), Alexa Fluor 647-conjugated anti-human CX3CR1 (Biolegend, 341608, PE/Cy7-conjugated anti-human CD11b (eBioscience, 25-0118-42).

2.7 Bulk RNA-seq

Total RNA was extracted using Arcturus PicoPure RNA Isolation kit (Thermofisher, KIT0204) according to manufacturer's protocol. All RNAs were analyzed on Labchip GX

system (Perkin Elmer, United States) for quality assessment with median RNA Quality Score
of 9.15. cDNA libraries were prepared using 2ng of total RNA and 1ul of a 1:50,000 dilution
of Ambion ERCC RNA Spike-in Controls (Thermofisher, 4456740) using the SMARTSeq
v2 protocol described [15] with the following modifications: 1. Use of 20mM TSO, 2. Use of
250pg of cDNA with 1/5 reaction of Nextera XT kit (Illumina, FC-131). The length
distribution of the cDNA libraries was monitored using DNA High Sensitivity Reagent Kit
(Perkin Elmer, CLS60672) on the Labchip GX system (Perkin Elmer). All samples were
subjected to an indexed PE sequencing run of 2x51 cycles on HiSeq 2000 (Illumina) at 16
samples per lane.

2.8 Bulk RNA-seq analysis

Paired-end raw reads were mapped to GRCh38 human genome build using STAR aligner
[16]. Genes were counted for reads that were mapped to genes using featureCounts [17]
based on GENCODEv29 gene annotation [18]. Log2 transformed counts per million mapped
read (log2CPM) and log2 transformed reads per kilobase per million mapped reads
(log2RPKM) were computed using edgeR Bioconductor package [19]. Across all samples,
genes with log2CPM inter-quartile range (IQR) less than 0.5 were filtered out from
subsequent differential expression gene (DEG) analysis. DEG analyses for culture condition
comparisons were all done using edgeR. Selection of DEGs was done with Benjamini-
Hochberg [20] which adjusted P-values <0.05. R function 'prcomp' was used to perform
Principal Component Analysis (PCA) on log2RPKM values. The pseudobulk values were
calculated using the 'AverageExpression' function in Seurat. The pearson correlation were
calculated using the 'cor' function in R. The GO enrichment was calculated using the
'enrichGO' function from the ClusterProfiler package.

2.9 Quantitative real time PCR (qPCR)

Total RNA extracted using RNeasy Plus Micro-kit (Qiagen, 74034) was quantified using a
NanoDropTM ND-1000 Spectrophotometer and converted to cDNA using iScript cDNA
synthesis kit (Bio-Rad Laboratories, 1708890). qPCR was performed in 7000 Fast Real-Time
PCR System (Applied Biosystems, Foster City, USA) with FastStart Universal SYBR Green
Master (Rox) (Roche, 04 913 850 001) and primers from GeneCopoeia, Inc. (Rockville, MD,
USA). Gyceraldehyde-3-phosphate dehydrogenase (GAPDH) served as internal control.
Accession numbers of tested genes are listed in Table below:

Target Gene	Gene Name	Accession
GAPDH	glyceraldehyde-3-phosphate dehydrogenase	NM_001256799.2
AFP	Alpha fetoprotein	NM_000295
AAT	Alpha-1 antitrypsin	NM_001134
ASGPR	Asialoglycoprotein Receptor	NM_001671
ALB	albumin	NM_000477.3
CYP1A2	Cytochrome P450 1A2	NM_000761.5
CYP3A4	Cytochrome P450 3A4	NM_017460.5
CYP2B6	Cytochrome P450 2B6	NM_000767.4

CYP2C9	Cytochrome P450 2C9	NM_000771.3
UGT1A3	UDP-glucuroosyltransferase 1A3	NM_019093.2
GST1A2	Glutathione S-Transferase	NM_000846.3
MRP2	Multidrug resistance-associated protein 2	NM_000392
CD14	Cluster of differentiation 14	NM_001174105.1
CD32	Cluster of differentiation 32	NM_001136219.1
CD68	Cluster of differentiation 68	NM_001251.2
CD163	Cluster of differentiation 163	NM_203416.3
ID1	Inhibitor of DNA-binding protein 1	NM_002165.3
ID3	Inhibitor of DNA-binding protein 3	NM_002167.4

2.10   Enzyme-linked immunosorbent assay (ELISA) for measurement of cytokines

Interleukin-6 (IL-6) levels in the media were measured human IL-6 ELISA kit (Abcam, ab178013) according to manufacturer's instructions. The cytokine production from dosed cells was normalized with the viability of cells measured from the same well. This ensures that changes in cytokines are not due to changes in cell numbers that might arise upon drug treatment. The normalized cytokine level was expressed as percentage of DMSO+LPS which allowed better comparison between different batches of experiments. This normalization approach has been used in previous studies [19].

2.11   Cell viability assay

Cell viability was examined with AlamarBlue^{TM} cell viability assay (Thermo Fisher Scientific Inc., Dal1025) according to manufacturer's instructions. Briefly, AlamarBlue was diluted 10-fold using PBS containing 2 mg/mL of glucose and this working solution was added to the cells and incubated for 1 hour. Fluorescence (Ex: 530nm, Em: 590nm) readings were obtained using Tecan Microplate Reader M1000 PRO.

2.12   Drug administration

Following set up of co-culture as described in 2.5, the model was subjected to treatments including 100 ng/ml LPS (Sigma Aldrich, L2654), and drugs (Sigma Aldrich) (with/without LPS) for 48hr. Stock solutions of drugs were prepared in dimethyl sulfoxide (DMSO) and diluted in medium prior to use. Medium containing 0.1% DMSO was used as vehicle control except. After 48 hr, supernatant was collected cytokine measurement and cell viability in the same wells were measured as described in 2.11.

2.13   Statistical analysis

Mean and standard deviation were obtained from at least three independent batches of cells.
Unpaired, paired student's T-test and one-way or two-way analysis of variance (ANOVA)
were performed accordingly at an overall confidence level of 95% using Prism software
(GraphPad, San Diego, CA, USA) and indicated below each figure.

2.14 **Data Availability**

Accession codes

Raw data from RNA-seq analysis have been deposited in the NCBI Gene Expression
Omnibus under accession number GSExxxxx.

3. Results

265
266

3.1 Optimization of culture conditions allow survival and function of iHeps and iMacs

In order to investigate the effects of co-culturing iHeps with iMacs, we first established and
optimized a co-culture system utilising IMR90 iPSC-derived iHeps and iMacs (**Figure 1A**).
To ensure survival and functionality of both cell types in our desired culture period (at least
one week), certain key factors had to be accounted for: Hepatocytes are high in metabolically
activity and often require frequent media change for replenishment of nutrients and removal
of waste. On the other hand, it would be ideal to allow the iHeps and iMacs to interact
through direct crosstalk as well as through secreted soluble factors without removing these
factors from the system via media exchange. To assess the performance of hepatocytes with
more infrequent media changes, expression of key hepatic markers was assessed when media
was changed every other day, every two days or every three days and compared to a daily
media change (**Supplementary Figure 1**). To compensate for the lower frequency of media
change, we tested the effects of 2X supplements; standard media volume (**Supplementary
Figure 1A**) and 2X supplements with 2X standard culture volume (**Supplementary Figure
1B**). Gene expression was compared as fold change to standard hepatocyte culture conditions
(daily media change with 1X supplement and standard media volume). Increasing
supplements alone could maintain the expression of ALB (albumin), AAT (alpha-1-
antitrypsin), cytochrome P450 (CYP)- 1A2, 3A4, 2B6 and UDP-glucuronosyltransferase
(UGT)1A3, but not CYP2C19 (36 - 68% decrease), glutathione S-transferase (GST)A2 (35-
66% decrease) and multidrug resistance protein-2 (MRP2; 7 to 18% decrease)
(**Supplementary Figure 1A**). In contrast, combining the increase in supplements with
increase in culture volume resulted in maintenance of all hepatic markers (**Supplementary
Figure 1B**). Albumin and urea production were also maintained to 1-1.4 pg/cell/48 hrs and
126-146 pg/cell/48 hrs (**Supplementary Figure 1C**) which is in the range of albumin and
urea production reported by us previously [11, 14, 21] and in the state-of-the-art [22-26].
Once iHep marker and functionality was confirmed, the cells in the co-culture system were
sorted via flow cytometry on Day 3 and Day 7. Flow cytometric analysis showed the
presence of CD45$^+$CD11b$^+$CD14$^+$ iMacs and CD45$^-$CD11b$^-$ iHeps at Day 3 and Day 7 of co-
culture (**Supplementary Figure 1D**).

3.2 Co-culture with iMacs improves iHep maturation and development

Cells sorted via flow cytometry on Day 3 and Day 7, and the co-cultured hepatocytes (co-iHeps) and co-cultured iMacs (co-iMacs) were then analysed by RNA sequencing in order to assess if such co-culture lead to transcriptomic changes reflecting improve differentiation and maturation. We also analysed Day 0 and Day 7 mono-culture controls, as well as iMacs cultured in PHCM for 7 days as control groups.

Principle component analysis (PCA) of the iHep samples revealed that that although the direction of change was the same for both the mono-cultured and co-cultured iHeps, there was a greater degree of change along the PC1 axis for co-iHeps (**Figure 1B**). We hypothesised that this might be due to the iMacs promoting further maturation and development of the iHeps, and performed a Pearson correlation between the mono-cultured and co-cultured iHeps against *in vivo* fetal liver hepatocytes from a published dataset [27], which revealed that the co-cultured iHeps were better correlated with the *in vivo* hepatocytes than the mono-cultured iHeps (**Supplementary Figure 1E**). Indeed, looking at the differentially expressed genes (DEGs) between the Day 7 iHeps and Day 7 co-iHeps revealed that the co-iHep upregulated genes associated with tissue morphogenesis and repair like Early Growth Response 1 (*EGR1*) [28], centrosome duplication such as the long non-coding RNA *CCDC14* [29] and angiogenesis-related gene like Maternally Expressed Gene 9 (*MEG9*) [30] (**Figure 1C, Supplementary Table 1**). On the other hand, iHeps cultured alone upregulated genes that inhibit proliferation and angiogenesis such as Histone Rich Glycoprotein (*HRG*) [31] and Kinogen 1 (*KNG1*) [32] (**Figure 1C, Supplementary Table 1**). We then performed pathway analysis using Ingenuity and discovered that the top upregulated pathway in Day 7 iHeps was acute phase response signalling, with the associated genes mainly relating to protein production such as Transthyretin (*TTR*) and Serpin Family D Member 1 (*SERPIND1*) (**Supplementary Table 2**), while the top upregulated pathway in Day 7 co-iHeps was related to nitric oxide and reaction oxygen species production, consisting mainly of metabolic and phosphatase genes like *c-FOS* and Serine/Threonine protein phosphatase 2A regulatory subunit B" beta (*PPP2R3B*) (**Figure 1D, Supplementary Table 3**). A closer look at the pathways upregulated in Day 7 iHeps and Day 7 co-iHeps also revealed that the Day 7 co-iHeps had a stronger tissue development and anabolic signature, upregulating pathways associated with tumor microenvironment, fibrosis and growth hormone signalling, while the Day 7 iHeps upregulated iron and cholesterol pathways.

Finally, we compared the expression of key genes across all the iHep samples (**Figure 1E**). In line with our hypothesis that co-culturing iMacs with iHeps would improve the maturation of the iHeps, we found that alpha-fetoprotein (*AFP*), a marker that is upregulated during fetal development and then downregulated as hepatocytes mature [33] followed a similar trend in our culture system, with lower expression in the Day 7 iHeps as compared to the Day 0 iHeps. Furthermore, the expression levels of *AFP* in Day 3 co-iHeps and Day 7 co-iHeps was even lower than that of Day 7 iHeps. Likewise Hepatocyte Nuclear Factor 4 (*HNFA4*), a transcription factor upregulated in hepatic progenitor cells [34], had reduced expression in the co-cultured iHeps. Insulin-like growth factor-2 (*IGF-2*), a key growth factor in fetal development [35], was upregulated in the co-cultured iHeps, with expression peaking 3 days after co-culture. Cytochrome P450 Family 2 Subunit C member 9 (*CYP2C9*), a cytochrome

involved in drug metabolism was also upregulated in the co-iHeps, while Aldolase Fructose-
Biphosphate B (*ALDOB*), an enzyme whose upregulation is associated with unregulated cell
proliferation and poor cancer prognosis [36] was downregulated. Taken together, our data
suggest that co-culturing the iHeps with iMacs increased their maturation and promotes
improved hepatocyte development.

3.3 Co-culture with iHeps imparts KC-like identity to iMacs

Next, we turned our attention towards the iMacs within our co-culture system to test if they
acquired KC features. PCA analysis showed that the co-iMacs clustered much tighter
together, while iMacs that were cultured alone or in hepatocyte conditioned media (cond-
iMac) were much more scattered (**Figure 2A**). This suggests that co-culturing iMacs in direct
contact with iHeps may help maintain the identity of the iMacs better, mirroring how *ex vivo*
resident tissue macrophages (RTMs) rapidly lose their identity when removed from their
tissue environment [37]. DEG analysis of all the iMac groups showed that both the Day 7
iMacs and the Day 7 cond-iMacs shared an activated and inflammatory profile, upregulating
genes such as TNFSF18 [38] and ACP5 [39] (**Figure 2B, Supplementary Table 4**). Day7
iMacs had a more migratory profile, uniquely expressing Chemokine Receptor 7 (*CCR7*) and
Chemokine Ligand 5 (*CCL5*), while Day 7 cond-iMacs uniquely expressed Dishevelled
Associated Activator of Morphogenesis 2 (*DAAM2*), a developmental regulator of the WNT
pathway [40].

As we were interested in how the presence of iHeps in such co-culture model might educate
and impart KC-like identity to the iMacs, we looked at the DEGs between Day 0 iMac and
Day 7 co-iMacs. Day 0 iMacs were more immunogenic, expressing Proteoglycan 2 Pro
Eosinophil Major Basic Protein 2/3 (*PRG2/PRG3*) and Carboxypeptidase A3 (*CPA3*) (**Figure
2C**). In contrast, Day 7 co-iMacs upregulated the angiogenic Angiopoietin-like 4 (*ANGPTL4*)
[41], matching the angiogenic signature observed in the Day 7 co-iHeps. Importantly,
Ingenuity pathway analysis revealed that the top pathway in Day 7 co-iMacs is the Liver X
Receptor/Farnesoid X Receptor (LXR/RXR) signalling pathway (**Figure 2D,
Supplementary Table 5**), a key regulator of KC identity [42], while the top pathway in Day
0 iMacs is the axonal guidance pathway, consisting of chemokines like C-X-C Motif
Chemokine Ligand 12 (*CXCL12*) and C-X-C Motif Chemokine Receptor 4 (*CXCR4*),
metalloproteases like ADAM Metallopeptidase with Thrombospondin Type 1 Motif 15
(*ADAMTS15*), and semaphorins like Semaphorin 3C (*SEMA3C*) (**Supplementary Table 6**).
This suggests that co-culturing iMacs with iHeps initiated a KC-specific programme, while
the Day 0 iMacs had a more general and unspecified identity lacking environmental cues. In
addition, Day 7 co-iMacs also upregulated pathways associated with tissue repair and
remodelling, as well as tissue growth factors, while migratory and immune signalling related
pathways were upregulated in the Day 0 iMacs (**Figure 2D**). Altogether, these data suggest
that co-culturing iMacs with iHeps is sufficient to impart a more tissue supportive and KC-
like identity to the iMacs.

The use of conditioned media as a surrogate of the *in vivo* tissue microenvironment is a
popular approach, and we wondered how different iMacs cultured in direct co-culture with
iHeps would be from iMacs cultured in conditioned media from primary hepatocytes
(PHCM). Comparing the DEGs between Day 7 cond-iMacs and Day 7 co-iMacs, the Day 7

cond-iMacs upregulated immune-related genes such as Toll-Like Receptor 4 (*TLR4*) and *CPA3*, while the Day 7 co-iMacs more highly expressed tissue residency-related Fc Gamma Receptor IIIa (*FCGR3A*) (**Figure 2E**). Pathway analysis showed that the top upregulated pathway in the Day 7 cond-iMacs was autophagy, which might indicate that conditioned media alone was insufficient to maintain proper macrophage biology in the absence of contact with other cells, although some hepatic related pathways were also upregulated as well (**Figure 2F, Supplementary Table 7**). On the other hand, the most highly upregulated pathway in the Day 7 co-iMacs was the FXR/RXR pathway, which along with the upregulation of other anabolic pathways related to hepatic health (**Supplementary Table 8**), suggest that co-culturing rather than conditioned media is superior at inducing KC-like identity in macrophages.

We then looked at the expression of selected key KC genes. Lymphatic Vessel Endothelial Hyaluronan Receptor 1 (*LYVE1*), DNA-binding protein inhibitor ID-1 (*ID1*) and DNA-binding protein inhibitor ID-3 (*ID3*), which are important markers of KC identity [42]. These were all upregulated in the co-iMacs but not in the cond-iMacs (**Figure 2G**). Retinoid X Receptor Alpha (*RXRA*), the obligate binding partner of LXR [43], was also upregulated in only the co-iMacs. Interestingly, we also found that the co-iMacs upregulated the expression of Heparin Binding EGF Like Growth Factor (*HBEGF*) [44] and Oncostatin M (*OSM*) [45], which are potent effectors of liver regeneration and development.

Finally, we compared the iMacs from our study with *in vivo* embryonic human liver monocytes and KCs [46]. Pearson correlation revealed that regardless of condition the *in vitro* derived iMacs were poorly correlated to the *in vivo* human liver monocyte/macrophages (**Supplementary Figure 2A**), suggesting that either the co-culture duration was insufficient to induce full KC commitment or that there might be missing cues from other cell types that were not present in our co-culture. However, pathway analysis on the DEGs between the embryonic human liver macrophages and either the Day 7 cond-iMacs (**Supplementary Figure 2B, Supplementary Table 9**) or Day 7 co-iMacs (**Supplementary Figure 2C, Supplementary Table 10**) revealed that pathways relating to cytokine production, sensory development and eye development were differentially regulated between the Day 7 cond-iMacs and embryonic liver macrophages, while there were no significant pathway differences between the Day 7 co-iMacs and embryonic liver macrophages. This suggests that direct co-culture might drive the macrophages to adopt a more tissue developmental or supportive phenotype.

Altogether, the data supports our hypothesis that co-culturing iMacs with iHeps induces a KC-like tissue supportive identity in the iMacs, which in turn promotes the maturation and development of the iHeps.

3.4 Validation of the acquisition of KC-like identity by iMacs and maturation of iHeps in co-culture

We confirmed the observations from the RNA sequencing analysis by analysing gene expression of macrophage and KC-specific markers in Day 3 co-iMacs and Day 7 co-iMacs via RT-PCR. Macrophage markers *CD68*, *CD163*, *CD11b*, *CD32* and *CD14* were all upregulated at Day 3 and Day 7 by 1.8 to 7.6 fold compared to iMac mono-culture control

(**Figure 3A**). Importantly, KC-specific markers *ID1* and *ID3* were upregulated by 3.8 fold
and 3.9 fold respectively at Day 3 and by 6.3 fold and 6.5 fold respectively at Day 7 (**Figure**
3B). This suggests that macrophage phenotype is maintained during the 7 days of co-culture
but KC-specific development increases in a time-dependent manner. When Day 7 co-iMacs
were compared to iMacs cultured in PHCM (Day 7 cond-iMacs; without the presence of
iHeps), macrophage marker upregulation compared to iMac mono-culture control were
similar in both conditions (**Figure 3C**) but KC-specific *ID1* and *ID3* were upregulated to a
lower extent in Day 7 cond-iMacs (2.6 fold and 2.5 fold respectively) compared to Day 7 co-
iMacs (6.3 fold and 6.5 fold respectively; **Figure 3D**).

Next, we confirmed the RNA sequence analysis of iHep via RT-PCR. With the exception of
Glutathione-S transferase1A2 (*GST1A2*) and Multidrug Resistance Associated Protein 2
(*MRP2*), co-culture with iMacs improved expression of several hepatic marker genes
including *ALB*, α-1 antitrypsin (*AAT*), cytochrome P450 enzymes (*CYP1A2*, *CYP3A4*,
CYP2C9, *CYP2B6*) and UDP-glucuronosyltransferases (*UGT1A3*) by 1.8-53.1 fold at Day 3
and Day 7 of co-culture (**Figure 3E**). Importantly, *AFP*, a marker that is expressed at high
levels in fetal hepatocytes and increases with hepatocyte maturation [30], had 37% lower
expression at Day 3 and 37% lower expression in Day 7 iHeps as compared to Day 0 iHeps
(**Figure 3E**). This is consistent with RNA sequencing data that showed a downregulation of
AFP (**Figure 1E**).

Overall, validation of the expression of key markers observed as modulated by RNA
sequencing analysis confirms that culturing iMacs with iHeps imparts KC-like phenotype in
iMacs and supports maturation and function of the iHeps.

3.5 Co-culture model accurately mimics clinical/*in vivo* immune responses of drugs

Drug-induced liver injury is a complex and critical clinical problem with an estimated
incidence rate of up to 19 cases per 100000 people in western countries [47]. In particular,
immune-mediated idiosyncratic drug-induced liver injuries remains especially poorly
understood, due in large part to the lack of appropriate human modelling systems [6]. To test
the application of the co-culture in detecting immune-mediated response to hepatotoxic
drugs, we carefully selected a group of 7 drugs based on the following criteria: 1) confirmed
reports of immune/inflammation-associated effects according to 3 databases [48-50]; 2)
available clinical data, or *in vivo* responses if the former is unavailable; and 3) known Cmax
(maximum serum concentration of drug) to guide the concentration of drug used in in the *in*
vitro model. Drug concentrations used in our system was no more than 20-fold of its Cmax
(Details of selected drugs are summarized in **Supplementary Table 11**). The drugs included
diclofenac (DIC), sulindac (SLD) leflunomide (LFM); amodiaquine (AQ), lamotrigine
(LTG), penicillin (PEN), pyrazinamide (PZA). DIC [51], SLD [52] and LFM [53, 54] have
been reported to cause a decrease in interleukin-6 (IL-6); whereas AQ and LTG have been
reported to cause an increase in IL-6 in patients with drug-induced liver injury [50]. PEN and
PZA were selected as negative controls as no immune-mediated effects of these drugs have
been reported.

Our co-culture model could correctly recapitulate IL-6 decrease in DIC (48%, 45% and 18%
at 15.7 μM, 50 μM and 150 μM respectively), SLD (46%, 11% and 8% at 67 μM, 200 μM

and 600 μM respectively) and LFM (65%, 46%, 38% at 32.5 μM, 125 μM and 500 μM
respectively). IL-6 increase upon AQ (157-200 fold) and LTG (155-322 fold) treatment was
also correctly recapitulated. No changes in IL-6 level in PEN and PZA-treated samples were
observed (**Figure 4**). Importantly, when iMac-derived iKC-like cells were replaced with
blood Monocytes/Macrophages, none of these cytokine changes were recapitulated with the
exception of SLD treatment (**Figure 5**). This shows, from a functional/application
perspective, how iMac-iKCs, but not MoMϕs, are indispensable for *in vitro* liver models that
can detect immune-mediated changes of hepatotoxicants.

In vivo, KCs have been known to release mediators that are early features of inflammatory
responses [55]. They are activated in response to inflammatory stimuli such as bacterial
lipopolysaccharide (LPS) that can damage the liver at large doses in a KC-dependent manner.
At smaller doses, LPS activates KCs (without causing liver injury) but sensitizes the liver to a
variety of other xenobiotics [56]. In our model, when the co-culture was co-treated with
paradigm hepatotoxicants and LPS, drug-dose-dependent changes in cytokine production
were observed (**Figure 4**), suggesting that the system can indeed mimic physiological
responses, demonstrating the importance of liver-specific imprinting of macrophages in
recapitulating responses in an *in vitro* liver model.

4. Discussion

IPSC-derived tissue cultures have tremendous potential in its ability to generate cells from all
intra-embryonic lineages and have greatly expanded both our knowledge of mammalian
tissue development and our technical toolbox for therapy and disease modelling [57].
However, a tissue is often more than the sum of its parts, and iPSC-derived tissue models
have so far proven to be limited in their ability to effectively recapitulate the functionality
and complexities of their presumed organs. In our study, we demonstrated how co-culturing
iMacs with iHeps leads to not only acquisition of resident tissue characteristics within the
iMac population, but also increased maturation of iHeps. We demonstrated that iPSC-derived
Heps can directly impart hepatic macrophage-like identity to iMacs, giving them an iKC-like
phenotype and genotype without the need for any additional soluble factors or components
present in conditioned media. Application of this model was demonstrated by the ability of
the system to mimic clinical IL-6 responses of 5 paradigm immune-mediated hepatotoxicants.
Two additional hepatotoxicants with no known immune-mediated responses did not show
any response in our model, demonstrating the specificity of the system. Finally, when the
immune-mediated drug effects were tested in a system with iHeps and PBMC-derived
macrophages, expected cytokine responses were not observed, indicating the importance of
iKC-iHep co-culture in predicting drug mediated-immune responses.

Animal models have been used to model immune-mediated liver injury [58-60]; however,
consistent liver injury was not observed upon treatment with drugs known to cause immune
response in humans, even at high doses [61]. In cases where injury was observed, the
characteristics were different from humans and possibly manifested through different
mechanisms [60]. Such interspecies variations, contradictory findings as well as cost and
ethical issues have spurred the development of *in vitro* models to supplement *in vivo* animal
models. *In vitro* models, possibly due to the lack of appropriate human immune cell sources,
have focused on soluble factor-driven single hepatocyte cultures in combination with

inflammatory mediators or animal cell co-cultures [6]. However, mono-cultures lack cell-cell
interactions, and animal *in vitro* models often fail to recapitulate *in vivo* and clinical findings.
In vitro hepatocyte-KC models have previously been used; however most of them have
utilised animal cells [62,63] which often prove inadequate in recapitulating *in vivo* or clinical
findings. In order to avoid interspecies variations and improve the predictivity of *in vitro*
models, appropriate human models still need to be developed and characterized.

The main challenge in studying human KCs in *in vitro* models is the cell source. The source
of commercial primary human KCs (PhKCs) is limited and costly and high donor-to-donor
variability is often observed [64]. Alternative human cell sources include THP-1 cells and
peripheral monocyte-derived macrophages; however their cytokine profiles are dramatically
different from human KCs [65, 66] and they lack liver-specific imprinting [42]. Even when
human KCs were used, the work mostly focused on co-culture dependent improvement of
hepatocyte function [6, 67], rather than interaction between the two cell-types and immune-
mediated effect of drugs. Moreover, these models are often non-isogenic, which further
confounds the inflammatory readouts. Hence, in our study, we aimed to first determine if the
presence of hepatocytes could impart a tissue-resident-like phenotype to iMacs and drive
them to be more Kupffer cell-like, then measured the systemic response to drug treatment
compared to using monocyte-derived macrophages. While all tested drugs showed expected
changes in cytokine production in the iMac-iHep co-culture (**Figure 4**), these changes were
not recapitulated when the iMacs were replaced by MoMφs (**Figure 5**) with the exception of
SLD. Interestingly, expected IL-6 increase upon treatment with LFM could not be
recapitulated in our previous model of iHep-iKCs co-culture where iKCs were generated
using conditioned media from primary hepatocytes (data not shown). However, in our current
model, LFM treatment resulted in the expected increase in IL-6, suggesting the potential
importance of direct cellular cross-talk in iKCs differentiation and function.

Development of a suitable model system is further complicated by the understanding that
KCs are derived from embryonic macrophages that seed the liver early on during
development [7-9]. This also implies that conventionally used human peripheral blood
MoMφs might not be able to fully mimic KC biology. Thankfully, iMacs more closely
resemble embryonic macrophages than MoMφs [10] and as demonstrated in our own
experiments, were able to acquire features of KCs upon co-culture with iHeps. Soluble
factors were only partially responsible for this tissue adaptation, as in contrast to the iMacs
that were cultured in hepatocyte conditioned media, the co-cultured iMacs upregulated genes
in the LXR/RXR pathway, which has been shown to be important for KC specification and
identity [42], further emphasising the importance of cell-cell contact for the acquisition of
RTM identity. Furthermore, we also observed that these co-cultured iMacs were upregulating
pathways associated with tissue repair and development, which might explain the increased
maturity of the iHeps within the co-culture system. This is particularly relevant not only for
hepatic model systems, but for other iPSC-derived model systems in general, as one of the
key limitations of these systems is the functional and phenotypic immaturity of the
differentiated cells [68]. Co-culturing iMacs with these immature iPSC-derived tissue model
systems might provide the missing developmental cues needed for more physiologically and
functionally relevant patient and disease models as well.

Acknowledgements

The authors of this paper are supported by the Agency for Science, Technology and Research (A*STAR) under its Industry Alignment Fund –Pre-Positioning Programme (IAF-PP) grant number H18/01/a0/D14 as part of the A*STAR Innovations in Food and Chemical Safety (IFCS) Programme as well as Institute of Bioengineering and Nanotechnology (IBN), Institute of Bioengineering and Nanotechnology (IBN) and Singapore Immunology Network (SIgN) core funding. F.G. is a European Molecular Biology Organization (EMBO) YIP awardee and supported by the Singapore National Research Foundation Senior Investigatorship (NRFI) NRF2016NRF-NRFI001-02. H.Y is supported by the National Medical Research Council (NMRC) CIRG21nov-0032 and CIRG22jul-0018, as well as funding from the Mechanobiology Institute of Singapore (MBI) A-0003467-00-00. C.Z.W.L. is supported by the A*STAR Graduate Scholarship.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

5. References

[1] Z. Shan, C. Ju, Hepatic macrophages in liver injury, Frontiers in immunology 11 (2020) 322.

[2] L.J. Dixon, M. Barnes, H. Tang, M.T. Pritchard, L.E. Nagy, Kupffer cells in the liver, Comprehensive Physiology 3(2) (2013) 785.

[3] O. Krenkel, F. Tacke, Liver macrophages in tissue homeostasis and disease, Nature Reviews Immunology 17(5) (2017) 306-321.

[4] C. Blériot, F. Ginhoux, Understanding the heterogeneity of resident liver macrophages, Frontiers in Immunology (2019) 2694.

[5] Y. Wen, J. Lambrecht, C. Ju, F. Tacke, Hepatic macrophages in liver homeostasis and diseases-diversity, plasticity and therapeutic opportunities, Cellular & molecular immunology 18(1) (2021) 45-56.

[6] F. Tasnim, X. Huang, C.Z.W. Lee, H. Yu, F. Ginhoux, Recent Advances in Models of Immune-mediated Drug-Induced Liver Injury, Frontiers in Toxicology 3 (2021) 22.

[7] E. Mass, I. Ballesteros, M. Farlik, F. Halbritter, P. Günther, L. Crozet, C. E. Jacome-Galarza, K. Händler, J. Klughammer, Y. Kobayashi, E. Gomez-Perdiguero, J. L. Schultze, C. Beyer, M., Bock, & F. Geissmann (2016). Specification of tissue-resident macrophages during organogenesis. *Science*, *353*(6304).

[8] C.Z.W. Lee, F. Ginhoux, Biology of resident tissue macrophages, Development 149(8) (2022) dev200270.

[9] Scott, C. L., Zheng, F., De Baetselier, P., Martens, L., Saeys, Y., De Prijck, S., Lippens, S., Abels, C., Schoonooghe, S., Raes, G., Devoogdt, N., Lambrecht, B. N., Beschin, A., & Guilliams, M. Bone marrow-derived monocytes give rise to self-renewing and fully differentiated Kupffer cells. *Nature Communications* (2016) *7*(1).

[10] K. Takata, T. Kozaki, C.Z.W. Lee, M.S. Thion, M. Otsuka, S. Lim, K.H. Utami, K. Fidan, D.S. Park, B. Malleret, Induced-pluripotent-stem-cell-derived primitive macrophages provide a platform for modeling tissue-resident macrophage differentiation and function, Immunity 47(1) (2017) 183-198. e6.

[11] F. Tasnim, J. Xing, X. Huang, S. Mo, X. Wei, M.-H. Tan, H. Yu, Generation of mature kupffer cells
from human induced pluripotent stem cells, Biomaterials 192 (2019) 377-391.
[12] Park, D. S., Kozaki, T., Tiwari, S. K., Moreira, M., Khalilnezhad, A., Torta, F., Olivié, N., Thiam, C.
H., Liani, O., Silvin, A., Phoo, W. W., Gao, L., Triebl, A., Tham, W. K., Gonçalves, L., Kong, W. T.,
Raman, S., Zhang, X. M., Dunsmore, G., . . . Ginhoux, F. (2023). iPS-cell-derived microglia promote
brain organoid maturation via cholesterol transfer. *Nature*, *623*(7986), 397–405.
[13] J. Cai, J. Fisher, A. Urick, T. Wagner, K.T.M. Cayo, M. Nagaoka, S.A. Duncan, Protocol for directed
differentiation of human pluripotent stem cells toward a hepatocyte fate, African Scientist 16(1)
(2015) 13-26.
[14] F. Tasnim, Y.-C. Toh, Y. Qu, H. Li, D. Phan, B.C. Narmada, A. Ananthanarayanan, N. Mittal, R.Q.
Meng, H. Yu, Functionally enhanced human stem cell derived hepatocytes in galactosylated cellulosic
sponges for hepatotoxicity testing, Molecular pharmaceutics 13(6) (2016) 1947-1957.
[15] S. Picelli, O.R. Faridani, Å.K. Björklund, G. Winberg, S. Sagasser, R. Sandberg, Full-length RNA-seq
from single cells using Smart-seq2, Nature protocols 9(1) (2014) 171-181.
[16] A. Dobin, C.A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T.R.
Gingeras, STAR: ultrafast universal RNA-seq aligner, Bioinformatics 29(1) (2013) 15-21.
[17] Y. Liao, G.K. Smyth, W. Shi, featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features, Bioinformatics 30(7) (2014) 923-930.
[18] J. Harrow, F. Denoeud, A. Frankish, A. Reymond, C.-K. Chen, J. Chrast, J. Lagarde, J.G. Gilbert, R.
Storey, D. Swarbreck, GENCODE: producing a reference annotation for ENCODE, Genome biology
7(1) (2006) 1-9.
[19] M.D. Robinson, D.J. McCarthy, G.K. Smyth, edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data, Bioinformatics 26(1) (2010) 139-140.
[20] Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: a practical and powerful
approach to multiple testing, Journal of the Royal statistical society: series B (Methodological) 57(1)
(1995) 289-300.
[21] F. Tasnim, D. Phan, Y.-C. Toh, H. Yu, Cost-effective differentiation of hepatocyte-like cells from
human pluripotent stem cells using small molecules, Biomaterials 70 (2015) 115-125.
[22] K.A. Rose, N.S. Holman, A.M. Green, M.E. Andersen, E.L. LeCluyse, Co-culture of hepatocytes
and Kupffer cells as an in vitro model of inflammation and drug-induced hepatotoxicity, Journal of
pharmaceutical sciences 105(2) (2016) 950-964.
[23] N. Shiraki, T. Yamazoe, Z. Qin, K. Ohgomori, K. Mochitate, K. Kume, S. Kume, Efficient
differentiation of embryonic stem cells into hepatic cells in vitro using a feeder-free basement
membrane substratum, PloS one 6(8) (2011) e24228.
[24] K. Takayama, K. Kawabata, Y. Nagamoto, K. Kishimoto, K. Tashiro, F. Sakurai, M. Tachibana, K.
Kanda, T. Hayakawa, M.K. Furue, 3D spheroid culture of hESC/hiPSC-derived hepatocyte-like cells for
drug toxicity testing, Biomaterials 34(7) (2013) 1781-1789.
[25] L. Xia, X. Hong, R.B. Sakban, Y. Qu, N.H. Singh, M. McMillian, S. Dallas, J. Silva, C. Sensenhauser,
S. Zhao, H.K. Lim, H. Yu, Cytochrome P450 induction response in tethered spheroids as a three-
dimensional human hepatocyte in vitro model, J Appl Toxicol 36(2) (2016) 320-9.
[26] M. Nishimura, M. Hagi, Y. Ejiri, S. Kishimoto, T. Horie, S. Narimatsu, S. Naito, Secretion of
albumin and induction of CYP1A2 and CYP3A4 in novel three-dimensional culture system for human
hepatocytes using micro-space plate, Drug metabolism and pharmacokinetics 25(3) (2010) 236-242.
[27] Popescu, D., Botting, R. A., Stephenson, E., Green, K., Webb, S., Jardine, L., Calderbank, E. F.,
Polanski, K., Goh, I., Efremova, M., Acres, M., Maunder, D., Vegh, P., Gitton, Y., Park, J., Vento-
Tormo, R., Miao, Z., Dixon, D., Rowell, R., . . . Haniffa, M. Decoding human fetal liver haematopoiesis.
Nature (2019), *574*(7778), 365–371.
[28] M.T. Pritchard, J.I. Cohen, S. Roychowdhury, B.T. Pratt, L.E. Nagy, Early growth response-1
attenuates liver injury and promotes hepatoprotection after carbon tetrachloride exposure in mice,
Journal of hepatology 53(4) (2010) 655-662.

[29] E.N. Firat-Karalar, N. Rauniyar, J.R. Yates III, T. Stearns, Proximity interactions among
centrosome components identify regulators of centriole duplication, Current biology 24(6) (2014)
664-670.
[30] C. Espinosa-Diez, R. Wilson, R. Mukherjee, M. Feltham, C. Hudson, R. Ruhl, S. Anand, DNA
damage dependent hypomethylation regulates the pro-angiogenic LncRNA MEG9, BioRxiv (2018)
442699.
[31] Q. Zhang, K. Jiang, Y. Li, D. Gao, L. Sun, S. Zhang, T. Liu, K. Guo, Y. Liu, Histidine-rich glycoprotein
function in hepatocellular carcinoma depends on its N-glycosylation status, and it regulates cell
proliferation by inhibiting Erk1/2 phosphorylation, Oncotarget 6(30) (2015) 30222.
[32] R.W. Colman, B.A. Jameson, Y. Lin, D. Johnson, S.A. Mousa, Domain 5 of high molecular weight
kininogen (kininostatin) down-regulates endothelial cell proliferation and migration and inhibits
angiogenesis, Blood, The Journal of the American Society of Hematology 95(2) (2000) 543-550.
[33] L.T. Ang, A.K.Y. Tan, M.I. Autio, S.H. Goh, S.H. Choo, K.L. Lee, J. Tan, B. Pan, J.J.H. Lee, J.J. Lum,
C.Y.Y. Lim, I.K.X. Yeo, C.J.Y. Wong, M. Liu, J.L.L. Oh, C.P.L. Chia, C.H. Loh, A. Chen, Q. Chen, I.L.
Weissman, K.M. Loh, B. Lim, A Roadmap for Human Liver Differentiation from Pluripotent Stem
Cells, Cell Rep 22(8) (2018) 2190-2205.
[34] P. Chaudhari, L. Tian, A. Deshmukh, Y.-Y. Jang, Expression kinetics of hepatic progenitor markers
in cellular models of human liver development recapitulating hepatocyte and biliary cell fate
commitment, Experimental Biology and Medicine 241(15) (2016) 1653-1662.
[35] G.D. Agrogiannis, S. Sifakis, E.S. Patsouris, A.E. Konstantinidou, Insulin-like growth factors in
embryonic and fetal growth and skeletal development, Molecular medicine reports 10(2) (2014)
579-584.
[36] Y.-C. Chang, Y.-C. Yang, C.-P. Tien, C.-J. Yang, M. Hsiao, Roles of aldolase family genes in human
cancers and diseases, Trends in Endocrinology & Metabolism 29(8) (2018) 549-559.
[37] D. Gosselin, D. Skola, N.G. Coufal, I.R. Holtman, J.C. Schlachetzki, E. Sajti, B.N. Jaeger, C.
O'Connor, C. Fitzpatrick, M.P. Pasillas, An environment-dependent transcriptional network specifies
human microglia identity, Science 356(6344) (2017).
[38] M. Croft, W. Duan, H. Choi, S.-Y. Eun, S. Madireddi, A. Mehta, TNF superfamily in inflammatory
disease: translating basic insights, Trends in immunology 33(3) (2012) 144-152.
[39] S.R. Räisänen, S.L. Alatalo, H. Ylipahkala, J.M. Halleen, A.I. Cassady, D.A. Hume, H.K. Väänänen,
Macrophages overexpressing tartrate-resistant acid phosphatase show altered profile of free radical
production and enhanced capacity of bacterial killing, Biochemical and biophysical research
communications 331(1) (2005) 120-126.
[40] N. de Alwis, S. Beard, N.K. Binder, N. Pritchard, J. Tu'uhevaha, S.P. Walker, O. Stock, K. Groom, S.
Petersen, A. Henry, DAAM2 is elevated in the circulation and placenta in pregnancies complicated by
fetal growth restriction and is regulated by hypoxia, Scientific reports 11(1) (2021) 1-12.
[41] S. Babapoor-Farrokhran, K. Jee, B. Puchner, S.J. Hassan, X. Xin, M. Rodrigues, F. Kashiwabuchi, T.
Ma, K. Hu, M. Deshpande, Angiopoietin-like 4 is a potent angiogenic factor and a novel therapeutic
target for patients with proliferative diabetic retinopathy, Proceedings of the National Academy of
Sciences 112(23) (2015) E3030-E3039.
[42] J. Bonnardel, W. T'Jonck, D. Gaublomme, R. Browaeys, C.L. Scott, L. Martens, B. Vanneste, S. De
Prijck, S.A. Nedospasov, A. Kremer, Stellate cells, hepatocytes, and endothelial cells imprint the
kupffer cell identity on monocytes colonizing the liver macrophage niche, Immunity 51(4) (2019)
638-654. e9.
[43] L. Yue, F. Ye, C. Gui, H. Luo, J. Cai, J. Shen, K. Chen, X. Shen, H. Jiang, Ligand-binding regulation of
LXR/RXR and LXR/PPAR heterodimerizations: SPR technology-based kinetic analysis correlated with
molecular dynamics simulation, Protein science 14(3) (2005) 812-822.
[44] D.T. Dao, L. Anez-Bustillos, R.M. Adam, M. Puder, D.R. Bielenberg, Heparin-binding epidermal
growth factor–like growth factor as a critical mediator of tissue repair and regeneration, The
American journal of pathology 188(11) (2018) 2446-2456.

[45] A. Miyajima, T. Kinoshita, M. Tanaka, A. Kamiya, Y. Mukouyama, T. Hara, Role of Oncostatin M
in hematopoiesis and liver development, Cytokine & growth factor reviews 11(3) (2000) 177-183.
[46] Bian, Z., Gong, Y., Huang, T., Lee, C. Z. W., Bian, L., Bai, Z., Shi, H., Zeng, Y., Liu, C., He, J., Zhou, J.,
Li, X., Li, Z., Ni, Y., Ma, C., Cui, L., Zhang, R., Chan, J. K. Y., Ng, L. G., . . . Liu, B. Deciphering human
macrophage development at single-cell resolution. *Nature* (2020), *582*(7813)
[47] Hoofnagle, J. H., & Björnsson, E. S. (2019). Drug-Induced Liver Injury — Types and phenotypes.
New England Journal of Medicine, *381*(3), 264–273. https://doi.org/10.1056/nejmra1816149
[48] N. Kaplowitz, Idiosyncratic drug hepatotoxicity, Nature reviews. Drug discovery 4(6) (2005) 489-
99.
[49] S. Ho, A. McLachlan, T. Chen, D. Hibbs, R. Fois, Relationships Between Pharmacovigilance,
Molecular, Structural, and Pathway Data: Revealing Mechanisms for Immune-Mediated Drug-
Induced Liver Injury, CPT: Pharmacometrics Systems Pharmacology 4(7) (2015) 426-441.
[50] S. Verma, N. Kaplowitz, Diagnosis, management and prevention of drug-induced liver injury, Gut
58(11) (2009) 1555-1564.
[51] A. Mahdy, H.F. Galley, M. Abdel-Wahed, K. El-Korny, S. Sheta, N.R. Webster, Differential
modulation of interleukin-6 and interleukin-10 by diclofenac in patients undergoing major surgery,
British Journal of Anaesthesia 88(6) (2002) 797-802.
[52] B.-S. Kang, E.-Y. Chung, Y.-P. Yun, M.K. Lee, Y.R. Lee, K.-S. Lee, K.R. Min, Y. Kim, Inhibitory effects
of anti-inflammatory drugs on interleukin-6 bioactivity, Biological Pharmaceutical Bulletin 24(6)
(2001) 701-703.
[53] H.-W. Yao, J. Li, J.-Q. Chen, S.-Y. Xu, Leflunomide attenuates hepatocyte injury by inhibiting
Kupffer cells, World Journal of Gastroenterology 10(11) (2004) 1608.
[54] S.K. Manna, B.B. Aggarwal, Immunosuppressive leflunomide metabolite (A77 1726) blocks TNF-
dependent nuclear factor-κB activation and gene expression, The Journal of Immunology 162(4)
(1999) 2095-2102.
[55] N.M. Steuerwald, D.M. Foureau, H.J. Norton, J. Zhou, J.C. Parsons, N. Chalasani, R.J. Fontana,
P.B. Watkins, W.M. Lee, K.R. Reddy, Profiles of serum cytokines in acute drug-induced liver injury
and their prognostic significance, Plos One 8(12) (2013).
[56] R.A. Roberts, P.E. Ganey, C. Ju, L.M. Kamendulis, I. Rusyn, J.E. Klaunig, Role of the Kupffer cell in
mediating hepatic toxicity and carcinogenesis, Toxicological Sciences 96(1) (2007) 2-15.
[57] Rowe, R. G., & Daley, G. Q. (2019). Induced pluripotent stem cells in disease modelling and drug
discovery. *Nature Reviews Genetics*, *20*(7), 377–388.
[58] L. Kuna, I. Bozic, T. Kizivat, K. Bojanic, M. Mrso, E. Kralj, R. Smolic, G.Y. Wu, M. Smolic, Models of
Drug Induced Liver Injury (DILI)–Current Issues and Future Perspectives, Current drug metabolism
19(10) (2018) 830-838.
[59] J.M. Shenton, J. Chen, J.P. Uetrecht, Animal models of idiosyncratic drug reactions, Chemico-
biological interactions 150(1) (2004) 53-70.
[60] W. Ng, A.R. Lobach, X. Zhu, X. Chen, F. Liu, I.G. Metushi, A. Sharma, J. Li, P. Cai, J. Ip, Animal
models of idiosyncratic drug reactions, Advances in pharmacology, Elsevier2012, pp. 81-135.
[61] J. Uetrecht, Mechanistic studies of idiosyncratic DILI: clinical implications, Frontiers in
pharmacology 10 (2019) 837.
[62] J.A. Bonzo, K. Rose, K. Freeman, E. Deibert, K.B. Amaral, S.S. Ferguson, M.E. Andersen, R.P.
Witek, E.L. LeCluyse, Differential effects of trovafloxacin on TNF-α and IL-6 profiles in a rat
hepatocyte–Kupffer cell coculture system, Applied In Vitro Toxicology 1(1) (2015) 45-54.
[63] F.F. Tukov, J.F. Maddox, D.E. Amacher, W.F. Bobrowski, R.A. Roth, P.E. Ganey, Modeling
inflammation–drug interactions in vitro: A rat Kupffer cell-hepatocyte coculture system, Toxicology
in vitro 20(8) (2006) 1488-1499.
[64] V. Kegel, E. Pfeiffer, B. Burkhardt, J.L. Liu, K. Zeilinger, A.K. Nussler, D. Seehofer, G. Damm,
Subtoxic Concentrations of Hepatotoxic Drugs Lead to Kupffer Cell Activation in a Human In Vitro
Liver Model: An Approach to Study DILI, Mediators Inflamm 2015 (2015) 640631.

[65] P.J. Shaw, M.J. Hopfensperger, P.E. Ganey, R.A. Roth, Lipopolysaccharide and trovafloxacin
coexposure in mice causes idiosyncrasy-like liver injury dependent on tumor necrosis factor-alpha,
Toxicological Sciences 100(1) (2007) 259-266.
[66] A. Kermanizadeh, D.M. Brown, V. Stone, The variances in cytokine production profiles from non-
or activated THP-1, Kupffer cell and human blood derived primary macrophages following exposure
to either alcohol or a panel of engineered nanomaterials, Scientific reports 14(8) (2019) e0220974.
[67] E. Novik, T.J. Maguire, P. Chao, K. Cheng, M.L. Yarmush, A microfluidic hepatic coculture
platform for cell-based drug metabolism studies, Biochemical Pharmacology 79(7) (2010) 1036-1044.
[68] M.X. Doss, A. Sachinidis, Current challenges of iPSC-based disease modeling and therapeutic
implications, Cells 8(5) (2019) 403.
[69] M. Chen, A. Suzuki, S. Thakkar, K. Yu, C. Hu, W. Tong, DILIrank: the largest reference drug list
ranked by the risk for developing drug-induced liver injury in humans, Drug Discov Today 21(4)
(2016) 648-653.
[70] S. Thakkar, T. Li, Z. Liu, L. Wu, R. Roberts, W. Tong, Drug-induced liver injury severity and toxicity
(DILIst): Binary classification of 1279 drugs by human hepatotoxicity, Drug discovery today 25(1)
(2020) 201-208.
[71] J.J. Xu, P.V. Henstock, M.C. Dunn, A.R. Smith, J.R. Chabot, D. de Graaf, Cellular imaging
predictions of clinical drug-induced liver injury, Toxicological sciences : an official journal of the
Society of Toxicology 105(1) (2008) 97-105.
[72] W. Zou, K.M. Beggs, E.M. Sparkenbaugh, A.D. Jones, H.S. Younis, R.A. Roth, P.E. Ganey, Sulindac
metabolism and synergy with tumor necrosis factor-α in a drug-inflammation interaction model of
idiosyncratic liver injury, Journal of Pharmacology Experimental Therapeutics 331(1) (2009) 114-121.
[73] J. Li, H.-W. Yao, Y. Jin, Y.-F. Zhang, C.-Y. Li, Y.-H. Li, S.-Y. Xu, Pharmacokinetics of leflunomide in
Chinese healthy volunteers, Acta Pharmacologica Sinica 23(6) (2002) 551-555.
[74] I. Agabeyoglu, T. Incecayir, Pharmacokinetic Modelling of Lamotrigine from Plasma
Concentrations in Healthy Volunteers, J Bioanal Biomed 1 (2009) 041-045.
[75] A.A. Ademisoye, J.O. Soyinka, S.O. Olawoye, S.I. Igbinoba, S.A. Olowookere, A.T. Ademisoye,
C.O. Onyeji, Induction of Amodiaquine Metabolism by Rifampicin Following Concurrent
Administration in Healthy Volunteers, Journal of Exploratory Research in Pharmacology 3(3) (2018)
71-77.
[76] S. Bareggi, R. Cerutti, R. Pirola, R. Riva, M. Cisternino, Clinical pharmacokinetics and metabolism
of pyrazinamide in healthy volunteers, Arzneimittel-Forschung 37(7) (1987) 849.
784
785

786
787

Figure 1: Model development and effects of iMacs on iHeps. A) Schematic of experimental layout. B) Principal Component Analysis of bulk RNA-seq data from Day 0 iHeps, Day 3 co-iHeps, Day 7 co-iHeps and Day 7 iHeps. C) Volcano plot showing differentially-expressed genes (DEGs) between Day 7 iHeps and Day 7 co-iHeps. D) Differentially-expressed upregulated pathways between Day 7 iHeps and Day 7 co-iHeps. E) Gene expression levels of *AFP*, *CYP2C9*, *IGF-2*, *EGR-1*, *ALDOB* and *HNF4A* in Day 0

iHeps, Day 3 co-iHeps, Day 7 co-iHeps and Day 7 iHeps. Student's T test was applied.
*p<0.05, **p<0.01

Figure 2. Effects of iHeps on iMacs. A) Principal Component Analysis of bulk RNA-seq
data from Day 0 iMacs, Day 3 co-iMacs, Day 7 co-iMacs, Day 7 iMacs and Day 7 cond-
iMacs. B) Heatmap showing the top 5 differentially (DEGs) and uniquely (UEGs) expressed
genes from culturing iMacs alone, with conditioned media (cond-iMacs) and after co-culture
with iHepatocytes (Day 7 co-iMacs). C) Volcano Plot showing differentially-expressed genes
between Day 0 iMacs and Day 7 co-iMacs. D) Differentially expressed upregulated pathways
between Day 0 iMacs and Day 7 co-iMacs. E) Volcano Plot showing DEGs between Day 7
cond-iMacs and Day 7 co-iMacs. F) Differentially expressed upregulated pathways between
Day 7 cond-iMacs and Day 7 co-iMacs. G) Gene expression levels of *LYVE1*, *ID1*, *ID3*,
RXRA, *HBEGF* and *OSM* in Day 0 iMacs, Day 3 co-iMacs, Day 7 co-iMacs, Day 7 iMacs
and Day 7 cond-iMacs. Student's T test was applied. *p<0.05, **p<0.01, ***p<0.001

Figure 3. Gene expression of iMac and iHep markers in co-cultures. A) Expression of
macrophage markers (left panel) and KC-specific markers (right panel) on day 3 and day 7 of
co-culture. B) Expression of macrophage markers (left panel) and KC-specific markers (right
panel) on day 7 of co-culture compared to iMac cultured in PHCM. C) Expression of hepatic
markers (left panel) on day 3 and day 7 of co-culture. * p<0.05.

Figure 4. Changes in IL-6 level upon treatment with 7 paradigm compounds when iMac-derived KCs were used. Cytokine production was assessed in iMac-derived iKC and iHep co-culture treated with the drug stated in the plot title. Data is expressed as the percentage of the LPS-treated vehicle control. Error bars represented S.D., n=3. One-way ANOVA was applied. *p<0.05, **p<0.01 between treatment and vehicle control.

Figure 5. Changes in IL-6 level upon treatment with 7 paradigm compounds without iMac-derived KCs. Cytokine production was assessed in blood monocyte-derived macrophages and iHep co-culture treated with the drug stated in the plot title. Data is expressed as the percentage of the LPS-treated vehicle control. Error bars represented S.D., n=3. One-way ANOVA was applied. *p<0.05, **p<0.01 between treatment and vehicle control.

Drug	Cmax/ (µM)	DILIrank [64]	DILIst [65]	Known immune-mediated effects [43-45]	Reported human responses	Reported *in vivo* (murine) responses
DIC	8.0 [69]	most	1	✓	IL-6↓ [51] IL-10↑ [51]	
SLD	32.0 [69]	most	1	✓	mechanistic IL-6↓ [52]	TNFα↑ [72]
LFM	24.8 [73]	most	1	✓		cytokines↓ via NF-kB [53,]
LTG	9.8 [74]	most	1	✓	IL-6↑[55]	
AQ	0.1 [75]	/	1	✓	IL-6↑ [55]	
PEN	/	less	1	×		
PZA	406.9 [76]	less	1	×		

Supplementary Table 11: Details of drugs used for testing

Supplementary Figure 1: Optimisation of co-culture conditions and correlation of iPSC-derived hepatocytes with or without co-culture with iPSC-derived macrophages. A & B) Fold change in expression of key hepatocyte genes in normal media volume and supplement concentration (A) or double media volume and supplement concentration (B), normalised to normal hepatocyte culture condition. C) Production of Albumin and Urea in co-culture system after 2 days, normalised to non-co-cultured hepatocytes. D) Flow cytometry of co-culture 3 and 7 days after co-culture. E) Pearson correlation of iHeps against fetal hepatocytes from Popescu et al, (2019).

A

B

C

Supplementary Figure 2: Correlation of iMacs with embryonic liver monocytes and
macrophages, and pathway analysis of the DEGs. A) Pearson correlation of iMacs under
control, conditioned media or co-culture conditions against embryonic liver monocytes and
macrophages from Bian *et al.*, (2020). B) Pathway analysis of DEGs between Day 7
conditioned media iMacs and embryonic liver macrophages. C) Pathway analysis of DEGs
between Day 7 co-cultured iMacs and embryonic liver macrophages

List of Supplementary Tables:

Supplementary Table 1: Differentially Expressed Genes between the iHeps Samples

Supplementary Table 2: Ingenuity Pathway Analysis for genes upregulated in Day 7 iHeps vs
Day 7 co-iHeps

Supplementary Table 3: Ingenuity Pathway Analysis for genes upregulated in Day 7 co-iHeps
vs Day 7 iHeps

Supplementary Table 4: Differentially Expressed Genes between the iMac Samples

Supplementary Table 5: Ingenuity Pathway Analysis for genes upregulated in Day 7 co-
iMacs vs Day 0 iMacs

Supplementary Table 6: Ingenuity Pathway Analysis for genes upregulated in Day 0 iMacs vs
Day 7 co-iMacs

Supplementary Table 7: Ingenuity Pathway Analysis for genes upregulated in Day 7 cond-
iMacs vs Day 7 co-iMacs

Supplementary Table 8: Ingenuity Pathway Analysis for genes upregulated in Day 7 co-
iMacs vs Day 7 cond-iMacs

Supplementary Table 9: GO enrichment results for DEGs between Day 7 conditioned media
iMacs and embryonic liver macrophages

Supplementary Table 10: GO enrichment results for DEGs between Day 7 co-cultured iMacs
and embryonic liver macrophages

Supplementary Table 11: Details of drugs used for testing

Figure 1: Model development and effects of iMacs on iHeps. A) Schematic of experimental layout. B) Principal Component Analysis of bulk RNAseq data from Day 0 iHeps, Day 3 co-iHeps, Day 7 co-iHeps and Day 7 iHeps. C) Volcano plot showing differentially-expressed genes between Day 7 iHeps and Day 7 co-iHeps. D) Differentially-expressed upregulated pathways between Day 7 iHeps and Day 7 co-iHeps. E) Gene expression levels of *AFP*, *CYP2C9*, *IGF-2*, *EGR-1*, *ALDOB* and *HNF4A* in Day 0 iHeps, Day 3 co-iHeps, Day 7 co-iHeps and Day 7 iHeps. Student's T test was applied. *p<0.05, **p<0.01

Supplementary Figure 1: Optimisation of co-culture conditions and correlation of iPSC-derived hepatocytes with or without co-culture with iPSC-derived macrophages. A & B) Fold change in expression of key hepatocyte genes in normal media volume and supplement concentration (A) or double media volume and supplement concentration (B), normalised to normal hepatocyte culture condition. C) Production of Albumin and Urea in co-culture system after 2 days, normalised to non-co-cultured hepatocytes. D) Flow cytometry of co-culture 3 and 7 days after co-culture. E) Pearson correlation of iHeps against fetal hepatocytes from Popescu et al, (2019).

Figure 2. Effects of iHeps on iMacs. A) Principal Component Analysis of bulk RNAseq data from Day 0 iMacs, Day 3 co-iMacs, Day 7 co-iMacs, Day 7 iMacs and Day 7 cond-iMacs. B) Heatmap showing the top 5 differentially and uniquely expressed genes from culturing iMacs alone, with conditioned media (cond-iMacs) and after co-culture with iHepatocytes (Day 7 co-iMacs). C) Volcano Plot showing differentially-expressed genes between Day 0 iMacs and Day 7 co-iMacs. D) Differentially expressed upregulated pathways between Day 0 iMacs and Day 7 co-iMacs. E) Volcano Plot showing differentially-expressed genes between Day 7 cond-iMacs and Day 7 co-iMacs. F) Differentially expressed upregulated pathways between Day 7 cond-iMacs and Day 7 co-iMacs. G) Gene expression levels of *LYVE1*, *ID1*, *ID3*, *RXRA*, *HBEGF* and *OSM* in Day 0 iMacs, Day 3 co-iMacs, Day 7 co-iMacs, Day 7 iMacs and Day 7 cond-iMacs. Student's T test was applied. *p<0.05, **p<0.01, ***p<0.001

A

B

C

Supplementary Figure 2: Correlation of iMacs with embryonic liver monocytes and macrophages, and pathway analysis of the DEGs. A) Pearson correlation of iMacs under control, conditioned media or co-culture conditions against embryonic liver monocytes and macrophages from Bian et al, (2020).
B) Pathway analysis of DEGs between Day 7 conditioned media iMacs and embryonic liver macrophages.
C) Pathway analysis of DEGs between Day 7 co-cultured iMacs and embryonic liver macrophages.

Figure 3. Gene expression of iMac and iHep markers in co-cultures. A) Expression of macrophage markers (left panel) and KC-specific markers (right panel) on day 3 and day 7 of co-culture. B) Expression of macrophage markers (left panel) and KC-specific markers (right panel) on day 7 of co-culture compared to iMac cultured in PHCM. C) Expression of hepatic markers (left panel) on day 3 and day 7 of co-culture. * $p < 0.05$.

Figure 4. Changes in IL-6 level upon treatment with 7 paradigm compounds when iMac-derived KCs were used. Cytokine production was assessed in iMac-derived iKC and iHep co-culture treated with the drug stated in the plot title. Data is expressed as the percentage of the LPS-treated vehicle control. Error bars represented S.D., n=3. One-way ANOVA was applied. *p<0.05, **p<0.01 between treatment and vehicle control.

Figure 5. Changes in IL-6 level upon treatment with 7 paradigm compounds without iMac-derived KCs. Cytokine production was assessed in blood monocyte-derived macrophages and iHep co-culture treated with the drug stated in the plot title. Data is expressed as the percentage of the LPS-treated vehicle control. Error bars represented S.D., n=3. One-way ANOVA was applied. *p<0.05, **p<0.01 between treatment and vehicle control.