Variations in H_2 thresholds and growth yields among hydrogenotrophic methanogens relate to different energy conservation strategies

Timothé Philippon [1], Jo Philips [1,2]*

[1]   Department of Biological and Chemical Engineering, University of Aarhus,

Gustav Wieds Vej 10, 8000 Aarhus C, Denmark

[2]   Novo Nordisk Foundation CO_2 Research Center (CORC), Gustav Wieds Vej 10,

8000 Aarhus C, Denmark

*Corresponding author: jo.philips@bce.au.dk

Abstract

Hydrogenotrophic methanogens are of high environmental and biotechnological relevance, as they convert CO_2 with H_2 into CH_4. Despite their common metabolism, variations among these methanogens likely exist. This study therefore determined the H_2 thresholds and growth yields of nine different hydrogenotrophic methanogens. The H_2 threshold, i.e. the H_2 partial pressure at which H_2 consumption halts, ranged over two orders of magnitude from 1.0 ± 0.5 Pa for *Methanobrevibacter arboriphilus* to 120 ± 10 Pa for *Methanosarcina mazei*. Growth yields ranged from 0.51 ± 0.28 $g_{DCW} \cdot (mol\ CH_4)^{-1}$ for *Methanococcus maripaludis* to 5.28 ± 1.25 $g_{DCW} \cdot (mol\ CH_4)^{-1}$. In addition, ATP gains, estimated from both the H_2 threshold and growth yield, correlated reasonably well, suggesting that these variations are due to differences in energy conservation strategies. Our results fitted with the classification previously proposed by Thauer[1], dividing hydrogenotrophic methanogens into two distinct groups: methanogens using cytochromes for energy conservation had a high H_2 threshold (> 20 Pa) and high growth yield (> 4.0 $g_{DCW} \cdot (mol\ CH_4)^{-1}$), whereas all methanogens without cytochromes had a low H_2 threshold (< 8 Pa) and low growth yield (< 1.7 $g_{DCW} \cdot (mol\ CH_4)^{-1}$). Moreover, our H_2 thresholds showed significant additional variations within both groups. Overall, this study found strong variations between hydrogenotrophic methanogens, which are important to understand their environmental prevalence and biotechnological applicability.

Introduction

Hydrogenotrophic methanogenesis is a metabolic pathway used by specific archaeal microbes to produce methane (CH_4) from carbon dioxide (CO_2) and dihydrogen (H_2) for the purpose of energy conservation through the following reaction:

$$CO_2 + 4\,H_2 + n\,ADP + n\,Pi \rightarrow CH_4 + 2\,H_2O + n\,ATP \qquad (1)$$

This metabolism is widely spread in the environment and plays an important role in the global carbon cycle, as methanogens are estimated to convert yearly about one gigaton of carbon into methane [1]. This production of methane is a major issue in the context of climate change, as methane is a potent greenhouse gas. Methanogenesis can also negatively impact steel infrastructure, as certain hydrogenotrophic methanogens induce strong corrosion [2]. Nevertheless, hydrogenotrophic methanogenesis is of huge interest for the development of energy conversion (e.g. power-to-gas) biotechnologies, as methane is a valuable fuel. For instance, bio-methanation is an already advanced technology, converting CO_2-containing gases into methane with the addition of H_2 gas (Reaction 1) [3]. Another notable technology based on hydrogenotrophic methanogens is electro-methanogenesis, in which an electrode is used to generate H_2 *in situ* of a bioreactor [4; 5]. Although the latter technology remains in the early stages of development, its potential is considerable [6; 7]. Because of the technological and environmental importance of methanogenesis, a deep understanding of its metabolism and insights into variations between different methanogenic species are crucial.

All hydrogenotrophic methanogens use a common pathway (Figure 1), consisting of a succession of steps reducing CO_2 to CH_4, using electrons obtained from the oxidation of H_2 and carried by electron shuttles, such as the redox cofactor F_{420}. Besides this common pathway, hydrogenotrophic methanogens vary in the involvement of cytochromes [1]. Two identified protein families named *vht* (or *vho*) and *hdrDE* (containing cytochromes) are only present in microbes that perform cytochromes dependent methanogenesis (Figure 1) [8].

Thauer [1] explained that the presence or absence of those cytochromes affects the amount of ATP (ATP gain or *n*) generated in Reaction 1, with a higher ATP gain when those cytochromes are present (Table 1). Different ATP gains in turn affect various metabolic

characteristics of methanogens, including their H_2 threshold and growth yield (Table 1).

The H_2 threshold quantifies the H_2 concentration at which H_2 oxidation and therefore

methanogenesis halts due to thermodynamic constraints. According to Thauer [1], the

H_2 threshold is lower for methanogens without cytochromes in comparison to

methanogens with cytochromes (Table 1). Accordingly, the growth yield, i.e. the amount

of biomass created per amount of H_2 and CO_2 consumed or CH_4 produced, is expected

to be higher with than without cytochromes (Table 1). So far, however, it remains unknown

whether additional variations in these metabolic characteristics exist, since a coherent

comparison of an extensive number of methanogens has not yet been performed.

Thauer had based his growth yield values on just few studies, studying each only one to

two methanogenic species (e.g. *Methanosarcina barkeri* [9], *Methanobrevibacter*

arboriphilus [10] or *Methanobacterium bryantii* [11; 12]). It is unclear whether these

studies resulted in comparable results, as different growth media and methods were

used. Concerning the H_2 threshold, various studies have been conducted on one to few

methanogens, while only one study compared four different methanogens, but could not

measure H_2 thresholds below 10 ppm [13]. Feldewert has compiled H_2 thresholds for

methanogens obtained by various studies [14], but as the H_2 threshold likely depends on

the experimental conditions [15], it is unsure if these values are comparable. Moreover,

recently strongly different H_2 thresholds were unexpectedly found for acetogens [16]. It

seems thus worthwhile revisiting the H_2 thresholds and growth yields of different

methanogens.

Furthermore, insights into the H_2 thresholds of different methanogens could be crucial to

assess their potential for electro-methanogenesis and role in microbial-induced

corrosion. In both cases, whether involving a cathode as in electro-methanogenesis or

steel as during corrosion, these surfaces act as extracellular electron donors, from which

methanogens can withdraw electrons, using H_2 as main intermediate [17]. Not all

hydrogenotrophic methanogens, however, are as effective in this extracellular electron

uptake [18]. Besides the involvement of extracellular hydrogenases [19], the H_2 partial

pressure was proposed to play a role in variations in electroactivity [15]. Microbes

maintaining low H_2 partial pressures likely favor H_2 evolution from the cathode or steel

surface, thereby increasing extracellular electron uptake rates. As the H_2 threshold

reflects the lowest H_2 partial pressure a strain can sustain, this parameter is thus likely

crucial to understand variations in electroactivity among methanogens.

This study therefore aimed to compare H_2 thresholds and growth yields of nine different

strains of mesophilic hydrogenotrophic methanogenic species in similar conditions

(Table 2). The selected methanogens include species with and without cytochromes,

originating from different environments and spanning various methanogenic genera. In

addition, two methanogenic strains known for their strong corrosive behavior were

included. For all these strains, the H_2 threshold, the growth yield, as well as the

assimilated CO_2 fraction, were determined. Moreover, ATP gain estimates based on either

the H_2 threshold or the growth yield were calculated and compared. All these different

variables are relevant to assess variations in energy conservation strategies among the

selected methanogens, as well as to evaluate their potential for extracellular electron

uptake.

Material and methods

Methanogenic strains and culture conditions

This study included nine different methanogens (Table 2). Seven methanogenic strains were ordered from the Leibniz Institute DSMZ, including *Methanococcus maripaludis* (DSMZ 2067 designated as type strain in this study), *Methanolacinia petrolearia* (DSMZ 11571), *Methanobacterium palustre* (DSMZ 3108), *Methanobacterium formicicum* (DSMZ 1535), *Methanosarcina mazei* (DSMZ 2053), *Methanosarcina barkeri* (DSMZ 800) and *Methanobrevibacter arboriphilus* (DSMZ 744). Once received, the strains were transferred to their recommended DSMZ medium (exact composition but without Na_2S, sludge fluid or fatty acid mixture) and stored frozen at – 80 °C in a 2 mL tube with 25% glycerol.

In addition, a culture containing *Methanobacterium* strain IM1 [20; 21] was obtained from Florin Musat (Aarhus University). The culture was inoculated in DSMZ medium 141 (without Na_2S) and was stored frozen at – 80 °C, as described for the other strains. Microscope imaging (Leica ICC50 W, Germany) showed that an additional coccus shaped microbe was present in this culture. Nevertheless, when cultivated in a medium with only H_2 as electron donor (no acetate), the IM1 strain (bacille shape [21]) became dominant. Acetate was thus further omitted from the medium. Therefore, we assume that the measured H_2 threshold was for *Methanobacterium* strain IM1.

Methanococcus maripaludis Mic1c10, was provided as living culture by Amelia-Elena Rotaru (Southern University of Denmark). This strain was cultured in DSMZ medium 141 (recommended for *Methanococcus maripaludis*, but without Na_2S) and stored at – 80 °C. Both *Methanococcus maripaludis* Mic1c10 and *Methanobacterium* strain IM1 are known corrosive methanogens, as previously documented [20; 22; 19; 23].

To start experiments, frozen stocks were revived in their respective DSMZ medium and transferred to the same medium once. When the late exponential growth phase was reached, 10 mL of culture was transferred to a sterile glass tube with conical bottom, which had first been hermetically sealed using a rubber stopper and flushed with N_2. These tubes were then centrifuged at 2500 rpm for 15 minutes. The cell pellet was washed two times using limited medium (described below), before being resuspended in

the same limited medium. For the H_2 threshold experiment, this cell suspension was diluted with limited medium until an optical density (OD) between 0.10 and 0.15 was reached. It must be noted that this procedure was used both for strains that grow planktonically or as aggregates (notably *M. barkeri* and *M. mazei*), even though similar OD values in this case likely do not correspond to comparable biomass densities.

The limited medium was derived from the DSMZ medium 141 and included 0.34 g of KCl, 3.45 g of $MgSO_4$ x 7 H_2O, 4 g of $MgCl_2$, 0.25 g of NH_4Cl, 0.14 g of $CaCl_2$, 0.14 g of K_2HPO_4, 0.2 mg of $Fe(NH_4)_2(SO_4)_2$, 0.2 g yeast extract powder (Y1625), 10 mL of Wolin mineral solution 10 mL of Wolin vitamins solution for a final volume of one liter. In addition, NaCl was added for a final concentration of 0.4 g per L or 18 g per L, respectively for freshwater methanogens (*M. palustre, M. formicicum, M. barkeri, M. arboriphilus* and *M. mazei*) and marine methanogens (*strain IM1*, *M. maripaludis* DSMZ 2067 and Mic1c10 strain and *M. petrolearia*) (Table 2). After autoclaving this basal medium, filter-sterile stocks solutions were added for a final concentration of 5 g of $NaHCO_3$ and 0.5 $g.L^{-1}$ of L-cysteine.

Measurement of the H_2 threshold

The H_2 threshold was measured for all methanogenic strains. A volume of 80 mL of the cell suspensions obtained after washing was brought into 250 mL serum bottles. For each strain, four replicates were prepared, while three additional bottles were filled with 80 mL of limited medium as abiotic controls. The experiment was initiated by flushing the headspace of the bottles with a gas mix composed of 2 % H_2, 78 % N_2 and 20 % CO_2. This mix was prepared using gas flow meters (Alicat®, MCW Series, USA) with two gas tanks, one containing 80% H_2 and 20% CO_2 and the other containing 80% N_2 and 20 % CO_2. Each serum bottle was pressurized for a total pressure of 2 bars (or 1 bar overpressure) of this gas mix. The bottles were incubated upright at 30 °C with horizontal orbital shaking at 100 rpm. The total pressure was measured with a digital manometer (OMEGA DPG108, UK) before each sampling event. The headspace of each bottle was frequently sampled with a syringe for H_2 analysis (1.5 mL). The H_2 threshold was considered reached when the H_2 partial pressure did not significantly decrease for at least four days with measurement intervals of maximally two days. The value of the H_2 threshold was calculated from the average of two to four measurements performed during those last four days. After the threshold was reached, the headspace was flushed and pressurized again with the gas

169 mix as described above. The headspace was sampled again until the H_2 threshold was

170 reached for a second time. The OD and the methane content in the headspace were

171 measured at the end of the experiment. For all methanogens, the amount of CH_4 at the

172 end of the H_2 threshold experiment was too low to be accurately quantified, even though

173 detectable by GC.

Growth yield measurements

In addition to the H_2 threshold, the growth yield of all methanogenic strains was

determined. For each methanogen, the inoculum was pre-grown and washed as

described above. Four replicated bottles (250 mL) with 80 mL of limited medium were

inoculated by injecting 0.5 mL of washed culture. These bottles were pressurized at 2

bars (overpressurize of 1 bar) using a gas mix with 80% H_2 and 20 % CO_2. The composition

of the headspace was quantified, and the concentration of dissolved inorganic carbon

(DIC) in the liquid phase was measured before the start of the incubation. The bottles

were incubated upright at 30 °C under horizontal and orbital shaking at 100 rpm. The

pressure was measured daily until at least one of the four replicates reached a negative

pressure, after which the incubation was ended for all replicates. Before biomass was

harvested, the analysis of the headspace composition and determination of the DIC

concentration were repeated. Next, the bottles were opened and 10 mL of each culture

was transferred into a 20 mL glass tube with conical bottom and centrifuged at 3000 rpm

for 15 minutes. The clear supernatant was discarded and an additional 10 mL of culture

was placed in the same tube, while keeping the cell pellet. This centrifugation procedure

was repeated until all 80 mL of each bottle was harvested. Each of the tubes containing

the stacked cell pellets were then dried at 100 °C for 48 hours, cooled down for one hour

in a desiccator and the dry weight was measured using an analytical balance (± 0.0001 g

precision) (VWR, LD17 4XN). The growth yield was calculated by dividing the dried cell

weight by the quantity of methane measured in the headspace at the end of the

experiment.

Analytical methods

Gaseous samples were analyzed for H_2, CH_4 and CO_2 using a Compact GC 4.0

(Interscience, Netherlands) with two different columns: one Molsieve 5A 30 m x 0.32 mm

and one Rt-QBond 3 m x 0.32 mm, using helium as gas carrier and a TCD detector [16].

Calibration was performed using calibrating gases mixed with N_2. The quantification limit

was 1 Pa or 10 ppm for H_2, 10 000 ppm for CH_4 and 50 ppm for CO_2.

For H_2 partial pressures below 50 ppm, a Peak Performer 1 (PP1) (Peak Laboratories, USA)

GC unit based on HgO–Hg conversion with a reducing compound photometer (RCP) was

also used. This GC measures trace H_2 levels (0.1–50 ppm of H_2) and has quantification

limit well below that of the CompactGC. Partial pressures were calculated by multiplying

the mol fraction, as measured by GC, with the total pressure. Dissolved concentrations

were calculated from the partial pressures using the Henry constants for H_2 at 30°C (7.8

x 10^{-4} mol.l^{-1}.bar^{-1}) [24].

Liquid samples were prepared for DIC analysis, as previously described [25]. A sample of

0.5 mL was mixed with 0.85 mL of 800 mM HCl in 20 mL closed Hungate tubes, previously

flushed with N_2 and kept at atmospheric pressure. The acidified sample was then

vortexed for 2 minutes and left to equilibrate for eight hours to convert bicarbonate into

CO_2. The CO_2 level in the headspace of the tube was then measured as described above.

A calibration curve was prepared prior to the experiment using sodium bicarbonate.

Optical densities (OD) were measured at 600 nm using a Denovix DS-11+

spectrophotometer.

Calculations

Estimation of the ATP gain from the H_2 threshold

As explained previously [16], H_2 thresholds can be used to calculate an estimate for the

ATP gain through calculating the critical Gibbs free energy, i.e. the Gibbs free energy

change when the H_2 threshold is reached [1].

Considering hydrogenotrophic methanogenesis:

223                    $$CO_2 + 4\,H_2 \rightarrow CH_4 + 2\,H_2O \qquad (2)$$

224   The critical Gibbs free energy (ΔG_c) becomes:

225           $$\Delta G_c = \Delta G^{0'}_{\text{methanogenesis}} + R \cdot T \cdot \ln \frac{p_{CH_4}}{p_{CO_2} * \theta_{H_2}{}^4} \qquad (3)$$

With $\Delta G^{0'}_{\text{methanogenesis}}$ the Gibbs free energy of methanogenesis (Reaction 2) in

physiological standard conditions (-131 kJ.mol^{-1} [1]), θ_{H_2} the H$_2$ threshold (in atm), p_{CO_2}

the partial pressure of CO$_2$ (in atm) and p_{CH_4} the partial pressure of methane (in atm) at

the H$_2$ threshold. As the methane partial pressures in the H$_2$ threshold experiment were

too low to be quantified, they were calculated from the amount of H$_2$ consumed,

according to the stoichiometric ratios of Reaction 2.

The critical Gibbs free energy at the H$_2$ threshold is not zero, because methanogenesis

should be considered as coupled to ADP to ATP phosphorylation (Reaction 1). At the H$_2$

threshold, it can be assumed that the total Gibbs free energy change of Reaction 1

considering both methanogenesis and ADP phosphorylation, equals zero [1]:

$$\Delta G_c + n \cdot \Delta G_{ATP} = 0 \tag{4}$$

Where ΔG_{ATP} is the Gibbs free energy change for the phosphorylation of ADP into ATP.

The value of this ΔG_{ATP} can vary between microorganisms and different values are used

in literature [1; 16], but here we used the value of -75 kJ.mol^{-1} [26]. The ATP gain n, i.e. the

number of ADP molecules phosphorylated for each completed methanogenesis

reaction, can finally be calculated using:

$$n = -\frac{\Delta G_c}{\Delta G_{ATP}} \tag{5}$$

Estimation the ATP gain from the growth yield

In parallel to the calculations from the H$_2$ thresholds, the ATP gain was also estimated

from the experimentally measured growth yields. Spormann [27] described a relationship

between the growth yield expressed per catabolic product (in our case methane) (Y^{CH_4},

g dry cell weight per mole of methane) and the growth yield expressed per ATP (Y^{ATP}, g

dry cell weight per mole ATP). This relationship can be used to estimate the ATP gain n

(mole ATP per mole methane):

$$n = \frac{Y^{CH_4}}{Y^{ATP}} \tag{6}$$

Y^{ATP} was estimated to be 5 g dry cell weight per mole of ATP for an anabolic metabolism

using CO$_2$ as carbon source [27].

Estimation of the fraction of assimilated carbon

The growth yield experiment also allowed to compare the fraction of CO_2 assimilated into biomass among the different methanogens. The total amount of inorganic carbon consumed by the methanogens was calculated from the headspace CO_2 levels and the DIC concentrations measured at the beginning and the end of the experiment. It was further assumed that methanogenic biomass has the molecular formula of $C_{4.9}H_{9.4}O_{2.9}$ N (molar weight of 129 g per mole), as proposed by Rittmann and McCarty [28]. The amount of moles of assimilated carbon was then calculated from the dry cell weight measured at the end of the experiment.

The fraction of assimilated carbon ($f_{C,biomass}$) was then obtained by dividing the moles of carbon assimilated in biomass ($n_{C,biomass}$) with the total moles of consumed CO_2 ($n_{C,total}$):

$$f_{C,biomass} = \frac{n_{C,biomass}}{n_{C,total}} \cdot 100\% \tag{7}$$

Statistical Analysis

Welch ANOVA test

Welch-Anova tests were performed to differentiate statistically significant groups using both GraphPad Prism® and RStudio (ver. 4.4.3). The Welch test was chosen because the variances were considered too different to perform a regular ANOVA test. All the tests were performed with $\alpha = 0.05$ and the p-value was set at 0.05 for all tests.

Error propagation

Standard deviations on the experimental results were calculated using Excel and were further propagated through the different calculations using available calculators [29; 30].

Results

H$_2$ threshold measurements

In this study, the H$_2$ threshold of nine different methanogenic strains was measured, as the H$_2$ partial pressure at which their H$_2$ consumption halts. In all abiotic controls, the H$_2$ partial pressure remained almost stable for the entire duration of the experiment (Figure 2, Figure S1), as only a small drop of the total pressure was observed, as the result of headspace sampling (Figure S1).

For all inoculated bottles, a clear drop in the H$_2$ partial pressure was observed (Figure 2), confirming H$_2$ consumption by the different methanogens. Moreover, H$_2$ consumption halted at a certain H$_2$ partial pressure for each strain. To confirm these H$_2$ thresholds, H$_2$ was reinjected, once stable H$_2$ partial pressures were reached. For most methanogenic strains, the H$_2$ thresholds reached after the re-addition of H$_2$ were comparable to those before the re-addition, except for *Methanosarcina mazei* and *Methanosarcina barkeri* (Figure S2). *M. mazei* consumed H$_2$ after the H$_2$ re-addition, but its H$_2$ consumption halted at a higher H$_2$ partial pressure (260 ± 49 Pa versus 120 ± 10 Pa before the re-addition). In contrast, *M. barkeri* no longer consumed H$_2$ after the re-addition. These findings are possibly explained by the exhaustion of these strains over the long incubation time (*M. mazei* reaches its H$_2$ threshold after 12 days of incubation, while *M. barkeri* required more than 28 days), while all other methanogens, required about five days or less to reach a H$_2$ partial pressure close to their H$_2$ threshold.

The three species belonging to the *Methanobacterium* genus displayed similar low H$_2$ thresholds of around 2 Pa (Figure 2, Figure 3, Table 2). Also, *Methanosarcina petrolearia* and *Methanobrevibacter arboriphilus* reached H$_2$ thresholds in this range. *M. arboriphilus* was the only species that reached a H$_2$ partial pressure of 1 Pa (Figure 2, Figure 3, Table 2). The Welch-ANOVA test gathered these five strains with the lowest H$_2$ thresholds in one statistical group (Figure 3). In contrast, all the other strains had higher H$_2$ thresholds and were classified as each a significantly different group (Figure 3). The two strains of *Methanococcus maripaludis* displayed H$_2$ thresholds slightly higher than the five strains with the lowest H$_2$ thresholds (Figure 2, Figure 3, Table 2). Interestingly, the corrosive strain *Mic1c10* had a significantly lower H$_2$ threshold (around 4 ± 1 Pa) than the type strain

(7 ± 1 Pa). The *Methanosarcina* species differed strongly in their H_2 thresholds, as *M.*

barkeri had a H_2 threshold of 21 Pa ± 2 Pa, while the highest H_2 threshold was recorded for

M. mazei (120 ± 10 Pa).

No significant growth was observed for any of the inoculated bottles over the course of

the experiment, as OD_{600} values remained stable (less than 0.1 increase). For the two

Methanosarcina strains, which grow both planktonically and as aggregates, the stable

OD_{600} cannot be used to claim the absence of growth, but as these strains consumed the

least H_2 (highest H_2 thresholds), their growth was likely limited. Methane was detected in

the headspace of the inoculated bottles at the end of the experiment, but its levels were

too low for an accurate quantification.

Growth yield measurements and estimation of the carbon assimilated as biomass

In parallel of the H_2 threshold study, all the methanogenic strains were grown in bottles

on 80% H_2 and 20% CO_2 to determine their growth yields, as well as their assimilated CO_2

percentage. In all these bottles methane levels significantly increased over the

incubation period, while H_2 and CO_2 were consumed and cell growth occurred (Table S1).

These experimental results were used to calculate the growth yield, i.e. the increase of

biomass expressed as gram dry cell weight (g_{DCW}) versus the amount of methane

produced. In addition, the assimilated CO_2 fraction, i.e. fraction of CO_2 assimilated in

biomass versus total amount of consumed CO_2, was calculated (Figure 4, Table 2).

Relatively high variabilities between the replicates were obtained for the growth yields

and the assimilated CO_2 percentages (large error bars in Figure 4). This is likely due to the

low total weight of dried biomass (in mg) obtained during the growth yield experiment

(Table S1), limiting the accuracy of this measurement. In addition, the centrifugation

steps were possibly not always as effective in removing cells from the culture

suspension, which could also explain the high variability between the replicates.

Despite the high variability in the dry cell weight, Welch-ANOVA tests identified two

significantly different groups in the growth yields (Figure 4, right). The two

Methanosarcina strains form one group, with high growth yields of 4.09 ± 1.16 and 5.28 ±

1.25 g_{DCW}/mole of CH_4 for *M. barkeri* and *M. mazei* respectively. All other methanogens

had significantly lower growth yields, ranging from 0.51 ± 0.28 g_{DCW}/mole of CH_4 (for *M.*

maripaludis type strain) to 1.69 ± 0.58 g_{DCW}/mole of CH_4 (for *M. formicicum*). Growth

yields were also expressed as g_{DCW} per mole H_2 or CO_2 consumed (Figure S3), leading two

the same statistical groups. Moreover, the same two significantly different groups were

also identified for the fraction of assimilated CO_2 percentages (Figure 4, right) with *M.*

mazei and *M. barkeri* having a high assimilation percentage, i.e. 22.7 ± 10.0 % for *M. mazei*

and t18.9 ± 3.0 % for and *M. barkeri*. All other methanogens displayed a significant lower

assimilation percentage, which was 7.15 ± 3.18% for *M. formicicum* and even lower for

the other strains.

Comparison ATP gains estimated from either the H_2 threshold or the growth yield

Both the H_2 threshold and the growth yield allow to estimate the ATP gain, i.e. the number

of ATP molecules obtained for one complete methanogenesis reaction (Reaction 1).

Using the H_2 threshold, the ATP gain estimates ranged from 0.32 ± 0.07 for *M. arboriphilus*

to 0.97 ± 0.01 for *M. mazei* (Table 2). Estimating ATP gains from the growth yields gave a

slightly wider range, starting at 0.10 ± 0.03 for *M. maripaludis* up to 1.06 ± 0.12 for *M.*

mazei. Despite the totally different experimental procedures and calculation methods,

both estimates thus led to a largely overlapping range, validating our experimental

approach. These ATP gains estimates were further correlated to each other, leading to a

clear but not perfect correlation (Figure 5, R^2 of 0.84). Only the two *Methanosarcina*

species, *M. mazei* and *M. barkeri,* fall on the 1:1 correlation line, while the correlation is

less clear for the other strains. For these other strains, the difference in ATP gain ranges

from 0.1 for *M. palustre* to 0.45 for *Methanococcus maripaludis type strain*. Moreover, for

these methanogens, higher ATP gains were calculated from the H_2 thresholds than from

the growth yield, demonstrating that there were discrepancies between our methods,

possibly because the growth yields were somewhat underestimated. This illustrates that

our ATP gains should be considered as estimates, rather than absolute values.

Nevertheless, the clear correlation (Figure 5) does suggest that the ATP gains differed

among the tested strains, reflecting differences in their energy conservation strategies.

Discussion

Variations in the H_2 thresholds and growth yields among methanogens correspond to different energy conservation strategies

This work evaluated the H_2 threshold, growth yield and assimilated CO_2 fraction for nine different methanogenic strains. Experimental H_2 thresholds varied two orders of magnitudes among methanogens, ranging from 1.0 ± 0.5 Pa for *M. arboriphilus* to 120 ± 10 Pa for *M. mazei* (Figure 3, Table 2), which is a similar range as previously recorded [31]. Recorded H_2 thresholds for hydrogenotrophic methanogens are usually not lower than 1 Pa [14], as in our study (Table 2). Only strict methylotrophic methanogens, which use H_2 for the reduction of methyl compounds (in contrast to CO_2 reduction as for hydrogenotrophic methanogens) have H_2 threshold as low as 0.1 Pa [14]. Intriguingly, widely varying H_2 thresholds were previously reported for *M. barkeri* (19 Pa to 150 Pa) [14; 13; 31]. Our experimental H_2 threshold for this strain (21 ± 2 Pa, Table 2) agrees with the lowest reported value. These variations in reported values are possibly due to different experimental conditions (e.g. different medium composition) [15]. In addition, the slow H_2 consumption by this specific strain (Figure 2) likely makes it hard to conclude when the H_2 threshold is reached. Here, we found that after the halt of H_2 consumption, *M. barkeri* could not be reactivated, while all other methanogens could (Fig S2). Our H_2 threshold for *M. barkeri* could thus still be somewhat overestimated. For *M. mazei*, we measured a six times higher H_2 threshold (120 ± 10 Pa, Table 2) than for *M. barkeri*. *M. mazei* was still active after the H_2 re-addition, so its significantly higher H_2 threshold is not due to an experimental artefact.

Overall, the experimental H_2 thresholds found in this study agree with the classification proposed by Thauer (Table 1), as we found H_2 thresholds below 10 Pa for all methanogens without cytochromes, while methanogens with cytochromes had H_2 thresholds above 10 Pa (Table 2). Nevertheless, in both these groups, we found methanogens with statistically different H_2 thresholds (Figure 3), showing that additional variations in H_2 thresholds exist.

Experimental growth yields varied one order of magnitude among methanogens, ranging from 0.51 ± 0.28 g_{DCW}/mol CH_4 for *M. maripaludis* to 5.28 ± 1.25 g_{DCW}/mol CH_4 for *M.*

mazei. These growth yields are coherent with what can be found in literature [1; 31]. For

instance, Weimer *et al.* reported a growth yield of 6.4 g_{DCW}/mol CH_4 for *M. barkeri*, which

is comparable to our growth yield (4.09 ± 1.16 g_{DCW}/mol CH_4, Table 2) [9]. Likewise, our

growth yield for *M. arboriphilus* (0.71 ± 0.41 g_{DCW}/mol CH_4) is comparable to the value

recorded by Morii et al. (up to 1.42 g_{DCW}/mole of CH_4) [10]. These values do not match

perfectly, possibly because growth yields of methanogens depend on the H_2 supply rate

[32; 33]. This again highlights the need to compare strains in similar experimental

conditions, as in our study.

Similar as for the H_2 thresholds, our experimental growth yields agree with the

classification proposed by Thauer (Table 1). We indeed found growth yields lower than 3

g_{DCW}/mol CH_4 for all methanogens without cytochromes, while the two *Methanosarcina*

species (with cytochromes) had growth yields above 3 g_{DCW}/mol CH_4 (Table 2). However,

no further significant differences in growth yields were identified in these two groups

(Figure 4, left), as was found for the H_2 thresholds (Figure 3). Even for *M. mazei* and *M.*

barkeri, which differ six times in their H_2 threshold, no significant different growth yields

were found (Figure 4). It is likely that this incoherence is due to the high variability of the

growth yield measurement (high error bars in Figure 4). Such a high variability is common

for growth yield measurements, as it was also observed in previous studies determining

methanogenic growth yield [34-37].

Thauer [1] did not evaluate possible differences in the fraction of CO_2 assimilated into

biomass between methanogens with and without cytochromes. In this study, we did

observe two statistically different groups in the assimilated CO_2 fraction, which

correlated perfectly with the presence or absence of cytochromes (Figure 4, right).

Methanogens without cytochromes assimilated a low fraction of CO_2 (ranging from 1.68

± 0.90% to 7.15 ± 3.18%). In contrast, methanogens with cytochromes assimilated a

higher fraction of the converted CO_2 (ranging from 18.97 ± 3.06% to 22.76 ± 10.04%). This

difference is not unexpected, as the assimilated CO_2 fraction is closely linked to the

growth yield.

The experimental H_2 thresholds and growth yields were furtherused to calculate

estimates for the ATP gain, to thereby evaluate differences in energy conservation among

the tested methanogens. Despite the two totally different experimental approaches and

the various assumptions underlying these calculations (Equations 5 and 6), an overlapping range and decent correlation was obtained between the two types of ATP estimates (Table 2, Figure 5), validating our approach.

Thauer [1] previously estimated an ATP gain of 1.5 ATP per mol CH_4 for *Methanosarcina barkeri* (methanogen with cytochrome), while we calculated this to be 0.75 ± 0.01 ATP per mol CH_4, when derived from the H_2 threshold. However, Thauer used a ΔG_{ATP} of - 50 kJ per mol, while we chose to use a ΔG_{ATP} of - 75 kJ per mole, as this also includes heat loss [26]. According to Thauer (Table 1), methanogens without cytochromes have an ATP gain of 0.5 per mole of CH_4, which corresponds with the range of ATP gain estimates (between 0.10 and 0.56 mol ATP per mol CH_4) determined here for methanogens without cytochromes (Table 2).

In addition to the clear distinctions between methanogens with and without cytochromes, our ATP gain estimates calculated from the H_2 thresholds suggest that additional variations in energy conservation exist, as additional significant differences in H_2 thresholds were recorded in both the groups of methanogens (Figure 3). Unfortunately, this was not confirmed by the ATP gains derived from the growth yields, again due to the high variability on the growth yield measurements. In conclusion, the presence or absence of cytochromes (use of hdrDE and vhtAGC, versus the use of hdrABC, Figure 1) thus represents two totally different mechanisms for energy conservation in hydrogenotrophic methanogens. Further minor variations likely exist in these energy conservation mechanisms, as reflected by our experimental H_2 thresholds. Further genetic and biochemical analyses will be required to unravel these variations in more detail.

Link between the H_2 threshold and extracellular electron uptake by methanogens

The H_2 threshold of methanogens is possibly a crucial parameter to understand variations in electroactivity among methanogens, as the rate of extracellular electron uptake from a cathode or steel surface is thought to depend on the H_2 pressure maintained by microbial H_2 consumption [15]. The experimental H_2 thresholds obtained

in this study (Table 2, Figure 3) thus allow to reevaluate previously reported differences in

electroactivity among methanogens.

Mayer et al. studied the electro-methanogenesis performance of several methanogens.

M. maripaludis and *M. petrolearia*, for which a low H_2 threshold was found here (Figure

3), increased cathodic currents (i.e. key signature of extracellular electron uptake), just

as several other methanogens without cytochromes. In contrast, *M. mazei*, for which the

highest H_2 threshold was recorded (Figure 3), was incapable of electro-methanogenesis

[18]. Similarly, Jaramillo *et al*. investigate the corrosion potential of different

methanogens [38]. Non-hydrogenotrophic methanogens were reported as non-

corrosive, confirming the important role of H_2 as intermediate in microbial corrosion. *M.*

mazei and *M. barkeri* (high H_2 threshold) were found to be mild corrosive, whereas

Methanobacterium spp. (low H_2 threshold) were classified as highly corrosive. Overall,

these correlations thus suggest that methanogens with a low H_2 threshold have a better

ability for extracellular electron uptake than methanogens with a high H_2 threshold.

Nevertheless, several other variables likely affect the extracellular electron uptake rate

of methanogens, in addition to the H_2 threshold. For instance, the H_2 pressure maintained

at the cathode or steel surface also depends on the rate of H_2 consumption. Fast H_2

consumption is thus likely also required to maintain low H_2 partial pressures. Moreover,

the H_2 evolution reaction on a cathode or steel surface does not only get stimulated by

low H_2 levels (thermodynamic effect), but also by extracellular enzymes, such as

hydrogenases (kinetic effect). *M. maripaludis* Mic1c10, for instance, has a unique

extracellular hydrogenase anchored to its cell surface [22], causing stronger corrosion

than other *M. maripaludis* strains [22]. Moreover, the cathodic current increase was

double as high for strain Mic1c10 than for a non-corrosive *M. maripaludis* strain [18; 39].

Here, we found that strain Mic1c10 had a significantly lower H_2 threshold than the *M.*

maripaludis wild type (Figure 3, Table 2). Kawachi et al. [22] also recorded that strain

Mic1c10 maintained slightly lower H_2 levels during steel corrosion than by the *M.*

maripaludis type strain. These differences, however, are possibly too small to explain the

strong corrosion rates caused by Mic1c10, which is thus likely mostly due to its

extracellular hydrogenases. Similarly, *Methanobacterium* -IM1 was more corrosive than

other tested methanogens [23], but here we did not find that strain IM1 had a lower H_2

threshold than other tested *Methanobacterium* strains (Figure 2, Table 3).

Methanobacterium IM1 thus likely also has an additional mechanism to enhance

extracellular electron uptake rate, which is likely similar to the one described for *M.*

maripaludis Mic1c10.

Overall, we can conclude that the extracellular electron uptake rates of methanogens are

inversely proportional to their H_2 threshold (lower H_2 thresholds correspond to higher

extracellular electron uptake rates), whereas few methanogens (e.g. *M. maripaludis*

Mic1c10, *Methanobacterium* IM1) have a special additional mechanism to increase

extracellular electron uptake rates.

Ecological and biotechnological implications of variations among methanogens

This study found important variations in the H_2 thresholds and growth yields among

hydrogenotrophic methanogens. We systematically found that methanogens with a low

H_2 threshold have a low growth yield, assimilate a low fraction of CO_2 and have a low ATP

gain (Table 2). In contrast, methanogens with a high H_2 threshold have a high growth yield,

assimilate a larger carbon fraction and have a higher ATP gain (Table 2). This illustrates

that hydrogenotrophic methanogens are confronted with a trade-off between the

capacity to consume H_2 at trace levels (methanogens without cytochromes), or the ability

to grow strongly (high growth yield) with a high energetic efficiency (high ATP gain)

(methanogens with cytochromes). This trade-off allows to understand the ecological

niche and biotechnological relevance of the different types of methanogens. For

instance, Mills *et al.* [40] found that the presence of the *Methanosarcina* genus in electro-

methanogenesis systems was correlated to high H_2 levels (reactors with low coulombic

efficiency), while the *Methanobacterium* genus was rather correlated to low H_2 levels

((high coulombic efficiency).

In addition, a correlation between the growth yield (high ATP gain) and the growth rate of

methanogens would be expected [16]. However, Thauer [1] described that methanogens

with cytochromes (high ATP gain) have a lower growth rate (high doubling time) than

methanogens without cytochromes (low ATP gain, but low doubling times). Comparing

doubling times, however, is possibly not adequate, since methanogens also strongly vary

in cell size (large clustered cells for *Methanosarcina* species versus small different

shaped cells for other methanogens) [41; 42]. Nevertheless, also in the H_2 threshold

experiment of this study, H_2 was faster consumed by methanogens without cytochromes,

than by methanogens with cytochromes (Figure 2). This observation possibly reflects that

Methanosarcina species are not well adapted to consume H_2. Indeed, these

methanogens are not strict hydrogenotrophic, but are also capable of acetoclastic

methanogenesis, in contrast to the other tested methanogens. Methanogens with

cytochromes are thus possibly mainly adapted to use other substrates than H_2 and only

shift to H_2 consumption at high H_2 partial pressures. Overall, future studies comparing

the growth and H_2 consumption rates of a wide range of different methanogens in

comparable conditions would be highly valuable to unravel these differences.

This study only included mesophilic methanogens. A similar comparison as in this study

could also be relevant for thermophilic methanogens, because of their high

biotechnological importance [43]. Their metabolic properties can most likely already be

assessed from the presence or absence of cytochromes in their genome. Interestingly,

Thauer (ref) stated that methanogens without cytochromes dominate at thermophilic

conditions, while methanogens with cytochromes rather occur at lower temperatures.

In this study, we found that H_2 thresholds of methanogens range over two orders of

magnitude (Figure 3). In comparison, we previously found three orders of magnitude

difference in the H_2 thresholds of acetogenic bacteria (6 Pa to 1990 Pa) [16]. It is often

assumed that acetogens cannot compete with methanogens for low H_2 levels, as the

latter have lower H_2 thresholds [44]. Our results, however, demonstrate that such a

competition rather should be evaluated on species level. For instance,

Methanobacterium strains indeed have lower H_2 thresholds than *Acetobacterium* spp,

but acetogenic *Sporomusa* spp. have H_2 thresholds in the range of those of methanogens

without cytochromes and well below those with cytochromes. The outcome of

competition for trace H_2 levels between methanogens and acetogens likely also depends

on the H_2 consumption rates. Recently, H_2 consumption by acetogens was described by

first order kinetics (increasing rates at increasing H_2 levels) [45], while the kinetics of H_2

consumption by methanogens rather follows Monod kinetics with low half-saturation

constants [46] or even zero-order kinetics [47]. Based on these kinetic differences,

methanogens (without cytochromes) consume H_2 likely faster at low H_2 levels, while

acetogens have a faster H_2 consumption at higher H_2 levels, even though H_2 consumption

rates can still widely vary within these groups [45]. Moreover, also the temperature and

pH affect the competition for H_2 between acetogens and methanogens [1; 46; 48; 49; 44;

50-52]. Detailed insights into the differences in H_2 consumption among methanogens

and between methanogens and acetogens are crucial to assess competition for H_2 in the

environment, as well as in mixed culture bioreactor systems, as in anaerobic digestion

MES and bio-methanation.

Conclusion

This study demonstrated that the H_2 thresholds, growth yields and assimilated CO_2

fractions of nine hydrogenotrophic methanogens varied widely. These differences are

related to variations in energy conservation and represent a trade-off among

hydrogenotrophic methanogens. Methanogens without cytochromes sustain

methanogenesis at low H_2 levels but have low growth yields and ATP gains, while

methanogens using cytochromes for energy conservation require higher H_2 levels but

achieve higher growth yields and ATP gains. In addition to this distinction based on

cytochromes, our results suggest additional metabolic variations, warranting further

investigations. Overall, the differences identified here are important to explain the

ecological distributions of hydrogenotrophic methanogens, as well as their ability to

increase microbial induced corrosion and cathodic electron uptake.

Acknowledgement

This study was financed by a Villum Young Investigator grant (VIL53083). We would like to thank Satoshi Kawaichi and Amelia-Elena Rotaru from the University of Southern Denmark for providing the living culture of *Methanococcus maripaludis* Mic1cc10 and Florin Musat from Aarhus University for providing the living culture of *Methanobacterium* IM1. We also acknowledge the Section for Microbiology at Aarhus University for lending their Peak Performer. Heidi Skov Johansen and Laura Mũnoz-Duarte are thanked for their practical support during the experiments and calculations. We also appreciate the advice of Sanne Sandberg Overby on the statistical tests.

Figures

Figure 1: Hydrogenotrophic methanogenesis metabolism. Black colored components and reactions are
common for all hydrogenotrophic methanogens. Red colored components are specific for cytochrome-
dependent methanogens, which have the two proteins complex *hrdDE* (methanophazine-dependent
heterodisulfide reductase) and *vhtAGC* (methanophenazine-reducing hydrogenase). Green colored
components only occur in methanogens without cytochromes using the *hdrABC* complex [8; 53; 54].

Table 1: Comparison of metabolic characteristics of hydrogenotrophic methanogens depending on the

presence or absence of cytochromes according to Thauer [1]:

	Methanogens without cytochromes	Methanogens with cytochromes
H_2 Threshold	Under 10 Pa	Above 10 Pa
Growth yield	Up to 3 g CDW per mol CH_4	Up to 7 g CDW per mol CH_4
ATP gain	0.5 mol ATP per mol CH_4	1.5 mol ATP per mol CH_4
Example of methanogenic orders	*Methanococcus, Methanobrevibacter, Methanobacterium*	*Methanosarcina*

Figure 2: Change of the H_2 partial pressure (Pa) over time in the H_2 threshold experiment for the nine methanogenic species and compared to the abiotic controls. The data points represent the average of four replicates; the error bars show the standard deviations. Only datapoints before the re-addition of H_2 are shown, later datapoints are shown in Figures S1 and S2.

Figure 3: Box plots summarizing the experimental H_2 threshold values for the nine methanogenic strains. The letters indicate statistically different groups, as determined using a Welch-ANOVA test.

Figure 4: (A) Comparison of the growth yields expressed as the increase of gram Dry Cell Weight (g_{DCW}) per mole of methane produced for the different methanogenic strains (B) Comparison of the estimated percentage of CO_2 assimilated into biomass for the different methanogenic strains. The letters indicate statistically different groups, as determined using a Welch-ANOVA test.

Figure 5: Correlation of the ATP gains estimations from the experimental H_2 thresholds versus the ATP gains
estimated from the experimental growth yields. The vertical and horizontal error bars reflect the standard
deviation on these estimates and were propagated from the experimental errors. The red line indicates a
1:1 correlation. The correlation coefficient (R2) was 0.84.

611 Table 2: Overview of the methanogenic strains examined in this study, as well as their experimentally determined H_2 thresholds, growth yields and assimilated CO_2

612 %. The table also includes the ATP gains estimated either from the experimental H_2 thresholds or the experimental growth yields. The H2 thresholds measured are

613 expressed as partial pressures in the headspace (both in ppm and Pa) and as concentration dissolved in the liquid phase (nM).

Methanogenic strain	Cytochromes	Salinity level	H_2 Threshold			Growth yield	Assimilated CO_2 %	ATP gain estimated from H_2 threshold	ATP gain estimated from growth yield
			ppm	Pa	nM	g_{DCW} (mol CH$_4$)$^{-1}$		mol ATP (mol CH$_4$)$^{-1}$	mol ATP (mol CH$_4$)$^{-1}$
Methanobrevibacter arboriphilus	No [55; 56]	Fresh water	7 ± 1	1.0 ± 0.5	8 ± 3	0.71 ± 0.47	2.82 ± 2.30	0.32 ± 0.07	0.14 ± 0.05
Methanolacinia petrolearia	No [55]	Marine water	10 ± 3	2.0 ± 0.5	16 ± 4	0.63 ± 0.31	1.86 ± 0.86	0.41 ± 0.03	0.13 ± 0.03
Methanobacterium palustre	No [55]	Fresh water	11 ± 2	2.0 ± 0.5	16 ± 4	1.57 ± 0.41	6.91 ± 2.15	0.41 ± 0.03	0.31 ± 0.04
Methanobacterium IM1	No [55]*	Marine water	9 ± 1	2.0 ± 0.5	16 ± 4	1.39 ± 0.36	5.12 ± 1.92	0.41 ± 0.03	0.28 ± 0.04
Methanobacterium formicicum	No [55]	Fresh water	10 ± 1	2.0 ± 0.5	16 ± 4	1.69 ± 0.58	7.15 ± 3.18	0.41 ± 0.03	0.34 ± 0.06
Methanococcus maripaludis Mutant strain Mic1c10	No [55; 53]	Marine water	17 ± 3	4.0 ± 1.0	31 ± 8	1.02 ± 0.55	5.27 ± 3.42	0.51 ± 0.03	0.20 ± 0.05
Methanococcus maripaludis Type strain	No [55; 53]	Marine water	33 ± 4	7.0 ± 1.0	47 ± 12	0.51 ± 0.28	1.68 ± 0.90	0.56 ± 0.01	0.10 ± 0.03
Methanosarcina barkeri	Yes [8; 55]	Fresh water	158 ± 18	21 ± 2	195 ± 31	4.09 ± 1.16	22.76 ± 10.04	0.75 ± 0.01	0.82 ± 0.11
Methanosarcina mazei	Yes [8; 55; 56]	Fresh water	720 ± 20	120 ± 10	936 ± 78	5.28 ± 1.25	18.97 ± 3.06	0.97 ± 0.01	1.06 ± 0.12

614 * The assertion that *Methanobacterium* IM1 does not possess cytochrome is based in the fact that no other *Methanobacterium* was found to have cytochromes.

References

[1] Thauer, Kaster, Seedorf, Buckel, and Hedderich (2008) Methanogenic archaea:
ecologically relevant differences in energy conservation. *Nat Rev Microbiol* **6**: 579-591.

[2] An, B.A., Deland, E., Sobol, O., Yao, J.Z., Skovhus, T.L., and Koerdt, A. (2021) The
differences in the corrosion product compositions of Methanogen-induced
microbiologically influenced corrosion (Mi-MIC) between static and dynamic growth
conditions. *Corrosion Science* **180**.

[3] Jensen, M.B., Ottosen, L.D.M., and Kofoed, M.V.W. (2021) H_2 gas-liquid mass transfer:
A key element in biological Power-to-Gas methanation. *Renewable & Sustainable Energy*
Reviews **147**.

[4] Cheng, S.A., Xing, D.F., Call, D.F., and Logan, B.E. (2009) Direct biological conversion
of electrical current into methane by Electromethanogenesis. *Environmental Science &*
Technology **43**: 3953-3958.

[5] Palacios, P.A., Philips, J., Bentien, A., and Kofoed, M.V.W. (2024) Relevance of
extracellular electron uptake mechanisms for electromethanogenesis applications.
Biotechnology Advances **73**.

[6] Pandit (2024) Wired for energy: Electromethanogenesis redefining anaerobic
digestion. *Process Safety and Environmental Protection*.

[7] Blasco-Gómez, R., Batlle-Vilanova, P., Villano, M., Balaguer, M.D., Colprim, J., and
Puig, S. (2017) On the edge of research and technological application: a critical review of
Electromethanogenesis. *International Journal of Molecular Sciences* **18**.

[8] Mand, T.D., and Metcalf, W.W. (2019) Energy conservation and hydrogenase function
in Methanogenic Archaea, in oarticular the genus *Methanosarcina*. *Microbiol Mol Biol Rev*
83.

[9] Weimer, P.J., and Zeikus, J.G. (1978) One carbon metabolism in Methanogenic bacteria
- cellular characterization and growth of *Methanosarcina barkeri*. *Archives of*
Microbiology **119**: 49-57.

[10] Morii, H., Koga, Y., and Nagai, S. (1987) Energetic analysis of the growth of
Methanobrevibacter Arboriphilus A2 in hydrogen-limited continuous cultures.
Biotechnology and Bioengineering **29**: 310-315.

[11] Karadagli, F., and Rittmann, B.E. (2005) Kinetic characterization of
Methanobacterium bryantii M.o.H. *Environmental Science & Technology* **39**: 4900-4905.

[12] Karadagli, F., Marcus, A., and Rittmann, B.E. (2023) Microbiological hydrogen (H_2)
thresholds in anaerobic continuous-flow systems: Effects of system characteristics.
Biotechnol Bioeng **120**: 1844-1856.

[13] Kral, T.A., Brink, K.M., Miller, S.L., and McKay, C.P. (1998) Hydrogen consumption by
methanogens on the early Earth. *Orig Life Evol Biosph* **28**: 311-319.

[14] Feldewert, C., Lang, K., and Brune, A. (2020) The hydrogen threshold of obligately
methyl-reducing methanogens. *FEMS Microbiol Lett* **367**.

[15] Philips, J. (2019) Extracellular electron uptake by acetogenic bacteria: Does H_2
consumption favor the H_2 evolution reaction on a cathode or metallic iron? *Front*
Microbiol **10**: 2997.

[16] Muñoz, L., and Philips, J. (2023) No acetogen is equal: Strongly different H_2
thresholds reflect diverse bioenergetics in acetogenic bacteria. *Environmental*
Microbiology **25**: 2032-2040.

[17] Pandit, S., Pandit, C., Mathuriya, A.S., Chatterjee, S., Jadhav, D.A., Yadav, K.K., and
Khalid, M. (2024) Wired for energy: Electromethanogenesis redefining anaerobic
digestion. *Process Safety and Environmental Protection* **185**: 588-601.

[18] Mayer, F., Enzmann, F., Lopez, A.M., and Hohmann, D. (2019) Performance of
different methanogenic species for the microbial electrosynthesis of methane from
carbon dioxide. *Bioresource Technology* **289**.

[19] Tsurumaru, H., Ito, N., Mori, K., Wakai, S., Uchiyama, T., Iino, T. et al. (2018) An
extracellular [NiFe] hydrogenase mediating iron corrosion is encoded in a genetically
unstable genomic island in *Methanococcus maripaludis*. *Sci Rep* **8**: 15149.

[20] Holmes, D.E., Woodard, T.L., Smith, J.A., Musat, F., and Lovley, D.R. (2024)
Electrobiocorrosion by microbes without outer-surface cytochromes. *mLife* **3**: 110-118.

[21] Tamisier, M., Schmidt, M., Vogt, C., Kummel, S., Stryhanyuk, H., Musat, N. et al.
(2022) Iron corrosion by methanogenic archaea characterized by stable isotope effects
and crust mineralogy. *Environ Microbiol* **24**: 583-595.

[22] Kawaichi, S., Kotoky, R., Fiutowski, J., and Rotaru, A.E. (2024) Adaptation of a
methanogen to Fe(0) corrosion via direct contact. *NPJ Biofilms Microbiomes* **10**: 100.

[23] Dinh, H.T., Kuever, J., Mussmann, M., Hassel, A.W., Stratmann, M., and Widdel, F.
(2004) Iron corrosion by novel anaerobic microorganisms. *Nature* **427**: 829-832.

[24] Sander, R. (2015) Compilation of Henry's law constants (version 4.0) for water as
solvent *Atmospheric Chemistry and Physics* **Volume 15**.

[25] Callander, G., Deutzmann, J.S., and Spormann, A.M. (2024) Alkaline
hydrogenotrophic methanogenesis in *Methanococcus vannielii* at low carbon dioxide
concentrations. *Journal of Co2 Utilization* **83**.

[26] Schink, B. (1997) Energetics of syntrophic cooperation in methanogenic degradation.
Microbiol Mol Biol Rev **61**: 262-280.

[27] Spormann, A.M., and SpringerLink (2023) nATP and Growth Yields Y_s and Y_{ATP}. In
Principles of Microbial Metabolism and Metabolic Ecology. Cham: Springer International
Publishing : Imprint: Springer, pp. 48-50.

[28] Rittmann, B.E., McCarty, P.L., and AccessEngineering (2020) An empirical formula
for microbial cells. In *Environmental biotechnology : principles and applications*. New
York, N.Y: McGraw-Hill Education, pp. 144-148.

[29] Lopes, J. URL https://errorpropagation.com/

[30] Truong, Q. URL https://uncertaintycalculator.com/

[31] Kaster, A.K., Moll, J., Parey, K., and Thauer, R.K. (2011) Coupling of ferredoxin and
heterodisulfide reduction via electron bifurcation in hydrogenotrophic methanogenic
archaea. *Proc Natl Acad Sci U S A* **108**: 2981-2986.

[32] Costa, K.C., Yoon, S.H., Pan, M., Burn, J.A., Baliga, N.S., and Leigh, J.A. (2013) Effects

of H_2 and formate on growth yield and regulation of methanogenesis in *Methanococcus*

maripaludis. *J Bacteriol* **195**: 1456-1462.

[33] de Poorter, L.M., Geerts, W.J., and Keltjens, J.T. (2007) Coupling of

Methanothermobacter thermautotrophicus methane formation and growth in fed-batch

and continuous cultures under different H_2 gassing regimens. *Appl Environ Microbiol* **73**:

740-749.

[34] Mladenovska, Z., and Ahring, B.K. (2000) Growth kinetics of thermophilic

spp. isolated from full-scale biogas plants treating animal manures. *Fems Microbiology*

Ecology **31**: 225-229.

[35] Richards, M.A., Lie, T.J., Zhang, J., Ragsdale, S.W., Leigh, J.A., and Price, N.D. (2016)

Exploring hydrogenotrophic methanogenesis: a genome scale metabolic reconstruction

of *Methanococcus maripaludis*. *J Bacteriol* **198**: 3379-3390.

[36] Laguillaumie, L., Peyre-Lavigne, M., Grimalt-Alemany, A., Gavala, H.N., Skiadas, I.V.,

Paul, E., and Dumas, C. (2023) Controlling the microbial competition between

hydrogenotrophic methanogens and homoacetogens using mass transfer and

thermodynamic constraints. *Journal of Cleaner Production* **414**.

[37] Leggieri, P.A., Kerdman-Andrade, C., Lankiewicz, T.S., Valentine, M.T., and O'Malley,

M.A. (2021) Non-destructive quantification of anaerobic gut fungi and methanogens in

co-culture reveals increased fungal growth rate and changes in metabolic flux relative to

mono-culture. *Microb Cell Fact* **20**: 199.

[38] Jaramillo, P.A.P. (2019) Microbial induced corrosion by methanogens and other

associated microbial groups. In *Nordcee, department of Biology*: University of Southern

Denmark, pp. 90 - 108.

[39] Mayer, F., Sabel-Becker, B., and Holtmann, D. (2022) Enhanced electron uptake and

methane production by corrosive methanogens during Electromethanogenesis.

Microorganisms **10**.

[40] Mills, S., Dessi, P., Pant, D., Farras, P., Sloan, W.T., Collins, G., and Ijaz, U.Z. (2022) A
meta-analysis of acetogenic and methanogenic microbiomes in microbial
electrosynthesis. *NPJ Biofilms Microbiomes* **8**: 73.

[41] Boone, D.R. Methanobacterium. In *Bergey's Manual of Systematics of Archaea and*
Bacteria, pp. 1-8.

[42] Sowers, K.R., Boone, J.E., and Gunsalus, R.P. (1993) Disaggregation of
Methanosarcina spp. and growth as single cells at elevated osmolarity. *Appl Environ*
Microbiol **59**: 3832-3839.

[43] Angelidaki, I., Karakashev, D., Batstone, D.J., Plugge, C.M., and Stams, A.J. (2011)
Biomethanation and its potential. *Methods Enzymol* **494**: 327-351.

[44] Tsapekos, P., Alvarado-Morales, M., and Angelidaki, I. (2022) H_2 competition between
homoacetogenic bacteria and methanogenic archaea during biomethanation from a
combined experimental-modelling approach. *Journal of Environmental Chemical*
Engineering **10**.

[45] Munoz-Duarte, L., Chakraborty, S., Gron, L.V., Bambace, M.F., Catalano, J., and
Philips, J. (2025) H_2 consumption by various acetogenic bacteria follows First-Order
Kinetics up to H_2 Saturation. *Biotechnol Bioeng* **122**: 804-816.

[46] Kotsyurbenko, O.R., Glagolev, M.V., Nozhevnikova, A.N., and Conrad, R. (2001)
Competition between homoacetogenic bacteria and methanogenic archaea for
hydrogen at low temperature. *FEMS Microbiology Ecology* **38**: 153-159.

[47] Ripoll, E., Lopez, I., and Borzacconi, L. (2020) Hydrogenotrophic activity: A tool to
evaluate the kinetics of methanogens. *J Environ Manage* **270**: 110937.

[48] Conrad, R., and Wetter, B. (1990) Influence of temperature on energetics of hydrogen
metabolism in homoacetogenic, methanogenic, and other anaerobic-bacteria. *Archives*
of Microbiology **155**: 94-98.

[49] Fu, B., Jin, X., Conrad, R., Liu, H., and Liu, H. (2019) Competition between
chemolithotrophic acetogenesis and hydrogenotrophic Methanogenesis for exogenous

H$_2$/CO$_2$ in anaerobically digested sludge: impact of temperature. *Front Microbiol* **10**:
2418.

[50] Molenaar, S.D., Saha, P., Mol, A.R., Sleutels, T.H.J.A., ter Heijne, A., and Buisman,
C.J.N. (2017) Competition between methanogens and acetogens in biocathodes: A
comparison between potentiostatic and galvanostatic control. *International Journal of*
Molecular Sciences **18**.

[51] Le Van, T.D., Robinson, J.A., Ralph, J., Greening, R.C., Smolenski, W.J., Leedle, J.A.,
and Schaefer, D.M. (1998) Assessment of reductive acetogenesis with indigenous
ruminal bacterium populations and *Acetitomaculum ruminis*. *Appl Environ Microbiol* **64**:
3429-3436.

[52] Leclerc, M., Bernalier, A., Donadille, G., and Lelait, M. (1997) H$_2$/CO$_2$ metabolism in
acetogenic bacteria isolated from the human colon. *Anaerobe* **3**: 307-315.

[53] Goyal, N., Zhou, Z., and Karimi, I.A. (2016) Metabolic processes of *Methanococcus*
maripaludis and potential applications. *Microb Cell Fact* **15**: 107.

[54] Spormann, A.M., and SpringerLink (2023) *Principles of Microbial Metabolism and*
Metabolic Ecology. Cham: Springer International Publishing : Imprint: Springer.

[55] Ou, Y.F., Dong, H.P., McIlroy, S.J., Crowe, S.A., Hallam, S.J., Han, P. et al. (2022)
Expanding the phylogenetic distribution of cytochrome b-containing methanogenic
archaea sheds light on the evolution of methanogenesis. *ISME J* **16**: 2373-2387.

[56] Kuhn, W., Fiebig, K., Hippe, H., Mah, R.A., Huser, B.A., and Gottschalk, G. (1983)
Distribution of cytochromes in methanogenic bacteria. *Fems Microbiology Letters* **20**:
407-410.

Methanosarcina mazei

Methanosarcina barkeri

Abiotic controls

Days