# Variations in $H_2$ thresholds and growth yields reveal bioenergetic diversity among hydrogenotrophic methanogens

**Timothé Philippon [1], Jo Philips [1,2]***

[1]  Department of Biological and Chemical Engineering, University of Aarhus,

Gustav Wieds Vej 10, 8000 Aarhus C, Denmark

[2]  Novo Nordisk Foundation $CO_2$ Research Center (CORC), Gustav Wieds Vej 10,

8000 Aarhus C, Denmark

*Corresponding author: jo.philips@bce.au.dk

## Abstract

Hydrogenotrophic methanogens are of high environmental and biotechnological importance, converting $CO_2$ with $H_2$ into $CH_4$. Despite their common metabolism, variations in the energy metabolism among these methanogens exist, likely affecting their $H_2$ thresholds and growth yields. However, a systematic comparison of these traits for a wide range of hydrogenotrophic methanogens has been lacking. Here, we measured the $H_2$ thresholds and growth yields of nine different hydrogenotrophic methanogens. The $H_2$ threshold, i.e. the $H_2$ partial pressure at which $H_2$ consumption halts, ranged over two orders of magnitude from $1.0 \pm 0.5$ Pa for *Methanobrevibacter arboriphilus* to $120 \pm 10$ Pa for *Methanosarcina mazei*. Growth yields in our experimental conditions ranged from $0.51 \pm 0.28$ $g_{DCW} \times (mol\ CH_4)^{-1}$ for *Methanococcus maripaludis* to $5.28 \pm 1.25$ $g_{DCW} \times (mol\ CH_4)^{-1}$ for *Methanosarcina mazei*. The ATP gains, estimated from both $H_2$ thresholds and growth yields, correlated reasonably well, confirming that these variations are due to differences in energy conservation strategies. Our results strongly differentiated the two previously proposed groups of hydrogenotrophic methanogens: methanogens with cytochromes had a high $H_2$ threshold ($\geq 21$ Pa) and high growth yield ($> 4.0$ $g_{DCW} \times (mol\ CH_4)^{-1}$), whereas methanogens without cytochromes had lower $H_2$ threshold ($\leq 7$ Pa) and low growth yield ($< 1.7$ $g_{DCW} \times (mol\ CH_4)^{-1}$). Moreover, our $H_2$ thresholds indicated that additional variations in energy metabolism exist within both groups. Overall, this study found strong variations between hydrogenotrophic methanogens, which are important to understand their environmental prevalence and biotechnological applicability.

## Importance

Hydrogenotrophic methanogens play key roles in natural ecosystems and biotechnological processes. Even though all hydrogenotrophic methanogens convert $CO_2$ with $H_2$ into $CH_4$, they can differ in their $H_2$ threshold and growth yield, due to variations in their energy metabolism. Here, we found that $H_2$ thresholds of hydrogenotrophic methanogens range over two orders of magnitude, while a ten-fold difference was observed in their growth yields. These strong variations in $H_2$ thresholds and growth yields demonstrate that hydrogenotrophic methanogens are confronted with

a trade-off between the capacity to grow at low $H_2$ levels (low $H_2$ threshold) and efficient

growth (high growth yield). Curiously, the observed $H_2$ thresholds also correlate with the

electroactivity of different hydrogenotrophic methanogens. Overall, by reporting $H_2$

thresholds and growth yields for a wide range of hydrogenotrophic methanogens, this

work provides new insights into the bioenergetic diversity of these microbes, as well as

their environmental prevalence and biotechnological applicability.

**Keywords**

Methanogenic archaea, $H_2$ threshold, growth yield, ATP gain, energy conservation,

extracellular electron uptake, ecological trade-off

## Introduction

Hydrogenotrophic methanogenesis is a metabolic pathway used by specific Archaea to produce methane ($CH_4$) from carbon dioxide ($CO_2$) and dihydrogen ($H_2$) for the purpose of energy conservation:

$$CO_2 + 4\,H_2 + n\,ADP + n\,Pi \rightarrow CH_4 + 2\,H_2O + n\,ATP \qquad (1)$$

This metabolism is widely spread in the environment and plays an important role in the global carbon cycle, as methanogens convert yearly about one gigaton of carbon into methane [1]. This production of methane is a major issue in the context of climate change, as methane is a potent greenhouse gas. Methanogenesis can also negatively impact steel infrastructure, as certain hydrogenotrophic methanogens induce strong corrosion [2]. Nevertheless, hydrogenotrophic methanogenesis is of huge interest for the development of energy conversion (e.g. power-to-gas) biotechnologies, as methane is a valuable fuel. Bio-methanation is already an advanced technology, converting $CO_2$-containing gases into methane with the addition of $H_2$ gas (Reaction 1) [3]. Another promising biotechnology is electro-methanogenesis, in which an electrode is used to generate $H_2$ *in situ* of a bioreactor [4, 5]. Because of the technological and environmental importance of hydrogenotrophic methanogens, variations between different methanogens need to be well understood.

All hydrogenotrophic methanogens use a common pathway for their energy metabolism, consisting of a succession of steps reducing $CO_2$ to $CH_4$, using electrons obtained from the oxidation of $H_2$ and generating a sodium ($Na^+$) gradient (Figure 1) [6]. Besides this common pathway, hydrogenotrophic methanogens vary in the involvement of cytochromes [1, 6]. Hydrogenotrophic methanogens without cytochromes (Figure 1A) generate ATP only through the $Na^+$ gradient [7]. In contrast, *M. barkeri* creates an additional $H^+$ gradient through two protein complexes (*Vht* and *hrdDE*), each containing cytochromes (Figure 1B) [6]. Methanogens with cytochromes thus have a higher ATP gain (amount of ATP generated ($n$) in Reaction 1) than methanogens without cytochromes (Table 1, Figure 1). Moreover, further variations in the energy conservation components of both groups of methanogens have been reported [6, 8, 9], suggesting that additional differences in ATP gains may exist.

83    Different ATP gains in turn affect various metabolic characteristics of methanogens, including their $H_2$ threshold and growth yield (Table 1). The $H_2$ threshold quantifies the $H_2$ concentration at which $H_2$ oxidation and therefore methanogenesis halts due to thermodynamic constraints. This $H_2$ threshold is generally lower for methanogens without cytochromes than for methanogens with cytochromes (Table 1 and S1). Accordingly, the growth yield, i.e. the amount of biomass created per amount of $CH_4$ produced, is higher for methanogens with cytochromes than for methanogens without cytochromes (Table 1, Table S2). Additional variations in these metabolic characteristics likely exist, but a coherent comparison of an extensive number of methanogens has not yet been performed.

93    Most studies measuring growth yields for hydrogenotrophic methanogens included just one to two methanogenic species, while each using different media and experimental setups (Table S2). However, growth yields strongly depend on the medium composition [10, 11], the gas supply [12], the dissolved $H_2$ concentration [13] and the growth rate [14] (Table S2). Growth yields determined in different studies are thus not directly comparable. Similarly, most studies measuring $H_2$ thresholds included only few methanogens, each using different conditions (Table S1). $H_2$ thresholds depend on the experimental conditions (e.g. temperature, total pressure, $CO_2$ and $CH_4$ levels and even trace element concentrations [15-17]), which likely explains why reported $H_2$ thresholds for e.g. *Methanosarcina barkeri* differ widely (from 13 to 150 Pa, Table S1). $H_2$ thresholds determined in different studies are thus neither directly comparable. Recently, strongly different $H_2$ thresholds were unexpectedly found for acetogens [18]. It seems thus worthwhile to revisit the $H_2$ thresholds and growth yields of different methanogens.

106   Insights into the $H_2$ thresholds of different methanogens are also crucial to assess their potential for electromethanogenesis and role in microbial-induced corrosion. In both cases, the cathode or steel surface acts as extracellular electron donor, from which methanogens withdraw electrons, using $H_2$ as main intermediate [5]. Not all hydrogenotrophic methanogens are as effective in this extracellular electron uptake [19]. Besides the involvement of extracellular hydrogenases [20], the $H_2$ partial pressure is proposed to play a role in electroactivity [15]. Microbes maintaining low $H_2$ partial pressures likely favor $H_2$ evolution from a cathode or steel surface. As the $H_2$ threshold

reflects the lowest $H_2$ partial pressure a strain can sustain, this parameter is thus likely

crucial to understand the electroactivity of methanogens.

This study aimed to compare $H_2$ thresholds and growth yields of nine strains of

mesophilic hydrogenotrophic methanogens in similar conditions (Table 2). The selected

methanogens include species with and without cytochromes, originating from different

environments and spanning various methanogenic genera. In addition, two

methanogenic strains known for their strong corrosive behavior were included. For all

these strains, the $H_2$ threshold and the growth yield were experimentally measured.

Moreover, ATP gains were estimated from either the $H_2$ threshold or the growth yield. All

these variables are relevant to assess variations among methanogens.

## Material and methods

### Methanogenic strains and culture conditions

This study included nine different methanogens (Table 2). Seven methanogenic strains were ordered from the Leibniz Institute DSMZ, including *Methanococcus maripaludis* (DSMZ 2067 designated as type strain in this study), *Methanolacinia petrolearia* (DSMZ 11571), *Methanobacterium palustre* (DSMZ 3108), *Methanobacterium formicicum* (DSMZ 1535), *Methanosarcina mazei* (DSMZ 2053), *Methanosarcina barkeri* (DSMZ 800) and *Methanobrevibacter arboriphilus* (DSMZ 744). Once received, the strains were transferred to their recommended DSMZ medium (exact composition but without $Na_2S$, sludge fluid or fatty acid mixture) and stored frozen at – 80 °C in a 2 mL tube with 25% glycerol.

In addition, a culture containing *Methanobacterium* strain IM1 [21, 22] was obtained from Florin Musat (Aarhus University). The culture was inoculated in DSMZ medium 141 (without $Na_2S$) and was stored frozen at – 80 °C. Microscope imaging (Leica ICC50 W, Germany) showed that an additional coccus shaped microbe was present in this culture. Nevertheless, when cultivated in a medium with only $H_2$ as electron donor (no acetate), the IM1 strain (bacille shape [22]) became dominant. Acetate was thus further omitted from the medium. Therefore, we assume that the measured $H_2$ threshold was for *Methanobacterium* strain IM1.

*Methanococcus maripaludis* Mic1c10, was provided as living culture by Amelia-Elena Rotaru (Southern University of Denmark). This strain was cultured in DSMZ medium 141 (recommended for *Methanococcus maripaludis*, but without $Na_2S$) and stored at – 80 °C. Both *Methanococcus maripaludis* Mic1c10 and *Methanobacterium* strain IM1 are known corrosive methanogens, as previously documented [20, 21, 23, 24].

To start experiments, frozen stocks were revived in their respective DSMZ medium and transferred to the same medium once. When the late exponential growth phase was reached, 10 mL of culture was transferred to a sterile glass tube with conical bottom, which had first been hermetically sealed using a rubber stopper and flushed with $N_2$. These tubes were then centrifuged at 2500 rpm for 15 minutes. The cell pellet was washed two times using limited medium (described below), before being resuspended in

the same limited medium. For the $H_2$ threshold experiment, this cell suspension was

diluted with limited medium until an optical density (OD) (method described below)

between 0.10 and 0.15 was reached. This procedure was used both for strains that grow

planktonically or as aggregates (notably *M. barkeri* and *M. mazei*), even though similar OD

values in this case do not correspond to comparable biomass densities.

All experiments were performed in limited medium. This limited medium was derived

from the DSMZ medium 141 and included 0.34 g of KCl, 3.45 g of $MgSO_4$ x 7 $H_2O$, 4 g of

$MgCl_2$, 0.25 g of $NH_4Cl$, 0.14 g of $CaCl_2$, 0.14 g of $K_2HPO_4$, 0.2 mg of $Fe(NH_4)_2(SO_4)_2$, 0.2 g

yeast extract powder (Y1625), 10 mL of Wolin mineral solution and 10 mL of Wolin

vitamins solution for a final volume of one liter. In addition, NaCl was added for a final

concentration of 0.4 g per L or 18 g per L, respectively for freshwater methanogens (*M.*

*palustre, M. formicicum, M. barkeri, M. arboriphilus* and *M. mazei*) and marine

methanogens (*strain IM1, M. maripaludis* DSMZ 2067 and Mic1c10 strain and *M.*

*petrolearia*) (Table 2). In contrast to regular DSMZ medium 141, this limited medium did

not contain acetate and had a ten-times lower yeast extract concentration. After

autoclaving this basal medium, filter-sterile stocks solutions were added for a final

concentration of 5 g of $NaHCO_3$ and 0.5 $g.L^{-1}$ of L-cysteine.

## Measurement of the $H_2$ threshold

A volume of 80 mL of the cell suspensions obtained after washing was brought into 250

mL serum bottles. For each strain, four replicates were prepared, while three additional

bottles were filled with 80 mL of limited medium as abiotic controls. The $H_2$ threshold

experiment was initiated by flushing the headspace of the bottles with a gas mix

composed of 2 % $H_2$, 78 % $N_2$ and 20 % $CO_2$. This mix was prepared using gas flow meters

(Alicat®, MCW Series, USA) with two gas tanks, one containing 80% $H_2$ and 20% $CO_2$ and

the other containing 80% $N_2$ and 20 % $CO_2$. Each bottle was pressurized at 2 bars (1 bar

overpressure). The bottles were incubated upright at 30 °C with horizontal orbital shaking

at 100 rpm. The total pressure was measured with a digital manometer (OMEGA DPG108,

UK) before each sampling event. The headspace of each bottle was frequently sampled

with a syringe for $H_2$ analysis (1.5 mL). The $H_2$ threshold was considered reached when

the $H_2$ partial pressure did not significantly decrease for at least four days with

measurement intervals of maximally two days. The value of the $H_2$ threshold was

calculated from the average of two to four measurements performed during those last four days. After the threshold was reached, the headspace was flushed and pressurized again with the gas mix as described above. The headspace was sampled again until the $H_2$ threshold was reached for a second time. The OD and the methane content in the headspace were measured at the end of the experiment. For all methanogens, the amount of $CH_4$ at the end of the $H_2$ threshold experiment was too low to be accurately quantified, even though detectable by GC.

**Growth yield measurements**

For each methanogen, the inoculum was pre-grown and washed as described above. Four replicated bottles (250 mL) with 80 mL of limited medium were inoculated by injecting 0.5 mL of washed culture. These bottles were pressurized at 2 bars (overpressure of 1 bar) using a gas mix with 80% $H_2$ and 20 % $CO_2$. The bottles were incubated upright at 30 °C under horizontal orbital shaking at 100 rpm. The pressure was measured daily until at least one of the four replicates reached a negative pressure, after which the incubation was ended for all replicates. Before biomass was harvested, the headspace composition was analyzed. Next, the bottles were opened and 10 mL of each culture was transferred into a 20 mL glass tube with conical bottom and centrifuged at 3000 rpm for 15 minutes. The clear supernatant was discarded and an additional 10 mL of culture was placed in the same tube, while keeping the cell pellet. This centrifugation procedure was repeated until all 80 mL of each bottle was harvested. Each of the tubes containing the stacked cell pellets were then dried at 100 °C for 48 hours, cooled down for one hour in a desiccator and the dry weight was measured using an analytical balance (± 0.0001 g precision) (VWR, LD17 4XN). The growth yield was calculated by dividing the dried cell weight by the quantity of methane measured in the headspace at the end of the experiment.

The effect of the medium composition on the growth yield was further investigated by measuring the growth yield of one strain (*M. palustre*) in limited medium (described above) and medium with additional yeast extract and sodium acetate (final concentration of 1 g·L$^{-1}$ each).

## Analytical methods

Gaseous samples were analyzed for $H_2$ and $CH_4$ using a Compact GC 4.0 (Interscience, Netherlands) with two different columns: one Molsieve 5A 30 m x 0.32 mm and one Rt-QBond 3 m x 0.32 mm, using helium as gas carrier and a TCD detector [18]. Calibration was performed using calibrating gases mixed with $N_2$. The quantification limit was 1 Pa or 10 ppm for $H_2$ and10 000 ppm for $CH_4$. For $H_2$ partial pressures below 50 ppm, a Peak Performer 1 (PP1) (Peak Laboratories, USA) GC unit based on HgO–Hg conversion with a reducing compound photometer (RCP) was also used. This GC measures trace $H_2$ levels (0.1–50 ppm of $H_2$) and has quantification limit well below that of the CompactGC. Partial pressures were calculated by multiplying the mol fraction, as measured by GC, with the total pressure. Dissolved concentrations were calculated from the partial pressures using the Henry constants for $H_2$ at 30°C (7.8 x 10$^{-4}$ mol.l$^{-1}$.bar$^{-1}$) [25]. Optical densities (OD) were measured at 600 nm using a Denovix DS-11+ spectrophotometer.

## Calculations

### Estimation of the ATP gain from the $H_2$ threshold

As explained previously [18], $H_2$ thresholds can be used to calculate an estimate for the ATP gain through calculating the critical Gibbs free energy, i.e. the Gibbs free energy change when the $H_2$ threshold is reached [1].

Considering hydrogenotrophic methanogenesis:

$$CO_2 + 4\,H_2 \rightarrow CH_4 + 2\,H_2O \tag{2}$$

The critical Gibbs free energy ($\Delta G_c$) becomes:

$$\Delta G_c = \Delta G^{0'}_{\text{methanogenesis}} + R \cdot T \cdot \ln \frac{p_{CH_4}}{p_{CO_2} * \theta_{H_2}^{\,4}} \tag{3}$$

With $\Delta G^{0'}_{\text{methanogenesis}}$ the Gibbs free energy of methanogenesis (Reaction 2) in physiological standard conditions (-131 kJ.mol$^{-1}$ [1]), $\theta_{H_2}$ the $H_2$ threshold (in atm), $p_{CO_2}$ the partial pressure of $CO_2$ (in atm) and $p_{CH_4}$ the partial pressure of methane (in atm) at the $H_2$ threshold. As the methane partial pressures in the $H_2$ threshold experiment were

too low to be quantified, they were calculated from the amount of $H_2$ consumed,
according to the stoichiometric ratios of Reaction 2.

The critical Gibbs free energy at the $H_2$ threshold is not zero, because methanogenesis
should be considered as coupled to ADP to ATP phosphorylation (Reaction 1). At the $H_2$
threshold, it can be assumed that the total Gibbs free energy change of Reaction 1
considering both methanogenesis and ADP phosphorylation, equals zero [1]:

$$\Delta G_c + n \cdot \Delta G_{ATP} = 0 \qquad (4)$$

Where $\Delta G_{ATP}$ is the Gibbs free energy change for the phosphorylation of ADP into ATP.
The value of this $\Delta G_{ATP}$ can vary between microorganisms and different values are used
in literature [1, 18], but here we used the value of -75 kJ.mol$^{-1}$ [26]. The ATP gain *n, i.e.* the
number of ADP molecules phosphorylated for each completed methanogenesis
reaction, can finally be calculated using:

$$n = -\frac{\Delta G_c}{\Delta G_{ATP}} \qquad (5)$$

**Estimation the ATP gain from the growth yield**

The ATP gain *n* can also be estimated from experimental growth yields ($Y^{CH_4}$, g dry cell
weight (DCW) per mol methane), using the biomass yield per mol ATP ($Y^{ATP}$, g dry cell
weight per mol ATP):

$$n = \frac{Y^{CH_4}}{Y^{ATP}} \qquad (6)$$

Different $Y^{ATP}$ values reported for methanogenesis are summarized in Table S3. Most of
these values are maximum values, meaning that the actual $Y_{ATP}$ must be lower. As our
experimental growth yields are observed (net) growth yields and not maximum growth
yields (requiring correction for maintenance energy), we selected a value lower than the
reported maximum and used 5 g dry cell weight per mole of ATP as $Y^{ATP}$ [27].

## Statistical Analysis

### Welch ANOVA test

Welch-Anova tests were performed to differentiate statistically significant groups using both GraphPad Prism® and RStudio (ver. 4.4.3). The Welch test was chosen because the variances were considered too different to perform a regular ANOVA test. All the tests were performed with $\alpha = 0.05$ and the p-value was set at 0.05 for all tests.

### Error propagation

Standard deviations on the experimental results were calculated using Excel and were further propagated through the different calculations using available calculators [28, 29].

## Results and Discussion

### H$_2$ threshold measurements

In this study, the H$_2$ threshold of nine methanogenic strains was measured, as the H$_2$ partial pressure at which their H$_2$ consumption halts. In all abiotic controls, the H$_2$ partial pressure remained almost stable for the entire duration of the experiment (Figure 2, Figure S1), as only a small drop of the total pressure was observed, as the result of headspace sampling (Figure S1).

For all inoculated bottles, a clear drop in the H$_2$ partial pressure was observed (Figure 2), confirming H$_2$ consumption by the different methanogens. Moreover, H$_2$ consumption halted at a certain H$_2$ partial pressure for each strain. To confirm these H$_2$ thresholds, H$_2$ was reinjected, once stable H$_2$ partial pressures were reached. For most methanogenic strains, the H$_2$ thresholds reached after the re-addition of H$_2$ were comparable to those before the re-addition, except for *Methanosarcina mazei* and *Methanosarcina barkeri* (Figure S2). *M. mazei* consumed H$_2$ after the H$_2$ re-addition, but its H$_2$ consumption halted at a higher H$_2$ partial pressure (260 ± 49 Pa versus 120 ± 10 Pa before the re-addition). In contrast, *M. barkeri* no longer consumed H$_2$ after the re-addition. Possibly, this is due to the loss of activity of these strains over the long incubation times (*M. mazei* reached its H$_2$ threshold after 12 days, while *M. barkeri* required 28 days). All other methanogens required only five days or less to reach their H$_2$ threshold.

The three *Methanobacterium* species displayed similar low H$_2$ thresholds around 2 Pa (Figure 2, Figure 3, Table 2). Also, *Methanosarcina petrolearia* and *Methanobrevibacter arboriphilus* reached H$_2$ thresholds in this range. *M. arboriphilus* was the only species that reached a H$_2$ partial pressure of 1 Pa (Figure 2, Figure 3, Table 2). The Welch-ANOVA test gathered these five strains with the lowest H$_2$ thresholds in one statistical group (Figure 3). In contrast, all the other strains had higher H$_2$ thresholds and were classified each as a significantly different group (Figure 3). The two strains of *Methanococcus maripaludis* displayed H$_2$ thresholds slightly higher than the five strains with the lowest H$_2$ thresholds (Figure 2, Figure 3, Table 2). Interestingly, the corrosive strain *Mic1c10* had a significantly lower H$_2$ threshold (around 4 ± 1 Pa) than the type strain (7 ± 1 Pa).

The *Methanosarcina* species differed strongly in their $H_2$ thresholds, as *M. barkeri* had a

$H_2$ threshold of 21 Pa ± 2 Pa, while the highest $H_2$ threshold was recorded for *M. mazei*

(120 ± 10 Pa). Previously, widely varying $H_2$ thresholds were reported for *M. barkeri* (Table

S1). These variations are possibly due to different experimental conditions (Table S1) or

the slow $H_2$ consumption by this strain. Our $H_2$ threshold for *M. barkeri* could be

somewhat overestimated, as this strain could not be reactivated after the halt of $H_2$

consumption. In contrast, *M. mazei* was still active after the $H_2$ re-addition, so its six-

times higher $H_2$ threshold in comparison to *M. barkeri* (Table 2) cannot be due to an

experimental artefact.

Overall, we obtained experimental $H_2$ thresholds ranging over two orders of magnitude

(Figure 3, Table 2), which is a similar range as previously recorded (Table S1). The $H_2$

thresholds also agree with the classification based on the presence of cytochromes

(Table 1). We found $H_2$ thresholds below 7 Pa for all methanogens without cytochromes,

while methanogens with cytochromes had $H_2$ thresholds above 20 Pa (Table 2). Our

results thus differentiate both groups more strongly than previous recordings (Table 1).

Moreover, in both groups, we found methanogens with statistically different $H_2$

thresholds (Figure 3), showing that additional variations in $H_2$ thresholds exist.

## Growth yield measurements

In parallel of the $H_2$ threshold study, all methanogenic strains were grown in bottles on 80% $H_2$ and 20% $CO_2$ to determine their growth yields. In all these bottles methane levels significantly increased over the incubation period, while $H_2$ was consumed and cell growth occurred (Table S4). These results were used to calculate the growth yield, i.e. the increase of biomass expressed as gram dry cell weight ($g_{DCW}$) versus the amount of methane produced (Figure 4, Table 2). For one strain, *M. palustre*, the growth yield was also determined with additional acetate and yeast extract in the medium (1 g·L$^{-1}$ each). A significantly higher growth yield was found with the addition of extra organic carbon sources (Figure S3), demonstrating the importance of using the same medium conditions when comparing growth yields.

Relatively high variabilities between the replicates were obtained for the growth yields (large error bars in Figure 4). This is likely due to the low total weight of dried biomass (in mg) obtained during the growth yield experiment (Table S4), limiting the accuracy of this measurement. In addition, the centrifugation steps were possibly not always as effective in removing cells from the culture suspension, which could also explain the high variability between the replicates and could have underestimated some of the growth yields. A high variability is fairly common for growth yield measurements, as this was also observed in other studies determining methanogenic growth yields [11, 30, 31].

Despite the high variability in the dry cell weight, Welch-ANOVA tests identified two significantly different groups in the growth yields (Figure 4, right). The two *Methanosarcina* strains form one group with high growth yields of 4.09 ± 1.16 and 5.28 ± 1.25 $g_{DCW}$/mole of $CH_4$ for *M. barkeri* and *M. mazei* respectively. All other methanogens had significantly lower growth yields, ranging from 0.51 ± 0.28 $g_{DCW}$/mole of $CH_4$ (for *M. maripaludis* type strain) to 1.69 ± 0.58 $g_{DCW}$/mole of $CH_4$ (for *M. formicicum*). Growth yields were also expressed as $g_{DCW}$ per mole $H_2$ consumed (Figure S4), leading to the same statistical groups.

Overall, our growth yields are consistent, but in the lower range, of previously reported growth yields (Table 2 and S2), likely because our medium did not contain organic carbon sources (Figure S3). Moreover, the two significantly different groups identified in the

growth yields agree with the classification based on the presence of cytochromes, while

our results differentiate both groups more strongly than previous findings (Table 1). In

both groups, no further significant differences in growth yields were identified. Even for

*M. mazei* and *M. barkeri,* which differ six times in their $H_2$ threshold, no significant

different growth yields were found (Figure 4). It is likely that this incoherence is due to the

high variability of the growth yield measurement (high error bars in Figure 4).

## Comparison of ATP gain estimates

Both the $H_2$ threshold and the growth yield allow to estimate the ATP gain. Using the $H_2$

threshold, the ATP gain estimates ranged from 0.32 ± 0.07 for *M. arboriphilus* to 0.97 ±

0.01 for *M. mazei* (Table 2). Estimating ATP gains from the growth yields gave a slightly

wider range, starting at 0.10 ± 0.03 for *M. maripaludis* up to 1.06 ± 0.12 for *M. mazei*.

Despite the totally different experimental procedures and the various assumptions

underlying these calculations (Equations 5 and 6), both estimates led to a largely

overlapping range, validating our approach. These ATP gain estimates were further

correlated to each other, leading to a clear but not perfect correlation (Figure 5, $R^2$ of

0.84). Only the two *Methanosarcina* species, *M. mazei* and *M. barkeri*, fall on the 1:1

correlation line. For all other strains, higher ATP gains were calculated from the $H_2$

thresholds than from the growth yield, demonstrating that there were discrepancies

between our methods, possibly because the growth yields were somewhat

underestimated. This illustrates that our ATP gains should be considered as estimates,

rather than absolute values. Nevertheless, the clear correlation (Figure 5) does suggest

that the ATP gains differed among the tested strains.

These ATP gain estimates also agree with the cytochrome-dependent classification of

methanogens (Table 1 and 2). Kaster et al. [7] estimated an ATP gain higher than 1 for *M.*

*barkeri*, while we calculated this to be 0.82 ± 0.11 ATP per mol $CH_4$, when derived from

the growth yield (Table 2). For methanogens without cytochromes, Kaster et al. [7]

proposed an ATP gain lower than 1, while Goyal et al. [32] estimated an ATP gain of 0.35

for *M. maripaludis*. Both correspond well with the range of ATP gain estimates (between

0.10 and 0.56 mol ATP per mol $CH_4$) determined here for methanogens without

cytochromes (Table 1 and 2).

In addition, our ATP gain estimates calculated from the $H_2$ thresholds suggest that

additional variations in energy conservation exist, as additional significant differences in

$H_2$ thresholds were recorded in both groups of methanogens (Figure 3). Unfortunately,

this was not confirmed by the ATP gains derived from the growth yields, likely again due

to the high variability on the growth yields. Besides the two different mechanisms for

energy conservation in methanogens (Figure 1), various additional variations in energy

conservation components are currently being unraveled [6, 8, 9, 33]. Moreover,

*Methanothermobacter marburgensis* was recently shown to have a flexible energy

conservation system, adapting to the nickel concentration in the medium [34]. We

therefore conclude that the measurement of $H_2$ thresholds could offer a valuable tool to

assess how variations in energy conservation affect the ATP gain.

## Link between the $H_2$ threshold and electroactivity of methanogens

The experimental $H_2$ thresholds obtained in this study allow to reevaluate previously reported differences in electroactivity among methanogens. Mayer et al [19] studied the electromethanogenesis performance of several methanogens. *M. maripaludis* and *M. petrolearia*, for which a low $H_2$ threshold was found here (Figure 3), increased cathodic currents (i.e. key signature of electroactivity), just as several other methanogens without cytochromes. In contrast, *M. mazei*, for which the highest $H_2$ threshold was recorded in this study (Figure 3), was incapable of electromethanogenesis [19]. Similarly, Jaramillo *et al*. investigate the corrosion potential of different methanogens [35]. Non-hydrogenotrophic methanogens were reported as non-corrosive, confirming the important role of $H_2$ as intermediate in microbial corrosion. *M. mazei* and *M. barkeri* (high $H_2$ threshold) were mild corrosive, whereas *Methanobacterium* spp. (low $H_2$ threshold) were highly corrosive. Overall, these correlations thus suggest that methanogens with a low $H_2$ threshold are more electroactive than methanogens with a high $H_2$ threshold.

Nevertheless, several other variables likely affect the electroactivity of methanogens, in addition to the $H_2$ threshold. The $H_2$ pressure maintained at the cathode or steel surface also depends on the rate of $H_2$ consumption. Moreover, the $H_2$ evolution reaction on a cathode or steel surface is stimulated by extracellular enzymes, such as hydrogenases. *M. maripaludis* Mic1c10 has a unique extracellular hydrogenase anchored to its cell surface [23], causing stronger corrosion than other *M. maripaludis* strains [23]. Moreover, its cathodic current increase was double as high than for a non-corrosive *M. maripaludis* strain [19, 36]. Here, we found that strain Mic1c10 had a significantly lower $H_2$ threshold than the *M. maripaludis* wild type (Figure 3, Table 2). Kawaichi et al [23] also recorded that strain Mic1c10 maintained slightly lower $H_2$ levels during steel corrosion than by the *M. maripaludis* type strain. These differences, however, are possibly too small to explain the strong corrosion rates caused by Mic1c10, which is thus likely mostly due to its extracellular hydrogenases. Similarly, *Methanobacterium* -IM1 was more corrosive than other tested methanogens [24], but here we did not find that strain IM1 had a lower $H_2$ threshold than other tested *Methanobacterium* strains (Figure 2, Table 3). *Methanobacterium* IM1 thus likely has an additional mechanism to enhance extracellular

429    electron uptake rate, which is possibly similar to the one described for *M. maripaludis*

430    Mic1c10.

431    Overall, we conclude that the electroactivity of methanogens is inversely proportional to

432    their $H_2$ threshold, whereas few methanogens (e.g. *M. maripaludis* Mic1c10,

433    *Methanobacterium* IM1) have additional mechanisms to increase their electroactivity.

## Ecological and biotechnological implications of variations among methanogens

We systematically found that methanogens with a low $H_2$ threshold have a low growth yield and a low ATP gain (Table 2). In contrast, methanogens with a high $H_2$ threshold have a high growth yield and have a higher ATP gain (Table 2). Hydrogenotrophic methanogens are thus confronted with a trade-off between the capacity to consume $H_2$ at trace levels or the ability to grow strongly (high growth yield) with a high energetic efficiency (high ATP gain). This trade-off allows to understand the ecological niche and biotechnological relevance of the different types of methanogens. For instance, Mills *et al.* [37] found that the presence of the *Methanosarcina* genus in electro-methanogenesis systems was correlated to high $H_2$ levels (reactors with low coulombic efficiency), while the *Methanobacterium* genus was rather correlated to low $H_2$ levels (high coulombic efficiency).

In addition, a correlation between the growth yield (high ATP gain) and the growth rate of methanogens would be expected [18]. However, Thauer [1] described that methanogens with cytochromes (high ATP gain) have a lower growth rate (high doubling time) than methanogens without cytochromes (low ATP gain, but low doubling times). Comparing doubling times is though not adequate, since methanogens strongly vary in cell size (large clustered cells for *Methanosarcina* species versus small different shaped cells for other methanogens) [38, 39]. Nevertheless, also in the $H_2$ threshold experiment of this study, $H_2$ was faster consumed by methanogens without cytochromes, than by methanogens with cytochromes (Figure 2). This observation possibly reflects that *Methanosarcina* species are not well adapted to consume $H_2$ at lower levels. These methanogens are also capable of acetoclastic and methylotrophic methanogenesis, in contrast to the other tested methanogens. Methanogens with cytochromes are thus possibly mainly adapted to use other substrates than $H_2$ and only shift to $H_2$ consumption at high $H_2$ partial pressures.

In this study, we found that $H_2$ thresholds of methanogens range over two orders of magnitude (Figure 3). In comparison, we previously found three orders of magnitude difference in the $H_2$ thresholds of acetogenic bacteria (6 Pa to 1990 Pa) [18]. It is often

assumed that acetogens cannot compete with methanogens for low $H_2$ levels, as the

latter have lower $H_2$ thresholds [40]. Our results, however, demonstrate that such a

competition rather should be evaluated on species level. For instance,

*Methanobacterium* strains indeed have lower $H_2$ thresholds than *Acetobacterium* spp,

but acetogenic *Sporomusa* spp. have $H_2$ thresholds in the range of those of methanogens

without cytochromes and well below those with cytochromes. The outcome of

competition for trace $H_2$ levels between methanogens and acetogens likely also depends

on the $H_2$ consumption rates. Recently, $H_2$ consumption by acetogens was described by

first order kinetics (increasing rates at increasing $H_2$ levels) [41], while the kinetics of $H_2$

consumption by methanogens rather follows Monod kinetics with low half-saturation

constants [42] or even zero-order kinetics [43]. Based on these kinetic differences,

methanogens (without cytochromes) consume $H_2$ likely faster at low $H_2$ levels, while

acetogens have a faster $H_2$ consumption at higher $H_2$ levels, even though $H_2$ consumption

rates can still widely vary within these groups [41].

Overall, we conclude that the trade-off between a low $H_2$ threshold and high growth yield

of hydrogenotrophic methanogens explains their ecological prevalence and affects their

biotechnological applicability.

## Acknowledgement

This study was financed by a Villum Young Investigator grant (VIL53083). We would like to thank Satoshi Kawaichi and Amelia-Elena Rotaru from the University of Southern Denmark for providing the living culture of *Methanococcus maripaludis* Mic1cc10 and Florin Musat from Aarhus University for providing the living culture of *Methanobacterium* IM1. We also acknowledge the Section for Microbiology at Aarhus University for lending their Peak Performer. Heidi Skov Johansen and Laura Mũnoz-Duarte are thanked for their practical support during the experiments and calculations. We also appreciate the advice of Sanne Sandberg Overby on the statistical tests.

**Data availability statement**

The data that supports the findings of this study are available in the supporting information file.

**Conflict of interest statement**

The authors declare no conflict of interest.

**Figures**

Figure 1: Energy metabolism for hydrogenotrophic methanogenesis without cytochromes (left) and with

cytochromes (for *M. barkeri*) (right). Black colored components and reactions are common for all

hydrogenotrophic methanogens, including the generation of a sodium ($Na^+$) gradient through the transfer

of the methyl group from tetrahydrometanopterin ($H_4PMT$) to Coenzyme M (CoM). Hydrogenotrophic

methanogens without cytochromes generate ATP only through this $Na^+$ gradient. They use flavin-based

electron bifurcation to couple the endergonic reduction of Ferredoxin (Fd) with $H_2$ as electron donor to the

exergonic reduction of CoM-CoB, through the action of the *Mvh-Hdr* ([NiFe]-hydrogenase - heterodisulfide

reductase) complex. In contrast, *M. barkeri* (with cytochromes) generates ATP through both a $Na^+$ and $H^+$

gradient. In addition to the $Na^+$ gradient, *M. barkeri* creates a $H^+$ gradient through two protein complexes

*Vht* (methanophenazine-reducing hydrogenase) and *hrdDE* (methanophazine-dependent heterodisulfide

reductase), each containing cytochromes and together forming an electron transport chain transporting

electrons from $H_2$ to CoM-CoB and pumping $H^+$ out of the cell. In addition, *M. barkeri* reduces Fd with the

electron donor $H_2$ through consumption of the $H^+$ gradient using an *Ech* (energy converting hydrogenase)

complex. Other abbreviations: methanofuran (MF), Coenzyme B (CoB)[6, 8].

Figure 2:  Change of the $H_2$ partial pressure (Pa) over time in the $H_2$ threshold experiment for the nine

methanogenic species and compared to the abiotic controls. The data points represent the average of four

replicates; the error bars show the standard deviations. Only datapoints before the re-addition of $H_2$ are

shown, later datapoints are shown in Figures S1 and S2.

Figure 3: Box plots summarizing the experimental $H_2$ threshold values for the nine methanogenic strains.

The letters indicate statistically different groups, as determined using a Welch-ANOVA test.

528  Figure 4: Comparison of the growth yields expressed as the increase of gram Dry Cell Weight ($g_{DCW}$) per

529  mole of methane produced for the different methanogenic strains. The letters indicate statistically different

530  groups, as determined using a Welch-ANOVA test.

Figure 5: Correlation of the ATP gains estimations from the experimental $H_2$ thresholds versus the ATP gains
estimated from the experimental growth yields. The vertical and horizontal error bars reflect the standard
deviation on these estimates and were propagated from the experimental errors. The red line indicates a
1:1 correlation. The correlation coefficient (R2) was 0.84.

537 Table 1: Comparison of metabolic characteristics of hydrogenotrophic methanogens depending on the

538 presence or absence of cytochromes according to previous literature, as well as this study.

| | Methanogens without cytochromes | Methanogens with cytochromes |
| --- | --- | --- |
| | Including *Methanococcus* *Methanobrevibacter* *Methanobacterium* *Methanolacinia* | Including *Methanosarcina* |
| $H_2$ threshold (Pa) | | |
| Previous literature (Table S1) | $\leq$20 | $\geq$13 |
| This study (Table 2) | $\leq$7.0 | $\geq$21 |
| Growth yield ($g_{DCW} \cdot mol^{-1} CH_4$) | | |
| Previous literature (Table S2) | 0.8 - 3.5 | 5.4 - 8.4 |
| This study (Table 2) | 0.5 - 1.7 | 4.1 - 5.3 |
| ATP gain (mol ATP$\cdot$mol$^{-1}$ CH$_4$) | | |
| [7] | <1 | >1 |
| This study, based on $H_2$ threshold (Table 2) | 0.32-0.56 | 0.75-0.95 |
| This study, based on growth yield (Table 2) | 0.10-0.34 | 0.82-1.06 |

547 Table 2: Overview of the methanogenic strains examined in this study, as well as their experimentally determined $H_2$ thresholds and growth yields. The table also
548 includes the ATP gains estimated either from the experimental $H_2$ thresholds or the experimental growth yields. The $H_2$ thresholds measured are expressed as partial
549 pressures in the headspace (both in ppm and Pa) and as concentration dissolved in the liquid phase (nM).

| Methanogenic strain | Cytochromes | Salinity level | $H_2$ Threshold | | | Growth yield | ATP gain estimated from $H_2$ threshold | ATP gain estimated from growth yield |
|---|---|---|---|---|---|---|---|---|
| | | | ppm | Pa | nM | $g_{DCw}\cdot(mol\ CH_4)^{-1}$ | mol ATP $(mol\ CH_4)^{-1}$ | mol ATP $(mol\ CH_4)^{-1}$ |
| *Methanobrevibacter arboriphilus* | No [33, 44] | Fresh water | 7 ± 1 | 1.0 ± 0.5 | 8 ± 3 | 0.71 ± 0.47 | 0.32 ± 0.07 | 0.14 ± 0.05 |
| *Methanolacinia petrolearia* | No [33] | Marine water | 10 ± 3 | 2.0 ± 0.5 | 16 ± 4 | 0.63 ± 0.31 | 0.41 ± 0.03 | 0.13 ± 0.03 |
| *Methanobacterium palustre* | No [33] | Fresh water | 11 ± 2 | 2.0 ± 0.5 | 16 ± 4 | 1.57 ± 0.41 | 0.41 ± 0.03 | 0.31 ± 0.04 |
| *Methanobacterium* IM1 | No [33]* | Marine water | 9 ± 1 | 2.0 ± 0.5 | 16 ± 4 | 1.39 ± 0.36 | 0.41 ± 0.03 | 0.28 ± 0.04 |
| *Methanobacterium formicicum* | No [33] | Fresh water | 10 ± 1 | 2.0 ± 0.5 | 16 ± 4 | 1.69 ± 0.58 | 0.41 ± 0.03 | 0.34 ± 0.06 |
| *Methanococcus maripaludis* Mutant strain Mic1c10 | No [32, 33] | Marine water | 17 ± 3 | 4.0 ± 1.0 | 31 ± 8 | 1.02 ± 0.55 | 0.51 ± 0.03 | 0.20 ± 0.05 |
| *Methanococcus maripaludis* Type strain | No [32, 33] | Marine water | 33 ± 4 | 7.0 ± 1.0 | 47 ± 12 | 0.51 ± 0.28 | 0.56 ± 0.01 | 0.10 ± 0.03 |
| *Methanosarcina barkeri* | Yes [6, 33] | Fresh water | 158 ± 18 | 21 ± 2 | 195 ± 31 | 4.09 ± 1.16 | 0.75 ± 0.01 | 0.82 ± 0.11 |
| *Methanosarcina mazei* | Yes [6, 33, 44] | Fresh water | 720 ± 20 | 120 ± 10 | 936 ± 78 | 5.28 ± 1.25 | 0.97 ± 0.01 | 1.06 ± 0.12 |

550 * The assertion that *Methanobacterium* IM1 does not possess cytochrome is based on the fact that no other *Methanobacterium* was found to have cytochromes.

# References

1. Thauer RK, Kaster AK, Seedorf H *et al*. Methanogenic archaea: Ecologically relevant differences in energy conservation. *Nat Rev Microbiol*. 2008;**6**:579-91 10.1038/nrmicro1931

2. An BA, Deland E, Sobol O *et al*. The differences in the corrosion product compositions of methanogen-induced microbiologically influenced corrosion (mi-mic) between static and dynamic growth conditions. *Corros Sci*. 2021;**180** 10.1016/j.corsci.2020.109179

3. Jensen MB, Ottosen LDM, Kofoed MVW. $H_2$ gas-liquid mass transfer: A key element in biological power-to-gas methanation. *Renew Sust Energ Rev*. 2021;**147** 10.1016/j.rser.2021.111209

4. Cheng SA, Xing DF, Call DF *et al*. Direct biological conversion of electrical current into methane by electromethanogenesis. *Environ Sci Technol*. 2009;**43**:3953-58 10.1021/es803531g

5. Palacios PA, Philips J, Bentien A *et al*. Relevance of extracellular electron uptake mechanisms for electromethanogenesis applications. *Biotechnol Adv*. 2024;**73** 10.1016/j.biotechadv.2024.108369

6. Mand TD, Metcalf WW. Energy conservation and hydrogenase function in methanogenic archaea, in particular the genus *methanosarcina*. *Microbiol Mol Biol Rev*. 2019;**83** 10.1128/MMBR.00020-19

7. Kaster AK, Moll J, Parey K *et al*. Coupling of ferredoxin and heterodisulfide reduction via electron bifurcation in hydrogenotrophic methanogenic archaea. *Proc Natl Acad Sci U S A*. 2011;**108**:2981-6 10.1073/pnas.1016761108

8. Downing BE, Nayak DD. Innovations in the electron transport chain fuel archaeal methane metabolism. *Trends in Biochemical Sciences*. 2025;**50**:425-37 10.1016/j.tibs.2025.02.004

9. Gilmore SP, Henske JK, Sexton JA *et al*. Genomic analysis of methanogenic archaea reveals a shift towards energy conservation. *BMC Genomics*. 2017;**18**:639 10.1186/s12864-017-4036-4

10. Costa KC, Yoon SH, Pan M *et al*. Effects of $h_2$ and formate on growth yield and regulation of methanogenesis in *methanococcus maripaludis*. *J Bacteriol*. 2013;**195**:1456-62 10.1128/JB.02141-12

11. Ferguson TJ, Mah RA. Effect of $h_2$-$co_2$ on methanogenesis from acetate or methanol in *methanosarcina* spp. *Applied and Environmental Microbiology*. 1983;**46**:348-55 10.1128/aem.46.2.348-355.1983

12. Schönheit P, Moll J, Thauer RK. Growth parameters ($k_s$, $\mu_{max}$, $y_s$) of *methanobacterium thermoautotrophicum*. *Arch Microbiol*. 1980;**127**:59-65 10.1007/BF00414356

13. de Poorter LM, Geerts WJ, Keltjens JT. Coupling of *methanothermobacter thermautotrophicus* methane formation and growth in fed-batch and continuous cultures under different $h_2$ gassing regimens. *Appl Environ Microbiol*. 2007;**73**:740-9 10.1128/AEM.01885-06

14. Morii H, Koga Y, Nagai S. Energetic analysis of the growth of *methanobrevibacter arboriphilus* a2 in hydrogen-limited continuous cultures. *Biotechnol Bioeng*. 1987;**29**:310-15 DOI 10.1002/bit.260290304

15.  Philips J. Extracellular electron uptake by acetogenic bacteria: Does h₂
consumption favor the h₂ evolution reaction on a cathode or metallic iron? *Front*
*Microbiol*. 2020;**10**:2997 10.3389/fmicb.2019.02997

16.  Conrad R, Wetter B. Influence of temperature on energetics of hydrogen
metabolism in homoacetogenic, methanogenic, and other anaerobic-bacteria.
*Arch Microbiol*. 1990;**155**:94-98 Doi 10.1007/Bf00291281

17.  Neubeck A, Sjöberg S, Price A *et al*. Effect of nickel levels on hydrogen partial
pressure and pethane production in methanogens. *Plos One*. 2016;**11**
10.1371/journal.pone.0168357

18.  Muñoz L, Philips J. No acetogen is equal: Strongly different h₂ thresholds reflect
diverse bioenergetics in acetogenic bacteria. *Environ Microbiol*. 2023;**25**:2032-40
10.1111/1462-2920.16429

19.  Mayer F, Enzmann F, Lopez AM *et al*. Performance of different methanogenic
species for the microbial electrosynthesis of methane from carbon dioxide.
*Bioresource Technol*. 2019;**289** 10.1016/j.biortech.2019.121706

20.  Tsurumaru H, Ito N, Mori K *et al*. An extracellular [nife] hydrogenase mediating iron
corrosion is encoded in a genetically unstable genomic island in *methanococcus*
*maripaludis*. *Sci Rep*. 2018;**8**:15149 10.1038/s41598-018-33541-5

21.  Holmes DE, Woodard TL, Smith JA *et al*. Electrobiocorrosion by microbes without
outer-surface cytochromes. *mLife*. 2024;**3**:110-18 10.1002/mlf2.12111

22.  Tamisier M, Schmidt M, Vogt C *et al*. Iron corrosion by methanogenic archaea
characterized by stable isotope effects and crust mineralogy. *Environ Microbiol*.
2022;**24**:583-95 10.1111/1462-2920.15658

23.  Kawaichi S, Kotoky R, Fiutowski J *et al*. Adaptation of a methanogen to fe(0)
corrosion via direct contact. *NPJ Biofilms Microbiomes*. 2024;**10**:100
10.1038/s41522-024-00574-w

24.  Dinh HT, Kuever J, Mussmann M *et al*. Iron corrosion by novel anaerobic
microorganisms. *Nature*. 2004;**427**:829-32 10.1038/nature02321

25.  Sander R. Compilation of henry's law constants (version 4.0) for water as solvent
*Atmospheric Chemistry and Physics* 2015;**Volume 15** /10.5194/acp-23-10901-
2023.

26.  Schink B. Energetics of syntrophic cooperation in methanogenic degradation.
*Microbiol Mol Biol Rev*. 1997;**61**:262-80 10.1128/mmbr.61.2.262-280.1997

27.  Spormann AM. *Principles of microbial metabolism and metabolic ecology*, 1st
2023. edn.: Springer International Publishing : Imprint: Springer, 2023.

28.  Lopes J. https://errorpropagation.com/

29.  Truong Q. https://uncertaintycalculator.com/

30.  Richards MA, Lie TJ, Zhang J *et al*. Exploring hydrogenotrophic methanogenesis: A
genome scale metabolic reconstruction of *methanococcus maripaludis*. *J*
*Bacteriol*. 2016;**198**:3379-90 10.1128/JB.00571-16

31.  Mladenovska Z, Ahring BK. Growth kinetics of thermophilic spp. Isolated from full-
scale biogas plants treating animal manures. *Fems Microbiology Ecology*.
2000;**31**:225-29 DOI 10.1111/j.1574-6941.2000.tb00687.x

32.  Goyal N, Zhou Z, Karimi IA. Metabolic processes of *methanococcus maripaludis*
and potential applications. *Microb Cell Fact*. 2016;**15**:107 10.1186/s12934-016-
0500-0

642 33. Ou YF, Dong HP, McIlroy SJ *et al*. Expanding the phylogenetic distribution of
643   cytochrome b-containing methanogenic archaea sheds light on the evolution of
644   methanogenesis. *ISME J*. 2022;**16**:2373-87 10.1038/s41396-022-01281-0

645 34. Nomura S, San Segundo-Acosta P, Protasov E *et al*. Electron flow in
646   hydrogenotrophic methanogens under nickel limitation. *Nature*. 2025;**644**:490-96
647   10.1038/s41586-025-09229-y

648 35. Palacios PA. Microbial induced corrosion by methanogens and other associated
649   microbial groups. PhD thesis, PhD thesis, University of Southern Denmark, 2019

650 36. Mayer F, Sabel-Becker B, Holtmann D. Enhanced electron uptake and methane
651   production by corrosive methanogens during electromethanogenesis.
652   *Microorganisms*. 2022;**10** 10.3390/microorganisms10112237

653 37. Mills S, Dessi P, Pant D *et al*. A meta-analysis of acetogenic and methanogenic
654   microbiomes in microbial electrosynthesis. *NPJ Biofilms Microbiomes*. 2022;**8**:73
655   10.1038/s41522-022-00337-5

656 38. Boone DR *Methanobacterium*. *Bergey's manual of systematics of archaea and
657   bacteria*, 1-8. Retreived from
658   https://onlinelibrary.wiley.com/doi/abs/10.1002/9781118960608.gbm00495

659 39. Sowers KR, Boone JE, Gunsalus RP. Disaggregation of *methanosarcina spp*. And
660   growth as single cells at elevated osmolarity. *Appl Environ Microbiol*.
661   1993;**59**:3832-9 10.1128/aem.59.11.3832-3839.1993

662 40. Tsapekos P, Alvarado-Morales M, Angelidaki I. $H_2$ competition between
663   homoacetogenic bacteria and methanogenic archaea during biomethanation
664   from a combined experimental-modelling approach. *J Environ Chem Eng*. 2022;**10**
665   10.1016/j.jece.2022.107281

666 41. Munoz-Duarte L, Chakraborty S, Gron LV *et al*. $H_2$ consumption by various
667   acetogenic bacteria follows first-order kinetics up to $h_2$ saturation. *Biotechnol
668   Bioeng*. 2025;**122**:804-16 10.1002/bit.28904

669 42. Kotsyurbenko OR, Glagolev MV, Nozhevnikova AN *et al*. Competition between
670   homoacetogenic bacteria and methanogenic archaea for hydrogen at low
671   temperature. *FEMS Microbiology Ecology*. 2001;**38**:153-59 10.1111/j.1574-
672   6941.2001.tb00893.x

673 43. Ripoll E, Lopez I, Borzacconi L. Hydrogenotrophic activity: A tool to evaluate the
674   kinetics of methanogens. *J Environ Manage*. 2020;**270**:110937
675   10.1016/j.jenvman.2020.110937

676 44. Kuhn W, Fiebig K, Hippe H *et al*. Distribution of cytochromes in methanogenic
677   bacteria. *Fems Microbiology Letters*. 1983;**20**:407-10