Bidirectional translocation of actomyosin drives epithelial invagination in ascidian siphon tube morphogenesis

Jinghan Qiao[1,#], Pengyu Yu[2,#], Hongzhe Peng[1], Wenjie Shi[1], Bo Li[2,*], and Bo Dong[1,3,*]

1 Fang Zongxi Center, MoE Key Laboratory of Marine Genetics and Breeding, College of Marine Life Sciences, Ocean University of China, Qingdao 266003, China.

2 Institute of Biomechanics and Medical Engineering, Applied Mechanics Laboratory, Department of Engineering Mechanics, Tsinghua University, Beijing 100084, China.

3 Institute of Evolution & Marine Biodiversity, Ocean University of China, Qingdao 266003, China.

# These authors contribute equally to this work

* Correspondence: bodong@ouc.edu.cn (B.D.); libome@tsinghua.edu.cn (B.L.)

Summary

How epithelia preform a spatiotemporal heterogeneous force generating program to drive a sequential tissue morphogenesis remains unclear, particularly the underlying precise mechanical mechanisms. This study investigated dynamic actomyosin reorganization between apical and lateral membrane cortex regions during two sequentially invaginated stages during ascidian atrial siphon tube morphogenesis. At the initial invagination stage, the originally lateral-located actomyosin translocated to the apical domains, while those actomyosin re-translocated back to lateral domains at the accelerated invagination stage. Using genetic mutants to modulate myosin activities, the initial invagination was strengthened or abolished, indicating invagination are apical constriction dependent. Optogenetic inhibition of myosin activities in lateral domains after initial invagination stage blocked the further processes, suggesting lateral constriction of actomyosin is required for the accelerated invagination. Vertex model simulations uncovered a coupled mechanism underlying epithelial invagination driven by apicobasal tension imbalance and lateral contraction. We thus propose an actomyosin translocation mechanical model: lateral actomyosin first translocate apically to drive apical constriction and shape the initial invagination, then apical actomyosin redistributes laterally to promote lateral contractility and accelerate invagination. Our findings discovery a bidirectional reorganization of actomyosin network as a central mechanism driving epithelial invagination, providing insights on epithelial invagination and the organ morphogenesis during development.

Keywords

Ciona; atrial siphon; epithelial invagination; translocation; vertex model

Introduction

Morphogenesis, by which cells and tissues acquire their shape and structure, is a central theme in developmental biology. The tissue shape transformation in morphogenesis is fundamentally driven by the spatiotemporally regulated biomechanical forces(Collinet & Lecuit, 2021; Matter & Balda, 2003; Štorgel et al., 2016). The generation and conduction of biomechanical forces play as an essential mechanism of morphogenesis, via regulating cell rearrangement, division, and deformation at both cellular and tissue scales(Chan et al., 2017; Lecuit et al., 2011; Matter & Balda, 2003; Nagafuchi, 2001; Osswald et al., 2022; Tepass, 2002). Among these, actomyosin contractility plays a pivotal structure generating biomechanical tension in subcellular level, which is the basic of cell and tissue-level morphogenesis(Chung et al., 2017; Even-Ram et al., 2007; Fristrom, 1988; Lee & Harland, 2007; Popkova et al., 2024; Vicente-Manzanares et al., 2009; Zartman & Shvartsman, 2010). Despite the widely acknowledged critical role of bio-mechanical forces in morphogenesis, the precise mechanisms by which these forces regulate cell and tissue remodeling, particularly how cells reorganize contractile domains in space and time to coordinate sequential morphogenetic events remains a significant challenge.

A typical example of morphogenesis is the transition from simple epithelial sheets to complex three-dimensional architectures(Collinet & Lecuit, 2021). A large number of previous researches demonstrated the function of apical constriction in epithelial sheet invagination, such as in the invagination process of the *Drosophila* embryonic trachea(Schottenfeld et al., 2010) and salivary gland placode(Pearl et al., 2017). However, more and more studies noticed that the apical constriction itself is not sufficient in shaping complex epithelial three-dimensional structures and various epithelial tube shapes(Kondo & Hayashi, 2013; Pearl et al., 2017). So, an interesting question is how do epithelial cells preform a spatiotemporal heterogeneous force generating program to drive a sequential epithelial tissue reshaping.

Marine ascidian *Ciona* is an excellent model for studying morphogenesis due to its simple developmental process, transparent embryos, and its close evolutionary relationship with vertebrates(Blair & Hedges, 2005; Delsuc et al.,

2006; Zhao et al., 2021). For example, asymmetrical actomyosin contractility in

notochord was demonstrated to provide force for tail bending(Lu et al., 2020;

Peng et al., 2020). Actomyosin sequentially localized on the apical and

basolateral cell surfaces to drive the endodermal invagination(Fiuza & Lemaire,

2021; Hotta et al., 2007; Kourakis et al., 2010). Local myosin activation

coupled with junctional rearrangements drives directional zippering(Hashimoto

& Munro, 2019; Hashimoto et al., 2015) in neural tube closure, and cell cortex

distribution and the stability of tight junctions (TJs) were essential for

notochord tube lumen opening and expansion(Shi et al., 2025).

Ciona and vertebrates share evolutionary developmental homology. There

may be a potential homology between the otic placode of vertebrates and the

atrial siphon primordium of *Ciona*(Kourakis et al., 2010). The atrial siphon

forms from a non-dividing region in the lateral-dorsal epidermis of the head

and follows a separate developmental trajectory(Hotta et al., 2020; Kourakis et

al., 2010). Invagination initially generates shallow pits, which later deepen and

connect to the gut lumen after metamorphosis, creating two atrial

siphons(Hotta et al., 2007; Kourakis et al., 2010). Eventually, the left and right

siphons fuse to form a single atrial siphon(Chiba et al., 2004). Despite these

observations, the early cellular events during atrial siphon tube morphogenesis

remain incompletely understood.

In this study, we investigated the morphogenesis of the *Ciona* atrial siphon.

Firstly, the detailed cellular process of *Ciona* atrial siphon formation through

initial and the accelerated stages was described by visualizing cell boundary

and nuclei. By combining actomyosin localization analysis, genetic

manipulation, and vertex model simulations, we demonstrated that lateral

actomyosin first translocated apically to drive apical constriction and

established the initial invagination, then redistributed laterally to promote cell

shortening and the further atrial siphon tube invagination. Disruption of myosin

activity altered invagination timing, while optogenetic inhibition of myosin

activity after initial apical constriction stage blocked the further invagination.

Furthermore, a cell-based vertex model was established to validate the

relationship between contractile force redistribution and epithelial invagination.

The coupled biomechanical mechanisms induced by apicobasal imbalance

and lateral contraction were uncovered as fundamental determinants of the

atrial siphon invagination. These results provide new insights into the biomechanical control of tissue morphogenesis, and highlight the value of marine model organisms for addressing fundamental questions in developmental biology.

Results

The cellular processes of *Ciona* atrial siphon invagination

To investigate the early cellular events during atrial siphon morphogenesis, we performed actin filament (F-actin) staining on the fixed samples to capture the detailed cellular dynamics (Figure 1A). The cell at the bottom center of the invaginating region was defined as the "center cell" (Figure 1B). Measurements taken at different stages included the height and apical-to-basal area ratio of the center cell, the invagination depth of the atrial siphon and the linear distance between the -3/-4 and +3/+4 cell junctions (Figure 1B-E, Figure 1—figure supplement 1). At the initial invagination stage (13.5-16.0 hpf), the center cell height increased, the apical-to-basal area ratio decreased, and the protruding cell surface became inward pit, but no significant invagination occurred (Figure 1A), as reflected by a low invagination slope (k = 0.2617). During the accelerated stage (16.0-18.0 hpf), the center cell height decreased, the apical-to-basal area ratio stabilized, and invagination progressed rapidly (k = 2.7920). EdU incorporation and TUNEL assays demonstrated that neither cell division nor apoptosis was involved in the whole invagination process (Figure 1—figure supplement 2).

Bidirectional translocation of actomyosin between apical and lateral domains during atrial siphon tube invagination

To analyze the forces driving atrial siphon tube invagination, we first visualized the distribution of F-actin and quantified its dynamics at different cellular domains (Figure 1—figure supplement 1A). During the initial invagination stage, before 15 hpf, F-actin concentration decreased at the lateral domains, while gradually increased around the apical membrane, resulting in a higher F-actin accumulation at the apical region. Subsequently, the F-actin concentration at lateral region started to increase and exceeded that in the

apical domain after 16 hpf, during which apical F-actin levels showed a gradual
decline (Figure 1—figure supplement 1A).

Using anti-pS19 MRLC antibody, we examined the spatial and temporal
patterns of myosin activity within the atrial siphon primordium. Quantification of
signal intensities from the apical, basal, and lateral regions during the initial
and the accelerated stages showed that the distribution pattern of active
myosin was consistent with that of F-actin (Figure 2A, B). Specifically, active
myosin and F-actin levels were higher at the apical region than the lateral one
during the initial stage, whereas the opposite pattern was observed during the
accelerated stage (Figure 2B).

Additionally, top view imaging showed that during the initial stage, atrial
siphon primordium cells were arranged in a circular pattern (Figure 2A). Active
myosin exhibited a similar ring-like localization, suggesting that the early
stages of atrial siphon morphogenesis are driven by circumferential contractile
forces generated by phosphorylated myosin, facilitating inward constriction of
the primordium (Figure 2A, B). Furthermore, quantification of the lateral cell
distance (Figure 1—figure supplement 1B) demonstrated a reduction in
spacing between peripheral cells, indicating that surrounding cells moved
centripetally toward the center position.

Integrating the localization of actomyosin with the observed cell shape
changes, we propose a hypothesis in which the bidirectional reorganization of
contractile forces at the apical and lateral regions drives epithelial invagination
during atrial siphon morphogenesis (Figure 2C). Initially, actomyosin
translocated from the lateral regions to the apical domains, generating
contractile forces for apical constriction (Figure 1—figure supplement 1A).
Meanwhile, the inward compression from surrounding cells facilitates cell
elongation, establishing the initial invaginated cell morphology and preparing
for subsequent morphogenesis (Figure 1—figure supplement 1B). Then,
actomyosin translocated from the apical domains to the lateral regions,
generating contractile forces that promoted center cell shortening and
accelerate the tissue invagination (Figure 2C).

Center cell height is coupled with invagination depth
To further verify our hypothesis and explore the relationship between myosin

contractility, force redistribution, and the progression of invagination, we
overexpressed MRLC (T18ES19E)::mCherry (a diphosphorylated mutant that
enhances myosin activity)(Espinoza-Fonseca et al., 2014), MRLC
(T18AS19A)::mCherry (an unphosphorylated mutant that reduces myosin
activity)(Iwasaki et al., 2001) and MRLC::mCherry (wild-type control) in the
atrial siphon primordium to modulate contractile force in embryos, respectively
(Figure 3A, B). The results showed that the invagination initiated earlier (15 hpf)
compared to control group in MRLC (T18ES19E)-expressed embryos. At 16
hpf, MRLC (T18ES19E) group exhibited a deeper invagination and shorter
center cell height. In contrast, in MRLC (T18AS19A) group, the invagination
initiation time was delayed. Even at 16 hpf, the center cell height did not
decrease and the invagination did not occur (Figure 3B). Notably, the increase
in invagination depth is strongly correlated with the reduction in center cell
height, with both changes occurring synchronously across experimental
groups with different myosin activities (Figure 3B). These results demonstrate
a strong coupling between the reduction in center cell height and the increase
in invagination depth, suggesting that this process is closely associated with
the redistribution of myosin localization from the apical to lateral regions as
invagination progresses.
200

201 Inhibition of myosin activity during apical-to-lateral redistribution
202 impedes invagination progression

To further examine the necessity of the redistribution of myosin between the
apical and lateral regions during invagination, we utilized an optogenetic
MLCP-BcLOV4(Berlew et al., 2021; Berlew et al., 2020; Berlew et al., 2022;
Glantz et al., 2019; Glantz et al., 2018; Yamamoto et al., 2021) system (Figure
4A) to specifically inhibit myosin II activity at the critical redistribution time point
between the initial invagination and the accelerated invagination stage (16-17
hpf). This system has been proven to effectively reduce the contractility of
epidermal cells in *Ciona* embryos(Qiao et al., 2023). Compared to the control
group (Figure 4—figure supplement 1) and dark treatment group, the
experimental group with 1 h light exposure, which activated the optogenetic
system, blocked further invagination (Figure 4B, Figure 4—video 1, 2), and
while invagination ceased to deepen, the center cell height did not show a

significant decrease (Figure 4C, D). These results confirm that the
redistribution of myosin contractility from the apical to lateral regions is
essential for the development of invagination.
218

Vertex model simulations recapitulate the mechanical process of
ascidian siphon tube invagination
To validate the hypothesis that the redistribution of contractile forces drives
invagination, we developed a cell-based vertex model(Alt et al., 2017) and
performed simulations for atrial siphon morphogenesis. The cross-section of
the epithelial cells is represented by two-dimensional polygons with vertices
and edges. The interaction between the epithelia and the other cells below,
which function as structural support, are simplified as a basal elastic force
(Figure 5A). Cell deformation and tissue morphology are simulated by the
motion of vertices, which is controlled by the constraint of cell area and cortical
contraction, tissue surface bending, and most importantly, active line tension
dynamics (Figure 5A, see Methods). The active tension $\Gamma_J^{a,b,l}$, on the apical,
lateral and basal edges of a specific cell J, are induced by dynamic production
and distribution of actomyosin $m_J^{a,b,l}$ on corresponding regions through a
tension coefficient k_m (Figure 5B). Here we use the apical and lateral F-actin
fluorescence intensities normalized by inactive basal intensity measured in
experiments (Figure 1—figure supplement 1A) to describe the evolution of
actomyosin intensity in the center cell of the model. The actomyosin intensity
in adjacent 9 cells (-4 to +4) are linearized from inactive to active $m_{exp}^{a,l}$ (see
Methods). We applied the periodic boundary conditions along x axis to avoid
the boundary effect, and reduced the periodic length in simulations to mimic
the decreased distance between adjacent cells obtained from experiments
(Figure 1—figure supplement 1B). The numerical simulations were conducted
based on an initial flat epithelial shape consisting of $N=25$ cells at 13 hpf to
examine the morphological processes driven by active tensions (see
Methods).
Our mechanical model successfully reproduced the invagination of the
atrial siphon observed in experiments (Figure 5C, D, Figure 5—Video 1).
During the initial stage, the rapid accumulation of actomyosin on the apical

surface gave rise to apical tension. The active tension induced the contraction

of central region and led the adjacent cells to migrate to the center cells, which

induced the cell elongation and apicobasal imbalance, breaking the apicobasal

symmetry and initializing the invagination into the basal side. After 15 hpf, the

lateral actomyosin intensity increased significantly and induced high tension on

the lateral surface, pulling cells and shortening the cell height. The coupling

apico-lateral contraction further facilitated the invagination, which gave rise to

the siphon-like epithelial shape formation at 18 hpf (Figure 5D). The

invagination depth and center cell height were quantified during this process

and were consistent with the experimentally measured data (Figure 5C). We

further varied both apical and lateral actomyosin intensities m by introducing

a scaling factor α to experimental measured m_{exp} as $m = \alpha m_{\mathrm{exp}}$ into the

model, to simulate unphosphorylated or diphosphorylated mutants whose

overall myosin activities in cells are altered in experiments (Figure 3). As

anticipated, simultaneous enhancement of apical and lateral actomyosin

activities significantly increased invagination depth while reducing center cell

height (and vice versa), closely corresponding with experimental mutant data

(Figure 5E, F). These results validate the validity of our model in capturing the

mechanical process of ascidian siphon tube invagination.

Coupled mechanism of apicobasal bending and lateral contraction via

actomyosin redistribution

To theoretically distinguish the effects of apical and lateral contraction on

morphogenesis, we performed perturbation on the intensity of regional

actomyosin. The apical actomyosin intensity m^{a} was modulated as $m^{\mathrm{a}} = \alpha_{\mathrm{a}} m_{\mathrm{exp}}^{\mathrm{a}}$,

where $m_{\mathrm{exp}}^{\mathrm{a}}$ represented the experimental data shown in Figure 5B. As a result,

the invagination depth increased significantly under higher α_{a} (Figure 6A),

while the central height remained relatively independent of α_{a} when $\alpha_{\mathrm{a}} \geq 0.6$

(Figure 6B). Intriguingly, the depth value became negative when $\alpha_{\mathrm{a}} \leq 0.2$,

which meant the invagination cannot occur but was replaced by evagination,

where the decrease of the central height after 15 hpf also disappeared. This

inversion of the tissue shape was attributed to the inversion of apico-basal

differential of actomyosin intensity (Figure 6C), in which a low α_{a} resulted in

apical actomyosin intensity falling below the reference basal intensity. These

results suggest a bending mode under an equivalent moment induced by

apico-basal tension imbalance, which breaks tissue symmetry and determines

the buckling direction, facilitating the invagination without obviously affecting

the cell height. In contrast, under modified lateral actomyosin, the central

height varied distinctly (Figure 6D) with mild invagination depth variation

(Figure 5B). This indicated that lateral tension primarily facilitates tissue

invagination by regulating cellular contraction along the thickness direction,

representing a lateral contraction mode. Collectively, these results

demonstrate that the invagination of the atrial siphon is mediated by the

coupled biomechanical mechanism of a bending mode induced by apicobasal

tension imbalance and a contraction mode driven by lateral tension.

Finally, to clarify the characteristics of actomyosin redistribution between

apical and lateral regions, we calculated the total actomyosin intensity

$S_c = m_c^{\text{a}} + m_c^{\text{l}}$ and the ratio of apical and lateral intensity $R_c = m_c^{\text{a}} / m_c^{\text{l}}$ in the center

cell (Figure 6E). During the whole morphogenesis, S_c increased persistently,

providing the driving force underling epithelial invagination. While a significant

peak value in the curve of R_c was observed during the initial stage,

representing the translocation from lateral to apical regions during 14-15 hpf

and a reverse translocation during 15-16 hpf (Figure 6E). To investigate the

translocation effect quantitively, we modified the translocation level after 14 hpf

in the model as $R_c = \beta(R_{\text{exp}} - 1) + 1$, where β is the translocation intensity

controlling the redistribution ratio. As shown in Figure 6F, varying total

actomyosin $S_c = \alpha S_{\text{exp}}$ and apical-lateral translocation R_c controlled by α

and β, respectively, were introduced into simulations. Computational results

indicated that the overall invagination magnitude (Figure 6G) is predominantly

regulated by the total actomyosin intensity within cells, whereas stronger

regional translocation and redistribution contribute to the rapid reduction in

height of invaginating cells (Figure 6H), thereby ensuring direct cell shape

changes that support robust atrial siphon formation in the ascidian.

Discussion

The morphogenesis of the *Ciona* atrial siphon provides a compelling model to dissect how biomechanical forces orchestrate tissue invagination. Our study revealed that actomyosin contractility underwent a dynamic spatial redistribution during this process, initially translocating from the lateral cortex to the apical region to promote apical constriction and establish the initial cell shape, and subsequently redistributing back to the lateral regions to promote center cell shortening and accelerate the tissue invagination. By combining actomyosin localization analysis, genetic perturbations, optogenetic manipulation, and vertex model, we established a mechanistic framework linking force dynamics to tissue remodeling. This work not only advances our understanding of siphon morphogenesis but also offers broader insights into the principles governing force-driven tissue shaping in developing organisms.

Our findings demonstrated that the atrial siphon invagination was driven by a coupled mechanism including apicobasal tension imbalance and lateral contraction in a two-stage process. During the initial stage (13.5-16.0 hpf), actomyosin translocated from the lateral cortex to the apical cortex, generated contractile forces that induced apicobasal tension imbalance and reduced apical cell area and elongated the primordium. This aligns with classical models of epithelial folding, where apical actomyosin networks drive tissue curvature through localized contraction, often accompanied by an increase in cell height, as observed in the invagination of the *Drosophila* embryonic trachea(Schottenfeld et al., 2010) and salivary gland placode(Pearl et al., 2017). Additionally, in traditional invagination models, various epithelial tissues use different mechanisms to drive deeper invagination. In *Drosophila* tracheal development, for instance, mitosis acts as a critical turning point in accelerating invagination(Kondo & Hayashi, 2013). During the early slow phase of tracheal invagination, apical constriction under EGFR signaling, forms a shallow pit. As invagination transitions into a rapid phase, mitotic entry induces cell rounding, which increases tension and epithelial buckling, thereby accelerating the invagination process and facilitating the internalization of placode cells(Kondo & Hayashi, 2013; Nishimura et al., 2007). Moreover, cell apoptosis can also expedite the invagination process, as seen in *Drosophila*

leg morphogenesis, where actomyosin cables formed in apoptotic cells help

trigger the invagination(Kiehart, 2015; Manjón et al., 2007; Monier et al., 2015).

Distinct from these classical models, the accelerated invagination stage of

Ciona atrial siphon morphogenesis was driven by a critical redistribution of

apical actomyosin to the lateral regions, during accelerated stage (16.0–18.0

hpf), providing a force-generating mechanism for invagination progression

without relying on cell proliferation or apoptosis to accelerate the process. The

lateral enrichment of p-MLC correlated with a reduction in center cell height

and the initiation of invagination, suggesting that lateral forces actively pulled

the tissue inward. This spatial redistribution of contractile forces in *Ciona* atrial

siphon invagination is reminiscent of mechanisms observed in certain

epithelial folding processes, such as the *Drosophila* wing disc, where

invagination is not solely driven by apical constriction but can also result from

increased lateral tension or reduced basal tension(Sui et al., 2018). Our

optogenetic inhibition of myosin activity during the apical-to-lateral

redistribution—using the MLCP-BcLOV4 system—directly confirmed that this

redistribution was not merely a passive consequence of invagination but was

an active driver of the process.

Our study revealed a strong coupling between cellular shape changes and

tissue remodeling during *Ciona* atrial siphon invagination, where the reduction

in center cell height was closely associated with invagination depth. Tissue

morphogenesis emerges from the integrated actions of multiple cellular

behaviors, such as contraction, adhesion, and migration, that work in concert

to remodel epithelial architecture(Chan et al., 2017). The transmission of local

cell-generated forces across tissues is further shaped by mechanical cues and

geometric constraints, enabling the transformation of cellular-scale changes

into organ-scale morphogenesis(Collinet & Lecuit, 2021). By perturbing

myosin activity, we found that this coupling was directly regulated:

dominant-negative myosin mutants delayed both processes, whereas

hyperactive myosin accelerated invagination. The quantitative agreement

between vertex model analysis and experimental measurements validated the

critical role of actomyosin contractility dynamics in linking cellular shape

changes to tissue remodeling. Furthermore, the model also uncoupled the

effects of the actomyosin in different domains, and predicted that stronger

apical-lateral translocation facilitates the rapid reduction in height of
invaginating cell. The *Ciona* atrial siphon, which underwent invagination
without cell proliferation or apoptosis, provided a minimal system for
investigating the role of biomechanical forces in tissue remodeling. Our
findings emphasized that the deformation of a few key cells under spatially
precise mechanical regulation, along with their coordination with surrounding
tissues, is crucial for proper morphogenesis.

While this study clarified the role of actomyosin redistribution in atrial
siphon invagination, several open questions remain regarding the molecular
mechanisms underlying this process. One key question is what molecular
signals drive the apical-to-lateral redistribution of contractility. Possible
candidates include Rho GTPase pathways(Martin et al., 2016), which regulate
myosin activation; apical-basal polarity(Peng et al., 2020), which coordinate
actin reorganization during morphogenesis; and mechanosensitive ion
channels that respond to tissue strain(Jin et al., 2020). Additionally, the role of
cell-cell adhesion and extracellular matrix remodeling in mediating force
transmission during invagination requires further investigation. Using tension
sensors or perturbing adhesion molecules may provide new insights into these
interactions. Although our vertex model effectively predicts the contribution of
force redistribution to invagination, the spatiotemporal evolution of actomyosin
could also be affected by mechanosensitive pathways while generating active
contraction in morphogenesis, which constructs a bidirectional loop of
actomyosin regulation. Future theoretical developments should focus on
integrating mechanochemical feedback linking molecular signaling with
tissue-level deformation to explain robust morphogenesis.

Materials and Methods

Experimental animals' preparation and electroporation

Ciona adults were collected from the coast of Qingdao and Rongcheng,
Shandong, China. They were maintained in the laboratory seawater circulation
system under constant light conditions to maintain their stability. Mature eggs
and sperm were separately collected from the dissected adults and
subsequently mixed for fertilization. The Fertilized eggs were dechorionated in

seawater containing 1 % sodium thioglycolate (T0632; Sigma), 0.05 %
protease E (P5147; Sigma) and 0.032 M NaOH. The dechorionated eggs were
then used for plasmid electroporation based on the previous technique
procedure(Christiaen et al., 2009). Finally, the embryos were cultured in an
agar-coated dish with microporous-filtered seawater in 18℃ for further
observation. Detailed key resource information is annotated in Appendix
1—table 1.

Plasmid construction
Ciona myosin II light chain (MRLC) was amplified with primers MRLC-F and
MRLC-R (Table S1). MRLC-mCherry(Dong et al., 2011);
MRLC(T18AS19A)-mCherry and MRLC(T18ES19E)-mCherry(Denker et al.,
2015); MLCP-BcLOV4 and MLCP(Qiao et al., 2023) were amplified with the
primers listed in Appendix 1—table 2.

Immunofluorescence
Ciona embryos were fixed with stationary liquid(Sherrard et al., 2010), which
consisted of 100 mM HEPES (pH 6.9); 100 mM EGTA (pH 7.0); 10 mM MgSO4;
2% formaldehyde; 0.1% glutaraldehyde; 300 mM dextrose and 0.2% Triton
X-100 for 40 m at room temperature. Following fixation, the embryos were
washed three times with PBS and subsequently incubated in PBST (PBS
supplemented with 0.1% Triton X-100) for 30 m to enhance permeability. To
reduce autofluorescence, embryos were treated with 0.1% sodium
borohydride in PBS for 20 m at room temperature. For immunostaining,
embryos were incubated with a 1:250 dilution of Phospho-Myosin Light Chain
2 (Ser19) antibody (#3671: Cell Signaling) at room temperature for 24 h. After
three additional washes with PBS, a 1:200 dilution of Alexa Fluor 568
anti-Rabbit IgG (A11011: Invitrogen) was added and incubated at room
temperature for 48 h. For cell boundary visualization, embryos were stained
with Alexa Fluor 488 Phalloidin (A12379: Invitrogen) at a 1:200 dilution. Finally,
after three washes with PBS, embryos were mounted in DAPI-containing
mounting medium and prepared for imaging.

Imaging and optogenetics

Live imaging, photoactivation experiments, and image acquisition were
performed using a Zeiss LSM 980 confocal microscope (Carl Zeiss). For
optogenetic experiments, *Ciona* embryos were placed in a 35 mm
glass-bottom dish for imaging. To activate the optogenetic system, the
designated region of interest was exposed to a 488 nm laser for 1 h. The
control group for dark treatment was placed in a dark box at the same
temperature for 1 h. All images were analyzed and quantified using ImageJ
(version 1.54p, NIH) and Imaris (version 9.0.1, Bitplane).

Vertex model simulations

Model description. The epithelial cells are represented by two-dimensional
polygons with vertices and edges. Vertices on the apical surface of cells are
subject to no boundary constraints, while vertices on the basal surface interact
with basal spring forces to simulate the structural support provided by
subjacent cells (Figure 5A). The potential energy of the tissue is formulated by

461
$$U = \sum_{J} \frac{1}{2} K_{A} \left(A_{J} - A_{0} \right)^{2} + \sum_{J} \frac{1}{2} K_{C} L_{J}^{2} + \sum_{i_{a,b}} \frac{1}{2} K_{B} (\cos\theta_{i} - 1)^{2}$$
$$+ \sum_{J} (\Gamma_{J}^{a} l_{J}^{a} + \Gamma_{J}^{b} l_{J}^{b} + \frac{1}{2} \Gamma_{J}^{l} l_{J}^{l}) + \sum_{i_{b}} k_{bm} (r_{yi} + \frac{h_{0}}{2})^{2},$$
(1)

where the six terms result from, respectively, cell area control, cell cortex
contractility, surface bending energy, active line tensions on apical, basal, and
lateral surfaces, and subjacent interactions. K_{A} is the cell area modulus, and
A_{J} and A_{0} are the current and preferred areas of the J th cell, respectively.
K_{C} denotes the passive cortical contraction coefficient of cells and L_{J}
represents the cell perimeter. K_{B} denotes the bending modulus of the
monolayer and θ_{i} is the angle between the neighboring apical or basal edges
of vertex i. Γ_{J}^{a}, Γ_{J}^{b}, and Γ_{J}^{l} are the active line tensions acting on the apical,
basal, and lateral surfaces of the J th cell with edge lengths l_{J}^{a}, l_{J}^{b} and l_{J}^{l},
respectively. k_{bm} is the elastic coefficient of basal interaction and h_{0} is the
initial cell height. $\mathbf{r}_{i} = (r_{xi}, r_{yi})$ denotes the Cartesian coordinates of vertex i.
$\sum_{J}$ stands for the summation over all cells and $\sum_{i_{a,b}}$ runs over all apical and
basal vertices.

To describe the cell actomyosin evolution during the morphogenesis, the

line tensions Γ_J^a, Γ_J^b, and Γ_J^l are proportional to actomyosin levels on the

apical, basal, and lateral surfaces of the cell J as

$$\Gamma_J^a = k_m m_J^a, \ \Gamma_J^b = k_m m_J^b, \ \Gamma_J^l = k_m m_J^l, \tag{2}$$

where k_m is the tension coefficient based on the myosin intensity m_J^a, m_J^b and

m_J^l. Given that the intensity of F-actin in the center cell has been measured as

$m_{exp}^{a,b,l}$ during the development, we define $m_{exp}^{a,b,l}$ as the actomyosin intensity of

the center cell in the model (Figure 5B).

Simulation scheme. In the simulations, we set the cell number $N = 25$

and a periodic boundary condition in the x direction with periodic box length

$L = NA_0 / h_0$. The central nine cells (index -4 to +4) are considered as active

cells under contractile forces whose actomyosin intensity varies in different

stages. The other cells are observed to be inactive whose actomyosin

intensities are always equal to $m_0 = m_{exp}^b = 1$. The actomyosin intensities of the

central nine cells (-4 to +4) are calculated through linear interpolation between

inactive m_0 of boundary cells and active $m_{exp}^{a,l}$ of the most center cell

according to the distance. The periodic length L in simulations is reduced

slowly to mimic the decreased distance between adjacent cells obtained from

experiments (Figure 1—figure supplement 1B). In our simulations, tissue

morphogenesis is represented by vertices motion and cell deformation. The

motion of vertex i obeys the Langevin equation

$$\eta \frac{d\mathbf{r}_i}{dt} = -\frac{\partial U}{\partial \mathbf{r}_i}, \tag{3}$$

where η denotes the damping coefficient. The equation was numerically

solved using the forward Euler method with a time step of 0.01, using MATLAB

R2021a. Simulations are run until $t = 400$, which corresponds to 18 hpf in

experimental development with the time scale $\tau = 45\,\text{s}$. The length scale

$\varepsilon = 8.91\,\mu\text{m}$ is determined by the normalization of initial height h_0 in model

under real center cell height. The other parameters are all normalized with the

time scale $\tau = 45\,\text{s}$, length scale $\varepsilon = 8.91\,\mu\text{m}$, and damping coefficient η.

Dimensionless parameters are set as follows: $K_A = 1$, $K_C = 0.01$, $K_B = 0.03$,

$k_{\mathrm{bm}} = 0.1$, $k_{\mathrm{m}} = 0.002$, $A_0 = 1$, and $h_0 = 1.25$. These parameter values are
determined by numerical simulations under which the simulated invagination
depth and center cell height imitates the experimental observations.
508

Acknowledgements

510 We are grateful to all members of the Fang Zongxi center for helpful
511 discussions.
512

Competing interests

514 The authors declare that they have no known competing financial interests or
515 personal relationships that could have appeared to influence the work reported
516 in this paper.
517

Author contributions

519 B.D., B.L. J.Q., P.Y. conceived of the study; B.D., J.Q., P.Y., B.L., H.P., and W.S.
520 analyzed data; P.Y. and B.L. designed the vertex model; B.D., J.Q., P.Y. and
521 B.L. wrote the original draft. All authors approved the final version of the article.
522

Funding

524 This work was supported by the Science & Technology Innovation Project of
525 Laoshan Laboratory (Nos. LSKJ202203204), the National Key Research and
526 Development Program of China (2022YFC2601302), and the Taishan Scholar
527 Program of Shandong Province, China (B.D.).

Data and resource availability

Lead contact

531 Requests for further resources should be directed to, and will be fulfilled by, the
532 lead contact, Bo Dong (bodong@ouc.edu.cn).

Materials availability

534 All unique/stable reagents generated in this study are available from the lead
535 contact with a completed materials transfer agreement.

Data and code availability

Model code and data are available upon request, and any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

References

Alt S, Ganguly P, Salbreux G (2017). **Vertex models: from cell mechanics to tissue morphogenesis** *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **372**:20150520. https://doi.org/10.1098/rstb.2015.0520

Berlew E E, Kuznetsov I A, Yamada K, Bugaj L J, Boerckel J D, Chow B Y (2021). **Single-component optogenetic tools for inducible RhoA GTPase signaling** *Advanced Biology* **5**:e2100810. https://doi.org/10.1002/adbi.202100810

Berlew E E, Kuznetsov I A, Yamada K, Bugaj L J, Chow B Y (2020). **Optogenetic Rac1 engineered from membrane lipid-binding RGS-LOV for inducible lamellipodia formation** *Photochemical and Photobiological Sciences* **19**:353-361. https://doi.org/10.1039/c9pp00434c

Berlew E E, Yamada K, Kuznetsov I A, Rand E A, Ochs C C, Jaber Z, Gardner K H, Chow B Y (2022). **Designing single-component optogenetic membrane recruitment systems: The Rho-Family GTPase signaling toolbox** *ACS Synthetic Biology* **11**:515-521. https://doi.org/10.1021/acssynbio.1c00604

Blair J E, Hedges S B (2005). **Molecular phylogeny and divergence times of deuterostome**

animals *Molecular* *Biology* *and* *Eevolution* **22**:2275-2284.

https://doi.org/10.1093/molbev/msi225

Chan C J, Heisenberg C P, Hiiragi T (2017). **Coordination of morphogenesis and cell-fate**

specification in development *Current* *Biology* **27**:R1024-r1035.

https://doi.org/10.1016/j.cub.2017.07.010

Chiba S, Sasaki A, Nakayama A, Takamura K, Satoh N (2004). **Development of** *Ciona*

intestinalis **juveniles (through 2nd ascidian stage)** *Zoological Science* **21**:285-298.

https://doi.org/10.2108/zsj.21.285

Christiaen L, Wagner E, Shi W, Levine M (2009). **Electroporation of transgenic DNAs in the**

sea squirt *Ciona* *Cold* *Spring* *Harbor* *Protocols* **2009**:pdb.prot5345.

https://doi.org/10.1101/pdb.prot5345

Chung S, Kim S, Andrew D J (2017). **Uncoupling apical constriction from tissue invagination**

eLife **6**:e22235. https://doi.org/10.7554/eLife.22235

Collinet C, Lecuit T (2021). **Programmed and self-organized flow of information during**

morphogenesis *Nature* *Reviews* *Molecular* *Cell* *Biology* **22**:245-265.

https://doi.org/10.1038/s41580-020-00318-6

Delsuc F, Brinkmann H, Chourrout D, Philippe H (2006). **Tunicates and not cephalochordates**

are the closest living relatives of vertebrates *Nature* **439**:965-968.

https://doi.org/10.1038/nature04336

Denker E, Sehring I M, Dong B, Audisso J, Mathiesen B, Jiang D (2015). **Regulation by a**

TGFβ-ROCK-actomyosin axis secures a non-linear lumen expansion that is essential

for tubulogenesis *Development* **142**:1639-1650. https://doi.org/10.1242/dev.117150

Dong B, Deng W, Jiang D (2011). **Distinct cytoskeleton populations and extensive crosstalk**

control *Ciona* **notochord tubulogenesis** *Development* **138**:1631-1641.

https://doi.org/10.1242/dev.057208

Espinoza-Fonseca L M, Colson B A, Thomas D D (2014). **Effects of pseudophosphorylation**

mutants on the structural dynamics of smooth muscle myosin regulatory light chain

Molecular Biosystems **10**:2693-2698. https://doi.org/10.1039/c4mb00364k

Even-Ram S, Doyle A D, Conti M A, Matsumoto K, Adelstein R S, Yamada K M (2007). **Myosin**

IIA regulates cell motility and actomyosin-microtubule crosstalk *Nature Cell Biology*

9:299-309. https://doi.org/10.1038/ncb1540

Fiuza U-M, Lemaire P (2021). **Mechanical and genetic control of ascidian endoderm**

invagination during gastrulation *Seminars in Cell and Developmental Biology*

120:108-118. https://doi.org/10.1016/j.semcdb.2021.08.001

Fristrom D (1988). **The cellular basis of epithelial morphogenesis. A review** *Tissue and Cell*

20:645-690. https://doi.org/10.1016/0040-8166(88)90015-8

Glantz S T, Berlew E E, Chow B Y (2019). **Synthetic cell-like membrane interfaces for probing**

dynamic protein-lipid interactions *Methods in Enzymology* **622**:249-270.

https://doi.org/10.1016/bs.mie.2019.02.015

Glantz S T, Berlew E E, Jaber Z, Schuster B S, Gardner K H, Chow B Y (2018). **Directly**

light-regulated binding of RGS-LOV photoreceptors to anionic membrane

phospholipids *Proceedings of the National Academy of Sciences of the United States*

of America **115**:E7720-e7727. https://doi.org/10.1073/pnas.1802832115

Hashimoto H, Munro E (2019). **Differential expression of a classic cadherin directs tissue-level**

contractile asymmetry during neural tube closure *Developmental Cell*

51:158-172.e154. https://doi.org/10.1016/j.devcel.2019.10.001

Hashimoto H, Robin F B, Sherrard K M, Munro E M (2015). **Sequential contraction and**

exchange of apical junctions drives zippering and neural tube closure in a simple

chordate *Developmental Cell* **32**:241-255.

https://doi.org/10.1016/j.devcel.2014.12.017

Hotta K, Dauga D, Manni L (2020). **The ontology of the anatomy and development of the**

solitary ascidian *Ciona*: **the swimming larva and its metamorphosis** *Scientific Reports*

10:17916. https://doi.org/10.1038/s41598-020-73544-9

Hotta K, Mitsuhara K, Takahashi H, Inaba K, Oka K, Gojobori T, Ikeo K (2007). **A web-based**

interactive developmental table for the ascidian *Ciona intestinalis*, **including 3D**

real-image embryo reconstructions: I. From fertilized egg to hatching larva

Developmental Dynamics **236**:1790-1805. https://doi.org/10.1002/dvdy.21188

Iwasaki T, Murata-Hori M, Ishitobi S, Hosoya H (2001). **Diphosphorylated MRLC is required for**

organization of stress fibers in interphase cells and the contractile ring in dividing cells

Cell Structure and Function **26**:677-683. https://doi.org/10.1247/csf.26.677

Jin P, Jan L Y, Jan Y N (2020). **Mechanosensitive ion channels: structural features relevant to**

mechanotransduction mechanisms *Annual Review of Neuroscience* **43**:207-229.

https://doi.org/10.1146/annurev-neuro-070918-050509

Kiehart D P (2015). **Epithelial morphogenesis: apoptotic forces drive cell shape changes**

Developmental Cell **32**:532-533. https://doi.org/10.1016/j.devcel.2015.02.020

Kondo T, Hayashi S (2013). **Mitotic cell rounding accelerates epithelial invagination** *Nature*

494:125-129. https://doi.org/10.1038/nature11792

Kourakis M J, Newman-Smith E, Smith W C (2010). **Key steps in the morphogenesis of a**

cranial placode in an invertebrate chordate, the tunicate *Ciona savignyi*

Developmental Biology 340:134-144. https://doi.org/10.1016/j.ydbio.2010.01.016

Lecuit T, Lenne P F, Munro E (2011). **Force generation, transmission, and integration during**

cell and tissue morphogenesis *Annual Review of Cell and Developmental Biology*

27:157-184. https://doi.org/10.1146/annurev-cellbio-100109-104027

Lee J Y, Harland R M (2007). **Actomyosin contractility and microtubules drive apical**

constriction in *Xenopus* **bottle cells** *Developmental Biology* **311**:40-52.

https://doi.org/10.1016/j.ydbio.2007.08.010

Lu Q, Gao Y, Fu Y, Peng H, Shi W, Li B, Lv Z, Feng X Q, Dong B (2020). *Ciona* embryonic tail

bending is driven by asymmetrical notochord contractility and coordinated by epithelial

proliferation *Development* 147:dev185868. https://doi.org/10.1242/dev.185868

Manjón C, Sánchez-Herrero E, Suzanne M (2007). **Sharp boundaries of Dpp signalling trigger**

local cell death required for *Drosophila* **leg morphogenesis** *Nature Cell Biology*

9:57-63. https://doi.org/10.1038/ncb1518

Martin K, Reimann A, Fritz R D, Ryu H, Jeon N L, Pertz O (2016). **Spatio-temporal**

co-ordination of RhoA, Rac1 and Cdc42 activation during prototypical edge protrusion

and retraction dynamics *Scientific Reports* **6**:21901. https://doi.org/10.1038/srep21901

Matter K, Balda M S (2003). **Functional analysis of tight junctions** *Methods* **30**:228-234.

https://doi.org/10.1016/s1046-2023(03)00029-x

Monier B, Gettings M, Gay G, Mangeat T, Schott S, Guarner A, Suzanne M (2015).

Apico-basal forces exerted by apoptotic cells drive epithelium folding *Nature*

518:245-248. https://doi.org/10.1038/nature14152

Nagafuchi A (2001). **Molecular architecture of adherens junctions** *Current Opinion in Cell*

Biology **13**:600-603. https://doi.org/10.1016/s0955-0674(00)00257-x

Nishimura M, Inoue Y, Hayashi S (2007). **A wave of EGFR signaling determines cell alignment**

and intercalation in the *Drosophila* **tracheal placode** *Development* **134**:4273-4282.

https://doi.org/10.1242/dev.010397

Osswald M, Barros-Carvalho A, Carmo A M, Loyer N, Gracio P C, Sunkel C E, Homem C C F,

Januschke J, Morais-de-Sá E (2022). **aPKC regulates apical constriction to prevent**

tissue rupture in the *Drosophila* follicular epithelium *Current Biology*

32:4411-4427.e4418. https://doi.org/10.1016/j.cub.2022.08.063

Pearl E J, Li J, Green J B (2017). **Cellular systems for epithelial invagination** *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences

372:20150526. https://doi.org/10.1098/rstb.2015.0526

Peng H, Qiao R, Dong B (2020). **Polarity establishment and maintenance in ascidian**

notochord *Frontiers in Cell and Developmental Biology* **8**:597446.

https://doi.org/10.3389/fcell.2020.597446

Popkova A, Andrenšek U, Pagnotta S, Ziherl P, Krajnc M, Rauzi M (2024). **A mechanical wave**

travels along a genetic guide to drive the formation of an epithelial furrow during

Drosophila **gastrulation** *Developmental Cell* **59**:400-414.e405.

https://doi.org/10.1016/j.devcel.2023.12.016

Qiao J, Peng H, Dong B (2023). **Development and application of an optogenetic manipulation**

system to suppress actomyosin activity in *Ciona* **epidermis** *International Journal of*

Molecular Sciences **24**:5707. https://doi.org/10.3390/ijms24065707

Schottenfeld J, Song Y, Ghabrial A S (2010). **Tube continued: morphogenesis of the**

Drosophila **tracheal system** *Current Opinion in Cell Biology* **22**:633-639.

https://doi.org/10.1016/j.ceb.2010.07.016

Sherrard K, Robin F, Lemaire P, Munro E (2010). **Sequential activation of apical and**

basolateral contractility drives ascidian endoderm invagination *Current Biology*

20:1499-1510. https://doi.org/10.1016/j.cub.2010.06.075

Shi W, Duclut C, Xu Y, Ma Y, Qiao J, Lin B, Yang D, Prost J, Dong B (2025). **Physics of**

notochord tube expansion in ascidians *Proceedings of the National Academy of*

Sciences   of   the   United   States   of   America   **122**:e2419960122.

https://doi.org/10.1073/pnas.2419960122

Štorgel N, Krajnc M, Mrak P, Štrus J, Ziherl P (2016). **Quantitative morphology of epithelial**

folds *Biophysical Journal* **110**:269-277. https://doi.org/10.1016/j.bpj.2015.11.024

Sui L, Alt S, Weigert M, Dye N, Eaton S, Jug F, Myers E W, Jülicher F, Salbreux G, Dahmann

C (2018). **Differential lateral and basal tension drive folding of** *Drosophila* **wing discs**

through   two   distinct   mechanisms *Nature   Communications*   **9**:4620.

https://doi.org/10.1038/s41467-018-06497-3

Tepass U (2002). **Adherens junctions: new insight into assembly, modulation and function**

Bioessays **24**:690-695. https://doi.org/10.1002/bies.10129

Vicente-Manzanares M, Ma X, Adelstein R S, Horwitz A R (2009). **Non-muscle myosin II takes**

centre stage in cell adhesion and migration *Nature Reviews Molecular Cell Biology*

10:778-790. https://doi.org/10.1038/nrm2786

Yamamoto K, Miura H, Ishida M, Mii Y, Kinoshita N, Takada S, Ueno N, Sawai S, Kondo Y,

Aoki K (2021). **Optogenetic relaxation of actomyosin contractility uncovers**

mechanistic roles of cortical tension during cytokinesis *Nature Communications*

12:7145. https://doi.org/10.1038/s41467-021-27458-3

Zartman J J, Shvartsman S Y (2010). **Unit operations of tissue development: epithelial folding**

Annual Review of Chemical and Biomolecular Engineering **1**:231-246.

https://doi.org/10.1146/annurev-chembioeng-073009-100919

Zhao L, Gao F, Gao S, Liang Y, Long H, Lv Z, Su Y, Ye N, Zhang L, Zhao C, Wang X, Song W,

Zhang S, Dong B (2021). **Biodiversity-based development and evolution: the**

emerging research systems in model and non-model organisms *Science China. Life*

Sciences **64**:1236-1280. https://doi.org/10.1007/s11427-020-1915-y

Figure legends

Figure 1. Cellular processes of *Ciona* atrial siphon tube invagination.

(A) Representative images of atrial siphon morphogenesis in *Ciona* embryos from 13.5 hpf to 18 hpf. Scale bar: 10 µm.

(B) Measurement parameters of the *Ciona* atrial siphon. The cell undergoing the most prominent apical constriction at the center of invagination was defined as the center cell (0). The adjacent cells on the left and right were defined sequentially as -1, -2, -3, -4 and +1, +2, +3, +4, respectively.

(C) Quantification of the invagination depth in the atrial siphon of *Ciona*

embryos. Red lines indicate linear regression fits of invagination depth during

the initial (13.5-16 hpf, k = 0.2617) and accelerated (16-18 hpf, k = 2.7920)

stages. n = 20.

(D, E) Quantification of the center cell height and apical-to-basal area ratio at

the atrial siphon of *Ciona* embryos. The blue-shaded region represents the

initial stage, while the orange-shaded region indicates the accelerated stage. n

= 20.

Figure 1—figure supplement 1. Quantification of F-actin intensity and

intercellular distance during *Ciona* atrial siphon morphogenesis.

(A) Normalized F-actin intensity at the apical and lateral regions of center cells

during *Ciona* atrial siphon morphogenesis (basal level set to 1).

(B) Quantification of the linear distance between the -3/-4 and +3/+4 cell

junctions at the apical or basal surface in the atrial siphon of *Ciona* embryos.

The blue-shaded region represents the initial stage, while the orange-shaded

region indicates the accelerated stage. Representative images are shown in

Figure 1A. n = 20.

Figure 1—figure supplement 2. EdU and TUNEL staining during *Ciona* atrial siphon morphogenesis.

(A) Representative images of EdU staining at 14-15 hpf. Orange: EdU-positive nuclei indicating cell proliferation. Blue: DAPI. No EdU signal was detected in the atrial siphon primordium (white dashed outline). Scale bar: 10 μm. n = 10.

(B1-3) Representative images of TUNEL staining at 15 hpf. (B1) Positive control: DNase I pretreatment (20 U/mL, 10 min) induced DNA fragmentation. (B2) Negative control: staining performed without terminal deoxynucleotidyl transferase (TdT) enzyme. (B3) Experimental group: no detectable TUNEL signal in the atrial siphon primordium (white dashed outline). Scale bar: 10 μm. n = 10.

805
806
Figure 2. Bidirectional translocation of actomyosin between apical and
lateral domains during atrial siphon tube invagination.
(A) Representative images of *Ciona* embryos stained for active myosin II
(anti-pS19 MRLC，red) and F-actin (green) at different stages of atrial siphon
invagination. The heatmap color scale represents the fluorescence intensity of
myosin II signal (8-bit grayscale range: 0-255), with red indicating the highest
intensity. Scale bar: 10 μm.
(B) Normalized fluorescence intensity of active myosin II (anti-pS19 MRLC)
and F-actin at the apical and lateral regions of center cells during different
stages of atrial siphon invagination (basal level set to 1). **p < 0.01, ***p <
0.001, ****p < 0.0001, n = 20.
(C) Schematic model illustrating the mechanical forces driving atrial siphon
primordium invagination. Brown arrows indicate the direction of contractile
forces; red areas depict active myosin II localization.
821

Figure 3. Modulation of atrial siphon invagination by overexpression of myosin mutants

(A) Representative images of *Ciona* atrial siphon primordium expressing wild-type and mutant MRLC constructs. Scale bar: 10 µm.

(B) Quantification of invagination depth and the center cell height in MRLC (T18ES19E), MRLC (T18AS19A), and MRLC groups. The blue-shaded region represents the initial stage, while the orange-shaded region indicates the accelerated stage. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Figure 4. Disruption of contractile forces during rapid invagination of the *Ciona* atrial siphon using an optogenetic system

(A) Schematic diagram depicting the structure and mechanism of the MLCP-BcLOV4 system. The PP1C::MYPT169::BcLOV4::mCherry::NES fusion protein is initially dispersed in the cytoplasm. Upon exposure to blue light, BcLOV4 undergoes a conformational change, allowing it to interact electrostatically with the plasma membrane. This leads to the recruitment of the PP1C and MYPT169 components, subunits of myosin light chain phosphatase (MLCP), to the membrane, where they reduce myosin activity.

(B) Representative images of developmental progression in the MLCP-BcLOV4 expression group exposed to blue light for 1 h, and in the dark control group maintained in darkness for 1 h. Scale bar: 10 μm.

(C, D) Quantification of invagination depth and the center cell height in the MLCP-BcLOV4 expression group, the dark control group and MLCP control group (Figure 4—figure supplement 1). *p < 0.05, ***p < 0.001, ****p < 0.0001.

849
850

Figure 4—figure supplement 1. MRCL control group in the optogenetic experiment

(A) Schematic diagram depicting the structure and mechanism of the MLCP control system. The PP1C::MYPT169::mCherry::NES fusion protein remained diffuse in the cytoplasm under light exposure and failed to function.

(B) Representative images of developmental progression in the MLCP control group exposed to blue light for 1 h. Scale bar: 10 μm.

Figure 4—video 1. The developmental processes of the MLCP control group under blue light illumination for 1 h. Scale bar: 10 μm.

Figure 4—video 2. The developmental processes of MLCP-BcLOV4-expressed group under blue light illumination for 1 h. Scale bar: 10 μm.

865
866

Figure 5. Simulations and analysis based on vertex model

(A) Schematic of the cell-based vertex model. The epithelial tissue is constructed by the apical, basal, and lateral vertices and edges. The effective energy U takes into cell area constraint (modulus K_A), passive cortical contraction (coefficient K_C), tissue surface bending (modulus K_B), basal interaction (coefficient k_{bm}), and apical, basal and lateral active tensions (Γ^a, Γ^l and Γ^b, respectively). $\mathbf{r}_i = (r_{xi}, r_{yi})$ denotes the Cartesian coordinates of vertex i. A_J, L_J, and $l_J^{a,b,l}$ are the area, perimeter, and length of apical, basal,

lateral edges of cell J. θ_i represents the angle between adjacent apical or basal edges connected to vertex i. A_0 is the reference area and h_0 is the initial cell height.

(B) Temporal evolution of actomyosin intensities at the apical, lateral, and basal domains of the center cell in the model, parameterized based on experimentally measured F-actin distribution. The blue-shaded and orange-shaded regions represent the apical constriction stage, and the invagination stage, respectively.

(C) Evolution curves of invagination depth (blue) and center cell height (orange) in simulations. Dots represent corresponding experimentally measured values.

(D) Snapshots at specific timepoints during simulation progression. The colors of edges represent actomyosin intensity.

(E, F) Invagination depth (E) and center cell height (F) regulated by changes in actomyosin intensities under the scaling factor α at both apical and lateral regions of all cells. The data points are from the mutant experiments in Figure 3B.

Figure 5—video 1. Vertex model simulation.

Top row: Progression of the vertex model simulation. Edge colors represent actomyosin intensity.

Bottom row (left to right):

1. Evolution curves of invagination depth;

2. Evolution curves of center cell height;

3. Temporal evolution of actomyosin intensities at the apical, lateral, and basal domains of the center cell in the model, parameterized based on experimentally measured F-actin distribution.

Figure 6. Mechanics of epithelial invagination revealed by vertex model simulations

(A, C) Invagination depth (A) and center cell height (C) under varying apical actomyosin intensity represented by a scaling factor α_a, with the lateral intensity unchanged. A Schematic illustration of the bending mode induced by apicobasal imbalance is presented. A snapshot of a specific inversed evagination modeling under $\alpha_a = 0.1$ is shown.

(B, D) Invagination depth (B) and center cell height (D) under varying lateral actomyosin intensity represented by a scaling factor α_l, with the apical intensity unchanged. A Schematic illustration of the contraction mode induced by lateral tension is presented.

(E) Temporal evolution of the sum S_{exp} and ratio R_{exp} of apical and lateral actomyosin intensities in the center cell in experiments.

(F) Varying sum S_c and ratio R_c of actomyosin intensities regulated by α and β in simulations.

(G, H) Invagination depth (G) and center cell height (H) under varying α and β.

942 **Appendix 1—table 1. Key resource table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
antibodies		
Phospho-Myosin Light Chain 2 (Ser19) Antibody	Cell Signaling	#3671
Alexa Fluor 647 anti-Mouse IgG	Invitrogen	#A28181
Chemicals		
EdU Cell Proliferation Kit with AF555	BeyoClick	C0075S
Alexa Fluor™ 488 Phalloidin	Thermo fisher	A12379
Protease E	sigma-Aldrich	P5147
Sodium thioglycolate	sigma-Aldrich	T0632
D-mannitol	sigma-Aldrich	M4125
MS-222	sigma-Aldrich	E10521
VECTASHIELD® with DAPI	Vectorlabs	H1200
Phanta Max Super-Fidelity DNA Polymerase	Vazyme	P505
ClonExpress MultiS One Step Cloning Kit	Vazyme	C113
TUNEL BrightRed Apoptosis Detection Kit	Vazyme	A113
Plasmids		
Epi1> MRLC::mCherry	This study	N/A
Epi1> MRLC(T18AS19A)::mCherry		N/A
Epi1> MRLC(T18ES19E)::mCherry		N/A
Epi1>PP1c::MYPT169::BcLOV4::mCherry:: NES		N/A
Epi1>PP1c::MYPT169::mCherry::NES		N/A
Software		
ImageJ	NIH Image	https://imagej.nih.gov/ij/
Prism 10 CN	Graphpad Software	https://www.graphpad.com/
Imaris 10	Oxford Instruments	https://imaris.oxinst.com/
MATLAB R2021a	MathWorks	https://www.mathworks.com/

950 **Appendix 1—table 2. Primer sequences used in this study**

Construct name	Primers	Sequence (5'→3')
MRLC	Forward	ATGTCG-AGCCGACGAACTAAA
	Reverse	CTAAATGTCATCTTTTTCTTTAGCG
MRLC(T18AS19A)	Forward	TGCAGCCGCCAATGTCTTTGCTATGTTCGATCAAGC
	Reverse	GACATTGGCGGCTGCACGCTGGGCGCGC
MRLC(T18ES19E)	Forward	TGCAGAAGAAAATGTCTTTGCTATGTTCGATCAAGCT
	Reverse	AGACATTTTCTTCTGCACGCTGGGCGCG
BcLOV4	Forward	ATGGCCACAGACGCAATCG
	Reverse	GATTATGATCTAGAGTC
PP1c	Forward	ATGGCGGACGGGGAGC
	Reverse	CCTTTTCTTCGGCGGA
MYPT169	Forward	ATGAAGATGGCGGACGCG
	Reverse	AGCTGCTTCTATATCAACCCCTTG