SPINK3-sperm interaction determines a stable sperm subpopulation with intact CatSper channel

SPINK3 stabilizes CatSper1 and membrane

Anabella R. Nicolli[1]*, Xiaofang Huang[2]*, Lucia Zalazar[1], Cintia Stival[4], Lucila R. Gomez-Olivieri[5], Gianluca Demare[1], Ana Romarowski[5], Mariano G. Buffone[5], Dario Krapf[4], Jean-Ju Chung[2,3] and Andreina Cesari[1]

*Equal contribution

1- Instituto de Investigaciones Biológicas (IIB-FCEyN/CONICET), Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Mar del Plata, Argentina.
2- Department of Cellular & Molecular Physiology, Yale University School of Medicine, New Haven, CT 06510, USA
3- Department of Ob/Gyn & Reproductive Sciences, Yale University School of Medicine, New Haven, CT 06510, USA
4- Cell Signal Transduction Networks Laboratory, Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET-UNR), Rosario, Argentina
5- Instituto de Biología y Medicina Experimental (IBYME-CONICET), Ciudad Autónoma de Buenos Aires, Argentina.

Corresponding author: Andreina Cesari; Funes 3250 3rd floor, Mar del Plata, Argentina; +5492235134227; andreinacesari@gmail.com

50
51
GRAPHICAL ABSTRACT
53
54
55

56
57
ABSTRACT
59

Sperm capacitation involves proteolytic remodeling of membrane proteins, including components of the CatSper calcium channel, which is essential for hyperactivation and male fertility. Here, we identify the seminal protease inhibitor SPINK3, a known decapacitation factor that suppresses premature capacitation in the female tract, as the first physiological inhibitor of CATSPER1 processing. In mouse sperm, SPINK3 blocks capacitation-induced CATSPER1 cleavage, preserving a subpopulation with intact CatSper channels and lacking

pTyr development in the flagellum. SPINK3 localizes to the outer surface of the sperm
principal piece membrane in a CatSper-dependent but non-quadrilateral pattern, stabilizes
membrane organization, and delays cholesterol efflux. These results reveal SPINK3 as a
multifunctional regulator of capacitation, shaping sperm subpopulations in the female
reproductive tract.

Keywords: sperm capacitation, SPINK3, CATSPER1, protease inhibition, membrane
stabilization, decapacitation factors

SIGNIFICANCE STATEMENT

This study unveils a novel physiological regulator from the male reproductive tract, SPINK3,
which extracellularly controls the processing of CATSPER1, a key subunit of the CatSper
calcium channel essential for sperm capacitation in mice. We show that SPINK3 binds to the
sperm surface and prevents the processing of CATSPER1 during capacitation. SPINK3-
sperm interaction is dependent on the presence of the sperm-specific CatSper channel. Our
findings reveal that SPINK3 stabilizes sperm membranes providing new insights into the
molecular regulation of sperm function and fertility.

INTRODUCTION

The sperm journey begins in the male reproductive tract, where millions of sperm undergo
surface modifications induced by male secretory molecules (A. RNicolli & Cesari, 2023).
After ejaculation, mammalian sperm must undergo biochemical and functional changes
within the female reproductive tract, a process collectively known as capacitation (Austin,
1952; Chang, 1951). This process involves membrane remodeling and the activation of
intracellular signaling pathways leading to the acquisition of fertilizing ability. Decapacitation
factors, typically originating from the male reproductive tract or seminal plasma, coat and
stabilize the sperm surface after ejaculation, preventing or delaying capacitation until the
sperm reaches an appropriate environment in the female reproductive tract (Dukelow et al.,
1967; Leahy & Gadella, 2011; Nicolli & Cesari, 2023).
CatSper is the primary Ca^{2+} channel in mammalian spermatozoa, consisting of four
pore-forming subunits (CATSPER1, 2, 3, 4) and at least ten additional auxiliary subunits
(Hwang & Chung, 2023). This channel is specifically located in the flagellar membrane of the
principal piece, exhibiting pH sensitivity and weak voltage dependence (Kirichok et al., 2006;
Ren et al., 2001) and is essential for the development of hyperactivated motility of sperm (Qi
et al., 2007; Quill et al., 2003). Infertility has been observed in male mice lacking the *Catsper*
genes encoding the transmembrane subunits (Chung et al., 2011; Huang et al., 2023;
Hwang, 2024; Qi et al., 2007; Quill et al., 2003; Ren et al., 2001). Capacitation induces
specific cleavage and processing of CATSPER1, one of the pore-forming subunits, in the
cytosolic N-terminus by unknown sperm enzyme(s), resulting in a graded distribution of
heterogeneous sperm populations with varying degrees of CATSPER1 integrity along the
female reproductive tract (Chung et al., 2014; Ded et al., 2020). Spermatozoa that reach the
ampulla*,* where fertilization takes place, mostly have intact CATSPER1 signal compared to
those that remain behind the utero-tubal junction (UTJ). Recently, it was reported that
another CatSper transmembrane subunit, CATSPERθ, which is essential for the CatSper
channel assembly and serves as a checkpoint for flagellar trafficking during spermiogenesis,
is also processed during sperm capacitation (Huang et al., 2023). This highlights the

proteolytic regulation of the CatSper channels as a sperm selection mechanism. Although
the cell-permeable potent proteasome inhibitor MG-132, as well as calpain inhibitors and the
phosphatase 1/2A inhibitor calyculin A, have been shown to inhibit CATSPER1 processing
(Chung et al., 2014; Ded et al., 2020), physiological inhibitors responsible for modulating the
processing of CatSper subunits remains unidentified.
SPINK3 (also known as P12 or SPINK1) is a small and highly conserved protein
consisting of 80 amino acids, naturally secreted by the male seminal vesicles of most
mammals (Dematteis et al., n.d.; Tiwari et al., 2020; Zalazar et al., 2020). It contains a Kazal
domain that acts as a substrate analogue by competitively and stoichiometrically binding to
the active site of its target protease, forming a stable protease–protease inhibitor complex,
such as kallikreins, trypsin and chymotrypsin-like proteases (Assis et al., 2013). SPINK3 also
has a sperm binding domain (Lin et al., 2006) and attaches to sperm during ejaculation
(Nicolli et al., 2022). SPINK3 is a known decapacitation factor regulating SRC tyrosine
kinase activation and membrane potential, apparently without affecting the tyrosine kinase or
PKA activity (Zalazar et al., 2020). The BSA-induced increase in intracellular Ca^{2+}
concentration ($[Ca^{2+}]_i$) was demonstrated to depend on CatSper channels. This $[Ca^{2+}]_i$
increase and CatSper activity are inhibited by the presence of SPINK3 (Zalazar et al., 2020),
suggesting an either direct or indirect blockade of calcium entry through the CatSper
channel. Similar to other decapacitation factors (Li et al., 2018; Lin et al., 2008; Silva et al.,
2021), SPINK3 has been demonstrated to bind sperm during ejaculation until detachment in
the female tract, following a specific but asynchronous spatiotemporal kinetics, and
contributing to sperm heterogeneity (Nicolli et al., 2022). The protease TESP1 is the only
natural target reported for SPINK3 on the sperm head and this interaction was found to be
blocked by the soybean trypsin inhibitor, SBTI (Ramachandran et al., 2021).
In this study, we show that SPINK3 regulates CATSPER1 processing and contributes
to the generation of sperm subpopulations with different capacitation states. This action is
accompanied by the stabilization of the sperm plasma membrane, suggesting that its
regulatory effect may involve protease inhibition and membrane-stabilizing mechanisms that
prevent protease access to CATSPER1 cleavage sites. These findings highlight a
multifaceted role for SPINK3 in modulating sperm capacitation.

MATERIALS AND METHODS
145
146   *Animals*
147
Female and male mice (*Mus musculus*) of BALB/c and C57BL/6 strains and sexually
mature (BALB/c x C57BL/6) F1 hybrids were maintained at 22°C with a photoperiod of 12 h
light:12 h darkness, food and water *ad libitum*. Sexually mature (2–3 months) male and
female mice were euthanized by cervical dislocation.
Catsper1-null mice were previously generated (Ren et al., 2001) and maintained on a
C57BL/6 background.
Mice were cared for in compliance with the guidelines approved by local Animal Care
and Use Committees (UNMdP RD 104/22), following the National Institutes of Health Guide
to the Care and Use of Laboratory Animals.
157
158   *Antibodies and reagents*
159

Rabbit polyclonal antibodies against CATSPER1 (Ren et al., 2001), CATSPERθ
(Huang et al., 2023) and SPINK3 (Sigma HPA027498) were used. A monoclonal anti-HA
antibody was purchased from Cell Signaling Technology (#3724), and a monoclonal anti-
tubulin antibody (T6094, 1:5000) was obtained from Sigma-Aldrich. HRP-conjugated goat
anti-rabbit IgG and goat anti-mouse IgG were used as secondary antibodies. Recombinant
SPINK3 protein was produced in *E. coli* and purified using HiTrap IMAC HP (GE Healthcare
Life Sciences).
Additional reagents included: Soybean Trypsin Inhibitor (SBTI) (1 µM), Fluo-3 AM
(Invitrogen USA, Molecular Probes, F1242), Pluronic acid (Invitrogen, USA), Laminin (100
µg/ml, BD Biosciences), PNA Alexa Fluor 594 lectin (1 µg/ml, Molecular Probes, L-32459),
Bodipy-Cholesterol (15 µM, Cayman, 24618), M540 (2.7 µM, Sigma-Aldrich, 323756), FM4-
64FX (100 µM, Invitrogen, F34653), CellBrite Fix 640 (2.5 µM, Biotium, 30089-T), Yo-Pro1
(50 nM, Invitrogen, Y3603), Trypsin (347 USP-u/mg, Biological industries, 4192025), PMSF
(Phenylmethylsulfonyl fluoride, 500µM, ICN Biomedicals Inc., 195381) and fluorogenic
protease substrates Z-CGGR-AMC (7-Amino-4-methylcoumarin), Suc-LLVY-AMC, and Z-
SLY-AMC (2 µM each; all from Sigma-Aldrich, Merck, Darmstadt, Germany) .
176
177	***Expression and purification of recombinant SPINK3***
178
179	For the production of SPINK3, the cDNA encoding the mature SPINK3 from *M.*
180	*musculus* (NCBI ID: NM 009258.5) was cloned into the pET-24b(+) (Novagen, Madison, WI,
181	USA) expression vector. Overexpression of SPINK3 was performed in *Escherichia coli*
182	Rosetta cells (Novagen) and the recombinant protein was purified to apparent homogeneity
183	as described (Assis et al., 2013, and Supplementary Information S1).
184
185	***Sperm preparation***
186
Epididymal spermatozoa from adult male mice were collected by swim-out from
caudal epididymis in M2 medium (EMD Millipore, MR-015-D) or HM medium (Modified Krebs
Ringer Bicarbonate medium: 20 mM Hepes, 119.3 mM NaCl, 4.7 mM KCl, 1.2 mM $MgSO_4$,
5.6 mM glucose, 1.2 mM KH_2PO_4, 0.5 mM Na pyruvate and 1.7 mM $CaCl_2$; pH 7.4).
Collected sperm were washed with fresh HM or PBS.
For capacitation, sperm were incubated in HMB (HM plus 15 mM $NaHCO_3$ and 5 mg
mL^{-1} BSA) or human tubular fluid HTF medium (EMD Millipore, MR-070-D) at (2-7 x 10^6 cells
mL^{-1}) w/o purified SPINK3 at 37°C, 5% CO_2 for 1 or 2 hr. SPINK3 concentrations (5-13 µM)
varied according to the assay.
In a set of experiments capacitation was performed in the presence of SPINK3 (13
µM) or SBTI (1 µM) (Noda & Ikawa, 2019; Samanta et al., 2018).
For the preparation of sperm lysates, sperm samples (10 × 10 cells) were
resuspended in 1 mL of 0.1% Triton X-100 in PBS. The suspension was sonicated three
times for 1 second each using a small ultracentrifuge tube. Samples were then agitated for 1
hour at 4 °C, followed by centrifugation at 15,000 × g for 10 minutes at 4 °C. The
supernatant was collected and used immediately. Lysates were always prepared freshly
before use.
204
205	***Seminal vesicle lysates***
206

Seminal vesicle homogenate was prepared from seminal vesicles obtained from
adult male mice. The tissues were placed in ice-cold PBS (pH 7.4) and homogenized using a
IKA T10 basic homogenizer under cold conditions. The homogenate was then centrifuged at
10000 × g for 10 minutes, and the supernatant was filtered (0.22µm) and stored on ice until
use. The homogenate was subsequently used in activity assays, adjusting the volume based
on the estimated concentration of native SPINK3 in seminal vesicle extracts.
213
214   ***Proteolytic activity of sperm lysates and inhibitory assay***
215
Proteolytic activity of sperm lysates was assessed over fluorogenic substrates: Z-
Arg-Gly-Arg-AMC (for trypsin-like proteases), Suc-LLVY-AMC (for chymotrypsin-like
proteases), and Z-SLY-AMC (for elastase-like proteases), each at a final concentration of
2 µM. All reactions were performed in 96-well black plates (Nunc) with a final volume (Fv) of
100 µl per well.
Reaction mixtures were prepared at room temperature with 40 µl of sperm lysate as
protease source. The reactions (Fv=100 µl) contained 50 mM Buffer phosphate pH 7.4, w/o
the following inhibitors: SPINK3 (1.3 µM final), SBTI (2 µg/ml final), seminal vesicle
homogenate as a control of native SPINK3 (6 µl) and PMSF (500 µM). The reaction was
initiated by adding 2 µM CGGR-AMC substrate, and fluorescence (Ex 355 nm / Em
436 nm) was measured every minute for 30 min at 37 °C with agitation (Fluoroskan
Ascent microplate reader, Thermo Electron Corporation, Waltham, MA, USA). Blank reaction
was conducted including 40 µl 0,1% Triton. Enzymatic activity was calculated as
Δfluorescence over time and expressed relative to the sperm lysate fluorescence. Inhibition
is evidenced as a decrease in proteolytic activity.

In vivo proteolysis of CATSPER1

Capacitation was performed w/o 13 µM SPINK3 or 1 µM SBTI for 2hrs in HTF media.
Western blots with whole sperm lysates (1 or 2 x 10^6 cells/lane) were performed and
developed with anti-CATSPER1 (Ren et al., 2001), anti-CATSPERθ (Huang et al., 2023),
anti-HA (CST, #3724) or anti-acetylated Tubulin (Sigma, T7451), followed by the
corresponding HRP-conjugated goat anti-rabbit or anti-mouse IgG.
237
In vitro proteolysis of CATSPER1

HEK293T cells were transiently transfected with construct encoding CATSPER1-HA
with Lipofectamine 2000 (Invitrogen), following the manufacturer's instruction. CATSPER1-
HA was solubilized with 0.1% Triton X-100 PBS (PBST) buffer, containing EDTA-free
protease inhibitor cocktail and pulled-down with anti-HA magnetic beads (Pierce, 88836) for
1 h. at RT. The enriched CATSPER1-HA protein beads were incubated with 30µL of sperm
lysates solubilized from $3.0x10^5$ sperm cells w/o 13 µM SPINK3 at 37ºC for the indicated
times. After incubation, the mixture was denatured by LDS sampling buffer supplemented
with 50 mM DTT at 75ºC for 15 min. Western Blot was performed as described before.

247
248   ***Time-lapse live cell calcium Imaging***
249

250    Mouse epididymal spermatozoa (n=3) (10 x 10^6 cells/ml) were loaded with Fluo-3 AM
251 (10 µM) (Invitrogen USA, Molecular Probes F1242) and 0.02% (v/v) pluronic acid (Invitrogen,
252 USA) as a surfactant at 37°C for 20 minutes in a non-capacitating medium (NC). Non-
253 capacitated sperm were then incubated w/o SPINK3 (13 µM) or SBTI (1 µM) for 15 minutes
254 at 37°C and then, adhered to a fluorescence compatible 96 well plate previously covered
255 with laminin (100 µg/ml, BD Biosciences). HM medium supplemented with 15 mM $NaHCO_3$
256 was added to the chamber and basal fluorescence was monitored under confocal
257 microscope (excitation 488 nm; emission 526 nm, Nikon C1SiR) at 400x magnification. BSA
258 (5 mg/ml) in a minimum volume was applied, and changes in the calcium signal were
259 recorded throughout the process. The viability of the spermatozoa was determined through
260 the motility of the flagellum. The intensity of the fluorescence signal in the head of
261 spermatozoa with motile flagella was quantified using the free software ImageJ 1.43
262 (National Institutes of Health, Bethesda, Maryland, United States). Data was normalized
263 using the equation (F-F0)/F0, where F is the fluorescence intensity at time t and F0 is the
264 mean fluorescence taken during the period before the addition of BSA (initial 10 seconds).
265 Then, the total series of (F-F0)/F0 was plotted against time, as well as the percentage of
266 positive response to the stimulus.
267
268   *Acrosomal reaction (AR) assay*
269
270    Mouse epididymal spermatozoa (n=3) were capacitated (7.5 x 10^6 cells/ml) for 60
271 minutes w/o SPINK3 (13 µM) or SBTI (1 µM). To induce the acrosomal reaction,
272 spermatozoa were incubated with progesterone (50 µM) (Zalazar et al., 2020) or Ca^{2+}
273 ionophore A23187 (10 µM) (Sigma-Aldrich, United States) for 15 min at 37°C after
274 capacitation. The cells were then incubated for 15 minutes at 37°C with the PNA Alexa fluor
275 594 lectin (1 µg/ml, Molecular Probes, L-32459), which allows AR to be detected by binding
276 to glycoproteins of the internal acrosomal membrane. The fluorescence was observed under
277 a microscope at a magnification of 400x. Sperm that lost their acrosomes (red fluorescence)
278 were expressed as a percentage of 100 cells per replicate.
279
280   *Cholesterol efflux assay*
281
282    Mouse epididymal spermatozoa (n=3) (15 x 10^6 cells/ml) were loaded with Bodipy-
283 Cholesterol (15 µM) (Cayman, 24618) for 30 minutes at 37°C and incubated 15 minutes at
284 37°C w/o SPINK3 (13 µM) or SBTI (1 µM). Cells were washed to remove excess reagent
285 and exposed for 60 minutes to either capacitation (CAP) or non-capacitation (NC)
286 conditions. The fluorescence of the labeled Bodipy cholesterol was observed in a Nikon
287 Eclipse T2000 epifluorescence microscope with a magnification of 400x, and the percentage
288 of cells that preserved the labeled cholesterol was quantified.
289
290   *Determination of lipid disorder*
291
292    Mouse epididymal sperm (n=3) (10x10^6 cells/ml) were incubated in the presence or
293 absence of rSPINK3 (13 µM) for 15 minutes at 37°C in HM medium. Subsequently, they
294 were incubated under CAP or NC conditions. After capacitation, 2.7 µM M540 (Ex 555/Em
295 578 nm, Sigma-Aldrich, 323756) was added to the sperm suspension (1x10^6 total cells) and
296 incubated for 15 minutes at room temperature in the dark. Cells were washed and Yo-Pro1
297 was added at a concentration of 50 nM (Ex 491/Em 509 nm., Invitrogen, Y3603). The

298 samples were analyzed by flow cytometry. The results were analyzed using the open access
299 Flowing 2 software. During acquisition, sperm were discriminated from debris. The results
300 are expressed as the percentage of cells with lipid disorder over the total live cells
301 (M540+/Yo-Pro1-).
302
303 ***Immunolabeling of SPINK3 and CATSPER1***
304
Epididymal spermatozoa from WT and *Catpser1*-/- mice (n=3) were incubated in non-
capacitating medium w/o SPINK3 (13 µM) for 15 minutes at 37°C. Each condition was
centrifuged twice at 500 × g for 5 minutes, and re-suspended in the same volume of fresh
PBS medium without SPINK3. Cells were seeded on 8-well glass slides and allowed to air-
dry. Sperm were fixed with 3.7% paraformaldehyde in PBS for 15 minutes at room
temperature, washed with PBS and permeabilized with 0.5% Triton X-100 for 5 minutes.
After washing with PBS, cells were blocked with 5% BSA in PBS for 1 h at room temperature
and followed by overnight incubation with anti-SPINK3 (1:50 in TBS supplemented with 1 %
BSA and 0.1 % tween, Sigma HPA027498) at 4 °C. After washing with PBS, anti-rabbit IgG-
Alexa 488 (1:1000, Abcam, ab15105) was added for 2 hours at room temperature. Finally,
the cells were washed with PBS and mounted to acquire images with the fluorescence
microscope NIKON ECLIPSE E800. Images from both conditions (WT and *Catpser1*-/-) were
equally processed and analyzed with ImageJ (v1.38, NIH). For data analysis, the percentage
of SPINK3-positive sperm under each condition was adjusted by subtracting the count of
false positive sperm observed in the basal immunofluorescence controls (i.e., WT or
Catpser1-/- sperm incubated without SPINK3).
To assess tyrosine-phosphorylated proteins (anti-pY), sperm were incubated w/o
SPINK3 (13 µM), incubated under CAP or NC conditions and then the cells were fixed with
4% formaldehyde, washed, and placed on slides. Once dried, they were fixed with 96%
ethanol, permeabilized with 10 µl of T-PBS (1x PBS, 0.5% Tween 20%) with 0.5% Triton 100
for 20 minutes. After washing, they were blocked with 3% BSA in T-PBS for 1 hour. The
samples were washed and incubated for 1 hour with the primary antibody anti-tyrosine
mouse monoclonal (anti-pY) (1:500, Clone 4G10, Millipore) or anti-SPINK3 (1:50, Sigma
Aldrich, HPA027498) in blocking buffer, followed by the secondary antibodies anti-rabbit
IgG-Alexa488 (1:1000, Invitrogen A11070) and anti-mouse IgG-Alexa455 (1:500, Invitrogen,
A21425). The preparations were mounted with glycerol:PBS (9:1) and observed under a
fluorescence microscope with a magnification of 400x (Ex 480/Em 525 nm).

Structured illumination microscopy

2D structured illumination microscopy (SIM) imaging was performed with Zeiss
LSM710 Elyra P1 using alpha Plan-APO 100X/1.46 oil objective lens. Samples were
prepared as described in the previous section (Sperm labelling with SPINK3 and
immunostaining with CATSPER1 antibody). Raw images were processed and rendered
using Zen 2012 SP2 software (Carl Zeiss).

Plasma membrane labeling
339
After obtaining the sperm sample and co-incubating it with SPINK3 as previously
described, the sample was incubated with 100 µM FM4-64 FX for 3 minutes at room
temperature or with 2.5 µM CellBrite Fix 640 for 15 minutes at 37°C, followed by the SPINK3

immunolabeling protocol as described above. Confocal images were acquired using a Zeiss
LSM-880 microscope equipped with an Alpha Plan-Apochromat 100x/1.46 objective. Images
were also captured in Super-Resolution mode using an Airyscan detector. A total of 10
frames were acquired, and the Mean-Shift Super Resolution (MSSR) algorithm (Torres-
García et al., 2022) was applied to enhance resolution.
In vitro fertilization
349
F1 hybrid females (female Balb/c x male C57BL/6) were superovulated by
intraperitoneal injection with 7.5U of PMSG and 48 hours later with 5U of hCG. Cumulus-
oocyte complexes (COCs) were obtained from the ampulla 15 hours post hCG and placed in
G-IVF plus medium droplets (LifeCell) under mineral oil (M8410, Sigma-Aldrich).
Subsequently, groups of COCs were transferred to a clean medium drop (90 µl) under oil.
Spermatozoa were obtained from C57BL/6 males in HM medium, incubated w/o SPINK3 (13
µM) for 15 minutes at 37°C and capacitated in G-IVF plus medium (LifeCell) (5 x 10^6
cells/ml). COCs were inseminated with 10 µl of the sperm suspension and incubated at
37°C, 5% CO2 for 4 hours. Afterward, COCs were washed, and putative zygotes were
incubated for 24 hours and 5 days (post-IVF) to assess the two-cell and blastocyst stages
under an inverted microscope with a magnification of 400x (Nikon Eclipse Ti-S). Remanent
sperm were prepared for immunostaining. Bar charts show the percentage of two-cell
embryos and blastocysts over the total fertilized oocytes.

Statistical analysis
364
Data were analyzed by GLMM (generalized linear mixed effect model) to determine
statistical significance between treatments and control (Zuur, 2019). The model included the
male variable as a random effect. Normality of residuals was assessed by plotting theoretical
quantiles vs standardized residuals (Q–Q plots). Homogeneity of variance was evaluated by
plotting residuals vs fitted values. A post hoc analysis was conducted with the 'lsmeans''
package (Lenth, 2016). In all cases, the analysis was performed using R software version
3.3.33, with the 'nlme' package for Gaussian models (Pinheiro et al., 2020). Statistically
significant differences were determined at $P < 0.05$. Graph bars indicate mean ± s.e.
373
374 **RESULTS**

375 **SPINK3 inhibits CATSPER1 processing induced by *in vitro* capacitation**
376
It has previously been reported that the cytoplasmic N-terminal domain of the
CATSPER1 undergoes proteolytic processing in sperm cells during capacitation, and the
progression of this processing is negatively correlated with the appearance of tyrosine
phosphorylation at single cell level *in vitro* and *in situ* (Ded et al., 2020). More recently,
another CatSper subunit, CATSPERθ, which plays a role in CatSper channel assembly and
serves as a checkpoint for CatSper flagellar trafficking, was also found to undergo
capacitation-induced degradation (Huang et al., 2023).
SPINK3 has previously been shown to inhibit CatSper channel activity (Zalazar et al.,
2020). As this small protein is not only a decapacitation factor but also a protease inhibitor,
we hypothesize that SPINK3 might be related to the regulation of CATSPER1 and/or
CATSPERθ processing. To test this idea, we evaluated capacitation-induced CATSPER1

and CATSPERθ protein degradation in mouse sperm incubated under capacitating
conditions, with or without SPINK3. Interestingly, we found that the presence of SPINK3
abolished the degradation of CATSPER1, but not CATSPERθ (**Figure 1A**). Consistent
results were seen with anti-CATSPER1 immunostaining in mouse spermatozoa (**Figure 1B,**
Supp. Figure 1). As sperm contains several proteases (Cesari et al., 2010), we assessed
the proteolytic activity of sperm lysates using synthetic substrates specific for trypsin,
chymotrypsin, and elastase, in order to identify the protease family targeted by SPINK3. Our
results confirmed that trypsin-like activity was predominant, and that this activity was
inhibited by SPINK3, by seminal vesicle lysates (which contain native SPINK3), and by
commercial inhibitors such as PMSF (phenylmethylsulfonyl fluoride) and SBTI (soybean
trypsin inhibitor) (**Figure 1C**). As a control, we capacitated mouse spermatozoa in the
presence of SBTI and observed the same blocking effect on CATSPER1 processing (**Supp.**
Figure 2). Heterologously expressed full-length recombinant CATSPER1 has also been
validated for processing by sperm lysate *in vitro* due to the presence of specific endogenous
protease(s) (Ded et al., 2020). Similarly, this *in vitro* CATSPER1 processing was also
inhibited by the presence of SPINK3 (**Figure 1D**). Taken together, these results show that
SPINK3 interferes with proteolytic processing of CATSPER1, by inhibiting sperm trypsin-like
protease(s). Furthermore, when sperm were preincubated with recombinant SPINK3 under
capacitating conditions, only spermatozoa lacking SPINK3 showed protein tyrosine
phosphorylation (pTyr) development in the flagella (**Figure 1E**), consistent with the
previously reported inverse correlation of pTyr and CatSper channel integrity (Chung et al.,
2014; Ded et al., 2020; Hwang & Chung, 2023). These results support the idea that SPINK3
binding results in a population of sperm with intact CATSPER1 and SPINK3 bound to the
flagellar surface.
412
413   **SPINK3 binds to the sperm flagellar membrane in a CatSper-dependent manner**
414
The CatSper channel forms four linear nanodomains along the flagellar membrane in
which the lipid raft marker caveolin1 resides (Chung et al., 2014). Recombinant SPINK3 can
bind to raft membrane domains in both acrosomal and tail membranes of non-capacitated
mature epididymal spermatozoa and this union is avoided by choleratoxin B supporting
binding to GM1 (A. Nicolli et al., n.d.; Zalazar et al., 2020). Given that SPINK3 is a secretory
protein, we tested whether SPINK3 and CatSper co-localize in the quadrilinear
nanodomains. Confocal microscopy shows that, unlike CATSPER1, which is localized
exclusively to the principal piece, SPINK3 is present in both principal piece and end piece of
the flagellum (**Figure 2A, upper panel**). The distinct linear CatSper nanodomain
arrangement visualized by Structured Illumination Microscopy (SIM) was not seen for
SPINK3 in the flagellar membrane (**Figure 2B**). These observations, combined with our
finding that SPINK3 inhibits the processing of the N-terminal cytoplasmic domain of
CATSPER1, prompted us to further investigate the spatial distribution of SPINK3 in relation
to the plasma membrane. Thus, SPINK3 immunostaining was performed together with
membrane labelling using the specific plasma membrane probes FM4-64 and CellBrite.
Images from Airyscan microscopy and MSSR indicate that SPINK3 is externally associated
with the plasma membrane (**Figure 2C, Supp. Movie 1**). Accordingly, SPINK3 signal spans
a wider area than the measured membrane-to-membrane distances (**Figure 2C, MSSR**
Images). Nevertheless, we observed that the SPINK3 association with sperm is
compromised in *Catsper1* KO sperm and there is also a pattern shift (**Figure 2A (lower**
panel), D; Supp. Figure 3). It is possible that in *Catsper1* KO sperm the loss of CatSper
complex from flagellum changed the membrane property and/or protease localization, thus

compromising the attachment of SPINK3. These results suggest that SPINK3 interacts with
the flagellar membrane in a CatSper-dependent manner but does not directly bind to
CatSper.

440
441 **SPINK3, but not SBTI, specifically affects sperm capacitation events**

442
443    Given that protection of CATSPER1 from proteolytic processing (*i.e.*, intact
CATSPER-mediated Ca^{2+} signaling) is required to control the timing of capacitation, we
analyzed whether the protease inhibitory activity of SPINK3 accounts for its function as a
decapacitation factor. To this end, we first tested whether the trypsin inhibitor SBTI could
mimic the known effects of SPINK3 on capacitation since SBTI can also inhibit CATSPER1
processing (**Supp. Figure 2**). In contrast to SPINK3, there was no difference in the BSA-
induced increase in $[Ca^{2+}]_i$ between SBTI-treated and untreated spermatozoa as assessed
by single-cell Ca^{2+} imaging with Fluo-3 AM (**Figure 3A**), nor in the percentage of acrosome-
reacted spermatozoa (**Figure 3B**). Notably, neither SPINK3 nor SBTI affected intrasperm
alkalinization (**Figure 3C**), which is known to increase CatSper currents (Ferreira et al.,
2021; Hwang et al., 2019; Kirichok et al., 2006). All these results suggest that among trypsin-
like protease inhibitors, SPINK3 specifically inhibits capacitation events such as rise in
$[Ca^{2+}]_i$ and AR likely through a mechanism distinct from its inhibition of CATSPER1
cleavage.

457
458 **SPINK3 alleviates capacitation-induced membrane perturbations**

459
460    Cholesterol efflux from the sperm plasma membrane is a crucial step in capacitation.
These changes promote an increase in membrane fluidity or decrease stability (Flesch et al.,
2001) and can be measured using BODIPY-cholesterol (BODIPY/Ch), a fluorescent
compound that can replace endogenous cholesterol in the membrane without disturbing the
lipid acyl chain region. Therefore, its loss can be measured by changes in fluorescence
intensity (Bernecic et al., 2019; Sankaranarayanan et al., 2011). By contrast, the subsequent
membrane remodeling can be assessed by the incorporation of anionic lipophilic fluorescent
dye Merocyanine 540 (M540) as the amount of dye per unit surface area increases in less-
ordered membranes (McEvoy et al., 1988; Steckler et al., 2015). Using these lipid indicators,
we show that capacitated spermatozoa contain lower cholesterol content (**Figure 4A**) and
increased lipid disorder (**Figure 4B**) compared to non-capacitated sperm, as previously
reported. In contrast, pre-incubation with SPINK3 blocked the capacitation-associated
cholesterol efflux and lipid disorder (**Figure 4A**, **B**), suggesting that SPINK3 binding to the
membrane hinders the membrane reordering during capacitation.

474
475 **SPINK3-exposed spermatozoa during capacitation reduce *in vitro* fertilization**

476
477    Pharmacological inhibition of CatSper during *in vitro* capacitation compromises
sperm fertilizing ability *in vitro* (Curci et al., 2021). *In vivo*, decapacitation factors are
gradually removed from the sperm surface within the female reproductive tract, thereby
allowing capacitation to proceed. In contrast, conventional *in vitro* fertilization (IVF) assays
are performed with epididymal sperm, which naturally lack seminal plasma proteins such as
SPINK3, and membrane-bound proteins might not be efficiently detached under these
conditions. To specifically evaluate whether exogenously added SPINK3 exerts an inhibitory

effect in vitro, epididymal sperm were incubated under capacitating conditions in the
absence or presence of 13 µM recombinant SPINK3 and subsequently used for
inseminating cumulus–oocyte complexes (COCs). We found that SPINK3 produced a
significant decrease in the percentage of fertilized eggs (**Figure 5A, B**). Most sperm that
were preincubated with SPINK3 retained SPINK3 immunostaining in the flagellum when
recovered 4 hours after insemination (**Figure 5C**).

490
491 **DISCUSSION**

492
493 **SPINK3 is involved in the regulation of CATSPER1 processing protease in mice**

494
We have previously reported that the small vesicular protein SPINK3, found in mouse
male secretions, acts as a decapacitation factor by inhibiting CatSper activity and the
increase of $[Ca^{2+}]_i$ during the early stages of capacitation induced by BSA and HCO_3^-
(Zalazar et al., 2020). We have also shown that SPINK3 functions as a Kazal-type serine
protease inhibitor that competitively inhibits trypsin-like proteases (Assis et al., 2013).
Concurrently, it has been shown that both CATSPER1 and CATSPERθ undergo
capacitation-induced degradation (Chung et al., 2014; Ded et al., 2020; Huang et al., 2023).
However, endogenous protein(s) responsible for this physiological regulation remains
unidentified. In this study, we show that SPINK3 inhibits sperm trypsin-like proteases and
specifically inhibits the capacitation-associated cleavage of CATSPER1, but not CATSPERθ,
suggesting a sophisticated regulatory role for SPINK3 in controlling CatSper channel activity.

It has been suggested that inhibitors of calpain, a family of calcium-dependent
cysteine proteases, also block CATSPER1 processing (Ded et al., 2020). However, SPINK3,
as a serine protease inhibitor, is unlikely to directly inhibit calpains. Given that SPINK3
specifically inhibits serine proteases and interferes with CATSPER1 processing, if this effect
results from a direct action of SPINK3 on a sperm protease, then the protease responsible
for CATSPER1 cleavage is likely a serine protease. Alternatively, CATSPER1 N-terminal
domain cleavage could be a multi-protease event involving different types of proteases.
Other sperm serine proteases have been reported in mouse sperm, such as ACROSIN
(Q3ZB06_MOUSE), PRS55 (Q14BX2_MOUSE), and TESP1 (Q3U132_MOUSE), but all are
localized in the sperm head, either as part of the membrane fraction (GPI-anchored or
transmembrane included) or soluble in the acrosome. However, no protease has been
described specifically in the sperm flagella, where CATSPER1 processing is thought to
occur. This requires further investigation.

SPINK3 modulates capacitation through mechanisms that extends beyond its
protease inhibitory activity

521
While the finding that SPINK3 inhibits CATSPER1 processing does not fully explain
the mechanism by which this protein regulates capacitation, we used soybean trypsin
inhibitor (SBTI) as an alternative trypsin-like protease inhibitor to further investigate this
process. SBTI had previously been shown to block the SPINK3 binding site on the sperm
head in mouse sperm (Ramachandran et al., 2021), suggesting a potential shared target
between these molecules in this region. Additionally, SBTI has been used to detect acrosin
exposure in other mammalian sperm (Sumigama et al., 2015), indicating its interaction with

acrosomal proteases. Given these findings, we selected SBTI to determine whether
inhibiting serine proteases alone is sufficient to regulate capacitation. Our results show that,
despite SBTI's ability to inhibit sperm proteases involved in CATSPER1 degradation, it does
not modulate capacitation as SPINK3 does, suggesting that SPINK3's role in capacitation
extends beyond its inhibitory function on serine proteases.

Our *in vitro* results demonstrate that both SPINK3 and seminal vesicles extracts
(containing native SPINK3) strongly inhibit sperm trypsin-like proteases, however evidence
shows that SPINK3 acts extracellularly. Therefore, it is possible to consider a role for surface
attached SPINK3 in stabilizing the membrane and CatSper arrangements in the principal
piece, by binding a membrane protease, whose substrate is CATSPER1, and likely to be
differently compartmentalized outside of CatSper nanodomains before capacitation.
Although there is no strict co-localization of SPINK3 and CATSPER1, sperm from *Catsper1*
KO mice, which lack the entire channel complex in the spermatozoa, displayed a different
SPINK3-sperm binding ability and pattern, suggesting that SPINK3 binds to the sperm
flagellum in a CatSper dependent manner. The different and compromised SPINK3 binding
with sperm may be because lipid domain arrangement was disrupted due to the lack of
CatSper channel in the sperm flagella. Given that TESP1 has previously been reported as
the ligand for SPINK3 in the acrosomal region (Ramachandran et al., 2021), it is possible
that the mechanisms governing SPINK3 binding to sperm vary across membrane
subregions. In sperm lacking the CatSper channel—and consequently exhibiting defective
molecular organization along the flagellum—SPINK3 attachment may be restricted to the
head, as the principal piece may no longer provide a suitable binding site. Additionally,
SPINK3 might play distinct roles in different sperm regions: one regulating calcium influx to
modulate hyperactivation and the other controlling the acrosome reaction. However, the
latter is beyond the scope of this study and requires further investigation.

SPINK3 might be involved in the development of sperm subpopulations in the
oviduct

SPINK3 is bound to the majority of sperm during ejaculation, and it is progressively
detached along the oviduct; however, this detachment is asynchronous, as it does not occur
simultaneously in all sperm (Nicolli et al., 2022). Besides, it is not clear what modulates
CatSper activity and how CatSper channel containing N-terminally processed CATSPER1
(Ded et al., 2020) alters its activity. As most sperm found in the ampulla has a molecular
barcode of acrosome reacted (AR +), tyrosine phosphorylation negative (pY -), and
CATSPER1 intact (CATSPER1 +), and our results show that pY- sperm are also SPINK3+,
one can speculate that late release of SPINK3 from the flagellar membrane contributes to
avoid premature cleavage of CATSPER1 and thus premature degeneration. Proteolysis of
membrane protein ectodomains acts as a post-translational modification that determines the
levels and activity of membrane proteins (Lichtenthaler et al., 2018). Would CATSPER1-
cleaved sperm show increased or reduced CatSper activity? Would intact CATSPER1 be
maintained in the fertilizing sperm or just until right before? In this work, our evidence shows
that SPINK3 prevents this proteolysis that occurs in capacitating media and also prevents
membrane reorganization, increase in $[Ca^{2+}]_i$, and acrosome exocytosis, but it is unclear
whether early or late release constitutes the destiny of sperm for fertilization. Recently, it has
been reported that seminal plasma spermine reversibly inhibits CatSper's temperature

gating, protecting against premature activation. May heterogeneity reflects a compilation of
the time- and space-dependent changes that sperm undergo along the oviduct? Could the
importance of tightly regulating CatSper function be such that multiple redundant molecules
operate in parallel to ensure fertilization success? All these questions serve ongoing and
future research directions.

While SPINK3 is gradually removed from sperm surface *in vivo* in female's
reproductive tract (Nicolli et al., 2022), the detachment is less efficient *in vitro*, as evidenced
by the intense SPINK3 labeling in the sperm recovered from the co-culture and fertility
significantly reduced, suggesting that the detachment of SPINK3 from sperm needs specific
physiological conditions or specific molecules in female's reproductive tract.

In conclusion, we demonstrate a role for SPINK3 in stabilizing the membrane and
preventing CATSPER1 processing, either indirectly or directly acting as an inhibitor of the
responsible protease that cleaves CATSPER1. Our findings provide molecular insights into
the dynamic regulation of capacitation by male-derived SPINK3 and provide the first
evidence of a natural molecule involved in modulating the CatSper channel. This work helps
to explain the mechanism for the graded subpopulations of sperm along the female
reproductive tract.

Acknowledgements
The authors would like to thank Viviana Daniel for their assistance during confocal
microscopy. The authors also acknowledge Hugo Nuñez for his assistance with the animal
handling and maintenance.
594

595 **Declaration of interest**
596 The authors declare that there is no conflict of interest that could be perceived as prejudicing
597 the impartiality of the research reported.
598
599 **Data availability statement**
600 All datasets used for this study are available upon request.
601

Funding
This work was supported by the National Scientific and Technical Research Council
(CONICET, Argentina, PIP 11220200100850CO) awarded to A.C., the National Institutes of
Health (R01HD096745) to J.-J.C, the Chan Zuckerberg Initiative (2021-240504), the
National Institutes of Health (R01HD106968), and the Agencia Nacional de Promoción de la
Investigación, el Desarrollo Tecnológico y la Innovación (Argentina, 2020-00988) to M.G.B,
and grants PICT 2019-1779 and PICT 2021-0102 to D.K.. X.H. is a postdoctoral fellowship
awardee of Male Contraceptive Initiative (MCI).
610

Author contribution statement
AC and JJ contributed to the conceptualization of the study and supervision. AN, XH, CS,
GD, AR and LG contributed to the design and development of the methodology. AN, XH,
DK, MB, LZ, AC and JJ contributed to the writing and editing of the manuscript. All the co-
authors were involved in the formal analysis, validation, visualization of the results and
revised the manuscript.

REFERENCES

Assis, D. M., Zalazar, L., Magno, D., 1#, A., Juliano, M. A., De Castro, R., & Cesari, A. (2013). Novel inhibitory activity for serine protease inhibitor Kazal type-3 (Spink3) on human recombinant kallikreins. *Ingentaconnect.Com*, *20*, 1098–1107. https://doi.org/10.2174/0929866511320100003

AUSTIN, C. R. (1952). The capacitation of the mammalian sperm. *Nature*, *170*(4321). https://doi.org/10.1038/170326a0

Bernecic, N. C., Zhang, M., Gadella, B. M., Brouwers, J. F. H. M., Jansen, J. W. A., Arkesteijn, G. J. A., de Graaf, S. P., & Leahy, T. (2019). BODIPY-cholesterol can be reliably used to monitor cholesterol efflux from capacitating mammalian spermatozoa. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-45831-7

Cesari, A., Monclus, M. de los A., Tejón, G. P., Clementi, M., & Fornes, M. W. (2010). Regulated serine proteinase lytic system on mammalian sperm surface: There must be a role. *Theriogenology*, *74*(5), 699-711.e5. https://doi.org/10.1016/J.THERIOGENOLOGY.2010.03.029

Chung, J. J., Navarro, B., Krapivinsky, G., Krapivinsky, L., & Clapham, D. E. (2011). A novel gene required for male fertility and functional CATSPER channel formation in spermatozoa. *Nature Communications*, *2*(1). https://doi.org/10.1038/ncomms1153

Chung, J. J., Shim, S. H., Everley, R. A., Gygi, S. P., Zhuang, X., & Clapham, D. E. (2014). Structurally distinct Ca2+ signaling domains of sperm flagella orchestrate tyrosine phosphorylation and motility. *Cell*, *157*(4). https://doi.org/10.1016/j.cell.2014.02.056

Curci, L., Carvajal, G., Sulzyk, V., Gonzalez, S. N., & Cuasnicú, P. S. (2021). Pharmacological Inactivation of CatSper Blocks Sperm Fertilizing Ability Independently of the Capacitation Status of the Cells: Implications for Non-hormonal Contraception. *Frontiers in Cell and Developmental Biology*, *9*. https://doi.org/10.3389/fcell.2021.686461

Ded, L., Hwang, J. Y., Miki, K., Shi, H. F., & Chung, J. J. (2020). 3D in situ imaging of female reproductive tract reveals molecular signatures of fertilizing spermatozoa in mice. *ELife*, *9*. https://doi.org/10.7554/eLife.62043

Dematteis, A., Miranda, S., … M. N.-B. of, & 2008, undefined. (n.d.). Rat caltrin protein modulates the acrosomal exocytosis during sperm capacitation. *Academic.Oup.Com*. Retrieved April 2, 2023, from https://academic.oup.com/biolreprod/article-abstract/79/3/493/2557586

Dukelow, W. R., Chernoff, H. N., & Williams, W. L. (1967). Properties of decapacitation factor and presence in various species. *Journal of Reproduction and Fertility*, *14*(3). https://doi.org/10.1530/jrf.0.0140393

Ferreira, J. J., Lybaert, P., Puga-Molina, L. C., & Santi, C. M. (2021). Conserved Mechanism of Bicarbonate-Induced Sensitization of CatSper Channels in Human and Mouse Sperm. *Frontiers in Cell and Developmental Biology*, *9*. https://doi.org/10.3389/fcell.2021.733653

Flesch, F. M., Brouwers, J. F. H. M., Nievelstein, P. F. E. M., Verkleij, A. J., Van Golde, L. M. G., Colenbrander, B., & Gadella, B. M. (2001). Bicarbonate stimulated phospholipid scrambling induces cholesterol redistribution and enables cholestrol depletion in the sperm plasma membrane. *Journal of Cell Science*, *114*(19). https://doi.org/10.1242/jcs.114.19.3543

Huang, X., Miyata, H., Wang, H., Mori, G., Iida-Norita, R., Ikawa, M., Percudani, R., &
Chung, J. J. (2023). A CUG-initiated CATSPERθ functions in the CatSper channel
assembly and serves as a checkpoint for flagellar trafficking. *Proceedings of the*
National Academy of Sciences of the United States of America, *120*(39).
https://doi.org/10.1073/pnas.2304409120
Hwang, J. Y. (2024). Analysis of Ca2+-mediated sperm motility to evaluate the functional
normality of the sperm-specific Ca2+ channel, CatSper. *Frontiers in Cell and*
Developmental Biology, *12*. https://doi.org/10.3389/fcell.2024.1284988
Hwang, J. Y., & Chung, J. J. (2023). CatSper Calcium Channels: 20 Years On. In *Physiology*
(Vol. 38, Issue 3). https://doi.org/10.1152/PHYSIOL.00028.2022
Hwang, J. Y., Mannowetz, N., Zhang, Y., Everley, R. A., Gygi, S. P., Bewersdorf, J., Lishko,
P. V., & Chung, J. J. (2019). Dual Sensing of Physiologic pH and Calcium by EFCAB9
Regulates Sperm Motility. *Cell*, *177*(6). https://doi.org/10.1016/j.cell.2019.03.047
Kirichok, Y., Navarro, B., & Clapham, D. E. (2006). Whole-cell patch-clamp measurements
of spermatozoa reveal an alkaline-activated Ca2+ channel. *Nature*, *439*(7077).
https://doi.org/10.1038/nature04417
Leahy, T., & Gadella, B. M. (2011). Sperm surface changes and physiological consequences
induced by sperm handling and storage. In *Reproduction* (Vol. 142, Issue 6).
https://doi.org/10.1530/REP-11-0310
Lenth, R. V. (2016). Least-squares means: The R package lsmeans. *Journal of Statistical*
Software, *69*. https://doi.org/10.18637/jss.v069.i01
Li, J., Roca, J., Pérez-Patiño, C., … I. B.-A. reproduction, & 2018, undefined. (n.d.). Is boar
sperm freezability more intrinsically linked to spermatozoa than to the surrounding
seminal plasma? *Elsevier*. Retrieved April 2, 2023, from
https://www.sciencedirect.com/science/article/pii/S0378432017309363
Lichtenthaler, S. F., Lemberg, M. K., & Fluhrer, R. (2018). Proteolytic ectodomain shedding
of membrane proteins in mammals—hardware, concepts, and recent developments.
The EMBO Journal, *37*(15). https://doi.org/10.15252/embj.201899456
Lin, H., Luo, C., Wang, C., Biophysical, Y. C.-B. and, & 2006, undefined. (n.d.). Epitope
topology and removal of mouse acrosomal plasma membrane by P12-targeted
immunoaggregation. *Elsevier*. Retrieved April 2, 2023, from
https://www.sciencedirect.com/science/article/pii/S0006291X06018377
Lin, M., Lee, R., Hwu, Y., Lu, C., … S. C.-, & 2008, undefined. (n.d.). SPINKL, a Kazal-type
serine protease inhibitor-like protein purified from mouse seminal vesicle fluid, is able to
inhibit sperm capacitation. *Rep.Bioscientifica.Com*. Retrieved April 2, 2023, from
https://rep.bioscientifica.com/view/journals/rep/136/5/559.xml
McEvoy, L., Schlegel, R. A., Williamson, P., & Del Buono, B. J. (1988). Merocyanine 540 as
a flow cytometric probe of membrane lipid organization in leukocytes. *Journal of*
Leukocyte Biology, *44*(5). https://doi.org/10.1002/jlb.44.5.337
Nature, M. C.-, & 1951, undefined. (n.d.). Fertilizing capacity of spermatozoa deposited into
the fallopian tubes. *Springer*. Retrieved April 2, 2023, from
https://link.springer.com/content/pdf/10.1038/168697b0.pdf
Nicolli, A., Alonso, C., Otamendi, C., … M. C.-, & 2022, undefined. (n.d.). How, where and
when is SPINK3 bound and removed from mouse sperm? *Rep.Bioscientifica.Com*.
Retrieved April 2, 2023, from https://rep.bioscientifica.com/view/journals/rep/163/5/REP-
21-0195.xml

Nicolli, A. R., & Cesari, A. (2023). The other side of capacitation: role of mouse male
molecules in the regulation of time and place of capacitation. *Reproduction*, *166*(6).
https://doi.org/10.1530/REP-23-0188
Noda, T., & Ikawa, M. (2019). Physiological function of seminal vesicle secretions on male
fecundity. In *Reproductive Medicine and Biology* (Vol. 18, Issue 3).
https://doi.org/10.1002/rmb2.12282
Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., & Van Willigen, B. (2020).
Package nlme: Linear and Nonlinear Mixed Effects Models. *R Package*.
Qi, H., Moran, M. M., Navarro, B., Chong, J. A., Krapivinsky, G., Krapivinsky, L., Kirichok, Y.,
Ramsey, I. S., Quill, T. A., & Clapham, D. E. (2007). All four CatSper ion channel
proteins are required for male fertility and sperm cell hyperactivated motility.
Proceedings of the National Academy of Sciences of the United States of America,
104(4). https://doi.org/10.1073/pnas.0610286104
Quill, T. A., Sugden, S. A., Rossi, K. L., Doolittle, L. K., Hammer, R. E., & Garbers, D. L.
(2003). Hyperactivated sperm motility driven by CatSper2 is required for fertilization.
Proceedings of the National Academy of Sciences of the United States of America,
100(25). https://doi.org/10.1073/pnas.2136654100
Ramachandran, S., … R. B.-M. H., & 2021, undefined. (n.d.). A mouse testis serine
protease, TESP1, as the potential SPINK3 receptor protein on mouse sperm acrosome.
Academic.Oup.Com. Retrieved April 2, 2023, from
https://academic.oup.com/molehr/article-abstract/27/10/gaab059/6370709
Ren, D., Navarro, B., Perez, G., Jackson, A. C., Hsu, S., Shi, Q., Tilly, J. L., & Clapham, D.
E. (2001). A sperm ion channel required for sperm motility and male fertility. *Nature*,
413(6856). https://doi.org/10.1038/35098027
Samanta, L., Parida, R., Dias, T. R., & Agarwal, A. (2018). The enigmatic seminal plasma: A
proteomics insight from ejaculation to fertilization. In *Reproductive Biology and
Endocrinology* (Vol. 16, Issue 1). https://doi.org/10.1186/s12958-018-0358-6
Sankaranarayanan, S., Kellner-Weibel, G., De La Llera-Moya, M., Phillips, M. C., Asztalos,
B. F., Bittman, R., & Rothblat, G. H. (2011). A sensitive assay for ABCA1-mediated
cholesterol efflux using BODIPY-cholesterol. *Journal of Lipid Research*, *52*(12).
https://doi.org/10.1194/jlr.D018051
Silva, A. A. S., Raimundo, T. R. F., Mariani, N. A. P., Kushima, H., Avellar, M. C. W.,
Buffone, M. G., Paula-Lopes, F. F., Moura, M. T., & Silva, E. J. R. (2021). Dissecting
EPPIN protease inhibitor domains in sperm motility and fertilizing ability: Repercussions
for male contraceptive development. *Molecular Human Reproduction*, *27*(12).
https://doi.org/10.1093/molehr/gaab066
Steckler, D., Stout, T. A. E., Durandt, C., & Nöthling, J. O. (2015). Validation of merocyanine
540 staining as a technique for assessing capacitation-related membrane
destabilization of fresh dog sperm. *Theriogenology*, *83*(9).
https://doi.org/10.1016/j.theriogenology.2015.01.019
Sumigama, S., Mansell, S., Miller, M., Lishko, P. V., Cherr, G. N., Meyers, S. A., & Tollner,
T. (2015). Progesterone accelerates the completion of sperm capacitation and activates
catsper channel in spermatozoa from the rhesus macaque. *Biology of Reproduction*,
93(6). https://doi.org/10.1095/biolreprod.115.129783
Tiwari, R., Manzar, N., Bhatia, V., Yadav, A., Nengroo, M. A., Datta, D., Carskadon, S.,
Gupta, N., Sigouros, M., Khani, F., Poutanen, M., Zoubeidi, A., Beltran, H., Palanisamy,
N., & Ateeq, B. (2020). Androgen deprivation upregulates SPINK1 expression and

potentiates cellular plasticity in prostate cancer. *Nature Communications*, *11*(1).

https://doi.org/10.1038/s41467-019-14184-0

Torres-García, E., Pinto-Cámara, R., Linares, A., Martínez, D., Abonza, V., Brito-Alarcón, E.,

Calcines-Cruz, C., Valdés-Galindo, G., Torres, D., Jabloñski, M., Torres-Martínez, H.

H., Martínez, J. L., Hernández, H. O., Ocelotl-Oviedo, J. P., Garcés, Y., Barchi, M.,

D'Antuono, R., Bošković, A., Dubrovsky, J. G., … Guerrero, A. (2022). Extending

resolution within a single imaging frame. *Nature Communications*, *13*(1).

https://doi.org/10.1038/s41467-022-34693-9

Zalazar, L., Stival, C., Nicolli, A. R., De Blas, G. A., Krapf, D., & Cesari, A. (2020). Male

Decapacitation Factor SPINK3 Blocks Membrane Hyperpolarization and Calcium Entry

in Mouse Sperm. *Frontiers in Cell and Developmental Biology*, *8*.

https://doi.org/10.3389/FCELL.2020.575126/FULL

Zuur. (2019). Mixed Effects Models and Extensions in Ecology with R by Zuur. In *springer*

(Vol. 53, Issue 9).

FIGURES AND LEGENDS

Figure 1. Capacitation-induced CATSPER1 processing by sperm proteases is
inhibited by SPINK3 *in vitro.* **A)** Mouse epididymal spermatozoa were incubated w/o
SPINK3 (13 µM) and exposed to capacitating conditions for either 1 or 2 hr., as indicated.
Immunoblot analysis of non-capacitated (NC) or capacitated (CAP) sperm lysates was
performed with anti-CATSPER1 antibody raised against the 1–150 aa region of CATSPER1
(7) and anti-CATSPERθ antibody, raised against the C-terminus of CATSPERθ (12).
Acetylated tubulin (AC-TUB) was probed as loading control (n=3). **B)** Immunostaining with
anti-CATSPER1 antibody was performed on non-capacitated sperm or capacitated sperm
that were previously incubated w/o SPINK3 for 30 min. Sperm with intact CATSPER1 signal
were quantified (n=3). **p < 0.001. **C)** Proteolytic activity of sperm lysates was measured
over three fluorogenic substrates—Z-Arg-Gly-Arg-AMC (CGGR; trypsin substrate), Suc-
LLVY-AMC (LLVY; chymotrypsin substrate), and Z-SLY-AMC (SLY; elastase substrate) in
the presence or absence of SPINK3 (13 µM), seminal vesicle homogenate (native SPINK3
source), SBTI (soybean trypsin inhibitor) (2 µg/ml), or PMSF (500µM) (serine protease
inhibitor) (n=3). Enzymatic activity was calculated as Δfluorescence over time and expressed
relative to the sperm lysate. **D)** FL-CATSPER1-HA was expressed in 293T cells and pulled-
down using magnetic beads conjugated with anti-HA antibody. The recombinant protein was
then incubated with sperm lysates w/o SPINK3 (13 µM) at 37 C for 0, 10, 30, or 60 min and
subjected to immunoblot with anti-HA antibody. **E)** Tyrosine phosphorylated proteins and
sperm-bound SPINK3 were detected by immunofluorescence, with anti-pY (clone 4G10,
green) and anti-SPINK3 (HPA027498, red) antibodies, respectively. Arrows indicate
spermatozoa exclusively stained with anti-pY or anti-SPINK3. Scale bar: 10 µm.
849

Figure 2. Immunodetection of recombinant SPINK3 binding to epididymal spermatozoa in wild type (WT) and *Catsper1* KO (*Catsper1-/-*) mice. A) Non-capacitated (NC) epididymal spermatozoa from WT and *Catsper1-/-* mice (n=3) were preincubated with SPINK3 (5 µM), and immunostaining was performed to detect SPINK3 (red) and CATSPER1 (green). Nuclei were counterstained with Hoechst (blue). Representative images from three independent experiments are shown. Scale bar: 10 µm. **B)** 2D Structured illumination microscopy (SIM) images reveal the plasma membrane localization of SPINK3 (red) on the WT NC sperm cells. The immunostaining of CATSPER1 (green) was used to indicate the

boundary of the plasma membrane. Three of the four lines of CatSper nanodomains (as
indicated by CATSPER1, green) were seen in this confocal microscopy image. **C)** (Left
panel) Airyscan- and MSSR-processed images show SPINK3 (red) localized on the plasma
membrane of the sperm principal piece, labeled with FM4-64 FX (green). At the level of the
yellow line in the FM4-64 FX MSSR image, the distance measured for FM4-64 FX is 0.409
µm, whereas for SPINK3, it is 0.736 µm. Orthogonal projections from a Z-stack are shown.
Scale bar: 1 µm. (Right panel) Airyscan- and MSSR-processed images show SPINK3 (red)
on the plasma membrane of the sperm principal piece, labeled with CellBrite Fix (cyan). The
midpiece plasma membrane exhibits stronger staining than the principal piece. At the level
of the yellow line in the CellBrite FixMSSR image, the distance measured for CellBrite is
0.442 µm, whereas for SPINK3, it is 0.752 µm. Orthogonal projections from a Z-stack are
shown. Scale bar: 1 µm. **D)** (Left panel) Quantification of SPINK3-positive cells in WT and
Catsper1-/- spermatozoa, analyzing both head and tail labeling. Data represent the mean ±
SEM (n=3). Significant differences ($p < 0.05$) between WT and *Catsper1*-/- groups are
indicated. (Right panel) Distribution of SPINK3 labeling in spermatozoa head and tail
regions, expressed as percentages for WT and *Catsper1*-/- samples.

Figure 3. The inhibitory effect of SPINK3 on capacitation is not mimicked by a standard trypsin inhibitor A) Non-capacitated (NC) sperm loaded with Fluo-3 AM were exposed to SPINK3 (13 µM) or SBTI (1 µM). After washing with fresh media containing NaHCO$_3$, sperm cells were attached on laminin-coated slides and calcium signal (F0) was monitored before and after BSA was added (left panels). The fluorescence increments of individual live cells (F-F0)/F0 after 24 sec after BSA addition were plotted for each condition: control (-), SPINK3 or SBTI (right panels). *p < 0.05 significant differences in the percentage of responsive cells (>0.5 fold), compared to control untreated sperm. **B)** Acrosome reaction was induced by 50 µM progesterone (P4) after capacitation. As control, NC sperm cells were incubated with the Ca^{2+} ionophore A23187 (10 µM). Data represent mean ± SE; N = 3. *p < 0.05 significant differences compared to NC sperm. **C)** Cells under capacitating (CAP, 60 min) or non-capacitating (NC) conditions were loaded with the pH-sensitive probe BCECF-

888 AM (2 µM), and fluorescence was measured (Ex 520/Em 605 nm). *p < 0.05 significant
889 differences compared to NC sperm.

890
**891 Figure 4. SPINK3 blocks capacitation-associated cholesterol efflux and membrane
destabilization.** Mouse epididymal spermatozoa were loaded with Bodipy Cholesterol
(BODIPY/Ch) **(A)** or M540 and Yo-Pro1 **(B)** with or without SPINK3 (13 µM) and incubated
under capacitating (CAP) or non-capacitating (NC) conditions. n=3 LD: low disordered
membrane (Yo-Pro1⁻/M540⁻). HD: high disordered membrane (Yo-Pro1⁻/M540⁺). Percentage
of M540⁺ cells is expressed relative to the total live cells (Yo-Pro1⁻). (*) Significant
differences among treatments, p < 0.05. Scale bar: 10 µm.

898
Figure 5. Effect of SPINK3 on *in vitro* fertilization. Mouse sperm were incubated in the
absence or presence of rSPINK3 (13 µM) under capacitating conditions (CAP). After

capacitation, sperm (5 thousand total cells, 10 µl) were added to groups of COCs (5-6
COCs, 90 µl) obtained from superovulated females, (n=4). **A)** Representative images of two-
cell embryos and blastocysts obtained 24 hs. or 5 days post-IVF, respectively. **B)**
Quantification of two-cell embryos and blastocysts relative to the total oocytes (n=113). (*)
Significant differences compared to the control without SPINK3, $p < 0.05$. **C)** Sperm
recovered after 4 hs. of co-culture were incubated with α-SPINK3 antibodies to detect
sperm-bound SPINK3 bound to sperm by immunofluorescence. Scale bar: 10 µm.

908
909 **Supplementary Figure 1.** Immunostaining was performed with anti-CATSPER1 antibody on
910 capacitated sperm that were previously incubated w/o SPINK3 for 30 min. Representative
911 images from three independent experiments are shown. Scale bar: 10 µm.
912

913
914
Supplementary Figure 2. *Effect of SPINK3 and SBTI on the proteolytic degradation of*
CATSPER1 in vivo. Mouse epididymal sperm (n=3) were incubated with or without SPINK3
(13 µM) or SBTI (1 µM) and exposed to CAP conditions. Cell lysis was performed on
aliquots of different treatments at 0 and 60 min. Proteins (0.5×10^6 cells/ml) were separated
on a 10% polyacrylamide gel and revealed by Western blot, using anti-CATSPER1 followed
by anti-Tubulin.

921
Supplementary Figure 3. *Immunodetection of recombinant SPINK3 incubated with*
epididymal spermatozoa from wild type and Catsper1 KO mice. Mouse epididymal NC
spermatozoa were incubated in the presence of SPINK3 (13 µM) and immunofluorescence
assays were performed using rabbit anti-SPINK3 (α-SPINK3) followed by anti-IgG-rabbit-
Alexa488 (green). (n=3) Scale bar: 10 µm. Images are representative of three independent
experiments. BF, bright field.

Supplementary Movie 1. *3D projection of SPINK3 and plasma membrane markers FM4-64*
FX and CellBrite Fix on the sperm flagellum, related to Figure 2. The video shows a rotating
view along the y-axis, providing a spatial visualization of SPINK3 (red) localization on the
plasma membrane of the sperm principal piece. The membrane is labeled with FM4-64 FX
(green) and CellBrite Fix (cyan), highlighting differences in staining intensity between the
midpiece   and   the   principal   piece.
https://drive.google.com/drive/folders/1nl2lqlkbBCWcofPsq-
asTwwj_lwKv36C?usp=drive_link