`Noah B Lemke, PhD
4921 Crawford St 3
Houston, Texas, USA 77004
E-mail: noahlemke22@gmail.com
Phone: US +1 630 347 3302

RESEARCH ARTICLE: Temporal Trends: Phase-shifted time-series analysis reveals highly correlated reproductive behaviors in the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae)

AUTHORS
Noah B Lemke 1, Chujun Li 2,3, Jeroen De Smet 4, Jeffery K Tomberlin 1

AFFILIATIONS
1 - Department of Entomology, 370 Olsen Blvd Suite 412, Texas A&M University, College Station, Texas, USA 77843
2 - Sun Yat-Sen University, Guangzhou, Guangdong, China
3 - Guangzhou Unique Biotechnology, Room 1717, 101 Science Avenue, Huanpu District, Guangzhou, Guangdong, China
4 – Research Group for Insect Production and Processing, Department of Microbial and Molecular Systems, KU Leuven, Campus Geel, Kleinhoefstraat 4, 2440 Geel, Belgium

* To whom correspondences should be addressed: noahlemke22@gmail.com

ENGLISH ABSTRACT (213 words)

For those researching reproduction of the black soldier fly (BSF), finding meaningful correlations

between mating and downstream reproductive fitness metrics, owing to complex reproductive

behavior and physiology, has been challenging. On the one hand, mating is not necessarily

deterministic of fertile eggs being produced, since females might lay unfertile eggs both before

and after mating. But conversely, the process is not 'random' either, since fertility is predicated on

the exchange of gametes during mating. To investigate the relationship between mating and other

reproductive metrics, a cross-correlation analysis between lagged time series was conducted on

data previously collected during a semi-outdoor greenhouse experiment. This process compared

the average percent correlation between un-lagged mating with oviposition in traps, oviposition

outside traps, total weight of eggs deposited, and hatch rate, which were lagged 0 to -6 days. This

revealed that average mating was 97.54% correlated with oviposition in traps lagged by -2 days.

For oviposition outside of traps, total egg weights, and hatch rate, these were also ~71.63%

correlated, 60.29% correlated, and 75.16% correlated with mating respectively, which were

optimized when the time series were lagged $n = -3$ days. These trends likely have a biological

basis since flies decrease their reproductive activity as they expend energy and senesce, aligning

with temporal patterns consistently observed during BSF rearing.

KEYWORDS

21   INTRODUCTION

Insect reproductive behavior is complex (Lemke et al. 2023, Meneguz et al. 2023,

Tomberlin et al. 2025) owing not only to the wide diversity in mating systems and reproductive

modes across species (Kokko et al. 2014, Normark 2014), but also the plethora of alternative

mating tactics, dynamic evolutionary optimums for fitness, phenotypes, and personas expressed

by populations or individuals of the same species (Shuker and Simmons 2014). As an emergent

phenomenon (Macklem 2008), reproductive behavior is influenced by several echelons of

processes occurring either at a lower organization level or earlier during the life of an insect. These

include social context hormonal state, physiology and gross anatomy, nutritional legacy, cellular

processes, genetics, eco-evolutionary history (Alcock and Thornhill 2014) and much more. On top

of this, empiricists must also consider the covariates and intermediate factors that can act on

behavioral outcomes, such as the testing environment or artifacts of the experimental design

(Tomberlin et al. 2025). As such, this level of complexity can require sophisticated analytical

techniques (Smetana et al. 2025), as well as a broad intuition of insect biology.

Take for example the black soldier fly (BSF), *Hermetia illucens* (L.) (Diptera:

Stratiomyidae), which is a large lekking fly (Tomberlin and Sheppard 2001) of neotropical origin

(Kaya et al. 2021). BSF has been recently domesticated (Rhode et al. 2020) and is now mass-

reared throughout the globe for various applications, such as waste management and subsequent

conversion of these materials into insect biomass that can serve as alternative feed ingredients for

pets and livestock. For this reason, the operation costs of mass-rearing facilities still need to be

driven down to make facilities more economic since the goal of a mass-culture program is to

produce the maximum number of fertile females in as short of time and inexpensively as possible

with minimal labor and space (Boller 1972).

Due to protandry, adult males emerge from pupae prior to females (Tomberlin et al. 2002,

Generalovic et al. 2025). However mating studies typically control for both age and sex by taking

a portion of reared flies and sorting them to create test populations (e.g., 100 males and 100

females). In addition, the labor to sort these flies can be extensive and time consuming, and so a

narrow range of ages can be expected (e.g., 0-3 days; 1-4 days). This approach can be contrasted

with the industrial process, in which one (or several) batch of pupae are placed directly inside a

cage, or in an area adjacent to the cage and allowed to emerge (Coudron et al. 2025). If multiple

batches of pupae are added to the cage continuously, then this can lead to the breeding cage

containing flies of many mixed ages and reproductive statuses (Dickerson et al. 2024, Lemke et

al. 2025), since post-reproductive adult flies might live several weeks or more (Harjoko et al.

2023), especially when supplemented with sugar-water or honey (Thinn and Kainoh 2022, Klüber

et al. 2023, Barrett et al. 2025), and mating has been shown to extend female longevity (Harjoko

et al. 2023) potentially through sperm digestion (Manas et al. 2024).

Irrespective of the exact method used to stock cages, what follows is that high densities of

individuals engage in 'lek' mating which typically plateaus 2-3 days post-emergence. Males are

drawn towards UV-light (Zhang et al. 2010), and females temporarily join these aerial swarms to

mate (Lemke et al. 2024). After an aerial courting ritual, copulation commences when male and

female lock genitalia and remain nearly motionless on the floor or walls of the cage for ~35 minutes

(Manas et al. 2025). Near the end of this time, sperm is transferred via the male's aedeagus into

the female (Manas et al. 2025), the two separate, and groom themselves (Laudani et al. 2024).

Following this, there is an intermittent period of several days (Tomberlin and Sheppard

2002) where sperm travels through the female's genital tract (Malawey et al. 2019) and is stored

in their spermathecae until the moment of oviposition (Manas et al. 2024), whereupon the eggs are

fertilized. To induce egg-laying, an attractant is then introduced (which can be a natural substrate,

or a synthetic blend pumped through the cage) to entice females to lay their eggs in a trap.

Oviposition activity typically then peaks between days 5-6 days post-emergence. To lay their eggs,

females normally seek out an area of dry cracks adjacent to a larval rearing substrate (but obviously

such a substrate may not actually be present if the attractant is a synthetic blend). Traps are then

replaced, and any deposited eggs are harvested by technicians. 2-5 days later, the neonates hatch

from their eggs and begin wandering towards food (Giannetti et al. 2022).

One major area complicating analysis is the temporal structure of the data. Mating and

oviposition activity peak several days apart (Tomberlin and Sheppard 2002, Dickerson et al. 2024,

Meyermans et al. 2025). Moreover, outside of these peaks there is very little activity, leading count

data to be highly zero-inflated (Smetana et al. 2025). To capture information about both,

observational experiments need to be run for several days (Chiabotto et al. 2024, Dickerson et al.

2024, Lemke et al. 2024), which presents a logistical hurdle because, unless remote videography

is used, human observations cannot accurately be made across this whole time period. A trade-off

then exists between capturing data across the entire cycle (e.g. during daylight hours for 1 week)

or focusing more narrowly on a particular period (e.g., the time-windows for mating or

oviposition). Furthermore, the temporal delay between copulation (i.e., mating), oviposition, and

eggs hatching has historically made predicting the values of one from the others difficult, leading

some reports to say that mating is only loosely related to the latter (Dickerson et al. 2024).

Hence, the objective of this study is to provide an additional statistical tool for those

wishing to monitor the success of their BSF colonies with regards to fertile egg production. This

study examined whether the correlation between BSF mating and other reproductive metrics could

be improved through phase shifting the time series data for the latter metrics. While the procedures

for time series analyses are formalized and widely used in the social sciences, biology, and

economics (Shumway and Stoffer 2017a), such analyses are quite new when it comes to

interpreting black soldier fly mating data (Lemke et al. 2024). We hypothesized that by adjusting

the lag between time series, that mating could be a useful predictor of downstream fitness metrics

such as oviposition, total egg weight, and hatch rate.

97   METHODS

Data Acquisition — Data for this analysis was originally collected as part of a semi-outdoor

greenhouse study conducted at the F.L.I.E.S. Facility at Texas A & M University throughout 2022

(Lemke et al. 2025). The original experiment investigated the interaction between BSF cohort age

and oviposition attractant provisioning on fitness but did conduct a time-series analysis. 4 trials

took place from (a) March 07-13, (b) April 25-May 01, (c) October 06-12, and (d) December 10-

16, each lasting 7 days. During these observation periods, count data for mating and oviposition

were recorded hourly from sunrise to sunset, while eggs traps were collected daily from cages at

noon and the total weight of collected eggs measured gravimetrically. After a 5-day incubation

period, hatch rate of the eggs for each collection was also assessed. A full description of the

experimental design can be found in Lemke et al., 2025; and similar procedures have been used in

other studies (Dickerson et al. 2024, Laursen et al. 2024, Lemke et al. 2024). During each trial, 4

experimental treatments (see below) were tested on the production of a 0.93 m3 breeding cage,

supplied with 100 male and 100 female BSF. Each of these 4 treatments was replicated 3 times,

for a total of 12 experimental units, yielding 336 cage-days of observations (4 treatments x 3

replicates x 2 blocks x 2 trials x 7 days).

Data Structure & Cleaning — The levels of the original experiment's treatment categories

followed a 2 × 2 factorial design, with attractant provisioning having four total combinations of

levels (i.e., (a) whether the attractant provided initially or (b) delayed until 1-days post peak

mating; and (c) whether the attractant contained substrate or (d) was an empty control).

Additionally, age group also had an additional two blocks (two trials each, of which all flies were

either "mixed" or "same" age). For the age groups, "young" flies were 1-4-day-old prior to their

introduction into mating cages, whereas the "mixed" cohort could have ages between 1-16-days.

However, because industrial producers of BSF do not delay the age of their flies prior to

release, for this analysis, we only considered data for the "young" treatment in which this artifact

was absent. Likewise, providing attractant boxes lacking attractant is also not representative of

industrial rearing conditions and was included in the original experiment as a negative control, but

was not taken into consideration for this analysis. Moreover, recent evidence has shown that virgin

females have no inherent preference towards a group of volatile organic compounds (VOCs) that

later caused attraction or avoidance behaviors in mated females (Klüber et al. 2024). And so, in

addition to "mixed" age flies, replicates for which there was never an attractant provided (labeled

as "IC" and "DC" respectively for Initial and Delayed Controls) were redacted from the data set.

This yielded 168 rows of data (2 trials × 12 cages × 7 days) (i.e., half of the original 336 cage-

days) that could further be grouped by cage position (i.e., #1-12). The reduced data frame

(containing only data from "young" flies that were provided attractant initially = "IO" or delayed

"DO") was then reformatted using the PivotTable function in MS Excel for Mac Version 16.97

(25051114), to make calculating the averages much easier.

Time Series Analysis — To examine temporal patterns, the data were plotted using MSExcel inbuilt

plotting function. By averaging values by day (of observations taken hourly from sunrise to sunset)

autoregression (as used by Lemke et al. 2024) was not necessary for this analysis because there

were no gaps in the data that needed to be filled. Visually inspecting the plots revealed that phase

shifting the data might lead to stronger cross-correlations, since mating and oviposition follow a

similar temporal pattern, albeit with oviposition lagging by two days.

Percent Correlation Coefficient Matrix — To make cross-correlations, a Pearson correlation

matrix was constructed which included all pairwise combinations for mating and oviposition

across treatments. For example, the mating-oviposition (in traps) matrix considered the mean

average of IO-mating, as correlated to the mean average of DO-mating, IO-oviposition, and DO-

oviposition. For the resulting matrix, only the lower triangle of correlation coefficients (i.e., cells

underneath the identity diagonal) were considered since the matrix was symmetrical. Values were

then squared (r-squared), which is also considered a measure of effect size (Rights and Sterba

2020), and multiplied by 100 to yield the percent correlation coefficient (PCC). From this, the

average Pearson PCC of the matrix was calculated (i.e., by summing all values and dividing by

the number of values within the lower triangle), as well as the standard deviation and standard

error (of n = 6 values under the diagonal). For this step, Pearson's correlation was chosen to

investigate when a linear increase in mating corresponded to a linear increase in oviposition, or

vice versa. However, using Spearman's rank correlation would allow a looser interpretation in

which an increase in mating corresponded to *any* increase in oviposition, not necessarily a linear

one (and was used in the Clustered Heat Map analysis below).

Phase Shifted PCC — Next, a series of PCC matrices were calculated to determine what among of

phase shifting optimized PCC. This was done for each pair of reproductive metrics (mating with

mating = autocorrelation, mating with oviposition in traps, mating with oviposition in cages,

mating with total egg weight, and mating with hatch rate). Data were phase shifted n = 0, −1, −2,

−3, −4, −5, and −6 days, yielding $y_t, y_{t-1}, y_{t-2}, \ldots y_{t-6}$. To phase shift the data, the first n-days of the

data were deleted and the remainder shifted forward, creating an empty tail that was filled with 0-

values in order to preserve the length of the time series. Mating was always kept 'un'-shifted,

primarily because it occurs first in the mating sequence (and so is more biologically relevant), but

also to reduce the total number of symmetrical combinations (e.g., when both mating and

oviposition were phase shifted equivalently). The average PCC across this continuum of phase

shifting was then plotted in MSExcel and examined visually to determine candidates for

optimization by finding local maxima/minima.

Clustered Heat Map — Finally, a clustered heat map was generated for all metrics combined. This

was constructed by first calculating the pairwise Spearman rank correlation between reproductive

metrics that were both unmodified as well as phase shifted. The number of days that each metric

was phase shifted was determined by taking the number of days that optimized the average PCC

in the above analysis and also using their neighboring values of ± 1 day (i.e., if an optimum was

found for oviposition with a phase shift of n = −2, then this along with n = −1 and n = −3 were

considered). The correlation matrix was calculated using the corr() function in RStudio (Wei and

Simko 2024), using Spearman's rank coefficient (since the functions above were not strictly

monotonic). These values were then plotted as a clustered heatmap using the corrplot() function in

RStudio.

184   RESULTS

Visual Analysis — As a first step, data for mating x day, oviposition x day, total egg weight x day,

and hatch rate x day was visualized by plotting in MS Excel. Raw values with ± SD can be found

in Table S1 and were plotted in MS Excel (Figures 1, 2). Doing this revealed average mating

peaked on day 1 (of the cycle), whereas average oviposition peaked on day 4. The trend for each

was to generally decrease following this peak, however mating was not strictly monotonic, in that

there was a second small peak between days 3-4. Cumulative average total egg weight and hatch

rate both plateaued by day 5, with daily values generally decreased after reaching a peak on day 3.

Qualitatively, it then appeared that mating would have the strongest correlation to oviposition

when the latter was shifted forward n = −2 days. However, because of the plateau, total egg weight

and hatch rate might be more strongly correlated with mating when phase shifted greater than −1

days.

Cross Correlation of Time Series Data — The initial analysis considering Mating x Oviposition

for each treatment was optimized when oviposition was lagged by −2 days, with individual values

within the matrix ranging from 97.56 - 98.66 PCC (Table S2), indicating that PCC could be

optimized by lagging data. Hence, for each reproductive metric, the average Pearson correlation

coefficient was calculated between mating, and that second metric phase shifted by n = 0, −1, −2,

−3, −4, −5, and −6 days. These were then plotted to visually inspect where (local) optima lie.

Mating autocorrelation (Figure 3) was minimized for a 1-day lag (34.48%), and highest for a 3-

day lag (96.10%). The comparison between mating and oviposition (in traps), average PCC was

maximized with a phase shift of n = −2 days. Doing this increased the PCC from 31.19% to

97.54%, a 3.12-fold difference (Figure 4). For the comparison between mating and oviposition (in

cage material) (Figure 5), average PCC was maximized with a −3-, −4-, or −5-day time lag (~72%)

and minimized with a 0-day time lag (19.71%). For the comparison between mating and total egg

weight (Figure 6), a local maximum was found with a −3-day lag (60.29%) up from 29.22% with

no lag, representing a 2.06-fold increase. Interestingly though, the highest value was for total egg

weight with −6-day time lag, though this is suspected to be a statistical artifact (see Discussion

below), and for this reason the local optima at n = −3 days is preferred. Lastly, the comparison

between mating and hatch rate (Figure 7) also had a maximized average PCC with a −3-day lag

(75.16%) up from 40.48%, representing a 1.85-fold increase.

Clustered Heat Map — A clustered heat map was constructed to reveal grouped relationships

between mating and downstream reproductive metrics that were either not phase shifted or those

that were phase shifted a number of days (±1), equal to the optima found in the previous step.

Within the clustered heat map, mating was found in a cluster together with oviposition (in traps)

phase shifted 2 and 3 days. This cluster was most closely related to a larger cluster containing total

egg weight phase shifted −2 and −3 days well as hatch rate phase shifted −2 days and −3 days.

Interestingly, the most distant cluster to mating contained the original unmodified oviposition in

traps, total egg weight, and hatch rate metrics, as well as oviposition in cages phase shifted −3

days.

227   DISCUSSION

The objective of this study was to examine whether correlations between time series data

of reproductive metrics recorded during a BSF breeding study could be improved by comparing

time series that were phase shifted/lagged. Since there is an obvious temporal component to mating

behaviors, this study greatly improves upon that of Lemke et al. 2025 who did not consider time

series analyses as part of the original study, focusing instead on statistical differences between

treatment effects. Indeed, the results of this analysis show that lagging the data forward in time

increased the Pearson percent correlation coefficient (PCC), i.e., effect size, between mating and

other downstream reproductive metrics substantially. By doing this, mating was 97.56 to 98.66%

correlated with oviposition in traps when the latter was lagged −2 days. Likewise, for oviposition

outside of traps, total egg weights, and hatch rate, these were ~71.63% correlated, 60.29%

correlated, and 75.16% correlated with mating respectively, which were optimized when the time

series were lagged n = −3 days. A clustered heat map confirmed this relationship, showing mating

was most closely related (i.e., had the strongest correlation) with a cluster of metrics that included

oviposition, total egg weight, and hatch rate, that were each phase shifted by these optima. In

addition, the clustered heat map showed that mating was also least correlated with these same

fitness metrics if they were unmodified. Thus, this experiment highlights the importance for BSF

producers to considering correlations between reproductive metrics measured at different time

points in production, as well as understanding whether these correlations are rooted in biology.

247  At first glance, the link between mating and these lags generally corresponded biologically

248 to temporal patterns within the data. The initial visual inspection of the reproductive metrics

249 (plotted by day) revealed that mating peaked on Day 1 (Figure 1). The trend for the mating time

250 series was then generally decreased but had a small 'hump' on days 3-4 (Figure 1). This suggested

251 that individuals can engage in multiple matings, as did the fact that mating autocorrelation was

252 maximized on these same days (Figure 3). Indeed, the multiple mating in BSF for both males and

253 females has now been verified by many recent experiments (Hoffmann et al. 2021, Chiabotto et

254 al. 2024, Muraro et al. 2024). Similarly, oviposition in traps peaked on day 3 and then decreased

255 monotonically (Figure 1), meaning the distance between peak mating and peak oviposition

256 (occurring on Days 1 and 3, respectively) was $n = -2$ days. This was equivalent to the optimized

257 time lag (Table S1, Figure 4).

258  Interestingly, the peak for total egg weight occurred prior to the peak for oviposition counts

259 in traps (Figures 1, 2), indicating that perhaps larger egg clutches are laid by females earlier in the

260 cycle, and smaller ones later in the cycle. A prior experiment reports multiple oviposition events,

261 but that subsequent clutches were smaller and infertile (Nakamura et al. 2016). Alternatively, it

262 could also mean that later in the cycle there are instances where females extend their ovipositors

263 but do not lay eggs (reflecting an inaccuracy in the original data set). Lastly, hatch rate peaked on

264 day 3 (i.e., eggs laid on day 3 had the highest hatch rate after their 5-day incubation period). The

265 hatch rate then decreased monotonically, suggesting that the process of sperm storage within the

266 female takes ~2 days (since before day 3 the hatch of any eggs collected is relatively low), aligning

267 with a recent report (Manas et al. 2024). Alternatively, because the data were gathered in an semi-

268 outdoor environment, it could also be that this peak in fertility corresponded to days with more

optimal environmental conditions, such as higher humidity that might reduce desiccation, or

ambient temperatures falling within thermal tolerances (Chia et al. 2018, Addeo et al. 2022).

Another interesting pattern in the data is that the PCC between mating and oviposition

(inside of cages) peaked earlier than did the PCC between mating and oviposition (outside of

cages), which seem to correspond to a 1-day earlier peak in oviposition in traps versus cages

(Figure 1). Since the data used here include only 1-4-day-olds, the uptick in oviposition outside of

cages can't easily be explained by delayed mating, as was the case for prior experiments which

did this to created aged test-populations (Dickerson et al. 2024, Lemke et al. 2025). Instead, the

pattern might indicate that the flies become less discriminating as they age. There could also be an

environmental effect, whereby the temperature happened to increase on those days (for both trials),

forcing the flies away from ovipositing in the trap, which absorbed/reflected a lot of heat (Lemke

et al. 2025). Another explanation could also be a snowball-effect whereby day 4-5 of the

experiment, some flies had haphazardly laid eggs in the cage material early on, and then the

volatiles from those eggs (Klüber et al. 2023) attracted additional flies to lay outside of the trap.

The temporal trends reported here generally align with what has been observed in previous

BSF breeding experiments (Tomberlin and Sheppard 2002, Zhang et al. 2010). For example, early

experiments on BSF showed that mating peaked on days 1-2, with oviposition peaking several

days later (Tomberlin and Sheppard 2002, Park et al. 2010). This has been a consistent trend in

mating experiments to this date (Jones and Tomberlin 2021, Dickerson et al. 2024, Lemke et al.

2024, Meyermans et al. 2025). In addition, more recent studies have demonstrated multiple mating

(Hoffmann et al. 2021, Chiabotto et al. 2024, Muraro et al. 2024), multiple oviposition (Muraro et

al. 2024), and physiological processes occurring during these 'lulls' in activity (Manas et al. 2024),

as well as the potential for a neurophysiological switch during this time for females to become

receptive to volatiles (Klüber et al. 2024). As such, it is reassuring to know that these patterns

continue to hold true after two decades of BSF rearing, both throughout the globe, as well as in

indoor and semi-outdoor rearing conditions.

However, because the original trend for mating in this experiment was to generally

decrease after day 1, the Pearson PCC between mating and any other reproductive metric appears

to have been optimized when the phase shift applied also transformed the time series to become a

monotonic decreasing function. Of course, this is reflective of the fact that the original experiment

only used a single batch of flies per cage, and it is expected that this trend will not hold true once

new flies are added to cages through continuous rearing (see below for suggested analysis to

perform in this case). For the batch-reared data analyzed here, although mating was not strictly

monotonic, in general its trend was to decrease. In effect, the high correlations may be a statistical

artifact (i.e., a spurious correlation) resulting from qualitatively similar trends between un-lagged

mating (i.e., decreasing) with the lagged oviposition or another lagged metric (i.e., transformed to

become decreasing). Such was also seen in the temporal autocorrelation function for mating, which

was lowest between days 0 and 1. Lagging the data n = -1 days effectively removed the peak on

day 1, and made the new mating peak occur between days 3-5 (shifted to days 2-4). Hence,

autocorrelation was highest on days 3-5, because lagging the function by this amount effectively

transformed it into a strictly decreasing (i.e., monotonic) function. Likewise, the PCC between

mating (unadjusted) and total egg weight was highest when egg weight was lagged by −6 days

(creating a strictly decreasing function to qualitatively match mating). Still, these trends have a

biological basis that resonates with long held understanding of temporal patterns in BSF

reproduction, in which reproductive activity decreases with time as flies exhaust their energy

stores, and senesce (Harjoko et al. 2023, Dickerson et al. 2024). Additionally, the cumulative

distribution for total egg weight (i.e., by adding each day's value to the previous sum) plateaus

around n = −6 days, corresponding with the global maximum PCC.

Of course, correlation does not imply causation, but to some extent we can infer

directionality since mating generally occurs prior to oviposition, oviposition before any eggs are

collected, and total egg weight necessarily occurring before any are hatched. In addition, all these

quantities intuitively must co-vary with one another, given that they are predicated on optimized

abiotic conditions (e.g., light, temperature, humidity).

Limitations and how to adapt this study for Industry — This analysis is limited largely because it

uses a dataset collected from two disjunct rearing cycles only lasting 7 days each. For industrial

programs that might need to produce high volumes of eggs/neonates, this analysis could be applied

but would need to be modified in one of the following ways:

First, for those practicing batch rearing (Coudron et al. 2025, Lemke et al. 2025), mating

might instead peak on day 2 or 3 of a breeding cycle rather than day 1 which would make sense

due to differences in the emergence time between males and females (i.e., protandry) or their initial

period of inactivity. In this case, since monitored mating activity would increase suddenly and then

decrease, the mating time series would no longer be monotonic, and might be qualitatively similar

to some of the other reproductive metrics. Similarly, the time series might be non-monotonic if

BSF experience several activity peaks as a result of multiple mating. In either case, a Spearman's

Rank correlation could be calculated instead of Pearson's to loosen some of the assumptions and

examine increases in non-monotonic functions, which is what was done to construct the clustered

heat map presented here (Figure 8, Table S2).

If instead, a producer operates their breeding cages continuously (Dickerson et al. 2024,

Coudron et al. 2025, Lemke et al. 2025), then it might be better to calculate a rolling window (i.e.,

a fixed time frame of e.g. 7 days, calculated each day) from which long term temporal patterns

could be ascertained (such as, whether mating on certain days is correlated with oviposition on

others, or whether there are weekly and seasonal trends across operations). Such patterns could be

also modeled arithmetically through an ARIMA model (Shumway and Stoffer 2017b), which are

traditionally used in stock price and commodities forecasting. Application of this technique will

be complex however if there is a non-stationary trend or autocorrelation between rearing cycles

(i.e., decreases over time due to genetic deterioration, or increases corresponding with sudden

increases in population density).

The extent to which these are practical to implement for BSF breeding will of course

depend on the elegance of behavioral assays and the information gleaned from them (Tomberlin

et al. 2025). Standard metrics like egg weight and hatch rate, which are simple enough to measure

gravi-, or volu-, metrically, can of course be formatted as Time Series data as has been done here.

In addition, there are also many remote detection methods proposed for use in insect agriculture

(Manoukis and Collier 2019, Nawoya et al. 2024), and some have started to be realized. These

include FlyCount (James et al. 2024), a program that monitors population density of adults passing

through a narrow opening prior to entering the breeding cage, as well as HistoNet (Sharma et al.

2020) to automatically assign size-classes to BSF. Counting the number of flies in a cage,

recording their sizes, and monitoring fertile egg production in relation to these factors

longitudinally (Dickerson et al. 2024) will be one step, but an important one, towards

understanding the system and increasing its predictability in a complex system.

ACKNOWLEDGEMENTS

We would like to thank our anonymous reviewers, as well as the group of researchers who

helped collect the original data set: Amy J. Dickerson, D. A. Salazar, J. Emmanuel Mendoza, Lisa

N. Rollinson, Erin Harris, Sienna McPeek, Rachel MacNeal, and Drs. Chelsea Miranda, and Casey

Flint.

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship under Grant No. 1746932. Any opinion, findings, and conclusions or

recommendations expressed in this material are those of the authors and do not necessarily reflect

the views of the National Science Foundation. This research was also partially supported by funds

from the US State Department and the Belgium-Luxembourg Fulbright Commission under the

2026 US-Belgium Fulbright Scholar Program. The content of this report is explicitly that of the

authors and does not reflect the views of the Fulbright Program, nor the governments of the United

States, Belgium, or Luxembourg.

CONFLICT OF INTEREST

The data analyzed in this study were derived from material purchased from EVO Conversion

Systems LLC, a company with which JK Tomberlin has historically had a significant financial

interest. This conflict of interest is managed by a plan submitted to and approved by Texas A&M

University and Texas A&M AgriLife.

REFERENCES

Addeo NF, Li C, Rusch TW, et al. 2022. Impact of age, size, and sex on adult black soldier fly
Hermetia illucens L. (Diptera: Stratiomyidae) thermal preference. Journal of Insects as Food and
Feed. 8(2):129–139. https://doi.org/10.3920/JIFF2021.0076

Alcock J, Thornhill R. 2014. The evolution of insect mating systems. In: Shuker DM, Simmons
LW, editors. Evolution of Insect Mating Systems. Illustrated. United Kingdom: Oxford
University Press. p. 78–91.

Barrett M, Patel N, McCarry B, et al. 2025. Dietary preferences and impacts of feeding on
behavior, longevity, and reproduction in adult black soldier flies (Diptera: Stratiomyidae;
Hermetia illucens). https://doi.org/10.31219/osf.io/p74xq_v1

Boller E. 1972. Behavioral aspects of mass-rearing of insects. Entomophaga. 17(1):9–25.
https://doi.org/10.1007/BF02371070

Chia SY, Tanga CM, Khamis FM, et al. 2018. Threshold temperatures and thermal requirements
of black soldier fly *Hermetia illucens*: Implications for mass production. PLOS ONE. 13(11):1–
26.

Chiabotto C, Grosso F, Doretto A, et al. 2024. Observation of mating behavior using marked flies
of black soldier fly (*Hermetia illucens*) under sunlight condition. Journal of Insects as Food and
Feed. 10(12):2017–2029. https://doi.org/10.1163/23524588-20230165

Coudron CL, Adamaki-Sotiraki C, Yakti W, et al. 2025. Bugbook: Basic information and good
practices on how to maintain stock populations for *Tenebrio molitor* and *Hermetia illucens* for
research. https://doi.org/10.1163/23524588-bja10240

Dickerson AJ, Lemke NB, Li C, et al. 2024. Impact of age on the reproductive output of
Hermetia illucens (Diptera: Stratiomyidae). Crippen T, editor. Journal of Economic
Entomology.:toae107. https://doi.org/10.1093/jee/toae107

Generalovic TN, Zhou W, Zhao LC, et al. 2025. Repeatable phenotypic but not genetic response
to selection on body size in the black soldier fly. :2025.02.25.640052.
https://doi.org/10.1101/2025.02.25.640052

Giannetti D, Schifani E, Reggiani R, et al. 2022. Do it by yourself: Larval locomotion in the
black soldier fly *Hermetia illucens*, with a novel "self-harvesting" method to separate prepupae.
Insects. 13(2):127. https://doi.org/10.3390/insects13020127

Harjoko DN, Hua QQH, Toh EMC, et al. 2023. A window into fly sex: mating increases female
but reduces male longevity in black soldier flies. Animal Behaviour. 200:25–36.
https://doi.org/10.1016/j.anbehav.2023.03.007

Hoffmann L, Hull KL, Bierman A, et al. 2021. Patterns of genetic diversity and mating systems
in a mass-reared black soldier fly colony. Insects. 12(6):480.
https://doi.org/10.3390/insects12060480

James A, Seth A, Marcireau A, et al. 2024. FlyCount: High-speed counting of black soldier flies
using neuromorphic sensors. IEEE Sensors Journal.:1–1.
https://doi.org/10.1109/JSEN.2024.3504289

Jones BM, Tomberlin JK. 2021. Effects of adult body size on mating success of the black soldier fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae). Journal of Insects as Food and Feed. 7(1):5–20. https://doi.org/10.3920/JIFF2020.0001

Kaya C, Generalovic TN, Ståhls G, et al. 2021. Global population genetic structure and demographic trajectories of the black soldier fly, *Hermetia illucens*. BMC Biol. 19(1):94. https://doi.org/10.1186/s12915-021-01029-w

Klüber P, Arous E, Jerschow J, et al. 2024. Fatty acids derived from oviposition systems guide female black soldier flies (*Hermetia illucens*) toward egg deposition sites. Insect Science. 31(4):1231–1248. https://doi.org/10.1111/1744-7917.13287

Klüber P, Arous E, Zorn H, et al. 2023. Protein- and carbohydrate-rich supplements in feeding adult black soldier flies (*Hermetia illucens*) affect life history traits and egg productivity. Life. 13(2):355. https://doi.org/10.3390/life13020355

Kokko H, Klug H, Jennions MG. 2014. Mating systems. In: Shuker DM, Simmons LW, editors. Evoltuion of Insect Mating Systems. Illustrated. United Kingdom: Oxford University Press. p. 78–91.

Laudani F, Campolo O, Latella I, et al. 2024. Does *Hermetia illucens* recognize sibling mates to avoid inbreeding depression? entomologia. 44(5):1225–1232. https://doi.org/10.1127/entomologia/2024/2746

Laursen SF, Flint CA, Bahrndorff S, et al. 2024. Reproductive output and other adult life-history traits of black soldier flies grown on different organic waste and by-products. Waste Management. 181:136–144. https://doi.org/10.1016/j.wasman.2024.04.010

Lemke NB, Dickerson AJ, Tomberlin JK. 2023. No neonates without adults. BioEssays. 45(1):2200162. https://doi.org/10.1002/bies.202200162

Lemke NB, Li C, Dickerson AJ, et al. 2025. Heterogeny in cages: Age-structure and attractant availability impacts fertile egg production in the black soldier fly, *Hermetia illucens*. Journal of Insects as Food and Feed. Online. https://doi.org/10.1163/23524588-bja10275

Lemke NB, Rollison LN, Tomberlin JK. 2024. Sex-specific perching: Monitoring of artificial plants reveals dynamic female-biased perching behavior in the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). Insects. 15(10):770. https://doi.org/10.3390/insects15100770

Macklem PT. 2008. Emergent phenomena and the secrets of life. Journal of Applied Physiology. 104(6):1844–1846. https://doi.org/10.1152/japplphysiol.00942.2007

Malawey AS, Mercati D, Love CC, et al. 2019. Adult reproductive tract morphology and spermatogenesis in the black soldier fly (Diptera: Stratiomyidae). Annals of the Entomological Society of America. 112(6):576–586. https://doi.org/10.1093/aesa/saz045

Manas F, Labrousse C, Bressac C. 2025. Plastic responses in sperm expenditure to sperm competition risk in black soldier fly (*Hermetia illucens*, Diptera) males. Journal of Insect Physiology. 161:104751. https://doi.org/10.1016/j.jinsphys.2025.104751

Manas F, Piterois H, Labrousse C, et al. 2024. Gone but not forgotten: dynamics of sperm storage and potential ejaculate digestion in the black soldier fly Hermetia illucens. Royal Society Open Science. 11(10):241205. https://doi.org/10.1098/rsos.241205

Manoukis NC, Collier TC. 2019. Computer vision to enhance behavioral research on insects. Annals of the Entomological Society of America. 112(3):227–235. https://doi.org/10.1093/aesa/say062

Meneguz M, Miranda CD, Cammack JA, et al. 2023. Adult behaviour as the next frontier for optimising industrial production of the black soldier fly *Hermetia illucens* (L.) (Diptera: Stratiomyidae). Journal of Insects as Food and Feed. 9(4):399–414. https://doi.org/10.3920/JIFF2022.0055

Meyermans R, Broeckx L, Mondelaers J, et al. 2025. Exploring the potential of crossbreeding to enhance black soldier fly (Hermetia illucens) production. https://doi.org/10.1163/23524588-00001391

Muraro T, Lalanne L, Pelozuelo L, et al. 2024. Mating and oviposition of a breeding strain of black soldier fly Hermetia illucens (Diptera: Stratiomyidae): polygynandry and multiple egg-laying. Journal of Insects as Food and Feed. 1(aop):1–13. https://doi.org/10.1163/23524588-20220175

Nakamura S, Ichiki RT, Shimoda M, et al. 2016. Small-scale rearing of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae), in the laboratory: Low-cost and year-round rearing. Appl Entomol Zool. 51(1):161–166. https://doi.org/10.1007/s13355-015-0376-1

Nawoya S, Ssemakula F, Akol R, et al. 2024. Computer vision and deep learning in insects for food and feed production: A review. Computers and Electronics in Agriculture. 216:108503. https://doi.org/10.1016/j.compag.2023.108503

Normark BB. 2014. Modes of reproduction. In: Shuker DM, Simmons LW, editors. Evoltuion of Insect Mating Systems. Illustrated. United Kingdom: Oxford University Press. p. 78–91.

Park K, Kim W, Lee S, et al. 2010. Seasonal pupation, adult emergence, and mating of black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae) in artificial rearing system. :3.

Rhode C, Badenhorst R, Hull KL, et al. 2020. Genetic and phenotypic consequences of early domestication in black soldier flies (*Hermetia illucens*). Animal Genetics. 51(5):752–762. https://doi.org/10.1111/age.12961

Rights JD, Sterba SK. 2020. New recommendations on the use of R-squared differences in multilevel model comparisons. Multivariate Behavioral Research. 55(4):568–599. https://doi.org/10.1080/00273171.2019.1660605

Sharma K, Gold M, Zurbruegg C, et al. 2020. HistoNet: Predicting size histograms of object
instances. In: 2020 IEEE Winter Conference on Applications of Computer Vision (WACV).
Snowmass Village, CO, USA: IEEE. p. 3626–3634.
https://doi.org/10.1109/WACV45572.2020.9093484

Shuker DM, Simmons LW, editors. 2014. The evolution of insect mating systems. University of
Oxford: Oxford University Press.

Shumway RH, Stoffer DS. 2017a. Time series analysis and Its applications: With R examples.
Cham: Springer International Publishing (Springer Texts in Statistics).
https://doi.org/10.1007/978-3-319-52452-8

Shumway RH, Stoffer DS. 2017b. ARIMA models. In: Shumway RH, Stoffer DS, editors. Time
Series Analysis and Its Applications: With R Examples. Cham: Springer International Publishing.
p. 75–163. https://doi.org/10.1007/978-3-319-52452-8_3

Smetana S, Coudron C, Deruytter D, et al. 2025. BugBook: Data analysis methods in studies of
insects for food and feed. https://doi.org/10.1163/23524588-bja10209

[Software] Wei T, Simko V. 2024. R package "corrplot": Visualization of a correlation matrix.
https://github.com/taiyun/corrplot.

Thinn AA, Kainoh Y. 2022. Effect of diet on the longevity and oviposition performance of black
soldier flies, Hermetia illucens (Diptera: Stratiomyidae). Japan Agricultural Research Quarterly:
JARQ. 56(2):211–217. https://doi.org/10.6090/jarq.56.211

Tomberlin J, Sheppard D. 2002. Factors influencing mating and oviposition of black soldier flies
(Diptera: Straiomyidae) in a colony. Journal of Entomological Science. 37(4):345–352.
https://doi.org/10.18474/0749-8004-37.4.345

Tomberlin JK, Klammsteiner T, Lemke N, et al. 2025. BugBook: Black soldier fly as a model to
assess behaviour of insects mass produced as food and feed. https://doi.org/10.1163/23524588-
bja10225

Tomberlin JK, Sheppard DC. 2001. Lekking behavior of the black soldier fly (Diptera:
Stratiomyidae). Florida Entomologist. 84(4):729–730.

Tomberlin JK, Sheppard DC, Joyce JA. 2002. Selected life-history traits of black soldier flies
(Diptera: Stratiomyidae) reared on three artificial diets. an. 95(3):379–386.
https://doi.org/10.1603/0013-8746(2002)095[0379:SLHTOB]2.0.CO;2

Zhang J, Huang L, He J, et al. 2010. An artificial light source influences mating and oviposition
of black soldier flies, Hermetia illucens. Journal of Insect Science. 10(202):1–7.
https://doi.org/10.1673/031.010.20201

FIGURE CAPTIONS AND ALT TEXT

FIGURE 1. Plot of time-series data for average mating and oviposition counts. Oviposition

counts include for both inside the traps ("OviTrap") and outside the traps ("OviCage"). The

experimental unit consisted of a 0.93 m3 cage supplied initially with 100 Male and 100 Female

BSF aged 0-4 days. Observations were made each hour from sunrise to sunset for 7 days. Hourly

data from 2 experimental treatments (with n = 3 replicate cages for each treatment) was summed

to give a total count for each day. Then the totals for the 6 replicates was averaged to give the

average total counts with respect to each day.

FIGURE 1 ALT TEXT: Line plot of time-series data for BSF reproductive behaviors. The red line

represents average total mating counts. The yellow line represents average total oviposition counts

inside traps. The purple line represents the average total oviposition counts outside traps. Below

the line plot is a table giving the exact values of each data point.

FIGURE 2. Plot of time-series data for total egg weight (mg) and hatch rate (%) observed in

egg traps. The experimental unit consisted of a 0.93 m3 cage supplied initially with 100 Male and

100 Female BSF aged 0-4 days. Eggs were harvested from cardboard traps harvested each day at

noon and then weighed. From this, hatch rate for each day's collected eggs was calculated after a

5-day incubation period. Egg weight data were summed to give total weight collected for each

day, then the average taken across 6 replicates (n = 2 treatments with n = 3 cages). Because hatch

rate was a percentage, instead the mean hatch rate for each day was calculated, then averaged

across the 6 replicates. Full description of egg collection, weighing, and hatch percentage

calculations are given in Lemke et al. (2025).

FIGURE 2 ALT TEXT. Line plot depicting average total egg weight and the average mean percent

hatch rate. The green line represents egg weight, whereas the blue line represents percent hatch.

Beneath the plot is a table giving exact values for each data point.

FIGURE 3. Temporal autocorrelation of mating. Average Pearson percent correlation

coefficient between mating (phase shift = 0 days) and mating (phase shifted n = 0, −1, −2, −3, −4,

−5, −6 days). Percent correlation was calculated by squaring the Pearson correlation coefficient

and multiplying the value by 100. The average of this was taken by summing all values in the

lower triangle of the correlation matrix. Error bars presented are ± Standard Error.

FIGURE 3 ALT TEXT: Line plot depicting the autocorrelation between the average Pearson

percent correlation between mating counts and mating counts that are phase shifted between 0 and

−6 days. Error bars show ± Standard Error.

FIGURE 4. Average Pearson percent correlation coefficient between mating (phase shift = 0

days) and oviposition (phase shifted n = 0, −1, −2, −3, −4, −5, −6 days). Percent correlation was

calculated by squaring the Pearson correlation coefficient and multiplying the value by 100. The

average of this was taken by summing all values in the lower triangle of the correlation matrix.

Error bars presented are ± Standard Error.

FIGURE 4 ALT TEXT: Line plot depicting the autocorrelation between the average Pearson

percent correlation between mating counts and oviposition counts in traps that are phase shifted

between 0 and −6 days. Error bars show ± Standard Error.

FIGURE 5. Average Pearson percent correlation coefficient between mating (phase shift = 0

days) and oviposition outside of traps (phase shifted n = 0, −1, −2, −3, −4, −5, −6 days). Percent

correlation was calculated by squaring the Pearson correlation coefficient and multiplying the

value by 100. The average of this was taken by summing all values in the lower triangle of the

correlation matrix. Error bars presented are ± Standard Error.

FIGURE 5 ALT TEXT: Line plot depicting the autocorrelation between the average Pearson

percent correlation between mating counts and oviposition counts outside traps that are phase

shifted between 0 and −6 days. Error bars show ± Standard Error.

FIGURE 6. Average Pearson percent correlation coefficient between mating (phase shift = 0

days) and total egg weight (phase shifted n = 0, −1, −2, −3, −4, −5, −6 days). Percent correlation

was calculated by squaring the Pearson correlation coefficient and multiplying the value by 100.

The average of this was taken by summing all values in the lower triangle of the correlation matrix.

Error bars presented are ± Standard Error.

FIGURE 6 ALT TEXT: Line plot depicting the autocorrelation between the average Pearson

percent correlation between mating counts and total egg weights that are phase shifted between 0

and −6 days. Error bars show ± Standard Error.

FIGURE 7. Average Pearson percent correlation coefficient between mating (phase shift = 0

days) and mean hatch rate (phase shifted n = 0, −1, −2, −3, −4, −5, −6 days). Percent correlation

was calculated by squaring the Pearson correlation coefficient and multiplying the value by 100.

The average of this was taken by summing all values in the lower triangle of the correlation matrix.

Error bars presented are ± Standard Error.

FIGURE 7 ALT TEXT: Line plot depicting the autocorrelation between the average Pearson

percent correlation between mating counts and mean hatch rates that are phase shifted between 0

and −6 days. Error bars show ± Standard Error.

FIGURE 8. Clustered Heat Map of Time Series Data. The numbers appended to the data labels

indicate how many days forward the time series was lagged. "Mating" = Mating. "OviTrap" =

Oviposition in traps. "OviCage" = Oviposition in Cages. "EggWeight" = Total Egg Weight.

"Hatch" = Hatch rate. Correlation matrix was calculated using Spearman's rank coefficient. Color

formatting is applied with the conditional rule as follows: $-1 = 0^{th}$ percentile (Orange), $0 = 50^{th}$

percentile (White), $+1 = 100^{th}$ percentile (Blue).

FIGURE 8 ALT TEXT: Clustered heat map indicating that the Spearman's Rank correlation

between mating and the other reproductive metrics is highest when those other metrics are phase

shifted −2 days for oviposition in traps and outside of traps, between −2 and −4 days for egg

weight, and −3 days for hatch rate.

Mating and Oviposition

	1	2	3	4	5	6	7
Mating	33.25	2.50	5.58	5.75	0.92	0.50	0.00
OviTrap	0.00	4.00	11.58	1.83	1.58	1.92	0.50
OviCage	0.00	0.00	3.33	5.92	1.42	0.08	0.25

Day of Experiment

	1	2	3	4	5	6	7
EggWeight	0.23	0.81	0.79	0.66	0.38	0.25	0.37
Hatch	0.17	0.52	0.58	0.59	0.55	0.14	0.08

Egg Weight and Hatch Rate

Egg Weight (Average Total mg), Hatch Rate (Average of Mean %)

Day of Experiment

Average PCC: Mating-Total Egg Weight