How does stochasticity in learning impact the accumulation of knowledge and the evolution of learning?

Ludovic Maisonneuve[1,*] & Laurent Lehmann[1]

1. Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

* ludovic.maisonneuve.2015@polytechnique.org.

Abstract

Learning behaviors are crucial for humans and other animals to adapt to their environment by acquiring and using knowledge that enhances survival and reproduction. In particular, some learning behaviors allow populations to accumulate knowledge across generations ('cumulative knowledge' or 'cumulative culture'). Here, we examine how individual stochasticity in the production and social acquisition of knowledge influences the evolution of learning behaviors and cumulative knowledge. Using a mathematical model where learning is a stochastic process, we show that stochasticity in learning enhances cumulative knowledge by generating variability in knowledge levels: individuals who acquire more knowledge by chance are more likely to survive and reproduce, and therefore more likely to transmit their knowledge to the next generation. As knowledge accumulates, social learning exemplars tend to possess more of it, favoring greater investment of time in social learning. Because social learning provides access to substantial amounts of knowledge, selection favors the evolution of more effective learning, at the expense of a fecundity cost. Moreover, when knowledge enhances fecundity but not survival, we find that selection favors learning from parents rather than unrelated adults. This is because parenthood signals reproductive success and, therefore, a higher likelihood of possessing fecundity-enhancing knowledge, unlike random adults.

1 Introduction

Individuals use knowledge to perform adaptive behaviors, that is, behaviors that enhance survival and reproduction. This knowledge can be acquired either through personal experience (individual learning), such as trial-and-error learning (Ghirlanda and Lind, 2017), or from other individuals (social learning), for example through imitation (Zentall, 2006; Bates and Byrne, 2010). Social learning can occur through a wide variety of pathways (Cavalli-Sforza and Feldman, 1981; Boyd and Richerson, 1985; Laland, 2004; Kendal et al., 2018; Camacho-Alpízar and Guillette, 2023), including, for instance, vertical transmission (from parent to offspring) and oblique transmission (from unrelated older individuals). The way individuals allocate resources to different learning behaviors across their lifetime affects the population dynamics of knowledge and can support its gradual accumulation and refinement across generations. This process, known as cumulative knowledge or cumulative culture, is widely regarded as a key factor in the ecological success of humans (Henrich, 2015; van Schaik, 2016), and growing evidence suggests that it may also contribute to increasingly adaptive behaviors in non-human animals (Hunt and Gray, 2003; Sasaki and Biro, 2017; Jesmer et al., 2018; Gunasekaram et al., 2024).

Learners' individual and social learning strategies influence changes in knowledge within a population by shaping *cultural deviation* and *cultural selection*, the two key mechanisms underlying population change in any cultural trait (i.e., a socially transmissible trait; see Henrich and Boyd, 2002; El Mouden et al., 2014; Aguilar and Akçay, 2018; Nettle, 2020; Mesoudi, 2021). First, cultural deviation occurs when the learning process, on average, leads to differences between the knowledge of learners and that of their exemplars. This mechanism can either promote the accumulation of knowledge, for example, when learners produce new or refine existing knowledge by individual learning after social learning (e.g., Enquist et al., 2008; Aoki et al., 2012a; Nakahashi, 2013; Wakano and Miura, 2014; Kempe et al., 2014; André and Baumard, 2020; Denton et al., 2023), or constrain it, as when learners tend to acquire a lower level of knowledge than their exemplars (e.g., Henrich and Boyd, 2002; Henrich, 2004). Second, cultural selection arises when some individuals serve as exemplars more frequently, amplifying the transmission of their specific knowledge or cultural trait and generating variation in the transmission success of different knowledge or cultural variants (Cavalli-Sforza and Feldman, 1981; Boyd and Richerson, 1985; Micheletti, 2020). Cultural selection is generally expected to support cumulative knowledge, as it tends to favor cultural variants that are more well-adapted to environmental conditions, for example, when individuals with more adaptive cultural variants are more likely

to be chosen as learning models (e.g., Henrich, 2004; Powell et al., 2009; Kobayashi and Aoki, 2012), or when such variants enhance survival and reproductive success, thereby increasing opportunities for oblique (e.g., Nakahashi, 2010) and vertical transmission (e.g., Tureček et al., 2019), respectively. Mathematical models have shown that the strength of cultural selection, and thus its potential to drive cumulative knowledge, increases with population variance in cultural traits (Cavalli-Sforza and Feldman, 1981; Boyd and Richerson, 1985).

In turn, knowledge dynamics shape what individuals can acquire through social learning, thereby influencing selection pressures on the allocation of resources across different types of learning behavior. This feedback between knowledge accumulation and learning strategies gives rise to complex coevolutionary dynamics, also shaped by trade-offs between learning and other functions essential for survival and reproduction (for empirical evidence on such trade-offs, see Mery and Kawecki, 2004; Burger et al., 2008; Snell-Rood et al., 2011; Jaumann et al., 2013; Christiansen et al., 2016; Evans et al., 2017). A large body of theoretical work has examined these dynamics and how ecological factors influence the allocation of resources to different forms of learning and the emergence of cumulative knowledge (Nakahashi, 2010; Aoki et al., 2012b; Lehmann et al., 2013; Nakahashi, 2013; Wakano and Miura, 2014; Kobayashi et al., 2015, 2016; Mullon and Lehmann, 2017; Ohtsuki et al., 2017; Maisonneuve et al., 2025). Nevertheless, the effect of cultural selection is generally neglected in these models (with few exceptions such as Kobayashi et al., 2016), often because these models assume deterministic knowledge dynamics at the individual level, which removes variation that can be selected upon.

However, cultural selection is likely to influence the coevolution of cumulative knowledge and learning strategies, as learning is inherently stochastic; for example, the success of trial-and-error learning often depends on chance discoveries of adaptive cultural variants, leading to individual variation in knowledge. While not their primary focus, Kobayashi et al. (2016) showed that stochasticity in individual learning promotes investment in social learning by amplifying the effect of cultural selection (caused by knowledge-based choice of social exemplars), thereby increasing overall population knowledge and increasing what can be acquired socially. However, in contrast to the assumptions of Kobayashi et al. (2016), individuals in natural populations may not always be able to reliably assess the knowledge of others and to use it to guide their social learning decisions (Argyle and McHenry, 1971; Lutz and Keil, 2002; Wood et al., 2012; Jiménez and Mesoudi, 2019; Hirel et al., 2025). This underscores the need to investigate how stochasticity in learning affects the coevolution of cumulative knowledge and learning strategies in the absence of knowledge-based choice of social exemplars. Furthermore, we still lack predictions about

how stochasticity in learning affects the choice among transmission pathways (e.g., vertical vs. oblique), or the trade-offs between learning and other functions essential for survival and reproduction.

In this study, we examine how stochasticity in learning influences the dynamics of knowledge accumulation. We then examine how it ultimately influences the evolution of learning strategies, specifically, the pathways individuals use to acquire information through social learning and the trade-off between investment in learning and reproduction. We develop an evolutionary model in which learning is described by a stochastic process. The model tracks the evolution of the overall investment of resources between learning and fecundity, as well as the allocation of time across different types of learning (vertical, oblique, and individual learning). By allowing knowledge to accumulate over generations, our framework captures the coevolutionary feedback between learning behaviors and the accumulation of knowledge.

2 Model

2.1 Life-cycle

We consider a large, asexual population where individuals acquire adaptive information, i.e., information that enhances fecundity and survival. This includes information such as the location of food sources, the edibility of different foods, or instructions on how to build and use a tool. Each individual possesses a quantity of adaptive information, referred to as knowledge, which is treated as a quantitative variable in our analysis. In each generation, the population goes through the following life-cycle events (see fig. 1a). (1) Adults produce offspring depending on their knowledge. (2) Offspring acquire knowledge socially from adults and through individual learning through a stochastic learning process. (3) Parents die. Offspring go through a density-dependent survival stage, where their knowledge affects survival. Those surviving become the adults of the next generation, and the cycle starts again.

We assume that during stage (2) of the life cycle, offspring perform sequentially three different types of learning: first, they learn from their parent (vertical learning), then from a randomly selected adult (oblique learning), and finally by themselves (individual learning; see fig. 1b). All individuals have a fixed time available for learning, which we normalize to 1. Accordingly, the time allocated to vertical, oblique, and individual learning must sum to one. Two evolving

traits shape how individuals allocate their time across the three different types of learning: the amount of time v spent learning vertically; the amount of time o spent learning obliquely; so that $1 - v - o$ is spent learning individually.

Moreover, we assume that an evolving trait λ controls the overall trade-off between investment in learning and fecundity. Higher values of λ enhance the efficiency of all learning types but at the cost of reduced fecundity. For instance, λ may reflect developmental costs or the metabolic expenditure associated with learning ability.

Next, we describe in detail how traits affect the three life cycle stages (Section 2.2) and then outline the approach used to analyze the cultural and the evolutionary dynamics (Section 2.3).

2.2 Traits effects and knowledge throughout the life-cycle

2.2.1 Adult reproduction

We start by considering a focal adult from an arbitrary generation, characterized by traits $(v_\bullet, o_\bullet, \lambda_\bullet)$ and amount of knowledge $k_{\mathrm{p}\bullet} \in \mathbb{R}$. This knowledge is taken to be a realization of the stochastic learning process it underwent as an offspring (detailed in the next section). The expected number of offspring of the focal adult, referred to as fecundity, is assumed to be given by

$$f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}) = (1 - \lambda_\bullet)^\theta \; (f_0 + \eta_{\mathrm{f}} \, k_{\mathrm{p}\bullet}), \tag{1}$$

where f_0 is the baseline fecundity in the absence of investment into learning, and η_{f} is a conversion factor translating knowledge into fecundity. An individual's fecundity is reduced by its investment $\lambda_\bullet$ in learning, reflecting costs associated with acquiring learning abilities, as well as the metabolic expenses involved in the learning process itself. The parameter θ controls the strength and shape of this fecundity penalty: higher values of θ amplify the fecundity cost of learning, while lower values make this cost more gradual.

2.2.2 Offspring learning

Next, we consider an offspring of the focal adult. This focal offspring is assumed to inherit the parent's traits $(v_\bullet, o_\bullet, \lambda_\bullet)$, barring mutation. Let us denote by $k_\bullet(a) \in \mathbb{R}$ the knowledge this offspring bears at age $a \in [0, 1]$ (where $a = 0$ is birth). Building on the models of Kobayashi et al. (2016) and Maisonneuve et al. (2025) (where similarities and differences with our model

are outlined in Appendix A.1), we assume that the knowledge $k_\bullet(a)$ of the focal offspring at each age $a \in [0, 1]$ is a realization (or sample path) of a stochastic learning process defined by the following stochastic differential equation

$$\frac{\mathrm{d}k_\bullet(a)}{\mathrm{d}a} = \begin{cases} \lambda_\bullet \left(\beta_{\mathrm{v}} + \sigma_{\mathrm{v}}\, \eta(a)\right) \left[(1 - \epsilon)\, k_{\mathrm{p}\bullet} - k_\bullet(a)\right] & \text{for } a \in [0, v_\bullet) \\ \lambda_\bullet \left(\beta_{\mathrm{o}} + \sigma_{\mathrm{o}}\, \eta(a)\right) \left[(1 - \epsilon)\, k_{\mathrm{a}\bullet} - \rho\, k_\bullet(v_\bullet) - \left(k_\bullet(a) - k_\bullet(v_\bullet)\right)\right] & \text{for } a \in [v_\bullet, v_\bullet + o_\bullet) \\ \lambda_\bullet \left(\alpha + \sigma_{\mathrm{i}}\, \eta(a)\right) & \text{for } a \in [v_\bullet + o_\bullet, 1], \end{cases}$$

(2)

where the initial condition is $k_\bullet(0) = 0$. The term $\eta(a)$ is a Brownian noise term (i.e., the formal derivative of a Brownian motion; Gardiner, 1985) that introduces random, unbiased fluctuations in the learning process at each age a, with no temporal correlation. Each realization of a Brownian noise term $\eta(a)$ over $a \in [0, 1]$ defines a corresponding realization of knowledge acquisition trajectory $k_\bullet(a)$ (e.g., fig. 1c).

Equation (2) says that the focal offspring first learns from its parent. Vertical learning occurs over a duration of length $v_\bullet$, during which the focal offspring acquires knowledge instantaneously at a rate proportional to a stochastic vertical transmission rate, $\lambda_\bullet \left(\beta_{\mathrm{v}} + \sigma_{\mathrm{v}}\, \eta(a)\right)$, which increases with the net investment in learning $\lambda_\bullet$. The term $\sigma_{\mathrm{v}}\, \eta(a)$ describes the impact of stochastic events during vertical learning at age a, where the parameter σ_{v} controls the magnitude of the impact of learning stochasticity. When $\sigma_{\mathrm{v}} = 0$, learning occurs with no stochastic fluctuations. We assume that σ_{v} is sufficiently small relative to β_{v} such that $\beta_{\mathrm{v}} + \sigma_{\mathrm{v}}\, \eta(a)$ is almost always positive, ensuring that the focal offspring cannot lose knowledge during vertical learning. The knowledge acquisition rate is assumed to be proportional to the difference $(1 - \epsilon)\, k_{\mathrm{p}\bullet} - k_\bullet(a)$, between the knowledge currently available $(1 - \epsilon)\, k_{\mathrm{p}\bullet}$ from the parent and the focal offspring's knowledge $k_\bullet(a)$, where $\epsilon \in [0, 1]$ is the proportion of knowledge that becomes obsolete between two generations. Indeed, as the amount of available knowledge decreases, the focal offspring is less likely, on average, to encounter new information during interactions with its parent, thereby slowing the learning process.

Secondly, the focal offspring learns obliquely from a random adult of the previous generation, whose knowledge is denoted by $k_{\mathrm{a}\bullet}$, for a duration of length $o_\bullet$. Similar to the parent's knowledge, $k_{\mathrm{a}\bullet}$ results from a realization of the stochastic learning process undergone by the oblique exemplar in the previous generation (see details below). During oblique learning, the instantaneous knowledge acquisition rate is proportional to the product of a stochastic oblique transmission rate, $\lambda_\bullet \left(\beta_{\mathrm{o}} + \sigma_{\mathrm{o}}\, \eta(a)\right)$ (with σ_{o} small enough to ensure this remains almost always

positive), and of the amount of knowledge held by the oblique exemplar that the focal offspring has yet acquired, which is assumed to be given by $(1 - \epsilon)\, k_{\mathrm{a}\bullet} - \rho\, k_\bullet(v_\bullet) - \big(k_\bullet(a) - k_\bullet(v_\bullet)\big)$. This expression accounts for both the partial overlap between the parent's and the exemplar's knowledge, and the knowledge the offspring has already acquired during the oblique learning phase. We now explain the assumptions underlying this decomposition. We assume that, on average, the knowledge held by two adults in the population overlaps by a proportion $\rho \in [0,1]$. Consequently, the knowledge of the parent and of the oblique exemplar also overlaps by the same proportion ρ. At the end of the vertical learning phase (i.e., at age $v_\bullet$), the focal offspring has acquired from its parent a quantity of knowledge $k_\bullet(v_\bullet)$, of which a quantity $\rho\, k_\bullet(v_\bullet)$ is also known by the oblique exemplar. The total amount of knowledge that the focal offspring can potentially acquire from the oblique exemplar is therefore $(1 - \epsilon)\, k_{\mathrm{a}\bullet} - \rho\, k_\bullet(v_\bullet)$. At age $a \geq v_\bullet$, the focal offspring has already acquired from the oblique exemplar a quantity of knowledge $k_\bullet(a) - k_\bullet(v_\bullet)$, which is precisely the amount gained beyond what was acquired from the parent. Hence, the remaining knowledge available from the oblique exemplar at age $a \geq v_\bullet$ is given by $(1 - \epsilon)\, k_{\mathrm{a}\bullet} - \rho\, k_\bullet(v_\bullet) - \big(k_\bullet(a) - k_\bullet(v_\bullet)\big)$.

Thirdly, the focal offspring learns individually at an instantaneous stochastic rate proportional to $\lambda_\bullet\, \big(\alpha + \sigma_{\mathrm{i}}\, \eta(a)\big)$ for the remaining $1 - v_\bullet - o_\bullet$ time. Importantly, during each learning phase, the rate of knowledge acquisition is proportional to the offspring's net investment in learning $\lambda_\bullet$. As a result, when there is no such investment (i.e., $\lambda_\bullet = 0$), the focal offspring does not acquire any knowledge at any age (i.e., $\forall a \in [0,1]$, $k_\bullet(a) = 0$). Unlike vertical and oblique learning, the term $\alpha + \sigma_{\mathrm{i}}\, \eta(a)$ can become negative during individual learning, implying that the offspring may lose knowledge. This could reflect situations where exploration leads to confusion, or replacement of previously held information by incorrect alternatives.

It is important to note that both the knowledge of the parent $k_{\mathrm{p}\bullet}$ and that of the oblique exemplar $k_{\mathrm{a}\bullet}$ of the focal offspring result from realizations of the stochastic learning process experienced by those individuals in the previous generation. Specifically, their knowledge is obtained as an instantiation of $k_\bullet(1)$, where $k_\bullet(a)$ denotes a realization of the stochastic differential equation defined in eq. (2), with the values of $k_{\mathrm{p}\bullet}$ and $k_{\mathrm{a}\bullet}$ on the right-hand side corresponding to the knowledge values of the parent's own parent and oblique exemplar, and, in the case of the focal's oblique exemplar, to those of its own learning exemplars. Since both reproduction and survival are affected by knowledge (eq. 1 and eq. 3), the realized parental knowledge $k_{\mathrm{p}\bullet}$ and oblique exemplar knowledge $k_{\mathrm{a}\bullet}$ among the offspring are not solely determined by the stochastic learning process defined in eq. (2), but also by reproduction and survival, and thus by offspring

survival, which we next specify.

2.2.3 Offspring survival

After completing their learning, the offspring enter a density-dependent survival stage. Let $k_{\mathrm{o}\bullet} = k_\bullet(1)$ denote the knowledge acquired by the focal offspring at the end of the learning phase, where $k_\bullet(1)$ is the outcome of the stochastic differential equation (2). The probability of survival of the focal offspring is assumed to be given by

$$s(k_{\mathrm{o}\bullet}, n_{\mathrm{o}}) = \frac{\tilde{s}(k_{\mathrm{o}\bullet})}{1 + \gamma\, n_{\mathrm{o}}}, \tag{3}$$

where n_{o} is the total number of offspring produced by all adults and $\gamma > 0$ is a parameter controlling the strength of density dependence. The numerator $\tilde{s}(k_{\mathrm{o}\bullet})$, which modulates survival probability, is assumed to be given by

$$\tilde{s}(k_{\mathrm{o}\bullet}) = s_0 + \eta_{\mathrm{s}}\, k_{\mathrm{o}\bullet}, \tag{4}$$

where s_0 is the baseline survival, and η_{s} is the conversion factor that translates knowledge into survival. Parameter values are chosen in our analyses such that $s(k_{\mathrm{o}\bullet}, n_{\mathrm{o}})$ remains between 0 and 1 for all individuals.

2.3 Analyses

Here, we detail the hypotheses and the method we employ to investigate the joint cultural and evolutionary dynamics. Assuming a small mutation rate and a large population size, evolutionary change proceeds more slowly than cultural dynamics. This timescale separation allows us to analyze cultural dynamics while treating population trait values as constant and consider the knowledge dynamics within lineages of individuals bearing the same trait values (Mullon and Lehmann, 2017).

2.3.1 Cultural dynamics

We first aim to describe the equilibrium probability density of knowledge for a member of an $\boldsymbol{x}_\bullet$-lineage, which is shaped both by the stochastic learning process defined in eq. (2), which enables both the production and intergenerational accumulation of knowledge, and by the effects

of knowledge on fecundity and survival given in eqs. (1) and (3), since individuals who survive and reproduce are more likely to transmit their knowledge. In general, the learning process is too complicated to obtain an explicit expression for the probability density of knowledge. To make the analysis tractable, we assume that the outcomes of social learning are deterministic (i.e., we set $\sigma_{\mathrm{v}} = \sigma_{\mathrm{o}} = 0$; a condition that will be relaxed in supplementary analyses using individual-based simulation), in contrast to individual learning, where we assume that producing knowledge is inherently stochastic. Under this assumption, we can solve the stochastic differential equation (2) (see Appendix A.2). Using this solution, along with the effects of knowledge on fecundity and survival given in eqs. (1) and (3), we derive a recursion for the expected knowledge and its variance for a member of the $\boldsymbol{x}_{\bullet}$-lineage (see Appendices A.3.1 and A.3.2). However, this recursion ultimately depends on the entire hierarchy of moments of the probability density of knowledge. To address this issue and be able to track the expected knowledge and its variance across generations, we employ a Gaussian closure approximation (see Appendix A.3.3). Note that the stationary distributions obtained from individual-based simulations suggest that a Gaussian approximation provides a good fit for the probability density of knowledge (see fig. S.1). With this, we can then fully characterize the cultural equilibrium in terms of the expected knowledge and its variance (see Appendix A.4).

2.3.2 Evolutionary dynamics

Under the assumptions of small mutation rate, small variance in knowledge, and a large population, the expected evolutionary dynamics can be inferred from the expected lineage fitness in a focal $\boldsymbol{x}_{\bullet}$-lineage, i.e., the expected fitness of a random individual in that lineage (Mullon and Lehmann, 2017). We show in Appendix B.1, that the expected lineage fitness in the $\boldsymbol{x}_{\bullet}$-lineage in a population with mean traits $\overline{\boldsymbol{x}}$ can be expressed as

$$W(\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}}) = \frac{f\big(\boldsymbol{x}_{\bullet}, \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\big)\, \tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big)}{f\big(\overline{\boldsymbol{x}}, \overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})\big)\, \tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\overline{\boldsymbol{x}})\big)}, \tag{5}$$

where $\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})$ and $\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})$ are the expected knowledge at cultural equilibrium for a random adult and a random offspring after learning is complete of the $\boldsymbol{x}_{\bullet}$-lineage.

The expected evolutionary dynamics can then be inferred from the selection gradient, defined as

$$\boldsymbol{s}(\overline{\boldsymbol{x}}) = \nabla W(\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})|_{\boldsymbol{x}_{\bullet} = \overline{\boldsymbol{x}}}, \tag{6}$$

where the operator ∇ acts such that, for any function of traits g, we have

$$\nabla g(\boldsymbol{x}_\bullet) = \begin{pmatrix} \frac{\partial g(\boldsymbol{x}_\bullet)}{\partial v_\bullet} \\ \frac{\partial g(\boldsymbol{x})}{\partial o_\bullet} \\ \frac{\partial g(\boldsymbol{x})}{\partial \lambda_\bullet} \end{pmatrix}. \tag{7}$$

Each entry of the selection gradient indicates whether natural selection favors an increase or a decrease in the corresponding trait. According to invasion analysis (Leimar, 2009), the mean trait values $\overline{\boldsymbol{x}}$ will eventually converge to a convergence stable trait vector denoted by $\overline{\boldsymbol{x}}^*$, which either satisfies $\boldsymbol{s}(\overline{\boldsymbol{x}}^*) = 0$ or lies on the boundary of the phenotypic space. We use the expression of the selection gradient to highlight the distinct selective pressures acting on each learning trait and to numerically estimate the trait values $\overline{\boldsymbol{x}}^*$ favored by selection starting from a population initially lacking any form of learning, that is, with ancestral traits $\overline{v} = \overline{o} = \overline{\lambda} = 0$ (see details in B.3).

To assess the robustness of our forthcoming analytical findings, we perform individual-based simulations that relax the previously stated assumptions (see details of individual-based simulations in Appendix D). In particular, we allow for small population sizes, allow $\sigma_{\rm v} > 0$ and $\sigma_{\rm o} > 0$, and no longer assume that the probability density of knowledge is normal.

3 Results

3.1 Cultural dynamics

3.1.1 Cultural selection enhances cumulative knowledge

We now characterize the expected value and variance of knowledge for a member of the $\boldsymbol{x}_\bullet$-lineage at cultural equilibrium, in order to reveal the effects of stochasticity in learning on cumulative knowledge. To achieve this, we first determine the outcome of the learning process. We show in Appendix A.2 that in the absence of stochasticity during social learning (i.e., when $\sigma_{\rm v} = \sigma_{\rm o} = 0$) the realized knowledge $k_{\rm o\bullet} = k_\bullet(1)$ acquired by a focal offspring with traits $\boldsymbol{x}_\bullet$, whose parent has knowledge $k_{\rm p\bullet}$ and who chooses an oblique exemplar with knowledge $k_{\rm a\bullet}$ is

$$k_{\rm o\bullet} = \overbrace{\omega_{\rm v}(\boldsymbol{x}_\bullet)\,(1-\epsilon)\,k_{\rm p\bullet}}^{\text{vertical learning}} + \overbrace{\omega_{\rm o}(\boldsymbol{x}_\bullet)\,[(1-\epsilon)\,k_{\rm a\bullet} - \rho\,\omega_{\rm v}(\boldsymbol{x}_\bullet)\,(1-\epsilon)\,k_{\rm p\bullet}]}^{\text{oblique learning}} + \overbrace{\lambda_\bullet\,\alpha\,(1-v_\bullet - o_\bullet) + \chi_{\rm k}}^{\text{individual learning}}, \tag{8}$$

where

$$\omega_{\mathrm{v}}(\boldsymbol{x_\bullet}) = 1 - \mathrm{e}^{-\lambda_\bullet \beta_{\mathrm{v}} v_\bullet} \quad \text{and} \quad \omega_{\mathrm{o}}(\boldsymbol{x_\bullet}) = 1 - \mathrm{e}^{-\lambda_\bullet \beta_{\mathrm{o}} o_\bullet}, \tag{9}$$

are the proportion of available knowledge at the start of the vertical and oblique learning phases, respectively, that is transmitted to the focal offspring. Since the offspring already acquired an amount $\omega_{\mathrm{v}}(\boldsymbol{x_\bullet})\,(1-\epsilon)\,k_{\mathrm{p}\bullet}$ of knowledge from its parents through vertical learning, and a fraction ρ of this overlaps with the oblique exemplar's knowledge, the amount of non-redundant knowledge available at the start of oblique learning is $(1-\epsilon)\,k_{\mathrm{a}\bullet} - \rho\,\omega_{\mathrm{v}}(\boldsymbol{x_\bullet})\,(1-\epsilon)\,k_{\mathrm{p}\bullet}$. Multiplying this non-redundant knowledge by $\omega_{\mathrm{o}}(\boldsymbol{x_\bullet})$ yields the knowledge acquired through oblique learning by the focal offspring. Finally, $\lambda_\bullet\,\alpha\,(1 - v_\bullet - o_\bullet) + \chi_{\mathrm{k}}$ is the realized knowledge acquired through individual learning, where χ_{k} is a realization of a Gaussian random variable with mean 0 and variance $\lambda_\bullet^2\,\sigma_{\mathrm{i}}^2\,(1 - v_\bullet - o_\bullet)$.

Let us now consider the offspring cohort produced by the adults of generation t in the $\boldsymbol{x_\bullet}$-lineage and denote by $\overline{k}_{\mathrm{a},t}(\boldsymbol{x_\bullet})$ and $V_{\mathrm{a},t}(\boldsymbol{x_\bullet})$ the mean and variance of the probability density of knowledge among these adults. By taking the expectation in eq. (8) with respect to all possible values of parental knowledge $k_{\mathrm{p}\bullet}$, oblique exemplar knowledge $k_{\mathrm{a}\bullet}$, and the stochastic outcomes of learning χ_{k}, we show in Appendix A.3.1 that the expected knowledge $\overline{k}_{\mathrm{o},t}(\boldsymbol{x_\bullet})$ of an offspring born to an adult of generation t in the $\boldsymbol{x_\bullet}$-lineage is given by

$$\overline{k}_{\mathrm{o},t}(\boldsymbol{x_\bullet}) = p_{\mathrm{il}}(\boldsymbol{x_\bullet}) + h_{\mathrm{ol}}(\boldsymbol{x_\bullet})\,\overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}}) + h_{\mathrm{vl}}(\boldsymbol{x_\bullet})\left(\overline{k}_{\mathrm{a},t}(\boldsymbol{x_\bullet}) + \frac{V_{\mathrm{a},t}(\boldsymbol{x_\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a},t}(\boldsymbol{x_\bullet})}\right), \tag{10}$$

where $p_{\mathrm{il}}(\boldsymbol{x_\bullet}) = \lambda_\bullet\,\alpha\,(1 - v_\bullet - o_\bullet)$, is the expected amount of knowledge produced through individual learning by the focal offspring, and

$$h_{\mathrm{ol}}(\boldsymbol{x_\bullet}) = (1 - \epsilon)\,\omega_{\mathrm{o}}(\boldsymbol{x_\bullet}), \tag{11}$$

and

$$h_{\mathrm{vl}}(\boldsymbol{x_\bullet}) = (1 - \epsilon)\,\omega_{\mathrm{v}}(\boldsymbol{x_\bullet})\,(1 - \rho\,\omega_{\mathrm{o}}(\boldsymbol{x_\bullet})). \tag{12}$$

are the oblique and vertical cultural heritabilities, respectively. The oblique and vertical cultural heritabilities, both lying between 0 and 1, are the proportion of knowledge from the oblique and vertical exemplar that is effectively transmitted to a learner with traits $\boldsymbol{x_\bullet}$.

The last two terms on the right-hand side of eq. (10) are the expected knowledge acquired from an oblique exemplar and from the parent. These terms depend on the corresponding cultural heritabilities, multiplied by the expected knowledge $\overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})$ of an oblique exemplar and

that $\overline{k}_{a,t}(\boldsymbol{x}_\bullet) + V_{a,t}(\boldsymbol{x}_\bullet)\,\eta_f / (f_0 + \eta_f\,\overline{k}_{a,t}(\boldsymbol{x}_\bullet))$ of a vertical exemplar. The expected knowledge of a vertical exemplar exceeds that of an average adult in the lineage. This occurs because individuals with greater knowledge are overrepresented among vertical exemplars, as they produce more offspring.

The term $h_{vl}(\boldsymbol{x}_\bullet)\,V_{a,t}(\boldsymbol{x}_\bullet)\,\eta_f / (f_0 + \eta_f\,\overline{k}_{a,t}(\boldsymbol{x}_\bullet))$ in (10) can be thought of as the response of expected knowledge to cultural selection (change in the expected knowledge in the offspring generation resulting from selection on knowledge in the parental generation, in analogy with the response to selection due to genetic inheritance; Lynch and Walsh, 1998), due to differences in fecundity. It arises from differences in vertical transmission caused by variation in fecundity: individuals who, by chance, have more knowledge tend to produce more offspring and, as a result, are more likely to transmit their knowledge via vertical transmission, leading to an increase in the expected knowledge within the lineage. The response to cultural selection is particularly pronounced when parents transmit a substantial portion of their knowledge to their offspring (i.e., $h_{vl}(\boldsymbol{x}_\bullet)$ is great), there is significant knowledge difference between adults (i.e., knowledge variance $V_{a,t}(\boldsymbol{x}_\bullet)$ is great), and when knowledge has a strong impact on fecundity (i.e., η_f is great). The strength of the response to cultural selection diminishes as $f_0 + \eta_f\,\overline{k}_{a,t}(\boldsymbol{x}_\bullet)$ increases, since relative differences in fecundity within the lineage become less pronounced.

After completing their learning, offspring go through the survival stage to reach adulthood in generation $t + 1$. We show in Appendix A.3.1 that the expected knowledge $\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet)$ of an adult of generation $t + 1$ in the $\boldsymbol{x}_\bullet$-lineage is given by

$$\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet) = \overline{k}_{o,t}(\boldsymbol{x}_\bullet) + \frac{V_{o,t}(\boldsymbol{x}_\bullet)\,\eta_s}{\tilde{s}(\overline{k}_{o,t}(\boldsymbol{x}_\bullet))}, \tag{13}$$

where $V_{o,t}(\boldsymbol{x}_\bullet)$ is the variance in the knowledge held by a randomly chosen offspring born to an adult of generation t in the $\boldsymbol{x}_\bullet$-lineage. Since offspring with above-average knowledge are more likely to survive, the expected knowledge among adults is higher than that among offspring (i.e., $\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet) > \overline{k}_{o,t}(\boldsymbol{x}_\bullet)$). This effect is captured by the selection differential $V_{o,t}(\boldsymbol{x}_\bullet)\,\eta_s / \tilde{s}(\overline{k}_{o,t}(\boldsymbol{x}_\bullet))$, which is the difference between mean knowledge of surviving and all offspring within the lineage. This selection differential is particularly pronounced when there is a significant knowledge difference between offspring (i.e., $V_{o,t}(\boldsymbol{x}_\bullet)$ is great) and a strong survival advantage conferred by knowledge (i.e., η_s is great). The selection differential diminishes with increasing values of $\tilde{s}(\overline{k}_{o,t}(\boldsymbol{x}_\bullet))$, as relative differences in survival within the lineage become less pronounced. The response of an individual's expected knowledge to cultural selection due to variation in survival depends on how knowledge is transmitted to the new offspring cohort. It is

given by $h_{\mathrm{ol}}(\boldsymbol{x_\bullet})\, V_{\mathrm{o},t}(\overline{\boldsymbol{x}})\, \eta_{\mathrm{s}}/\tilde{s}\big(\overline{k}_{\mathrm{o},t}(\overline{\boldsymbol{x}})\big) + h_{\mathrm{vl}}(\boldsymbol{x_\bullet})\, V_{\mathrm{o},t}(\boldsymbol{x_\bullet})\, \eta_{\mathrm{s}}/\tilde{s}\big(\overline{k}_{\mathrm{o},t}(\boldsymbol{x_\bullet})\big)$ (obtained by substituting eq. (13) with $\boldsymbol{x_\bullet} = \boldsymbol{x_\bullet}$ and $\boldsymbol{x_\bullet} = \overline{\boldsymbol{x}}$ into eq. (10) with $t = t+1$ and identifying the terms corresponding to cultural selection).

3.1.2 Learning stochasticity enhances knowledge by increasing knowledge variance

Equations (10) and (13) highlight the role of cultural selection on mean knowledge. Taken together, these equations allow us to characterize the expected knowledge of an adult of the $\boldsymbol{x_\bullet}$-lineage at equilibrium. We show in Appendix A.4.1 that the expected knowledge of the focal adult then stabilizes at a value that satisfies

$$
\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x_\bullet}) = \frac{\overbrace{p_{\mathrm{il}}(\boldsymbol{x_\bullet}) + h_{\mathrm{ol}}(\boldsymbol{x_\bullet})\, \overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})}^{\substack{\text{knowledge aquired through individual} \\ \text{learning and from an oblique exemplar}}} + \overbrace{\dfrac{h_{\mathrm{vl}}(\boldsymbol{x_\bullet})\, V_{\mathrm{a}}^{*}(\boldsymbol{x_\bullet})\, \eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x_\bullet})} + \dfrac{V_{\mathrm{o}}^{*}(\boldsymbol{x_\bullet})\, \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x_\bullet})\big)}}^{\text{cultural selection}}}{\underbrace{1 - h_{\mathrm{vl}}(\boldsymbol{x_\bullet})}_{\substack{\text{knowledge accumulation} \\ \text{within the lineage due} \\ \text{to vertical transmission}}}}, \tag{14}
$$

where the superscript * indicates that the quantities are evaluated at equilibrium (see also fig. 1d). The first two terms in the numerator of eq. (14) are the expected amount of knowledge acquired through individual and from an oblique exemplar. The remaining terms in the numerator are the increase in expected knowledge driven by cultural selection resulting from differences in fecundity and survival within the lineage due to variation in knowledge. The denominator of eq. (14) captures the inter-generational accumulation of knowledge within the lineage due to vertical transmission. This can be seen by expressing $1/\big(1 - h_{\mathrm{vl}}(\boldsymbol{x_\bullet})\big)$ as $1 + \sum_{t=1}^{\infty} h_{\mathrm{vl}}(\boldsymbol{x_\bullet})^{t}$, where $h_{\mathrm{vl}}(\boldsymbol{x_\bullet})^{t}$ is the proportion of knowledge acquired by an ancestor t generations ago that is effectively transmitted to the focal individual.

We observe that the mean knowledge of offspring $\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x_\bullet})$ is generally close to that of adults $\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x_\bullet})$ (see the first line of eq. (A.53) and fig. S.2). However, because offspring with above-average knowledge are more likely to survive and become adults, the expected knowledge among adults is slightly higher than that among offspring (i.e., $\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x_\bullet}) > \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x_\bullet})$; see fig. S.2). In contrast, when knowledge does not affect offspring survival to adulthood (i.e., $\eta_{\mathrm{s}} = 0$), the expected equilibrium knowledge is the same for adults and offspring (i.e., $\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x_\bullet}) = \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x_\bullet})$; see fig. S.2).

Equation (14) reveals that cultural selection promotes knowledge accumulation. Since the impact of cultural selection increases with knowledge variance (see eq. (14)), we derive in Appendix

A.4.2 a condition that the equilibrium variance in the knowledge held by an adult from the $\boldsymbol{x_\bullet}$-lineage must satisfy, and that is

$$V_{\mathrm{a}}^*(\boldsymbol{x_\bullet}) = \frac{\overbrace{\lambda_\bullet^2\,\sigma_{\mathrm{i}}^2\,(1 - v_\bullet - o_\bullet)}^{\substack{\text{variance produced by}\\\text{stochasticity in learning}}} + \overbrace{h_{\mathrm{ol}}(\boldsymbol{x_\bullet})^2\,V_{\mathrm{a}}^*(\overline{\boldsymbol{x}})}^{\substack{\text{variance transmitted}\\\text{from an oblique exemplar}}} \overbrace{- \left(\frac{h_{\mathrm{vl}}(\boldsymbol{x_\bullet})\,V_{\mathrm{a}}^*(\boldsymbol{x_\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\boldsymbol{x_\bullet})}\right)^2 - \left(\frac{V_{\mathrm{o}}^*(\boldsymbol{x_\bullet})\,\eta_{\mathrm{s}}}{\tilde{s}\bigl(\overline{k}_{\mathrm{o}}^*(\boldsymbol{x_\bullet})\bigr)}\right)^2}^{\text{cultural selection reducing knowledge variance}}}{\underbrace{1 - h_{\mathrm{vl}}(\boldsymbol{x_\bullet})^2}_{\substack{\text{variance accumulation}\\\text{within the lineage due}\\\text{to vertical transmission}}}}.$$

$$(15)$$

The first term in the numerator of eq. (15) is the knowledge variance produced by stochasticity in individual learning. If learning were entirely deterministic (i.e., $\sigma_{\mathrm{i}} = 0$), no knowledge variance would be produced, and the only solution to eq. (15) would be $V_{\mathrm{a}}^*(\boldsymbol{x_\bullet}) = 0$ (see proof in Appendix A.4.3). The second term in the numerator accounts for variance transmitted from an oblique exemplar. The remaining terms in the numerator are the effect of cultural selection, which reduces knowledge variance. Because survival to adulthood reduces knowledge variance, the equilibrium variance in the knowledge held by an adult $V_{\mathrm{a}}^*(\boldsymbol{x_\bullet})$ is generally slightly lower than that of an offspring $V_{\mathrm{o}}^*(\boldsymbol{x_\bullet})$ (see fig. S.2). When knowledge has no effect on survival (i.e., $\eta_{\mathrm{s}} = 0$), the two variances are equal (i.e., $V_{\mathrm{a}}^*(\boldsymbol{x_\bullet}) = V_{\mathrm{o}}^*(\boldsymbol{x_\bullet})$; see fig. S.2). Finally, the denominator of eq. (15) describes the accumulation of knowledge variance within the $\boldsymbol{x_\bullet}$-lineage through vertical transmission.

Altogether eqs. (14) and (15) reveal how stochasticity in individual learning impacts knowledge equilibrium. Since stochasticity in individual learning produces knowledge variance (see eq. (15)) and the effect of cultural selection increases with knowledge variance (see eq. (14)), stochasticity in individual learning ultimately enhances the accumulation of knowledge. By numerically estimating $V_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ and $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ (see Appendix A.4.4 for details on the procedure) for different values of σ_{i}, we find that increased stochasticity in individual learning σ_{i} increase population knowledge variance $V_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$, which in turn lead to an increase in mean knowledge $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ (more than doubling; fig. 2a).

Remarkably, stochasticity in individual learning σ_{i}, can foster knowledge accumulation even in populations with low average knowledge-production ability (e.g., $\alpha = 0.001$; see fig. 2b). When stochasticity is particularly high, cultural selection becomes the dominant force driving knowledge acquisition. As a result, populations with different average knowledge-production abilities tend to carry similar levels of mean knowledge, whereas, in the absence of stochasticity, they exhibit marked differences in mean knowledge (see fig. 2b). Additionally, because knowledge

improves both fecundity and survival, greater stochasticity in individual learning σ_i is associated with larger population size (see fig. S.3). The effect of cultural selection in enhancing the mean knowledge $\overline{k}_a^*(\overline{\boldsymbol{x}})$ is stronger when knowledge has a stronger impact on fecundity and survival (i.e., when η_f and η_s are higher; see fig. S.4), and when vertical or oblique cultural heritability is higher (fig. S.5).

3.2 Evolutionary dynamics

3.2.1 Trade-off between fecundity and learning

With mean knowledge at equilibrium characterized, we now use the selection gradient to investigate the selection pressures shaping the evolution of the learning traits. By substituting eq. (5) into eq. (6), we obtain the direction of selection on the traits

$$
\boldsymbol{s}(\overline{\boldsymbol{x}}) = \overbrace{\frac{\nabla(1 - \lambda_\bullet)^\theta\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}}{(1 - \overline{\lambda})^\theta}}^{\text{learning traits effects on fecundity}} + \overbrace{\frac{\eta_f \; \nabla\overline{k}_a^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}}{f_0 + \eta_f \, \overline{k}_a^*(\overline{\boldsymbol{x}})} + \frac{\eta_s \; \nabla\overline{k}_o^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}})\big)}}^{\text{learning traits effects lineage members' expected knowledge}} \;, \quad (16)
$$

where the gradient operator ∇ is defined in eq. (7).

The first term in eq. (16) describes selection arising from the fecundity costs associated with investment in learning. The remaining terms describe the effect of selection resulting from the impact of learning traits on the expected knowledge of members of a lineage. The strength of these selection pressures diminishes with increasing values of $f_0 + \eta_f \, \overline{k}_a^*(\overline{\boldsymbol{x}})$ and $\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}})\big)$, as relative differences in fecundity and survival among lineages driven by differences in knowledge become less pronounced.

Overall, eq. (16) indicates that the evolution of learning traits, and consequently the trait values $\overline{\boldsymbol{x}}^* = (\overline{v}^*, \overline{o}^*, \overline{\lambda}^*)$ favored by selection in the long-term, which can be either interior equilibria satisfying $\boldsymbol{s}(\overline{\boldsymbol{x}}^*) = (0, 0, 0)$ or values lying on the boundary of the phenotypic space, result from a trade-off between the fecundity costs incurred and the knowledge gained. We focus on evaluating the trait values favored by selection starting from a population initially expressing trait $\overline{\boldsymbol{x}} = (0, 0, 0)$ (see Appendix B.3 for details on the procedure). Numerical estimates of the average learning traits at $\overline{\boldsymbol{x}}^*$ reveal that the fecundity costs significantly influence the evolution of the learning traits: when the fecundity costs associated with learning investment are higher (i.e., θ is higher), individuals invest less in learning (i.e., $\overline{\lambda}^*$ is lower; fig. S.6a). This rapidly

limits the evolution of both types of social learning (i.e., $\overline{v}^* = \overline{o}^* = 0$; fig. S.6a). Indeed, when the fecundity cost exponent θ is higher, individuals allocate more resources to reproduction and less to learning at $\overline{\boldsymbol{x}}^*$ (i.e., $\overline{\lambda}^*$ is lower; fig. S.6a), which restricts knowledge production and consequently reduces the benefits of social learning. In the following numerical analyses, we set $\theta = 0.1$, a value that permits the evolution of social learning.

3.2.2 Trade-off between different types of learning

To better understand the selection pressures on the learning traits, we decompose their effect on the expected knowledge of adults in the $\boldsymbol{x}_\bullet$-lineage. We show in Appendix B.2, that $\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ satisfies

$$
\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\Big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}} = \frac{1}{1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}})} \times
$$

$$
\left[\overbrace{\nabla p_{\mathrm{il}}(\boldsymbol{x}_\bullet) + \nabla h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)\, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}) + \nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet) \left(\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}) + \frac{V_{\mathrm{a}}^*(\overline{\boldsymbol{x}})\, \eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})} \right)}^{\text{learning traits effects on learning}} \right.
$$

$$
\left. + \underbrace{h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\, \nabla \frac{V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\, \eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)} + \nabla \frac{V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\, \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)}}_{\text{learning traits effects on cultural selection}} \right]\Bigg|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}. \quad (17)
$$

(Appendix B.2). The terms on the second line describe the impact of learning traits on knowledge acquisition. This effect depends on how traits v, o, and λ influence the knowledge produced individually, as well as that acquired from oblique and vertical exemplars. The marginal knowledge gained from oblique and vertical exemplars depends on the effect of the learning traits on the oblique $\nabla h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ and vertical $\nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ cultural heritability, multiplied by the expected knowledge of a parent and an oblique exemplar. The marginal knowledge gained through individual learning is given by $\nabla p_{\mathrm{il}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$. Since the traits v and o enhance the time allocated to vertical and oblique learning, but reduce the time available for individual knowledge production, their evolution is shaped by trade-offs between different types of learning. The terms on the third line describe the impact of learning traits on cultural selection acting on the expected adult knowledge in the lineage. Learning traits can impact cultural selection by impacting the variance in potential knowledge, as well as the expected fecundity and survival within the lineage.

The selection pressure scales as $1/\big(1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\big)$, which captures the inter-generational accumu-

lation of the effects of learning traits on knowledge acquisition and on cultural selection, due to vertical transmission. This can be seen by expressing this factor as $1 + \sum_{t=1}^{\infty} h_{\mathrm{vl}}(\overline{\boldsymbol{x}})^t$, where $h_{\mathrm{vl}}(\overline{\boldsymbol{x}})^t$ is the population mean proportion of knowledge initially acquired by an ancestor that is transmitted to a direct descendant t generations later. This formulation highlights that a change in knowledge in one generation, driven by a change in learning traits, influences the knowledge of all future descendants within the lineage. By capturing the cumulative effects of these interactions, which occur among relatives, the evolutionary dynamics thus incorporate multi-generational kin selection effects.

To understand selection on the learning traits, we also need to examine $\nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ (see eq. (16)). The condition satisfied by $\nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ is identical to eq. (17) with $\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}} = \nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ on the left-hand side, except for a factor $h_{\mathrm{vl}}(\overline{\boldsymbol{x}})$ multiplying the term capturing the effect of learning traits on cultural selection due to differences in survival (see eq. (B.23)). This difference arises because offspring have not yet undergone survival themselves. As a result, the effect of cultural selection on offspring knowledge due to within-lineage survival differences comes only from the knowledge of parents who survived and who transmit an average proportion $h_{\mathrm{vl}}(\overline{\boldsymbol{x}})$ of that knowledge to their offspring.

Altogether eqs. (16) and (17) reveals that the evolution of learning traits, and therefore the trait values $\overline{\boldsymbol{x}}^* = (\overline{v}^*, \overline{o}^*, \overline{\lambda}^*)$ favored by selection, are influenced by the trade-off between knowledge acquisition from different learning types. Numerically estimating $\overline{\boldsymbol{x}}^*$ reveals that, when knowledge remains relevant over time (i.e., low ϵ; e.g., in a stable environment), individuals allocate significant time to both vertical and oblique learning (i.e., high $\overline{v}^*$ and $\overline{o}^*$; see fig. S.6b) because they can acquire substantial knowledge socially. Since social learning becomes more efficient in stable environments, individuals tend to invest significant resources into learning (i.e., high $\overline{\lambda}^*$; see fig. S.6b). The trade-off between vertical and oblique learning depends on the overlap in knowledge between two adults ρ. Since we assume it is easier to learn from parents (i.e., $\beta_{\mathrm{v}} > \beta_{\mathrm{o}}$), individuals tend to skip the oblique learning phase at $\overline{\boldsymbol{x}}^*$ (i.e., $\overline{o}^* = 0$) when there is a high knowledge overlap between adults (i.e., high ρ; see fig. S.6c). In such cases, oblique exemplars provide similar knowledge to parents, but it is harder to learn from them. Conversely, when there is a low knowledge overlap between adults (i.e., low ρ), at $\overline{\boldsymbol{x}}^*$ individuals allocate time to oblique learning after learning vertically (i.e., $\overline{o}^* > 0$), enabling them to acquire knowledge not available from their parents (see fig. S.6c). This outcome is consistent with the results of Maisonneuve et al. (2025) (see AppendixB.4 for details). Although it strongly influences the learning schedule, the overlap in knowledge between two adults ρ has little effect on how

resources are allocated between learning and fecundity at the reached evolutionary equilibrium (i.e., little effect on $\overline{\lambda}^*$; see fig. S.6c).

3.2.3 How learning stochasticity promotes social learning

We now investigate the impact of stochasticity in individual learning, quantified by the parameter σ_i, on the average learning traits at $\overline{\boldsymbol{x}}^*$. When σ_i is high, the knowledge variance $V_a^*(\overline{\boldsymbol{x}})$ in adults and $V_o^*(\overline{\boldsymbol{x}})$ in offspring increases (see fig. 2a and eqs. (15) and (A.51)), boosting cultural selection and leading to a rise in expected knowledge of the exemplar individual during vertical $\overline{k}_a^*(\overline{\boldsymbol{x}}) + V_a^*(\boldsymbol{x}_\bullet)\,\eta_f / \big(f_0 + \eta_f\,\overline{k}_a^*(\overline{\boldsymbol{x}})\big)$ and oblique $\overline{k}_a^*(\overline{\boldsymbol{x}})$ learning (as $\overline{k}_a^*(\overline{\boldsymbol{x}})$ increases with $V_a^*(\overline{\boldsymbol{x}})$ and $V_o^*(\overline{\boldsymbol{x}})$; see eq. (14)). Consequently, with higher stochasticity in individual learning, individuals are expected to rely more on social learning, as offspring can acquire substantial knowledge from exemplars, who, under more stochastic learning conditions, tend to hold more knowledge.

This prediction is supported by numerical estimates of $\overline{\boldsymbol{x}}^* = (\overline{v}^*, \overline{o}^*, \overline{\lambda}^*)$, which show that as σ_i increases, individuals tend to engage more in both vertical and oblique learning (i.e., $\overline{v}^*$ and $\overline{o}^*$ increase; see fig. 3a). This is because, as anticipated, high σ_i allows the population to accumulate substantial knowledge (i.e., high $\overline{k}_a^*(\overline{\boldsymbol{x}}^*)$; see fig. 3g), resulting in cultural exemplars possessing greater knowledge. Under high σ_i, the increased efficiency of social learning then drives individuals to invest more resources in learning at $\overline{\boldsymbol{x}}^*$ (i.e., high $\overline{\lambda}^*$; see fig. 3d). This effect is especially pronounced in populations with low knowledge-production ability: stochasticity in learning can drive the evolution of great investment in learning (e.g., $\overline{\lambda}^* \approx 0.95$ when $\alpha = 0.025$ and $\sigma_i = 0.25$; see fig. S.7), whereas in its absence, investment in learning would be low (e.g., $\overline{\lambda}^* \approx 0.2$ when $\alpha = 0.025$ and $\sigma_i = 0$; see fig. S.7). However, in populations with very low knowledge-production ability, individuals evolve to allocate all resources to fecundity and none to learning, so greater stochasticity in learning has no effect (e.g., $\alpha = 0.001$; see fig. S.7). The impact of σ_i on learning traits at $\overline{\boldsymbol{x}}^*$ also weakens when the obsolescence rate is high, as rapid knowledge decay across generations limits the accumulation of knowledge (e.g., $\epsilon = 0.25$; fig. S.6d). Although higher σ_i generally leads to investing more in learning at the expense of fecundity (i.e., higher $\overline{\lambda}^*$), the resulting enhancement in population mean knowledge still leads to larger population sizes at $\overline{\boldsymbol{x}}^*$ (see fig. 3g).

Individual-based simulations, which notably relax the assumption of a Gaussian probability density of knowledge, confirm that greater stochasticity in individual learning promotes both

increased time allocated to social learning and greater investment in learning at $\overline{\boldsymbol{x}}^*$ and leads to higher population mean knowledge and population size (see fig. S.9). In addition, simulations that relax the assumption of non-stochastic social learning (i.e., allowing $\sigma_{\mathrm{v}} > 0$ or $\sigma_{\mathrm{o}} > 0$) reveal similar patterns under increased stochasticity in vertical and oblique learning (see fig. S.10).

However, when adults knowledge significantly overlaps, increase in σ_{i} results in a greater allocation of time to vertical learning and a reduced allocation to oblique learning at $\overline{\boldsymbol{x}}^*$ (e.g., increase in $\overline{v}^*$ but not in $\overline{o}^*$ when $\rho = 0.4$; see fig. S.6e). This occurs because offspring can acquire many similar knowledge units from either their parents or other adults, leading individuals to favor the most effective social learning mode for acquiring shared knowledge. Under higher stochasticity in individual learning, the average knowledge of parents $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}) + V_{\mathrm{a}}^*(\overline{\boldsymbol{x}})\,\eta_{\mathrm{f}} / (f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}))$ increases more than that of oblique exemplars $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ (see fig. S.8). This is because, within the adult population with varying levels of knowledge, parents tend to have higher fertility and, consequently, above-average knowledge. As a result, when learning is more stochastic, offspring acquire more of the shared knowledge from their parents.

3.2.4   Selection promotes learning from selected individuals

The above results show that stochasticity in learning promotes knowledge accumulation and increases the benefits of social learning. We now turn to the question of from whom individuals should learn when knowledge varies across the population. Specifically, we investigate which social learning exemplars are favored by natural selection when knowledge improves survival and/or fecundity.

When knowledge enhances fecundity but does not affect survival (i.e., when $\eta_{\mathrm{f}} > 0$ and $\eta_{\mathrm{s}} = 0$), increased stochasticity in individual learning (i.e., increased σ_{i}) leads to allocating more time to vertical learning rather than oblique learning at $\overline{\boldsymbol{x}}^*$ (i.e., a rise in $\overline{v}^*$ but not in $\overline{o}^*$; see fig. 3b). In that case, because stochasticity in individual learning leads to uncertainty in the knowledge held by social learning exemplars, being a parent is a cue of high fecundity and, therefore, greater knowledge. By contrast, when knowledge increases survival but has no effect on fecundity (i.e., when $\eta_{\mathrm{f}} = 0$ and $\eta_{\mathrm{s}} > 0$), increased stochasticity in individual learning (i.e., increased σ_{i}) leads to allocating more time to both vertical and oblique learning at $\overline{\boldsymbol{x}}^*$ (i.e., a rise in $\overline{v}^*$ and $\overline{o}^*$; see fig. 3c). It follows that because adults who are available as social learning exemplars, whether during vertical or oblique learning, have survived until adulthood, their survival is a cue of great knowledge. In summary, when stochasticity in learning renders the knowledge of social learning

exemplars unpredictable, natural selection favors learning from individuals who exhibit cues of possessing great knowledge (e.g., being a parent or having survived to adulthood).

3.2.5 Selection promotes stochasticity in learning

The results so far show that stochasticity in individual learning, quantified by σ_i, markedly affects both the cultural and evolutionary dynamics, and in perhaps counterintuitive ways. Yet, these results may not have traction if σ_i itself is selected away. We now relax the assumption of fixed σ_i by introducing an additional evolving trait $s \in [0, +\infty)$ that increases stochasticity (i.e., $\forall s \in [0, +\infty)$, $\sigma_i'(s) > 0$) and show that this will be favored by selection. For this, we assume that learning is inherently stochastic, such that even in the absence of any trait promoting stochasticity (i.e., when $s = 0$), a small baseline level of stochasticity remains (i.e., $\sigma_i(0) > 0$, but small). Each individual is now characterized by a vector of traits $\boldsymbol{x_\bullet} = (v_\bullet, o_\bullet, \lambda_\bullet, s_\bullet)$. We study how selection may favor the emergence of the trait s that increases stochasticity in learning, starting from a population with mean trait vector $\overline{\boldsymbol{x}}_0 = (\overline{v}, \overline{o}, \overline{\lambda}, 0)$. Using eq. (16), we show in Appendix C that the trait s emerge if

$$\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\; \frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\bigg|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0} + \frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\; \frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\bigg|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0} > 0, \tag{18}$$

where

$$\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\bigg|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0}$$
$$\approx \frac{1}{1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)}\left[\frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\; \frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\bigg|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0} + \frac{\eta_{\mathrm{s}}}{s_0 + \eta_{\mathrm{s}}\, \overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)}\; \frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\bigg|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0}\right]. \tag{19}$$

This latter expression shows that, in the $\boldsymbol{x_\bullet}$-lineage, $s_\bullet$ affects the expected adult knowledge by impacting knowledge variance in both adults and offspring, which in turn modulates the effect of cultural selection that increases the expected adult knowledge in the lineage. This effect is further amplified under strong vertical transmission (i.e., large $h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)$), as increases in expected knowledge driven by cultural selection accumulate across generations within the lineage. Since $s_\bullet$ increases knowledge variance in both adults and offspring in the lineage (i.e., $\partial V_{\mathrm{a}}^*(\boldsymbol{x_\bullet})/\partial s_\bullet|\, \boldsymbol{x_\bullet} = \overline{\boldsymbol{x}}_0 > 0$ and $\partial V_{\mathrm{o}}^*(\boldsymbol{x_\bullet})/\partial s_\bullet|\, \boldsymbol{x_\bullet} = \overline{\boldsymbol{x}}_0 > 0$; see Appendix C), it strengthens the effect of cultural selection and leads to higher expected adult knowledge (i.e., $\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\big|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0} > 0$). For the same reason, $s_\bullet$ also increases mean offspring knowledge (i.e., $\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x_\bullet})}{\partial s_\bullet}\big|_{\boldsymbol{x_\bullet}=\overline{\boldsymbol{x}}_0} > 0$; see

Appendix C). As a result, selection consistently favors the emergence of traits s that increase stochasticity in individual learning. These results indicate that, by increasing the effect of cultural selection, greater stochasticity in learning is favored by selection, thus confirming the potent role of σ_i on both the cultural and evolutionary dynamics.

4 Discussion

Here, we first derive the dynamics of individual knowledge in a population using a microscopic stochastic model of learning that links generations through vertical and oblique transmission. We show that as long as the learning process has an element of stochasticity, generating variability in knowledge, and knowledge is transmitted across generations, cultural selection, driven by differences in transmission linked to individual knowledge levels, will inevitably shape the dynamics of knowledge. By estimating the population mean of individual knowledge at equilibrium, our model shows that such cultural selection enhances cumulative knowledge. Those who, by chance, acquire greater knowledge are more likely to survive to adulthood, allowing them to interact with potential learners from the next generation, and to produce more offspring, thereby increasing opportunities for vertical transmission. As a result, individuals with higher knowledge are more likely to pass it on, thereby promoting the accumulation of knowledge across generations. Our results show that stochasticity in learning amplifies the action of natural selection by promoting variation in knowledge levels.

Our findings are consistent with previous theoretical work in which cultural selection arises because learners preferentially choose more knowledgeable exemplars (e.g., Henrich, 2004; Powell et al., 2009; Kobayashi and Aoki, 2012). However, we extend these results by showing that cultural selection should operate even when learners cannot directly assess exemplar knowledge, as long as individuals with greater knowledge are still more likely to transmit. Thus, cultural selection is likely to contribute to the improvement of any form of adaptive knowledge that enhances fecundity or survival, including ones in which the knowledgeability of exemplars is hard to assess. This may be particularly relevant in pre-industrial societies, where transmission often occurs in domains such as food processing, medicine, rituals, ecological knowledge, or institutional knowledge, whose benefits are not directly observable because they may be delayed, probabilistic, or rely on hidden causal mechanisms.

Stochasticity in learning has been extensively studied in reinforcement learning through the effects of random exploration of novel 'actions' (in the terminology of reinforcement learn-

ing, which can be thought of as cultural variants in our framework) on learning (Kaelbling et al., 1996; Ladosz et al., 2022; Hao et al., 2024). By increasing the chance of discovering high-reward actions, random exploration enhances the efficiency of selecting and retaining such actions, thereby improving reinforcement learning. While this role of stochasticity is well established, our results demonstrate that it can also enhance the efficiency of social learning, provided knowledge-biased transmission. In stationary environments, the advantages of random exploration in reinforcement learning generally decline as learners approach the optimal action (e.g., Tokic, 2010; Zhang et al., 2023). In contrast, in our model, greater stochasticity remains consistently advantageous because knowledge is unbounded. If we were instead to consider a cultural trait with an optimum (e.g., an ideal tool design or behavior suited to a specific ecological challenge), then, much like in reinforcement learning, stochasticity would likely promote early cultural improvement by enabling the discovery of more adaptive variants, but slow convergence as the population's traits approach their optimal value. For example, combining individual learning through trial-and-error with social transmission, trait variability across trials can speed convergence when the cultural trait is simple, but increases the distance from the optimum at equilibrium (Lehmann and Wakano, 2013).

Our results also show that, by enhancing cumulative knowledge, the cultural selection introduced by stochasticity in learning favors the evolution of extended social learning phases, as individuals can acquire substantial knowledge from exemplars. In addition, since social learning can provide a large amount of knowledge under these conditions, stochasticity in learning also promotes the evolution of more efficient learning, despite associated fecundity costs. This enables individuals to acquire a greater amount of knowledge during the social learning phase. Our findings may contribute to understanding how the mode of knowledge acquisition differs across knowledge types that vary in the extent of stochasticity in their production. For example, the production of opaque knowledge, such as adaptive taboos (Henrich and Henrich, 2010), adaptive supernatural beliefs (Lightner and Hagen, 2022), or food preparation methods that trigger complex chemical reactions (Beck, 1992), may be particularly stochastic, as their adaptive value is not readily apparent and it is difficult to intentionally produce knowledge that reliably enhances survival and fecundity. Due to this opacity, such knowledge likely became adaptive through a gradual accumulation of random improvements over time. As a result, individuals can acquire far more of this type of knowledge through social learning than they could by independently discovering it. Our results, therefore, predict that opaque knowledge would be primarily acquired socially. This could be the case for knowledge related to food preparation among the Aka Pygmies. Indeed, interviewed individuals reported that they had acquired all

of their knowledge related to the preparation of koko, magnoc, and palm wine through social learning (Hewlett and Cavalli-Sforza, 1986). By contrast, other forms of knowledge, such as basic hunting techniques, navigating terrain, or learning to avoid predators, may rely on more transparent causal relationships. As a result, their acquisition may involve less stochasticity and is more likely to occur through individual learning.

Our results reveal that whether knowledge affects fecundity or survival shapes the evolution of learning traits. For instance, when knowledge enhances only fecundity, natural selection favors individuals who mainly learn from their parents rather than from unrelated adults, consistent with the findings of McElreath and Strimling (2008). This is because, when stochasticity introduces variability in knowledge levels, parents tend to have above-average reproductive success and are therefore more likely to possess above-average knowledge. In contrast, randomly chosen adults tend to have average reproductive success and are not particularly likely to possess higher levels of knowledge. Therefore, individuals can acquire more knowledge by learning from their parents than from random adults. One interpretation is that parenthood serves as a cue for possessing knowledge that enhances fecundity, whereas simply being an adult provides no such indication. These findings suggest that knowledge enhancing fecundity should be preferentially acquired from parents. This pattern is supported by empirical evidence from the Aka Pygmies: interviewed individuals reported acquiring, on average, 85.6% of their knowledge about infant care from their parents, compared to 80.7% for knowledge across all domains (Hewlett and Cavalli-Sforza, 1986).

Our results rely on several assumptions that merit further discussion. In particular, we assume that knowledge has a linear effect on survival and fecundity. However, if this relationship were nonlinear, it could influence knowledge dynamics and the evolution of learning traits. For instance, if knowledge yields increasing returns on survival and fecundity, individuals who, by chance, possess above-average knowledge would gain greater advantages as knowledge accumulates in the population. This would amplify knowledge accumulation, since variation in knowledge would more strongly translate into differences in survival to adulthood and fecundity, thereby increasing the likelihood that higher levels of knowledge are transmitted to the next generation. In such a case, selection would favor greater reliance on social learning, as potential exemplars would carry a higher level of knowledge. By contrast, if knowledge yields diminishing returns, we would expect the opposite outcome: knowledge accumulation would slow, and the evolutionary pressure for social learning would weaken. We also assume that learning occurs sequentially through vertical, oblique, and individual phases. Although this is a

simplification, it aligns with a general developmental pattern: social learning is more common early in life, while individual learning becomes more prevalent with age (Reader and Laland, 2001; Biesmeijer and Seeley, 2005; Noble et al., 2014; Carr et al., 2015). In humans, the sources of social learning also tend to shift, from primarily vertical transmission during childhood, to greater reliance on oblique transmission in later life stages (Hewlett et al., 2011; Demps et al., 2012; Garfield et al., 2016).

Our study highlights the central role of stochasticity in learning in favoring cumulative knowledge and driving the evolution of learning behaviors. Such stochasticity may be widespread, not only because some level is inevitable, but also because our results show that selection favors increased stochasticity, as it increases the expected knowledge of lineage members when vertical transmission is non-negligible. However, fully understanding the evolution of traits underlying stochasticity in learning requires considering potential trade-offs; for example, a greater tendency to explore may be energetically costly or increase exposure to risk. These insights point to promising directions for future research on the evolutionary dynamics of traits impacting knowledge acquisition and their role in shaping cumulative cultural knowledge.

Acknowledgments

LM would like to thank Arthur Weyna and Cédric Perret for the useful discussions.

Author Contributions

LM and LL conceived and designed the study. LM developed the codes and performed the analyses. LM wrote the manuscript with contributions from LL.

Financial Support

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflicts of interest

The authors declare no conflicts of interest.

Code availability

All the codes used in this study are accessible at

github.com/Ludovic-Maisonneuve/stochasticity_evolution_learning

AI declaration

LM has used AI tools to identify grammatical and spelling errors and to enhance the overall fluency of the text.

Figures and Table

Figure 1: **Model overview.** **a** Illustration of the life cycle. **b** Illustration of the learning process. A focal individual can obtain knowledge (e.g., the skill set to crack nuts open, denoted by $k_{o\bullet} = k_\bullet(1)$ and represented here as a round set) by learning from three sources: (i) vertically from its parent (with knowledge $k_{p\bullet}$; blue arrow); (ii) obliquely from a randomly selected adult (with knowledge $k_{a\bullet}$, the knowledge of the parent $k_{p\bullet}$ and the oblique exemplar $k_{a\bullet}$ can also overlap and thus be redundant; green arrow); and (iii) individually, when it produces its own knowledge (in pink). See main text section 2.2.2 for more details. **c** A realisation of knowledge accumulation with a lifetime: individual knowledge $k_\bullet(a)$ of a focal offspring against its age a (realisation of the stochastic process defined by eq. (2) with traits $v_\bullet = 0.34$, $o_\bullet = 0.2$ and $\lambda_\bullet = 0.92$ for the offspring; and parameters $\beta_v = 1.4$, $\beta_o = 1.3$, $\alpha = 0.1$, $\epsilon = 0.05$, $\rho = 0.05$, $\sigma_v = \sigma_o = 0.1$, $\sigma_i = 0.25$, $k_{p\bullet} = k_{a\bullet} = 0.13$). The dashed line shows knowledge accumulation in the absence of stochasticity in learning, that is, when $\sigma_v = \sigma_o = \sigma_i = 0$. **d** Knowledge accumulation within a lineage: mean's adult knowledge $\overline{k}_{a,t}(\boldsymbol{x}_\bullet)$ within an $\boldsymbol{x}_\bullet$-lineage at each generation t (obtained from an individual-based simulation using the same parameters as in panel c, with trait mutation turned off and starting with a population of one ancestral individual with no knowledge, we set $k_{p\bullet} = k_{a\bullet} = 0$ for the ancestral individual, with $\gamma = 0.01$, $f_0 = 5$, $s_0 = 1$, $\eta_f = 25$, $\eta_s = 5$, $\theta = 0.1$); see Appendix D for more detail on individual-based simulations). The shaded area corresponds to cumulative knowledge (where individuals, on average, possess more knowledge than they could acquire through individual learning alone, i.e., where $\overline{k}_{a,t}(\boldsymbol{x}_\bullet) > \lambda_\bullet \alpha$). The dashed line shows the expected knowledge of a random adult of the $\boldsymbol{x}_\bullet$-lineage at equilibrium $\overline{k}_a^*(\boldsymbol{x}_\bullet)$ predicted by our analysis (see Section 2.3).

Table S1: Key symbols and their definitions

Model parameters

ϵ	Rate of knowledge obsolescence across generations
ρ	Average overlap in knowledge between two adults
$\beta_\mathrm{v}, \beta_\mathrm{o}$	Vertical and oblique transmission rate per fraction of investment allocated to learning
α	Individual learning rate per fraction of investment allocated to learning
$\sigma_\mathrm{v}, \sigma_\mathrm{o}, \sigma_\mathrm{i}$	Intensity of stochastic effects in vertical, oblique, and individual learning
θ	Fecundity cost exponent
f_0, s_0	Baseline fecundity and baseline survival modulator
$\eta_\mathrm{f}, \eta_\mathrm{s}$	Conversion factor that translates knowledge into fecundity and survival benefits
γ	Strength of density-dependent competition

Variables for traits

$\boldsymbol{x}_\bullet = (v_\bullet, o_\bullet, \lambda_\bullet)$	Trait values of a focal individual: vertical learning time ($v_\bullet$), oblique learning time ($o_\bullet$), and net investment in learning ($\lambda_\bullet$)
$\overline{\boldsymbol{x}} = (\overline{v}, \overline{o}, \overline{\lambda})$	Mean trait values in the population
$\overline{\boldsymbol{x}}^* = (\overline{v}^*, \overline{o}^*, \overline{\lambda}^*)$	Mean trait values favored by selection starting from a population initially expressing $\overline{\boldsymbol{x}} = (0,0,0)$
$\boldsymbol{G}$	Variance/covariance matrix of trait values in the population

Variables for knowledge and its transmission

$k_\bullet(a)$	Realized knowledge of the focal individual at age $a \in [0,1]$ (eq. (2))
$k_{\mathrm{o}\bullet}, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}$	Realized knowledge of a focal offspring, its parent, and its oblique exemplar at the end of learning (i.e, when $a = 1$)
$\phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)$	Stationary probability density that an adult from the $\boldsymbol{x}_\bullet$-lineage carries knowledge $k_{\mathrm{p}\bullet}$ at cultural equilibrium
$\overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet)$	Expected knowledge at cultural equilibrium for a random adult from generation t and for a random offspring produced by an adult of generation t in the $\boldsymbol{x}_\bullet$-lineage (eqs. (10) and (13))
$\overline{k}_\mathrm{a}^*(\boldsymbol{x}_\bullet), \overline{k}_\mathrm{o}^*(\boldsymbol{x}_\bullet)$	Expected knowledge at cultural equilibrium for a random adult and a random offspring of the $\boldsymbol{x}_\bullet$-lineage (eqs. (14) and (A.53))
$\omega_\mathrm{v}(\boldsymbol{x}_\bullet), \omega_\mathrm{o}(\boldsymbol{x}_\bullet)$	Proportion of available knowledge at the start of the vertical and oblique learning phases that is transmitted to a learner with traits $\boldsymbol{x}_\bullet$ (eq. (9))
$h_\mathrm{vl}(\boldsymbol{x}_\bullet), h_\mathrm{ol}(\boldsymbol{x}_\bullet)$	Vertical and oblique cultural heritability in the $\boldsymbol{x}_\bullet$-lineage (eqs. (12) and (11))
$V_{\mathrm{a},t}(\boldsymbol{x}_\bullet), V_{\mathrm{o},t}(\boldsymbol{x}_\bullet)$	Variances in the knowledge held by a random adult from generation t and for a random offspring produced by an adult of generation t in the $\boldsymbol{x}_\bullet$-lineage (eqs. (A.41) and (A.51))
$V_\mathrm{a}^*(\boldsymbol{x}_\bullet), V_\mathrm{o}^*(\boldsymbol{x}_\bullet)$	Equilibrium variances in the knowledge held by a random adult and a random offspring from the $\boldsymbol{x}_\bullet$-lineage (eqs. (15) and (A.53))

Variables for fitness

$f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})$	Fecundity of an adult with traits $\boldsymbol{x}_\bullet$ and realized knowledge $k_{\mathrm{p}\bullet}$ (eq. (1))
$s(k_{\mathrm{o}\bullet}, n_\mathrm{o})$	Survival probability of an offspring with realized knowledge $k_{\mathrm{o}\bullet}$ in a cohort of n_o competing offspring (eq. (3))
$\tilde{s}(k_{\mathrm{o}\bullet})$	Modulator of survival probability of an offspring with realized knowledge $k_{\mathrm{o}\bullet}$ (eq. (4))
$\overline{s}_\mathrm{o}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})$	Expected probability of survival for a random offspring of an adult with traits $\boldsymbol{x}_\bullet$ and knowledge $k_{\mathrm{p}\bullet}$
$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}})$	Expected lineage fitness in the $\boldsymbol{x}_\bullet$-lineage in a population with mean traits $\overline{\boldsymbol{x}}$ (eq. (5))
$\boldsymbol{s}(\overline{\boldsymbol{x}})$	Selection gradient (eq. (6))
$n_\mathrm{a}^*(\overline{\boldsymbol{x}}^*)$	Equilibrium number of adults when population mean traits are at evolutionary equilibrium $\overline{\boldsymbol{x}}^*$

Figure 2: **The accumulation of knowledge. a** Population mean knowledge $\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ and population knowledge variance $V_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ at cultural equilibrium according to intensity of stochastic effects in individual learning σ_{i} (left axis gives the scale of cumulative knowledge, and right axis gives the scale of knowledge variance). **b** Population mean knowledge $\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ according to intensity of stochastic effects in individual learning σ_{i} for different individual learning rate per fraction of investment allocated to learning α. Default parameters are: $f_0 = 5$, $s_0 = 1$, $\beta_{\mathrm{v}} = 1.4$, $\beta_{\mathrm{o}} = 1.3$, $\alpha = 0.1$, $\epsilon = 0.05$, $\rho = 0.05$, $\sigma_{\mathrm{i}} = 0.1$, $\eta_{\mathrm{f}} = 25$, $\eta_{\mathrm{s}} = 5$, $\overline{v} = 0.3$, $\overline{o} = 0.2$, $\overline{\lambda} = 0.9$.

Figure 3: **Evolution of learning traits. a-c** Learning schedule (y-axis) at $\overline{\boldsymbol{x}}^*$ against the intensity of stochastic effects in individual learning σ_i (x-axis) for different values of the conversion factor that translates knowledge into fecundity η_f and survival benefits η_s. Blue, green, and pink areas represent time spent performing vertical, oblique, and individual learning, respectively. **d-f** Investment in learning $\overline{\lambda}^*$ at $\overline{\boldsymbol{x}}^*$ corresponding to panels a-c. **g-j** Population mean knowledge $\overline{k}_a^*(\overline{\boldsymbol{x}}^*)$ (blue) and adult population size $n_a^*(\overline{\boldsymbol{x}}^*)$ (black) at $\overline{\boldsymbol{x}}^*$ corresponding to panels a-c (left axis gives scale of knowledge, and right axis gives scale of population size, with $\gamma = 10^{-4}$). Default parameters are the same as in fig. 2.

References

S. Ghirlanda and J. Lind. 'aesop's fable' experiments demonstrate trial-and-error learning in birds, but no causal understanding. *Animal Behaviour*, 123:239–247, 2017.

T. R. Zentall. Imitation: definitions, evidence, and mechanisms. *Animal Cognition*, 9(4):335–353, 2006.

L. A. Bates and R. W. Byrne. Imitation: what animal imitation tells us about animal cognition. *WIREs Cognitive Science*, 1(5):685–695, 2010.

L. L. Cavalli-Sforza and M. W. Feldman. *Cultural transmission and evolution: A quantitative approach.* Number 16 in Monographs in Population Biology. Princeton University Press, 1981.

R. Boyd and P. J. Richerson. *Culture and the Evolutionary Process.* University of Chicago Press, Chicago, 1985.

K. N. Laland. Social learning strategies. *Learning and Behavior*, 32:4–14, 2004.

R. L. Kendal, N. J. Boogert, L. Rendell, K. N. Laland, M. Webster, and P. L. Jones. Social learning strategies: Bridge-building between fields. *Trends in Cognitive Sciences*, 22(7):651–665, 2018.

A. Camacho-Alpízar and L. M. Guillette. From whom do animals learn? a meta-analysis on model-based social learning. *Psychonomic Bulletin & Review*, 30(3):863–881, 2023.

J. Henrich. The secret of our success: How culture is driving human evolution, domesticating our species, and making us smarter. In *The secret of our success*. princeton University press, 2015.

C. P. van Schaik. *The Primate Origin of Human Behavior.* Wiley-Blackwell, New Jersey, 2016.

G. R. Hunt and R. D. Gray. Diversification and cumulative evolution in new caledonian crow tool manufacture. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1517):867–874, 2003.

T. Sasaki and D. Biro. Cumulative culture can emerge from collective intelligence in animal groups. *Nature Communications*, 8(1):15049, 2017.

B. R. Jesmer, J. A. Merkle, J. R. Goheen, E. O. Aikens, J. L. Beck, A. B. Courtemanch, M. A. Hurley, D. E. McWhirter, H. M. Miyasaki, K. L. Monteith, and M. J. Kauffman. Is ungulate migration culturally transmitted? evidence of social learning from translocated animals. *Science*, 361(6406):1023–1025, 2018.

C. Gunasekaram, F. Battiston, O. Sadekar, C. Padilla-Iglesias, M. A. van Noordwijk, R. Furrer, A. Manica, J. Bertranpetit, A. Whiten, C. P. van Schaik, L. Vinicius, and A. B. Migliano. Population connectivity shapes the distribution and complexity of chimpanzee cumulative

culture. *Science*, 386(6724):920–925, 2024.

J. Henrich and R. Boyd. On Modeling Cognition and Culture: Why cultural evolution does not require replication of representations. *Journal of Cognition and Culture*, 2:87–112, 2002.

C. El Mouden, J. André, O. Morin, and D. Nettle. Cultural transmission and the evolution of human behaviour: a general approach based on the price equation. *Journal of Evolutionary Biology*, 27(2):231–241, 2014.

E. G. Aguilar and E. Akçay. Gene-culture coinheritance of a behavioral trait. *The American Naturalist*, 192(3):311–320, 2018. PMID: 30125232.

D. Nettle. Selection, adaptation, inheritance and design in human culture: the view from the price equation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375 (1797):20190358, 2020.

A. Mesoudi. Cultural selection and biased transformation: two dynamics of cultural evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1828):20200053, 2021.

M. Enquist, S. Ghirlanda, A. Jarrick, and C. A. Wachtmeister. Why does human culture increase exponentially? *Theoretical Population Biology*, 74:46–55, 2008.

K. Aoki, J. Y. Wakano, and L. Lehmann. Evolutionarily stable learning schedules in discrete generation models. *Theoretical Population Biology*, 81:300–309, 2012a.

W. Nakahashi. Evolution of improvement and cumulative culture. *Theoretical Population Biology*, 83:30–38, 2013.

J. Y. Wakano and C. Miura. Trade-off between learning and exploitation: The Pareto-optimal versus evolutionarily stable learning schedule in cumulative cultural evolution. *Theoretical Population Biology*, 91:37–43, 2014.

M. Kempe, S. J. Lycett, and A. Mesoudi. From cultural traditions to cumulative culture: parameterizing the differences between human and nonhuman culture. *Journal of Theoretical Biology*, 359:29–36, 2014.

J.-B. André and N. Baumard. Cultural evolution by capital accumulation. *Evolutionary Human Sciences*, 2:e18, 2020.

K. K. Denton, Y. Ram, and M. W. Feldman. Conditions that favour cumulative cultural evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 378(1872): 20210400, 2023.

J. Henrich. Demography and cultural evolution: how adaptive cultural processes can produce maladaptive losses: the Tasmanian case. *American Antiquity*, 69:197–214, 2004.

A. J. C. Micheletti. Modelling cultural selection on biological fitness to integrate social trans-

mission and adaptive explanations for human behaviour. *Evolutionary Human Sciences*, 2: e10, 2020.

A. Powell, S. Shennan, and M. G. Thomas. Late Pleistocene demography and the appearance of modern human behavior. *Science*, 324(5932):1298–1301, 2009.

Y. Kobayashi and K. Aoki. Innovativeness, population size and cumulative cultural evolution. *Theoretical Population Biology*, 82(1):38–47, 2012.

W. Nakahashi. Evolution of learning capacities and learning levels. *Theoretical Population Biology*, 78:211–224, 2010.

P. Tureček, J. Slavík, M. Kozák, and J. Havlíček. Non-particulate inheritance revisited: evolution in systems with parental variability-dependent inheritance. *Biological Journal of the Linnean Society*, 127(2):518–533, 2019.

F. Mery and T. J. Kawecki. An operating cost of learning in drosophila melanogaster. *Animal Behaviour*, 68(3):589–598, 2004.

J. M. S. Burger, M. Kolss, J. Pont, and T. J. Kawecki. Learning ability and longevity: A symmetrical evolutionary trade-off in drosophila. *Evolution*, 62(6):1294–1304, 2008.

E. C. Snell-Rood, G. Davidowitz, and D. R. Papaj. Reproductive tradeoffs of learning in a butterfly. *Behavioral Ecology*, 22(2):291–302, 2011.

S. Jaumann, R. Scudelari, and D. Naug. Energetic cost of learning and memory can cause cognitive impairment in honeybees. *Biology Letters*, 9(4):20130149, 2013.

I. C. Christiansen, S. Szin, and P. Schausberger. Benefit-cost trade-offs of early learning in foraging predatory mites amblyseius swirskii. *Scientific Reports*, 6(1):23571, 2016.

L. J. Evans, K. E. Smith, and N. E. Raine. Fast learning in free-foraging bumble bees is negatively correlated with lifetime resource collection. *Scientific Reports*, 7(1):496, 2017.

K. Aoki, J. Y. Wakano, and L. Lehmann. Evolutionarily stable learning schedules and cumulative culture in discrete generation models. *Theoretical Population Biology*, 81(4):300–309, 2012b.

L. Lehmann, J. Y. Wakano, and K. Aoki. On optimal learning schedules and the marginal value of cumulative cultural evolution. *Evolution*, 67:1435–1445, 2013.

Y. Kobayashi, J. Y. Wakano, and H. Ohtsuki. A paradox of cumulative culture. *Journal of Theoretical Biology*, 379:79–88, 2015.

Y. Kobayashi, H. Ohtsuki, and J. Y. Wakano. Population size vs. social connectedness — a gene-culture coevolutionary approach to cumulative cultural evolution. *Theoretical Population Biology*, 111:87–95, 2016.

C. Mullon and L. Lehmann. Invasion fitness for gene-culture co-evolution in family-structured

populations and an application to cumulative culture under vertical transmission. *Theoretical Population Biology*, 116:33–46, 2017.

H. Ohtsuki, J. Y. Wakano, and Y. Kobayashi. Inclusive fitness analysis of cumulative cultural evolution in an island-structured population. *Theoretical Population Biology*, 2017.

L. Maisonneuve, L. Lehmann, and C. Mullon. The coevolution of learning schedules and teaching enhances cumulative knowledge and drives a teacher–innovator syndrome. *Proceedings of the Royal Society B: Biological Sciences*, 292(2040):20242470, 2025.

M. Argyle and R. McHenry. Do spectacles really affect judgements of intelligence? *British Journal of Social and Clinical Psychology*, 10(1):27–29, 1971.

D. J. Lutz and F. C. Keil. Early understanding of the division of cognitive labor. *Child Development*, 73(4):1073–1084, 2002.

L. A. Wood, R. L. Kendal, and E. G. Flynn. Context-dependent model-based biases in cultural transmission: children's imitation is affected by model age over model knowledge state. *Evolution and Human Behavior*, 33(4):387–394, 2012.

Á. V. Jiménez and A. Mesoudi. Prestige-biased social learning: current evidence and outstanding questions. *Palgrave Communications*, 5(1):20, 2019.

M. Hirel, H. Meunier, R. Mundry, H. Rakoczy, J. Fischer, and S. Keupp. Choose your partner: Social evaluation of skillfulness at cooperative co-action tasks in tonkean macaques (Macaca tonkeana). *Animal Behavior and Cognition*, 12(3):330–359, 2025.

C. Gardiner. *Handbook of Stochastic Methods for Physics, Chemistry, and the Natural Sciences*. Proceedings in Life Sciences. Springer-Verlag, 1985.

O. Leimar. Multidimensional convergence stability. *Evolutionary Ecology Research*, 11(2):191–208, 2009.

M. Lynch and B. Walsh. *Genetics and Analysis of Quantitative Traits*. Sinauer, Massachusetts, 1998.

L. P. Kaelbling, M. L. Littman, and A. W. Moore. Reinforcement learning: A survey. *Journal of artificial intelligence research*, 4:237–285, 1996.

P. Ladosz, L. Weng, M. Kim, and H. Oh. Exploration in deep reinforcement learning: A survey. *Information Fusion*, 85:1–22, 2022.

J. Hao, T. Yang, H. Tang, C. Bai, J. Liu, Z. Meng, P. Liu, and Z. Wang. Exploration in deep reinforcement learning: From single-agent to multiagent domain. *IEEE Transactions on Neural Networks and Learning Systems*, 35(7):8762–8782, 2024.

M. Tokic. Adaptive ϵ-greedy exploration in reinforcement learning based on value differences. In R. Dillmann, J. Beyerer, U. D. Hanebeck, and T. Schultz, editors, *KI 2010: Advances in*

Artificial Intelligence, pages 203–210, Berlin, Heidelberg, 2010. Springer Berlin Heidelberg.

S. Zhang, H. Li, M. Wang, M. Liu, P.-Y. Chen, S. Lu, S. Liu, K. Murugesan, and S. Chaudhury. On the convergence and sample complexity analysis of deep q-networks with ϵ-greedy exploration. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023.

L. Lehmann and J. Y. Wakano. The handaxe and the microscope: individual and social learning in a multidimensional model of adaptation. *Evolution and Human Behavior*, 34(2):109–117, 2013.

J. Henrich and N. Henrich. The evolution of cultural adaptations: Fijian food taboos protect against dangerous marine toxins. *Proceedings of the Royal Society B: Biological Sciences*, 277 (1701):3715–3724, 2010.

A. D. Lightner and E. H. Hagen. All models are wrong, and some are religious: Supernatural explanations as abstract and useful falsehoods about complex realities. *Human Nature*, 33 (4):425–462, 2022.

W. Beck. Aboriginal preparation of cycas seeds in australia. *Economic Botany*, 46(2):133–147, 1992.

B. S. Hewlett and L. L. Cavalli-Sforza. Cultural transmission among aka pygmies. *American Anthropologist*, 88(4):922–934, 1986.

R. McElreath and P. Strimling. When natural selection favors imitation of parents. *Current Anthropology*, 49:307–316, 2008.

S. M. Reader and K. N. Laland. Primate innovation: sex, age and social rank differences. *Internation Journal of Primatology*, 22:787–805, 2001.

J. C. Biesmeijer and T. D. Seeley. The use of waggle dance information by honey bees throughout their foraging careers. *Behavioral Ecology and Sociobiology*, 59:133–142, 2005.

D. W. A. Noble, R. W. Byrne, and M. J. Whiting. Age-dependent social learning in a lizard. *Biology Letters*, 10(7), 2014.

K. Carr, R. L. Kendal, and E. G. Flynn. Imitate or innovate? children's innovation is influenced by the efficacy of observed behaviour. *Cognition*, 142:322–332, 2015.

B. S. Hewlett, H. N. Fouts, A. H. Boyette, and B. L. Hewlett. Social learning among Congo Basin hunter-gatherers. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 366(1567):1168–1178, 2011.

K. Demps, F. Zorondo-Rodríguez, C. García, and V. Reyes-García. Social learning across the life cycle: cultural knowledge acquisition for honey collection among the Jenu Kuruba India. *Evolution and Human Behavior*, 33:460–470, 2012.

Z. H. Garfield, M. J. Garfield, and B. S. Hewlett. *A Cross-Cultural Analysis of Hunter-Gatherer Social Learning*, pages 19–34. Springer Japan, Tokyo, 2016.

Appendix to "How does stochasticity in learning impact the accumulation of knowledge and the evolution of learning?"

Ludovic Maisonneuve[1,*], Laurent Lehmann[1]

1. Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

∗ ludovic.maisonneuve.2015@polytechnique.org.

Appendix A: Cultural dynamics and equilibrium

Here, we begin by highlighting the similarities and differences between our learning model and previous works (section A.1). We then characterize the outcome of the learning process described by eq. (2) in the absence of stochasticity in social learning (section A.2), and derive the recurrence equations for the expected knowledge and knowledge variance within an $x_\bullet$-lineage using a Gaussian moment closure (section A.3). Next, we characterize the cultural equilibrium and develop a numerical procedure for computing the population mean knowledge and knowledge variance (section A.4).

A.1 Connection to previous models

Our learning model, described in the main text and formalized in eq. (2), builds on previous work by Kobayashi et al. (2016) and Maisonneuve et al. (2025), sharing some core assumptions while differing in key aspects. Following Kobayashi et al. (2016), we model stochasticity in both individual and social learning using Brownian noise (see eq. (2) in the main text and eq. (15) in the appendix of Kobayashi et al., 2016). As in Maisonneuve et al. (2025), we assume that learning occurs sequentially through vertical, oblique, and individual phases. A key difference between our model and that of Maisonneuve et al. (2025) lies in the structure of oblique learning:

while Maisonneuve et al. (2025) allows individuals to learn from all adults, we assume they learn from only one, reflecting the realistic constraint that offspring cannot interact with everyone. Learning from the entire adult population can inflate the available knowledge, especially under stochastic learning, where some individuals, by chance, accumulate significantly more knowledge than others.

A.2   Knowledge acquisition without stochasticity in social learning

In this section, we derive eq. (8), which provides an expression for computing the realized knowledge acquired by a focal offspring at the end of its learning process, under the assumption that social learning occurs without stochasticity (i.e., $\sigma_{\mathrm{v}} = \sigma_{\mathrm{o}} = 0$). When we neglect stochasticity during social learning, the realized knowledge of a focal offspring at each age $a \in [0, 1]$ with traits $\boldsymbol{x}_\bullet = (v_\bullet, o_\bullet, \lambda_\bullet)$, who learn from a parent with knowledge $k_{\mathrm{p}\bullet}$ and an oblique exemplar with knowledge $k_{\mathrm{a}\bullet}$, is a realization of the following stochastic differential equation (obtained by substituting $\sigma_{\mathrm{v}} = \sigma_{\mathrm{o}} = 0$ into eq. (2))

$$\frac{\mathrm{d}k_\bullet(a)}{\mathrm{d}a} = \begin{cases} \lambda_\bullet \, \beta_{\mathrm{v}} \, [(1 - \epsilon) \, k_{\mathrm{p}\bullet} - k_\bullet(a)] & \text{for } a \in [0, v_\bullet) \\ \lambda_\bullet \, \beta_{\mathrm{o}} \, \big[(1 - \epsilon) \, k_{\mathrm{a}\bullet} - \rho \, k_\bullet(v_\bullet) - \big(k_\bullet(a) - k_\bullet(v_\bullet)\big)\big] & \text{for } a \in [v_\bullet, v_\bullet + o_\bullet) \\ \lambda_\bullet \, \big(\alpha + \sigma_{\mathrm{i}} \, \eta(a)\big) & \text{for } a \in [v_\bullet + o_\bullet, 1], \end{cases} \tag{A.1}$$

with $k_\bullet(0) = 0$.

For the final phase of learning, where the dynamics include a stochastic component, eq. (A.1) is interpreted in the Itô sense (Gardiner, 1985, Sec. 4.2, pp. 83–84)

$$k_\bullet(a) = k_\bullet(v_\bullet + o_\bullet) + \int_{v_\bullet + o_\bullet}^{a} \lambda_\bullet \, \alpha \, \mathrm{d}s + \int_{v_\bullet + o_\bullet}^{a} \lambda_\bullet \, \sigma_{\mathrm{i}} \, \mathrm{d}B_s, \tag{A.2}$$

where B_s is a standard Brownian motion.

Accordingly, the knowledge accumulated by the focal offspring at the end of the learning period is given by

$$k_\bullet(1) = k_\bullet(v_\bullet + o_\bullet) + \int_{v_\bullet + o_\bullet}^{1} \lambda_\bullet \, \alpha \, \mathrm{d}s + \int_{v_\bullet + o_\bullet}^{1} \lambda_\bullet \, \sigma_{\mathrm{i}} \, \mathrm{d}B_s, \tag{A.3}$$

which simplifies to

$$k_\bullet(1) = k_\bullet(v_\bullet + o_\bullet) + \lambda_\bullet \, \alpha \, (1 - v_\bullet - o_\bullet) + \lambda_\bullet \, \sigma_{\mathrm{i}} \, B_{1 - v_\bullet - o_\bullet}. \tag{A.4}$$

Substituting the explicit expression for $k_\bullet(v_\bullet + o_\bullet)$, obtained by solving eq. (A.1) over the deterministic phases, into eq. (A.4) yields

$$k_\bullet(1) = \overbrace{\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})}^{\text{expected knowledge acquired}} + \overbrace{\chi_{\mathrm{k}}}^{\substack{\text{stochastic fluctuations} \\ \text{in knowledge acquired}}}, \tag{A.5}$$

where

$$\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) = \omega_{\mathrm{v}}(\boldsymbol{x}_\bullet)\,(1-\epsilon)\,k_{\mathrm{p}\bullet} + \omega_{\mathrm{o}}(\boldsymbol{x}_\bullet)\,[(1-\epsilon)\,k_{\mathrm{a}\bullet} - \rho\,\omega_{\mathrm{v}}(\boldsymbol{x}_\bullet)\,(1-\epsilon)\,k_{\mathrm{p}\bullet}] + \lambda_\bullet\,\alpha\,(1-v_\bullet-o_\bullet), \tag{A.6}$$

is the expected knowledge acquired. The terms $\omega_{\mathrm{v}}(\boldsymbol{x}_\bullet)$ and $\omega_{\mathrm{o}}(\boldsymbol{x}_\bullet)$, given by

$$\omega_{\mathrm{v}}(\boldsymbol{x}_\bullet) = 1 - \mathrm{e}^{-\lambda_\bullet\,\beta_{\mathrm{v}}\,v_\bullet} \quad \text{and} \quad \omega_{\mathrm{o}}(\boldsymbol{x}_\bullet) = 1 - \mathrm{e}^{-\lambda_\bullet\,\beta_{\mathrm{o}}\,o_\bullet}. \tag{A.7}$$

are the proportion of available knowledge at the start of the vertical and oblique learning phases, respectively, that is effectively transmitted to the focal offspring.

The term

$$\chi_{\mathrm{k}} = \lambda_\bullet\,\sigma_{\mathrm{i}}\,B_{1-v_\bullet-o_\bullet}, \tag{A.8}$$

captures the effect of stochastic fluctuations in individual learning. Since $B_{1-v_\bullet-o_\bullet}$ is a realization of a Gaussian random variable with mean 0 and variance $1 - v_\bullet - o_\bullet$ (Bass, 2011), it follows that χ_{k} can be obtained as a realization of a Gaussian variable with mean 0 and variance $\lambda_\bullet^2\,\sigma_{\mathrm{i}}^2\,(1 - v_\bullet - o_\bullet)$.

Substituting the expression for $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ from eq. (A.6) into eq. (A.5) gives eq. (8) of the main text.

To simplify the notation in what follows, we rewrite eq. (A.6) as

$$\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) = h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)\,k_{\mathrm{p}\bullet} + h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)\,k_{\mathrm{a}\bullet} + p_{\mathrm{il}}(\boldsymbol{x}_\bullet), \tag{A.9}$$

where terms $h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)$, $h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)$ and $p_{\mathrm{il}}(\boldsymbol{x}_\bullet)$ are given by

$$h_{\mathrm{vl}}(\boldsymbol{x}_\bullet) = (1-\epsilon)\,\omega_{\mathrm{v}}(\boldsymbol{x}_\bullet)\,(1 - \rho\,\omega_{\mathrm{o}}(\boldsymbol{x}_\bullet)), \tag{A.10}$$

$$h_{\mathrm{ol}}(\boldsymbol{x}_\bullet) = (1-\epsilon)\,\omega_{\mathrm{o}}(\boldsymbol{x}_\bullet), \tag{A.11}$$

and

$$p_{\text{il}}(\boldsymbol{x}_\bullet) = \lambda_\bullet \, \alpha \, (1 - v_\bullet - o_\bullet). \tag{A.12}$$

A.3 Knowledge dynamics

In this section, we derive expressions for the change in the expected knowledge (section A.3.1) and knowledge variance (section A.3.2) in an $\boldsymbol{x}_\bullet$-lineage. Then we apply a Gaussian closure approximation to derive a closed dynamical system for tracking across generations the expected knowledge and knowledge variance (section A.3.3).

A.3.1 Expected knowledge dynamics

Here, we detail how we get eqs. (10) and (13) from the main text, which give the dynamics of the expected knowledge in an $\boldsymbol{x}_\bullet$-lineage. Let $\overline{k}_{\text{a},t}(\boldsymbol{x}_\bullet)$ denote the expected knowledge of a random adult from the $\boldsymbol{x}_\bullet$-lineage at generation t for all $t \geq 1$ (here and throughout, the subscript $_{\text{a},t}$ denotes quantities computed among adults of generation t; see fig. A.1). By definition,

$$\overline{k}_{\text{a},t+1}(\boldsymbol{x}_\bullet) = \int k_{\text{o}\bullet} \, \phi_{\text{a},t+1}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{\text{o}\bullet}, \tag{A.13}$$

where $\phi_{\text{a},t+1}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)$ denotes the probability density that a random adult from the $\boldsymbol{x}_\bullet$-lineage of generation $t + 1$ possesses knowledge $k_{\text{o}\bullet}$. Note that we denote the knowledge of a focal adult of generation $t + 1$ as $k_{\text{o}\bullet}$ because here, we view adults as the surviving offspring from the previous generation.

Figure A.1: **Notation specifying the stage in the life cycle and generation at which variables are calculated.**

The probability density that a random adult from the $\boldsymbol{x}_\bullet$-lineage of generation $t + 1$ possesses knowledge $k_{\text{o}\bullet}$, $\phi_{\text{a},t+1}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)$, can be derived from the probability density that a random offspring produced by an adult from the $\boldsymbol{x}_\bullet$-lineage of generation t possesses knowledge $k_{\text{o}\bullet}$, $\phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)$ (here and throughout, the subscript $_{\text{o},t}$ denotes quantities computed among an

offspring produced by an adult of generation t after learning is complete and before the survival stage; see fig. A.1), who will potentially become one of the adults of generation $t + 1$. The probability density of knowledge of a random offspring is then weighted by the relative survival probability associated with each possible knowledge:

$$\forall k_{\text{o}\bullet}, \quad \phi_{\text{a},t+1}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet) = \frac{s(k_{\text{o}\bullet}, n_{\text{o},t})}{\overline{s}_{\bullet,t}} \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet), \tag{A.14}$$

where $n_{\text{o},t}$ is the number of offspring produced by adults of generation t, $s(k_{\text{o}\bullet}, n_{\text{o},t})$ is the survival probability associated with traits $\boldsymbol{x}_\bullet$ and knowledge $k_{\text{o}\bullet}$, and $\overline{s}_{\bullet,t}$ is the expected survival probability of an offspring produced by an adult of the $\boldsymbol{x}_\bullet$-lineage of generation t, given by

$$\overline{s}_{\bullet,t} = \int s(k_{\text{o}\bullet}, n_{\text{o},t}) \, \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet) \, dk_{\text{o}\bullet}. \tag{A.15}$$

Noting from eqs. (3) and (4) that $s(k_{\text{o}\bullet}, n_{\text{o},t})$ depends linearly on $k_{\text{o}\bullet}$ and using that $\int \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet) \, dk_{\text{o}\bullet} = 1$ we have

$$\overline{s}_{\bullet,t} = s\Big(\int k_{\text{o}\bullet} \, \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet) \, dk_{\text{o}\bullet}, n_{\text{o},t} \Big) = s(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet), n_{\text{o},t}). \tag{A.16}$$

By substituting the expressions of $\phi_{\text{a},t+1}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)$ and $\overline{s}_{\bullet,t}$ from eqs. (A.14) and (A.16) into eq. (A.13) we obtain

$$\overline{k}_{\text{a},t+1}(\boldsymbol{x}_\bullet) = \int k_{\text{o}\bullet} \, \frac{s(k_{\text{o}\bullet}, n_{\text{o},t})}{s(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet), n_{\text{o},t})} \, \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet) \, dk_{\text{o}\bullet}. \tag{A.17}$$

Next, we rewrite $s(k_{\text{o}\bullet}, n_{\text{o},t})/s(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet), n_{\text{o},t})$. Using the expression for $s(k_{\text{o}\bullet}, n_{\text{o},t})$ and $s(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet), n_{\text{o},t})$ from eq. (3) with $n_{\text{o}} = n_{\text{o},t}$ and with $k_{\text{o}\bullet} = \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)$ and $n_{\text{o}} = n_{\text{o},t}$ we obtain

$$\frac{s(k_{\text{o}\bullet}, n_{\text{o},t})}{s(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet), n_{\text{o},t})} = \frac{\tilde{s}(k_{\text{o}\bullet})}{\tilde{s}(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet))}. \tag{A.18}$$

By substituting $k_{\text{o}\bullet} = \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet) + \xi_{k_{\text{o}\bullet},t}$, where $\xi_{k_{\text{o}\bullet},t} = k_{\text{o}\bullet} - \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)$, into eq. (A.18) and noting from eq. (4) that $\tilde{s}(k_{\text{o}\bullet})$ depends linearly on $k_{\text{o}\bullet}$ with a linear coefficient η_{s}, we obtain

$$\frac{s(k_{\text{o}\bullet}, n_{\text{o},t})}{s(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet), n_{\text{o},t})} = 1 + \xi_{k_{\text{o}\bullet},t} \frac{\eta_{\text{s}}}{\tilde{s}(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet))}. \tag{A.19}$$

Substituting $k_{o\bullet} = \overline{k}_{o,t}(\boldsymbol{x}_\bullet) + \xi_{k_{o\bullet},t}$ and eq. (A.19) into eq. (A.17) gives

$$\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet) = \overline{k}_{o,t}(\boldsymbol{x}_\bullet) + \frac{\eta_s}{\tilde{s}\big(\overline{k}_{o,t}(\boldsymbol{x}_\bullet)\big)} \int \xi_{k_{o\bullet},t}^2 \, \phi_{o,t}(k_{o\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{o\bullet}. \tag{A.20}$$

Noting that the integral is the variance in the knowledge held by a random offspring produced by an adult from the $\boldsymbol{x}_\bullet$-lineage at generation t, denoted as $V_{o,t}(\boldsymbol{x}_\bullet)$, we obtain

$$\boxed{\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet) = \overline{k}_{o,t}(\boldsymbol{x}_\bullet) + \frac{V_{o,t}(\boldsymbol{x}_\bullet)\,\eta_s}{\tilde{s}\big(\overline{k}_{o,t}(\boldsymbol{x}_\bullet)\big)},} \tag{A.21}$$

which is eq. (13) from the main text.

Note that, in eq. (A.21), the term $V_{o,t}(\boldsymbol{x}_\bullet)\,\eta_s/\tilde{s}\big(\overline{k}_{o,t}(\boldsymbol{x}_\bullet)\big)$ captures the effect of cultural selection arising from differential survival among lineage members, due to variation in knowledge.

To obtain the expected knowledge dynamics across one generation, we need to determine the expected knowledge of a random offspring produced by an adult of generation t, $\overline{k}_{o,t}(\boldsymbol{x}_\bullet)$. By definition,

$$\overline{k}_{o,t}(\boldsymbol{x}_\bullet) = \int k_{o\bullet} \, \phi_{o,t}(k_{o\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{o\bullet}. \tag{A.22}$$

For each offspring, the knowledge acquired depends on the knowledge of both the parent and the oblique exemplar, as well as on the outcome of the stochasticity in the learning process. By integrating over the probability density of parental knowledge, the probability density of knowledge among potential oblique exemplars, and the probability density of stochastic learning outcomes, we obtain

$$\overline{k}_{o,t}(\boldsymbol{x}_\bullet) = \iiint k_{o\bullet} \, \nu(k_{o\bullet} \mid \boldsymbol{x}_\bullet, k_{p\bullet}, k_{a\bullet}) \, \phi_{a,t}(k_{a\bullet} \mid \overline{\boldsymbol{x}}) \, \frac{f(\boldsymbol{x}_\bullet, k_{p\bullet})}{\overline{f}_{\bullet,t}} \, \phi_{a,t}(k_{p\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{o\bullet} \, \mathrm{d}k_{a\bullet} \, \mathrm{d}k_{p\bullet}, \tag{A.23}$$

where $\overline{f}_{\bullet,t}$ is the mean fecundity of adults belonging to the $\boldsymbol{x}_\bullet$-lineage at generation t and is equal to

$$\overline{f}_{\bullet,t} = \int f(\boldsymbol{x}_\bullet, k_{p\bullet}) \, \phi_{a,t}(k_{p\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{p\bullet} = f\big(\boldsymbol{x}_\bullet, \overline{k}_{a,t}(\boldsymbol{x}_\bullet)\big). \tag{A.24}$$

The integral over $k_{p\bullet}$ encompasses all possible knowledge value of parents, weighted by the probability density $f(\boldsymbol{x}_\bullet, k_{p\bullet})/\overline{f}_{\bullet,t} \, \phi_{a,t}(k_{p\bullet} \mid \boldsymbol{x}_\bullet)$ that a parent has knowledge $k_{p\bullet}$. Each parent's contribution depends on its fecundity relative to the lineage's mean fecundity, $f(\boldsymbol{x}_\bullet, k_{p\bullet})/\overline{f}_{\bullet,t}$, because lineage members with higher fecundity are more likely to transmit their traits and

knowledge.

The integral over $k_{\mathrm{a}\bullet}$ encompasses all possible knowledge values of oblique exemplars in generation t, weighted by the probability density $\phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})$ that an oblique exemplar has knowledge $k_{\mathrm{a}\bullet}$. Since the variances of population traits are small, we approximate the population as being quasi-monomorphic for the mean traits $\overline{\boldsymbol{x}}$, and thus assume that offspring acquire knowledge obliquely from adults with traits $\overline{\boldsymbol{x}}$.

Finally, the integral over $k_{\mathrm{o}\bullet}$ encompasses all possible outcomes of the learning process, weighted by the probability density of the offspring's knowledge $k_{\mathrm{o}\bullet}$ at the end of the learning process $\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$.

From eq. (A.5) with $k_\bullet(1) = k_{\mathrm{o}\bullet}$, we have that $k_{\mathrm{o}\bullet} = \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) + \chi_{\mathrm{k}}$ where $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ is deterministic and χ_{k} is a realization a Gaussian random variable with mean 0 and variance $\lambda_\bullet^2 \sigma_{\mathrm{i}}^2 (1 - v_\bullet - o_\bullet)$. Since $k_{\mathrm{o}\bullet}$ is the sum of a deterministic term $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ and a random fluctuation χ_{k}, its probability density has the same shape as that of χ_{k}, but centered around $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ instead of 0. This means the probability density of $k_{\mathrm{o}\bullet}$ is obtained by shifting the density of χ_{k} horizontally so that its mean aligns with $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$

$$\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) = \nu_{\chi_{\mathrm{k}}}\big(k_{\mathrm{o}\bullet} - \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) \mid \boldsymbol{x}_\bullet\big), \tag{A.25}$$

where $\nu_{\chi_{\mathrm{k}}}$ is a Gaussian probability function with mean 0 and variance $\lambda_\bullet^2 \sigma_{\mathrm{i}}^2 (1 - v_\bullet - o_\bullet)$. We obtain eq. (A.25) by noting that $k_{\mathrm{o}\bullet} - \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) = \chi_{\mathrm{k}}$, so the probability that $k_{\mathrm{o}\bullet}$ takes a particular value is the same as the probability that χ_{k} takes the value $k_{\mathrm{o}\bullet} - \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$.

By substituting the expressions of $\overline{f}_{\bullet,t}$ and $\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ from eqs. (A.24) and (A.25) into eq. (A.23) and by performing a change of variable $\chi_{\mathrm{k}} = k_{\mathrm{o}\bullet} - \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ we obtain

$$\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \iiint \bigg[(\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) + \chi_{\mathrm{k}})$$
$$\times \nu_{\chi_{\mathrm{k}}}(\chi_{\mathrm{k}} \mid \boldsymbol{x}_\bullet) \, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}}) \, \frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)\big)} \, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \bigg] \mathrm{d}\chi_{\mathrm{k}} \, \mathrm{d}k_{\mathrm{a}\bullet} \, \mathrm{d}k_{\mathrm{p}\bullet}. \tag{A.26}$$

Equation (A.26) can be rearranged as follows

$$\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \iint \left[\left(\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) + \int \chi_{\mathrm{k}}\, \nu_{\chi_{\mathrm{k}}}(\chi_{\mathrm{k}} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}\chi_{\mathrm{k}} \right) \right.$$
$$\left. \times\, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))}\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \right]\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}, \quad \text{(A.27)}$$

and, because χ_{k} is centered at 0 it simplifies to

$$\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \iint \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})\, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))}\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}. \quad \text{(A.28)}$$

Next, we expand $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})/f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))$. By substituting $k_{\mathrm{p}\bullet} = \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{p}\bullet},t}$, where $\xi_{k_{\mathrm{p}\bullet},t} = k_{\mathrm{p}\bullet} - \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)$, and noting from eq. (1) that $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})$ depends linearly on $k_{\mathrm{p}\bullet}$, we obtain

$$\frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))} = 1 + \xi_{k_{\mathrm{p}\bullet},t}\, \frac{(1 - \lambda_\bullet)^\theta\, \eta_{\mathrm{f}}}{f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))}. \quad \text{(A.29)}$$

By substituting the expression of $f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))$ from eq. (1) with $k_{\mathrm{p}\bullet} = \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)$ into the right-hand side of eq. (A.29) we obtain

$$\frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet))} = 1 + \xi_{k_{\mathrm{p}\bullet},t}\, \frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)}. \quad \text{(A.30)}$$

We now rewrite $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$. Because $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ is linear on $k_{\mathrm{p}\bullet}$ and $k_{\mathrm{a}\bullet}$ with coefficient of linearity $h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)$ and $h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)$ (see eq. (A.9)), by substituting $k_{\mathrm{p}\bullet} = \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{p}\bullet},t}$ and $k_{\mathrm{a}\bullet} = \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}}) + \xi_{k_{\mathrm{a}\bullet},t}$, where $\xi_{k_{\mathrm{a}\bullet},t} = \left(k_{\mathrm{a}\bullet} - \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right)$, we obtain

$$\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) = \mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) + \xi_{k_{\mathrm{p}\bullet},t}\, h_{\mathrm{vl}}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{a}\bullet},t}\, h_{\mathrm{ol}}(\boldsymbol{x}_\bullet). \quad \text{(A.31)}$$

Substituting eqs. (A.30) and (A.31) into eq. (A.28), and noting that $\xi_{k_{\mathrm{p}\bullet},t}$ and $\xi_{k_{\mathrm{a}\bullet},t}$ are centered around zero and are independent, we obtain

$$\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) + h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)\, \frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \int \xi_{k_{\mathrm{p}\bullet},t}^2\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}. \quad \text{(A.32)}$$

Noting that the integral is the variance in the knowledge held by a random adult from the $\boldsymbol{x}_\bullet$-lineage at generation t, denoted as $V_{\mathrm{a},t}(\boldsymbol{x}_\bullet)$, we obtain

$$\boxed{\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) + \frac{h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)\, V_{\mathrm{a},t}(\boldsymbol{x}_\bullet)\, \eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)}.} \quad \text{(A.33)}$$

By substituting the expression of $\mathcal{L}\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\big)$ from eq. (A.9) with $k_{\mathrm{p}\bullet} = \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)$ and $k_{\mathrm{a}\bullet} = \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})$ gives eq. (10) from the main text.

Note that, in eq. (A.33), the term $h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)\, V_{\mathrm{a},t}(\boldsymbol{x}_\bullet)\, \eta_{\mathrm{f}}/\big(f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)\big)$ captures the effect of cultural selection arising from differential fecundity among lineage members, due to variation in knowledge.

Equations (A.21) and (A.33) give the dynamics of the expected knowledge within the $\boldsymbol{x}_\bullet$-lineage.

A.3.2   Knowledge variance dynamics

Equations (A.21) and (A.33) enable us to monitor changes in expected knowledge within a lineage across a single generation. However, to iterate these equations also requires tracking the variance in the knowledge held by a lineage member simultaneously. In this section, we derive the expression of the dynamics of the variance in the knowledge held by a random adult within the $\boldsymbol{x}_\bullet$-lineage.

By definition, the variance in the knowledge held by a random adult from the $\boldsymbol{x}_\bullet$-lineage at generation $t + 1$, $V_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet)$, is given by

$$V_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) = \int \big(k_{\mathrm{o}\bullet} - \overline{k}_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet)\big)^2 \, \phi_{\mathrm{a},t+1}(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{\mathrm{o}\bullet}. \tag{A.34}$$

Using $k_{\mathrm{o}\bullet} - \overline{k}_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) = \xi_{k_{\mathrm{o}\bullet},t} - \big(\overline{k}_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet)\big)$ and $\int \xi_{k_{\mathrm{o}\bullet},t} \, \phi_{\mathrm{a},t+1}(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{\mathrm{o}\bullet} = \overline{k}_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet)$ we can expand eq. (A.34) as follows

$$V_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) = \int \xi_{k_{\mathrm{o}\bullet},t}^2 \, \phi_{\mathrm{a},t+1}(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{\mathrm{o}\bullet} - \big(\overline{k}_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet)\big)^2. \tag{A.35}$$

By substituting the expression of $\phi_{\mathrm{a},t+1}(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet)$ from eq. (A.14) into eq. (A.35), we obtain

$$V_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) = \int \xi_{k_{\mathrm{o}\bullet},t}^2 \, \frac{s(k_{\mathrm{o}\bullet}, n_{\mathrm{o},t})}{\overline{s}_{\bullet,t}} \, \phi_{\mathrm{o},t}(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{\mathrm{o}\bullet} - \big(\overline{k}_{\mathrm{a},t+1}(\boldsymbol{x}_\bullet) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet)\big)^2. \tag{A.36}$$

Substituting the expression of $s(k_{\mathrm{o}\bullet}, n_{\mathrm{o},t})/\overline{s}_{\bullet,t}$ from eq. (A.19) (with $s(\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet), n_{\mathrm{o},t}) = \overline{s}_{\bullet,t}$) into

eq. (A.36) we find

$$V_{\text{a},t+1}(\boldsymbol{x}_\bullet) = \int \xi_{k_{\text{o}\bullet},t}^2 \left(1 + \xi_{k_{\text{o}\bullet},t} \frac{\eta_{\text{s}}}{\tilde{s}\big(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)}\right) \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)\,\text{d}k_{\text{o}\bullet} - \big(\overline{k}_{\text{a},t+1}(\boldsymbol{x}_\bullet) - \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)^2 .$$
(A.37)

By expanding the integral in eq. (A.37) we find

$$V_{\text{a},t+1} = \int \xi_{k_{\text{o}\bullet},t}^2\,\phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)\,\text{d}k_{\text{o}\bullet} + \frac{\eta_{\text{s}}}{\tilde{s}\big(\overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)} \int \xi_{k_{\text{o}\bullet},t}^3\,\phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)\,\text{d}k_{\text{o}\bullet}$$
$$- \big(\overline{k}_{\text{a},t+1}(\boldsymbol{x}_\bullet) - \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)^2 . \quad \text{(A.38)}$$

Noting that

$$\int \big(k_{\text{o}\bullet} - \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)^2 \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)\,\text{d}k_{\text{o}\bullet} = V_{\text{o},t}(\boldsymbol{x}_\bullet),$$
(A.39)

and

$$\int \big(k_{\text{o}\bullet} - \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)^3 \phi_{\text{o},t}(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet)\,\text{d}k_{\text{o}\bullet} = C_{k^3,\text{o},t}(\boldsymbol{x}_\bullet),$$
(A.40)

where $C_{k^3,\text{o},t}(\boldsymbol{x}_\bullet)$ is the skewness in the knowledge held by a random offspring produced by an adult from the $\boldsymbol{x}_\bullet$-lineage at generation t, and by substituting the expression for $\overline{k}_{\text{a},t+1}(\boldsymbol{x}_\bullet)$ from eq. (A.21) into eq. (A.38), we obtain

$$\boxed{V_{\text{a},t+1}(\boldsymbol{x}_\bullet) = V_{\text{o},t}(\boldsymbol{x}_\bullet) + \frac{C_{k^3,\text{o},t}(\boldsymbol{x}_\bullet)\,\eta_{\text{s}}}{\tilde{s}\big(\overline{k}_{\text{o}}^*(\boldsymbol{x}_\bullet)\big)} - \left(\frac{V_{\text{o},t}(\boldsymbol{x}_\bullet)\,\eta_{\text{s}}}{\tilde{s}\big(\overline{k}_{\text{o}}^*(\boldsymbol{x}_\bullet)\big)}\right)^2 .}$$
(A.41)

We now compute the variance in the knowledge held by a random offspring produced by an adult from the $\boldsymbol{x}_\bullet$-lineage at generation t, $V_{\text{o},t}(\boldsymbol{x}_\bullet)$. The variance $V_{\text{o},t}(\boldsymbol{x}_\bullet)$ can be obtained by integrating over the probability density of parental knowledge, the probability density of knowledge among potential oblique exemplars, and the probability density of all stochastic learning outcomes to obtain

$$V_{\text{o},t}(\boldsymbol{x}_\bullet) = \iiint \left[\big(k_{\text{o}\bullet} - \overline{k}_{\text{o},t}(\boldsymbol{x}_\bullet)\big)^2 \right.$$
$$\left. \times \nu(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\text{p}\bullet}, k_{\text{a}\bullet})\,\phi_{\text{a},t}(k_{\text{a}\bullet} \mid \overline{\boldsymbol{x}}) \frac{f(\boldsymbol{x}_\bullet, k_{\text{p}\bullet})}{\overline{f}_{\bullet,t}}\,\phi_{\text{a},t}(k_{\text{p}\bullet} \mid \boldsymbol{x}_\bullet)\right] \text{d}k_{\text{o}\bullet}\,\text{d}k_{\text{a}\bullet}\,\text{d}k_{\text{p}\bullet}. \quad \text{(A.42)}$$

By substituting the expressions of $\overline{f}_{\bullet,t}$ and $\nu(k_{\text{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\text{p}\bullet}, k_{\text{a}\bullet})$ from eqs. (A.24) and (A.25) into

eq. (A.42) and by performing a change of variable $\chi_{\mathrm{k}} = k_{\mathrm{o}\bullet} - \mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ we obtain

$$V_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \iiint \Bigg[\big(\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet}) + \chi_{\mathrm{k}} - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) \big)^2$$
$$\times \nu_{\chi_{\mathrm{k}}}(\chi_{\mathrm{k}} \mid \boldsymbol{x}_\bullet)\, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)\big)}\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \Bigg]\, \mathrm{d}\chi_{\mathrm{k}}\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}. \quad \text{(A.43)}$$

By substituting the expression of $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})/f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)\big)$ and $\mathcal{L}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ from eqs. (A.30) and (A.31) into eq. (A.43) we obtain

$$V_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = \iiint \Bigg[\big(\mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{p}\bullet},t}\, h_{\mathrm{vl}}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{a}\bullet},t}\, h_{\mathrm{ol}}(\boldsymbol{x}_\bullet) + \chi_{\mathrm{k}} \big)^2$$
$$\times \left(1 + \xi_{k_{\mathrm{p}\bullet},t}\, \frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \right) \nu_{\chi_{\mathrm{k}}}(\chi_{\mathrm{k}} \mid \boldsymbol{x}_\bullet)\, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \Bigg]\, \mathrm{d}\chi_{\mathrm{k}}\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}. \quad \text{(A.44)}$$

Expanding the square in eq. (A.44), and noting that χ_{k} is centered at zero, we get

$$V_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = V_{\mathrm{l}}(\boldsymbol{x}_\bullet) + \iiint \Bigg[\big(\mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{p}\bullet},t}\, h_{\mathrm{vl}}(\boldsymbol{x}_\bullet) + \xi_{k_{\mathrm{a}\bullet},t}\, h_{\mathrm{ol}}(\boldsymbol{x}_\bullet) \big)^2$$
$$\times \left(1 + \xi_{k_{\mathrm{p}\bullet},t}\, \frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \right) \nu_{\chi_{\mathrm{k}}}(\chi_{\mathrm{k}} \mid \boldsymbol{x}_\bullet)\, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \Bigg]\, \mathrm{d}\chi_{\mathrm{k}}\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}, \quad \text{(A.45)}$$

where

$$V_{\mathrm{l}}(\boldsymbol{x}_\bullet) = \int \nu_{\chi_{\mathrm{k}}}(\chi_{\mathrm{k}} \mid \boldsymbol{x}_\bullet)\, \chi_{\mathrm{k}}^2\, \mathrm{d}\chi_{\mathrm{k}} = \lambda_\bullet^2\, \sigma_{\mathrm{i}}^2\, (1 - v_\bullet - o_\bullet). \quad \text{(A.46)}$$

Expanding the integral in eq. (A.45) and noting that $\xi_{k_{\mathrm{p}\bullet},t}$ and $\xi_{k_{\mathrm{a}\bullet},t}$ are centered around zero and independent, we find

$$V_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = V_{\mathrm{l}}(\boldsymbol{x}_\bullet) + h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)^2 \int \xi_{k_{\mathrm{p}\bullet},t}^2\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet} + h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)^2 \int \xi_{k_{\mathrm{a}\bullet},t}^2\, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \mathrm{d}k_{\mathrm{a}\bullet}$$
$$+ \big(\mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) \big)^2 + h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)^2\, \frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \int \xi_{k_{\mathrm{p}\bullet},t}^3\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}$$
$$+ 2 \big(\mathcal{L}\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right) - \overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet) \big)\, h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)\, \frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \int \xi_{k_{\mathrm{p}\bullet},t}^2\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}.$$
$$\text{(A.47)}$$

Noting that

$$\int \xi_{k_{\mathrm{p}\bullet},t}^2\, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet} = V_{\mathrm{a},t}(\boldsymbol{x}_\bullet), \quad \text{(A.48)}$$

$$\int \xi_{k_{\mathrm{a}\bullet},t}^2 \, \phi_{\mathrm{a},t}(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}}) \, \mathrm{d}k_{\mathrm{a}\bullet} = V_{\mathrm{a},t}(\overline{\boldsymbol{x}}), \tag{A.49}$$

and

$$\int \xi_{k_{\mathrm{p}\bullet},t}^3 \, \phi_{\mathrm{a},t}(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \, \mathrm{d}k_{\mathrm{p}\bullet} = C_{k^3,\mathrm{a},t}(\boldsymbol{x}_\bullet), \tag{A.50}$$

where $C_{k^3,\mathrm{a},t}(\boldsymbol{x}_\bullet)$ is the skewness in the knowledge held by a random adult from the $\boldsymbol{x}_\bullet$-lineage at generation t and by substituting the expression for $\overline{k}_{\mathrm{o},t}(\boldsymbol{x}_\bullet)$ from eq. (A.33) into eq. (A.47) we obtain

$$
\begin{aligned}
V_{\mathrm{o},t}(\boldsymbol{x}_\bullet) = V_{\mathrm{l}}(\boldsymbol{x}_\bullet) + h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)^2 \, V_{\mathrm{a},t}(\boldsymbol{x}_\bullet) + h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)^2 \, V_{\mathrm{a},t}(\overline{\boldsymbol{x}}) + \frac{h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)^2 \, C_{k^3,\mathrm{a},t}(\boldsymbol{x}_\bullet) \, \eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}} \, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \\
- \left(\frac{h_{\mathrm{vl}}(\boldsymbol{x}_\bullet) \, V_{\mathrm{a},t}(\boldsymbol{x}_\bullet) \, \eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}} \, \overline{k}_{\mathrm{a},t}(\boldsymbol{x}_\bullet)} \right)^2 .
\end{aligned}
\tag{A.51}
$$

Equations (A.41) and (A.51) give the dynamics of knowledge variance.

A.3.3   Gaussian closure approximation

Equations (A.21), (A.33), (A.41), and (A.51) describe the dynamics of the expected knowledge and knowledge variance in a lineage. However, tracking the expected knowledge and knowledge variance requires tracking knowledge skewness, which in turn necessitates tracking fourth-order moments of knowledge probability density. Since the dynamics of each moment depend on higher-order moments, tracking the cultural dynamics entails following the dynamics of the infinite sequence of moments. To circumvent this issue, we apply a Gaussian closure approximation, assuming that the probability density of knowledge within a lineage can be approximated by a Gaussian probability density. This assumption allows us to track the evolution of the probability density of knowledge across generations by tracking only its mean and variance.

Under the Gaussian closure approximation, the knowledge skewness is zero. By substituting $C_{k^3,\mathrm{a},t}(\boldsymbol{x}_\bullet) = C_{k^3,\mathrm{o},t}(\boldsymbol{x}_\bullet) = 0$ into the system of equations formed by eqs. (A.21), (A.33), (A.41)

and (A.51) we obtain

$$
\begin{cases}
\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet) = \overline{k}_{o,t}(\boldsymbol{x}_\bullet) + \frac{V_{o,t}(\boldsymbol{x}_\bullet)\,\eta_s}{\tilde{s}\big(\overline{k}_o^*(\boldsymbol{x}_\bullet)\big)} \\[2mm]
\overline{k}_{o,t}(\boldsymbol{x}_\bullet) = \mathcal{L}\big(\boldsymbol{x}_\bullet, \overline{k}_{a,t}(\boldsymbol{x}_\bullet), \overline{k}_{a,t}(\overline{\boldsymbol{x}})\big) + \frac{h_{vl}(\boldsymbol{x}_\bullet)\,V_{a,t}(\boldsymbol{x}_\bullet)\,\eta_f}{f_0 + \eta_f\,\overline{k}_{a,t}(\boldsymbol{x}_\bullet)} \\[2mm]
V_{a,t+1}(\boldsymbol{x}_\bullet) = V_{o,t}(\boldsymbol{x}_\bullet) - \left(\frac{V_{o,t}(\boldsymbol{x}_\bullet)\,\eta_s}{\tilde{s}\big(\overline{k}_o^*(\boldsymbol{x}_\bullet)\big)}\right)^2 \\[2mm]
V_{o,t}(\boldsymbol{x}_\bullet) = V_l(\boldsymbol{x}_\bullet) + h_{vl}(\boldsymbol{x}_\bullet)^2\,V_{a,t}(\boldsymbol{x}_\bullet) + h_{ol}(\boldsymbol{x}_\bullet)^2\,V_{a,t}(\overline{\boldsymbol{x}}) - \left(\frac{h_{vl}(\boldsymbol{x}_\bullet)\,V_{a,t}(\boldsymbol{x}_\bullet)\,\eta_f}{f_0 + \eta_f\,\overline{k}_{a,t}(\boldsymbol{x}_\bullet)}\right)^2 ,
\end{cases}
\tag{A.52}
$$

which characterized the dynamics of the expected knowledge and knowledge variance of the $\boldsymbol{x}_\bullet$-lineage member.

A.4   Probability density of knowledge at cultural equilibrium

In this section, we first characterize the equilibrium conditions for the expected knowledge and knowledge variance at cultural equilibrium (sections A.4.1 and A.4.2). Next, we demonstrate that in the absence of stochasticity in learning, there is no knowledge variance at the cultural equilibrium (section A.4.3). Finally, we describe the numerical method used to estimate the equilibrium values of mean knowledge and knowledge variance in the population (section A.4.4).

A.4.1   Characterization of the expected knowledge at cultural equilibrium

Here, we derive eq. (14) from the main text, which specifies the condition satisfied by the expected adults knowledge $\overline{k}_a^*(\boldsymbol{x}_\bullet)$ at equilibrium (here and hereafter the notation * denotes a variable at equilibrium).

At equilibrium, the expected knowledge and the variance in knowledge for randomly chosen adult and offspring remain unchanged across generations. By substituting $\overline{k}_{a,t+1}(\boldsymbol{x}_\bullet) = \overline{k}_{a,t}(\boldsymbol{x}_\bullet) = \overline{k}_a^*(\boldsymbol{x}_\bullet)$, $\overline{k}_{o,t}(\boldsymbol{x}_\bullet) = \overline{k}_o^*(\boldsymbol{x}_\bullet)$, $V_{a,t+1}(\boldsymbol{x}_\bullet) = V_{a,t}(\boldsymbol{x}_\bullet) = V_a^*(\boldsymbol{x}_\bullet)$ and $V_{o,t}(\boldsymbol{x}_\bullet) = V_o^*(\boldsymbol{x}_\bullet)$ into the dynamical system eq. (A.52), we obtain the following conditions that these equilibrium

values must satisfy

$$
\begin{cases}
\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) = \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) + \frac{V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big)} \\[2mm]
\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) = \mathcal{L}\big(\boldsymbol{x}_{\bullet}, \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), \overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})\big) + \frac{h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})} \\[2mm]
V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) = V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) - \left(\frac{V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big)}\right)^{2} \\[2mm]
V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) = V_{\mathrm{l}}(\boldsymbol{x}_{\bullet}) + h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})^{2}\,V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) + h_{\mathrm{ol}}(\boldsymbol{x}_{\bullet})^{2}\,V_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}}) - \left(\frac{h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})}\right)^{2},
\end{cases}
\tag{A.53}
$$

where $\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ and $V_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ are the population's mean and variance of adult knowledge, which verifies eq. (A.53) with $\boldsymbol{x}_{\bullet} = \overline{\boldsymbol{x}}$.

By replacing $\mathcal{L}\big(\boldsymbol{x}_{\bullet}, \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), \overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})\big)$ with its expression from eq. (A.9) (with $k_{\mathrm{p}\bullet} = \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})$ and $k_{\mathrm{a}\bullet} = \overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$) in the second line of eq. (A.53), we obtain

$$
\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) = h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) + h_{\mathrm{ol}}(\boldsymbol{x}_{\bullet})\,\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}}) + p_{\mathrm{il}}(\boldsymbol{x}_{\bullet}) + \frac{h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})}.
\tag{A.54}
$$

By substituting the expression for $\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})$ from eq. (A.54) into the first line of eq. (A.53), we obtain

$$
\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) = h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) + h_{\mathrm{ol}}(\boldsymbol{x}_{\bullet})\,\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}}) + p_{\mathrm{il}}(\boldsymbol{x}_{\bullet}) + \frac{h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})} + \frac{V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big)}.
\tag{A.55}
$$

This can be rewritten as

$$
\boxed{\ \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) = \frac{p_{\mathrm{il}}(\boldsymbol{x}_{\bullet}) + h_{\mathrm{ol}}(\boldsymbol{x}_{\bullet})\,\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}}) + \frac{h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})\,V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})} + \frac{V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big)}}{1 - h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})}.\ }
\tag{A.56}
$$

Equation (A.56) is eq. (14) from the main text.

A.4.2   Characterization of the knowledge variance at cultural equilibrium

Here, we derive eq. (15) from the main text, which specifies the condition satisfied by the variance in the knowledge held by a random adult $V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})$ at equilibrium. By substituting the expression for $V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})$ from the fourth line of eq. (A.53) into the third line of the same equation, we obtain

$$V_a^*(\boldsymbol{x_\bullet}) = V_l(\boldsymbol{x_\bullet}) + h_{vl}(\boldsymbol{x_\bullet})^2 \, V_a^*(\boldsymbol{x_\bullet}) + h_{ol}(\boldsymbol{x_\bullet})^2 \, V_a^*(\overline{\boldsymbol{x}})$$

$$- \left(\frac{h_{vl}(\boldsymbol{x_\bullet}) \, V_a^*(\boldsymbol{x_\bullet}) \, \eta_f}{f_0 + \eta_f \, \overline{k}_a^*(\boldsymbol{x_\bullet})} \right)^2 - \left(\frac{V_o^*(\boldsymbol{x_\bullet}) \, \eta_s}{\tilde{s}(\overline{k}_o^*(\boldsymbol{x_\bullet}))} \right)^2, \quad \text{(A.57)}$$

which can be rearranged as

$$V_a^*(\boldsymbol{x_\bullet}) = \frac{V_l(\boldsymbol{x_\bullet}) + h_{ol}(\boldsymbol{x_\bullet})^2 \, V_a^*(\overline{\boldsymbol{x}}) - \left(\frac{h_{vl}(\boldsymbol{x_\bullet}) \, V_a^*(\boldsymbol{x_\bullet}) \, \eta_f}{f_0 + \eta_f \, \overline{k}_a^*(\boldsymbol{x_\bullet})} \right)^2 - \left(\frac{V_o^*(\boldsymbol{x_\bullet}) \, \eta_s}{\tilde{s}(\overline{k}_o^*(\boldsymbol{x_\bullet}))} \right)^2}{1 - h_{vl}(\boldsymbol{x_\bullet})^2}. \quad \text{(A.58)}$$

By substituting the expression for $V_l(\boldsymbol{x_\bullet})$ from eq. (A.46) into eq. (A.58) gives eq. (15) from the main text.

A.4.3 Knowledge variance in the absence of stochasticity in learning

Here, we show that if individual learning is deterministic (i.e., $\sigma_i = 0$), the equilibrium variance of adult knowledge is zero (i.e., $V_a^*(\boldsymbol{x_\bullet}) = 0$).

We assume that, initially, all individuals have no knowledge, so that the knowledge variance is null: $V_0(\overline{\boldsymbol{x}}) = 0$. We aim to prove that this remains true for all generations t, i.e., $V_{a,t}(\overline{\boldsymbol{x}}) = 0$ for all t. Using mathematical induction, this is equivalent to showing that $V_{a,t}(\overline{\boldsymbol{x}}) = 0$ implies $V_{a,t+1}(\overline{\boldsymbol{x}}) = 0$.

Assume that $V_{a,t}(\overline{\boldsymbol{x}}) = 0$. Substituting $\sigma_i = 0$ into eq. (A.46), we obtain $V_l(\boldsymbol{x_\bullet}) = 0$. Then, substituting both $V_{a,t}(\overline{\boldsymbol{x}}) = 0$ and $V_l(\boldsymbol{x_\bullet}) = 0$ into the fourth line of eq. (A.52), we find that $V_{o,t}(\overline{\boldsymbol{x}}) = 0$. Finally, substituting $V_{o,t}(\overline{\boldsymbol{x}}) = 0$ into the third line of eq. (A.52) yields $V_{a,t+1}(\overline{\boldsymbol{x}}) = 0$. This completes the inductive step and shows that $V_{a,t}(\overline{\boldsymbol{x}}) = 0$ for all t, which implies that at equilibrium, $V_a^*(\boldsymbol{x_\bullet}) = 0$. This concludes the proof.

A.4.4 Numerical estimation of population's mean knowledge and variance in knowledge

To numerically determine the mean and variance of knowledge within the population at cultural equilibrium, $\overline{k}_a^*(\overline{\boldsymbol{x}})$ and $V_a^*(\overline{\boldsymbol{x}})$, we iterated the dynamics described by eq. (A.52) with $\boldsymbol{x_\bullet} = \overline{\boldsymbol{x}}$ across generations, starting from the initial conditions $\overline{k}_{a,1}(\overline{\boldsymbol{x}}) = V_{a,1}(\overline{\boldsymbol{x}}) = 0$. The expressions for $\tilde{s}(\overline{k}_{o,t}(\overline{\boldsymbol{x}}))$, $f(\overline{\boldsymbol{x}}, \overline{k}_{a,t}(\overline{\boldsymbol{x}}))$, $\mathcal{L}(\overline{\boldsymbol{x}}, \overline{k}_{a,t}(\overline{\boldsymbol{x}}), \overline{k}_{a,t}(\overline{\boldsymbol{x}}))$, $h_{vl}(\overline{\boldsymbol{x}})$, $h_{ol}(\overline{\boldsymbol{x}})$, $p_{il}(\overline{\boldsymbol{x}})$ and $V_l(\boldsymbol{x_\bullet})$ are given

by eqs. (4) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$ and $k_{\mathrm{o}\bullet} = \overline{k}_{\mathrm{o},t}(\overline{\boldsymbol{x}})$), (1) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$ and $k_{\mathrm{p}\bullet} = \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})$), (A.9) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$ and $k_{\mathrm{p}\bullet} = k_{\mathrm{a}\bullet} = \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})$), (A.10) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$), (A.11) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$), (A.12) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$) and (A.46) (with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$) respectively, treating $\overline{\boldsymbol{x}}$ as a parameter. We iterate this dynamics until $\sqrt{\left(\overline{k}_{\mathrm{a},t+1}(\overline{\boldsymbol{x}}) - \overline{k}_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right)^2 + \left(V_{\mathrm{a},t+1}(\overline{\boldsymbol{x}}) - V_{\mathrm{a},t}(\overline{\boldsymbol{x}})\right)^2} < 10^{-6}$.

Appendix B: Evolutionary dynamics

In Appendix B, we derive eq. (5) from the main text, which provides a simple expression for the lineage fitness (section B.1). We then derive eq. (17) from the main text, which establishes the conditions that $\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ and $\nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ satisfy (section B.2). Lastly, we detail the numerical procedure for computing the values of mean traits favored by selection (section B.3).

B.1   Lineage fitness

In this section, we detail how to obtain eq. (5) from the main text, which provides an expression for the lineage fitness $W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}})$ of an $\boldsymbol{x}_\bullet$-lineage introduced into a population with mean trait values $\overline{\boldsymbol{x}}$. The lineage fitness satisfies

$$\log(W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}})) = \lim_{t \to \infty} \frac{1}{t} \log(n_{\bullet,t}), \tag{B.1}$$

where the sequence $(n_{\bullet,t})_{t \geq 0}$ is a realization of a stochastic process, where $n_{\bullet,t}$ describes the random number of individuals in the $\boldsymbol{x}_\bullet$-lineage at time t following the introduction of the $\boldsymbol{x}_\bullet$-mutant (Cohen, 1979; Caswell, 2000). This number changes according to the recursion

$$n_{\bullet,t+1} = r_{\bullet,t}\, n_{\bullet,t}, \tag{B.2}$$

with initial condition $n_{\bullet,0} = 1$. Here, $r_{\bullet,t}$ is the realized growth rate, i.e., the average number of viable offspring produced by individuals in the $\boldsymbol{x}_\bullet$-lineage, t generations after the introduction of the $\boldsymbol{x}_\bullet$-mutant. This quantity depends on multiple factors, including the knowledge individuals carry, the number of offspring they produce, the learning outcomes of the offspring, and their survival based on their knowledge. Since offspring acquire knowledge from both their parent and randomly selected adults in the population, the probability density of $r_{\bullet,t}$ depends in part on the distribution of knowledge within the lineage and across the population in the parental generation.

Using eq. (B.2) with $t = h$, for $h \in \{0, \ldots, t-1\}$, we obtain

$$\log(W(\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})) = \lim_{t \to \infty} \frac{1}{t} \log \left(\prod_{h=0}^{t-1} r_{\bullet,h} \right), \tag{B.3}$$

$$= \lim_{t \to \infty} \frac{1}{t} \sum_{h=0}^{t-1} \log (r_{\bullet,h}). \tag{B.4}$$

We assume that the stochastic process underlying the lineage process $\boldsymbol{x}_{\bullet}$ growth reaches a stationary regime, such that $r_{\bullet,t}$ converges in distribution to a random variable $r_{\bullet}$. By applying the ergodic theorem for population growth (Caswell, 2000), eq. (B.3) can then be expressed as

$$\log(W(\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})) = \mathrm{E}\left(\log(r_{\bullet}) | \boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}}\right), \tag{B.5}$$

where E denotes the expectation under the stationary probability density of $r_{\bullet}$. It depends on the trait values within the lineage, $\boldsymbol{x}_{\bullet}$, as these directly influence learning and fecundity, and also determine the stationary probability density of knowledge within the lineage. It also depends on the population mean trait values $\overline{\boldsymbol{x}}$, which determine the equilibrium population size and the stationary probability density of knowledge in the population.

The variance in $r_{\bullet,t}$ stems from generational differences in the knowledge held by members of the lineage. We assume that the variability in knowledge is small, resulting in limited fluctuations in $r_{\bullet,t}$ at stationarity. Using a Taylor expansion of $\log(r_{\bullet})$ around $\mathrm{E}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})$, we obtain

$$\log(W(\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})) \approx \log\left(\mathrm{E}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})\right) - \frac{\mathrm{Var}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})}{2\,\mathrm{E}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})^2}, \tag{B.6}$$

where Var denotes the variance under the stationary probability density of $r_{\bullet,t}$. By neglecting the term proportional to $\mathrm{Var}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})$, we obtain

$$W(\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}}) \approx \mathrm{E}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}}). \tag{B.7}$$

While this is an approximation of lineage fitness, the estimated trait values reached at the evolutionary equilibrium using this approximation (see section B.3 for details) are consistent with those observed in individual-based simulations (see fig. S.9), suggesting that eq. (B.7) provides a reliable expression of lineage fitness.

The expected growth rate of the lineage $\mathrm{E}(r_{\bullet}|\boldsymbol{x}_{\bullet}, \overline{\boldsymbol{x}})$ equals the expected number of viable offspring for a random lineage member under the stationary distribution. Accordingly, the

lineage fitness $W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}})$ can be written as

$$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \int f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})\, \overline{s}_{\mathrm{o}}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}, \tag{B.8}$$

where $\phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)$ is the stationary probability density that an adult with traits $\boldsymbol{x}_\bullet$ carries knowledge value $k_{\mathrm{p}\bullet}$ at cultural equilibrium, and the integral is taken over all possible values of $k_{\mathrm{p}\bullet}$.

The term $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})$ (whose expression is given in eq. (1)) is the expected number of offspring produced by an adult with traits $\boldsymbol{x}_\bullet$ and knowledge $k_{\mathrm{p}\bullet}$. The term

$$\overline{s}_{\mathrm{o}}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}) = \iint \frac{\tilde{s}(k_{\mathrm{o}\bullet})}{1 + \gamma\, n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})}\, \nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})\, \phi^*(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \mathrm{d}k_{\mathrm{o}\bullet}\, \mathrm{d}k_{\mathrm{a}\bullet}, \tag{B.9}$$

is the expected probability of survival for a random offspring of an adult with traits $\boldsymbol{x}_\bullet$ and knowledge $k_{\mathrm{p}\bullet}$. The survival probability of a descendant offspring depends on its trait, which is $\boldsymbol{x}_\bullet$ in the $\boldsymbol{x}_\bullet$-lineage, and on its knowledge, $k_{\mathrm{o}\bullet}$, which is a random variable. To account for all possible survival outcomes, we must consider the full probability density of possible offspring knowledge values.

Since offspring's knowledge results from learning from an oblique exemplar, first we integrate over the probability density of the exemplar's knowledge $k_{\mathrm{a}\bullet}$. Given that the variances of population traits are small, we approximate the population as being quasi-monomorphic for the mean traits $\overline{\boldsymbol{x}}$. Consequently, we assume that offspring acquire knowledge obliquely from adults with traits $\overline{\boldsymbol{x}}$, which carry knowledge $k_{\mathrm{a}\bullet}$ with probability density $\phi^*(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})$. Then, for each possible oblique exemplar knowledge value $k_{\mathrm{a}\bullet}$, we integrate over all possible offspring knowledge values $k_{\mathrm{o}\bullet}$, weighted by the probability density $\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ that an offspring with trait $\boldsymbol{x}_\bullet$, parental knowledge $k_{\mathrm{p}\bullet}$, and exemplar knowledge $k_{\mathrm{a}\bullet}$ acquires knowledge $k_{\mathrm{o}\bullet}$. Since $k_{\mathrm{o}\bullet} = k_\bullet(1)$, where $k_\bullet(a)$ is a realization (or sample path) of the stochastic differential equation eq. (2), the probability density $\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ is entirely determined by eq. (2) (see, for example, the expression of $\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$ given in eq. (A.25) when $\sigma_{\mathrm{v}} = \sigma_{\mathrm{o}} = 0$). Finally, $n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})$ is the expected number of offspring produced at demographic and cultural equilibrium in a population with mean traits $\overline{\boldsymbol{x}}$.

Using the expression for $\tilde{s}(k_{\mathrm{o}\bullet})$ and $\overline{s}_{\mathrm{o}}(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})$ from eqs. (4) and (B.9), we rewrite eq. (B.8) as

$$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \int \left[f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}) \right.$$

$$\left. \times \left(s_0 + \frac{\eta_{\mathrm{s}}}{1 + \gamma\, n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})} \iint k_{\mathrm{o}\bullet}\, \nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})\, \phi^*(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \mathrm{d}k_{\mathrm{o}\bullet}\, \mathrm{d}k_{\mathrm{a}\bullet}\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet) \right) \right] \mathrm{d}k_{\mathrm{p}\bullet},$$
(B.10)

which can be rewritten as

$$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \frac{s_0 \int f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}}{1 + \gamma\, n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})}$$

$$+ \frac{\eta_{\mathrm{s}} \iiint f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})\, k_{\mathrm{o}\bullet}\, \nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})\, \phi^*(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{o}\bullet}\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}}{1 + \gamma\, n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})}. \quad (B.11)$$

Because $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})$ depends linearly on $k_{\mathrm{p}\bullet}$ and $\int \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet} = 1$ we have

$$\int f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet} = f\!\left(\boldsymbol{x}_\bullet, \int k_{\mathrm{p}\bullet}\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}\right). \quad (B.12)$$

By definition, $\int k_{\mathrm{p}\bullet}\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{p}\bullet}$ is the expected knowledge of a random adult at cultural equilibrium in the $\boldsymbol{x}_\bullet$-lineage, which we denote as $\overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)$. Using this notation and eq. (B.12), and factorizing by $f\!\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\right)/\left(1 + \gamma\, n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})\right)$, eq. (B.11) reduces to

$$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \frac{f\!\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\right)}{1 + \gamma\, n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})} \left[s_0 + \eta_{\mathrm{s}}\, \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet) \right], \quad (B.13)$$

where

$$\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet) = \iiint k_{\mathrm{o}\bullet}\, \nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})\, \phi^*(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})\, \frac{f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})}{f\!\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\right)}\, \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)\, \mathrm{d}k_{\mathrm{o}\bullet}\, \mathrm{d}k_{\mathrm{a}\bullet}\, \mathrm{d}k_{\mathrm{p}\bullet}. \quad (B.14)$$

The term $\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)$ is the expected knowledge of a random offspring once it has completed learning within the $\boldsymbol{x}_\bullet$-lineage at cultural equilibrium, taking into account the probability density of parental knowledge $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})/f\!\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\right) \phi^*(k_{\mathrm{p}\bullet} \mid \boldsymbol{x}_\bullet)$, the probability density of knowledge among potential oblique exemplars $\phi^*(k_{\mathrm{a}\bullet} \mid \overline{\boldsymbol{x}})$, and the probability density of stochastic learning outcomes $\nu(k_{\mathrm{o}\bullet} \mid \boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet}, k_{\mathrm{a}\bullet})$. Note that the probability density of parental knowledge is proportional to $f(\boldsymbol{x}_\bullet, k_{\mathrm{p}\bullet})/f\!\left(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\right)$ because lineage members with higher fecundity are overrepresented among parents and, consequently, among vertical exemplars.

By replacing $s_0 + \eta_{\mathrm{s}}\,\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)$ by $\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)$ (obtained using eq. (4) with $k_{\mathrm{o}\bullet} = \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)$) we obtain

$$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \frac{f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big)\,\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)}{1 + \gamma\,n_{\mathrm{o}}^*(\overline{\boldsymbol{x}})}. \tag{B.15}$$

By substituting $n_{\mathrm{o}}^*(\overline{\boldsymbol{x}}) = f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big)\,n_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ into eq. (B.15) we obtain

$$W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \frac{f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big)\,\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)}{1 + f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big)\,n_{\mathrm{a}}^*(\overline{\boldsymbol{x}})}. \tag{B.16}$$

To move forward, we derive an expression for $n_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$. At demographic equilibrium, each adult, on average, produces one viable offspring to replace itself, so the mean fitness in the population remains equal to 1. When trait variances are small, the mean fitness can be approximated by $W(\overline{\boldsymbol{x}}, \overline{\boldsymbol{x}})$, so $n_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ is characterized by

$$W(\overline{\boldsymbol{x}}, \overline{\boldsymbol{x}}) = 1. \tag{B.17}$$

By substituting the expression for $W(\overline{\boldsymbol{x}}, \overline{\boldsymbol{x}})$ from eq. (B.16) with $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$ into eq. (B.17) and solving for $n_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ we obtain

$$n_{\mathrm{a}}^*(\overline{\boldsymbol{x}}) = \frac{f\big(\overline{\boldsymbol{x}}, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})\big)\,\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}})\big) - 1}{\gamma\,f\big(\overline{\boldsymbol{x}}, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})\big)}, \tag{B.18}$$

where $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ and $\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}})$ are the expected knowledge at cultural equilibrium for a random adult and a random offspring after learning is complete in an $\overline{\boldsymbol{x}}$-lineage. Because trait variances are small, $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ and $\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}})$ are equal to the mean adult and offspring knowledge in the population at cultural equilibrium.

By substituting the expression for $n_{\mathrm{a}}^*(\overline{\boldsymbol{x}})$ from eq. (B.18) into eq. (B.16), we obtain

$$\boxed{W(\boldsymbol{x}_\bullet, \overline{\boldsymbol{x}}) = \frac{f\big(\boldsymbol{x}_\bullet, \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big)\,\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)}{f\big(\overline{\boldsymbol{x}}, \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}})\big)\,\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}})\big)},} \tag{B.19}$$

which is eq. (5) of the main text.

B.2 Learning traits effect on lineage knowledge

Here, we derive the conditions that $\nabla\overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ and $\nabla\overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}}$ satisfy.

We first derive eq. (17) from the main text, which give the conditions that $\nabla \overline{k}_a^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}$ satisfies. Differentiating both sides of eq. (A.55) with respect to the traits $\boldsymbol{x}_\bullet$ and evaluating at $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$, we obtain

$$
\begin{aligned}
\nabla \overline{k}_a^*(\boldsymbol{x}_\bullet)\Big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} = \\
h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\ \nabla \overline{k}_a^*(\boldsymbol{x}_\bullet)\Big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} + \nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \overline{k}_a^*(\overline{\boldsymbol{x}}) + \nabla h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \overline{k}_a^*(\overline{\boldsymbol{x}}) + \nabla p_{\mathrm{il}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} \\
+ \nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \frac{V_a^*(\overline{\boldsymbol{x}})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_a^*(\boldsymbol{x}_\bullet)} + h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\ \nabla \frac{V_a^*(\boldsymbol{x}_\bullet)\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_a^*(\boldsymbol{x}_\bullet)}\bigg|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} + \nabla \frac{V_o^*(\boldsymbol{x}_\bullet)\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_o^*(\boldsymbol{x}_\bullet)\big)}\bigg|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}. \quad\text{(B.20)}
\end{aligned}
$$

Equation (B.20) can be rewritten as

$$
\boxed{
\begin{aligned}
\nabla \overline{k}_a^*(\boldsymbol{x}_\bullet)\Big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} = \frac{1}{1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}})} \times \\
\bigg[\nabla p_{\mathrm{il}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} + \nabla h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \overline{k}_a^*(\overline{\boldsymbol{x}}) + \nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \left(\overline{k}_a^*(\overline{\boldsymbol{x}}) + \frac{V_a^*(\overline{\boldsymbol{x}})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_a^*(\boldsymbol{x}_\bullet)} \right) \\
+ h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\ \nabla \frac{V_a^*(\boldsymbol{x}_\bullet)\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_a^*(\boldsymbol{x}_\bullet)}\bigg|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} + \nabla \frac{V_o^*(\boldsymbol{x}_\bullet)\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_o^*(\boldsymbol{x}_\bullet)\big)}\bigg|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} \bigg],
\end{aligned}
}
\quad\text{(B.21)}
$$

which is eq. (17) of the main text.

We now derive the conditions that $\nabla \overline{k}_o^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}$ satisfies. Differentiating both sides of eq. (A.54) with respect to the traits $\boldsymbol{x}_\bullet$ and evaluating at $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}$, we obtain

$$
\begin{aligned}
\nabla \overline{k}_o^*(\boldsymbol{x}_\bullet)\Big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} = \\
h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\ \nabla \overline{k}_a^*(\boldsymbol{x}_\bullet)\Big|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} + \nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \overline{k}_a^*(\overline{\boldsymbol{x}}) + \nabla h_{\mathrm{ol}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \overline{k}_a^*(\overline{\boldsymbol{x}}) + \nabla p_{\mathrm{il}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}} \\
+ \nabla h_{\mathrm{vl}}(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}\ \frac{V_a^*(\overline{\boldsymbol{x}})\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_a^*(\boldsymbol{x}_\bullet)} + h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\ \nabla \frac{V_a^*(\boldsymbol{x}_\bullet)\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_a^*(\boldsymbol{x}_\bullet)}\bigg|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}}. \quad\text{(B.22)}
\end{aligned}
$$

By substituting the expression of $\overline{k}_a^*(\boldsymbol{x}_\bullet)$ from the first line of eq. (A.53) into eq. (B.22) and rearranging we obtain

$$
\boxed{
\begin{aligned}
\nabla \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\Big|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}} &= \frac{1}{1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}})} \times \\
&\Bigg[\nabla p_{\mathrm{il}}(\boldsymbol{x}_{\bullet})|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}} + \nabla h_{\mathrm{ol}}(\boldsymbol{x}_{\bullet})|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}}\, \overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}}) + \nabla h_{\mathrm{vl}}(\boldsymbol{x}_{\bullet})|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}} \left(\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}}) + \frac{V_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})\,\eta_{\mathrm{f}}}{f_{0} + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})} \right) \\
&+ h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\, \nabla \frac{V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{f}}}{f_{0} + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})}\Bigg|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}} + h_{\mathrm{vl}}(\overline{\boldsymbol{x}})\, \nabla \frac{V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big)}\Bigg|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}} \Bigg],
\end{aligned}
}
\tag{B.23}
$$

which gives a condition satisfied by $\nabla \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\Big|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}}$.

B.3   Numerical estimation of the average learning traits favored by selection

Let $\overline{\boldsymbol{x}}_{t}$ denote the mean trait values in generation t. The expected change in these mean trait values from one generation to the next is given by

$$
\Delta \overline{\boldsymbol{x}}_{t+1} = \overline{\boldsymbol{x}}_{t} + \boldsymbol{G} \cdot \boldsymbol{s}(\overline{\boldsymbol{x}}_{t}),
\tag{B.24}
$$

where $\boldsymbol{G}$ is the traits covariance matrix, assumed to be constant here (Iwasa et al., 1991; Mullon and Lehmann, 2019).

To numerically determine the mean trait values favored by selection $\overline{\boldsymbol{x}}^{*}$, we first need to obtain the full expression for the selection gradient $\boldsymbol{s}(\overline{\boldsymbol{x}}_{t})$. This requires deriving an exact expression for $\nabla \overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet})\Big|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}_{t}}$ and $\nabla \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\Big|_{\boldsymbol{x}_{\bullet}=\overline{\boldsymbol{x}}_{t}}$.

To simplify notation, we rewrite eq. (A.53) in functional form

$$
\begin{cases}
\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) = \mathcal{F}_{\overline{k}_{\mathrm{a}}^{*}}\big(\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big) \\[4pt]
\overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) = \mathcal{F}_{\overline{k}_{\mathrm{o}}^{*}}\big(\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big) \\[4pt]
V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}) = \mathcal{F}_{V_{\mathrm{a}}^{*}}\big(\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big) \\[4pt]
V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}) = \mathcal{F}_{V_{\mathrm{o}}^{*}}\big(\overline{k}_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), \overline{k}_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{a}}^{*}(\boldsymbol{x}_{\bullet}), V_{\mathrm{o}}^{*}(\boldsymbol{x}_{\bullet})\big),
\end{cases}
\tag{B.25}
$$

where the functions $\mathcal{F}_{\overline{k}_{\mathrm{a}}^{*}}$, $\mathcal{F}_{\overline{k}_{\mathrm{o}}^{*}}$, $\mathcal{F}_{V_{\mathrm{a}}^{*}}$ and $\mathcal{F}_{V_{\mathrm{o}}^{*}}$ are defined by the right-hand sides of eq. (A.53).

Thus, by differentiating both sides of each line in the system (B.25) with respect to $\boldsymbol{x}_{\bullet}$ and

evaluating at $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t$, we obtain the system

$$
\begin{cases}
\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} = \nabla \mathcal{F}_{\overline{k}_{\mathrm{a}}^*}\big(\overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} \\
\nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} = \nabla \mathcal{F}_{\overline{k}_{\mathrm{o}}^*}\big(\overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} \\
\nabla V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} = \nabla \mathcal{F}_{V_{\mathrm{a}}^*}\big(\overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} \\
\nabla V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t} = \nabla \mathcal{F}_{V_{\mathrm{o}}^*}\big(\overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet), V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}
\end{cases} . \tag{B.26}
$$

By solving eq. (B.26) for $\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}$, $\nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}$, $\nabla V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}$ and $\nabla V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}$ using the *Solve* function in Wolfram Mathematica 13.0.0, we obtain an exact expression for $\nabla \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}$ and $\nabla \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_t}$. These results enable us to derive an exact expression for the selection gradient $\boldsymbol{s}(\overline{\boldsymbol{x}}_t)$ (see the Mathematica Notebook in the supplementary material).

With the expression of the selection gradient in hand, we can now numerically track the evolution of mean traits across generations. We iterated the dynamics described by eq. (B.24) with $\boldsymbol{G} = 0.1\boldsymbol{I}$ where $\boldsymbol{I}$ is the identity matrix, starting from the initial conditions $\overline{\boldsymbol{x}}_1 = (0, 0, 0)$. Initially, the values of $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_1)$, $V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_1)$ are computed using the procedure described in section A.4.4, with $\overline{\boldsymbol{x}} = \overline{\boldsymbol{x}}_1$. For each subsequent generation $t > 1$, $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_t)$ and $V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_t)$ are recalculated using the same procedure, setting $\overline{\boldsymbol{x}} = \overline{\boldsymbol{x}}_t$ and initializing the procedure with the equilibrium values of $\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_{t-1})$ and $V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_{t-1})$ from the previous generation. We iterate this dynamics until $\sqrt{\left(\overline{v}_{t+1} - \overline{v}_t\right)^2 + \left(\overline{o}_{t+1} - \overline{o}_t\right)^2 + \left(\overline{\lambda}_{t+1} - \overline{\lambda}_t\right)^2} < 10^{-7}$.

B.4 Link to the findings of Maisonneuve et al. (2025)

Here, we present the connection with the findings of Maisonneuve et al. (2025). Maisonneuve et al. (2025) showed that oblique learning can evolve even if vertical learning is more efficient, because individuals can access knowledge from other adults that differs from their parents. Oblique learning evolves when the parameter ρ, defined in Maisonneuve et al. (2025) as the probability that two individuals from the same generation produce simultaneously different knowledge during individual learning, is low, thereby limiting overlap in adult knowledge. In our model, we explicitly control the degree of knowledge overlap among adults using the parameter ρ as well. Although ρ is not defined identically in both models, in both cases, low values of ρ correspond to low overlap in the knowledge held by adults. Consistent with previous findings, we show that when adult knowledge does not fully overlap (i.e., low value of ρ), oblique learning can evolve, as it enables individuals to acquire information unavailable from their parents.

Appendix C: Selection pressure on stochasticity in learning

In this section, we introduce and derive a condition for the emergence of an additional trait $s \in [0, +\infty)$ that affects stochasticity in individual learning $\sigma_i(s)$. We assume that $\forall s \in [0, +\infty)$, $\sigma_i'(s) > 0$, meaning that higher values of s correspond to greater stochasticity in learning. We also assume that $\sigma_i(0) > 0$, with $\sigma_i(0)$ remaining small, to reflect the idea that learning is inherently stochastic even in the absence of a specific trait promoting it.

Each individual is now characterized by a vector of traits $\boldsymbol{x}_\bullet = (v_\bullet, o_\bullet, \lambda_\bullet, s_\bullet)$. We study how selection may favor the emergence of the trait s that increases stochasticity in learning, starting from a population with mean trait vector $\overline{\boldsymbol{x}}_0 = (\overline{v}, \overline{o}, \overline{\lambda}, 0)$. A trait located at its lower boundary will emerge if selection favors an increase in its population mean value, that is, if the corresponding component of the selection gradient is positive. Using eq. (16) we obtain that the trait s emerge if

$$
\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\,\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\bigg|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0} + \frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\,\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\bigg|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0} > 0, \tag{C.1}
$$

which is eq. (18) from the main text.

To assess the emergence of s, we need to determine in the $\boldsymbol{x}_\bullet$-lineage the effect of $s_\bullet$ on the expected knowledge of adults and offspring. Using equations (B.21) and (B.23), we find that

$$
\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\bigg|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0} = \frac{1}{1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)} \times
$$

$$
\left[\frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\,\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0}\,\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)} - \frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\,V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\,\eta_{\mathrm{f}}^2\,\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0}}{\big(f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\big)^2} \right.
$$

$$
\left. + \frac{\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0}\,\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)} - \frac{V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\,\eta_{\mathrm{s}}^2\,\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\big|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)^2} \right], \tag{C.2}
$$

and

$$\left.\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} = \frac{1}{1-h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)} \times$$

$$\left[\frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ \left.\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)} - \frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\ \eta_{\mathrm{f}}^2\ \left.\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}}{\left(f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\right)^2}\right.$$

$$\left.+\ \frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ \left.\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)} - \frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\ \eta_{\mathrm{s}}^2\ \left.\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)^2}\right]. \quad \text{(C.3)}$$

Solving eqs. (C.2) and (C.3) for $\left.\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$ and $\left.\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$ we obtain

$$\left.\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} = \frac{1}{1-h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\left(1-\left(\frac{\eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^2 V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\right)\left(1-\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\right)}$$

$$\times\left[\frac{\left(1-\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\right) h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ \left.\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)} + \frac{\left.\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right], \quad \text{(C.4)}$$

and

$$\left.\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} = \frac{1}{1-h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\left(1-\left(\frac{\eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^2 V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\right)\left(1-\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\right)}$$

$$\times h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\left[\frac{\left.\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)} + \frac{\left(1-\left(\frac{\eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^2 V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\right) \left.\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right]. \quad \text{(C.5)}$$

Since $\sigma_{\mathrm{i}}(0)$ is small at $\overline{\boldsymbol{x}}_0$, the variance in knowledge of both adults and offspring is small in each lineage. In this regime, eqs. (C.4) and (C.5) simplify to

$$\boxed{\left.\frac{\partial \overline{k}_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} \approx \frac{1}{1-h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)}\left[\frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ \left.\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)} + \frac{\left.\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right],} \quad \text{(C.6)}$$

and

$$\boxed{\left.\frac{\partial \overline{k}_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial \boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} \approx \frac{1}{1-h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)} \times \left[\frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ \left.\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{f}}}{f_0+\eta_{\mathrm{f}}\ \overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)} + \frac{h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\ \left.\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}\ \eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right].}$$
$$\text{(C.7)}$$

Equation (C.6) is eq. (19) from the main text.

To proceed further, we determine the effect of $s_\bullet$ on the variance of knowledge held by a randomly chosen adult and offspring within the $\boldsymbol{x}_\bullet$-lineage. By differentiating the third and fourth lines of eq. (A.53) with respect to $s_\bullet$ and evaluating at $\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0$ we obtain

$$\left.\frac{\partial V_a^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} = \left.\frac{\partial V_o^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} - \frac{2\,\eta_s^2\,V_o^*(\overline{\boldsymbol{x}}_0)\,\left.\frac{\partial V_o^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}}_0)\big)^2} + \frac{2\,\eta_s^3\,V_o^*(\overline{\boldsymbol{x}}_0)^2\,\left.\frac{\partial\overline{k}_o^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}}_0)\big)^3},$$

(C.8)

and

$$\left.\frac{\partial V_o^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} = \left.\frac{\partial V_l(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} + h_{vl}(\overline{\boldsymbol{x}}_0)^2\,\left.\frac{\partial V_a^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$$

$$h_{vl}(\overline{\boldsymbol{x}}_0)^2\left[-\frac{2\,\eta_f^2\,V_a^*(\overline{\boldsymbol{x}}_0)\,\left.\frac{\partial V_a^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}}{\big(f_0 + \eta_f\,\overline{k}_a^*(\overline{\boldsymbol{x}}_0)\big)^2} + \frac{2\,\eta_f^3\,V_a^*(\overline{\boldsymbol{x}}_0)^2\,\left.\frac{\partial\overline{k}_a^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}}{\big(f_0 + \eta_f\,\overline{k}_a^*(\overline{\boldsymbol{x}}_0)\big)^3}\right]. \quad \text{(C.9)}$$

By substituting the expression of $\left.\frac{\partial\overline{k}_a^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$ and $\left.\frac{\partial\overline{k}_o^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$ from eqs. (C.6) and (C.7) into eqs. (C.8) and (C.9) and by solving for $\left.\frac{\partial V_a^*(\boldsymbol{x}_\bullet)}{\partial\boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$ and $\left.\frac{\partial V_o^*(\boldsymbol{x}_\bullet)}{\partial\boldsymbol{x}_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}$ we obtain

$$\left.\frac{\partial V_a^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} \approx \frac{1}{D}\left[h_{vl}(\overline{\boldsymbol{x}}_0)\left(2\left(\frac{\eta_s}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}}_0)\big)}\right)^4 V_o^*(\overline{\boldsymbol{x}}_0)^2 + 2\left(\frac{\eta_s}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_o^*(\overline{\boldsymbol{x}}_0) - 1\right)\right.$$

$$\left. - 2\left(\frac{\eta_s}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_o^*(\overline{\boldsymbol{x}}_0) + 1\right]\left.\frac{\partial V_l(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}, \quad \text{(C.10)}$$

and

$$\left.\frac{\partial V_o^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0} \approx \frac{1}{D}\left[1 - 2h_{vl}(\overline{\boldsymbol{x}}_0)\frac{\eta_f}{f_0 + \eta_f\,\overline{k}_a^*(\overline{\boldsymbol{x}}_0)}\left(\frac{\eta_s}{\tilde{s}\big(\overline{k}_o^*(\overline{\boldsymbol{x}}_0)\big)}\right)^3 V_o^*(\overline{\boldsymbol{x}}_0)^2 - h_{vl}(\overline{\boldsymbol{x}}_0)\right]$$

$$\times \left.\frac{\partial V_l(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet=\overline{\boldsymbol{x}}_0}, \quad \text{(C.11)}$$

where

$$D = 1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)\left[1 + 2\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^3 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)^2\right]$$

$$+ h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)^2\left[-2\left(\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^3\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right) V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)^2\right.$$

$$\left. + \left(\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^2 V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\left(2 - 4\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\right) + 2\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0) - 1\right]$$

$$h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)^3\left(1 - 2\left(\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^4 V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)^2 - 2\left(\frac{\eta_{\mathrm{f}}}{f_0 + \eta_{\mathrm{f}}\,\overline{k}_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)}\right)^2 V_{\mathrm{a}}^*(\overline{\boldsymbol{x}}_0)\right)$$

$$\times\left(1 - 2\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^4 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)^2 - 2\left(\frac{\eta_{\mathrm{s}}}{\tilde{s}\big(\overline{k}_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\big)}\right)^2 V_{\mathrm{o}}^*(\overline{\boldsymbol{x}}_0)\right). \quad \text{(C.12)}$$

Given that the variance in knowledge of both adults and offspring is small, and using the expression of $V_{\mathrm{l}}(\boldsymbol{x}_\bullet)$ from eq. (A.46) with $\sigma_{\mathrm{i}} = \sigma_{\mathrm{i}}(s_\bullet)$, we obtain

$$\boxed{\left.\frac{\partial V_{\mathrm{a}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0} \approx \left.\frac{\partial V_{\mathrm{o}}^*(\boldsymbol{x}_\bullet)}{\partial s_\bullet}\right|_{\boldsymbol{x}_\bullet = \overline{\boldsymbol{x}}_0} \approx \frac{2\overline{\lambda}^2\,\sigma_{\mathrm{i}}(0)\sigma_{\mathrm{i}}'(0)\,(1 - \overline{v} - \overline{o})}{1 - h_{\mathrm{vl}}(\overline{\boldsymbol{x}}_0)^2}.} \quad \text{(C.13)}$$

Since the right-hand side of eq. (C.13) is always positive, it shows that, in the $\boldsymbol{x}_\bullet$-lineage, the trait $s_\bullet$ consistently increases the variance of knowledge among randomly chosen adults and offspring within the lineage. Moreover, from eqs. (C.6) and (C.7), we infer that $s_\bullet$ also consistently increases their expected knowledge. Consequently, eq. (C.1) always holds, and selection systematically favors the emergence of the trait s.

Appendix D: Individual-based simulations

Our individual-based simulations track the evolution of the trait distribution, the distribution of individual knowledge, and the population size at each generation for a fixed number of generations following the life cycle described in the main text. Each individual i at each generation is characterised by the vector of its three traits (v_i, o_i, λ_i), as well as its knowledge k_i at the end of the learning period, the knowledge of its parent $k_{\mathrm{p,i}}$ and its oblique exemplar $k_{\mathrm{a,i}}$. At the first generation, the values of v_i, o_i, λ_i, $k_{\mathrm{p,i}}$ and $k_{\mathrm{a,i}}$ are by default all initialized to 0 for each individual $i \in \{1, \ldots, n_1\}$, where $n_1 = 100$ is the number of individuals generation 1.

At each generation $t \geq 1$, the following occurs:

(i) *Reproduction.* We first determine the fecundity f_i of each individual $i \in \{1, ..., n_t\}$, where n_t is the number of adult individuals at generation t. The fecundity of each individual i, is calculated using $f_i = f((v_i, o_i, \lambda_i), k_i)$, where the expression of $f((v_i, o_i, \lambda_i), k_i)$ is given in eq. (1) with $\boldsymbol{x}_\bullet = (v_i, o_i, \lambda_i)$ and $k_{p\bullet} = k_i$. Each individual i then produces a number of offspring that is sampled from a Poisson probability density with mean equal to f_i. In total, they produce a total number of offspring that we denote as $n_{o,t}$.

(ii) *Mutation.* With probability $1 - \mu$, an offspring inherits the same traits as its parent. Otherwise, with probability μ, traits mutate; we model this by adding an effect sampled from a normal probability density with mean 0 and variance σ^2 to each parental trait value. If necessary, the resulting trait values are truncated to remain in $\{(v, o, \lambda) : 0 \leq v \leq 1, 0 \leq o \leq 1, 0 \leq v + o \leq 1, 0 \leq \lambda \leq 1\}$. In all simulations, we set $\mu = 0.01$ and $\sigma = 0.01$.

(iii) *Learning.* For each offspring, i, an oblique exemplar is randomly sampled from the adults, with its knowledge denoted as $k_{a,i}$. The knowledge k_i of each offspring i after learning is complete, for $i \in \{1, ..., n_{o,t}\}$, is determined by $k_i = k_\bullet(1)$ where $k_\bullet(1)$ is computed by solving numerically the stochastic differential equation eq. (2) with $\boldsymbol{x}_\bullet = (v_i, o_i, \lambda_i)$, $k_{p\bullet} = k_{p,i}$ and $k_{a\bullet} = k_{a,i}$ using the *sdeint.itoint* function from the Python package *sdeint*. To reduce computation time when $\sigma_v = \sigma_o = 0$, we instead compute $k_\bullet(1)$ using eq. (A.5) with $\boldsymbol{x}_\bullet = (v_i, o_i, \lambda_i)$, $k_{p\bullet} = k_{p,i}$ and $k_{a\bullet} = k_{a,i}$, which provides an explicit expression for the final knowledge after learning, thereby avoiding numerical integration of the stochastic differential equation eq. (2).

(iv) *Survival.* All parents die and each offspring i in $\{1, ..., n_{o,t}\}$ then survives till adulthood with probability $s(k_i, n_{o,t})$ (and otherwise dies). The survival probability of each offspring i is calculated using the expression of $s(k_i, n_{o,t})$ given by substituting the expression of $\tilde{s}(k_i)$ from eq. (4) with $k_{o\bullet} = k_i$ into eq. (3) with $k_{o\bullet} = k_i$ and $n_o = n_{o,t}$. This results in the number n_{t+1} of adults of the next generation (with $n_{t+1} \leq n_{o,t}$).

We repeat steps (i)-(iv) for a fixed number of generations (see figure legends for parameter values).

References

Y. Kobayashi, H. Ohtsuki, and J. Y. Wakano. Population size vs. social connectedness — a gene-culture coevolutionary approach to cumulative cultural evolution. *Theoretical Population Biology*, 111:87–95, 2016.

L. Maisonneuve, L. Lehmann, and C. Mullon. The coevolution of learning schedules and teaching enhances cumulative knowledge and drives a teacher–innovator syndrome. *Proceedings of the Royal Society B: Biological Sciences*, 292(2040):20242470, 2025.

C. Gardiner. *Handbook of Stochastic Methods for Physics, Chemistry, and the Natural Sciences.* Proceedings in Life Sciences. Springer-Verlag, 1985.

R. F. Bass. *Stochastic Processes.* Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2011.

J. E. Cohen. Long-run growth rates of discrete multiplicative processes in Markovian environments. *Journal of Mathematical Analysis and Applications*, 69:243–251, 1979.

H. Caswell. *Matrix Population Models.* Sinauer Associates, Massachusetts, 2000.

Y. Iwasa, A. Pomiankowski, and S. Nee. The evolution of costly mate preferences II. The 'handicap' principle. *Evolution*, 45:1431–1442, 1991.

C. Mullon and L. Lehmann. An evolutionary quantitative genetics model for phenotypic (co)variances under limited dispersal, with an application to socially synergistic traits. *Evolution*, 73(9):1695–1728, 2019.

E Supplementary figures

Figure S.1: **Stationary distribution of adults knowledge from individual-based simulation.** The histogram shows the distribution of adult knowledge over 100,000 generations at equilibrium, based on individual-based simulations. The population was first allowed to evolve for 100,000 generations to ensure that both cultural and evolutionary equilibria were reached. The red curve represents the distribution that would be expected if adult knowledge were normally distributed, with the mean and variance of the Gaussian distribution matching those observed in the simulation, illustrating that a Gaussian approximation provides a good fit for the distribution of knowledge in the population. Parameters are: $f_0 = 5$, $s_0 = 1$, $\beta_v = 1.4$, $\beta_o = 1.3$, $\alpha = 0.1$, $\epsilon = 0.05$, $\rho = 0.05$, $\sigma_v = \sigma_o = 0$, $\sigma_i = 0.05$, $\eta_f = 25$, $\eta_s = 5$, $\gamma = 0.01$.

Figure S.2: **The impact of the conversion factors that translate knowledge into survival benefits η_s on the expected knowledge and variance in knowledge at cultural equilibrium for a random adult and offspring of the $\boldsymbol{x_\bullet}$-lineage. a** $\bar{k}_a^*(\boldsymbol{x_\bullet})$ (blue solid line) and $\bar{k}_o^*(\boldsymbol{x_\bullet})$ (blue dashed line) and **b** $V_a^*(\boldsymbol{x_\bullet})$ (blue solid line) and $V_o^*(\boldsymbol{x_\bullet})$ (blue dashed line) according to η_s. This shows that the expected knowledge in adults is slightly higher than in offspring, while the variance in adult knowledge is slightly lower than in offspring. Default parameters are the same as in fig. 2 with $\boldsymbol{x_\bullet} = \bar{\boldsymbol{x}}$.

Figure S.3: **The impact of stochasticity in individual learning on population size.** Equilibrium adult population size $n_a^*(\overline{\boldsymbol{x}})$ according to σ_i. This shows that the population size increases with stochasticity in individual learning. Default parameters are the same as in fig. 2 with $\gamma = 10^{-4}$.

Figure S.4: **Impact of selective pressure on the population mean knowledge.** Population mean knowledge at cultural equilibrium $\overline{k}_a^*(\overline{\boldsymbol{x}})$ according to η_f and η_s. This shows that the effect of selection in enhancing the mean knowledge $\overline{k}_a^*(\overline{\boldsymbol{x}})$ is stronger when η_f and η_s are higher. Default parameters are the same as in fig. 2.

Figure S.5: **Impact of transmission on the population mean knowledge.** Population mean **a** vertical cultural heritability $h_{\mathrm{vl}}(\overline{\boldsymbol{x}})$ and **b** knowledge at cultural equilibrium $\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ according to β_{v}. Population mean **c** oblique cultural heritability $h_{\mathrm{ol}}(\overline{\boldsymbol{x}})$ and **d** knowledge at cultural equilibrium $\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ according to β_{o}. This shows that the effect of selection in enhancing the mean knowledge $\overline{k}_{\mathrm{a}}^{*}(\overline{\boldsymbol{x}})$ is stronger when intergenerational transmission is higher. Default parameters are the same as in fig. 2.

Figure S.6: **Factors impacting the evolution of learning traits.** Learning schedule and investment in learning $\overline{\lambda}^*$ at $\overline{x}^*$ against **a** θ, **b** ϵ, **c** ρ, and **d-e** σ_i. Blue, green, and pink areas represent time spent performing vertical, oblique, and individual learning, respectively. Default parameters are the same as in fig. 3, except in panel d, where $\epsilon = 0.25$ and in panel e, where $\rho = 0.4$.

Figure S.7: **Impact of stochasticity in individual learning σ_i on investment in learning $\overline{\lambda}^*$ and the population mean knowledge $\overline{k}_a^*(\overline{x}^*)$ at $\overline{x}^*$ for different values of α.** This shows that stochasticity in learning can promote both higher average knowledge and greater investment in learning, except in populations with very low values of α. Default parameters are the same as in fig. 3.

Figure S.8: **Impact of stochasticity in individual learning σ_i on the average knowledge of parents $\overline{k}_a^*(\overline{x}^*) + V_a^*(\overline{x}^*)\,\eta_f / \left(f_0 + \eta_f\,\overline{k}_a^*(\overline{x}^*)\right)$ and of oblique exemplars $\overline{k}_a^*(\overline{x}^*)$ at $\overline{x}^*$.** This shows that under higher stochasticity in individual learning, the average knowledge of parents increases more than that of oblique exemplars. Default parameters are the same as in fig. 3.

Figure S.9: **Impact of stochasticity in individual learning on the stationary distribution of mean traits, average adults knowledge, and adult population size in individual-based simulations.** Box plots show distributions of **a** mean traits $\overline{x} = (\overline{v}, \overline{o}, \overline{\lambda})$ and **b** average adults knowledge $\overline{k}_a$ and adult population size n_a over 400,000 generations at equilibrium from individual-based simulations for different values of the strength σ_i of stochasticity in individual learning. For each value of σ_i, the population was first allowed to evolve for 100,000 generations to ensure equilibrium was reached. The box shows the interquartile range (IQR) with the median marked inside; whiskers extend to the smallest and largest values within $1.5 \times$ IQR from Q1 and Q3, respectively. Dashed lines show the predicted values derived from evolutionary analyses. The close match between simulation outcomes and analytical predictions suggests a good level of agreement between the two approaches. Default parameters are the same as in fig. 3 with $\sigma_v = \sigma_o = 0$ and $\gamma = 0.01$.

Figure S.10: **Impact of stochasticity in the different types of learning on the stationary distribution of mean traits, average adults knowledge, and adult population size in individual-based simulations.** Box plots show distributions of **a** mean traits $\overline{\boldsymbol{x}} = (\overline{v}, \overline{o}, \overline{\lambda})$ and **b** average adults knowledge $\overline{k}_a$ and adult population size n_a over 100,000 generations at equilibrium from individual-based simulations for different values of strength of stochasticity in vertical σ_v, oblique σ_o, and individual σ_i learning. For each combination of values for σ_v, σ_o, and σ_i, the population was first allowed to evolve for 100,000 generations to ensure equilibrium was reached. Because individual-based simulations run significantly slower when $\sigma_v > 0$ or $\sigma_o > 0$ (due to the need to numerically solve the stochastic differential equation eq. (2)), we initialize trait values for each individual i in the first generation as $(v_i, o_i, \lambda_i) = (0.13404, 0.0762715, 0.876229)$. These correspond to the mean trait values predicted at $\overline{\boldsymbol{x}}^*$ in the absence of stochasticity in learning (i.e., when $\sigma_v = \sigma_o = \sigma_i = 0$). The box shows the interquartile range (IQR) with the median marked inside; whiskers extend to the smallest and largest values within $1.5 \times$ IQR from Q1 and Q3, respectively. Dashed lines show the mean values obtained for individual-based simulation in the absence of stochasticity in learning. This shows that, similarly to individual learning, greater stochasticity in vertical and oblique learning promotes the accumulation of knowledge, an increase in population size, and a greater allocation of time to social learning. Default parameters are the same as in fig. 3 with $\gamma = 0.01$.