MBOAT7 and CDS2 coordinate lipid droplet dynamics and cholesterol metabolism in liver

Jiesi Xu[1, 2, 3, *], Jianxin Zhang[3, 4, *], Wei Wang[3], Honggang Su[3, 4], Siyu Chen[5], Mei Ding[2, 3] and Xun Huang[2, 3, 6, #]

1. The Key Laboratory of Geriatrics, Beijing Institute of Geriatrics, Institute of Geriatric Medicine, Chinese Academy of Medical Sciences, Beijing Hospital/National Center of Gerontology of National Health Commission, Beijing, 100730, China

2. State Key Laboratory of Metabolism and Regulation in Complex Organisms, Taikang Center for Life and Medical Sciences, School of Basic Medical Sciences, Wuhan University, Wuhan, Hubei 430071, China

3. Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

4. University of Chinese Academy of Sciences, Beijing 100049, China

5. Medical Research Center, Beijing Institute of Respiratory Medicine and Beijing Chao-Yang Hospital, Capital Medical University, Beijing 100020, China

6. Tianjian Laboratory of Advanced Biomedical Sciences, Zhengzhou University, Zhengzhou 450001, China

* Equal contribution

To whom correspondence should be addressed at: State Key Laboratory of Metabolism and Regulation in Complex Organisms, Taikang Center for Life and Medical Sciences, School of Basic Medical Sciences, Wuhan University, Wuhan, Hubei 430071, China

Email: huangxun@whu.edu.cn; xhuang@genetics.ac.cn

Abstract

Metabolic dysfunction-associated steatotic liver disease (MASLD) is a highly

prevalent disorder worldwide characterized by the accumulation of hepatic lipid

droplets (LDs). However, the mechanisms by which hepatocytes adapt to the dynamic

changes in LD morphology to maintain metabolic homeostasis remains poorly

understood. Here, we show that MBOAT7, a known MASLD-associated protein,

cooperates with CDS2 to regulate lipid droplet dynamics and cholesterol metabolism.

Under physiological conditions, their interaction in the endoplasmic reticulum (ER)

maintains metabolic homeostasis. Disruption of this interaction, such as through

CDS2 knockdown or loss of function, triggers an adaptive response wherein

MBOAT7 translocates to ER-LD contact sites in a RAB1-dependent manner. This

redistribution reduces LD size and activates the AMPK signaling, thereby promoting

cholesterol excretion. Conversely, impairment of this compensatory axis results in

hepatic cholesterol accumulation, a key driver of MASLD progression. Our findings

uncover a mechanism that integrates inter-organelle communication with metabolic

regulation, offering new insights into MASLD pathogenesis.

Introduction

Metabolic dysfunction-associated steatotic liver disease (MASLD) has emerged as the most prevalent chronic liver disorder worldwide, affecting 20%-25% of the general population (1, 2). Despite its significant clinical burden, the mechanisms that drive disease onset and progression remain incompletely understood, limiting the development of effective therapeutic strategies.

A central hallmark of MASLD is the excessive accumulation of hepatic lipid droplets (LDs), dynamic organelles that function as central hubs for neutral lipid storage and metabolism (3). Under pathological conditions, hepatocyte LDs can undergo substantial expansion in size, leading to macrovesicular steatosis. In response to lipid overload, hepatocytes can adapt by remodeling LD morphology toward small LDs. This shift is generally considered protective, as smaller LDs present a larger surface area-to-volume ratio and are more accessible for lipolysis and lipophagic degradation (4). Nevertheless, while small LDs are more metabolically active and facilitate lipid mobilization, their persistence can also increase the generation of lipotoxic intermediates. To mitigate potential lipotoxicity from sustained LD turnover, hepatocytes employ several compensatory mechanisms, such as directing free fatty acids toward oxidation, re-esterification, or export. Moreover, this adaptive process involves active remodeling of inter-organelle contacts between LDs and other organelles—such as the endoplasmic reticulum, mitochondria, and lysosomes—further integrating lipid handling with cellular metabolic homeostasis. Thus, LD remodeling represents a crucial adaptive process for maintaining hepatic metabolic homeostasis. Although LD-associated proteins are known to modulate LD size, their specific roles during the adaptive phase of hepatic steatosis are not fully understood.

The MBOAT7 (membrane-bound O-acyltransferase 7) variant rs641738 (C>T, loss of function mutation) has been robustly linked to all stages of MASLD, including steatosis, inflammation and fibrosis (5-8). MBOAT7 catalyzes the incorporation of arachidonic acid into phosphatidylinositol (PI) through the PI remodeling Lands' cycle (9). Parallel to this pathway, PI can also be synthesized *de novo* via the Kennedy

pathway, in which the conversion of phosphatidic acid to CDP-diacylglycerol

(CDP-DAG) by CDS (CDP-DAG synthase) constitutes a critical step (9, 10). Notably,

CDS2, one of the two CDS proteins in mammals, is selectively inhibited by

arachidonoyl-containing PI species (PI-AA), suggesting a regulatory feedback loop

that links MBOAT7-mediated PI remodeling with CDS2-mediated *de novo* PI

synthesis (11). This interplay is biologically significant: *Mboat7* deficiency has been

shown to promote hepatic lipid accumulation and steatohepatitis, partly through

increasing CDS2 activity (12, 13). Conversely, deletion of *Cds2* leads to the formation

of large LDs and spontaneous development of metabolic dysfunction-associated

steatohepatitis (MASH) (14-16), providing a valuable model for studying hepatocyte

adaptations to drastic LD morphological changes. Although MBOAT7 has been

implicated in modulating LD lipid composition (13, 17), its precise subcellular

localization and functional relationship with CDS2 in regulating LD dynamics remain

unclear.

Here, we uncover a coordinated mechanism in which MBOAT7 and CDS2 jointly

regulate LD dynamics and cholesterol metabolism. We show that deletion of CDS2

results in adaptive relocation of MBOAT7 to ER-LD contact sites via RAB1, where it

restricts LD expansion and activates the AMPK-LXR axis to promote cholesterol

excretion, thereby protecting hepatocytes from lipotoxicity. Conversely, disruption of

this regulatory circuit leads to cholesterol accumulation-a central feature of MASLD

progression. These findings establish MBOAT7 and CDS2 as key regulators of a

membrane-metabolic crosstalk that integrates inter-organelle communication with

metabolic control, revealing a robust metabolic adaptation to hepatic steatosis in

hepatocytes.

Results

MBOAT7 and CDS2 regulate cholesterol and fatty acid metabolism in liver

To investigate the connection between MBOAT7 and CDS2, we analyzed hepatic

transcriptome datasets and GEPIA (Gene Expression Profiling Interactive Analysis).

This showed that *MBOAT7* and *CDS2* expression levels in human liver samples are

positively correlated (Figure 1A-C). Both *MBOAT7* and *CDS2* are upregulated in liver

tumor samples compared with non-tumor liver samples (Figure S1A). To determine

the roles of MBOAT7 and CDS2 in the regulation of hepatic lipid metabolism, we

specifically knocked down these enzymes in the liver by injecting C57BL/6 mice with

AAV-TBG-sh*Mboat7*, AAV-TBG-sh*Cds2* or

AAV-TBG-sh*Mboat7*+AAV-TBG-sh*Cds2* (Figure 1D). Consistent with previous

report (18), knockdown of *Mboat7* in liver did not affect plasma lipid content but

increased hepatic triglyceride and cholesterol content (Figure 1E-F and Figure S1B-C).

Knockdown of *Cds2* did not affect plasma triglyceride content, but increased plasma

and hepatic cholesterol and hepatic triglyceride content (Figure 1E-F and Figure

S1B-C). Knockdown of both *Mboat7* and *Cds2* did not further enhance triglyceride

and cholesterol content (Figure 1E-F and Figure S1B-C), indicating that combined

knockdown of *Mboat7* and *Cds2* did not exert additive effects on the increased lipid

levels. Consistent with the above results, H&E staining showed that knockdown of

Mboat7, *Cds2*, and *Mboat7* + *Cds2* increased hepatic steatosis (Figure 1G-H).

    To examine the pathways regulated by MBOAT7 and CDS2, we performed RNA

sequencing (RNAseq) analysis on liver from the sh*Mboat7*, sh*Cds2*, and sh*Mboat7* +

sh*Cds2* groups. When we compared the results, we noticed an extensive overlap

between the three datasets: 266 genes were significantly upregulated and 301 genes

were significantly downregulated in all three groups (Figure 1I). Kyoto Encyclopedia

of Genes and Genomes (KEGG) analysis revealed significant changes in cholesterol

and fatty acid metabolism pathways among the genes with downregulated expression

in all three groups (Figure 1J). For the genes with upregulated expression in all three

groups, KEGG analysis revealed significant alterations in steroid biosynthesis and

non-alcoholic fatty liver pathways (Figure S1D). LXR, a key regulator of cholesterol

efflux, controls the transcription levels of genes such as *Abca1* and *Abcg5/8*. Our

qPCR analysis showed that genes involved in cholesterol excretion (*Lxr, Abcg8* and

Abca1) were downregulated in liver (Figure 1K), indicating that reduced expression

of *Mboat7* or *Cds2* impairs cholesterol excretion. Knockdown of *Mboat7* and *Cds2*

increased the expression of *Fasn*, which regulates fatty acid biosynthesis (Figure 1K).

Accordingly, LXR protein levels were reduced in the liver of sh*Mboat7*, sh*Cds2*, and

sh*Mboat7*+sh*Cds2* mice (Figure 1L and Figure S1E). Nuclear SREBP2 and HMGCR

levels were diminished, likely due to feedback inhibition of cholesterol biosynthesis

by increased cellular cholesterol levels (Figure 1L and Figure S1E). ALT and AST

measurements revealed that knockdown of *Mboat7* or *Cds2* or both did not exacerbate

liver inflammation (Figure S1F and G). Overall, these data suggests that MBOAT7

and CDS2 coordinate fatty acid and cholesterol metabolism.

MBOAT7 is localized to the ER-LD contacts in lipid-loaded CDS2 KO cells

To investigate how MBOAT7 and CDS2 cooperatively regulate lipid metabolism, we

focused on LDs. Previous studies have shown that CDS2 knockout increases LD size

(15). We examined the localization and roles of MBOAT7 in LD regulation.

MBOAT7-EGFP exhibited an ER-like localization pattern in both wild-type and

CDS2 KO HeLa cells (Figure S2A and B). Interestingly, in CDS2 KO cells treated

with oleic acid (OA) to induce LD formation, MBOAT7-EGFP (Figure 2A-B) and

MBOAT7-mCherry (Figure S2C-D) surrounded LDs stained by LipidTOX and

BODIPY, respectively. In contrast, MBOAT7-EGFP did not show colocalization with

LDs in OA-treated control HeLa cells (Figure 2A-B). Furthermore, in OA-treated

CDS2 KO cells, MBOAT7-EGFP was colocalized with the LD marker

mScarlet-PLIN1, whereas no such colocalization was observed in control cells

(Figure S2E). Similarly, in OA-treated HepG2 cells, MBOAT7-EGFP remained

localized to the ER, but enclosed LDs when CDS2 was knocked down (Figure S2F).

We also analyzed the localization of endogenous MBOAT7. In OA-treated CDS2 KO

cells, endogenous MBOAT7 colocalized with LDs, but this was not observed in

OA-treated control cells (Figure 2C-D). 3D reconstruction of the LD surface

confirmed that MBOAT7 localized to the LD surface in OA-treated CDS2 KO cells

(Figure 2E). Since MBOAT7 is a multipass transmembrane protein, it is likely

localized to the ER regions adjacent to LDs, rather than to the LD membrane.

The colocalization of MBOAT7 with LDs in CDS2 KO cells led us to test

whether CDS2 depletion affects the targeting of other ER proteins to LDs. The ER

marker mScarlet-SEC61b did not colocalize with LDs in OA-loaded CDS2 KO cells

(Figure 2F-G). Since CDS2 protein levels are downregulated in MASLD mouse

models (16), we also tested whether MBOAT7 targets to LDs in the liver of MASLD

model mice, in which MASLD is induced by the feeding with the

fat/fructose/cholesterol-rich Gubra-Amylin NASH (GAN) diet. Confocal imaging

showed that endogenous MBOAT7 was more enriched at the LD periphery in liver

sections from GAN diet-induced MASLD mice compared to those from normal mice

(Figure 2H-I). These findings suggest that MBOAT7 specifically localizes to the

ER-LD contacts in lipid-overloaded CDS2-deficient cells but not in control cells.

The N-terminus of MBOAT7 is required for localization of MBOAT7 to ER-LD

contacts in CDS2 KO cells

MBOAT7 consists of 11 transmembrane (TM) helices organized into two bundles

(TM1-5 and TM9-11) that tilt apart, creating a central catalytic chamber (19) (Figure

3A). To identify the region required for the localization of MBOAT7 to ER-LD

contacts, we generated three EGFP-tagged truncated versions of MBOAT7,

containing the N-terminal region (amino acids 1-135), the central region (180-310),

and the C-terminal region (311-472) (Figure 3A). Only the N-terminal region

localized to ER-LD contacts in CDS2 KO cells, and MBOAT7 with a deletion of the

N-terminal 99 amino acids did not localize to ER-LD contacts (Figure 3B). We further

investigated the specific TM regions critical for LD targeting in CDS2 KO cells.

Deleting TM1-2 slightly disrupted MBOAT7's ring-like pattern on LDs, while

deleting TM3-4 completely abolished its LD anchoring (Figure 3C). To validate the

importance of TM3-4 for ER-LD localization, we generated chimeric proteins by

swapping TM3 or TM4 with the corresponding regions from MBOAT5, which

belongs to the membrane-bound O-acyltransferase (MBOAT) family and carries out

re-acylation of phosphatidylserine (PS) and phosphatidylcholine (PC) on ER.

MBOAT7 and MBOAT5 have similar structure (19). These chimeric proteins failed to

localize to LDs (Figure 3D). Therefore, the N-terminus of MBOAT7, particularly

TM3-4, is required for the localization of MBOAT7 to ER-LD contacts in CDS2 KO

cells.

MBOAT7 interacts with CDS2

Next, we examined why MBOAT7 is localized to the ER-LD contacts in CDS2 KO

cells but not in control cells. Similar to MBOAT7, CDS2 is localized to the ER and

MAMs (16). We hypothesized that CDS2 retains MBOAT7 at the ER, while

MBOAT7 relocates to the ER-LD contacts in the absence of CDS2.

Co-immunoprecipitation (co-IP) assays demonstrated that Flag-CDS2 precipitated

MBOAT7-EGFP (Figure 3E), and conversely, MBOAT7-EGFP precipitated

Flag-CDS2 (Figure 3F). Using CRISPR-based knock-in (KI) of EGFP at the

N-terminus of endogenous CDS2, we found that endogenous CDS2 coprecipitated

with endogenous MBOAT7 (Figure 3G). To examine the direct binding between these

two proteins, we purified His-MBOAT7 from *E. coli* and Flag-CDS2 from

mammalian cells. Western blot analysis confirmed their direct interaction (Figure 3H).

Overall, these results indicate that CDS2 interacts with MBOAT7, retaining it at the

ER.

RAB1 is required for localization of MBOAT7 at ER-LD contacts in CDS2 KO
cells

Next, we investigated how MBOAT7 targets to the ER-LD contacts in CDS2 KO cells.

ER proteins typically target the LD monolayer surface, which requires membrane

continuity between the ER outer leaflet and the LD monolayer (20). This targeting can

occur either during early LD formation or to mature LDs after their formation. These

ER proteins associate with the LD monolayer through hydrophobic hairpins, which

insert midway into the lipid bilayers of the ER, with both the N- and C-termini facing

the cytosol (21). MBOAT7, however, is a multispanning transmembrane protein with

11 transmembrane domains, making it unlikely that MBOAT7 localizes directly to the

LD monolayer. Since MBOAT7 lacks a hairpin structure, we hypothesized that

ER-localized MBOAT7 associates with LDs in CDS2 KO cells via interaction with

LD-associated proteins.

We immunoprecipitated MBOAT7-EGFP from control and CDS2 KO cells and identified its interacting proteins using LC-MS/MS (Figure 4A). We hypothesized that MBOAT7 would interact with LD proteins in CDS2 KO cells but not in control cells, so we selected candidates whose interaction with MBOAT7 in CDS2 KO cells was at least 2-fold stronger than in control cells. Based on previously published LD proteomes (22-24), we focused on LD proteins. Five proteins from our dataset were shared with all the published LD proteomes (Figure 4B). Interestingly, of these five candidates, four of them are RAB proteins (RAB1A, RAB7A, RAB14 and RAB18) (Figure 4C). RAB1 mediates targeting triglyceride biosynthetic enzymes to LDs (25, 26). Our results showed that knockdown of RAB1A impaired the LD targeting of MBOAT7 in OA-loaded CDS2 KO cells (Figure 4D-F). Co-IP assays showed that MBOAT7-EGFP interacted with Flag-RAB1A in CDS2 KO cells but not in control cells, suggesting that RAB1A helps MBOAT7 target to the ER-LD contacts in CDS2 KO cells (Figure 4G). Furthermore, expressing HA-tagged CDS2 in CDS2 KO cells inhibited the interaction between MBOAT7 and RAB1A (Figure 4H). Together, these results suggest that RAB1A is essential for targeting of MBOAT7 to the ER-LD contacts in CDS2 KO cells, and that CDS2 inhibits this targeting by competing with RAB1A for binding to MBOAT7.

MBOAT7 reduces LD size in CDS2 KO cells by inhibiting the targeting of DGAT2 to LDs

We next investigated the cellular and physiological functions of ER-LD contact-localized MBOAT7. Overexpression of MBOAT7-EGFP did not affect LD size in control cells, but it reduced LD size in CDS2 KO cells (Figure 5A-B). Overexpression of an MBOAT7 mutant with reduced catalytic activity (MBOAT7(N321A, H356A)-EGFP) still reduced the LD size in CDS2 KO cells (Figure 5A-B). This indicates that the effect of MBOAT7 on LD size reduction in CDS2 KO cells is independent of its enzymatic activity.

To explore the underlying mechanism, we considered previous findings suggesting that the lipid biosynthetic enzymes DGAT2 and GPAT4 target LDs earlier

in CDS2 KO cells than in control cells, promoting LD growth (14). Knockdown of

DGAT2 or GPAT4 reduced the LD size in CDS2 KO cells (14). We hypothesized that

MBOAT7 may impede the LD targeting of DGAT2 or GPAT4 by competitively

interacting with RAB1, thus limiting LD growth in CDS2 KO cells. Consistent with

this idea, we found that both Flag-DGAT2 and Flag-GPAT4 bound to GFP-tagged

RAB1A (Figure 5C and D). The interaction between GPAT4 and RAB1A was

unaffected by MBOAT7 overexpression (Figure S3A). Importantly, MBOAT7-EGFP

expression reduced the binding between Flag-DGAT2 and GFP-RAB1A in CDS2 KO

cells (Figure 5E). Conversely, knockdown of MBOAT7 enhanced the binding between

Flag-DGAT2 and GFP-RAB1A in CDS2 KO cells (Figure 5F). Confocal images

showed that EGFP-DGAT2 was targeted to LDs in CDS2 KO cells, but

overexpression of MBOAT7-mCherry blocked this targeting (Figure 5G). Both

Flag-DGAT2 and Flag-MBOAT7 interact with full-length GFP-RAB1A. However,

their interactions are markedly reduced when RAB1A is truncated at either the

N-terminus or the C-terminus (Figure 5H-I). In response to OA treatment, triglyceride

synthesis enzymes like DGAT2 or GPAT4 are more likely to target to LDs to promote

LD growth (27). Since this process requires RAB1, we then tested whether OA

treatment affects the interaction of these enzymes with RAB1. OA treatment enhanced

DGAT2 binding to RAB1A but did not affect GPAT4 binding to RAB1A (Figure 5J

and Figure S3B). Interestingly, MBOAT7 binding to RAB1A increased after OA

treatment in CDS2 KO cells (Figure 5K). Taken together, our data suggest that CDS2

retains MBOAT7 at the ER. When CDS2 is absent, MBOAT7 binds to RAB1 and

localizes to the ER-LD contacts. Furthermore, MBOAT7 reduces the size of LDs by

inhibiting the targeting of DGAT2 to LDs.

MBOAT7 regulates LD size and lipolysis in the absence of CDS2

The studies above reveal the regulatory roles of MBOAT7 and CDS2 in controlling

LD size in cultured cells. We next examined how MBOAT7 and CDS2 functionally

interact and regulate LD size *in vivo*. Since MBOAT7 and CDS2 are bound to each

other, we tested whether CDS2 affects the acyltransferase activity of MBOAT7 using

MBOAT7 substrates like lyso-PI and [14C]-arachidonic acid. We analyzed the activity of MBOAT7 in the liver from control mice (*Cds2^f/f*), mice with liver-specific depletion of CDS2 (*Cds2^f/f;AlbCre*), and hepatic CDS2-depleted mice with AAV-mediated hepatic re-expression of CDS2 (*Cds2^f/f;AlbCre* + AAV-CDS2). The results showed that CDS2 deficiency did not significantly impact the enzymatic activity of MBOAT7 (Figure 6A).

To determine how MBOAT7 and CDS2 regulate LDs *in vivo*, *Cds2^f/f* and *Cds2^f/f;AlbCre* mice were injected with AAV-TBG-Con or AAV-TBG-*Mboat7* (Figure 6B). Hepatic overexpression of *Mboat7* did not affect plasma lipids in either control or *Cds2^f/f;AlbCre* mice (Figure S4A-B). Hepatic overexpression of *Mboat7* did not change hepatic triglyceride and cholesterol levels in *Cds2^f/f* mice, but it reduced hepatic cholesterol levels in *Cds2^f/f;AlbCre* mice (Figure 6C and D). H&E staining showed that liver sections from *Cds2^f/f;AlbCr*e mice contained "supersized" LDs, while overexpression of *Mboat7* reduced the LD size in *Cds2^f/f;AlbCre* mice, resulting in multiple small-sized LDs in hepatocytes (Figure 6E). This reduction in LD size was also confirmed in *Cds2*-deficient primary hepatocytes (Figure 6F and G). Cell fractionation assays revealed that overexpression of *Mboat7* reduced the protein level of DGAT2 in the LD fraction from the livers of *Cds2^f/f;AlbCre* mice, while increasing the protein level of ATGL (also called PNPLA2) (Figure 6H), a rate-limiting enzyme for lipolysis. To test if MBOAT7 regulates lipolysis, we used isoprenaline (ISO) to stimulate lipolysis. The results showed that overexpression of MBOAT7 increased free fatty acid (FFA) release in both DMSO and ISO-treated *Cds2*-deficient primary hepatocytes (Figure 6I).

Overall, our data suggest that MBOAT7 controls LD size and lipolysis by inhibiting DGAT2 targeting to LDs and promoting ATGL targeting to LDs in both CDS2-deficient cell lines and primary hepatocytes.

MBOAT7 and CDS2 regulate cholesterol metabolism through the AMPK pathway

We next investigated the physiological significance of ER-LD contact-localized

MBOAT7. The hepatic RNAseq results showed that knockdown of *Mboat7* reduced

the expression levels of genes in the liver X receptor (LXR) pathway, suggesting

impaired cholesterol excretion (Figure 1K-L). Heatmap analysis further confirmed the

downregulation of genes involved in cholesterol excretion (Figure S4C). We then

explored how cholesterol excretion is regulated by MBOAT7 and CDS2. Activation of

the AMPK pathway is known to inhibit SREBP and enhance LXR protein levels (28).

Interestingly, AMPK signaling was downregulated in the liver of sh*Mboat7*, sh*Cds2*,

or sh*Mboat7* + sh*Cds2* mice, as indicated by heatmap analysis and western blotting

(Figure S4C-E). In contrast, overexpression of *Mboat7* increased p-AMPK and

p-ACC levels in *Cds2^{f/f};AlbCre* mice, suggesting that AMPK pathway activity was

enhanced (Figure 7A-B). This was accompanied by reduced nuclear SREBP2 levels

and increased expressions of LXR and its target ABCA1 in *Cds2^{f/f};AlbCre* mice

(Figure 7A-B). Accordingly, mRNA analysis revealed that downstream targets of

LXR (*Abca1*, *Abcg5* and *Abcg8*) were upregulated in the liver of

Mboat7-overexpressing *Cds2^{f/f};AlbCre* mice (Figure S4F). To confirm whether

MBOAT7 regulates cholesterol excretion via the AMPK pathway in *Cds2^{f/f};AlbCre*

mice, we treated primary hepatocytes from *Cds2^{f/f};AlbCre* mice with Ad-EGFP and

Ad-MBOAT7-EGFP and the AMPK inhibitor dorsomorphin. The results showed that

dorsomorphin blocked the ability of *Mboat7* to elevate LXR protein levels (Figure

7C). This demonstrates that MBOAT7 regulates cholesterol excretion via the

AMPK-LXR axis.

We then explored how overexpression of MBOAT7 increases AMPK activity in

Cds2^{f/f};AlbCre mice. Lipolysis regulates key intracellular nutrient-sensing pathways,

such as the AMPK pathway (29). Since MBOAT7 overexpression promotes lipolysis

in *Cds2*-deficient primary hepatocytes (Figure 6I), we examined this process in

primary hepatocytes. Activation of lipolysis with isoprenaline (ISO) enhanced AMPK

activity in wild-type primary hepatocytes under both Ad-EGFP and

Ad-MBOAT7-EGFP conditions (Figure S4G-H). Conversely, inhibiting lipolysis with

the ATGL inhibitor Atglistatin (ATGLi) reduced AMPK signaling (Figure S4I-J),

which demonstrates that lipolysis regulates AMPK signaling in primary hepatocytes.

In *Cds2*-deficient primary hepatocytes, overexpression of MBOAT7 activated AMPK

signaling, which was further enhanced by ISO treatment (Figure 7D-E). Importantly,

ATGLi treatment diminished AMPK activation caused by MBOAT7 overexpression

in *Cds2*-deficient primary hepatocytes (Figure 7F-G), demonstrating that MBOAT7

regulates AMPK activity at least partially by promoting lipolysis. These findings

suggest that *Mboat7* reduces LD size and cholesterol content in *Cds2*-deficient

conditions by activating the AMPK-LXR axis.

Discussion

Here, we elucidate that spatial localization of MBOAT7 to ER-LD contact

sites-orchestrated by CDS2-represents a fundamental cellular adaptation to lipid

overload. This redistribution is critical for its role in regulating LD dynamics and

cholesterol metabolism. At the ER, CDS2 binds to MBOAT7, but in the absence of

CDS2, MBOAT7 locates to the ER-LD contacts. This relocation promotes lipolysis

and inhibits the targeting of DGAT2 to LDs in a RAB1-dependent manner.

Physiologically, MBOAT7-mediated LD size reduction activates the AMPK-LXR axis,

enhancing cholesterol excretion (Figure 7H).

LDs play an essential role in adaptation to fluctuating metabolic states, through

controlling their size, subcellular distribution and interaction with other organelles

(30). In response to lipid overload, hepatocytes sequester free fatty acid as triglyceride

within the core of LDs to prevent lipotoxicity and oxidative stress. Although the LD

size varies considerably across hepatic sinusoids, the physiological relevance of this

LD heterogeneity is largely undetermined. LD proteins can integrate nutrient signals

and regulate LD size (31), suggesting the existence of an adaptive mechanism that

maintains LD dimensions within a degradable range and minimizes lipotoxicity. Our

study identifies MBOAT7 and CDS2 as key coordinators of LD size and cholesterol

homeostasis. In MASLD mouse livers, MBOAT7 is more concentrated on LDs

compared to normal controls, indicating a compensatory redistribution that

counteracts steatosis-driven LD expansion. The translocation of MBOAT7 from the

ER to ER-LD contacts in the absence of CDS2 depends on RAB1A, consistent with

established mechanisms involving Rab GTPases, tethering complexes, and SNAREs

in ER-to-LD protein trafficking (18-20).

Previous evidence showed that CDS2 deficiency resulted in "supersized" LD. We

also observed large LD in *Cds2* deficient hepatocytes (15, 16). We further

demonstrate that MBOAT7 expression normalizes LD size in CDS2-KO cells by

preventing DGAT2 localization to LDs via a RAB1-dependent mechanism. This

process reflects an adaptive response orchestrated by the MBOAT7–RAB1–DGAT2

axis to restrict LD expansion. Of note, MBOAT7 antagonizes RAB1-mediated

DGAT2 recruitment to LDs. However, this only occurs when CDS2 is absent,

suggesting that RAB1 has a stronger binding affinity for DGAT2 than for MBOAT7.

Interestingly, GPAT4-mediated LD growth remains unaffected in CDS2 KO cells,

highlighting the specificity of this regulatory node.

In *Cds2* deficient condition, the compensatory redistribution of MBOAT7 to

ER-LD contacts exerts significant physiological effects. It stimulates ATGL-mediated

lipolysis, which reduces LD size and elevates cytosolic free fatty acids and cholesterol.

Hepatocytes counteract the resulting lipotoxic stress by activating export pathways.

We showed that reduced LD size-regulated by ER-LD localized MBOAT7- triggers an

energy-consuming futile cycle of triglyceride hydrolysis and resynthesis, raising AMP

levels and activating AMPK signaling (29). This in turn enhances fatty acid oxidation

and cholesterol excretion, which helps remove toxic free fatty acids and free

cholesterol, thereby safeguarding liver health. Given the strong association between

MBOAT7 loss-of-function variants and MASLD across various populations (32, 33),

therapeutic activation of AMPK may represent a promising strategy for the MBOAT7

deficiency-related MASLD.

In summary, our findings reveal a previously unrecognized axis of inter-organelle

communication and metabolic control, orchestrated by MBOAT7, CDS2, and RAB1

that adaptively responds to lipid overload. We demonstrate that LD size directly

regulates hepatic cholesterol content through AMPK-LXR signaling, providing a new

mechanistic link between organelle membrane dynamics and metabolic homeostasis

in MASLD pathogenesis.

Material and Methods

Mice

Cds2^{f/f} and *Cds2^{f/f};AlbCre* mice were generated as described (16). C57BL/6J mice were purchased from GemPharmatech Co., Ltd. For liver-specific gene knockdowns, C57BL/6J mice were injected with AAV-TBG-sh*Con*, AAV-TBG-sh*Mboat7*, AAV-TBG-sh*Cds2* intravenously (i.v) with 3×10^{11} genomic copies of AAVs. The target sequence for *Mboat7* is 5'-CCTGTTGGCTTCCTCTTTAAG-3'. The target sequence for *Cds2* is 5'-GTCCGAAGCAAAGCTAGAT-3'. The control sequence is 5'-GAAGTCGTGAGAAGTAGAA-3'. For liver-specific overexpression of *Mboat7*, C57BL/6J mice were injected with AAV-TBG-*Con* or AAV-TBG-*Mboat7* intravenously (i.v) with 1×10^{11} genomic copies of AAVs. Mice were housed at room temperature (22°C) and fed a standard rodent chow diet (Beijing HFK Bioscience Co., Ltd #1032). Unless otherwise stated, 8-week-old male mice were used and all mice were fasted for 5-6 hours prior to euthanization. In some experiments, C57BL/6J mice were fed a chow or GAN diet (Research Diets Inc, D09100310) for 26 weeks. All animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at the Institute of Genetics and Developmental Biology, Chinese Academy of Sciences (IGDB-CAS).

Cell culture

The *CDS2* knockout HeLa cell line was generated using the CPISPR/Cas9 system according to published literature (34). The target sequence was 5'-TATTTACTGAGAATCCGCAA-3'. HepG2 cells (ATCC, HB-8065) and HeLa cells were cultured in DMEM with 10% FBS. In certain experiments, HeLa cells were transfected with siRNAs. The targeting sequence for RAB1A is 5'-CAGCATGAATCCCGAATAT-3'. The targeting sequence for CDS2 is 5'-GGCCACCTTTGTCAATGTA-3'. Mouse primary hepatocytes were isolated from *Cds2^{f/f}* and *Cds2^{f/f};AlbCre* mice and infected with Ad-EGFP or Ad-MBOAT7-EGFP with MOI=50. All cells were grown at 37°C in a 5% CO_2 incubator.

Plasmids

Human *MBOAT7* cDNA was cloned into the *pcDNA3.1-EGFP* and *pCMV-C-mCherry* vectors. Human *MBOAT5* was cloned into the *pcDNA3.1-EGFP* vector. Human *CDS2* cDNA was cloned into the *pCMV-Tag2b* and *pCS2-HA* vectors. Human *RAB1A*, *DGAT2* and *GPAT4* cDNAs were cloned into both the *pEGFP-N1* and *pCMV-Tag2b* vectors. For His pulldown assays, human *MBOAT7* cDNA was cloned into the *pET-28a* vector.

MBOAT7 activity assay

The activity of MBOAT7 was measured as described with minor modifications (19). Briefly, 100 μg of protein from liver was incubated in 200 μl of medium (10 mM Tris/HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA) with 50 μM lysophosphatidylinositol (LPI), 25 μM arachidonoyl-CoA (20:4-CoA), and 5 μM (0.2 μCi) radiolabeled arachidonoyl[1-14C]CoA. The reaction was incubated for 15-30 minutes at 37 ºC and then stopped by adding 1 mL of chloroform:methanol (2:1, vol/vol) and 0.4 mL of 2% orthophosphoric acid. After centrifugation (3,500 g, 15 min, 4 ºC), the lower organic phase, containing the lipids, was collected and dried under a gentle nitrogen stream. Lipids were reconstituted in chloroform. Total lipids were extracted using the Bligh and Dyer method and separated by one-dimensional TLC on silica gel 60 plates using chloroform/methyl acetate:1-propanol:methanol:0.25% KCl (25:25:25:10:9, vol/vol). The radioactivity in the separated bands were quantified using a Typhoon FLA 9000 (GE Healthcare Japan). The corresponding areas of radiolabeled phospholipids on TLC plates were cut out, and the radiolabeled products were measured by liquid scintillation counting.

Lipid analysis

Lipids were extracted based on the chloroform/methanol (2:1, v/v) lipid extraction method (3). In brief, 50 mg of liver tissue were extracted with 6 mL of chloroform/methanol (2:1, v/v), then vortexed and refrigerated overnight. 3 mL of 0.043% magnesium chloride was then added to separate the aqueous phase from the chloroform phase which contains the lipids. After separation of the two phases, the upper water/methanol phase was removed. A 0.5-mL aliquot of the bottom phase was combined with 1 mL of 0.5% Triton X-100 in chloroform and dried. Triglyceride and

total cholesterol levels were measured using commercially available kits (BioSino, Beijing, China). Plasma free fatty acids (FFA) were measured using commercially available kits (Sigma-Alrich, MAK044).

Immunoprecipitation

Cells were transfected with the indicated plasmids using Lipofectamine 3000 (ThermoFisher, L3000001). Proteins were extracted using RIPA lysis buffer, containing100 mM Tris Base, 1% deoxycholate, 0.1% SDS, 5 mM EDTA, 1% Triton-X-100, 150 mM NaCl, pH 8.0), supplemented with 100 mM PMSF, protease inhibitor cocktail (Roche, P8340) and 100 mM sodium vanadate. Lysates were incubated on ice for 15 min, then centrifuged at 12,000 rpm for 15 minutes at 4 °C. Supernatants were incubated overnight at 4 °C with anti-Flag M2 affinity gel (Sigma-Aldrich, A2220) or anti-GPF agarose beads (Lablead Biotech Co., Ltd). After four washes with lysis buffer, beads were denatured in 2X SDS sample buffer for 10 min at 100 °C for western blotting. For the pulldown assay, MBOAT7-His was pulled down using Ni-NTA agarose beads (Qiagen, 30210), then washed three times with PBS + PMSF + 1% Triton buffer. 10 μg of *Flag-CDS2* was expressed in 293T cells, immunoprecipitated using anti-Flag M2 affinity gel, and eluted with 3X Flag peptide. The eluted Flag-CDS2 (1.5 mL) was then incubated with the Ni-NTA agarose-bound MBOAT7-His for 2 hours. The immunoprecipitate was washed six times with PBS + PMSF + 1% Triton buffer and denatured with 2X SDS sample buffer for 10 min before western blotting.

Confocal imaging

The images were acquired using a LSM 980 with Airyscan 2 (Zeiss) or a confocal microscope (Leica SP8). Images were analyzed by Image J.

Subcellular fractionation

Subcellular fractionation of liver tissue was performed as described (35).

Antibodies

Rabbit antibodies against CDS2 (1:500, 13175-1-AP), SREBP2 (1:1000, 28212-1-AP), LDLR (1:1000, 10785-1-AP), SR-B1 (1:1000, 21277-1-AP) and MBOAT7 (1:100, 83546-3-RR) were purchased from Proteintech. Rabbit antibody

against MBOAT7 (1:1000, AV49811) was purchased from Sigma-Aldrich. Mouse antibody against Flag (1:1000, F1804) was purchased from Sigma-Aldrich. Rabbit antibody against Flag (1:2500, PM020) was purchased from MBL. Rabbit antibodies against ATGL (1:1000, 2138), phospho-AMPKα(Thr172)(40H9) (1:1000, 2535), AMPKα (1:1000, 2532), phospho-ACC (1:1000, 3661), ACC (1:1000, 3662), and ABCA1 (1:1000, 96292) were purchased from Cell Signaling. Mouse antibody against LXR (1:1000, ab41902) was purchased from Abcam. Rabbit antibodies against DGAT2 (1:1000, ab237613) and HMGCR (1:1000, ab174830) were purchased from Abcam.

Statistics

Results with error bars are mean±SEM. Comparisons between 2 groups were performed using the two-tailed Student's t-test. ANOVA was used for comparisons between more than 2 groups.

Acknowledgments

We thank Xixian Li from the Center for Biological Imaging at the Institute of Biophysics, CAS, for analyzing images. This work was supported by grants 2024YFA1306101, 2023YFA1800500 from National Key R&D Program of China and 92354301, 32230044, 32371230 and 32321004 from the National Natural Science Foundation of China, and BJ-2024-235 from National High Level Hospital Clinical Research Funding.

Figure 4A, Figure 6I and Figure 7H were created with BioRender (*https://BioRender.com/ k88h902*, *https://BioRender.com/s73x778, and https://BioRender.com/e57c295*).

Declaration of Interests

The authors have declared that no conflict of interest exists.

Author Contributions

J Xu and X Huang designed the experiments. J Xu, J Zhang, W Wang and H Su performed the experiments. J Xu, J Zhang, S Chen and M Ding analyzed the data. J Xu and X Huang wrote the manuscript.

References

1. Llovet JM, Willoughby CE, Singal AG, Greten TF, Heikenwälder M, El-Serag HB, Finn RS, et al.
Nonalcoholic steatohepatitis-related hepatocellular carcinoma: pathogenesis and treatment. Nature
Reviews Gastroenterology & Hepatology 2023;20:487-503.

2. Xiang L, Li X, Luo Y, Zhou B, Liu Y, Li Y, Wu D, et al. A multi-omic landscape of steatosis-to-NASH
progression. Life Metab 2022;1:242-257.

3. Walther TC, Chung J, Farese RV, Jr. Lipid Droplet Biogenesis. Annu Rev Cell Dev Biol
2017;33:491-510.

4. Schott MB, Weller SG, Schulze RJ, Krueger EW, Drizyte-Miller K, Casey CA, McNiven MA. Lipid
droplet size directs lipolysis and lipophagy catabolism in hepatocytes. J Cell Biol 2019;218:3320-3335.

5. Buch S, Stickel F, Trepo E, Way M, Herrmann A, Nischalke HD, Brosch M, et al. A genome-wide
association study confirms PNPLA3 and identifies TM6SF2 and MBOAT7 as risk loci for alcohol-related
cirrhosis. Nat Genet 2015;47:1443-1448.

6. Thangapandi VR, Knittelfelder O, Brosch M, Patsenker E, Vvedenskaya O, Buch S, Hinz S, et al.
Loss of hepatic Mboat7 leads to liver fibrosis. Gut 2021;70:940-950.

7. Varadharajan V, Massey WJ, Brown JM. Membrane-bound O-acyltransferase 7 (MBOAT7)-driven
phosphatidylinositol remodeling in advanced liver disease. J Lipid Res 2022;63:100234.

8. Alharthi J, Bayoumi A, Thabet K, Pan Z, Gloss BS, Latchoumanin O, Lundberg M, et al. A metabolic
associated fatty liver disease risk variant in MBOAT7 regulates toll like receptor induced outcomes. Nat
Commun 2022;13:7430.

9. Caddeo A, Spagnuolo R, Maurotti S. MBOAT7 in liver and extrahepatic diseases. Liver Int
2023;43:2351-2364.

10. Liu Y, Wang W, Shui G, Huang X. CDP-diacylglycerol synthetase coordinates cell growth and fat
storage through phosphatidylinositol metabolism and the insulin pathway. PLoS Genet
2014;10:e1004172.

11. D'Souza K, Kim YJ, Balla T, Epand RM. Distinct properties of the two isoforms of
CDP-diacylglycerol synthase. Biochemistry 2014;53:7358-7367.

12. Moore MP, Wang X, Kennelly JP, Shi H, Ishino Y, Kano K, Aoki J, et al. Low MBOAT7 expression, a
genetic risk for MASH, promotes a profibrotic pathway involving hepatocyte TAZ upregulation.
Hepatology 2024;81:576-590.

13. Tanaka Y, Shimanaka Y, Caddeo A, Kubo T, Mao Y, Kubota T, Kubota N, et al. LPIAT1/MBOAT7
depletion increases triglyceride synthesis fueled by high phosphatidylinositol turnover. Gut
2021;70:180-193.

14. Xu Y, Mak HY, Lukmantara I, Li YE, Hoehn KL, Huang X, Du X, et al. CDP-DAG synthase 1 and 2
regulate lipid droplet growth through distinct mechanisms. J Biol Chem 2019;294:16740-16755.

15. Qi Y, Kapterian TS, Du X, Ma Q, Fei W, Zhang Y, Huang X, et al. CDP-diacylglycerol synthases
regulate the growth of lipid droplets and adipocyte development. J Lipid Res 2016;57:767-780.

16. Xu J, Chen S, Wang W, Man Lam S, Xu Y, Zhang S, Pan H, et al. Hepatic CDP-diacylglycerol synthase
2 deficiency causes mitochondrial dysfunction and promotes rapid progression of NASH and fibrosis.
Sci Bull (Beijing) 2022;67:299-314.

17. Helsley RN, Varadharajan V, Brown AL, Gromovsky AD, Schugar RC, Ramachandiran I, Fung K, et al.
Obesity-linked suppression of membrane-bound O-acyltransferase 7 (MBOAT7) drives non-alcoholic
fatty liver disease. Elife 2019;8.

18. Meroni M, Dongiovanni P, Longo M, Carli F, Baselli G, Rametta R, Pelusi S, et al. Mboat7

down-regulation by hyper-insulinemia induces fat accumulation in hepatocytes. EBioMedicine 2020;52:102658.

19. Wang K, Lee CW, Sui X, Kim S, Wang S, Higgs AB, Baublis AJ, et al. The structure of phosphatidylinositol remodeling MBOAT7 reveals its catalytic mechanism and enables inhibitor identification. Nat Commun 2023;14:3533.

20. Olarte MJ, Kim S, Sharp ME, Swanson JMJ, Farese RV, Jr., Walther TC. Determinants of Endoplasmic Reticulum-to-Lipid Droplet Protein Targeting. Dev Cell 2020;54:471-487

21. Dhiman R, Caesar S, Thiam AR, Schrul B. Mechanisms of protein targeting to lipid droplets: A unified cell biological and biophysical perspective. Semin Cell Dev Biol 2020;108:4-13.

22. Pataki CI, Rodrigues J, Zhang L, Qian J, Efron B, Hastie T, Elias JE, et al. Proteomic analysis of monolayer-integrated proteins on lipid droplets identifies amphipathic interfacial alpha-helical membrane anchors. Proc Natl Acad Sci U S A 2018;115:E8172-E8180.

23. Bersuker K, Peterson CWH, To M, Sahl SJ, Savikhin V, Grossman EA, Nomura DK, et al. A Proximity Labeling Strategy Provides Insights into the Composition and Dynamics of Lipid Droplet Proteomes. Dev Cell 2018;44:97-112 e117.

24. Mejhert N, Kuruvilla L, Gabriel KR, Elliott SD, Guie MA, Wang H, Lai ZW, et al. Partitioning of MLX-Family Transcription Factors to Lipid Droplets Regulates Metabolic Gene Expression. Mol Cell 2020;77:1251-1264 e1259.

25. Song J, Mizrak A, Lee CW, Cicconet M, Lai ZW, Tang WC, Lu CH, et al. Identification of two pathways mediating protein targeting from ER to lipid droplets. Nat Cell Biol 2022;24:1364-1377.

26. Malis Y, Armoza-Eilat S, Nevo-Yassaf I, Dukhovny A, Sklan EH, Hirschberg K. Rab1b facilitates lipid droplet growth by ER-to-lipid droplet targeting of DGAT2. Sci Adv 2024;10:eade7753.

27. Wilfling F, Wang H, Haas JT, Krahmer N, Gould TJ, Uchida A, Cheng JX, et al. Triacylglycerol synthesis enzymes mediate lipid droplet growth by relocalizing from the ER to lipid droplets. Dev Cell 2013;24:384-399.

28. Wang JQ, Li LL, Hu A, Deng G, Wei J, Li YF, Liu YB, et al. Inhibition of ASGR1 decreases lipid levels by promoting cholesterol excretion. Nature 2022;608:413-420.

29. Chen G, Zhou G, Lotvola A, Granneman JG, Wang J. ABHD5 suppresses cancer cell anabolism through lipolysis-dependent activation of the AMPK/mTORC1 pathway. J Biol Chem 2021;296:100104.

30. Klemm RW, Carvalho P. Lipid Droplets Big and Small: Basic Mechanisms That Make Them All. Annu Rev Cell Dev Biol 2024;40:143-168.

31. Bersuker K, Olzmann JA. Establishing the lipid droplet proteome: Mechanisms of lipid droplet protein targeting and degradation. Biochim Biophys Acta Mol Cell Biol Lipids 2017;1862:1166-1177.

32. Mancina RM, Dongiovanni P, Petta S, Pingitore P, Meroni M, Rametta R, Boren J, et al. The MBOAT7-TMC4 Variant rs641738 Increases Risk of Nonalcoholic Fatty Liver Disease in Individuals of European Descent. Gastroenterology 2016;150:1219-1230 e1216.

33. Xu X, Xu H, Liu X, Zhang S, Cao Z, Qiu L, Du X, et al. MBOAT7 rs641738 (C>T) is associated with NAFLD progression in men and decreased ASCVD risk in elder Chinese population. Front Endocrinol (Lausanne) 2023;14:1199429.

34. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the CRISPR-Cas9 system. Nature Protocols 2013;8:2281-2308.

35. Ding Y, Zhang S, Yang L, Na H, Zhang P, Zhang H, Wang Y, et al. Isolating lipid droplets from multiple species. Nat Protoc 2013;8:43-51.

 Figure Legends

Figure 1. MBOAT7 and CDS2 regulate lipid metabolism in liver.

(A-B) The correlation between *MBOAT7* and *CDS2* transcript levels in the gene

expression datasets GSE48452 (NASH samples) and GSE66676, respectively. (C)

The correlation between *MBOAT7* and *CDS2* transcript levels in the GEPIA website.

(D-F) Procedure for liver-specific knockdown of *Mboat7* and/or *Cds2* in mice. Mice

were injected with AAV-TBG-shCon, AAV-TBG-sh*Mboat7*, AAV-TBG-sh*Cds2*, or

AAV-TBG-sh*Mboat7* + AAV-TBG-sh*Cds2* and fed a chow diet for 4 weeks before

analysis. (D). Hepatic triglyceride (E) and cholesterol (F) levels were measured in the

mice in (D). N=6-7 mice per group. (G) H&E staining of liver sections from mice in

(D). Scale bar represents 50 μm in main images and 10 μm in enlarged images. (H)

Hepatic steatosis score based on liver sections of mice in (D). N=6-7 mice per group.

(I) RNAseq was performed on liver samples from the mice in (D). The numbers of

genes up- or down-regulated in sh*Mboat7* vs sh*Con*, sh*Cds2* vs sh*Con* and sh*Mboat7*

+ sh*Cds2* vs sh*Con* are shown. (J) KEGG analysis results showing the pathways

related to the 301 common downregulated genes in (I). (K) qRT-PCR analysis of the

mRNA levels of genes that regulate cholesterol and fatty acid metabolism in the liver.

N=5-8 mice per group. (L) Western blot analysis of liver lysates from mice in (D).

N=3 mice per group. * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 2. MBOAT7 is localized to the ER-LD contacts in CDS2 KO cells.

(A) CDS2 KO and control HeLa cells were transfected with MBOAT7-EGFP and

treated with 100 μM OA for 12 hours, followed by LD labeling with LipidTOX. (B)

Fluorescence colocalization was analyzed along the white line in the main confocal

image. (C) CDS2 KO and control HeLa cells were treated with 100 μM OA for 12

hours and immunostained with anti-MBOAT7 antibody. (D) Fluorescence

colocalization was analyzed along the white line in the main confocal image. (E) A

3D-SIM image reveals that MBOAT7-EGFP enwraps a LD in an OA-treated CDS2

KO cell. Scale bar: 2 μm. (F) CDS2 KO and control HeLa cells were transfected with

mCherry-SEC61b (an ER marker), then treated with 100 μM OA for 12 hours. LDs

were labeled with BODIPY. (G) Fluorescence colocalization was analyzed along the

white line in the main confocal image. (H) C56BL/6 mice were fed a chow (control)

or GAN (MASLD) diet (40 kcal% fat, 20 kcal% fructose and 2% cholesterol) for 26

weeks. Liver sections were immunostained with anti-MBOAT7 antibody. White

arrows indicate MBOAT7 enclosing LDs. (I) The number of LDs surrounded by

MBOAT7 was quantified by analyzing 1,000 LDs in confocal images. N=9-11 liver

sections per group. Except for (E), scale bars represent 10 μm in main images and 2

μm in enlarged images.

Figure 3. The N-terminus of MBOAT7 is essential for LD targeting of MBOAT7

and MBOAT7 interacts with CDS2.

(A) Top: 3D structure of MBOAT7 in the ER membrane, highlighting its

transmembrane (TM) domains. Bottom: The 11 TM domains of MBOAT7 (boxes)

and the three regions of the protein expressed as truncations. (B) CDS2 KO cells

transfected with truncated versions of MBOAT7 were treated with 100 μM OA for 12

hours, followed by LD labeling with LipidTOX. (C) CDS2 KO cells transfected with

MBOAT7-EGFP containing specific TM domain deletions were treated with 100 μM

OA for 12 hours. LDs were labeled with LipidTOX. (D) CDS2 KO cells transfected

with MBOAT7-EGFP chimeras containing TM3 or TM4 from MBOAT5 were treated

with 100 μM OA for 12 hours and labeled with LipidTOX to stain LDs. (E) Co-IP

assay showing that Flag-CDS2 precipitates MBOAT7-EGFP. (F) Co-IP assay showing

that MBOAT7-EGFP precipitates Flag-CDS2. (G) Co-IP assay demonstrating that

endogenous CDS2 precipitates MBOAT7. (H) *In vitro* pull-down assay indicating that

Flag-CDS2 directly binds to MBOAT7-His. Scale bars: 10 μm in main images and 2

μm in enlarged images.

Figure 4. RAB1A is required for targeting of MBOAT7 to LDs in CDS2 KO cells.

(A) Schematic of the workflow for identifying MBOAT7-interacting proteins in

control and CDS2 KO cells. (B) Venn diagram illustrating the overlap between the

MBOAT7-interacting proteins from (A) and LD proteins from various LD proteomes.

(C) Five MBOAT7-interacting proteins identified from the Venn diagram in (B). (D)

CDS2 KO cells were transfected with siNC or siRAB1A, and MBOAT7-EGFP, then

treated with 100 μM OA for 12 hours. LDs were labeled with LipidTOX. (E) *RAB1A*

mRNA levels in HeLa cells transfected with control siRNA or siRAB1A. (F)

Quantification of LDs surrounded by MBOAT7-EGFP in control (siNC) or RAB1A

knockdown (siRAB1A) cells. (G) Co-IP assay showing that Flag-RAB1A precipitates

MBOAT7-EGFP in CDS2 KO cells. (H) Co-IP assay showing that Flag-RAB1A

precipitates MBOAT7-EGFP in CDS2 KO cells, and expression of HA-CDS2 in

CDS2 KO cells disrupts the RAB1A/MBOAT7 interaction.

Figure 5. MBOAT7 reduces LD size in CDS2 KO cells by competing with

DGAT2 for targeting to LDs.

(A) Control HeLa or CDS2 KO cells were transfected with MBOAT7-EGFP or the

MBOAT7(N321A, H356A)-EGFP mutant, then treated with 100 μM OA for 12 hours.

LDs were labeled by LipidTOX. Scale bar: 10 μm. (B) Quantification of LD sizes

from (A). N=200 per group. (C) 293T cells were transfected with GFP-RAB1A and

Flag-DGAT2 or Flag control plasmids. Co-IP assay shows that Flag-DGAT2

precipitates GFP-RAB1A. (D) 293T cells were transfected with GFP-RAB1A and

Flag-GPAT4 or Flag control plasmid. Co-IP assay shows that Flag-GPAT4 precipitates

GFP-RAB1A. (E) CDS2 KO cells were transfected with GFP-RAB1A and

Flag-DAGT2 with or without MBOAT7-EGFP. Co-IP shows that Flag-DGAT2

precipitates GFP-RAB1A. (F) CDS2 KO cells were transfected with Flag-DGAT2 and

GFP-RAB1A with siRNA control (siCon) or siMBOAT7. Co-IP assay shows that

Flag-DGAT2 precipitates GFP-RAB1A. (G) Confocal images showing that

EGFP-DGAT2 targets to LDs in CDS2 KO cells. MBOAT7-mCherry expression

blocks LD targeting of DGAT2. Scale bar: 10 μm in main images and 2 μm in

enlarged images. (H) Co-IP assay showing that Flag-DGAT2 precipitates full-length

GFP-RAB1A as well as N-terminal and C-terminal truncated forms. (I) Co-IP assay

showing that Flag-MBOAT7 precipitates full length GFP-RAB1A as well as

N-terminal and C-terminal truncated forms. (J) Co-IP assay showing that

Flag-DGAT2 precipitates GFP-RAB1A in CDS2 KO cells treated with or without OA.

(K) Co-IP assay showing that Flag-MBOAT7 precipitates GFP-RAB1A in CDS2 KO

cells treated with or without OA. **** p < 0.0001.

708   Figure 6. MBOAT7 regulates LD size and lipolysis.

(A) MBOAT7 enzymatic activity was measured in *Cds2^{f/f}*, *Cds2^{f/f};AlbCre* mice, and

Cds2^{f/f};AlbCre mice injected with AAV-TBG-*Cds2*. N=5-6 mice per group. (B)

Schematic showing the procedure for hepatic overexpression of MBOAT7 in the liver

of mice with and without hepatic depletion of CDS2. *Cds2^{f/f}* and *Cds2^{f/f};AlbCre* mice

were injected with AAV-TBG-Con or AAV-TBG-*Mboat7* and fed a chow diet for 4

weeks before analysis. (C and D) Hepatic cholesterol (C) and triglyceride (D) levels

were analyzed in the mice from (B). N=7-8 mice per group. (E) H&E staining of liver

sections from *Cds2^{f/f}* and *Cds2^{f/f};AlbCre* mice. Scale bar represents 50 μm in main

images and 10 μm in enlarged images. (F) Primary hepatocytes isolated from *Cds2^{f/f}*

and *Cds2^{f/f};AlbCre* mice were infected with Ad-EGFP or Ad-MBOAT7-EGFP and

treated with 100 μM OA for 12 hours. LDs were stained with LipdTOX before

confocal imaging. Scale bar represents 10 μm in main images and 2 μm in enlarged

images. (G) Quantification of LD sizes in (F). (H) Western blot analysis of LD, ER

and cytosol fractions from the liver of *Cds2^{f/f};AlbCre* mice that are injected with

AAV-TBG-Con or AAV-TBG-*Mboat7*. (I) Primary hepatocytes from *Cds2^{f/f}* and

Cds2^{f/f};AlbCre mice were infected with Ad-EGFP or Ad-MBOAT7-EGFP, then treated

with 10 μM ISO or DMSO (control) for the indicated times. Free fatty acids in the

medium were analyzed. * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001.

728   Figure 7. MBOAT7 and CDS2 regulate cholesterol metabolism through the

729   AMPK pathway.

(A) Western blot of *Cds2^{f/f}* and *Cds2^{f/f};AlbCre* mice injected with AAV-TBG-Con or

AAV-TBG-*Mboat7*, and fed a chow diet for 4 weeks before analysis. (B)

Quantifications of proteins from (A). (C) Primary hepatocytes isolated from

Cds2^{f/f};AlbCre mice were infected with Ad-EGFP or Ad-MBOAT7-EGFP. Cells were

serum-starved for 4 hours, then treated with 10 μM dorsomorphin in no-glucose

DMEM + 10% FBS medium for 4 hours. Western blot analysis was then performed.

(D) Primary hepatocytes from *Cds2^{f/f};AlbCre* mice were infected with Ad-EGFP or

Ad-MBOAT7-EGFP and treated with 10 μM ISO or DMSO (control) for 6 hours.

Western blot analysis was then performed. (E) Quantifications of proteins from (D).

(F) Primary hepatocytes from *Cds2^{f/f};AlbCre* mice were infected with Ad-EGFP or

Ad-MBOAT7-EGFP. Cells were pretreated with DMSO (control) or Atglistatin

(ATGLi; 20 μM) for 30 minutes, then treated with 10 μM ISO for 6 hours. Protein

levels were analyzed by western blotting. (G) Quantifications of proteins from (F). (H)

A model illustrating how MBOAT7 and CDS2 regulate LD dynamics and cholesterol

metabolism in hepatocytes. CDS2 and MBOAT7 physically interact in the ER

membrane and both regulate PI biosynthesis. MBOAT7 catalyzes the generation of

PI-AA, which inhibits CDS2 activity. In the absence of CDS2, MBOAT7 relocates to

the ER-LD contacts via binding to RAB1. CDS2 deficiency causes increased LD size

accompanied by increased recruitment of DGAT2 to LDs, whereas overexpression of

MBOAT7 in CDS2-deficient conditions reduces DGAT2 targeting to LDs, increases

ATGL recruitment to the LD surface, and reduces LD size. The increased lipolysis

induced by MBOAT7 expression activates AMPK, enhancing LXR protein levels and

expression of its downstream target genes, thus stimulating cholesterol metabolism. *

$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

A
Control +OA · CDS2KO +OA
MBOAT7-EGFP LipidTOX

B
Control +OA · CDS2KO +OA
— MBOAT7-EGFP
— LipidTOX

C
Control +OA · CDS2KO +OA
MBOAT7 BODIPY

D
Control +OA · CDS2KO +OA
— MBOAT7
— BODIPY

F
Control +OA · CDS2KO +OA
mCherry-SEC61b BODIPY

H
Control · MASLD
MBOAT7 LipidTOX DAPI

E
CDS2KO +OA
MBOAT7-EGFP LipidTOX

G
Control +OA · CDS2KO +OA
— mCherry-SEC61b
— BODIPY

I
MBOAT7+ LD/1000 LDs
***
Control · MASLD

A

Transfect with EGFP or MBOAT7-EGFP — Control
Transfect with EGFP or MBOAT7-EGFP — CDS2KO

IP:GFP

Immunoprecipitate

MS/MS

B

Pataki et al. 2018
Bersuker et al. 2018
Mejhert et al. 2020 (SUM159 cell)
Mejhert et al. 2020 (THP-1 macrophage)

C

Protein	Fold change
RAB7A	10.4
RAB1A	7.2
RAB18	6.2
RAB14	4.8
CYB5R3	4

D

siNC

siRAB1A

MBOAT7-EGFP LipidTOX

E

RAB1A

Relative mRNA

siNC   siRAB1A

F

MBOAT7+ LD/total LD (per cell)

****

siNC   siRAB1A

G

Con   CDS2KO

Flag	+	−	+	−
Flag-RAB1A	−	+	−	+
MBOAT7-EGFP	+	+	+	+

IB: GFP —70kDa
IB: Flag —25kDa   IP: Flag

IB: GFP —70kDa
IB: Flag —25kDa   input

H

CDS2KO

Flag	+	−	−
Flag-RAB1A	−	+	+
MBOAT7-EGFP	+	+	+
HA-CDS2	−	−	+

IB: GFP —70kDa
IB: Flag —25kDa   IP: Flag

IB: GFP —70kDa
IB: Flag —25kDa   input