Variation in Synaptic Inputs Drives Functional Versatility for Sound Encoding in Globular Bushy Cells

(Abbreviated title): Synaptic Input Variation in Globular Bushy Cells

Chunjian Wang[1,2], Go Ashida[3,4], Christian Keine[1,2], Ivan Milenkovic[1,2,4]

[1]Department of Human Medicine, Carl von Ossietzky Universität Oldenburg, Germany
[2]Research Center Neurosensory Science, Carl von Ossietzky Universität Oldenburg, Germany
[3]Department of Neuroscience, Carl von Ossietzky Universität Oldenburg, Germany
[4]Cluster of Excellence "Hearing4all", Carl von Ossietzky Universität Oldenburg, Germany

ORCIDs:

Chunjian Wang   0000-0002-3399-3489

Go Ashida: 0000-0002-8886-1213

Christian Keine: 0000-0003-0995-9667

Ivan Milenkovic: 0000-0003-1431-9031

Summary

Synaptic convergence is fundamental to neuronal circuit function, enabling diverse processing such as coincidence detection and reliable signal transmission. In sensory systems, the architecture of convergence and strengths of synaptic inputs are pivotal for extracting distinct features of the sensory stimulus. In auditory system, globular bushy cells (GBC) in the cochlear nucleus receive multiple axosomatic endbulb of Held terminals from the auditory nerves, which vary considerably in size, even among inputs targeting the same cell. However, the functional consequences of this input strength variation for sound encoding remain unclear. Here, we investigated how synaptic input variation shapes sound encoding in GBC of Mongolian gerbils, using in vitro conductance-clamp recordings and computational modeling, in which synaptic inputs with varying strengths were simulated. We found that input variation critically shapes GBC sound encoding by influencing temporal precision and firing rates. Low input variation allows GBC to act as coincidence detectors, thereby improving temporal precision. Conversely, high input variation increases overall firing rates and enhances the encoding of amplitude modulations, albeit at the expense of temporal precision. These findings suggest that inherent endbulb strength heterogeneity allows GBC to operate along a functional continuum. This variation in input strength may provide the basis for generating diverse information streams to downstream targets including the medial nucleus of the trapezoid body (MNTB).

Keywords: Cochlear nucleus; globular bushy cells; temporal precision; endbulb of Held; coincidence detection; conductance-clamp; auditory nerve; Mongolian gerbil; auditory brainstem; functional heterogeneity; computational modeling

Introduction

Neuronal circuits rely on intricate patterns of synaptic communication to process information and guide behavior. A core organizational principle of these circuits is synaptic convergence, where individual neurons integrate inputs from multiple presynaptic sources. The extent of such convergence varies considerably across neuron types and brain regions, reflecting distinct operational modes tailored to specific computational needs.

This diversity is particularly evident in sensory systems, where parallel pathways are often specialized to extract distinct stimulus features (Blackburn and Sachs, 1989; Nagayama et al., 2004; Doucet and Ryugo, 2006; Huang and Chacron, 2016; Ashida et al., 2019). For instance, within the auditory brainstem, the degree of synaptic convergence varies dramatically. Neurons in the medial nucleus of the trapezoid body (MNTB) are innervated by a single massive axosomatic terminal – the calyx of Held – which ensures fast and reliable signal transmission (Spirou et al., 1990; Smith et al., 1998; von Gersdorff and Borst, 2002; Schneggenburger and Forsythe, 2006; Englitz et al., 2009; Borst and Soria van Hoeve, 2012). In contrast, octopus and D-stellate cells in the cochlear nucleus (CN) integrate inputs from dozens of auditory nerve (AN) fibers, responding to the coincident firing of their inputs (Oertel, 1999; Cao and Oertel, 2010; McGinley et al., 2012; Xie and Manis, 2017; Kreeger et al., 2025). Such heterogeneity in input number also occurs among cells of the same type. For instance, retinal ganglion cells (RGCs) in the primate fovea receive input from only a few bipolar cells, whereas peripheral RGCs integrate signals from hundreds of inputs (Calkins et al., 1994; Jacoby et al., 2000; Kolb and Marshak, 2003). Similarly, in the avian CN homolog, the nucleus magnocellularis, neurons processing low-frequency sounds receive numerous small inputs, while high-frequency neurons are innervated by fewer, larger inputs (Köppl, 1994; Fukui and Ohmori, 2004), enabling frequency-dependent specialization of temporal processing (Oline et al., 2016). These examples illustrate how convergence patterns can be finely tuned to match specific functional requirements across diverse neural systems.

Beyond input number, the variation in synaptic strength among converging inputs represents

another critical, yet less understood, factor influencing neuronal circuit function (Sherman and

Guillery, 1998; Rouiller and Welker, 2000; Groh et al., 2008; Wolff et al., 2021; Govindaiah et al.,

2023). In multiple brain regions, including the cerebellum, thalamus, and cortex, some neurons

integrate many similarly sized subthreshold inputs, while others are driven by one or two

suprathreshold "driver inputs" alongside multiple weaker ones (Sherman and Guillery, 1998; Groh

et al., 2008; Wolff et al., 2021). Understanding how such variations in synaptic strength affect

neuronal processing is essential for deciphering neural circuit computations and understanding

the fundamental aspects of sensory operation.

In this study, we used globular bushy cells (GBC) in the mammalian CN as a model system to

explore the functional consequences of synaptic input variation. GBC, located near the AN root in

the caudal anteroventral (AVCN) and posteroventral CN (PVCN) (Osen, 1969; Cant and Morest,

1979), receive excitatory inputs via large endbulb of Held terminals from a varying number of AN

fibers, ranging from 5-12 in mice (Cao and Oertel, 2010; Spirou et al., 2023) to over 20 in cats

(Liberman, 1991; Ostapoff and Morest, 1991; Spirou et al., 2005). Functionally, GBC are crucial

for high-fidelity temporal processing of auditory information, and their temporal precision can

exceed that of AN inputs (Blackburn and Sachs, 1989; Joris et al., 1994; Wei et al., 2017). Recent

detailed morphological analysis has revealed significant heterogeneity in the size of endbulbs

converging onto individual GBC with some receiving inputs from many similarly sized subthreshold

endbulbs, while others are dominated by a few large suprathreshold endbulbs alongside smaller

ones (Nicol and Walmsley, 2002; Spirou et al., 2023). Despite this documented morphological

heterogeneity, the computational role of endbulb input variation in shaping GBC output during

sound processing remains poorly understood.

To address this knowledge gap, we combined *in vitro* whole-cell conductance-clamp recordings

from GBC in Mongolian gerbils with computational modeling. By systematically manipulating the

variation in synaptic input strength among simulated AN fibers, we assessed its impact on GBC

firing rates and temporal precision. Our findings reveal that the degree of input variation shapes

GBC output characteristics. Specifically, while overall firing rates increased with input strength

variation, temporal precision in encoding the stimulus fine-structure deteriorated. These findings

suggest that the inherent morphological diversity of endbulb inputs plays a key role in regulating

GBC encoding strategies and may contribute to the control of information flow to multiple sound

processing circuits in the downstream regions of the auditory brainstem.

Results

Synaptic input variation enhances GBC firing but reduces temporal precision

To study the impact of synaptic input variation in GBC, we targeted them using their established anatomical and electrophysiological characteristics. Located in the caudal AVCN near the AN root and PVCN (Osen, 1969), GBC are characterized by round-to-oval cell bodies with a few short dendrites and typically fire only one or a few action potentials (APs) in response to suprathreshold current injection (Fig. 1A). Functionally, GBC are specialized for high-fidelity temporal processing of auditory information, often exceeding the temporal precision of their AN inputs, and can sustain high firing rates up to a few hundred spikes per second for low-frequency sound stimulation (Joris et al., 1994; Wei et al., 2017, 2023).

Using an established auditory periphery model (Bruce et al., 2018), we generated spike trains for 10 converging AN fibers responding to 200-ms pure tones (10 repetitions) at frequencies of 350 Hz, 1 kHz, and 3.5 kHz. These frequencies were chosen to encompass the range where AN fibers exhibit significant phase-locking (350 and 1000 Hz) and to extend beyond the typical upper limit, where phase-locking diminishes in rodents (Wei et al., 2017, 2023). The resulting AN spike trains were convolved with an experimentally derived EPSC waveform to generate conductance trains, which were then injected into the recorded GBC. By systematically varying the synaptic input strength distribution across these 10 simulated AN fibers, we created three distinct conditions: 'no variation' (all inputs of equal strength: coefficient of variation $v = 0$), 'medium variation' ($v = 0.4$), and 'high variation' ($v = 1$) (Fig. 1B). For the 'high variation' condition, the largest input was near or above the spike threshold, while for the 'no' and 'medium' variation condition, all inputs were subthreshold.

First, we analyzed how input variation affected average firing rates. We found that increasing input variation elevated GBC firing rates during stimulation at 1 kHz and 3.5 kHz (1 kHz: no variation 132 ± 48 Hz vs. high variation 171 ± 34 Hz, p = 7.9e-6, n = 15), while firing rates were moderately

decreased by higher input variation during 350 Hz stimulation (no variation = 245 ± 52 Hz vs. high

variation = 228 ± 39 Hz, p = 0.01, Fig. 1C1).

Next, we assessed the impact of input variation on temporal precision, quantified by vector

strength (VS), which measures the neuron's ability to phase-lock to the periodic waveform of an

acoustic stimulus. The simulated AN inputs exhibited VS values of 0.75 and 0.77 at stimulation

frequencies of 350 and 1000 Hz, respectively, consistent with *in vivo* observations (Johnson, 1980;

Palmer and Russell, 1986; Heeringa et al., 2020; Heeringa, 2024). In postsynaptic GBC, robust

phase-locking was observed at 350 Hz and 1 kHz but not at 3.5 kHz, in line with *in vivo* data (Wei

et al., 2017). Notably, increasing input variation led to a significant reduction in GBC phase-

locking, an effect that was particularly robust at 350 Hz (main effect variation: p = 3.7e-6; 350 Hz:

no variation = 0.86 ± 0.02 vs. high variation = 0.81 ± 0.01, p = 3.3e-8, Fig. 1C2). These results

indicate that higher input variation leads to a trade-off between increasing firing rates and

degrading temporal precision.

Effects of input variation are well reproduced in a coincidence-counting model

To investigate whether these experimental observations could be generalized and to ascertain if

they arose primarily from the input statistics, we employed a computational model for GBC, which

was first developed for cats and adapted for gerbils in the present study (Ashida et al., 2019). This

model generates an output spike when the summed inputs exceed an adaptive threshold. The

model was driven with simulated AN spike trains identical to those used in the *in vitro* experiments.

A wider configuration of input variations was adopted, including the 'no', 'medium', and 'high'

variations tested in our conductance-clamp experiments (Fig. 2A). Notably, the model reproduced

the key experimental findings, namely increased firing rates (Fig. 2B) and reduced temporal

precision with higher input variation (Fig. 2C). This suggests that the interaction between the

synaptic input statistics and a simplified cellular integration mechanism is sufficient to replicate the

experimentally observed phenomena.

Higher input variation shortens AP latency but increases temporal jitter

Given that increased synaptic input variation enhances overall firing rates (Fig. 1C1), we next investigated whether this effect differentially impacts the onset and sustained phases of GBC responses. To isolate these components, we constructed peristimulus time histograms (PSTH) from responses to 200 repetitions of a 3.5 kHz pure-tone stimulus, a frequency selected to minimize confounding phase-locking effects. Consistent with *in vivo* recordings (Britt and Starr, 1976; Ritz and Brownell, 1982; Blackburn and Sachs, 1989), both the simulated AN inputs and GBC responses exhibited primary-like PSTHs, characterized by a sharp onset peak followed by adaptation to a lower sustained rate (Fig. 3A). In addition, the PSTH exhibited a 'notch' with lower activity immediately following the onset peak, which is characteristic for GBC (Smith and Rhode, 1987; Blackburn and Sachs, 1989) A qualitative comparison of the 'no variation' and 'high variation' conditions indicated that higher input variation tended to reduce peak onset firing while elevating the sustained rate (Fig. 3A2 inset). Quantitative analysis confirmed these distinct effects. Onset spike probability, defined as the fraction of trials with at least one spike within the initial 1.5 ms of the response, was generally high but significantly decreased with increasing input variation (no variation = 0.97 ± 0.02 vs. high variation = 0.94 ± 0.02, p = 7.3e-5, n = 17, Fig. 3B2). Conversely, both sustained (100–200 ms) and spontaneous firing rates significantly increased with higher input variation (sustained: no variation = 72 ± 27 Hz vs. high variation = 131 ± 28 Hz, p = 3.2e-11, Fig. 3B3; spontaneous: no variation = 25 ± 14 Hz vs. high variation = 65 ± 15 Hz, p = 4.9e-11, Fig. 3B4). To further probe the reliability and timing of the response onset, we analyzed first-spike latency and jitter. Visual inspection of superimposed spikes suggested that while some initial spikes occurred earlier under the 'high variation' condition, their timing across repetitions appeared more variable compared to responses generated with no input variation (Fig. 3B1). Indeed, quantitative analysis confirmed that the average first-spike latency significantly decreased with higher input variation (Fig. 3B5, no variation = 3 ± 0.14 ms vs. high variation =

2.92 ± 0.15 ms, p = 0.007). However, this shorter latency was accompanied by reduced temporal

precision, as evidenced by a significant increase in first-spike jitter (Fig. 3B6, no variation =

0.42 ± 0.12 ms vs. high variation = 0.63 ± 0.21 ms, p = 7.4e-5). These experimental findings were

replicated in the computational model (Fig. 3C) which reproduced the characteristic PSTH shapes

(Fig 3A3) and provided additional insight. While most response characteristics changed gradually,

the median spontaneous rate increased sharply at input variations of 0.7-0.8 (Fig. 3C4),

corresponding to the point where the largest input transitions from subthreshold to suprathreshold,

enabling it to generate GBC spikes on its own. The slight decrease in first-spike latency and the

increased first-spike jitter were also observed in the model and comparable to the experimental

results (Fig. 3C5, 3C6). However, the simulated distributions of first-spike latency and jitter were

narrower than those of the experimental data, likely due to the existence of various additional

noise sources in the experimental setting that were not included in the model (e.g., channel and

thermal fluctuations). In summary, both experimental and modeling results demonstrated that

input variability differentially sculpts GBC responses by enhancing sustained firing rates while

simultaneously degrading both the reliability and temporal precision of the response onset.

Input variation enhances firing rate tracking of amplitude-modulated sounds

The results from pure-tone stimulation suggest that the degree of synaptic input variation can

modulate GBC function, potentially shifting its operational emphasis: low input variation favors

high temporal precision, whereas high input variation enhances firing rates. To further explore this

phenomenon in the context of more complex, naturalistic stimuli, we examined how input variation

impacts the encoding of sinusoidal amplitude-modulated (SAM) sounds, which mimic the temporal

fluctuations present in speech and environmental sounds (Van Tasell et al., 1987; Drullman et al.,

1996; Joris et al., 2004). Effective SAM encoding requires both precise spike timing relative to the

modulation cycle (temporal code) and the ability of firing rates to track changes in sound level

(rate code). GBC are known to effectively follow rapid amplitude modulations (Palmer et al., 1986;

Rhode and Greenberg, 1994; Wang and Sachs, 1994).

We simulated the spiking responses of AN fibers to a 13 kHz pure-tone carrier, 100% amplitude-

modulated at frequencies of 200 Hz, 600 Hz, and 2000 Hz. Consistent with previous in vivo

observations (Smith and Brachman, 1980; Joris and Yin, 1992), the firing rates of simulated AN

fibers were modulated by the stimulus envelope (Fig. 4A1). This envelope phase-locking was

evident for 200 and 600 Hz, but not at 2000 Hz. Injecting the corresponding conductance trains

into GBC under no, medium, and high variation conditions yielded robustly modulated firing rates

(Fig. 4A2), which were also reproduced by the GBC model (Fig. 4A3). In both the experimental

data and the model, responses appeared qualitatively sharper and more temporally restricted

under the 'no variation' condition compared to the 'high variation' condition.

Consistent with our pure-tone data, overall firing rates during SAM stimulation significantly

increased with input variation for all tested modulation frequencies (main effect of variation: $p = 3e\text{-}$

10; 600 Hz: no variation $= 67 \pm 50$ Hz vs. high variation $= 115 \pm 38$ Hz, $p = 1.5e\text{-}7$, $n = 15$,

Fig. 4B1). Next, we assessed temporal coding fidelity using two complementary metrics: VS and

correlation index (CI). The VS, which measures synchronicity of spikes to the modulation cycle,

indicated robust phase-locking for 200 Hz and 600 Hz modulation frequencies and the VS value

significantly decreased with increasing input variation (main effect of variation: $p = 5.8e\text{-}14$,

600 Hz: no variation $= 0.51 \pm 0.06$ vs. high variation $= 0.37 \pm 0.03$, $p = 7.2e\text{-}10$, Fig. 4B2). This is

consistent with the reduced temporal precision observed during pure-tone stimulation, indicating

poorer encoding of the modulation fine-structure. Similarly, the CI, which quantifies spike time

precision across stimulus repetitions (Joris et al., 2006; Kessler et al., 2021), also significantly

decreased with higher input variation (main effect of variation: $p = 5.9e\text{-}13$, 600 Hz: no variation $=$

1.53 ± 0.12 vs. high variation $= 1.26 \pm 0.04$, $p = 3e\text{-}8$, Fig. 4B3). Together, these measures

indicate that increased input variation degrades temporal precision relative to the stimulus cycle

and reduces spike time consistency across trials. Concurrently, the entrainment index, quantifying

the proportion of interspike intervals within $0.5 - 1.5$ times the stimulus period (Joris et al., 1994),

significantly increased with input variation at low modulation frequencies (main effect of variation:

p =4.7e-6, 200 Hz: no variation = 0.6 ± 0.17 vs. high variation = 0.68 ± 0.1, p = 0.01, Fig. 4B4). At

200 Hz, the distributions of the measured and simulated entrainment index were broader than

those of the vector strength (Fig 4B2), because in contrast to phase-locking, entrainment is linked

to firing rate which was also broadly distributed (Fig 4B1).

Next, we investigated how well GBC firing rates tracked the stimulus envelope, reflecting their

rate-coding capacity. By calculating the normalized cross-correlation (CorrNorm) between the

stimulus envelope and the periodically oscillating firing rate, we quantified the GBC's ability to

follow the stimulus envelope. Firing rates tracked envelope modulations at 200 Hz and 600 Hz,

and CorrNorm increased significantly with input variation at the 200 Hz modulation frequency,

indicating an improved neuronal representation of the stimulus envelope (200 Hz: no variation =

0.87 ± 0.02 vs. high variation = 0.93 ± 0.01, p = 2.5e-10, Fig. 4B5). In summary, during SAM

stimulation, increased input variation shifts the GBC coding strategy by degrading fine-structure

temporal precision (Figs. 1C2, 2C) while simultaneously enhancing envelope representation

through firing rate modulation. This suggests that GBC receiving different convergent patterns of

AN inputs might be differentially suited to either favor precise temporal representation, requiring

highly coincident inputs, or emphasize envelope tracking via firing rate modulation and generally

higher firing rates (Rothman and Young, 1996; Joris and Smith, 2008; Koert and Kuenzel, 2021).

Impact of input variation is consistent with morphologically plausible input sizes

As demonstrated by our present results, the degree of variation among endbulb inputs can

influence GBC coding motifs, potentially shifting the operational focus between temporal precision

(i.e., coincidence detection) and firing rate encoding (latency-first mode). As these interpretations

were based on simulated input amplitudes drawn from a gamma distribution, we further evaluated

their *in vivo* relevance by testing whether the observed effects can also be verified with the known

morphological variability of GBC synaptic inputs. Therefore, we integrated recent morphological

data quantifying endbulb size distributions onto mouse GBC (Spirou et al., 2023). Using these

estimates allowed us to constrain synaptic input variations within a plausible morphological range.

Specifically, we tested two conditions of synaptic input variation corresponding approximately to

our previously employed 'medium' ($v \approx 0.4$, 'morph$_{MED}$') and 'high' ($v \approx 0.7$, morph$_{HIGH}$) variation

conditions.

GBC responses to pure tones under morphologically constrained input variations were consistent

with our previous observations. The 'morph$_{HIGH}$' condition significantly increased firing rates at 1

kHz and 3.5 kHz stimulation frequencies compared to the 'morph$_{MED}$' condition (1 kHz: morph$_{MED}$

= 90 ± 28 Hz vs. morph$_{HIGH}$ = 116 ± 32 Hz, p = 1.2-9, n = 18, Fig. 5A). Concurrently, temporal

precision (VS) at 350 Hz and 1 kHz was significantly reduced under high input variation (1 kHz:

morph$_{MED}$ = 0.65 ± 0.04 vs morph$_{HIGH}$ = 0.62 ± 0.05, p = 1.9e-5, Fig. 5B). Analysis of onset

responses revealed reduced first-spike latencies (1 kHz: morph$_{MED}$ = 4.1 ± 0.14 ms vs. morph$_{HIGH}$

= 3.99 ± 0.17 ms, p = 4.1e-8, Fig. 5C) but increased temporal jitter at the high input variation

condition (1 kHz: morph$_{MED}$ = 0.55 ± 0.11 ms vs. morph$_{HIGH}$ = 0.79 ± 0.1, p = 1.3e-12, Fig. 5D),

mirroring the pattern observed with gamma-distributed variations (Figs. 1 and 3). These effects

were reproduced with the model incorporating morphological input size estimates. These results

suggest that the capacity of input variation to shape GBC output is a direct functional consequence

of the morphological heterogeneity among synaptic inputs.

Morphologically plausible input variation impacts encoding of amplitude-modulated sounds

Finally, to confirm the impact of realistic input variations on the encoding of amplitude-modulated sounds, we repeated the SAM stimulation experiments using two morphologically constrained input variations. Consistent with the results from gamma-distributed variations (Fig. 4), higher input variations elevated overall firing rates across modulation frequencies (main effect of variation: p = 2.9e-10, e.g. 600 Hz: $morph_{MED}$ = 96 ± 38 Hz vs $morph_{HIGH}$ = 125 ± 34 Hz, p = 7.4e-9, n = 19, Fig. 6A), whereas temporal precision was reduced (VS: main effect of variation: p = 3.8e-15, e.g. 600 Hz: $morph_{MED}$ = 0.46 ± 0.04 vs $morph_{HIGH}$ = 0.37 ± 0.02, p = 8.1e-12; CI: main effect of variation: p = 2.5e-18, e.g. 600 Hz: $morph_{MED}$ = 1.43 ± 0.07 vs $morph_{HIGH}$ = 1.26 ± 0.03, p = 1.2e-10, Fig. 6B and C). Rate-coding fidelity (CorrNorm) was enhanced at 200 Hz modulation frequency under high variation (main effect of variation: p = 3.8e-11, 200 Hz: medium variation = 0.89 ± 0.01 vs high variation = 0.93 ± 0.01 Hz, p = 2.9e-17, Fig. 6D). These findings further support the conclusion that the degree of morphological heterogeneity among GBC endbulb inputs dictates the balance between temporal and rate-coding strategies during the processing of both simple and complex sounds.

Discussion

The heterogeneity of synaptic inputs converging onto neurons is a fundamental aspect of neural circuit design. The strength and variability of these inputs profoundly shape neuronal output and, consequently, determine a neuron's computational role within its circuit. In this study, we investigated how such input strength variation affects GBC which receive multiple, variably sized endbulb inputs from the AN. Our findings demonstrated that the degree of variation among these endbulb inputs critically determines GBC firing characteristics. The present results revealed a functional trade-off: increased input variability enhances overall firing rates and improves the encoding of stimulus amplitude modulations. Simultaneously it degrades temporal precision in representing the stimulus fine-structure and increases onset spike jitter. The modeling results, which extended the experimental observations by testing a wider range of input variations, demonstrated that this functional transition occurs continuously. This suggests that morphological variation in synaptic strength allows GBC to operate along a functional continuum shaped by their respective input configurations.

Limitations and biological context

Endbulb terminals are generally well-spaced across the GBC's somatic surface, likely minimizing the interaction between individual inputs (Nicol and Walmsley, 2002; Spirou et al., 2023). Accordingly, our experimental and modeling approaches assumed a linear summation of statistically independent synaptic inputs. While this approach effectively characterized the core phenomena and revealed a robust effect across individual neurons, additional complexities that are not captured by our current approach might exist. For instance, multiple endbulb inputs originating from the same AN fiber or exhibiting highly correlated activity might functionally act as a single, large input, potentially reducing the effective degree of input variation onto the postsynaptic cells. Additionally, while GBC dendrites are typically short, some endbulb arborizations may synapse onto them (Ostapoff and Morest, 1991; Spirou et al., 2005; Gómez-Nieto and Rubio, 2009). Active dendritic properties could provide slower modulation or alter

integration dynamics in a state-dependent manner, which is not currently accounted for in our

model (Häusser and Mel, 2003; Koert and Kuenzel, 2021; Spirou et al., 2023). However, the robust

reproduction of key experimental findings using a simple coincidence-counting model suggests

that input variation is the most fundamental driver of the observed effects. Future work should

investigate how these effects might be altered by somatic and dendritic inhibition and other

modulatory factors, which have been shown to impact temporal precision and sound encoding

(Kopp-Scheinpflug et al., 2002; Dehmel et al., 2010; Nerlich et al., 2014; Keine and Rübsamen,

2015; Goyer et al., 2016; Keine et al., 2016, 2017).

Species-dependent differences in input convergence

The number of endbulbs converging onto GBC varies across species. Our recordings were

performed in Mongolian gerbils, for which detailed anatomical data on input convergence and

variation are currently unavailable. However, the observed effects of input variation were

consistent across recorded cells, suggesting a mechanism independent of the exact number of

endbulbs terminating on a single neuron. Furthermore, the ability of our model to replicate these

effects suggests that the underlying mechanism is robust across species-specific differences in

input number and variation. By constraining input variations with morphological data from mouse

GBC (Spirou et al., 2023), we demonstrated that biologically plausible variations in input sizes are

sufficient to induce the observed effects on GBC coding.

The source of this variation is not well understood. In mice, BC inputs were found to originate from

different spiral ganglion neuron (SGN) subtypes, each exhibiting differential responses to sound

(Wang et al., 2021; Wong et al., 2025). In cats, endbulb inputs onto each individual BC appear to

originate predominantly from the same SGN subtype, with endbulbs originating from type Ia SGN

typically being larger than those from type Ic SGN (Rouiller et al., 1986; Sento and Ryugo, 1989;

Liberman, 1991). If different SGN subtypes systematically provide larger or smaller endbulb inputs

in gerbils, this could introduce synaptic heterogeneity that is not explicitly covered by our current

approach. However, in mice, SGN inputs onto the same BC exhibit comparable short-term

plasticity, and no differences in firing threshold or spontaneous firing rate have been observed in

BCs contacted by different proportions of SGN subtypes (Wong et al., 2025). This finding suggests

that intrinsic variations in synapse size, potentially independent of SGN subtype, may be a

significant factor driving the observed effects.

Functional implications for auditory processing

GBCs provide the primary excitatory input to the MNTB, where their large-diameter axons form

the calyx of Held synaptic terminals. The calyx–MNTB synapse functions as a crucial relay (Moore

and Caspary, 1983; Wenthold et al., 1987; Wu and Kelly, 1993; Borst and Soria van Hoeve, 2012;

Keine and Englitz, 2025), providing well-timed inhibition to several key nuclei in the superior olivary

complex (SOC) involved in sound localization (e.g. medial superior olive, MSO; lateral superior

olive, LSO) and gap detection (superior paraolivary nucleus, SPN) (Spangler et al., 1985; Sommer

et al., 1993; Smith et al., 1998; Srinivasan et al., 2004; Kopp-Scheinpflug et al., 2011).

Our findings, together with simulations based on detailed EM reconstructions (Spirou et al., 2023),

suggest that the GBC population is not functionally uniform but exhibits considerable diversity

driven by AN input variations. GBC receiving endbulb inputs of similar sizes could act as high-

fidelity temporal relays, preserving and even enhancing temporal precision essential for binaural

computations on a sub-millisecond timescale required for sound localization (Yin and Chan, 1990;

Tollin and Yin, 2005; Grothe et al., 2010; Spirou et al., 2023). Conversely, GBC receiving highly

variable inputs, characterized by one or two large suprathreshold endbulbs alongside smaller

ones, may sacrifice some temporal precision to emphasize a more robust rate code. This

enhances their ability to track stimulus envelopes and provides short response latencies, albeit

with increased temporal jitter. Indeed, morphological data from mice, combined with computational

modeling, suggest that approximately half of the GBC operate in a coincidence detection mode,

while the other half show a mixed-mode of operation (Spirou et al., 2023).

The implications of this GBC heterogeneity on downstream targets are intriguing. Although MNTB neurons are typically considered a homogenous group of cells, their calyx of Held inputs exhibit considerable structural and functional heterogeneity independent of tonotopic location. For example, MNTB cells contacted by simple calyces tend to exhibit stronger EPSC depression than those contacted by complex calyces (Grande and Wang, 2011; Fekete et al., 2019), indicating potential differences in release probability that may impact their response to sound. It is tempting to speculate that the observed morphological and functional heterogeneity in GBC input variation might be linked to the variations found in their MNTB targets and, ultimately, relate to the specific functional requirements of the auditory nuclei targeted by MNTB neurons. For instance, MNTB inputs to the MSO might require high temporal precision for interaural time difference (ITD) coding, whereas inputs to the LSO might emphasize firing rate encoding for the calculation of interaural level differences (ILDs). Although direct evidence linking GBC input variation to specific calyx subtypes or SOC targets is currently lacking, our results suggest that input variation is essential for functional heterogeneity within these auditory pathways.

Computational mechanisms of synaptic integration and input variation

Computationally, the effects of input variation can be understood by considering how individual inputs interact with the GBC's AP threshold. High input variation, characterized by a few strong inputs, allows a GBC to operate near the AP threshold driven by a small number of coincident or even individual strong inputs (Xu-Friedman and Regehr, 2005a). While this increases overall spike probability and firing rate, it may also result in stimulus-independent or less precisely timed "false-positive" spikes, contributing to an increased temporal jitter. Conversely, configurations with many uniform small inputs require the near-coincident arrival of multiple inputs, thereby enforcing strict coincidence detection and higher temporal fidelity, albeit at the expense of longer response latencies and a reduced overall firing rate (Rothman et al., 1993; Joris et al., 1994; Rothman and Young, 1996; Xu-Friedman and Regehr, 2005b). This principle of integrating strong and weak synaptic inputs to shape neuronal output is a common strategy observed in various brain regions,

including the cortex and thalamus (Sherman and Guillery, 1998; Rouiller and Welker, 2000; Groh

et al., 2008; Wolff et al., 2021; Govindaiah et al., 2023). Such heterogeneous neuronal populations

have been shown to enhance learning of tasks with temporal structure (Perez-Nieves et al., 2021).

Our findings place GBC within this broader framework, demonstrating how synaptic input variation

tunes synaptic integration to support diverse coding strategies, even at this early stage of sensory

processing (Ashida et al., 2019; Spirou et al., 2023). This heterogeneity would ultimately be critical

for tailoring neuronal information coding to the various computational demands of downstream

auditory circuits.

Resource Availability

Lead contact

Requests for further information and resources should be directed and will be fulfilled by Ivan Milenkovic (ivan.milenkovic@uol.de) or Go Ashida (go.ashida@uol.de)

Materials availability

No unique reagents have been generated in this study.

Data and code availability

All numerical data used to generate figures are available at [DOI will be added] under the terms of Creative Commons Attribution 4.0 License (CC BY 4.0).

Acknowledgments

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2177/1 - Project ID 390895286. and intramural funding to CK (Carl von Ossietzky Universität Oldenburg; FP 2024-099)

Author contributions:

CW: Investigation; **GA:** Conceptualization, Methodology, Software, Verification, Formal Analysis, Resources, Data Curation, Writing – Review & Editing; **CK:** Conceptualization, Methodology, Software, Verification, Formal Analysis, Resources, Data Curation, Writing – Original Draft, Writing – Review & Editing, Visualization, Supervision, Funding acquisition; **IM:** Conceptualization, Resources, Writing – Review & Editing, Supervision, Project administration

Declaration of interests

The authors declare no competing interests

STAR Methods

Experimental Model and Subject Details

Animals and ethical approval

All experiments were performed at the Physiology Laboratories of the Department of Human Medicine at the University of Oldenburg. Experimental protocols were approved and conducted according to the German Animal Welfare Act (TSchG §4) and European Union Directive 2010/63/EU. Mongolian gerbils (*Meriones unguiculatus*) were group-housed under a standard 12h light/dark cycle with unrestricted access to food and water. Experiments were performed on young adult gerbils (postnatal day P20–P25) of both sexes (35 animals, 12 female). All available measures were taken to minimize the number of animals used and alleviate any potential pain or distress.

Method details

Acute brain slice preparation

Following rapid decapitation, acute parasagittal slices (200 µm thickness) containing the CN were prepared using a vibrating microtome (Leica VT1200 S) equipped with zirconia ceramic blades (EF-INZ10, Cadence Blades). Slicing was performed in ice-cold, low-calcium artificial cerebrospinal fluid (aCSF) containing (in mM): 125 NaCl, 2.5 KCl, 0.1 $CaCl_2$, 3 $MgCl_2$, 1.25 NaH_2PO4, 25 $NaHCO_3$, 10 glucose, 3 myo-inositol, 2 sodium pyruvate, and 0.4 ascorbic acid. The aCSF was continuously oxygenated with carbogen gas (5% CO_2 and 95% O_2) and maintained at pH 7.4. Slices were then transferred to a holding chamber containing standard recording solution at near-physiological temperature (36–37°C; identical to the slicing aCSF, except $CaCl_2$ and $MgCl_2$ were adjusted to 1.2 mM and 1 mM, respectively), allowed to recover for 30–45 min, and stored

thereafter until used for recordings. All solutions were prepared using ultrapure water

(0.055 µS/cm).

Electrophysiology

For electrophysiological recordings, slices were placed in a recording chamber continuously

perfused with oxygenated aCSF (~1 mL/min) and maintained at physiological temperature

(~37°C). Both the recording chamber and perfusion inlet were temperature-controlled using a dual

automatic temperature controller (TC-344B, Warner Instruments), and the bath temperature was

monitored throughout the experiments. Neurons within the CN were visualized using an upright

microscope (Axioskop 2 FS plus, Zeiss) equipped with a water-immersion 63x objective (W Plan-

Apochromat 63x/1.0, Cat# 421480-9900-000, Zeiss) and a CCD camera (IR-1000, DAGE-MTI).

Recording pipettes were pulled from borosilicate glass (GB-150F-8P, Science Products) to a

resistance of 3–5 MΩ when filled with an intracellular solution containing (in mM): 120 K-gluconate,

20 KCl, 5 EGTA, 10 HEPES, 0.05 $CaCl_2$, 5 Na_2-phosphocreatine, 4 Mg-ATP, and 0.3 Na-GTP,

supplemented with 0.2% biocytin for post-hoc morphological verification. Whole-cell recordings

were made using a Multiclamp 700B amplifier (Molecular Devices), digitized at a 100 kHz sampling

rate with a Digidata 1440A interface (Molecular Devices) and acquired using pClamp software

(RRID:SCR_011323, version 11.2.2, Molecular Devices). In current-clamp mode, the resting

membrane potential was held near -60 mV using a small bias current (<140 pA). Series resistance

was typically <10 MΩ and was fully compensated via bridge balance. Bridge balance and pipette

capacitance neutralization were carefully adjusted and monitored throughout each recording. All

experiments were performed at 36–37°C.

Simulating auditory nerve fiber input

The spiking activity of AN fibers was simulated using the 2018 version of the auditory periphery

model (Bruce et al., 2018). We used the 'cat' configuration in the model, as AN fiber responses in

gerbils share many similarities with those in cats (Versteegh et al., 2011; Huet et al., 2016;

Heeringa et al., 2023). Based on observations that GBC primarily receive inputs from high-

spontaneous rate (SR) type Ia SGN (Liberman, 1991; Wang et al., 2021; Wong et al., 2025), we

set the SR of the model fibers to 70 spikes/s. Refractory periods were fixed at the midpoint of the

model's default parameter range, representing typical physiological values (absolute: 0.45 ms,

relative: 0.5125 ms).

Based on anatomical studies, showing that GBC receive around 5-12 endbulb inputs in mice and

around 20 in cats (Spirou et al., 2005, 2023), we simulated the convergence of 10 AN inputs for

each target GBC. Because GBC exhibit narrow frequency tuning similar to that of AN fibers

(Rhode and Smith, 1986), all converging simulated AN fibers were assumed to share the same

characteristic frequency (CF). For pure-tone stimuli, the fibers' CF matched the tone frequency;

for amplitude-modulated (AM) sounds, the CF matched the carrier frequency. Sound intensity was

set to 70 dB SPL for pure tones and 25 dB SPL for AM sounds.

Controlling synaptic input variation

To investigate the impact of heterogeneity among converging inputs, we systematically varied the

strength of the 10 simulated AN fibers. Synaptic input variation (v) was quantified as the coefficient

of variation, defined as the standard deviation (s) of the 10 AN input amplitudes divided by their

mean (m; $v = s/m$). For our primary manipulations, sets of 10 AN amplitudes were drawn from a

gamma distribution (shape parameter $a = 1/v^2$, scale parameter $b = mv^2$) to create three distinct

variation conditions: 'no variation' ($v = 0$), 'medium variation' ($v = 0.4$), and 'high variation' ($v = 1$)

(Fig. 1B). These levels were chosen to encompass a broad range of synaptic size heterogeneity,

including the range reported from anatomical estimates of GBC inputs (Spirou et al., 2023).

Average input strength was varied between cells to result in GBC firing rates of >100 spikes/s

during 350 Hz stimulation and kept constant across input variation conditions.

To confirm the biological relevance of our findings, we also implemented synaptic strength

distributions based on the morphological data from (Spirou et al., 2023). Two representative GBC

input data sets were selected: one (GBC21) previously characterized as fitting the coincidence

detection model (CV ≈ 0.33, approximating our 'medium variation', $morph_{MED}$) and another

(GBC18L) fitting the mixed-mode integration model (CV ≈ 0.8, approximating our 'high variation',

$morph_{HIGH}$). To facilitate direct comparison between these two morphologically derived input

variations, we standardized the number of AN inputs to eight for both conditions.

Conductance-clamp implementation

Synaptic conductance traces (G_{syn}) were generated by convolving the simulated AN spike trains

with a stereotypical EPSC waveform. This waveform was modeled as an alpha function with a

time constant of 0.2 ms, and the EPSC amplitude for each of the 10 inputs was scaled according

to the predefined input variation condition. The total G_{syn} represented the linear summation of the

conductances from all simulated AN fibers with the average input strength kept constant between

input variation conditions.

These pre-calculated G_{syn} waveforms were implemented using a conductance-clamp system, as

described previously (Yang et al., 2015). Briefly, the instantaneous membrane potential (V_m) of

the recorded GBC was continuously monitored and digitized using a PCIe-6361 data acquisition

card (National Instruments) connected via a BNC-2120 interface (National Instruments). This V_m

was then fed into a separate computer running mafDC (courtesy of M. A. Xu-Friedman) within Igor

Pro (RRID:SCR_000325, version 8, Wavemetrics). The appropriate AMPA receptor-mediated

current (I_{syn}) was then calculated in real-time (latency: 15 μs, AMPAR cycle time: 8 μs), according

to Ohm's law, assuming a reversal potential (E_{syn}) of 0 mV, and injected back into the GBC via the

recording electrode:

528
$$I_{syn} = G_{syn}(V_m - E_{syn})$$

 Post-hoc morphological identification of GBC

To confirm the identity of recorded cells as GBC based on their morphology and location within

the CN, slices containing biocytin-filled neurons were processed for morphological analysis

following electrophysiological recordings. Immediately after recording, slices were transferred to

a culture dish and fixed with 4% paraformaldehyde (PFA) in 1x phosphate-buffered saline (PBS)

overnight at 4°C. Following fixation, slices were thoroughly rinsed with 1x PBS (6 × 5 min) followed

by 1x PBS containing 0.3% TritonX-100 (PBST, 6 × 5 min). To visualize the biocytin-filled GBC,

slices were incubated for 2.5 h at room temperature with Cy5-conjugated streptavidin (5 µg/mL,

Cat#016-170-084, Jackson ImmunoResearch) diluted in PBST. Final rinses consisted of PBST (6

× 5 min), PBS (2 × 5 min) and ultrapure water (2x5 min).

All incubation and rinsing steps were performed on a tilting shaker in the dark. The slices were

then mounted on glass slides using Aqua-Poly/Mount mounting medium (Cat#18606,

Polysciences, Inc.). Confocal images of recorded neurons were acquired using a Leica SP8

confocal scanning microscope equipped with a 10x (0.3 NA, Cat#11506505) dry objective and a

63x (1.4 NA, Cat#11506350) oil-immersion objective using a 647 nm excitation wavelength.

Globular bushy cell model

Computational simulations of GBC responses were performed using the "adaptive coincidence-

counting model" described previously (Ashida et al., 2019). In this model, each incoming spike

from a simulated AN fiber evoked a simplified rectangular postsynaptic response (Fig. 2A). The

duration (W) of this rectangle remained constant for all inputs, whereas its amplitude (A) was

varied across the 10 inputs based on the specified input variation condition described in an earlier

section. These individual postsynaptic responses were linearly summed to produce the total

integrated input u(t). The model generated an output spike when u(t) exceeded an adaptive

threshold $\theta(t)$. Following an output spike, the model entered an absolute refractory period (R)

during which no further spikes could be triggered.

The adaptive threshold consisted of static and dynamic components: $\theta(t) = \theta_S + \theta_D(t)$. The static

component θ_S was fixed at 1, so that postsynaptic response amplitudes (A) were considered

normalized to this static input threshold. The dynamic component θ_D changed according to the

total input amplitude: $T(d\theta_D/dt) = -\theta_D(t) + S \cdot (u(t))^2$, where T is the time constant and S is the

strength of adaptation (Ashida et al., 2019). The quadratic exponent of u(t) was designed to mimic,

in a simplified manner, the nonlinear activation of various adaptive mechanisms in BC, such as

low-threshold K^+ channels and inhibitory feedback from other CN neurons. This adaptive threshold

mechanism allowed the model to reject poorly synchronized inputs, preventing them from reaching

the threshold (Fig. 2A). The spike generation mechanism implemented in the GBC model is

deterministic and does not produce spike time jitter. To simulate this intrinsic variability, a spike-

generation delay $D_{spike} = D_{fix} + D_{rand}$ was added to each output spike timing. The fixed component

D_{fix} was set to 0.2 ms, and the random component D_{rand} was drawn from an alpha distribution (Xu-

Friedman and Regehr, 2005a) with a time constant of 0.1 ms. This modification was based on the

spike latency data from pure-tone stimulation (Fig 3B5 and 3B6) and resulted in a mean increase

in response latency of 0.4 ms (SD = 0.14 ms).

Model parameters and validation

In addition to the input variation described above, the GBC model used in this study had five free

parameters (Table 1). First, a default parameter set was established that generated exemplary

responses to pure-tone stimuli resembling those observed in our dynamic-clamp experiments

(e.g., PSTHs and raster plots in Fig. 3). To confirm the robustness of the modeling results, we

systematically varied the modeling parameter values around the default set (Table 1, right

column). Each "model instance", representing a unique parameter combination, was tested with

pure-tone stimulation. Only those instances exhibiting GBC-like responses, as defined by the

quantitative criteria described below, were adopted for further simulations.

The adoption criteria were based on our experimental observations (Figs. 1 and 3). Model

instances were required to satisfy the following criteria: (1) spontaneous rate of 10-60 spikes/s;

(2) for 350 Hz stimulation: steady-state driven rate >180 spikes/s and vector strength >0.85; (3)

for 3500 Hz stimulation: steady-state driven rate of 80-180 spikes/s and onset spike probability

>0.95. Furthermore, the PSTH shape near the onset of the 3500 Hz tone was examined to ensure

a primary-like-with-notch (PL_N) spiking pattern, characterized by a sharp peak followed by a notch

(pause of spiking activity) and a smooth recovery towards the steady-state spiking response (Fig.

3A2). Specific criteria for the PL_N response were as follows: (4) the notch appeared within 3 ms

after the onset peak; (5) the notch amplitude (minimum rate in PSTH) was below 50% of the

steady-state spike rate; (6) a second peak in the PSTH was absent or below 300 spikes/s; (7) the

average spike rate at 6-10 ms after the stimulus onset was >90% of the steady-state spike rate;

and (8) a second notch was absent or higher than 50% of the steady-state spike rate. The last

criterion was used to distinguish PL_N-type PSTHs from chopper-type PSTHs (Ashida et al., 2019).

Of the 28,812 model instances tested, 313 (1.1%) met all of these physiological response criteria

to be considered 'GBC' and were used in all subsequent simulations.

Quantification and statistical analysis

Individual GBC were treated as independent biological samples. Aggregated data are presented

as mean ± standard deviation, unless otherwise noted. All comparisons involved within-cell

manipulations (i.e., stimulus type and input variation for the same cell) and were analyzed using

a multifactorial repeated-measures ANOVA (two-way RM ANOVA). Post-hoc multiple

comparisons were corrected using the Bonferroni method (Bonferroni, 1936). A p-value <0.05 was

deemed statistically significant for all interpretations. No significant difference was detected

between sexes regarding firing rate (p = 0.25) or temporal precision (p = 0.86); therefore, data

from both sexes were pooled for further analysis. Sample sizes were determined based on

previous similar studies and the typical variance observed for these electrophysiological

measures. No statistical methods were used to pre-determine sample sizes prior to data collection.

Electrophysiological data analysis

All data analyses were performed using custom-written scripts in MATLAB (RRID:SCR_001622,

Version 2024b, The Mathworks). Action potentials (APs) were detected offline using a voltage

threshold crossing criterion of -20 mV. Detected events were visually inspected and curated to

exclude potential artifacts. For all subsequent analyses, the time of the AP peak was used as the

spike time. Average firing rates were calculated during the period of acoustic stimulation (evoked

rate, 150–200 ms duration) or from periods outside the stimulation window (spontaneous rate, 150

ms duration). Onset probability was assessed within 1.5 ms of the stimulus-evoked response

onset and calculated as the fraction of trials with at least one AP within that time window.

Peristimulus time histograms (PSTH) were constructed by binning spike times (bin width = 0.5 ms)

across 100–200 identical stimulus repetitions. Phase-locking to periodic stimuli (pure tones or

SAM) was quantified with vector strength (VS) (Goldberg and Brown, 1969). To assess spike

timing precision independent of the stimulus waveform, the correlation index (CI) was calculated

as the normalized peak height of the shuffled autocorrelogram, constructed across identical

stimulus repetitions (Joris et al., 2006). For SAM stimuli, the representation of the stimulus

envelope in the GBC firing rate was quantified using the normalized cross-correlation (CorrNorm)

between the stimulus envelope and the corresponding response PSTH, after correcting for the

neuron's response latency (Tolnai et al., 2008; Keine et al., 2016). Note that CorrNorm values

range from 0 to 1; unmodulated responses typically yield a CorrNorm value around 0.82, while

values >0.9 indicate robust tracking of the stimulus envelope.

Figure Legends

Fig. 1: Synaptic input variability increases firing rates at the expense of temporal precision.

A) GBC identification based on morphological and electrophysiological features. **A1)** Left: Low-magnification image of cochlear nucleus parasagittal slice containing a biocytin-filled GBC near the AN root (arrow). AVCN: anteroventral cochlear nucleus; PVCN: posteroventral cochlear nucleus; DCN: dorsal cochlear nucleus; AN: auditory nerve. Right: High-magnification image of the GBC shown on the left, displaying its characteristic round soma and a small number of short dendrites. **A2)** Electrophysiological signature of a GBC (top) responding with one or a few APs at the onset of suprathreshold depolarizing current injection (bottom) and voltage sag in response to hyperpolarizing current injection. **B)** Schematic of the experimental paradigm: Acoustic stimuli (top) were processed by an auditory nerve model to generate spike trains for 10 AN fibers (ANF1-10). These spike trains were convolved with the EPSC waveform to generate conductance templates with varying synaptic strengths ranging from 'no variation' (blue) to high variation (orange) and injected into the GBC. Histograms illustrate input variations for the different conditions used. The GBC response (bottom right) was recorded, and the number and timing of the APs (gray triangles) were analyzed. **C1)** Population data (n = 15) of the average GBC firing rate in response to pure-tone stimuli at 350, 1000, and 3500 Hz under three input conditions (blue: no variation, green: medium variation, orange: high variation). The gray triangles indicate the average values for the AN input. Firing rates increased with input variation for 1000 Hz and 3500 Hz stimulation. **C2)** Temporal precision quantified as vector strength decreases with input variation, particularly at lower stimulation frequencies. Boxplots show median, IQR with whiskers extending to 1.5 × IQR. Individual cells are represented by gray circles. P-values were calculated using 2-way RM ANOVA with Bonferroni post-hoc tests.

Fig. 2: Adaptive coincidence-counting model reproduces experimental data. **A)** Schematic illustration of the GBC model. Top: Example spike trains of four AN fibers highlighting different spike synchrony. Middle: Model responses to the AN fiber inputs with no (blue) and high (orange) input variations. The adaptive threshold changes dynamically according to the recent history of the summed inputs (internal state). GBC model output spikes are generated when the summed inputs exceed the adaptive threshold (blue and orange circles). Note that in the 'high variation' condition spikes can be generated with fewer and less synchronous AN fiber inputs due to the presence of strong individual inputs. **B)** Average firing rates for 313 GBC model instances in response to pure-tone stimuli at 350 Hz, 1000 Hz, and 3500 Hz across input variations. Colored data indicate the input variations used also for our *in vitro* experiments (blue: no variation; green: medium variation; orange: high variation). The model reproduces the experimental trend of higher firing rates for larger input variations at 1000 and 3500 Hz pure-tone stimulation. **C)** Temporal precision quantified as vector strength was slightly reduced with increasing input variation, mirroring the experimental results. Box plots show the range of values from all 313 model instances with thick, medium, and thin boxes indicating 25-75, 10-90, and 0-100 percentiles, respectively.

Fig. 3: Synaptic input variation increases sustained firing rates but reduces onset spike-time precision. **A1)** Peristimulus time histogram (PSTH) of the AN inputs in response to a 3500 Hz pure-tone stimulus. Note the characteristic sharp onset followed by an adapting sustained rate. **A2)** Representative GBC output PSTH in response to the AN input shown in A1 for no variation (blue) and high variation condition (orange). **A3)** Output of the GBC model in response to the same stimulus under no variation (blue) and high variation (orange) conditions, mirroring the experimental data. Insets in A2-A3 show a magnified view of the response onset, highlighting the reduced onset peak under the 'high variation' condition. **B)** Experimental population data (n=17) quantifying GBC response properties for different input variations (blue: no variation, green: medium variation, orange: high variation). **B1)** Raster plot (15 repetitions) of the first spike within 1.5 ms of stimulus onset during 'no variation' (top, blue) and 'high variation' (bottom, orange) conditions. Note the slightly earlier but more variable first spike in the 'high variation' condition. **B2)** Onset spike probability was significantly reduced with input variation, whereas **B3)** sustained firing rates (50-200 ms of the GBC response, and **B4**) spontaneous firing rates increased with input variation. Gray triangles indicate the mean values of the AN input. **B5)** The average first-spike latency decreased with input variation, whereas **B6)** first-spike jitter, defined as the standard deviation of the first-spike latencies, increased with input variation. **C1-C6**). Output of the GBC model quantifying the same parameters as for the experimental data in B1-B6. Colored box plots indicate the input variations used in the *in vitro* experiments. In B) box plots show median, IQR with whiskers extending to 1.5 × IQR. Individual cells are represented by gray circles. P-values were calculated using 2-way RM ANOVA with Bonferroni post-hoc tests. In C), box plots for the modeling data show the range of values from all 313 model instances with thick, medium, and thin boxes indicating 25-75, 10-90, and 0-100 percentiles, respectively.

Fig. 4: Synaptic input variation shapes GBC encoding of amplitude-modulated sounds. A1) Example of simulated AN fiber response to a 200-Hz amplitude-modulated sound (top). The AN firing rate was robustly modulated by the stimulus envelope. Note that the stimulus was aligned with the AN firing rate modulation for display purposes. **A2)** PSTH of representative GBC in response to the AN input shown in A1 under the conditions of 'no variation' (blue) and 'high variation' (orange). Note the broader peaks in the 'high variation' condition. **A3)** GBC model response to the same input variations as shown in A2. The general PSTH shape, including broader peaks in the 'high variation' condition, was well reproduced in the model. **B)** Population data (n=15) quantifying various aspects of the response to amplitude-modulated sounds across three input variations (blue: no variation, green: medium variation, orange: high variation) for modulation frequencies of 200 Hz, 600 Hz and 2000 Hz. For each panel the experimental data are shown on the left and the modeling responses on the right. **B1)** The average firing rates generally increased with input variation across all modulation frequencies. **B2-B3)** Vector strength and correlation index decreased significantly with input variation. **B4-B5)** Entrainment index as well as CorrNorm increased with input variation at low modulation frequencies and remained unchanged at high modulation frequencies. Box plots containing experimental data show median, IQR with whiskers extending to 1.5 × IQR. Individual cells are represented by gray circles. P-values were calculated using 2-way RM ANOVA with Bonferroni post-hoc tests. Box plots for the modeling data show the range of values from all 313 model instances with thick, medium, and thin boxes indicating 25-75, 10-90, and 0-100 percentiles, respectively.

Fig. 5: **Morphological input variation drives GBC functional heterogeneity**. Population data

(n=18) comparing the responses to pure-tone stimulation under two input variation conditions

derived from morphological estimates: 'medium variation' (CV ≈ 0.33, green, GBC21) and 'high

variation' (CV ≈ 0.8, orange, GBC18L) (Spirou et al., 2023). Histogram illustrates AN fiber strength

distribution of eight AN inputs. **A)** Left: High input variation enhanced GBC firing at 1000 Hz and

3500 Hz with no effect found at 350 Hz. Right: GBC model responses using the morphological

input distribution corroborated the experimental results. **B)** Temporal precision was reduced when

GBC receives highly variable inputs. **C)** First-spike latency is slightly reduced at the expense of

an increase in first-spike jitter **(D)** with high input variations. Box plots containing experimental

data show median, IQR with whiskers extending to 1.5 × IQR. Individual cells are represented by

gray circles. P-values were calculated using 2-way RM ANOVA with Bonferroni post-hoc tests.

Box plots for the modeling data show the range of values from all 313 model instances with thick,

medium, and thin boxes indicating 25-75, 10-90, and 0-100 percentiles, respectively.

Fig. 6: Morphologically constrained input variation alters GBC encoding of amplitude-modulated sounds. Population data (n=19) comparing GBC responses to SAM stimuli at different modulation frequencies under two morphologically constrained input variations: 'medium variation' (CV ≈ 0.33, green, GBC21) and 'high variation' (CV ≈ 0.8, orange, GBC18L) (Spirou et al., 2023). High input variations enhanced firing rates **(A)**, decreased vector strength **(B)** and correlation index **(C)**, and improved representation of the stimulus envelope (CorrNorm, **D**) in both the experimental and the model data. Box plots containing experimental data show median, IQR with whiskers extending to 1.5 × IQR. Individual cells are represented by gray circles. P-values were calculated using 2-way RM ANOVA with Bonferroni post-hoc tests. Box plots for the modeling data show the range of values from all 313 model instances with thick, medium, and thin boxes indicating 25-75, 10-90, and 0-100 percentiles, respectively.

Tables

Table 1: Model parameters. The total number of parameter combinations was 12×7×7×7×7 = 28,812. The default
parameter values were used for Figs. 3A3, 3C1, 4A3.

Parameter	Default value	Ranges tested
(mean) Input amplitude *A*	0.38	0.28 − 0.50 (step 0.02)
Input duration (ms) *W*	0.6	0.2 − 1.4 (step 0.2)
Refractory period (ms) *R*	1.5	0.8 − 2.0 (step 0.2)
Adaptation strength *S*	0.8	0.4 − 1.6 (step 0.2)
Adaptation time constant (ms) *T*	0.7	0.3 − 2.1 (step 0.3)

Table 2: Summary of statistical analysis for gamma-distributed input variations

			Variation				
		Stimulus	**no**	**medium**	**high**	**p-value RM ANOVA**	**post-hoc comparisons**
Figure 1 (n=15)	**Firing Rate (Hz)**	**350 Hz**	245 ± 52	239 ± 52	228 ± 39	variation: p=4.6e-5	no vs medium: p=0.1 no vs high: p=0.01 medium vs. high: p=0.05
		1000 Hz	132 ± 48	139 ± 46	171 ± 34		no vs medium: p=0.03 no vs high: p=7.9e-6 medium vs. high: p=6.6e-6
		3500 Hz	119 ± 47	130 ± 48	170 ± 33		no vs medium: p=5.5e-5 no vs high: p=8.9e-7 medium vs. high: p=1.4e-5
	Vector Strength	**350 Hz**	0.86 ± 0.02	0.85 ± 0.01	0.81 ± 0.01	variation: p=3.7e-6	no vs medium: p=0.001 no vs high: p=3.3e-8 medium vs. high: p=4.3e-8
		1000 Hz	0.61 ± 0.08	0.59 ± 0.09	0.57 ± 0.06		no vs medium: p=0.43 no vs high: p=0.005 medium vs. high: p=0.29
		3500 Hz	0.06 ± 0.03	0.05 ± 0.03	0.05 ± 0.02		no vs medium: p=0.57 no vs high: p>0.99 medium vs. high: p>0.99
Figure 3 (n=15)	**Onset probability**	**3500 Hz**	0.97 ± 0.02	0.96 ± 0.02	0.94 ± 0.02	p=3.5e-6	no vs medium: p=0.42 no vs high: p=7.3e-5 medium vs. high: p=0.0004
	Sustained firing rate (Hz)	**3500 Hz**	71.7 ± 26.8	86.6 ± 29.7	130.8 ± 28.2	p=8.7e-16	no vs medium: p=5.8e-5 no vs high: p=3.2e-11 medium vs. high: p=2.7e-12
	Spontaneous firing rate (Hz)	**3500 Hz**	24.83 ± 14.31	34.82 ± 17.8	64.96 ± 14.64	p=2.4e-17	no vs medium: p=0.0009 no vs high: p=4.9e-11 medium vs. high: p=1.8e-9
	First-spike latency (ms)	**3500 Hz**	3 ± 0.14	2.95 ± 0.12	2.92 ± 0.15	p=0.001	no vs medium: p=0.009 no vs high: p=0.007 medium vs. high: p=0.21
	First-spike jitter (ms)	**3500 Hz**	0.42 ± 0.12	0.52 ± 0.17	0.63 ± 0.21	p=5.84e-6	no vs medium: p=0.001 no vs high: p=7.4e-5 medium vs. high: p=3e-4
Figure 4 (n=15)	**Firing Rate (Hz)**	**200 Hz**	130 ± 36	131 ± 33	145 ± 23	variation: p=3e-10	no vs medium: p>0.99 no vs high: p=0.01 medium vs. high: p=0.002
		600 Hz	67 ± 50	77 ± 50	115 ± 38		no vs medium: p=3e-4 no vs high: p=1.47e-7 medium vs. high: p=3.5e-6
		2000 Hz	65 ± 50	71 ± 46	111 ± 43		no vs medium: p=0.12 no vs high: p=7.8e-9 medium vs. high: p=1.4e-9
	Vector Strength	**200 Hz**	0.84 ± 0.03	0.83 ± 0.02	0.76 ± 0.01	variation: p=5.7e-14	no vs medium: p=0.006 no vs high: p=9.3e-10 medium vs. high: p=1.6e-12
		600 Hz	0.51 ± 0.06	0.48 ± 0.05	0.37 ± 0.03		no vs medium: p=2.2e-6 no vs high: p=7.2e-10 medium vs. high: p=9.6e-10
		2000 Hz	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0		no vs medium: p=0.84 no vs high: p>0.99

					medium vs. high: p=0.1
Correlation Index	200 Hz	3.06 ± 0.19	2.93 ± 0.1	2.39 ± 0.06	no vs medium: p=4e-4 no vs high: p=3.7e-10 medium vs. high: p=1.3e-12
	600 Hz	1.53 ± 0.12	1.46 ± 0.1	1.26 ± 0.04	no vs medium: p=6.2e-6 no vs high: p=3e-8 medium vs. high: p=2.6e-8
	2000 Hz	1.01 ± 0.02	1.01 ± 0.02	1 ± 0.01	no vs medium: p>0.99 no vs high: p=0.5 medium vs. high: p=0.06
Entrainment Index	200 Hz	0.6 ± 0.17	0.62 ± 0.17	0.68 ± 0.1	no vs medium: p=0.87 no vs high: p=0.01 medium vs. high: p=0.04
	600 Hz	0.01 ± 0.01	0.01 ± 0.03	0.04 ± 0.05	no vs medium: p=0.36 no vs high: p=0.005 medium vs. high: p=0.001
	2000 Hz	0 ± 0	0 ± 0	0 ± 0	no vs medium: p=0.44 no vs high: p=0.64 medium vs. high: p>0.99
CorrNorm	200 Hz	0.87 ± 0.02	0.88 ± 0.01	0.93 ± 0.01	no vs medium: p=0.004 no vs high: p=2.5e-10 medium vs. high: p=6.3e-11
	600 Hz	0.99 ± 0.01	0.99 ± 0	0.98 ± 0.01	no vs medium: p=0.027 no vs high: p=0.5 medium vs. high: p=0.003
	2000 Hz	0.82 ± 0.01	0.82 ± 0.01	0.82 ± 0	no vs medium: p>0.99 no vs high: p=0.42 medium vs. high: p=0.26

The variation column values: Correlation Index: variation: p=5.9e-13; Entrainment Index: variation: p=4.7e-6; CorrNorm: variation: p=1.8e-10.

744 *Table 3: Summary of statistical analysis for morphologically-derived input variations*

		Stimulus	morph$_{MED}$	morph$_{HIGH}$		post-hoc comparison (morph$_{MED}$ vs morph$_{HIGH}$)
Figure 5 (n=18)	Firing Rate (Hz)	350 Hz	149 ± 40	155 ± 31	p=4.6e-11	p=0.1
		1000 Hz	90 ± 28	116 ± 32		p=1.2e-9
		3500 Hz	87 ± 28	119 ± 30		p=3.5e-11
	Vector Strength	350 Hz	0.86 ± 0.01	0.82 ± 0.01	p=5.9e9	p=4.1e-11
		1000 Hz	0.65 ± 0.04	0.62 ± 0.05		p=1.9e-5
		3500 Hz	0.02 ± 0.01	0.02 ± 0.01		p=0.38
	First-spike latency (ms)	350 Hz	4.67 ± 0.17	4.57 ± 0.14	p=1.2e-6	p=0.0015
		1000 Hz	4.09 ± 0.14	3.99 ± 0.17		p=4.1e-8
		3500 Hz	3.05 ± 0.09	3.04 ± 0.12		p=0.51
	First-spike jitter (ms)	350 Hz	0.46 ± 0.2	0.68 ± 0.14	p=7.1e-12	p=2e-4
		1000 Hz	0.55 ± 0.11	0.79 ± 0.11		p=1.3e-12
		3500 Hz	0.34 ± 0.03	0.45 ± 0.08		p=3e-5
Figure 6 (n=19)	Firing Rate (Hz)	200 Hz	134 ± 30	142 ± 27	p=2.9e-10	p=2e-5
		600 Hz	96 ± 38	125 ± 34		p=7.4e-9
		2000 Hz	89 ± 36	125 ± 31		p=6.2e-11
	Vector Strength	200 Hz	0.81 ± 0.02	0.75 ± 0.02	p=3.8e-15	p=2e-17
		600 Hz	0.46 ± 0.04	0.37 ± 0.02		p=8.1e-12
		2000 Hz	0.01 ± 0.01	0.01 ± 0		p=0.07
	Correlation index	200 Hz	2.78 ± 0.11	2.33 ± 0.1	p=2.5e-18	p=3.7e-19
		600 Hz	1.43 ± 0.07	1.26 ± 0.03		p=1.2e-10
		2000 Hz	1 ± 0.01	1 ± 0		p=0.19
	CorrNorm	200 Hz	0.89 ± 0.01	0.93 ± 0.01	p=3.8e-11	p=2.9e-17
		600 Hz	0.99 ± 0.01	0.98 ± 0		p=0.002
		2000 Hz	0.81 ± 0.01	0.82 ± 0.01		p=0.04

References

Ashida G, Heinermann HT, Kretzberg J (2019) Neuronal population model of globular bushy cells covering unit-to-unit variability. PLoS Comput Biol 15:e1007563.

Blackburn CC, Sachs MB (1989) Classification of unit types in the anteroventral cochlear nucleus: PST histograms and regularity analysis. J Neurophysiol 62:1303–1329.

Bonferroni CE (1936) Teoria statistica delle classi e calcolo delle probabilità. Pubblicazioni del R Istituto Superiore di Scienze Economiche e Commerciali di Firenze 8:3–62.

Borst JGG, Soria van Hoeve J (2012) The calyx of Held synapse: from model synapse to auditory relay. Annu Rev Physiol 74:199–224.

Britt R, Starr A (1976) Synaptic events and discharge patterns of cochlear nucleus cells. I. Steady-frequency tone bursts. J Neurophysiol 39:162–178.

Bruce IC, Erfani Y, Zilany MSA (2018) A phenomenological model of the synapse between the inner hair cell and auditory nerve: Implications of limited neurotransmitter release sites. Hear Res 360:40–54.

Calkins DJ, Schein SJ, Tsukamoto Y, Sterling P (1994) M and L cones in macaque fovea connect to midget ganglion cells by different numbers of excitatory synapses. Nature 371:70–72.

Cant NB, Morest DK (1979) Organization of the neurons in the anterior division of the anteroventral cochlear nucleus of the cat. Light-microscopic observations. Neuroscience 4:1909–1923.

Cao X-J, Oertel D (2010) Auditory nerve fibers excite targets through synapses that vary in convergence, strength, and short-term plasticity. J Neurophysiol 104:2308–2320.

Dehmel S, Kopp-Scheinpflug C, Weick M, Dörrscheidt GJGJ, Rübsamen R (2010) Transmission of phase-coupling accuracy from the auditory nerve to spherical bushy cells in the Mongolian gerbil. Hear Res 268:234–249.

Doucet JR, Ryugo DK (2006) Structural and functional classes of multipolar cells in the ventral cochlear nucleus. Anat Rec A Discov Mol Cell Evol Biol 288:331–344.

Drullman R, Festen JM, Houtgast T (1996) Effect of temporal modulation reduction on spectral contrasts in speech. J Acoust Soc Am 99:2358–2364.

Englitz B, Tolnai S, Typlt M, Jost J, Rübsamen R (2009) Reliability of synaptic transmission at the synapses of held in vivo under acoustic stimulation. PLoS One 4:e7014.

Fekete A, Nakamura Y, Yang Y-M, Herlitze S, Mark MD, DiGregorio DA, Wang L-Y (2019) Underpinning heterogeneity in synaptic transmission by presynaptic ensembles of distinct morphological modules. Nat Commun 10:826.

Fukui I, Ohmori H (2004) Tonotopic gradients of membrane and synaptic properties for neurons of the chicken nucleus magnocellularis. J Neurosci 24:7514–7523.

Goldberg JM, Brown PB (1969) Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: some physiological mechanisms of sound localization. J Neurophysiol 32:613–636.

Gómez-Nieto R, Rubio ME (2009) A bushy cell network in the rat ventral cochlear nucleus. J Comp Neurol 516:241–263.

Govindaiah G, Fox MA, Guido W (2023) Pattern of driver-like input onto neurons of the mouse ventral lateral geniculate nucleus. eNeuro 10:ENEURO.0386-22.2022.

Goyer D, Kurth S, Gillet C, Keine C, Rübsamen R, Kuenzel T (2016) Slow cholinergic modulation of spike probability in ultra-fast time-coding sensory neurons. eNeuro 3 Available at: http://eneuro.org/cgi/doi/10.1523/ENEURO.0186-16.2016.

Grande G, Wang L-Y (2011) Morphological and functional continuum underlying heterogeneity in the spiking fidelity at the calyx of Held synapse in vitro. J Neurosci 31:13386–13399.

Groh A, de Kock CPJ, Wimmer VC, Sakmann B, Kuner T (2008) Driver or coincidence detector: modal switch of a corticothalamic giant synapse controlled by spontaneous activity and short-term depression. J Neurosci 28:9652–9663.

Grothe B, Pecka M, McAlpine D (2010) Mechanisms of sound localization in mammals. Physiol Rev 90:983–1012.

Häusser M, Mel B (2003) Dendrites: bug or feature? Curr Opin Neurobiol 13:372–383.

Heeringa AN (2024) Single-unit data for sensory neuroscience: Responses from the auditory nerve of young-adult and aging gerbils. Sci Data 11:1–13.

Heeringa AN, Teske F, Ashida G, Köppl C (2023) Cochlear aging disrupts the correlation between spontaneous rate and sound-level coding in auditory nerve fibers. J Neurophysiol 130:736–750.

Heeringa AN, Zhang L, Ashida G, Beutelmann R, Steenken F, Köppl C (2020) Temporal Coding of Single Auditory Nerve Fibers Is Not Degraded in Aging Gerbils. J Neurosci 40:343–354.

Huang CG, Chacron MJ (2016) Optimized parallel coding of second-order stimulus features by heterogeneous neural populations. J Neurosci 36:9859–9872.

Huet A, Batrel C, Tang Y, Desmadryl G, Wang J, Puel J-L, Bourien J (2016) Sound coding in the auditory nerve of gerbils. Hear Res 338:32–39.

Jacoby RA, Wiechmann AF, Amara SG, Leighton BH, Marshak DW (2000) Diffuse bipolar cells provide input to OFF parasol ganglion cells in the macaque retina. J Comp Neurol 416:6–18.

Johnson DH (1980) The relationship between spike rate and synchrony in responses of auditory-nerve fibers to single tones. J Acoust Soc Am 68:1115–1122.

Joris PX, Carney LH, Smith PH, Yin TCT (1994) Enhancement of neural synchronization in the anteroventral cochlear nucleus. I. Responses to tones at the characteristic frequency. J Neurophysiol 71:1022–1036.

Joris PX, Louage DH, Cardoen L, van der Heijden M (2006) Correlation index: a new metric to quantify temporal coding. Hear Res 216–217:19–30.

Joris PX, Schreiner CE, Rees A (2004) Neural processing of amplitude-modulated sounds. Physiol Rev 84:541–577.

Joris PX, Smith PH (2008) The volley theory and the spherical cell puzzle. Neuroscience 154:65–76.

Joris PX, Yin TCT (1992) Responses to amplitude-modulated tones in the auditory nerve of the cat. J Acoust Soc Am 91:215–232.

Keine C, Englitz B (2025) Cellular and synaptic specializations for sub-millisecond precision in the mammalian auditory brainstem. Front Cell Neurosci 19:1568506.

Keine C, Rübsamen R (2015) Inhibition shapes acoustic responsiveness in spherical bushy cells. J Neurosci 35:8579–8592.

Keine C, Rübsamen R, Englitz B (2016) Inhibition in the auditory brainstem enhances signal representation and regulates gain in complex acoustic environments. Elife 5:19295.

Keine C, Rübsamen R, Englitz B (2017) Signal integration at spherical bushy cells enhances representation of temporal structure but limits its range. Elife 6:29639.

Kessler D, Carr CE, Kretzberg J, Ashida G (2021) Theoretical relationship between two measures of spike synchrony: Correlation index and vector strength. Front Neurosci 15:761826.

Koert E, Kuenzel T (2021) Small dendritic synapses enhance temporal coding in a model of cochlear nucleus bushy cells. J Neurophysiol 125:915–937.

Kolb H, Marshak D (2003) The midget pathways of the primate retina. Doc Ophthalmol 106:67–81.

Köppl C (1994) Auditory nerve terminals in the cochlear nucleus magnocellularis: differences between low and high frequencies. J Comp Neurol 339:438–446.

Kopp-Scheinpflug C, Dehmel S, Dörrscheidt GJ, Rübsamen R (2002) Interaction of excitation and inhibition in anteroventral cochlear nucleus neurons that receive large endbulb synaptic endings. J Neurosci 22:11004–11018.

Kopp-Scheinpflug C, Tozer AJB, Robinson SW, Tempel BL, Hennig MH, Forsythe ID (2011) The sound of silence: ionic mechanisms encoding sound termination. Neuron 71:911–925.

Kreeger LJ, Honnuraiah S, Maeker S, Shea S, Fishell G, Goodrich L (2025) An anatomical and physiological basis for flexible coincidence detection in the auditory system. Elife 13 Available at: https://elifesciences.org/reviewed-preprints/100492.

Liberman MC (1991) Central projections of auditory-nerve fibers of differing spontaneous rate. I. Anteroventral cochlear nucleus. J Comp Neurol 313:240–258.

McGinley MJ, Liberman MC, Bal R, Oertel D (2012) Generating synchrony from the asynchronous: compensation for cochlear traveling wave delays by the dendrites of individual brainstem neurons. J Neurosci 32:9301–9311.

Moore MJ, Caspary DM (1983) Strychnine blocks binaural inhibition in lateral superior olivary neurons. J Neurosci 3:237–242.

Nagayama S, Takahashi YK, Yoshihara Y, Mori K (2004) Mitral and tufted cells differ in the decoding manner of odor maps in the rat olfactory bulb. J Neurophysiol 91:2532–2540.

Nerlich J, Kuenzel T, Keine C, Korenic A, Rübsamen R, Milenkovic I (2014) Dynamic fidelity control to the central auditory system: synergistic glycine/GABAergic inhibition in the cochlear nucleus. J Neurosci 34:11604–11620.

Nicol MJ, Walmsley B (2002) Ultrastructural basis of synaptic transmission between endbulbs of Held and bushy cells in the rat cochlear nucleus. J Physiol 539:713–723.

Oertel D (1999) The role of timing in the brain stem auditory nuclei of vertebrates. Annu Rev Physiol 61:497–519.

Oline SN, Ashida G, Burger RM (2016) Tonotopic optimization for temporal processing in the cochlear nucleus. J Neurosci 36: 8500–8515.

Osen KK (1969) Cytoarchitecture of the cochlear nuclei in the cat. J Comp Neurol 136:453–484.

Ostapoff EM, Morest DK (1991) Synaptic organization of globular bushy cells in the ventral cochlear nucleus of the cat: a quantitative study. J Comp Neurol 314:598–613.

Palmer a. R, Russell IJ (1986) Phase-locking in the cochlear nerve of the guinea-pig and its relation to the receptor potential of inner hair-cells. Hear Res 24:1–15.

Palmer AR, Winter IM, Darwin CJ (1986) The representation of steady-state vowel sounds in the temporal discharge patterns of the guinea pig cochlear nerve and primarylike cochlear nucleus neurons. J Acoust Soc Am 79:100–113.

Perez-Nieves N, Leung VCH, Dragotti PL, Goodman DFM (2021) Neural heterogeneity promotes robust learning. Nat Commun 12:5791.

Rhode WS, Greenberg S (1994) Encoding of amplitude modulation in the cochlear nucleus of the cat. J Neurophysiol 71:1797–1825.

Rhode WS, Smith PH (1986) Encoding timing and intensity in the ventral cochlear nucleus of the cat. J Neurophysiol 56:261–286.

Ritz LA, Brownell WE (1982) Single unit analysis of the posteroventral cochlear nucleus of the decerebrate cat. Neuroscience 7:1995–2010.

Rothman JS, Young ED, Manis PB (1993) Convergence of auditory nerve fibers onto bushy cells in the ventral cochlear nucleus: implications of a computational model. J Neurophysiol 70:2562–2583.

Rothman JSS, Young EDD (1996) Enhancement of neural synchronization in computational models of ventral cochlear nucleus bushy cells. Aud Neurosci 2:47–62.

Rouiller EM, Cronin-Schreiber R, Fekete DM, Ryugo DK (1986) The central projections of intracellularly labeled auditory nerve fibers in cats: an analysis of terminal morphology. J Comp Neurol 249:261–278.

Rouiller EM, Welker E (2000) A comparative analysis of the morphology of corticothalamic projections in mammals. Brain Res Bull 53:727–741.

Schneggenburger R, Forsythe ID (2006) The calyx of Held. Cell Tissue Res 326:311–337.

Sento S, Ryugo DK (1989) Endbulbs of held and spherical bushy cells in cats: morphological correlates with physiological properties. J Comp Neurol 280:553–562.

Sherman SM, Guillery RW (1998) On the actions that one nerve cell can have on another: distinguishing "drivers" from "modulators." Proc Natl Acad Sci U S A 95:7121–7126.

Smith PH, Joris PX, Yin TCT, J.oris PX (1998) Anatomy and physiology of principal cells of the medial nucleus of the trapezoid body (MNTB) of the cat. J Neurophysiol 79:3127–3142.

Smith PH, Rhode WS (1987) Characterization of HRP-labeled globular bushy cells in the cat anteroventral cochlear nucleus. J Comp Neurol 266:360–375.

Smith RL, Brachman ML (1980) Response modulation of auditory-nerve fibers by AM stimuli: effects of average intensity. Hear Res 2:123–133.

Sommer I, Lingenhöhl K, Friauf E (1993) Principal cells of the rat medial nucleus of the trapezoid body: an intracellular in vivo study of their physiology and morphology. Exp Brain Res:223–239.

Spangler KM, Warr WB, Henkel CK (1985) The projections of principal cells of the medial nucleus of the trapezoid body in the cat. J Comp Neurol 238:249–262.

Spirou GA, Brownell WE, Zidanic M (1990) Recordings from cat trapezoid body and HRP labeling of globular bushy cell axons. J Neurophysiol 63:1169–1190.

Spirou GA, Kersting MG, Carr S, Razzaq B, Alves-Pinto CY, Dawson M, Ellisman MH, Manis PB (2023) High-resolution volumetric imaging constrains compartmental models to explore synaptic integration and temporal processing by cochlear nucleus globular bushy cells. Elife 12:1–44.

Spirou GA, Rager J, Manis PB (2005) Convergence of auditory-nerve fiber projections onto globular bushy cells. Neuroscience 136:843–863.

Srinivasan G, Friauf E, Löhrke S (2004) Functional glutamatergic and glycinergic inputs to several superior olivary nuclei of the rat revealed by optical imaging. Neuroscience 128:617–634.

Tollin DJ, Yin TCT (2005) Interaural phase and level difference sensitivity in low-frequency neurons in the lateral superior olive. J Neurosci 25:10648–10657.

Tolnai S, Englitz B, Kopp-Scheinpflug C, Dehmel S, Jost J, Rübsamen R (2008) Dynamic coupling of excitatory and inhibitory responses in the medial nucleus of the trapezoid body. Eur J Neurosci 27:3191–3204.

Van Tasell DJ, Soli SD, Kirby VM, Widin GP (1987) Speech waveform envelope cues for consonant recognition. J Acoust Soc Am 82:1152–1161.

Versteegh CPC, Meenderink SWF, van der Heijden M (2011) Response characteristics in the apex of the gerbil cochlea studied through auditory nerve recordings. J Assoc Res Otolaryngol 12:301–316.

von Gersdorff H, Borst JGG (2002) Short-term plasticity at the calyx of Held. Nat Rev Neurosci 3:53–64.

Wang M, Zhang C, Lin S, Wang Y, Seicol BJ, Ariss RW, Xie R (2021) Biased auditory nerve central synaptopathy is associated with age-related hearing loss. J Physiol 599:1833–1854.

Wang X, Sachs MB (1994) Neural encoding of single-formant stimuli in the cat. II. Responses of anteroventral cochlear nucleus units. J Neurophysiol 71:59–78.

Wei L, Karino S, Verschooten E, Joris PX (2017) Enhancement of phase-locking in rodents. I. An axonal recording study in gerbil. J Neurophysiol 118:2009–2023.

Wei L, Verschooten E, Joris PX (2023) Enhancement of phase-locking in rodents. II. An axonal recording study in chinchilla. J Neurophysiol 130:751–767.

Wenthold RJ, Huie D, Altschuler R a., Reeks K a. (1987) Glycine immunoreactivity localized in the cochlear nucleus and superior olivary complex. Neuroscience 22:897–912.

Wolff M, Morceau S, Folkard R, Martin-Cortecero J, Groh A (2021) A thalamic bridge from sensory perception to cognition. Neurosci Biobehav Rev 120:222–235.

Wong NF, Brongo SE, Forero EA, Sun S, Cook CJ, Lauer AM, Müller U, Xu-Friedman MA (2025) Convergence of type 1 spiral ganglion neuron subtypes onto principal neurons of the anteroventral cochlear nucleus. J Neurosci 45:e1507242024.

Wu SH, Kelly JB (1993) Response of neurons in the lateral superior olive and medial nucleus of the trapezoid body to repetitive stimulation: Intracellular and extracellular recordings from mouse brain slice. Hear Res 68:189–201.

Xie R, Manis PB (2017) Radiate and planar multipolar neurons of the mouse anteroventral cochlear nucleus: Intrinsic excitability and characterization of their auditory nerve input. Front Neural Circuits 11:77.

Xu-Friedman MA, Regehr WG (2005a) Dynamic-clamp analysis of the effects of convergence on spike timing. II. Few synaptic inputs. J Neurophysiol 94:2526–2534.

Xu-Friedman MA, Regehr WG (2005b) Dynamic-clamp analysis of the effects of convergence on spike timing. I. Many synaptic inputs. J Neurophysiol 94:2512–2525.

Yang Y, Adowski T, Ramamurthy B, Neef A, Xu-Friedman MA (2015) High-speed dynamic-clamp interface. J Neurophysiol 113:2713–2720.

Yin TCT, Chan JCK (1990) Interaural time sensitivity in medial superior olive of cat. J Neurophysiol 64:465–488.