Distinct sub-MIC kill kinetics of Cu and Ag in *Escherichia coli*

Merilin Rosenberg[1], Sigrit Umerov[1], Carmen Marianne Teär[1], Angela Ivask[1]

[1] Institute of Molecular and Cell Biology, University of Tartu, Estonia

Correspondence: merilin.rosenberg@ut.ee

Abstract

Copper and silver are well-known and widely used antimicrobial metals that are often considered to employ diverse overlapping biocidal mechanisms of action and induce corresponding bacterial defense responses. Exposure to antimicrobial metals at concentrations below the minimal inhibitory concentration (sub-MIC) are widespread in natural, clinical and built environments and can shape evolutionary trajectories of the affected microbes that could lead to antimicrobial tolerance and/or resistance. By analyzing growth and kill kinetics *of Escherichia coli* under copper or silver exposure we observed that sub-MIC copper concentrations resulted in a lasting dose-dependent slowing of exponential growth with reduced yield while silver seemed to cause dose-dependent growth delay without substantially affecting exponential growth or yield. Time-kill experiments revealed minimal loss of viability in early copper exposure while in case of silver a rapid dose-dependent transient killing followed by normal exponential regrowth of the survivors was observed, underlying the seemingly dose-dependently extended lag phase durations. Distinguishing conditions that select for antimicrobial resistance (sustained growth) versus tolerance (survival without growth) is essential for antimicrobial stewardship as acquiring tolerance is considered a steppingstone towards developing resistance. Our results suggest that short-term survival of the initial killing by silver is sufficient for selective advantage while maintaining energy-intensive enhanced growth in the presence of copper is needed to gain competitive benefit over the general population. The findings highlight new and known challenges in antimicrobial characterization and risk assessment of metal-based formulations by using wide-spread non-kinetic endpoint assays such as MIC.

Main text

Silver and copper are widely used in various antimicrobial applications with the intent to kill or limit the spread of potentially pathogenic microbes, yet their mechanisms of action are not fully understood. Cellular targets of the antibacterial activity of copper and silver as well as the corresponding bacterial defenses are often considered overlapping due to similar coordination chemistry and sulfhydryl reactivity, leading to shared efflux and sequestration mechanisms (1–5). *E.g.*, the CusCFBA system exports both Cu(I) and Ag(I) and is induced by CusRS (5). Unlike toxic silver, copper is an essential micronutrient tightly regulated by homeostasis but becomes toxic in excess (1, 6). A key difference is Cu(I)/Cu(II) redox activity, which generates reactive oxygen species and causes direct damage such as lipid peroxidation (7). We have previously found that silver and copper also differ in their relative toxicity to bacteria depending on exposure conditions. *E. coli* tolerates higher copper ion concentrations

(millimolar range) than silver ion concentrations (micromolar range) in liquid exposure media but survives longer on silver surfaces than on copper surfaces in semi-dry exposure conditions (8, 9). These contrasts suggest distinct antibacterial mechanisms or condition-dependent defense responses.

Antibacterial activity is typically measured and compared via static endpoints not kinetic assays. Minimal inhibitory concentration (MIC) indicates presence or absence of growth after standardized exposure while minimal biocidal concentration (MBC) only reflects endpoint mortality (>99.9%) (10, 11). However, endpoint assays can mask regrowth after transient killing. To capture dynamics, where an antimicrobial agent is quickly detoxified and/or causes transient killing that allows survivors to regrow before the assay endpoint, growth and kill kinetics assays in well-defined experimental conditions should be used. Implications of the strain, media and exposure conditions selection in the current study are further discussed in Supplementary Note 1.

Silver and copper affect *E. coli* growth kinetics differently. To study the effects of copper and silver on the growth and viability of *E. coli*, exponential cultures were exposed to 0.0625 to1 times the MIC value increments of the respective metal salts in MOPS-buffered defined medium (Fig. 1; detailed methods of the growth and time-kill experiments presented can be found in Supplementary Note 2). At comparable sub-MIC concentration ranges, $CuSO_4$ increased doubling time and reduced biomass yield of *E. coli* in a dose-dependent manner (Fig. 1A, Supplementary Fig. S1 and S2). In contrast, $AgNO_3$ seemingly prolonged lag time with little effect on doubling time or yield (Fig. 1B, Supplementary Fig. S1 and S2).

Silver and copper cause distinctly different sub-MIC kill kinetics during the ostensible lag phase. Copper exposure resulted in minimal early killing in addition to lasting growth inhibition (Fig. 1A and 1C), whereas silver caused rapid dose-dependent up to over 4-log reduction in viable counts prior to resumption of normal exponential growth (Fig. 1B and 1D). Prolonged physiological lag time can signal a stress response (12, 13) and, without the time-kill data in Fig. 1D, could plausibly account for the delayed growth in Fig. 1B. A decrease in inoculum density also increases population-level lag phase duration - a phenomenon commonly known as the inoculum effect (12, 14). The latter was also evident for *E. coli* under our growth-assay conditions (Supplementary Figs. S1 and S4). Therefore, dose-dependently increased lag phase duration under antimicrobial exposures as in case of silver in Fig 1B, could be primarily attributable to sub-MIC killing and reduced viable cell count at resumption of growth as illustrated on Fig. 1D. In case of observed lag shifts, growth kinetics should be complemented by sub-MIC tolerance assays to discriminate partial killing from physiologically prolonged lag phase.

Figure 1. Growth (A, B) and kill (C, D) kinetics of *Escherichia coli* BW25113 exposed to $CuSO_4$ (A, C) or $AgNO_3$ (B, D) in the range of 0.0625-1 xMIC value of 8 mM Cu or 1.95 μM Ag. Mean and SD of 3 (C, D) to 4 (A, B) biological replicates are shown. All growth curves from individual experiments represented on panels A and B can be found in Supplementary Fig S3, correlation between metal concentration and growth parameters in Supplementary Fig S2 and numerical data in Supplementary Data file. Grey shading on A and B denotes standard deviation and on C and D value range below post-exposure colony counting detection limit with no colonies detected.

The results suggest that repeated or prolonged sub-MIC copper and silver exposures could differentially select for tolerance and resistance. Tolerance—survival without growth—is easier to develop than resistance, which requires enhanced growth in the presence of an antimicrobial agent. Tolerance is thus considered a precursor to resistance (15, 16), offering survivors time to acquire adaptive traits. While both silver and copper are used in commercial antimicrobial products (*e.g.* wound care, textiles, touch surface materials) copper resistance is more studied in the context of environmental pollution or colonization of the indoor environment. Silver resistance, on the other hand, is a well-known challenge in *e.g.* burn wound care (17, 18), necessitating combined treatment strategies. Our results suggest that short-term survival of the initial killing by silver is sufficient to gain competitive benefit over the general population while in the presence of copper overcoming growth inhibition and continuous upkeep of the energy-intensive advantage is needed. This raises the question of whether silver-based antimicrobial applications generally pose a higher risk for the development of tolerance and resistance compared to copper-based applications, due to the distinct growth inhibition and kill kinetics of copper and silver. It also prompts further investigation into how these differences influence the co-selection of antibiotic tolerance and

development of antibiotic resistance in medical applications. In conclusion, our observations highlight new and known challenges in assessing antimicrobial characteristics by using wide-spread endpoint assays such as MIC to make decisions guiding the design and risk assessment of antimicrobial formulations in different fields of application.

Acknowledgements

This study was supported by Estonian Research Council (PRG1496) and European Commission Twinning project FAST-Real (101159721). The study received funding from the Estonian Ministry of Education and Research under projects TK210 and TEM-TA55.

References

1. Giachino A, Waldron KJ. 2020. Copper tolerance in bacteria requires the activation of multiple accessory pathways. Mol Microbiol 114:377–390.

2. Terzioğlu E, Arslan M, Balaban BG, Çakar ZP. 2022. Microbial silver resistance mechanisms: recent developments. World J Microbiol Biotechnol 38:158.

3. Zuily L, Lahrach N, Fassler R, Genest O, Faller P, Sénèque O, Denis Y, Castanié-Cornet M-P, Genevaux P, Jakob U, Reichmann D, Giudici-Orticoni M-T, Ilbert M. 2022. Copper Induces Protein Aggregation, a Toxic Process Compensated by Molecular Chaperones. mBio 13:e03251-21.

4. Li H, Xu H. 2024. Mechanisms of bacterial resistance to environmental silver and antimicrobial strategies for silver: A review. Environ Res 248:118313.

5. Delmar JA, Su C-C, Yu EW. 2015. Heavy metal transport by the CusCFBA efflux system. Protein Sci Publ Protein Soc 24:1720–1736.

6. Hyre A, Casanova-Hampton K, Subashchandrabose S. 2021. Copper Homeostatic Mechanisms and Their Role in the Virulence of Escherichia coli and Salmonella enterica. EcoSal Plus 9:eESP00142020.

7. Hong R, Kang TY, Michels CA, Gadura N. 2012. Membrane Lipid Peroxidation in Copper Alloy-Mediated Contact Killing of Escherichia coli. Appl Environ Microbiol 78:1776–1784.

8. Kaur H, Rosenberg M, Kook M, Danilian D, Kisand V, Ivask A. 2023. Antibacterial activity of solid surfaces is critically dependent on relative humidity, inoculum volume and organic soiling. FEMS Microbes xtad022.

9. Rosenberg M, Park S, Umerov S, Ivask A. 2025. Experimental evolution of *Escherichia coli* on semi-dry silver, copper, stainless steel, and glass surfaces. Microbiol Spectr 13:e02173-24.

10. Balouiri M, Sadiki M, Ibnsouda SK. 2016. Methods for in vitro evaluating antimicrobial activity: A review. J Pharm Anal 6:71–79.

11. Kadeřábková N, Mahmood AJS, Mavridou DAI. 2024. Antibiotic susceptibility testing using minimum inhibitory concentration (MIC) assays. Npj Antimicrob Resist 2:37.

12. Hamill PG, Stevenson A, McMullan PE, Williams JP, Lewis ADR, S S, Stevenson KE, Farnsworth KD, Khroustalyova G, Takemoto JY, Quinn JP, Rapoport A, Hallsworth JE. 2020. Microbial lag phase can be indicative of, or independent from, cellular stress. Sci Rep 10:5948.

13. Bertrand RL. 2019. Lag Phase Is a Dynamic, Organized, Adaptive, and Evolvable Period That Prepares Bacteria for Cell Division. J Bacteriol 201.

14. Ardré M, Doulcier G, Brenner N, Rainey PB. 2022. A leader cell triggers end of lag phase in populations of Pseudomonas fluorescens. microLife 3:uqac022.

15. Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ. 2017. Antibiotic tolerance facilitates the evolution of resistance. Science 355:826–830.

16. Sulaiman JE, Lam H. 2021. Evolution of Bacterial Tolerance Under Antibiotic Treatment and Its Implications on the Development of Resistance. Front Microbiol 12:617412.

17. Percival SL, Bowler PG, Russell D. 2005. Bacterial resistance to silver in wound care. J Hosp Infect 60:1–7.

18. Evans A, Kavanagh KA. 2021. Evaluation of metal-based antimicrobial compounds for the treatment of bacterial pathogens. J Med Microbiol 70:001363.

Supplementary Notes and Figures for:

Distinct sub-MIC kill kinetics of Cu and Ag in *Escherichia coli*

Merilin Rosenberg, Sigrit Umerov, Carmen Marianne Teär, Angela Ivask

Supplementary Note 1 on Strain selection, growth and exposure conditions.

Strain and inoculum selection

We first observed the effects described in this study for *E. coli* ATCC 8739 (unpublished data), a strain that is routinely used in antimicrobial testing. However, the strain tends to rapidly acquire silver resistance in liquid medium granted by mutations in the cryptic genomic *sil* locus [1–4] that interferes with studying the effects of antimicrobial metals on the wild-type strain. *E. coli* K-12 derivative BW25113 [5] was used for the current study as a better described model strain of *E. coli* lacking the *sil* locus. Exponential phase inoculum was used in the study similarly to standardized MIC and MBC assays and to avoid potential biases caused by inoculating stationary or biofilm cultures with different tolerance profiles into fresh growth-supporting exposure medium. Due to the limited growth phase coverage, the results cannot be generalized to stationary and colony-biofilm cultures in which cases additional experiments would be necessary.

Exposure medium selection

It has been demonstrated before that transferring *E. coli* from pH=7.0 to pH=4.2 medium can substantially decrease viable counts [6] and pH=5.5 also seems to both decrease growth rate and prolong lag phase at population level compared to pH=7.5 [7]. In addition, during exponential growth phase *E. coli* culture itself can further decrease pH of a glucose-supplemented medium by 1.5-2 pH units [8,9]. In the context of antimicrobial metals, the addition of 365 ppm (5.7 mM) or 650 ppm (10 mM) copper to regular unbuffered medium such as LB has been shown to decreases pH of the medium from 7.2 to 5.3 or 4.7 accordingly [10], an acidity range that impairs growth of *E. coli* also without excess copper. Similarly, also in our experience supplementing LB (Lennox salt) with $CuSO_4$ reduces pH while the pH change is expectedly much smaller in the buffered MOPS Minimal Medium (MMM), figure below. Additionally, acid resistance, pH recovery and growth dynamics of *E. coli* subjected to near-lethal acid stress in unbuffered LB also depends on inoculum density with complete growth inhibition at initial pH≤4.0 and MIC assay cell density while increasing pH to above ~4.4 is needed before resuming growth [11]. This further complicates interpretation of results with varying cell counts or potentially partial initial killing by either acidity or copper or both. Additional toxic effect of acidification due to copper in the medium might spark ideas for combined synergistic use of antimicrobial agents. However, the slow-release copper sources such as CuO nanoparticles do not seem to have a similar effect on medium pH.

MMM [12,13] was selected for metal exposures due to its buffering capacity to reduce pH bias while comparing antimicrobial activity of copper and silver. Buffered medium counteracts acidification upon addition of $CuSO_4$ and helps to maintain the pH near the optimal growth range of *E. coli* (pH ~6.5-7.5) at sub-MIC copper concentrations. Metal ion toxicity is heavily affected by exposure medium [14,15]. MMM was also used due to its defined characteristics and lower organic content to control the metal ion speciation effects on bioavailability and thus toxicity to bacteria (*e.g.* formation of insoluble salts and complexing of metal ions by thiol groups). Both silver and copper ions can be complexed by organic contents, especially cysteine residues, of the medium with variable affinity, while only silver forms practically insoluble AgCl in the presence of ~60 mM chloride ions in MMM. However, similar situations would be encountered in clinically relevant conditions. Chloride content of MMM is lower than in LB (Miller 10 g/L NaCl → 171 mM; Lennox 5g /L NaCl → 86 mM), PBS (140 mM) or human blood plasma (~100 mM).

Re-growth conditions

We have previously observed that copper exposure can cause slower colony growth substantially affecting results of agar MIC assays [2] possibly also due to copper-induced pH changes. Therefore, growth and kill kinetics were recorded for 48 h instead of 24 h. Drop-plated colonies from the kill curve experiments were counted after 14-16 h for optimal countable colony size and re-checked after 24, 48 and 72 h. Spread-plates from kill curves were counted after 24 h and checked after 48 and 72 h for possible slower growth from exposures with higher metal concentration. No delayed emergence of colonies was observed. 96-well exposure plates from the growth and kill kinetics experiments were kept at 37°C and 150 rpm for 5 days after the 48-h data collection and visually confirmed that delayed growth (emergence of visible turbidity) was not observed for metal concentrations at which no OD increase had been detected during 48 h.

Supplementary Note 2 on Method Details

Escherichia coli K-12 derivative BW25113 was used for the current study as a well-described model strain of *E. coli*. Lysogeny broth (LB: 5 g/L yeast extract, 10 g/L tryptone, 5 g/L NaCl with optional 15 g/L bacteriological agar) and 37°C incubation with or without 150 rpm orbital shaking was used for all precultures and phosphate-buffered saline

(PBS: 8 g/L NaCl, 0.2 g/L KCl, 1.44 g/L Na_2HPO_4, 0.2 g/L KH_2PO_4; pH 7.1) for washing and serial dilutions of cultures. Metal exposures used MOPS Minimal Medium (MMM)[12,13] supplemented with 0.4% glucose, 0.4% casamino acids, and 20 µg/L tryptophan (~60 mM chloride in 1× MMM as described above).

Second subculture of *E. coli* on LB was used to inoculate overnight liquid LB cultures. The stationary culture was 100-fold diluted into fresh LB and grown to exponential phase (OD_{600}=0.5-0.7), then washed twice with cold PBS (4°C, 4400 g, 10 min). Pellets were resuspended in 2× MMM and adjusted to OD_{600} = 0.002 (~10^6 CFU/mL). Inoculum was added to twofold dilution series of $AgNO_3$ or $CuSO_4$ or mixed 1:1 with deionized water and serially diluted in 1× MMM to test inoculum effects. The final volume of exposures in 96-well suspension culture plates was 150 µl for growth curves and 200 µl for kill curves. Sterile water was pipetted between the wells and plates sealed with parafilm to reduce drying during long incubations.

Growth curves were measured at OD_{600} every 15 min at 37°C with double-orbital mixing (BioTek Synergy H1, Agilent). Blank-corrected data were analyzed with Dashing Growth Curves at https://dashing-growth-curves.ethz.ch/ [16].

For kill curves, exposure plates were incubated at 37°C, 150 rpm in the dark. At each point, three concentration series per metal were serially diluted in PBS and drop-plated on LB; 150 µL of undiluted exposures were also spread-plated on 10 cm LB plates. Survival was expressed as log_{10}-transformed survivor fraction (post-/pre-exposure viable counts).

Statistical analysis was done in GraphPad Prism 10.4.1 using correlation, linear regression and one-way ANOVA with post-hoc testing for multiple comparisons at α = 0.05 across 3–4 biological replicates, as detailed in figure legends.

Supplementary Figure S1. Growth parameters of *E. coli* BW25113 exposed to $CuSO_4$ (orange) or $AgNO_3$ (blue) or inoculums with different viable cell densities (grey). Values are calculated from growth curves on Supplementary Figure S3 and S4. Nominal metal concentration or inoculum fold dilution is marked on X-axis where appropriate. Single data points with mean and SD are shown. Lower case letters denote similarity groups based on one-way ANOVA followed by Tukey post-hoc test for multiple comparisons at α=0.05.

Supplementary Figure S2. Dose-dependent changes in lag phase duration (left), doubling time (middle) and yield (right) in response to sub-MIC copper (orange) or silver (blue) concentration. Statistically significant correlations are marked in bold. Growth parameter values are calculated from growth curves on Supplementary Figure S3. Results from 4 biological replicates are presented.

Supplementary Figure S3. Growth curves of *E. coli* BW25113 in the presence or absence of $CuSO_4$ (left, orange) or $AgNO_3$ (right, blue). Individual curves from 4 experiments underlying the mean curves on Figure 1 are shown. Non-exposed controls (black) on both panels are pooled across all experiments.

Supplementary Figure S4. Effect of inoculum density of *E. coli* BW25113 on growth curves (left) and observed strong negative correlation between the lag phase duration and inoculum density (right). Dotted line denotes linear regression. Results from 4 biological replicates are presented.

References for Supplementary Notes and Figures

1. Randall, C. P., Gupta, A., Jackson, N., Busse, D. & O'Neill, A. J. Silver resistance in Gram-negative bacteria: a dissection of endogenous and exogenous mechanisms. *J. Antimicrob. Chemother*. **70**, 1037–1046 (2015).

2. Rosenberg, M., Park, S., Umerov, S. & Ivask, A. Experimental evolution of *Escherichia coli* on semi-dry silver, copper, stainless steel, and glass surfaces. *Microbiol. Spectr*. **13**, e02173-24 (2025).

3. Woolley, C. A., Sutton, J. M. & Wand, M. E. Mutations in SilS and CusS/OmpC represent different routes to achieve high level silver ion tolerance in Klebsiella pneumoniae. *BMC Microbiol*. **22**, 113 (2022).

4. Blanco Massani, M. *et al*. Chromosomal Sil system contributes to silver resistance in E. coli ATCC 8739. *BioMetals* **31**, 1101–1114 (2018).

5. Baba, T. *et al*. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol*. **2**, 2006.0008 (2006).

6. Xu, Y. *et al*. An acid-tolerance response system protecting exponentially growing Escherichia coli. *Nat. Commun*. **11**, 1496 (2020).

7. Zhang, W. *et al*. Escherichia coli Increases its ATP Concentration in Weakly Acidic Environments Principally through the Glycolytic Pathway. *Genes* **11**, 991 (2020).

8. Sánchez-Clemente, R. *et al*. Study of pH Changes in Media during Bacterial Growth of Several Environmental Strains. in *Environment, Green Technology, and Engineering International Conference* 1297 (MDPI, 2018). doi:10.3390/proceedings2201297.

9. Vivijs, B., Moons, P., Aertsen, A. & Michiels, C. W. Acetoin Synthesis Acquisition Favors Escherichia coli Growth at Low pH. *Appl. Environ. Microbiol.* **80**, 6054–6061 (2014).

10. Gunawan, C., Teoh, W. Y., Marquis, C. P. & Amal, R. Cytotoxic Origin of Copper(II) Oxide Nanoparticles: Comparative Studies with Micron-Sized Particles, Leachate, and Metal Salts. *ACS Nano* **5**, 7214–7225 (2011).

11. Segura Munoz, R. R. & Sourjik, V. Collective dynamics of *Escherichia coli* growth under near-lethal acid stress. *mBio* e01932-25 (2025) doi:10.1128/mbio.01932-25.

12. Neidhardt, F. C., Bloch, P. L. & Smith, D. F. Culture Medium for Enterobacteria. *J. Bacteriol.* **119**, 736–747 (1974).

13. MOPS Minimal Medium.

14. Levard, C. *et al.* Effect of Chloride on the Dissolution Rate of Silver Nanoparticles and Toxicity to *E. coli*. *Environ. Sci. Technol.* **47**, 5738–5745 (2013).

15. Rewak-Soroczynska, J., Dorotkiewicz-Jach, A., Drulis-Kawa, Z. & Wiglusz, R. J. Culture Media Composition Influences the Antibacterial Effect of Silver, Cupric, and Zinc Ions against Pseudomonas aeruginosa. *Biomolecules* **12**, 963 (2022).

16. Reiter, M. A. & Vorholt, J. A. Dashing Growth Curves: a web application for rapid and interactive analysis of microbial growth curves. *BMC Bioinformatics* **25**, 67 (2024).