PlantCAD2: A Long-Context DNA Language Model for Cross-Species Functional Annotation in Angiosperms

Jingjing Zhai[1*+], Aaron Gokaslan[2*], Sheng-Kai Hsu[1], Szu-Ping Chen[3], Zong-Yan Liu[3], Edgar Marroquin[2], Eric Czech[5], Betsy Cannon[5], Ana Berthel[1], M. Cinta Romay[1,3], Matt Pennell[4], Volodymyr Kuleshov[2+], Edward S. Buckler[1,3,6+]

Institute for Genomic Diversity, Cornell University, Ithaca, NY USA 14853

Department of Computer Science, Cornell University, Ithaca, NY, USA 14853

Section of Plant Breeding and Genetics, Cornell University, Ithaca, NY USA 14853

Department of Computational Biology, Cornell University, Ithaca, NY, USA 14853

Open Athena AI Foundation, New York, NY, USA 10001

USDA-ARS; Ithaca, NY, USA 14853

* These authors contributed equally to this work

+ To whom correspondence may be addressed. Email: jz963@cornell.edu, vk379@cornell.edu and ed.buckler@usda.gov

Abstract

Understanding how DNA sequence encodes biological function remains a fundamental challenge in biology. Flowering plants (angiosperms), the dominant terrestrial clade, exhibit maximal biochemical complexity, extraordinary species diversity (over 100,000 species), relatively recent origins (~160 million years), ~200-fold variation in genome size and relative compact coding regions compared with other eukaryotes. These features present both a unique challenge and opportunity for pre-training DNA language models to understand plant-specific evolutionary conservation, regulatory architectures and genomic functions. Here, we introduce PlantCAD2, a long-context, plant-specific DNA language model with single-nucleotide resolution, pre-trained on 65 angiosperm genomes, together with a series of public benchmarks for evaluation. Comprehensive zero-shot testing shows that PlantCAD2 (676 million parameters) efficiently captures evolutionary conservation, surpassing the 7-billion-parameter Evo2 model in 10 of 12 tasks. With parameter-efficient fine-tuning, PlantCAD2 also outperforms the 1-billion-parameter AgroNT across seven cross-species tasks including chromatin accessible region, gene expression and protein translation. Moreover, its 8,192bp context window substantially improves accessible chromatin prediction in large genomes such as maize (AUPRC increasing from 0.587 to 0.711), underscoring the importance of long-range context for modeling distal regulation. Together, these results establish PlantCAD2 as a powerful, efficient, and versatile foundation model for plant genomics, enabling accurate genome annotation across diverse species.

Introduction

Deciphering how DNA sequence encodes molecular functions, phenotypes and fitness remains a fundamental goal in biology. The rapid decline in sequencing costs has enabled large-scale initiatives such as the Darwin Tree of Life project [1], the Earth BioGenome Project [2], the Vertebrate Genomes Project [3], and the 10KP Plant Genome Project [4], which collectively aim to sequence tens of thousands of species across the tree of life, with plants alone contributing over a thousand assembled genomes [5]. While genomic data accumulates exponentially, functional annotations lag far behind, particularly in plants where labeled data exists for only a few model species and crops [6]. This gap highlights the critical need for computational models that can learn from raw sequences alone and transfer knowledge to other plant species.

Recent advances in foundation models pre-trained with self-supervised strategy have opened new possibilities for interpreting genomic sequences at scale [7]. Unlike traditional supervised machine learning approaches, which typically require large amounts of labeled data, foundation models are pre-trained on vast collections of unlabeled sequence data. This is particularly advantageous in biology, especially plant biology, where high-quality labeled datasets are often limited. Foundation models can then be fine-tuned on specific downstream tasks using only a small number of labeled examples. This approach has achieved significant success in protein science, where models such as ESM [8–10], ProtTrans [11], and ProBERT [12] have demonstrated strong performance in predicting protein function [13], structure [14], and variant effect [15].

In contrast, genomic LMs are still rapidly evolving, with recent developments spanning DNA [16–25], RNA [26–28] and transcriptomes [29–31]. Among DNA LMs, early work such as the DNABERT [16] pre-trained BERT [32] model on the human genome showed improved performance in predicting regulatory sequence elements compared to supervised models such as DeepSEA [33], DanQ [34] and Basset [35]. Subsequently, more general-purpose DNA LMs have emerged: Evo (pre-trained on all prokaryotic and phage genomes) [18], Evo2 (pre-trained on genomes across tree of life) [19], and Nucleotide Transformer (pre-trained on 850 genomes excluding plants) [25] have demonstrated success across a wide range of tasks, from regulatory element discovery to evolutionary constraint prediction. Notably, multi-species pretraining has proven particularly important for learning evolutionary conservation [17,20,25,36]. Plant-specific models have also emerged. GPN [20], pre-trained on eight Brassicales genomes using a 25-layer convolutional neural network, demonstrated strong performance in variant effect prediction. While AgroNT [22], pre-trained on 48

genomes and modeled regulatory sequences using longer context windows with a non-

overlapping k-mer encoding method. To further improve plant-specific genome modeling, we

previously developed PlantCaduceus [23] (PlantCAD), a DNA LM pre-trained on 16 divergent

angiosperm genomes. It uses the Caduceus [24] architecture, a Mamba-based [37] design that

efficiently models both DNA strands simultaneously. PlantCAD achieved up to a 7-fold

improvement over the next-best model in cross-species gene annotation tasks and variant effect

prediction tasks.

However, PlantCAD is limited by its context window of 512 base pairs, restricting its ability to

model many biological processes that depend on long-range sequence information [38]. Many

regulatory elements can influence gene expression over tens to hundreds of kilobases and they

are key contributors to phenotypic variation [39–41], yet remain challenging to capture with short-

context models. Therefore, we hypothesize that PlantCAD would struggle to model chromatin

accessibility and gene expression patterns across diverse genomic contexts due to the lack of

comprehension in long-range regulatory interactions. While AgroNT extends the context window

to 6,000 bp, its non-overlapping k-mer tokenization strategy sacrifices single-nucleotide resolution,

making it unsuitable for tasks requiring base-level precision such as variant effect prediction [20,22,23].

In contrast, Evo2 is a general-purpose DNA language model with single-nucleotide resolution, but

its massive size (7-40 billion parameters) limits accessibility and, more importantly, its pre-training

across the entire tree of life makes it less-suited to capture the plant-specific regulatory patterns.

Unlike animal genomes which rely on complex long-range regulation, plants have larger and more

variable genomes with expanded gene families and diverse metabolic repertoires. Flowering

plants (angiosperms) [42], the dominant terrestrial clade [43], exhibit maximal biochemical complexity,

extraordinary species diversity, and wide genome size variation, yet maintain relatively compact

coding regions, making them both a challenge and an ideal testbed for plant-specific DNA

language models.

In this paper, we introduce PlantCAD2, an improved DNA LM tailored to angiosperm genomes.

PlantCAD2 is pre-trained using a masked language modeling objective on 65 curated flowering

plant genomes. PlantCAD2 is built on the efficient Mamba2 architecture [44], which scales linearly

with sequence length instead of quadratically such as transformers [45]. It supports 8,192-bp input

windows and reverse-complement equivariance, allowing the model to capture long-range,

strand-invariant regulatory features. To reduce pretraining bias, we applied sampling strategies

that both down-weight repetitive sequences and emphasize coding and regulatory regions.

Subsequently, we first evaluated PlantCAD2 on 12 comprehensive benchmarks using a zero-shot
strategy, demonstrating its efficiency and capacity to capture evolutionary conservation (**Table 1**).
We then fine-tuned the model on seven functional genomics tasks including chromatin
accessibility, gene expression, and protein abundance to further demonstrate its state-of-the-art
cross-species predictive ability (**Table 2**). Together, these results highlight PlantCAD2's ability to
generalize across species and tasks, and to serve as a versatile foundation model for plant
genome interpretation.

Table 1. Zero-shot evaluation summary compared with the best-performing benchmark models.
For each task, the bold and underscored value indicates the highest score.

Category	Task	Description	Metric	PlantCAD2 vs best benchmark
Cross-species evolutionary conservation (Figure 2)	Conservation within Andropogoneae (Genome-wide)	Predict conserved vs non-conserved sites using alignments within 35 Andropogoneae genomes (n = 19,030 vs 19,030)	AUROC	**0.725** vs 0.691
	Conservation within Poaceae (non-TIS)	Predict conserved vs non-conserved coding sites (excluding TIS) within Poaceae (n=103,368 vs 80,317)		0.713 vs **0.822**
	Conservation within Poaceae (TIS)	Predict conserved vs non-conserved TIS sites (n=26,650 vs 10,012)		**0.670** vs 0.551
Key junction recovery (Figure 3)	Translation initiation site (maize)	Recover masked ATG start codon (n = 39,035)	Accuracy	**0.657** vs 0.447
	Translation termination site (maize)	Recover masked TAG/TAA/TGA stop codon (n = 39,035)		**0.410** vs 0.256
	Splice donor (maize)	Recover masked GT motif (n = 153,869)		**0.910** vs 0.741
	Splice acceptor (maize)	Recover masked AG motif (n = 153,869)		**0.900** vs 0.738
Within-species conservation (Figure S4)	Translation initiation site (maize)	Predict core TIS vs non-core TIS in maize (n = 28,291 vs 8,118)	AUROC	**0.710** vs 0.624
	Translation termination site (maize)	Predict core TTS vs non-core TTS in maize (n = 28,291 vs 8,118)		0.618 vs **0.628**
	Splice donor (maize)	Predict core splice donor vs non-core splice donor in maize (n = 123,183 vs 21,367)		**0.808** vs 0.754
	Splice acceptor (maize)	Predict core splice acceptor vs non-core splice acceptor in maize (n = 123,183 vs 21,367)		**0.836** vs 0.761
Structural variant effect (Figure 4)	Structural variant effect prediction	Predict conserved deletions vs non-conserved deletions (n = 7,662 vs 10,413)	AUPRC	**0.841** vs 0.771

112
113 **Note**: For key junction recovery and within-species conservation, comparable results were observed in
114 tomato.

115   **Table 2.** Fine-tuning evaluation summary compared with the best-performing benchmark
116   models. For each task, the bold and underscored value indicates the highest score.

Category	Task	Dataset (train→test)	Task Type	Metric	PlantCAD2 vs best benchmark
Chromatin accessibility (Figure 5)	Cross-species accessible regions	Arabidopsis→ 10 species	Binary classification	AUPRC (mean across 10)	**0.409** vs 0.340
	Cross-species accessible regions (multi-species)	9 species→ 2 species	Binary classification	AUPRC (mean across 2)	**0.570** vs 0.499
	Cell-type-specific accessible regions	Maize (hold-out chr10)	Multi-label classification	AUPRC (mean across 92 cell types)	**0.662** vs 0.650
Gene Expression (Figure 6)	Cross-species leaf gene on/off prediction	15 Andropogoneae → 26 NAM genomes	Binary classification	AUROC	**0.854** vs 0.819
	Cross-species leaf absolute gene expression	15 Andropogoneae → 26 NAM genomes	Regression	Spearman correlation	**0.633** vs 0.616
Protein Translation (Figure 6)	Cross-species leaf translation on/off prediction	Arabidopsis → Maize	Binary classification	AUROC	**0.692** vs 0.597
	Cross-species leaf absolute translation abundance	Arabidopsis → Maize	Regression	Spearman correlation	**0.321** vs 0.181

117   Note: sample sizes (n) for training and test sets are provided in Supplementary Tables.

Results

PlantCAD2: a long-context DNA language model for angiosperms

PlantCAD2 builds on the original PlantCAD [23] DNA language model, preserving its single-nucleotide tokenization and masked language modeling objective, while introducing four major improvements: architectural efficiency, context length, parameter scale, and phylogenetic breadth (**Figure 1A**). First, PlantCAD2 retains the Caduceus [24] architecture with its bidirectional, reverse-complement-equivariant design, but replaces the original Mamba [37] blocks with Mamba2 blocks [44]. Mamba2 has substantial advantages over Mamba1, leveraging structured state space duality for more efficient parallel training and simplifying recurrence computations to reduce memory usage (see Methods). Compared to traditional transformer architectures [45,46], PlantCAD2 model architecture shows a much slower increase in inference time than modernBERT models under the same input and output dimensions (**Figure S1A**), due to the inherent efficiency of state space models in handling long sequences [37,47]. Exploiting this efficiency, PlantCAD2 takes 8,192 base pair (bp) windows, which is a 16-fold increase over the 512-bp windows used in PlantCAD. Second, to evaluate the effect of model sizes on performance, we trained a series of depth-scaled PlantCAD2 models of 88M, 311M and 694M parameters (**Figure 1A-1B**), which we named PlantCAD2-S, PlantCAD2-M, and PlantCAD2-L respectively. As expected, following pre-training, the largest model (PlantCAD2-L) demonstrated the best masked token prediction accuracy (0.657), followed by PlantCAD2-M (0.641) and PlantCAD2-S (0.598), when evaluated on hold-out test set by randomly masking 15% of nucleotides per sequence (**Figure S1B**). However, the largest model also shows slowest inference speed at ~32,238 nucleotides per second on a NVIDIA H100 80 GB PCIe GPU (**Figure 1C**) and uses ~27G GPU memory with a of 32 (**Table 3**), reflecting the typical trade-off between accuracy and computational efficiency. The smallest PlantCAD2 model, PlantCAD2-S, achieves a 4.5-fold (**Figure 1C**) higher inference throughput while requiring considerably less GPU memory at equivalent batch sizes (**Table 3**), highlighting its practicality for high-throughput analyses. Despite differences in model size, the three models showed high correlation in their per-species prediction accuracies (r > 0.97), suggesting consistent learning patterns across scales (**Figure S1C; Supplemental Table 1**). Third, to assess the effect of input length on pretraining accuracy, we varied the context window size from 512bp to 8,192bp and evaluated performance by masking the central token. All three models showed improved masked token prediction accuracy with longer contexts, underscoring the importance of extended context for modeling kilobase-scale genomic dependencies (**Figure 1D**). Lastly, we

expanded the evolutionary diversity of the training dataset from 16 to 65 angiosperm genomes (**Figure 1E; Supplemental Table 1**), selecting one representative species per genus to maximize phylogenetic breadth. When analyzing pre-training performance across species, we found a weak positive correlation between genome size and masked token accuracy (r = 0.525; **Figure S1D, Supplemental Table 1**). This relationship is likely driven by the fact that larger genomes tend to contain more repetitive sequences [48]. Since the masked language modeling objective can predict repetitive elements more easily than non-repetitive elements even after applying down-sampling and down-weighting (see Methods), the amount of repeats in the test set could inflate accuracy [20,23,36]. Consistent with our expectation, we also detected a positive correlation between the number of repeats in the test set and masked language modeling accuracy (**Figure S1E**), this also highlights the importance of down-sampling and down-weighting repetitive sequences [36] in pre-training DNA language models.

Figure 1. Overview of the PlantCAD2 model. (A) Comparison of PlantCAD1 and PlantCAD2 model configurations. PlantCAD2 introduces a longer context window, upgraded architecture (Mamba2), expanded pre-training species set, and scaled model sizes (small: 88M, medium: 311M, large: 694M parameters), while maintaining single-nucleotide tokenization. **(B)** Schematic of the PlantCAD2 architecture based on Mamba2 with reverse-complement (RC) equivariance, convolutional and state space modules

(SSM), and a masked language modeling objective applied to 8,192 bp input sequences. **(C)** Total throughput (sequences/second) of PlantCAD2 models on NVIDIA H100 80 GB PCIe GPU across batch sizes (1–64). Values on the bar represent mean throughput across batch sizes. **(D)** Effect of context window length on model performance. The y-axis shows the prediction accuracy of three models when masking the single central token in the held-out test set. **(E)** Phylogenetic distribution of the 65 angiosperm genomes across flowering plant orders. Numbers in parentheses indicate the number of species included from each order.

Table 3. Inference throughput and GPU memory usage of PlantCAD2 models on an NVIDIA H100 80 GB PCIe GPU.

Model	Seq Len (bp)	Batch size	Peak memory (GB)	Seq/s	Tokens/s
PlantCAD2-S	8192	1	0.79	19.58	160,422
		4	1.94	20.12	164,812
		8	3.55	19.63	160,829
		16	6.56	19.61	160,653
		32	12.76	19.26	157,767
		64	24.89	19	155,649
PlantCAD2-M	8192	1	2.5	5.46	44,761
		4	3.49	7.02	57,489
		8	5.54	6.88	56,388
		16	9.62	6.88	56,386
		32	17.62	6.76	55,342
		64	33.62	6.79	55,636
PlantCAD2-L	8192	1	4.76	3.98	32,634
		4	5.75	4.01	32,847
		8	8.83	3.94	32,264
		16	14.89	3.92	32,111
		32	26.95	3.87	31,741
		64	51.09	3.89	31,833

PlantCAD2 accurately predicts evolutionary conservation with a zero-shot strategy

Evolutionary conservation, commonly estimated from multiple sequence alignment (MSA), is widely used to identify deleterious mutations that may reduce organismal fitness [49–52]. However, estimating a genome-wide MSA is particularly challenging in plants due to extensive transposable element (TE) insertions and their high turnover rate, which obscure orthologous relationships outside of conserved coding regions [53]. This limitation highlights the need for alignment-free

approaches to assess conservation across diverse plant genomes. Given that PlantCAD2 is pre-
trained on 65 evolutionary distant species, we hypothesize that PlantCAD2 can be used to predict
evolutionary conservation without an MSA. We first evaluated how accurate PlantCAD2 is to
distinguish highly conserved sites versus less conserved sites using a zero-shot strategy. As
illustrated in **Figure 2A,** we used the masked nucleotide/token prediction accuracy from the frozen
model to represent per-base conservation, which means highly conserved bases would receive
higher predicted probabilities for the reference allele, whereas less conserved bases would yield
lower confidence scores. We benchmarked the performance of PlantCAD2 against three
baselines: its predecessor PlantCAD [23], GPN [20] (a plant specific DNA LM trained on Brassicales
genomes), and Evo2 [19], a general-purpose DNA language model pre-trained using a causal
language modeling (CLM) objective, also known as next-token prediction. Unlike masked
language modeling, which enables access to both upstream and downstream context, CLM
imposes a strict left-to-right constraint, meaning Evo2 is inherently unidirectional. Therefore, we
input the entire sequences without masking for Evo2 and use the likelihood of the model to
represent conservation. Notably, Evo2 was trained at a substantially greater scale, with 7 billion
parameters and 9.3 trillion nucleotides, which is over 310 times more training data than used for
PlantCAD2, therefore providing a rigorous benchmark for assessing the efficiency and
representational power of our models. We excluded AgroNT from zero-shot evaluation as its non-
overlapping k-mer tokenization strategy prevents single-nucleotide resolution tasks, and we
previously demonstrated its limited zero-shot capabilities [23].

We assessed this strategy in three independent tasks. First, we performed cross-species
alignments of 34 Andropogoneae genomes [23] to the sorghum reference genome, and identified
highly conserved and less conserved sites based on alignment coverage and identity (see
Methods). PlantCAD2 consistently outperformed PlantCAD in distinguishing highly conserved
from less conserved sites in the sorghum genome, with the largest PlantCAD2 achieving the
highest AUROC (**Figure 2B; Supplemental Table 2**). Notably, PlantCAD2-M achieved slightly
better performance than Evo2 (AUROC 0.708 vs 0.691) despite being ~22-fold smaller (311M vs
7B parameters), while PlantCAD2-L, being ~11-fold smaller (694M parameters), further improved
to 0.73. This demonstrates that our PlantCAD2 models can match or exceed Evo2's performance
with far fewer parameters. Given that PlantCAD2 is pre-trained with a context window of 8192 bp,
we also examined the effect of context length on conservation prediction. AUROC scores
increased with longer sequence contexts, plateauing at 4096 bp for all PlantCAD2 models (**Figure
S2A; Supplemental Table 2**). These findings indicate that evolutionary constraint signals benefit

from broader sequence context and that larger models with extended receptive fields are better
suited to capture these dependencies.
222
In the second task, we used multiple sequence alignments from coding sequences of 325
Poaceae genomes to calculate phyloP scores and define highly conserved sites (phyloP > 5) and
less conserved sites (phyloP < 1.5). While the relationship between selection and phyloP scores
can be nuanced [54], restricting phyloP calculation to coding regions helps mitigate alignment noise
caused by the very high transposable element turnover rate in plant genomes [55], providing a more
reliable benchmark for conservation prediction. Given Evo2's unidirectional nature, we
hypothesized it might struggle with features requiring bidirectional context, particularly translation
initiation sites (TIS), where both upstream regulatory motifs in the 5′ UTR and downstream coding
sequence context critically influence start codon recognition and conservation [56]. To test this
hypothesis, we separately evaluated performance on TIS versus non-TIS positions within coding
sequences. For non-TIS sites, PlantCAD2 models showed lower performance compared to Evo2
(**Figure 2C; Supplemental Table 2**), potentially because Evo2's training data included mature
mRNA sequences while PlantCAD2 was trained exclusively on genomic DNA, giving Evo2 an
advantage in coding sequence conservation tasks. We also note that the original PlantCAD
slightly outperforms all PlantCAD2 models on this specific task (**Figure 2C**). This likely reflects
differences in input–pretraining alignment: PlantCAD was pre-trained on short 512 bp windows
enriched for coding regions, and the CDS-centered inputs used here closely match that training
setting. In contrast, PlantCAD2 was pre-trained on 8 kb genomic windows that systematically
include intronic and intergenic contexts. Consequently, restricting the input to CDS-centered
sequences removes intronic context and deviates from PlantCAD2's pre-training setting,
attenuating the advantages conferred by its long receptive field. Nevertheless, despite this
mismatch, PlantCAD2 models still achieve competitive performance, indicating that their learned
representations remain informative even under constrained input conditions. Interestingly, when
evaluating TIS conservation, we observed a strong bias in Evo2: its AUROC dropped drastically
to 0.534, barely above random. In contrast, PlantCAD2 maintained robust performance (AUROC:
0.632–0.670; **Figure 2D; Supplemental Table 2**). Even the 65M-parameter GPN outperformed
the 7B-parameter Evo2 on this task (AUROC of 0.551), further highlighting Evo2's architectural
limitations for TIS prediction (**Figure 2D; Supplemental Table 2**). This TIS-specific weakness in
Evo2 validates our hypothesis: without access to coding sequence contexts that are more
evolutionary constrained, Evo2 cannot properly assess the conservation patterns at translation
start sites. Consistent with the previous task, we observed only modest improvements from longer

context windows, indicating that evolutionary conservation signals are predominantly local (**Figure S2B-S2C**). These results demonstrate that PlantCAD2 provides more consistent and unbiased conservation predictions across different genomic contexts.

To further assess whether PlantCAD2 generalizes to lineages absent from pre-training. We evaluated its zero-shot conservation predictions in Solanaceae, a major angiosperm family not represented in the 65 pre-training species. This evaluation used SNPs from the potato (*Solanum tuberosum* L.) genome, with GERP scores [57] calculated from a multiple sequence alignment of 95 Solanaceae genomes [58]. Conserved sites were defined as SNPs with GERP scores greater than 1.5, and neutral sites as SNPs with GERP scores lower than 0. As shown in **Figure 2E**, PlantCAD2 models achieved higher AUROC than both Evo2 and GPN, despite multiple Solanum genomes being part of the Evo2 pre-training corpus but absent from PlantCAD2 pre-training. Additionally, to understand functional constraints across the entire genome, we stratified the evaluation into six distinct genomic contexts: CDS, upstream (promoters), downstream, introns, intergenic regions, and transposable elements (TEs) (**Figure S3**). As expected, accuracy was higher in coding versus non-coding regions (**Figure S3**), likely because the stronger functional constraints on CDS make these patterns easier for the model to capture. When comparing models across the five non-coding categories, PlantCAD2 consistently showed the best performance in four categories (upstream, downstream, intergenic, and TEs), with GPN showing only a marginal advantage in introns. Notably, PlantCAD2 consistently surpassed Evo2 across all six genomic contexts. The strong consistent performance across three phylogenetically distant angiosperm lineages indicates that PlantCAD2 captures fundamental conservation principles shared across flowering plants rather than lineage-specific patterns. This enables accurate alignment-free conservation inference even in lineages unseen during pre-training.

Figure 2. PlantCAD2 accurately predicts evolutionary conservation using zero-shot strategy. (A)
Illustration of the zero-shot conservation approach. For each masked token (nucleotide) in the reference
sequence, the frozen PlantCAD2 model outputs the probability of the reference allele; higher probabilities
indicate higher evolutionary constraint. **(B)** AUROC for predicting conserved versus neutral sites in sorghum,
representing conservation within the Andropogoneae tribe. **(C-D)** AUROC for predicting conservation within
Poaceae using sites in coding regions, evaluated separately for non-TIS **(C)** and TIS **(D)**. **(E)** AUROC for

PlantCAD2 accurately predicts within-species conserved transcriptional and translational junction sites with zero-shot strategy

We next quantified how well PlantCAD2 captured the sequence context that defines key transcriptional and translational junctions. Using a similar zero-shot strategy, we designed four tasks to recapitulate motifs (**Figure 3**). Instead of masking one base pair, for each annotated junction, we replaced the annotated motif with consecutive [MASK] tokens. Given the non-canonical motifs do exist in plants [59], we retained all annotated sites without filtering for canonical signals. Consequently, the evaluation included both canonical and non-canonical variants as defined in the reference annotation. For example, translation initiation sites included both canonical ATG and non-canonical start codons such as GTG and TTG. We then extracted a fixed 8,192-bp window centered on the masked motif and presented the entire masked sequence to the model without fine-tuning. A prediction was considered correct if the model's top-1 reconstruction exactly matched the ground truth motif found in the genome.

As above, we benchmarked the performance of PlantCAD2 against PlantCAD, GPN and Evo2. Given what we observed in **Figure 2D** that Evo2 is limited with its poor TIS conservation prediction, we evaluated it using two configurations: (1) forward sequences (Evo2-fwd), where the model uses upstream context to predict the junction, and (2) reverse-complement sequences (Evo2-rc), where the model uses downstream context (reverse complemented) to predict the junction. For GPN and PlantCAD, which are both limited to context windows of 512 bp, we used 512-bp windows centered on the junctions for evaluation.

When evaluated on both maize (*Zea mays*), which was included in pre-training and tomato (*Solanum lycopersicum*), which was not included in pre-training, PlantCAD2 consistently outperformed PlantCAD1 across all junction types (**Figure 3; Supplemental Table 3**). Notably, even the smallest PlantCAD2 model (88M parameters) outperformed the original PlantCAD (311M parameters), demonstrating that architectural improvements and expanded phylogenetic diversity (65 vs. 16 genomes) provide substantial benefits beyond parameter scaling alone. Accuracy increased with model size, following expected scaling law, with the largest model achieving the highest masked-motif prediction accuracy across both species.

As expected, Evo2 showed strong directional effects. For junctions where downstream context is
more informative (TIS and splice acceptor), Evo2-rc performed better, as the reverse-complement
orientation allows the model to 'see' the downstream coding sequences that provide stronger
signals. Conversely, for junctions where upstream context matters more (TTS and splice donor),
Evo2-fwd showed superior performance. The sharp contrast in performance between forward
sequences and reverse complemented sequences reflects a fundamental limitation of causal
language models: their unidirectional nature prevents them from accessing both upstream and
downstream signals simultaneously. In contrast, PlantCAD2's bidirectional and reverse-
complement equivariant design achieved robust performance regardless of sequence orientation,
consistently leveraging both upstream and downstream context for all junction types.

Figure 3. PlantCAD2 accurately predicts transcriptional and translational junction sites using zero-shot masked motif prediction. Left panels show the masking strategy where canonical motifs are replaced with [MASK] tokens and models predict the correct sequence. Right panels show prediction accuracy for each model on maize (left, included in training) and tomato (right, excluded from training). **(A)** Translation initiation sites (ATG masking). **(B)** Translation termination sites (TAG/TGA/TAA masking). **(C)** Splice donor sites (GT masking). **(D)** Splice acceptor sites (AG masking).

While recovering canonical junction motifs demonstrates basic sequence understanding, we next tested whether PlantCAD2 captures deeper evolutionary signals that distinguish core genes

(evolutionarily constrained and present across taxa) from non-core genes (rapidly evolving and

taxa-specific). We extracted each model's log-likelihood of the canonical motif as a conservation

score and evaluated binary classification performance using AUROC (**Figure S4**). For Evo2, we

selected the optimal orientation based on junction type (forward for TTS/donor, reverse-

complement for TIS/acceptor) as determined above. Remarkably, even though the tomato

genome was excluded from PlantCAD2's 65 pre-training genomes, PlantCAD2 consistently

outperformed Evo2—which did include tomato during pre-training. This demonstrates strong

cross-species generalization: PlantCAD2 learned transferable conservation patterns from other

angiosperms that effectively predict functional constraints in unseen species. These results

highlight PlantCAD2's ability to capture fundamental evolutionary principles instead of just

recognizing simple motif recognition.

PlantCAD2 predicts functional structural variants with zero-shot strategy

In addition to single-nucleotide conservation, we investigated whether PlantCAD2 can generalize

to predicting the functional impact of structural variants, such as small deletions, using a zero-

shot approach. To do this, we simulated a set of deletions in the Arabidopsis genome and

computed the Δlog P score, defined as the log-likelihood ratio between the mutated and reference

sequences surrounding the deletion site, averaged across the deletion window (**Figure 4A**). To

assess how well Δlog P reflects the functional deletions, we utilized two complementary

conservation metrics derived from multiple sequence alignments of 63 genomes [60]: phyloP [61],

which measures conservation or acceleration at individual sites, and phastCons [61] which identifies

conserved elements by analyzing patterns across adjacent sites. We classified deletions as either

"highly conserved" or "less conserved" based on the average scores from each metric.

Given more negative Δlog P represents higher conservation, we observed PlantCAD2's negative

Δlog P scores showed strong positive correlation with both phyloP-based and phastCons

constraint scores (**Figure 4B; Figure S5**), with the model assigning higher likelihoods to

mutations in evolutionarily constrained regions. To quantify this relationship, we binarized

deletions into "highly conserved" and "less conserved" categories based on either phyloP (**Figure

4A**) or phastCons scores (**Figure S5A**) and evaluated whether Δlog P could discriminate between

them. PlantCAD2 achieved robust classification performance either using phyloP (**Figure 4C;

Supplemental Table 4**) or phastCons (**Figure S5**), with even the 88M-parameter PlantCAD2-S

outperforming the 7B-parameter Evo2, highlighting the advantage of plant-specific training over

general-purpose models again. Additionally, we also observed classification performance is

saturated with just 20 bp of flanking sequence on each side (**Figure S6**), indicating that local
sequence context sufficiently captures the functional impact of small deletions. This strong
performance is particularly impressive given that PlantCAD2 was never explicitly trained on
structural variants, suggesting it learned generalizable sequence constraint patterns during pre-
training. These results suggest that PlantCAD2's learned representations generalize beyond
single-nucleotide changes, capturing broader sequence dependencies relevant to noncoding
structural variation. This provides a scalable alternative to traditional alignment-based
conservation methods. Notably, this also represents one of the first efforts to use DNA LMs for
estimating indel effects in plant genomes, underscoring the potential of foundation models in
addressing complex variant interpretation challenges.

Figure 4. PlantCAD2 predicts functional impact of structural variants using zero-shot
strategy. (A) ΔlogP calculation approach for deletion variants and phyloP score distribution for
classification. **(B)** Scatter plot showing the positive correlation between PlantCAD2's -ΔlogP

scores and phyloP-based conservation scores. **(C)** AUROC performance distinguishes highly
conserved from less conserved deletions.

Transcription factor binding sites revealed by high-confidence predictions of PlantCAD2

To evaluate the interpretability and regulatory grammar capture of PlantCAD2, we first assessed
whether the model's high-confidence predictions align with experimental verified transcription
factor binding sites (TFBS) using a zero-shot approach [20]. We masked each position within
DAP-Seq-defined DREB2 peak regions [62] and generated importance scores from the model's
output distribution, then clustered these into motifs using TF-MoDISco [63]. DREB2 (AP2/ERF
family) was selected as a representative transcription factor because AP2/ERF TF family is
prevalent and evolutionary conserved across angiosperms [64].
396
The de novo motif discovered by TF-MoDISco with PlantCAD2 predictions showed a highly
significant match to the canonical AP2/ERF profile (https://jaspar.elixir.no/matrix/MA1222.1) in
the JASPAR database (Tomtom p-value = 0.0001; **Figure S7**). In contrast, while the Evo2
model also detected the motif, the match was substantially less significant (Tomtom p-
value=0.005), suggesting that PlantCAD2 captures plant-specific regulatory features with higher
fidelity.
403
We then extended this strategy to performing de novo motif discovery across all 1.5-kb regions
upstream of translation initiation sites in the Arabidopsis genome. This genome-wide analysis
was designed to systematically identify representative cis-regulatory motifs captured by the pre-
trained PlantCAD2 model beyond known DAP-Seq peaks. In total, 104 high-confidence motifs
were identified. Representative examples, highlighted in **Figure S8** include BPC1 (AT2G01930),
involved in ovule identity [65]; TSO1 (AT3G22780), a regulator of floral cell division [66];
SRS7/SPRI2 (AT1G19790), which modulates auxin-dependent reproductive barriers [67]; and
ZML1 (AT3G21175), a GAGA-type factor mediating photoprotection [69]. These results may
suggest these TFBS exhibit profound evolutionary conservation and indicate PlantCAD2 could
be a useful tool to recover both established and potentially novel conserved regulatory
elements. All discovered motifs and their statistics are available on GitHub:
https://plantcad.github.io/TFBS.

Fine-tuning PlantCAD2 accurately predicts cross-species chromatin accessible regions and cell-type-specific accessible regions

We next investigated whether PlantCAD2 learned chromatin states by assessing its performance in predicting genome-wide chromatin accessibility across multiple plant species. We formulated this as a binary classification task, in which the model predicts whether a given genomic region corresponds to an accessible chromatin region, as defined by ATAC-seq (**Methods**). We used recently published ATAC-seq data including 11 diverse plant species, including both dicots and monocots [67]. In this task, positive examples correspond to accessible peaks from ATAC-Seq, while negative examples were from genomic background regions. We used 600-bp genomic windows for all models, as this resolution captures the typical size of ATAC-seq peaks while providing sufficient sequence context for regulatory element prediction. Due to the biological reality of accessible regions comprising only a small fraction of the entire genome, this task is highly imbalanced (**Supplementary Table 5**). For example, less than 1% of regions in the maize genome are labeled as positive.

To effectively leverage the pre-trained foundation model, we used a Low-Rank Adaptation (LoRA) [70] fine-tuning strategy for PlantCAD2, which inserts small trainable rank-decomposition matrices into the feedforward layers while keeping the rest of the model frozen (**Figure 5A**). This approach updates only a small fraction of parameters, enabling efficient task-specific adaptation with minimal risk of overfitting or forgetting the pre-trained knowledge. To assess the contribution of pre-training, we compared this approach to two supervised baselines: (1) a fully supervised version of PlantCAD2-S from scratch, where all model parameters were randomly initialized and updated during training; and (2) a commonly used CNN+LSTM [33] architecture trained from scratch. We also benchmarked against AgroNT [23], another plant-specific DNA LM. We excluded GPN and PlantCAD due to their limited 512-bp context window. For other DNA LMs, we tailored our benchmarking strategy to the specific characteristics of each model. Specifically, we benchmarked against AgroNT using LoRA, as our previous study [68] demonstrated that AgroNT yields suboptimal performance with frozen embeddings but gains significant accuracy through fine-tuning, which is computationally feasible given its 1-billion-parameter size. In contrast, while fine-tuning the 7-billion-parameter Evo2 model was computationally expensive, we sought to strictly evaluate the quality of its learned representations against PlantCAD2. Therefore, we implemented a dual-probing strategy using frozen embeddings extracted from the last hidden state of both models. To ensure a rigorous comparison, we evaluated embeddings derived from

three distinct input configurations: the forward sequence, the reverse-complement sequence, and
the average of both. This comprehensive approach ensures that we captured the maximal
information content available from Evo2, accounting for potential strand biases or invariant
features. We then trained two distinct classifiers: (1) a simple linear layer to assess linear
separability; and (2) XGBoost [71] to capture complex non-linear feature interactions. This ensures
a comprehensive evaluation of the information content encoded by each model without requiring
full parameter updates. All models were trained using Arabidopsis and validated on hold-out
chromosomes within Arabidopsis as well as on 10 additional test species spanning a broad
phylogenetic range.

Given the strong class imbalance of this task, we measured model performance using the area
under the precision–recall curve (AUPRC), which is more informative than AUROC in imbalanced
classification settings. LoRA fine-tuned PlantCAD2 consistently achieved the best performance
in both within-species evaluation and cross-species generalization, outperforming supervised
baselines and AgroNT across all test species (**Figure 5B; Supplementary Table 5**). Furthermore,
in the comparison using fixed (frozen) embeddings against the 7-billion-parameter Evo2,
PlantCAD2 consistently achieved higher AUPRC scores across the 11 species, regardless of
whether a linear or non-linear classifier was used (**Figure S9**). While the averaged embedding
(combining forward and reverse-complement strands) proved to be the most robust configuration
for Evo2, it was still consistently outperformed by all PlantCAD2 models across all test species.
Remarkably, even our smallest model, PlantCAD2-S, surpassed the 7B-parameter Evo2. On
average, PlantCAD2-L yielded an AUPRC improvement of 0.1142 and 0.1735 over Evo2's best
performance using Linear and XGBoost classifiers, respectively (**Figure S9**). This result highlights
the efficiency of PlantCAD2 as an angiosperm DNA foundation model, demonstrating that
domain-specific pre-training contributes more to representation quality than simply scaling up
model parameters.

Across the diverse species tested, we observed a strong negative correlation between genome
size of AUPRC, which reflects the increasing difficulty of distinguishing sparse regulatory
elements in large intergenic regions (**Figure S10**). Despite this challenge, fine-tuned DNA LMs
consistently outperformed supervised models trained from scratch (including CNN+LSTM and
Supervised PlantCAD2-S), indicating that pre-training enables better learning of chromatin states,
particularly when transferring knowledge across species (**Figure 5B**). Specifically, the supervised
models (whether using CNN+LSTM or Supervised PlantCAD2-S) trained on Arabidopsis

generalized reasonably well to closely related dicots, but their performance declined substantially
when applied to evolutionarily distant monocots such as maize and barley. In contrast, fine-tuned
PlantCAD2 retained strong predictive accuracy across both dicots and monocots, demonstrating
its ability to capture regulatory features conserved across deep evolutionary divergence. These
results underscore the power of combining self-supervised pre-training with parameter-efficient
fine-tuning for plant regulatory genomics.

Figure 5. PlantCAD2 predicts chromatin accessibility across species and cell types. (A)
LoRA fine-tuning approach for binary accessibility prediction using ATAC-seq peaks versus
genomic background. **(B)** Cross-species AUPRC performance when trained on Arabidopsis,
showing superior generalization of PlantCAD2 models compared to supervised baselines across
evolutionary distances. **(C)** Multi-species training performance on held-out barley and maize. **(D)**
Effect of context window length on accessibility prediction accuracy for PlantCAD2-S. **(E)** Multi-
label classification approach for cell-type-specific accessibility prediction. **(F)** Performance
comparison across models for 92 cell types . **(G)** Context window effects on cell-type-specific
prediction accuracy for PlantCAD2-S.

While the foundation model fine-tuned on Arabidopsis already demonstrated clear advantages in
cross-species prediction, we further fine-tuned a multi-species version of PlantCAD2 using
accessible chromatin regions from multiple plant genomes to enhance its robustness across
diverse lineages. With maize and barley held out as test species, this multi-species model
achieved impressive AUPRC scores of 0.691 for maize and 0.449 for barley (**Figure 5C;**
Supplemental Table 6). To investigate whether extended sequence context could further
improve prediction accuracy, we maintained the same 600-bp labels but varied the input window
size by including different amounts of flanking sequence. For computational efficiency, we
conducted this analysis using PlantCAD2-S. Performance consistently improved with longer
context windows, with AUPRC increasing from 0.587 to 0.711 for maize and from 0.398 to 0.479
for barley when extending from 600 bp to 4,600 bp (**Figure 5D**). This substantial improvement
suggests that distal regulatory elements and broader chromatin context beyond the immediate
peak boundaries contribute to accessibility prediction, highlighting the advantage of PlantCAD2's
long-context architecture. These fine-tuned models serve as a robust predictor of chromatin
accessibility across flowering plants and are publicly available to the community as a ready-to-
use resource for regulatory annotation in non-model species.

To further assess whether PlantCAD2 can resolve cell-type–specific regulatory landscapes, we
tested its ability to predict accessible chromatin regions identified through single-cell ATAC-seq
(scATAC-seq) in maize [72]. In contrast to the binary classification task used for genome-wide
accessibility, this task was framed as a multi-label classification problem, where each genomic
window could be accessible in one or more cell types (**Figure 5E**). We curated high-confidence
cell-type–specific peaks across major maize tissues from published scATAC-seq datasets
(**Methods**), using them as labels for multi-label fine-tuning and evaluation.

We applied the same LoRA fine-tuning strategy used in prior experiments, adapting PlantCAD2
to predict cell-type–specific accessibility using only a small number of trainable parameters. As in

previous sections, we compared performance against two supervised baselines: a CNN+LSTM
model trained from scratch and a fully supervised version of PlantCAD2. All models were trained
on all maize cell types and evaluated on held-out chromosomes, with performance measured
using micro-averaged precision-recall curves across cell types. Despite the complexity and
subtlety of cell-type–specific regulatory signatures, LoRA fine-tuned PlantCAD2 can still achieve
very high accuracy and outperformed other baselines (**Figure 5F; Supplemental Table 7**).
Similar to our genome-wide accessibility results, extending the input context window beyond the
core 600-bp peak region further improved cell-type specificity, with AUPRC increasing from 0.665
to 0.707 when using 4,600-bp windows (**Figure 5G**). This suggests that cell-type–specific
regulatory programs are influenced by broader chromatin context and distal regulatory
interactions. The model captured both shared and lineage-specific accessibility patterns,
demonstrating that pre-trained DNA representations can be effectively adapted to fine-grained
regulatory annotations. These results suggest that PlantCAD2 is not only effective at modeling
general chromatin accessibility across species but is also capable of distinguishing cell-type–
specific regulatory programs within a single genome.

Fine-tuning PlantCAD2 predict cross-species gene expression and protein abundance

To evaluate PlantCAD2's ability to capture gene regulatory signals, we fine-tuned PlantCAD2
models using LoRA for two complementary tasks: gene expression and leaf translation (**Figure
6A, 6D**). Each task involved both classification (on/off status) and regression (absolute
expression/translation level) objectives. For gene expression, we used promoter and terminator
sequences (1024 bp each) as input; for translation, we used 500 bp upstream sequences.
Following the same strategy as the accessible chromatin prediction task, we fine-tuned
PlantCAD2 with LoRA and compared its performance with supervised PlantCAD2-S, CNN+LSTM,
and AgroNT.
549
For cross-species gene expression modeling, we fine-tuned PlantCAD2 on a diverse panel of 15
plant species and evaluated predictions in 26 maize Nested Association Mapping (NAM)
genomes [73] (Methods). We used three tasks to evaluate different models' performance to predict
gene expression: i) a binary classification task to predict if a gene is expressed or not expressed
in leaf tissue; ii) a regression task to predict the absolute expression in leaf tissue; iii) a regression
task to predict the maximum expression across tissues which represents a gene's maximal

transcriptional potential and avoids dilution from inactive tissues (**Figure 6A**). Across these tasks,

PlantCAD2 consistently outperformed established baselines such as AgroNT and supervised

CNN+LSTM. When testing in 26 NAM genomes, even the smallest PlantCAD2 (PlantCAD2-S,

88M parameters) outperformed the performance of the much larger AgroNT model (1B

parameters) (**Figure 6B-6D; Supplemental Table 8**), demonstrating the efficiency of our

foundation model framework. The largest model, PlantCAD2-L, achieved the best AUROC for

binary leaf expression and the highest Spearman correlation for absolute expression prediction.

To also rigorously assess predictive fit for regression tasks (ii and iii), we also reported Pearson

correlation, R^2, MAE (mean absolute error) and MSE (mean squared error) (**Supplemental Table

8**), observing consistent model performance rankings across these metrics. To further validate

the model's generalization capabilities beyond maize, we extended our evaluation to rice and

tomato, where PlantCAD2 maintained the best cross-species performance (**Figure S11**).

Given that regulatory information also lies outside the proximal promoter [70–72], we evaluated the

effect of varying input window sizes on gene expression prediction. Increasing the window from

1 kb to 4 kb both upstream of the transcription start site and downstream of the transcription stop

site resulted in measurable improvements, raising the AUROC from 0.8221 to 0.8455 for binary

leaf expression task and Spearman correlation from 0.6296 to 0.6455 for absolute expression

prediction task on the NAM test set (**Figure S12**). To confirm that these improvements represent

a true biological signal rather than stochastic variance, we performed repeated evaluations with

different random seeds. The results exhibited high stability with minimal standard deviation

(average SD = 0.0025 for binary leaf expression task and SD = 0.0032 for absolute expression

prediction task; **Supplemental Table 9**; **Figure S12**), confirming that the performance gain from

extended context—though numerically modest—is statistically significant and consistently

reproducible. These improvements highlight the role of distal enhancers and long-range motifs in

shaping expression. However, previous studies in both humans and plants have shown that

current deep learning models lack the resolution to capture allele-specific effects [73–75]. We

therefore tested whether fine-tuning a foundation model could mitigate this limitation by evaluating

per-orthogroup correlations within the NAM population, a comparison sensitive to allelic

differences. Consistent with prior findings [73–75], only marginal improvements were observed for

leaf absolute expression prediction, with the median Spearman correlation increasing from 0.112

(supervised CNN+LSTM) to 0.140 (PlantCAD2-S) (**Figure S13**). These small gains likely reflect

both biological and data-related limitations of allele-specific modeling. Cis-regulatory elements

such as transcription factor binding sites often exhibit high turnover rate [76], context-dependent,

and effects driven by variants that are not explicitly represented in the training objective. In
addition, robust allele-specific evaluation requires extensive data curation—such as accurate
partitioning of cis and trans effects [77,78] and access to large-scale hybrid datasets with phased
haplotypes or experimentally validated cis-regulatory variants. Such resources remain limited,
and assembling them requires substantial experimental and computational curation, making it
challenging for sequence-to-function models to learn and generalize allele-specific regulatory
signatures. Together, these results suggest that achieving robust allele-specific expression
prediction will require future models to incorporate explicit cis-variant modeling (e.g.,
VariantFormer [79]) or fine-tuning on curated regulatory variant datasets [77,80].

Translation prediction was based on ribosome profiling (ribo-seq), a sequencing-based approach
that estimates translation activity by mapping ribosome-protected mRNA fragments. Given high-
quality ribo-seq datasets are scarce in plants, we restricted training to Arabidopsis and tested
both within-species performance and cross-species transfer to maize. Interestingly, although
supervised PlantCAD2-S (88M) is much larger than the CNN-LSTM (~1.7M), the latter performed
better, suggesting that large supervised models are prone to overfitting when trained on limited
data (**Figure 6G; Supplemental Table 10**). By contrast, fine-tuned PlantCAD2 with parameter-
efficient LoRA maintained robust performance without signs of overfitting. However, cross-
species regression of absolute translation levels was less effective (**Figure 6G**), likely due to noise
inherent in current ribo-seq–based quantitative estimates. We anticipate that as more high-quality
cross-species ribo-seq datasets become available, fine-tuning will demonstrate significantly
improved robustness for quantitative prediction. Despite this limitation, we were encouraged by
the strong transfer observed for binary classification of leaf translation. We next tested whether
gene expression could similarly be transferred from Arabidopsis to maize. Using a separate
Arabidopsis gene expression dataset [81] with 1,024 bp upstream and downstream sequences, we
found that direct transfer performed poorly (AUROC: 0.786 in Arabidopsis vs. 0.631 in maize)
compared to protein abundance prediction (0.790 in Arabidopsis vs. 0.692 in maize) (**Figure S14**).
This contrast may highlight a fundamental difference between regulatory layers: translational
control appears to be more evolutionarily conserved than transcriptional regulation. Consistent
with evidence that protein abundance is under stronger selective constraint than transcript levels
[63], these results explain why translation prediction transfers effectively across species, whereas
accurate gene expression prediction requires phylogenetically diverse training data.

Figure 6. PlantCAD2 predicts gene expression and translation across species. (A) Gene expression prediction pipeline using promoter and terminator sequences (1024 bp each) for binary classification and regression tasks. **(B-D)** Cross-species gene expression performance on maize NAM population for binary on/off prediction (B), absolute expression levels (C) and maximum gene expression across tissues. **(D)** Translation prediction pipeline using 500 bp upstream sequences. **(E-G)** Translation prediction performance trained on Arabidopsis and tested cross-species on maize for binary on/off prediction (**F**) and absolute translation levels (**G**).

Discussions

In this work, we present PlantCAD2, a long-context window DNA language model that substantially advances the sequence-to-function modeling in plant genomics. Building on the

foundation laid by PlantCAD, PlantCAD2 features a model architecture that is three times larger, a 16-fold longer context window (8,192bp vs. 512bp), and a pre-training dataset that is evenly distributed across angiosperm orders to better capture phylogenetic diversity. Through comprehensive zero-shot and fine-tuned evaluations, we demonstrate that PlantCAD2 not only exhibits strong cross-species generalization, but also achieves superior performance across a wide range of sequence-to-function tasks, including evolutionary conservation prediction, functional important junction sites prediction involved in both transcription and translation, variant (including indels) effect estimation, cis-regulatory activity, gene expression, and protein translation.

A key design consideration for PlantCAD2 is the careful curation of 65 angiosperm species to maximize phylogenetic diversity while minimizing data bias. Rather than using all publicly available plant genomes, we selected a single representative genome per genus. This balanced strategy avoids two major issues. First, pre-training on all available genomes would dramatically increase computational cost. As shown by scaling laws in large language models, increasing dataset size alone while keeping model size fixed leads to diminishing returns unless model capacity and training compute are expanded proportionally [82]. In practice, simply adding more genomes would not necessarily improve performance unless the model were enlarged substantially, which would reduce its efficiency and complicate deployment. Second, public genome resources are heavily skewed toward a few lineages such as grasses and Brassicaceae [83]. Such skewed distributions have been widely documented across NLP [84,85] and protein language models [86]. These studies consistently show that large models tend to learn disproportionately from overrepresented groups in the training data, resulting in biased representations and reduced generalization to underrepresented categories. This highlights the importance of deliberate data curation to ensure balanced phylogenetic coverage and to prevent the model from overfitting to a few dominant plant lineages.

Furthermore, we emphasize that phylogenetic diversity is often more important than the absolute number of species used for pre-training [25]. This has been observed in models such as NTv2 [38], which achieve strong performance with a few hundred phylogenetically diverse species, outperforming models trained on much larger but with 1,000 human genomes. This distinction is particularly critical when comparing PlantCAD2 to models trained on broader evolutionary scales, such as Evo2. While Evo2 incorporates genomes from across all domains of life (representing >100k genomes), this extreme heterogeneity can dilute lineage-specific regulatory

information. By strictly focusing on angiosperms—which diverged around 160 million years ago

and share a coherent regulatory grammar—PlantCAD2 captures generalizable constraints

specific to flowering plants without the "noise" of unrelated genomic architectures (e.g.,

prokaryotic or mammalian regulatory logic). This targeted approach allows PlantCAD2 to achieve

superior performance on plant-specific tasks while maintaining computational efficiency.

The choice of an 8,192 bp context window reflects a deliberate balance between biological

resolution and computational feasibility. Although distal regulatory interactions in plants can

extend over tens of kilobases [84], our survey of the 65 training genomes shows that the median

gene length is 2,422 bp (**Figure S15**), therefore, an 8 kb window is sufficient to encompass most

gene bodies together with their core regulatory regions. Regions far beyond this range are

frequently dominated by transposable elements rather than functional cis-regulatory modules [87],

reducing their contribution to prediction accuracy. From a technical perspective, even with efficient

state-space architectures such as Mamba, both computation time and GPU memory increase

approximately linearly with context length [85,86], making substantially longer windows (e.g., 16–32

kb) impractical for many users and typical hardware settings. Importantly, PlantCAD2 enables

flexible chunking strategies that allow the model to integrate distal information across adjacent

windows when needed. Together, these considerations make 8 kb a pragmatic choice that

maximizes biological utility while maintaining accessibility and efficiency.

PlantCAD2 represents a significant step toward a foundational model for plant genomics. Rather

than building task-specific models for each application, it enables unified modeling of sequence-

to-function relationships that can be efficiently adapted across cell types, tissues, and species.

Fine-tuning PlantCAD2 enables accurate de novo genome annotation [88], further expanding its

utility for genome interpretation. This paradigm shift opens new opportunities to integrate deep

learning into practical breeding applications. For example, PlantCAD2 could assist in prioritizing

causal variants in GWAS studies, interpreting SVs in noncoding regions, or guiding sequence

design for synthetic promoters with desired expression patterns. Its ability to transfer knowledge

across evolutionarily distant species further enhances its utility for crop improvement, particularly

in non-model organisms where high-quality training data are limited but genomic sequences are

available.

Despite its advances, PlantCAD2 also presents new challenges. First, its large model size may

limit deployment in GPU-constrained environments. To directly assess the lower bound of model

size, we additionally trained and evaluated a substantially smaller PlantCAD2 model (~20M parameters) using the same pre-training data and zero-shot evaluation protocols. In contrast to the PlantCAD2-S, this lightweight model showed a pronounced drop in performance across most tasks (**Supplementary Table 11**). For example, in predicting evolutionary conservation within Andropogoneae, the AUROC decreased from 0.725 (PlantCAD2-L) to 0.593 (20M model), indicating a clear capacity threshold below which biologically meaningful sequence representations cannot be reliably learned. These results suggest that the smallest released model (88M parameters) already represents a near-minimal configuration that balances predictive accuracy and computational efficiency, and that further reductions in parameter count will likely require model compression or distillation strategies rather than naive scaling. Second, there are still technical barriers that may limit accessibility to wet-lab biologists or breeders. Building intuitive interfaces, pretrained APIs, and end-to-end pipelines will be crucial to broaden the use of PlantCAD2 in broader plant science communities, and these are obvious areas for future development. Third, while the 8,192-bp context window allows PlantCAD2 to model distal regulatory elements, further extending this capability would be valuable for capturing long-range interactions such as enhancer–promoter loops. For example, in maize, the teosinte branched 1 (*tb1*) enhancer [89,90] and Vegetative to generative transition 1 (*Vgt1*) [91] are located approximately 70kb and 60kb upstream of their target genes, respectively. However, capturing such interactions will likely require novel tokenization or compression strategies that can represent long, repetitive sequences without sacrificing resolution.

Looking forward, future directions include combining PlantCAD2 with multi-modal data such as DNA methylation and chromatin states will provide trans-factors to the genome. In addition, diffusion-based sequence generation models could also be promising coupled with synthetic biology. Ultimately, we envision PlantCAD2 and its successors as key building blocks for a sequence-to-function foundation model capable of enabling predictive genomics and rational genome design in diverse plant species.

Methods

Preparing pre-training genomes

A total of 65 genomes were selected for pre-training from the Phytozome database. To ensure taxonomic relevance and minimize redundancy, we applied a series of manual filtering steps. First, non-angiosperm species were excluded. For each remaining species, we retained only the most

recent genome assembly version. In cases where two haplotypes were available for a species,

we selected the haplotype with the higher N50 value; if N50 values were comparable, we retained

the assembly with fewer scaffolds to prioritize less fragmented genomes. Taxonomic information,

including order, family, and genus, was appended to each genome to facilitate downstream

analyses, and their relationships were visualized using a published time-callibrated phylogeny [92].

For each selected genome, we extracted genomic sequences centered on each annotated gene,

extending 5 kilobases (kb) upstream and 5 kb downstream from the gene center. These ±5 kb

gene-centered regions were then segmented into overlapping windows of 8,192 bp with a step

size of 4,096 bp, ensuring comprehensive coverage of regulatory and genic features while

maintaining continuity across sequence boundaries. These windows served as input sequences

for model pre-training.

PlantCAD2 model architecture and pre-training

PlantCAD2 builds upon the Caduceus architecture [23] used in PlantCAD [37], retaining its key design

principles while incorporating architectural improvements. Like PlantCAD1, PlantCAD2 maintains

three core features: (1) bidirectional sequence processing, where sequences are processed both

forward and reverse with outputs summed together; (2) reverse-complement (RC) equivariance,

ensuring the model commutes with RC operations; and (3) parameter-efficient bidirectional

implementation through shared linear projections between forward and reverse passes.

The primary architectural improvement in PlantCAD2 is the replacement of Mamba1 blocks [44]

with Mamba2 blocks [89]. Mamba2 introduces a structured state space duality that recasts the

selective state space computation into an equivalent convolutional form using structured matrices,

improving parallelism and hardware efficiency. This dual representation enables significantly

faster training (up to 2–4× in some scenarios) while retaining the input-dependent selection

mechanism that allows the model to dynamically modulate state updates based on sequence

content. These advances allow PlantCAD2 to efficiently handle 8,192 bp sequences with linear

computational complexity.

For the pre-training of PlantCAD2, each model was trained for 240,000 steps using a Decoupled

AdamW optimizer [93] with the global batch size of 2,048. The learning rate is 2E-4 with a cosine

decay scheduler, and 6% of the training duration was dedicated to warm up. The learning rate

decayed to 4E-6 by the end of training. The default BERT [20,23] masking recipe was used with a

masking probability of 15%. For each masked token: i) there is an 80% probability it will be

replaced by a special token ([MASK]), ii) a 10% probability it will be replaced by a random token,

and iii) a 10% probability it will remain unchanged. Unless otherwise specified, all models were

trained using a sequence length of 8192 base pairs. A weight decay of 1E-5 was applied

throughout the training process. Pre-training of the 676M-parameter PlantCAD2-L model required

approximately 28 days on 64 NVIDIA H100 GPUs, while the 88M-parameter model completed

training in approximately 3 days under the same hardware configuration.

Repeat annotation and loss re-weighting

To mitigate the influence of highly repetitive sequences during pre-training, we explicitly down-

weighted the contribution of repetitive regions to the masked language modeling loss. This loss has

been found in other DNA models [90] to improve performance on downstream tasks and better calibrate

likelihoods between repetitive and nonrepetitive DNA. We employed a de-novo pipeline to annotate

highly repetitive sequences [94] (https://github.com/baoxingsong/dCNS), then repetitive sequences are

down-weighted during pre-training as demonstrated important in previous studies [91]. Then the loss is

calculated as:

$$\mathcal{L} = \frac{\sum_{i=1}^{N} w_i \cdot \ell(x_i, y_i)}{\sum_{j=1}^{N} w_j},$$

Where w_i represents the weight of i-th nucleotide,

$$w_i = \begin{cases} 0.1 \ if \ position \ i \ is \ in \ repetitive \ region \\ 1.0 \ \ otherwise \end{cases}$$

Evolutionary constraint prediction using the zero-shot strategy

To evaluate the extent to which PlantCAD2 captures evolutionary conservation signals, a zero-

shot strategy was applied to predict constrained genomic regions. Two independent tasks were

used. The first task focused on *Sorghum bicolor*, using conservation estimates from the

Andropogoneae tribe, a large clade of approximately 1,200 grass species that descended from a

common ancestor approximately 18 million years ago [95]. To generate conservation labels, 34

high-quality genomes [96] were aligned to the Sorghum bicolor reference genome using

AnchorWave [97]. Per-base conservation was quantified using alignment identity scores across all

species. Sites with high-quality coverage (i.e., aligned in at least 34 out of 35 species) were

retained for analysis. Among these, positions with an identity score ≥34 were labeled as

conserved, while those with identity scores <15 were labeled as neutral. Sites with intermediate

identity scores or insufficient coverage were excluded from evaluation to ensure high-confidence

labels [94].

The second task utilized conservation scores derived from multiple sequence alignments (MSAs)

of orthologous coding sequences from 325 Poaceae genomes, a high-quality subset of the

recently published set of 727 genomes. Using *Pharus latifolius* as an outgroup species, gap

columns were removed prior to conservation estimates. Using PHAST [98], PhyloP scores were

calculated per site based on a neutral model derived from fourfold degenerate sites and "LRT"

methods with the mode "CONACC". Sites with phyloP scores above 5 were classified as

conserved, while those below 1.5 were considered neutral. Sites with intermediate scores were

excluded to maintain label clarity. For TIS sites, we retained all 36,668 (26,653 conserved vs

10,015 less conserved) sites given their biological importance. For non-TIS sites, we

downsampled to 183,687 sites (103,369 conserved versus 80,318 neutral) for computational

efficiency while maintaining the conserved-to-neutral ratio.

For both tasks, the evaluated site was centered (i.e., 4096^{th} of 8,192bp) within an input sequence,

and the reference base at that position was masked. The model's predicted likelihood of the

reference allele was extracted and used as the zero-shot conservation score. Higher likelihoods

were hypothesized to reflect stronger conservation. Model performance was assessed using

AUROC, comparing scores between conserved and neutral sites.

817   Core and non-core gene classification using the zero-shot strategy

To assess the ability of PlantCAD2 to distinguish between core and non-core genes in a population, a

zero-shot strategy was applied to classify within species conservation of genes in maize and tomato.

For maize, the pangene table derived from 26 Nested Association Mapping genomes [99] was used.

Core genes were defined as those present in all 26 NAM genomes, whereas non-core genes included

both dispensable genes (present in 2-23 genomes) and private genes (present in only one genome).

For genes with multiple transcripts, the canonical transcript specified in the annotation was used. For

tomato, the pan-genome dataset assembled from 586 high-quality genomes [100] was used. Genes

present in all 586 accessions were defined as core genes, and non-core genes consist of dispensable

(present in 6-580 accessions) and private (present in less than 5 accessions). The longest transcript

was selected to represent each gene across all analyses.

To quantify the model's prediction at each functional junction, a masked motif accuracy score was
defined. For example, to evaluate translation initiation sites, the canonical ATG start codon was
masked, and the model's predicted likelihoods for the three masked nucleotides were extracted and
averaged. A similar approach was applied to other junction types, including translation termination
sites (TAA, TAG, TGA), splice donor sites (GT), and splice acceptor sites (AG), by masking the
corresponding motifs and calculating average token likelihoods.
835
836 To evaluate performance, core genes were treated as positive and non-core genes as negative, and
837 AUROC was calculated based on the masked motif accuracy scores for each gene.

Accessible chromatin region prediction

839 To evaluate the capability of PlantCAD2 to capture regulatory sequence features, fine-tuning
840 experiments were performed using Low-Rank Adaptation (LoRA) [70] on two accessible chromatin
841 prediction tasks: (1) cross-species accessible regions (ACRs) prediction, and (2) cell-type-specific
842 ACR prediction.
843
For the cross-species task, ATAC-seq peak regions from 12 plant species [67] were downloaded
from NCBI (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE128434). We followed the
data processing pipeline as described by Wrightsman et al [101]. For each species, peak regions
were processed by extracting the midpoint of each peak and symmetrically extending it by half
the target input window size (300bp, 600bp and 1,000bp) in both directions to generate positive
observations. To reflect the real-world scenario in which most of the genome is inaccessible, the
rest of the genome was used as negative examples, ensuring no overlap with known peaks.
851
852 For the cell-type-specific task, we used the single-cell ATAC-Seq [72] and used a similar
853 preprocessing pipeline, but each genomic region could be associated with accessibility across 92
854 cell types. As such, the task was framed as a multi-label classification problem, where each region
855 was assigned a binary accessibility label for each of the 92 cell types based on its overlap with
856 experimentally identified peaks.

Gene expression prediction in leaf

To evaluate the models' ability to predict gene expression, we designed two tasks: (1) leaf
absolute expression and (2) leaf on/off expression classification. The training dataset was derived
from 15 Andropogoneae species [73]. For validation, we held out two species closest to *Zea mays*—

Tripsacum zopilotense and *Zea diploperennis*—both members of the Tripsacinae subtribe, which

diverged from maize approximately 0.6 to 4 million years ago. This setup enabled evaluation of

the models' cross-species generalization to closely related taxa. For the leaf absolute expression

task, the $\log_{10}$TPM values were used as regression targets during fine-tuning. For the on/off

expression task, genes with TPM > 1 were labeled as expressed (positive), and those with TPM

≤ 1 were considered non-expressed (negative). This setup enabled evaluation of the models'

ability to generalize expression predictions across closely related species within the clade.

Leaf protein abundance prediction task

To evaluate the models' ability to predict protein abundance, we designed two tasks analogous

to the gene expression analysis: absolute abundance and on/off classification. Ribo-Seq data

were obtained from Arabidopsis [102] and *Z. mays* [103]. Raw reads were downloaded from NCBI, and

Trimmomatic [104] was used to trim adapters and filter low-quality reads. Cleaned reads were first

aligned to rRNA reference sequences using Bowtie [105] to remove contaminating rRNA. The

remaining reads were then mapped to the reference genomes of Arabidopsis and maize using

STAR [106]. Gene-level translation abundance was quantified using StringTie [107] based on uniquely

mapped reads. We then designed two tasks analogous to the expression prediction setup: (1)

absolute protein abundance, where the log10-transformed Ribo-Seq expression values were

used as regression targets, and (2) on/off classification, where genes with TPM > 1 were

considered expressed (positive) and those with TPM ≤ 1 were labeled as non-expressed

(negative).

Fine-tuning PlantCAD2

To adapt the pre-trained PlantCAD2 model to downstream tasks, we employed Low-Rank

Adaptation (LoRA) [70], a parameter-efficient fine-tuning strategy that inserts trainable low-rank

matrices into the attention layers of the transformer. This approach enables effective adaptation

while keeping the vast majority of model parameters frozen. Fine-tuning was performed using the

PEFT library [108] with LoRA rank = 8, α = 32, and dropout = 0.1, targeting the "x_proj", "in_proj", and

"out_proj" modules. Models were trained using the Hugging Face Trainer with a learning rate of

1e−4, a global batch size of 128, and one training epoch. BF16 precision and linear learning rate

scheduling with 50 warm-up steps were used. Over 98% of the model parameters remained

frozen, enabling efficient and scalable fine-tuning across tasks. All tasks were fine-tuned for a

single epoch without hyperparameter tuning to ensure stability and consistency across

experiments. Fine-tuning objectives for all models were a binary cross entropy loss for

classification tasks and a mean squared error loss for regression tasks.

Fine-tuning AgroNT
895

To directly compare the performance of fine-tuned PlantCAD2 with AgroNT [45], we applied the

same parameter-efficient fine-tuning strategy using LoRA. All LoRA hyperparameters were kept

consistent with those used for PlantCAD2, including rank = 8, α = 32, and dropout = 0.1. For

AgroNT, LoRA adapters were inserted into the "query" and "key" projection layers of the

transformer [34], reflecting its architecture. This setup ensured a fair comparison between models

under matched fine-tuning conditions.

Supervised CNN + LSTM baseline
903

904 To benchmark against traditional supervised models, we implemented a CNN+LSTM architecture

905 based on DanQ [19], a widely used hybrid model for DNA sequence classification. For each task,

906 the model was trained from scratch using one-hot encoded sequences. We used the Adam

907 optimizer with a learning rate of 0.01, a batch size of 2,048, and trained for up to 200 epochs, with

908 early stopping after 20 epochs without validation improvement.

Supervised PlantCAD2 baseline
910

911 To assess the contribution of pretraining, we trained a small PlantCAD2 model from scratch for

912 each downstream task. This supervised baseline used the same architecture and

913 hyperparameters as the fine-tuned version but was initialized without pretrained weights—by

914 loading only the Hugging Face model configuration.

Zero-shot evaluation of PlantCAD2, PlantCAD and GPN models
916

917 All three models were pre-trained with masked language modeling. For PlantCAD, we used the

918 largest available model "kuleshov-group/PlantCaduceus_l32" for evaluation. For GPN, we used

"songlab/gpn-brassicales". Due to the 512 bp context window limitation of both PlantCAD and GPN, we cropped input sequences to 512 bp centered on the target position. All other evaluation configurations remained identical to those used for PlantCAD2.

Zero-shot evaluation of Evo2 model

All zero-shot tasks were also benchmarked using the Evo2 [19] model ("evo2_7b") for comparison. Since Evo2 is autoregressive (predicting the next token rather than masked tokens), masked token accuracy could not be directly computed. Therefore, for the evolutionary constraint task, we fed the full input sequence into the model and extracted the likelihood of the reference allele at the target site as the conservation score. To ensure a fair comparison, we used an 8,192 bp context window for Evo2, matching the input length used for PlantCAD2 evaluations. The same approach was applied for benchmarking structural variants.

For the masked motif accuracy task, we evaluated Evo2 using two configurations to compensate for its unidirectional architecture: (1) forward sequences (Evo2-fwd), where the model uses upstream context to predict the junction—for example, for TIS prediction, we used the 4,094 bp upstream of the TIS as a prompt for Evo2 to generate the next three tokens; and (2) reverse-complement sequences (Evo2-rc), where the model uses downstream context (reverse complemented) to predict the junction in the opposite direction.

Code availability

All pre-trained models, datasets, and benchmark tasks are available at https://huggingface.co/collections/kuleshov-group/plantcad2-67e437e241a382671371a572. Fine-tuning pipelines and code are available at https://github.com/plantcad/plantcad.

Acknowledgements

This work is funded by a cooperative agreement from US Department of Agriculture Agricultural Research Service to Cornell University (MCR), NSF grant (#2240888), NSF CAREER grant (#2145577), NIH Maximizing Investigators' Research Awards (#1R35GM151243-01 and 5R35GM151348). SWe thank Travis Wrightsman (Cornell University) for sharing gene expression tasks, Michelle C. Stitzer (Cornell University) for plotting the tree, Hai Wang (China Agricultural University) for sharing max expression data of 17 species, Yaoyao Wu (Nanjing Agricultural University) for providing conserved and neutral SNP datasets, Sara Miller (Cornell

948 University) for helpful comments, and all members of Cornell's Institute of Genomic Diversity
949 (MCR & ESB) for helpful discussions. We would also like to thank the SCINet project, Texas
950 Advanced Computing Center at The University of Texas at Austin, and MosaicML for providing
951 compute resources for pretraining and fine-tuning experiments.

Author contributions

953 J.Z., A.G., V.K., and E.S.B. designed research; J.Z., A.G., S.-K.H., Z.-Y.L., S.-P.C., E.M., E.C.,
954 B.C., A.B., M.C.R., M.P., V.K., and E.S.B. performed research and analyses; J.Z., E.M., S.-K.H.,
955 M.P., and E.S.B. wrote the manuscript with all other authors' suggestions and comments.
956

Competing interests

958 The authors declare no competing interests.

Supplemental Information

Supplemental Table 1. Pretraining species and masked language modeling performance across
65 angiosperm genomes
Supplemental Table 2. Cross-species evolutionary conservation prediction performance
Supplemental Table 3. Masked motif prediction accuracy for transcriptional and translational
junction sites
Supplemental Table 4. Zero-shot structural variant impact prediction performance
Supplemental Table 5. Cross-species chromatin accessibility prediction trained on Arabidopsis
Supplemental Table 6. Multi-species chromatin accessibility prediction performance
Supplemental Table 7. Cell-type-specific chromatin accessibility prediction in maize
Supplemental Table 8. Gene expression prediction performance across species
Supplemental Table 9. Gene expression prediction performance with five random seeds
Supplemental Table 10. Translation prediction performance across species
Supplemental Table 11. The zero-shot performance of the tiny PlantCAD2 model with 20M
parameters
975

References

1. Vancaester, E., and Blaxter, M. (2023). Phylogenomic analysis of Wolbachia genomes from the Darwin Tree of Life biodiversity genomics project. PLoS Biol. *21*, e3001972.

2. Lewin, H.A., Robinson, G.E., Kress, W.J., Baker, W.J., Coddington, J., Crandall, K.A., Durbin, R., Edwards, S.V., Forest, F., Gilbert, M.T.P., et al. (2018). Earth BioGenome Project: Sequencing life for the future of life. Proc. Natl. Acad. Sci. U. S. A. *115*, 4325–4333.

3. Rhie, A., McCarthy, S.A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., et al. (2021). Towards complete and error-free genome assemblies of all vertebrate species. Nature *592*, 737–746.

4. Cheng, S., Melkonian, M., Smith, S.A., Brockington, S., Archibald, J.M., Delaux, P.-M., Li, F.-W., Melkonian, B., Mavrodiev, E.V., Sun, W., et al. (2018). 10KP: A phylodiverse genome sequencing plan. Gigascience *7*, 1–9.

5. Sun, Y., Shang, L., Zhu, Q.-H., Fan, L., and Guo, L. (2022). Twenty years of plant genome sequencing: achievements and challenges. Trends Plant Sci. *27*, 391–401.

6. Fu, L.-Y., Zhu, T., Zhou, X., Yu, R., He, Z., Zhang, P., Wu, Z., Chen, M., Kaufmann, K., and Chen, D. (2022). ChIP-Hub provides an integrative platform for exploring plant regulome. Nat. Commun. *13*, 3413.

7. Bommasani, R., Hudson, D.A., Adeli, E., Altman, R., Arora, S., von Arx, S., Bernstein, M.S., Bohg, J., Bosselut, A., Brunskill, E., et al. (2021). On the opportunities and risks of foundation models. arXiv [cs.LG].

8. Rives, A., Meier, J., Sercu, T., Goyal, S., Lin, Z., Liu, J., Guo, D., Ott, M., Zitnick, C.L., Ma, J., et al. (2021). Biological structure and function emerge from scaling unsupervised learning to 250 million protein sequences. Proc. Natl. Acad. Sci. U. S. A. *118*, e2016239118.

9. Lin, Z., Akin, H., Rao, R., Hie, B., Zhu, Z., Lu, W., Smetanin, N., Verkuil, R., Kabeli, O., Shmueli, Y., et al. (2023). Evolutionary-scale prediction of atomic-level protein structure with a language model. Science *379*, 1123–1130.

10. Hayes, T., Rao, R., Akin, H., Sofroniew, N.J., Oktay, D., Lin, Z., Verkuil, R., Tran, V.Q., Deaton, J., Wiggert, M., et al. (2025). Simulating 500 million years of evolution with a language model. Science *387*, eads0018.

11. Elnaggar, A., Heinzinger, M., Dallago, C., Rehawi, G., Wang, Y., Jones, L., Gibbs, T., Feher, T., Angerer, C., Steinegger, M., et al. (2022). ProtTrans: Toward Understanding the Language of Life Through Self-Supervised Learning. IEEE Trans. Pattern Anal. Mach. Intell. *44*, 7112–7127.

12. Brandes, N., Ofer, D., Peleg, Y., Rappoport, N., and Linial, M. (2022). ProteinBERT: a universal deep-learning model of protein sequence and function. Bioinformatics *38*, 2102–2110.

13. Kulmanov, M., Guzmán-Vega, F.J., Duek Roggli, P., Lane, L., Arold, S.T., and Hoehndorf, R. (2024). Protein function prediction as approximate semantic entailment. Nat. Mach. Intell.

1014   *6*, 220–228.

14. Chowdhury, R., Bouatta, N., Biswas, S., Floristean, C., Kharkar, A., Roy, K., Rochereau, C., Ahdritz, G., Zhang, J., Church, G.M., et al. (2022). Single-sequence protein structure prediction using a language model and deep learning. Nat. Biotechnol. *40*, 1617–1623.

15. Brandes, N., Goldman, G., Wang, C.H., Ye, C.J., and Ntranos, V. (2023). Genome-wide prediction of disease variant effects with a deep protein language model. Nat. Genet. *55*, 1512–1522.

16. Ji, Y., Zhou, Z., Liu, H., and Davuluri, R.V. (2021). DNABERT: pre-trained Bidirectional Encoder Representations from Transformers model for DNA-language in genome. Bioinformatics *37*, 2112–2120.

17. Zhou, Z., Ji, Y., Li, W., Dutta, P., Davuluri, R.V., and Liu, H. (2023). DNABERT-2: Efficient Foundation Model and Benchmark For Multi-Species Genomes.

18. Nguyen, E., Poli, M., Durrant, M.G., Kang, B., Katrekar, D., Li, D.B., Bartie, L.J., Thomas, A.W., King, S.H., Brixi, G., et al. (2024). Sequence modeling and design from molecular to genome scale with Evo. Science *386*, eado9336.

19. Brixi, G., Durrant, M.G., Ku, J., Poli, M., Brockman, G., Chang, D., Gonzalez, G.A., King, S.H., Li, D.B., Merchant, A.T., et al. (2025). Genome modeling and design across all domains of life with Evo 2. bioRxiv. https://doi.org/10.1101/2025.02.18.638918.

20. Benegas, G., Batra, S.S., and Song, Y.S. (2023). DNA language models are powerful predictors of genome-wide variant effects. Proc. Natl. Acad. Sci. U. S. A. *120*, e2311219120.

21. Benegas, G., Albors, C., Aw, A.J., Ye, C., and Song, Y.S. (2025). A DNA language model based on multispecies alignment predicts the effects of genome-wide variants. Nat. Biotechnol., 1–6.

22. Mendoza-Revilla, J., Trop, E., Gonzalez, L., Roller, M., Dalla-Torre, H., de Almeida, B.P., Richard, G., Caton, J., Lopez Carranza, N., Skwark, M., et al. (2024). A foundational large language model for edible plant genomes. Commun. Biol. *7*, 835.

23. Zhai, J., Gokaslan, A., Schiff, Y., Berthel, A., Liu, Z.-Y., Lai, W.-Y., Miller, Z.R., Scheben, A., Stitzer, M.C., Romay, M.C., et al. (2025). Cross-species modeling of plant genomes at single-nucleotide resolution using a pretrained DNA language model. Proc. Natl. Acad. Sci. U. S. A. *122*, e2421738122.

24. Schiff, Y., Kao, C.-H., Gokaslan, A., Dao, T., Gu, A., and Kuleshov, V. (2024). Caduceus: Bi-Directional Equivariant Long-Range DNA Sequence Modeling. arXiv [q-bio.GN].

25. Dalla-Torre, H., Gonzalez, L., Mendoza-Revilla, J., Lopez Carranza, N., Grzywaczewski, A.H., Oteri, F., Dallago, C., Trop, E., de Almeida, B.P., Sirelkhatim, H., et al. (2025). Nucleotide Transformer: building and evaluating robust foundation models for human genomics. Nat. Methods *22*, 287–297.

26. Wang, N., Bian, J., Li, Y., Li, X., Mumtaz, S., Kong, L., and Xiong, H. (2024). Multi-purpose RNA language modelling with motif-aware pretraining and type-guided fine-tuning. Nat. Mach. Intell. *6*, 548–557.

27. Shen, T., Hu, Z., Sun, S., Liu, D., Wong, F., Wang, J., Chen, J., Wang, Y., Hong, L., Xiao, J., et al. (2024). Accurate RNA 3D structure prediction using a language model-based deep learning approach. Nat. Methods *21*, 2287–2298.

28. Yu, H., Yang, H., Sun, W., Yan, Z., Yang, X., Zhang, H., Ding, Y., and Li, K. (2024). An interpretable RNA foundation model for exploring functional RNA motifs in plants. Nat. Mach. Intell. *6*, 1616–1625.

29. Hao, M., Gong, J., Zeng, X., Liu, C., Guo, Y., Cheng, X., Wang, T., Ma, J., Zhang, X., and Song, L. (2024). Large-scale foundation model on single-cell transcriptomics. Nat. Methods *21*, 1481–1491.

30. Zeng, Y., Xie, J., Shangguan, N., Wei, Z., Li, W., Su, Y., Yang, S., Zhang, C., Zhang, J., Fang, N., et al. (2025). CellFM: a large-scale foundation model pre-trained on transcriptomics of 100 million human cells. Nat. Commun. *16*, 4679.

31. Cui, H., Wang, C., Maan, H., Pang, K., Luo, F., Duan, N., and Wang, B. (2024). scGPT: toward building a foundation model for single-cell multi-omics using generative AI. Nat. Methods *21*, 1470–1480.

32. Devlin, J., Chang, M.-W., Lee, K., and Toutanova, K. (2018). BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding. arXiv [cs.CL].

33. Zhou, J., and Troyanskaya, O.G. (2015). Predicting effects of noncoding variants with deep learning-based sequence model. Nat. Methods *12*, 931–934.

34. Quang, D., and Xie, X. (2016). DanQ: a hybrid convolutional and recurrent deep neural network for quantifying the function of DNA sequences. Nucleic Acids Res. *44*, e107.

35. Kelley, D.R., Snoek, J., and Rinn, J.L. (2016). Basset: learning the regulatory code of the accessible genome with deep convolutional neural networks. Genome Res. *26*, 990–999.

36. Benegas, G., Ye, C., Albors, C., Li, J.C., and Song, Y.S. (2025). Genomic language models: opportunities and challenges. Trends Genet. *41*, 286–302.

37. Gu, A., and Dao, T. (2023). Mamba: Linear-Time Sequence Modeling with Selective State Spaces. arXiv [cs.LG].

38. Schmitz, R.J., Grotewold, E., and Stam, M. (2022). Cis-regulatory sequences in plants: Their importance, discovery, and future challenges. Plant Cell *34*, 718–741.

39. Marand, A.P., Eveland, A.L., Kaufmann, K., and Springer, N.M. (2023). Cis-regulatory elements in plant development, adaptation, and evolution. Annu. Rev. Plant Biol. *74*, 111–137.

40. Engelhorn, J., Snodgrass, S.J., Kok, A., Seetharam, A.S., Schneider, M., Kiwit, T., Singh, A., Banf, M., Khaipho-Burch, M., Runcie, D.E., et al. (2024). Genetic variation at transcription factor binding sites largely explains phenotypic heritability in maize. bioRxiv, 2023.08.08.551183. https://doi.org/10.1101/2023.08.08.551183.

41. Marand, A.P., Jiang, L., Gomez-Cano, F., Minow, M.A.A., Zhang, X., Mendieta, J.P., Luo, Z., Bang, S., Yan, H., Meyer, C., et al. (2025). The genetic architecture of cell type-specific cis regulation in maize. Science *388*, eads6601.

42. Magallón, S., and Castillo, A. (2009). Angiosperm diversification through time. Am. J. Bot.
96, 349–365.

43. Bar-On, Y.M., Phillips, R., and Milo, R. (2018). The biomass distribution on Earth. Proc. Natl.
Acad. Sci. U. S. A. *115*, 6506–6511.

44. Dao, T., and Gu, A. (2024). Transformers are SSMs: Generalized models and efficient
algorithms through structured state space duality. arXiv [cs.LG].

45. Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A.N., Kaiser, L., and
Polosukhin, I. (2017). Attention Is All You Need. arXiv [cs.CL].

46. Warner, B., Chaffin, A., Clavié, B., Weller, O., Hallström, O., Taghadouini, S., Gallagher, A.,
Biswas, R., Ladhak, F., Aarsen, T., et al. (2024). Smarter, better, faster, longer: A modern
bidirectional encoder for fast, memory efficient, and long context finetuning and inference.
arXiv [cs.CL].

47. Gu, A., Goel, K., and Ré, C. (2021). Efficiently Modeling Long Sequences with Structured
State Spaces. arXiv [cs.LG].

48. Novák, P., Guignard, M.S., Neumann, P., Kelly, L.J., Mlinarec, J., Koblížková, A., Dodsworth,
S., Kovařík, A., Pellicer, J., Wang, W., et al. (2020). Repeat-sequence turnover shifts
fundamentally in species with large genomes. Nat Plants *6*, 1325–1329.

49. Sun, S., Wang, B., Li, C., Xu, G., Yang, J., Hufford, M.B., Ross-Ibarra, J., Wang, H., and
Wang, L. (2023). Unraveling Prevalence and Effects of Deleterious Mutations in Maize Elite
Lines across Decades of Modern Breeding. Mol. Biol. Evol. *40*.
https://doi.org/10.1093/molbev/msad170.

50. Lozano, R., Gazave, E., dos Santos, J.P.R., Stetter, M.G., Valluru, R., Bandillo, N.,
Fernandes, S.B., Brown, P.J., Shakoor, N., Mockler, T.C., et al. (2021). Comparative
evolutionary genetics of deleterious load in sorghum and maize. Nature Plants *7*, 17–24.

51. Lye, Z., Choi, J.Y., and Purugganan, M.D. (2022). Deleterious Mutations and the Rare Allele
Burden on Rice Gene Expression. Mol. Biol. Evol. *39*.
https://doi.org/10.1093/molbev/msac193.

52. Mezmouk, S., and Ross-Ibarra, J. (2014). The pattern and distribution of deleterious
mutations in maize. G3 *4*, 163–171.

53. Song, B., Buckler, E.S., and Stitzer, M.C. (2024). New whole-genome alignment tools are
needed for tapping into plant diversity. Trends Plant Sci. *29*, 355–369.

54. Huber, C.D., Kim, B.Y., and Lohmueller, K.E. (2020). Population genetic models of GERP
scores suggest pervasive turnover of constrained sites across mammalian evolution. PLoS
Genet. *16*, e1008827.

55. Bennetzen, J.L., and Wang, H. (2014). The contributions of transposable elements to the
structure, function, and evolution of plant genomes. Annu. Rev. Plant Biol. *65*, 505–530.

56. Kozak, M. (1986). Point mutations define a sequence flanking the AUG initiator codon that
modulates translation by eukaryotic ribosomes. Cell *44*, 283–292.

57. Davydov, E.V., Goode, D.L., Sirota, M., Cooper, G.M., Sidow, A., and Batzoglou, S. (2010).
Identifying a high fraction of the human genome to be under selective constraint using
GERP++. PLoS Comput. Biol. *6*, e1001025.

58. Wu, Y., Li, D., Hu, Y., Li, H., Ramstein, G.P., Zhou, S., Zhang, X., Bao, Z., Zhang, Y., Song,
B., et al. (2023). Phylogenomic discovery of deleterious mutations facilitates hybrid potato
breeding. Cell *186*, 2313-2328.e15.

59. Li, Y.-R., and Liu, M.-J. (2020). Prevalence of alternative AUG and non-AUG translation
initiators and their regulatory effects across plants. Genome Res. *30*, 1418–1433.

60. Tian, F., Yang, D.-C., Meng, Y.-Q., Jin, J., and Gao, G. (2020). PlantRegMap: charting
functional regulatory maps in plants. Nucleic Acids Res. *48*, D1104–D1113.

61. Hubisz, M.J., Pollard, K.S., and Siepel, A. (2011). PHAST and RPHAST: phylogenetic
analysis with space/time models. Brief. Bioinform. *12*, 41–51.

62. O'Malley, R.C., Huang, S.-S.C., Song, L., Lewsey, M.G., Bartlett, A., Nery, J.R., Galli, M.,
Gallavotti, A., and Ecker, J.R. (2016). Cistrome and epicistrome features shape the
regulatory DNA landscape. Cell *165*, 1280–1292.

63. Shrikumar, A., Tian, K., Avsec, Ž., Shcherbina, A., Banerjee, A., Sharmin, M., Nair, S., and
Kundaje, A. (2018). Technical note on Transcription Factor Motif Discovery from Importance
Scores (TF-MoDISco) version 0.5.6.5. arXiv [cs.LG].
https://doi.org/10.48550/arXiv.1811.00416.

64. Kerstens, M.H.L., Schranz, M.E., and Bouwmeester, K. (2020). Phylogenomic analysis of the
APETALA2 transcription factor subfamily across angiosperms reveals both deep
conservation and lineage-specific patterns. Plant J. *103*, 1516–1524.

65. Kooiker, M., Airoldi, C.A., Losa, A., Manzotti, P.S., Finzi, L., Kater, M.M., and Colombo, L.
(2005). BASIC PENTACYSTEINE1, a GA binding protein that induces conformational
changes in the regulatory region of the homeotic Arabidopsis gene SEEDSTICK. Plant Cell
17, 722–729.

66. Liu, Z., Running, M.P., and Meyerowitz, E.M. (1997). TSO1 functions in cell division during
Arabidopsis flower development. Development *124*, 665–672.

67. Fujii, S., Yamamoto, E., Ito, S., Tangpranomkorn, S., Kimura, Y., Miura, H., Yamaguchi, N.,
Kato, Y., Niidome, M., Yoshida, A., et al. (2023). SHI family transcription factors regulate an
interspecific barrier. Nat. Plants *9*, 1862–1873.

68. Shaikhali, J., de Dios Barajas-Lopéz, J., Ötvös, K., Kremnev, D., Garcia, A.S., Srivastava,
V., Wingsle, G., Bako, L., and Strand, Å. (2012). The CRYPTOCHROME1-dependent
response to excess light is mediated through the transcriptional activators ZINC FINGER
PROTEIN EXPRESSED IN INFLORESCENCE MERISTEM LIKE1 and ZML2 in Arabidopsis.
Plant Cell *24*, 3009–3025.

69. Lu, Z., Marand, A.P., Ricci, W.A., Ethridge, C.L., Zhang, X., and Schmitz, R.J. (2019). The
prevalence, evolution and chromatin signatures of plant regulatory elements. Nat. Plants *5*,
1250–1259.

70. Hu, E.J., Shen, Y., Wallis, P., Allen-Zhu, Z., Li, Y., Wang, S., Wang, L., and Chen, W. (2021).
LoRA: Low-Rank Adaptation of large language models. arXiv [cs.CL].

71. Chen, T., and Guestrin, C. (2016). XGBoost: A Scalable Tree Boosting System. arXiv [cs.LG].

72. Marand, A.P., Chen, Z., Gallavotti, A., and Schmitz, R.J. (2021). A cis-regulatory atlas in
maize at single-cell resolution. Cell *184*, 3041-3055.e21.

73. Wrightsman, T., Ferebee, T.H., Romay, M.C., Seetharam, A.S., AuBuchon-Elder, T., Phillips,
A.R., Syring, M., Hufford, M.B., Kellogg, E.A., and Buckler, E.S. (2024). Current genomic
deep learning architectures generalize across grass species but not alleles. bioRxiv,
2024.04.11.589024. https://doi.org/10.1101/2024.04.11.589024.

74. Huang, C., Shuai, R.W., Baokar, P., Chung, R., Rastogi, R., Kathail, P., and Ioannidis, N.M.
(2023). Personal transcriptome variation is poorly explained by current genomic deep
learning models. Nat. Genet. *55*, 2056–2059.

75. Tang, Z., Toneyan, S., and Koo, P.K. (2023). Current approaches to genomic deep learning
struggle to fully capture human genetic variation. Nat. Genet. *55*, 2021–2022.

76. Hale, C.O., Hsu, S.-K., Zhai, J., Schulz, A.J., Aubuchon-Elder, T., Costa-Neto, G., Gelfond,
A., El-Walid, M., Hufford, M., Kellogg, E.A., et al. (2025). Widespread turnover of a conserved
cis-regulatory code across 589 grass species. Mol. Biol. Evol., msaf324.

77. Jaganathan, K., Ersaro, N., Novakovsky, G., Wang, Y., James, T., Schwartzentruber, J.,
Fiziev, P., Kassam, I., Cao, F., Hawe, J., et al. (2025). Predicting expression-altering
promoter mutations with deep learning. Science *389*, eads7373.

78. Giri, A., Khaipho-Burch, M., Buckler, E.S., and Ramstein, G.P. (2021). Haplotype associated
RNA expression (HARE) improves prediction of complex traits in maize. PLoS Genet. *17*,
e1009568.

79. Ghosal, S., Barhomi, Y., Ganapathi, T., Krystosik, A., Krishnan, L., Guntury, S., Li, D.,
Alzheimer's Disease Neuroimaging Initiative, Casale, F.P., and Karaletsos, T. (2025).
VariantFormer: A hierarchical transformer integrating DNA sequences with genetic variations
and regulatory landscapes for personalized gene expression prediction. bioRxiv,
2025.10.31.685862. https://doi.org/10.1101/2025.10.31.685862.

80. Liu, T., Zhang, X., Ying, R., and Zhao, H. (2025). Pre-training genomic language model with
variants for better modeling functional genomics. bioRxiv, 2025.02.26.640468.
https://doi.org/10.1101/2025.02.26.640468.

81. Li, T., Xu, H., Teng, S., Suo, M., Bahitwa, R., Xu, M., Qian, Y., Ramstein, G.P., Song, B.,
Buckler, E.S., et al. (2024). Modeling 0.6 million genes for the rational design of functional
cis-regulatory variants and de novo design of cis-regulatory sequences. Proc. Natl. Acad.
Sci. U. S. A. *121*, e2319811121.

82. Kaplan, J., McCandlish, S., Henighan, T., Brown, T.B., Chess, B., Child, R., Gray, S.,
Radford, A., Wu, J., and Amodei, D. (2020). Scaling laws for neural language models. arXiv
[cs.LG]. https://doi.org/10.48550/arXiv.2001.08361.

83. Marks, R.A., Hotaling, S., Frandsen, P.B., and VanBuren, R. (2021). Representation and

participation across 20 years of plant genome sequencing. Nat Plants *7*, 1571–1578.

84. Bender, E.M., Gebru, T., McMillan-Major, A., and Shmitchell, S. (2021). On the dangers of
stochastic parrots: Can language models be too big? 🦜. In Proceedings of the 2021 ACM
Conference on Fairness, Accountability, and Transparency (ACM).
https://doi.org/10.1145/3442188.3445922.

85. Geirhos, R., Jacobsen, J.-H., Michaelis, C., Zemel, R., Brendel, W., Bethge, M., and
Wichmann, F.A. (2020). Shortcut learning in deep neural networks. Nat. Mach. Intell. *2*, 665–
673.

86. Ding, F., and Steinhardt, J. (2024). Protein language models are biased by unequal sequence
sampling across the tree of life. bioRxiv, 2024.03.07.584001.
https://doi.org/10.1101/2024.03.07.584001.

87. Stitzer, M.C., Anderson, S.N., Springer, N.M., and Ross-Ibarra, J. (2021). The genomic
ecosystem of transposable elements in maize. PLoS Genet. *17*, e1009768.

88. Liu, Z.-Y., Berthel, A., Czech, E., Stitzer, M., Hsu, S.-K., Pennell, M., Buckler, E.S., and Zhai,
J. (2025). GeneCAD: Plant genome annotation with a DNA foundation model. bioRxivorg,
2025.10.31.685877. https://doi.org/10.1101/2025.10.31.685877.

89. Clark, R.M., Wagler, T.N., Quijada, P., and Doebley, J. (2006). A distant upstream enhancer
at the maize domestication gene tb1 has pleiotropic effects on plant and inflorescent
architecture. Nat. Genet. *38*, 594–597.

90. Studer, A., Zhao, Q., Ross-Ibarra, J., and Doebley, J. (2011). Identification of a functional
transposon insertion in the maize domestication gene tb1. Nat. Genet. *43*, 1160–1163.

91. Tu, X., Mejía-Guerra, M.K., Valdes Franco, J.A., Tzeng, D., Chu, P.-Y., Shen, W., Wei, Y.,
Dai, X., Li, P., Buckler, E.S., et al. (2020). Reconstructing the maize leaf regulatory network
using ChIP-seq data of 104 transcription factors. Nat. Commun. *11*, 5089.

92. Smith, S.A., and Brown, J.W. (2018). Constructing a broadly inclusive seed plant phylogeny.
Am. J. Bot. *105*, 302–314.

93. Loshchilov, I., and Hutter, F. (2017). Decoupled Weight Decay Regularization. arXiv [cs.LG].

94. Song, B., Buckler, E.S., Wang, H., Wu, Y., Rees, E., Kellogg, E.A., Gates, D.J., Khaipho-
Burch, M., Bradbury, P.J., Ross-Ibarra, J., et al. (2021). Conserved noncoding sequences
provide insights into regulatory sequence and loss of gene expression in maize. Genome
Res. *31*, 1245–1257.

95. Welker, C.A.D., McKain, M.R., Estep, M.C., Pasquet, R.S., Chipabika, G., Pallangyo, B., and
Kellogg, E.A. (2020). Phylogenomics enables biogeographic analysis and a new subtribal
classification of Andropogoneae (Poaceae—Panicoideae). J. Syst. Evol. *58*, 1003–1030.

96. Stitzer, M.C., Seetharam, A.S., Scheben, A., Hsu, S.-K., Schulz, A.J., AuBuchon-Elder, T.M.,
El-Walid, M., Ferebee, T.H., Hale, C.O., La, T., et al. (2025). Extensive genome evolution
distinguishes maize within a stable tribe of grasses. bioRxivorg, 2025.01.22.633974.
https://doi.org/10.1101/2025.01.22.633974.

97. Song, B., Marco-Sola, S., Moreto, M., Johnson, L., Buckler, E.S., and Stitzer, M.C. (2022).
AnchorWave: Sensitive alignment of genomes with high sequence diversity, extensive
structural polymorphism, and whole-genome duplication. Proc. Natl. Acad. Sci. U. S. A. *119*.
https://doi.org/10.1073/pnas.2113075119.

98. Pollard, K.S., Hubisz, M.J., Rosenbloom, K.R., and Siepel, A. (2010). Detection of nonneutral
substitution rates on mammalian phylogenies. Genome Res. *20*, 110–121.

99. Hufford, M.B., Seetharam, A.S., Woodhouse, M.R., Chougule, K.M., Ou, S., Liu, J., Ricci,
W.A., Guo, T., Olson, A., Qiu, Y., et al. (2021). De novo assembly, annotation, and
comparative analysis of 26 diverse maize genomes. Science *373*, 655–662.

100.Gao, L., Gonda, I., Sun, H., Ma, Q., Bao, K., Tieman, D.M., Burzynski-Chang, E.A., Fish,
T.L., Stromberg, K.A., Sacks, G.L., et al. (2019). The tomato pan-genome uncovers new
genes and a rare allele regulating fruit flavor. Nat. Genet. *51*, 1044–1051.

101.Wrightsman, T., Marand, A.P., Crisp, P.A., Springer, N.M., and Buckler, E.S. (2022).
Modeling chromatin state from sequence across angiosperms using recurrent convolutional
neural networks. Plant Genome *15*, e20249.

102.Wu, H.-Y.L., Ai, Q., Teixeira, R.T., Nguyen, P.H.T., Song, G., Montes, C., Elmore, J.M.,
Walley, J.W., and Hsu, P.Y. (2024). Improved super-resolution ribosome profiling reveals
prevalent translation of upstream ORFs and small ORFs in Arabidopsis. Plant Cell *36*, 510–
539.

103.Zhu, W., Miao, X., Qian, J., Chen, S., Jin, Q., Li, M., Han, L., Zhong, W., Xie, D., Shang, X.,
et al. (2023). A translatome-transcriptome multi-omics gene regulatory network reveals the
complicated functional landscape of maize. Genome Biol. *24*, 60.

104.Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics *30*, 2114–2120.

105.Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. Genome Biol. *10*, R25.

106.Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*,
15–21.

107.Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.-C., Mendell, J.T., and Salzberg, S.L.
(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.
Nat. Biotechnol. *33*, 290–295.

108.Mangrulkar, S., Gugger, S., Debut, L., Belkada, Y., Paul, S., and Bossan, B. (2022). Peft:
State-of-the-art parameter-efficient fine-tuning methods. In Peft: State-of-the-art parameter-
efficient fine-tuning methods.

Figure S1. PlantCAD2 model performance across different scales and species. (A) Inference speed benchmarking shows that the Caduceus architecture exhibits significantly faster CUDA and wall-clock runtime than ModernBERT across increasing sequence lengths. **(B)** Masked language modeling accuracy across the three models, evaluated by randomly masking 15% of nucleotides per sequence, showing improved performance with increasing model size. **(C)** Correlation matrix of per-species masked language modeling accuracy of three PlantCAD2 models. **(D)** Relationship between genome size and masked language modeling accuracy for PlantCAD2-L. **(E)** Relationship between the number of repetitive sequences in the test set and masked language modeling accuracy.

Figure S2. Effect of context window length on AUROC performance. Evaluation across three PlantCAD2 models for conservation within Andropogoneae **(A)**, Poaceae non-TIS **(B)**, Poaceae TIS **(C)**, and Solanaceae **(D)**.

Figure S3. AUROC of different genomic contexts for conservation prediction in Solanaceae.

Figure S4. PlantCAD2 distinguishes core from non-core junctions with zero-shot strategy. ROC curves showing binary classification performance for distinguishing evolutionarily constrained core junctions from rapidly evolving non-core junctions using model log-likelihood scores at junction sites. Left panels show maize (included in pre-training), right panels show tomato (excluded from training). **(A)** Translation initiation sites. **(B)** Translation termination sites. **(C)** Splice donor sites. **(D)** Splice acceptor sites.

Figure S5. PlantCAD2 predicts functional impact of structural variants using zero-shot strategy. (A) Distribution of phastCons. (B) Scatter plot showing the positive correlation between PlantCAD2's -ΔlogP scores and phyloP-based conservation scores. (C) AUROC performance distinguishes highly conserved from less conserved deletions.

Figure S6. Effect of flanking sequence length on deletion classification performance. AUROC performance for distinguishing conserved from less conserved deletions by flanking sequence length.

Figure S7. Motifs generated by TF-MoDISco from PlantCAD2 and Evo2 high-confidence predictions for DREB2 (AP2/ERF family).

Figure S8. Example motifs generated by TF-MoDISco from PlantCAD2 high-confidence predictions in Arabidopsis promoter regions.

Figure S9. Accessible region prediction task comparison with Evo2 using frozen embeddings. **(A)** XGBoost classifier, **(B)** neural network with single linear layer.

Figure S10. Relationship between genome size and accessibility prediction performance.

Figure S11. Gene expression prediction performance in rice and tomato. (A) Leaf expression on/off prediction, **(B)** leaf absolute expression prediction.

Figure S12. Effect of context window length on gene expression prediction performance.
Performance improvement with increasing context window size for binary leaf expression
classification **(A)** and absolute expression regression **(B)** on the maize NAM dataset. Error bars
represent five replicates with different random seeds.

Figure S13. Per-orthogroup expression correlation performance within the maize NAM population. Spearman correlation coefficients (SCC) for predicting allele-specific expression differences within orthogroups.

Figure S14. Cross-species transfer performance for gene expression prediction. ROC curves showing performance when training on Arabidopsis and testing cross-species transfer to maize.

Figure S15. Gene length distribution of 65 pretraining genomes.