1 ***Plasmodium* ARK1 regulates spindle formation during atypical mitosis and forms a**
2 **divergent chromosomal passenger complex**
3

4 Annu Nagar[1+], Ryuji Yanase[2+], Mohammad Zeeshan[2+], David J. P. Ferguson[3], Steven Abel[4],
5 Sarah L. Pashley[2], Akancha Mishra[2], Antonius Eze[2], Edward Rea[2], Declan Brady[2], Andrew R.
6 Bottrill[5], Sue Vaughan[3], Karine G. Le Roch[4], David S. Guttery[2], Anthony A. Holder[6], Eelco C.
7 Tromer[7&], Pushkar Sharma[1&], Rita Tewari[2&]
8

[1]Eukaryotic Gene Expression Laboratory, National Institute of Immunology, New Delhi, India-
10067.
[2] School of Life Sciences, University of Nottingham, Nottingham, UK.
[3]Department of Biological and Medical Sciences, Oxford Brookes University, Oxford, UK.
[4]Department of Molecular, Cell and Systems Biology, University of California Riverside, 900
University Ave., Riverside, USA.
[5]School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry, UK.
[6]Molecular Parasitology Laboratory, The Francis Crick Institute, London, UK.
[7]Faculty of Science and Engineering, Groningen Biomolecular Sciences and Biotechnology
Institute, Cell Biochemistry, University of Groningen, Groningen, The Netherlands.

19
20 [+] Contributed equally
21 [&] joint corresponding author:
22 Tromer, E.C; e.c.tromer@rug.nl
23 Pushkar Sharma; pushkar@nii.ac.in
24 Rita Tewari; rita.tewari@nottingham.ac.uk (Lead)

Abstract

Mitosis in *Plasmodium* spp., the causative agent of malaria, is fundamentally different from model eukaryotes, proceeding via a bipartite microtubule organising centre (MTOC) and lacking canonical regulators such as Polo and Bub1 kinases. During schizogony, asynchronous nuclear replication produces a multinucleate schizont, while rapid male gametogony generates an octaploid nucleus before gamete formation. Here, we identify Aurora-related kinase 1 (ARK1) as a key component of inner MTOC and spindle formation, controlling kinetochore dynamics and driving mitotic progression. Conditional ARK1 depletion disrupts spindle biogenesis, kinetochore segregation, karyokinesis and cytokinesis in both stages, and affects parasite transmission. Interactome analysis reveals ARK1 as the catalytic core of a non-canonical chromosomal passenger complex (CPC) containing two divergent inner centromere proteins (INCENPs) but lacking Survivin and Borealin. Comparative genomics indicates this CPC architecture arose early in Apicomplexa, replacing canonical centromere-targeting modules. These findings uncover a distinct mitotic machinery in *Plasmodium* and identify the ARK1–INCENP interface as a potential multistage target for malaria therapeutic intervention.

Keywords: *Plasmodium*; Aurora kinase; chromosomal passenger complex (CPC); INCENP; spindle; karyokinesis; male gametogony; MTOC; cytokinesis; transmission

Introduction

Plasmodium spp., the causative agent of malaria, is a haploid unicellular eukaryote of the phylum Apicomplexa, which is evolutionarily distant from most model organisms and has some unusual divergent features of cell division. During its life cycle, mitosis is closed and typified by two distinct forms: schizogony within vertebrate host cells, and male gametogony within the female mosquito gut (Fig. 1a). During asexual proliferation within host red blood cells (schizogony), repeated asynchronous nuclear division, without chromosome condensation, produces a multinucleate schizont. Subsequent budding produces invasive merozoites to continue the cycle. In contrast, during male gametogony three rounds of rapid genome replication to octaploid (1N to 8N) occur in 8 minutes within the intact nucleus. Karyokinesis and cytokinesis are synchronised with axoneme dynamics by replication and separation of a bipartite microtubule-organising centre (MTOC), chromosome condensation, and complementary de novo basal body formation, leading to the budding of eight flagellated male gametes. These events occur within 12 to 15 min of gametocyte activation, which is probably the most rapid mitosis amongst eukaryotes[1]. Compared to schizogony, mitosis in male gametogony is unusual, with atypical continuous spindles and an MTOC that spans the nuclear envelope connecting the spindle microtubules with those of the axoneme. Male gametes are the only flagellated stage in the parasite's life cycle, and therefore the only stage where centriole formation and de novo replication of the basal body occurs[1-4].

Aurora-related kinases (ARKs) are multi-functional mitotic kinases that regulate eukaryotic cell division. They act on spindle pole duplication, kinetochore-microtubule attachment, spindle assembly, mitotic checkpoints and cytokinesis[5-8]. Budding yeast and the amoeba *Dictyostelium discoideum*[9] have a single ARK to execute all functions, suggesting these to be ancestral to all eukaryotes[7]. However, multiple different lineages (e.g. vertebrates, plants, kinetoplastids) have one or more paralogs that resulted from parallel duplications, which subdivided functions over the different paralogs[7,10]. One paralog (Aurora B in humans) is the catalytic centre of the chromosomal passenger complex (CPC), a key regulator of mitotic events at the cell equator, which initially hitchhikes on the inner centromere and finally remains at the central spindle to orchestrate cytokinesis. In addition to the single Aurora kinase, the CPC consists of inner centromere protein (INCENP), Survivin and Borealin[11,12] that guide its localisation to kinetochores, spindle and the central spindle. It is also critical for mitotic histone phosphorylation, fixing errors in kinetochore-spindle attachment, and correct cytokinesis (Fig. 1b).

In *Plasmodium*, there are also three ARKs, which have divergent sequence and mostly lack canonical scaffolds[10,13,14]. ARK1, 2 and 3 are refractory to gene disruption and indispensable during asexual blood stage schizogony in both *P. berghei* and *P. falciparum*[13-15]. ARK1 associates with the spindle pole body during schizogony[16,17], but its function is not understood during this process and its location and function in sexual stages remains unknown. Our recent studies on ARK2 revealed its association with spindle microtubules (MTs) forming a divergent protein complex with EB1, myosin K and MISFIT[10], but little is known of ARK1 and 3 functions during male gametogony or whether they are essential for parasite transmission. ARK1 function has been studied in another apicomplexan, *Toxoplasma gondii*, where it interacts with INCENP1-2 and is important for asexual replication, but its function during male gametogony is difficult to study because the sexual cycle of these parasites only takes place in the gut of cats (Fig. 1b)[18].

Here, we have investigated *Plasmodium* ARK1, integrating studies of location, function and molecular interactions across two life-cycle stages using *P. falciparum* (Pf, for asexual blood-stage schizogony) and *P. berghei* (Pb, for male gametogony). High-resolution live-cell imaging and expansion microscopy revealed ARK1 associated with the mitotic spindle, kinetochores, and the inner MTOC. Conditional gene knockout using rapamycin-DiCre in *P. falciparum*[19] and gene knockdown using a promoter-trap approach in *P. berghei*[20] revealed that ARK1 regulates spindle biogenesis, karyokinesis and cytokinesis during both schizogony and male gametogony, suggesting that ARK1 is a critical regulator of *Plasmodium* mitosis. Interactome analysis also identified *Plasmodium* ARK1 as a component of a non-canonical Chromosomal Passenger Complex (CPC) comprising the two divergent INCENP1 and INCENP2, but without the canonical subunits Survivin and Borealin. These data underscore the evolutionary divergence of the *Plasmodium* mitotic machinery from that of its host, and therefore its potential as a target for therapeutic intervention at multiple stages of the lifecycle.

Results

ARK1 is present in asexual stage parasites undergoing mitotic division

We studied the location of PfARK1 in asexual blood stages, using a rapamycin-inducible dimerizable Cre recombinase (DiCre)-based strategy to knockout the ARK1 gene in the 1G5DC parasite line (Extended Data Fig. 1)[19]. As a first step in this strategy, we generated a HA-PfARK1-loxP parasite line, and these parasites were used to locate ARK1 during parasite

division using an antibody to the HA-tag. Indirect immunofluorescence (IFA) indicated that

ARK1 is expressed in trophozoite and schizont stages, but not detectable in late schizonts and

all developing merozoites (Fig. 2a). ARK1 was present as a punctum proximal to the MTOC

also known as a centriolar plaque, as identified by staining with an anti-centrin antibody,

present only in cells with two or more associated centrin puncta, indicative of ongoing mitosis

(Fig. 2a). To gain a deeper insight into ARK1 dynamics, Ultrastructure Expansion

Microscopy (U-ExM) was performed using an anti-tubulin antibody to track spindle

biogenesis. As reported previously[21], parasites possessed hemispindles prior to the onset of

division, but PfARK1 was not expressed in these parasites or in parasites containing interpolar

spindles (Fig. 2b, c). Strikingly, PfARK1 was specifically found associated with mitotic

spindles and PfARK1-positive double puncta seemed to be present at the polar ends of the

mitotic spindles (Fig. 2b, c). These data strongly indicate that ARK1 is expressed transiently,

and only when parasites undergo mitotic division.

These observations were also validated using the rodent malaria parasite, *P. berghei*, a well-

established model for studying gametogony and parasite development within the mosquito host.

The subcellular location of PbARK1 in asexual stages was examined by generating an

endogenous, C-terminal GFP-tagged PbARK1 (PbARK1-GFP) transgenic parasite line

(Extended Data Fig. 2a-c and Supplementary Notes). Live cell imaging of PbARK1-GFP in

asexual blood stages revealed a diffuse location in early ring stages. During trophozoite and

early schizont stages, PbARK1-GFP re-localised to discrete focal points in the nucleus

(Extended Data Fig. 2d). As schizogony progressed, the focal points of fluorescence divided

into two closely associated puncta, which followed subsequent asynchronous nuclear divisions

and eventually became diffuse during the later stages (Extended Data Fig. 2d, e). These

observations are consistent with the ARK1 location in *P. falciparum* as described above (Fig.

2a).

PbARK1 is located in the nucleoplasm, at the spindle poles and at microtubules during

male gametogony

In male gametocytes, PbARK1-GFP had a diffuse distribution predominantly within the

nucleoplasm. Within one-min post-activation this was concentrated to a single discrete focal

point adjacent to the DNA (Fig. 2d), and then the distribution rapidly extended to form an arc-

like structure, which within three-min post-activation divided into two distinct puncta

positioned at opposite sides of the nucleus (Fig. 2d and Supplementary Video 1). The two

PbARK1-GFP foci elongated again, forming two arc-like structures along the periphery of the

nucleus (Fig. 2d and Supplementary Video 2). By 6- to 8-min post-activation, a subsequent

round of extension and division had produced eight discrete PbARK1-GFP foci, coincident

with the completion of three successive rounds of mitosis (Fig. 2d and Supplementary Video

3).

The arc-like pattern of PbARK1 observed in live cell imaging coincided with the pattern of

anti-tubulin staining in IFA and U-ExM, suggesting an association with the mitotic spindle and

spindle poles (Fig 2e and Extended Data Fig. 2f). To further investigate ARK1 location,

dynamics, and association with mitotic spindles and kinetochore segregation, we compared the

location of PbARK1 with that of the kinetochore marker NDC80, spindle marker ARK2 and

cytoplasmic axonemal protein kinesin-8B. We generated dual fluorescent parasite lines

expressing PbARK1-GFP (green) and either PbNDC80-mCherry (magenta), PbARK2-

mCherry (magenta) or Pbkinesin-8B-mCherry (magenta) by genetic cross as described

previously[4]. Live-cell imaging showed that both PbARK1-GFP and PbNDC80-mCherry were

located next to the Hoechst-stained DNA, partially overlapping at the spindle but not at the

spindle poles (Fig. 2f left panel, Extended Data Fig. 2g, and Supplementary Video 4). 3D-

structured illumination microscopy (SIM) at one-min post-activation showed a clear

association of PbNDC80-mCherry with PbARK1-GFP at the spindle but not at the spindle

poles (Fig. 2f right panel). In parasite lines expressing PbARK1-GFP and PbARK2-mCherry,

these proteins were co-located at the spindle (Fig. 2g left panel, Extended Data Fig. 2h, and

Supplementary Video 5). Further 3D-SIM at one-min post-activation showed an association of

PbARK1-GFP with PbARK2-mCherry at the spindle, with PbARK1-GFP more outside at the

spindle poles (Fig. 2g right panel). However, in a parasite line expressing PbARK1-GFP and

Pbkinesin-8B-mCherry (a cytoplasmic marker) there was no overlap between these two

proteins (Fig. 2h left panel and Extended Data Fig. 2i), an observation which was further

substantiated by 3D-SIM imaging (Fig. 2h right panel).

During meiotic stages of development post-fertilisation, PbARK1-GFP was diffuse in the

cytoplasm, with transient nuclear foci forming in zygotes and disappearing in mature ookinetes

(Extended data Fig. 3a and Supplementary Notes). A diffuse distribution was also observed

throughout oocyst and sporozoite stages, including those from salivary glands (Extended Data

Fig. 3b and Supplementary Notes). In liver stages (in HepG2-infected cells), the PbARK1-GFP

signal remained diffuse even at 48 hours post infection (h.p.i), with slight perinuclear

enrichment at 64 h.p.i (Extended Data Fig. 3c and Supplementary Notes).

PfARK1 regulates spindle formation and karyokinesis

The functional role of PfARK1 in asexual blood stage development was assessed by examining

the effect of deleting the gene. Rapamycin (RAP) treatment of HA-PfARK1-loxP parasites

efficiently excised the floxed sequence (Extended Data Fig. 1c) and caused depletion of

PfARK1 protein (Extended Data Fig. 1d). Synchronized ring stage cultures of 1G5DC-

SkipFlox (control) (Extended Data Fig. 4a) or HA-PfARK1-loxP parasite lines were treated

with RAP or DMSO (control) and parasite growth was monitored every 24 h. In cycle 0 (RAP-

treatment) there was little effect on the parasitaemia of HA-PfARK1-loxP parasites. However,

a significant growth defect was observed at the beginning of the next cycle (cycle 1) as

indicated by a marked reduction in parasitaemia (Fig. 3a and Extended Data Fig. 4b), whilst

there was almost no effect of RAP treatment on 1G5DC-SkipFlox (Extended Data Fig. 4a).

This growth defect was further accentuated in subsequent cycles. Microscopic examination of

Giemsa-stained blood smears suggested normal ring to trophozoite development in cycle 1

DMSO-treated parasites (Extended Data Fig. 4c). However, following RAP treatment and

PfARK1 depletion, a significant number of parasites were pyknotic (at ~96 hpi) and fewer rings

were found in cycle 2 (Fig. 3b). Schizont development following treatment with E64 to block

late schizont maturation and merozoite egress was analysed: a significant number PfARK1-

depleted cycle 1 schizonts had abnormal morphology, with unsegregated nuclei (Fig. 3c and

Extended Data Fig. 4d), which is consistent with the role of PfARK1 in nuclear division. This

suggestion, that the decrease in parasitaemia resulting from PfARK1 depletion was due to its

role in nuclear division and/or schizont maturation, was investigated further.

To begin to understand how PfARK1 regulates asexual cell division, MTOC duplication and

spindle formation were examined. U-ExM revealed the formation of various forms of spindles

as described earlier with one or two centrin-labelled foci on the cytoplasmic side of the nuclear

envelope (Fig. 3d) and associated with NHS-ester labelled MTOCs or associated with mitotic

spindles[21,22]. In PfARK1-depleted parasites there were fewer spindles indicated by either

absent or speckled tubulin staining (Fig. 3d and Extended figure 4e, grey arrow). As mentioned

above, the nuclei appeared unsegregated with associated multiple centrin foci, and centrin

aggregates were seen away from the nuclei and disorganized (Fig. 3d and Extended figure 4e,

square). Collectively, these data suggested that while the MTOC duplication was unaltered, it

was disorganized and spindle formation and karyokinesis were impaired in PfARK1-depleted

parasites.

PfARK1 depletion impairs segmentation of daughter merozoites

At the end of schizogony, segmentation of the schizont by invagination of the parasite plasma

membrane leads to formation of merozoites prior to their egress. IFAs were performed using

antibody to MSP1, a parasite surface marker, to assess segmentation following ARK1

depletion. MSP1 fluorescence with a bunch of grape-like contours, characterized the membrane

of control parasites, but in contrast, PfARK1-depleted parasites either lacked MSP1 staining,

or contained large aggregations that encompassed multiple merozoites or unsegregated nuclei

(Fig. 3e, cyan arrow).

U-ExM of mature (segmented) schizonts with anti-GAP45 antibody-labelling revealed inner

membrane complex (IMC) and tubulin labelled subpellicular MT, and rhoptries labeled with

NHS-ester (Fig. 3f). However, for the PfARK1-depleted parasites, several unsegregated nuclei

were present and these cells either lacked GAP45 staining, or the parasite pellicle (the plasma

membrane/IMC) was disorganized (Fig. 3f, cyan arrow). NHS-ester labelled rhoptries were

found with some merozoites, typically the ones with GAP45 staining. In some parasites

rhoptries were either not formed properly or were absent, which may be due to arrested division

of these parasites prior to rhoptry formation. In addition, subpellicular MT were either not

observed in the cell or were mislocalised. These observations suggest impaired cytokinesis of

daughter merozoites upon PfARK1 depletion.

To study the effect of PfARK1 depletion on merozoite release and egress, time-lapse

microscopy was performed[23]. For this purpose, parasites were treated with protein kinase G

(PKG)-inhibitor ML-10 to allow schizont maturation but prevent egress, then its removal

allowed egress, which was captured by time-lapse microscopy[24,25]. Egress and dispersal of

merozoites for control parasites was observed (Fig. 3g and Supplementary Video 6), and while

egress from the erythrocyte initially appeared almost normal in PfARK1-depleted parasites,

the merozoites either remained attached to the residual body or clumped together (Fig. 3g,

magenta arrows, and Supplementary Videos 7, 8) even after prolonged duration. These defects

likely result from the abnormal nuclear segregation and cytokinesis in PfARK1-depleted

parasites.

PbARK1 knockdown reveals an essential role during male gametogony and parasite transmission

Previous analysis by gene disruption had shown that PbARK1 is probably essential for asexual blood-stage development[14]. To be able to manipulate its expression in sexual stages we used a promoter trap strategy, replacing the *ark1* promoter with the promoter from *ama1,* resulting in $P_{ama1}ark1$ hereafter referred as (*ark1PTD*), a gene which is not transcribed in gametocytes but is highly expressed in asexual blood stages (Extended Data Fig. 5a)[26]. Correct genetic integration in the transgenic parasite line was confirmed by PCR (Extended Data Fig. 5b), and *ark1* transcription was shown to be downregulated in *ark1PTD* gametocytes by qRT-PCR (Fig. 4a). A phenotypic analysis of these *ark1PTD* parasites was then performed at different stages of parasite development within the mosquito.

Phenotypic analysis revealed that male gametogony (as scored by exflagellation centres) was severely affected in the *ark1PTD* line compared to GFP control (wild-type-GFP; WT-GFP) parasite (Fig. 4b) [27], and therefore significantly fewer ookinetes (differentiated zygotes after fertilisation) were formed (Fig. 4c). In mosquitoes fed with *ark1PTD* parasites, we observed significantly fewer oocysts in the midgut at day 7 post infection, in contrast to the hundreds of oocysts in mosquitoes fed WT-GFP parasites (Fig. 4d). The number of *ark1PTD* oocysts was further reduced by day 14 and day 21 post infection compared to WT-GFP oocysts (Fig. 4d). The *ark1PTD* oocysts were also significantly smaller than the WT-GFP oocysts 7 days post infection and diminished further by day 21 post infection (Fig. 4e, f). The number of salivary glands sporozoites was significantly reduced in *ark1PTD* parasites on day 21 (Extended Data Fig. 5c). To further confirm the absence of sporogony, mosquitoes that had been fed with *ark1PTD* or WT-GFP parasites were allowed to feed on susceptible naïve mice. Mosquitoes infected with WT-GFP parasites successfully transmitted the parasite, with blood-stage infection detected in naïve mice after 4 days. In contrast, *ark1PTD* parasites failed to transmit in most experiments (five out of eight) (Extended Data Fig. 5d). However, on three occasions, they were able to establish infection in bite-back experiments, although the prepatent period was markedly prolonged, with blood-stage parasites first detected 14 days post-infection (Extended Data Fig. 5d).

To investigate the effect of *Pbark1* downregulation at the genome wide level, we performed RNA-seq analysis in triplicate, starting with total RNA extracted from purified *ark1PTD* and WT-GFP gametocytes at two time points: before activation (non-activated, 0 min) and at 30

min post-activation (mpa). Differences in gene expression between the correspondent samples were detected. In non-activated gametocytes, only a small number of differentially expressed genes were identified in the *ark1PTD* sample, with 10 genes up- and 8 genes downregulated, respectively (Extended Data Fig. 5e and Supplementary Table 1). Those genes included several fam a and b proteins known to be highly variable during life cycle progression due to their involvement in host immune response modulation[28]. They were therefore not considered as significant. At 30 mpa, 23 genes were upregulated including several conserved *Plasmodium* proteins with unknown function as well as PBANKA_1206900, a tubulin beta chain and putatively critical during gametocyte formation[29]. A significantly higher number of genes (229) were downregulated (Extended Data Fig. 5f and Supplementary Table 1). These genes included several protein kinases, e.g. PBANKA_1228800 and calcium-dependent protein kinase 5 (CDPK5 – PBANKA_1351500), as well as genes involved in lipid biosynthesis including PBANKA_0201300, a putative lysophospholipase, PBANKA_1127000, and a choline/ethanolaminephosphotransferase. Also, down-regulated were myosin essential light chain (ELC) and skeleton-binding protein 1 (PBANKA_1101300), as well as several enzymes known to be critical for sexual differentiation. Gene ontology (GO) analysis also revealed that several genes detected as downregulated were associated with ribosome biogenesis (Extended Data Fig. 5g). These data indicate not only a potential cell cycle arrest but also most likely stress sensing and translational inhibition as a result of PbARK1 depletion.

ark1PTD parasites have defective spindle formation, MTOC separation, and axoneme formation during male gametogony

To further investigate the function of PbARK1 during male gametogony, WT-GFP and *ark1PTD* male gametocytes were observed during gametogony using U-ExM. In WT-GFP male gametocytes at 8 mpa, individual MTOCs were distinct and separated, whereas in *ark1PTD* male gametocytes multiple MTOCs were clumped and aggregated together (Fig. 4g and Extended Data Fig. 6a). At 15 mpa, WT-GFP male gametocytes were observed undergoing exflagellation, whereas in *ark1PTD* male gametocytes the MTOCs were still clumped together and no exflagellation was observed (Fig. 4g and Extended Data Fig. 6a).

We examined whether the lack of MTOC separation in the *ark1PTD* line led us to examine whether this was caused by a defect in spindle formation. At 8 mpa, long spindles with clearly separated MTOCs were present in WT-GFP parasites, but in *ark1PTD* lines, short spindles,

with the two MTOCs remaining close together (Fig. 5a, b) were present. Measurement showed that *ark1PTD* spindles were significantly shorter (~0.8 μm) than WT-GFP spindles (~2.5 μm; Fig. 5c). These findings corroborated studies on PfARK1 during schizogony as depletion of the kinase resulted in abnormal spindle formation and accumulation of MTOCs around unsegregated nuclei (Fig. 3d).

TEM observations further revealed structural defects in *ark1PTD* male gametocytes. At 8 mpa, in the *ark1PTD* line, MTOCs formed a normal bipartite structure consisting of an inner spindle MTOC and an outer basal body MTOC, similar to those in WT-GFP male gametocytes, and normal axoneme formation was observed (Fig. 5dA-D, I-L and Extended Data Fig. 6bA-D), However, we also observed MTOCs that appeared detached from the nucleus in the *ark1 PTD* line (Fig. 5dI, J and Extended Data Fig. 6bA, B, D). As revealed by expansion microscopy, multiple basal body MTOCs were often aggregated and clumped together (Fig. 5dL). Within the cytoplasm, we observed singlet or doublet MT that may have been axoneme precursors and likely represent failures in axoneme formation (Fig. 5dK, L).

At 15 mpa, WT-GFP gametocytes had started to form gametes with flagellum and nucleus (Fig. 5dE, F, G, H), while in the *ark1PTD* line there was incomplete axoneme formation and exflagellation rarely occurred (Fig. 5dM-P and Extended Data Fig. 6bE-H). In addition, two basal body MTOCs remained close together without proper separation (Fig. 5dM, N, P), with short spindle MT visible between them (Fig. 5dP and Extended Data Fig. 6c).

PbARK1 is part of a divergent chromosome passenger complex (CPC) with two INCENP proteins

To explain the ARK1 phenotypes and confirm the logic of the observed subcellular localisations, we wanted to understand what protein scaffolds or modulator proteins might direct ARK1 activity and localisation during both asexual and sexual stages of the parasite. We therefore prepared lysates of *P. berghei* schizonts and male gametocytes from PbARK1-GFP and WT-GFP control parasites, immunoprecipated the protein with GFP-Trap and then identified bound proteins using mass spectrometry. At both stages we identified two divergent inner centromere proteins (INCENP1/2) as major ARK1 interactors (Fig. 6a and Extended Data Fig. 7). Although the differences were not significant (n = 2 for each sample), in the male gametocyte sample we observed co-variation of various spindle and kinetochore proteins

amongst the pulldowns (Supplementary Table 2): for example, the putative ARK2-based spindle complex including MISFIT-MyoK-EB1, several kinetochore subunits e.g. Apicomplexan Kinetochore proteins (AKiTs) 1, 3, 5 and 7 and the condensin-II-specific subunits CapH2-G2-D3. Strikingly, we saw no survivin/borealin-like proteins in the interactome, confirming earlier predictions that *Plasmodium* spp., and more generally apicomplexans, lack other conserved components of the CPC (Fig. 6c). Since the spatiotemporal dynamics of ARK1 suggest that it may not act as the classic passenger complex on chromosomes, it is possible that the chromatin-directing function of Survivin and Borealin has become obsolete for apicomplexans and that the structural motifs for localisation and activation of ARK1 by INCENP1 and 2 might be revealed upon detailed comparative sequence analyses.

T. gondii, another apicomplexan, sister to *Plasmodium* spp. also has two INCENP paralogs that interact with TgARK1, so we sought to reconstruct the evolutionary history of the INCENP proteins with a focus on alveolates and apicomplexans (Fig. 6b, c). The duplication that gave rise to INCENP1 and INCENP2 can be traced back to the common ancestor of all Apicomplexa (sensu stricto). We found parallel but separate duplications in other different alveolate lineages such as apicomplexpan-like Squirmida, Chrompodellida and dinoflagellate-related lineages concomitant with the loss of borealin/survivin (Fig. 6c), suggesting INCENP duplications might compensate for loss of these chromatin-directing subunits. Interestingly, INCENP2 is absent from Piroplasmida, lineages which have also lost ARK2 and ARK3, suggesting that INCENP2 might also play a role in ARK2/3-based regulatory pathways. A co-fold model of *P. berghei* INCENP1/2 and ARK1 using AlphaFold3, suggests that the conserved IN-box region involved in kinase interaction is flanked in all apicomplexan proteins by N- and C-terminal extensions that form a beta-sheet – folding INCENP1/2 around ARK1 (Fig. 6b and Extended Data Fig. 8). In addition, the C-terminal extension has a conserved CxC motif, possibly involved in posttranslational modification (Fig. 6b). We could only predict a coiled-coil in the N-terminus of INCENP1, and no further structural features could be discerned for INCENP2 using AlphaFold3. Such absence of any feature makes it hard to predict the binding mode of INCENP2 either as a single ARK1-INCENP2 interaction or forming an INCENP2-ARK1-INCENP1 trimeric complex.

Finally, we sought whether any non-canonical ARK1/INCENP features might indicate alternative CPC functions in *Plasmodium* spp., such as the highly conserved

autophosphorylation sites in both Aurora B and INCENP, which are crucial for the kinase

activity of the CPC in vertebrates[30]. We found that both the broadly conserved

autophosphorylation site in the activation loop of Aurora kinases (RxTxCGT) and the

activating phosphosite on INCENP (RTSS in human) are markedly different (GxSH in ARK1)

and (RTSS absent from INCENP1/2). However, some Aurora consensus sites can be found in

the vicinity among INCENP1/2 paralogs (Fig. 6b – green boxes). ARK2 paralogs have a

conserved Aurora activation loop, while ARK3 has a serine-rich activation loop with multiple

candidate activating phosphorylation sites (Fig. 6c). The absence of clear conserved auto/trans-

phosphorylation sites between ARK1 and INCENP1/2 in *Plasmodium* spp. indicate CPC kinase

activity might be regulated in a different fashion, possibly by input of another kinase.

Discussion

Our cross-stage analysis identified ARK1 as a central organiser of *Plasmodium* mitosis using

P. falciparum and *P. berghei* model systems. First, ARK1 localises to the spindle and inner

MTOC specifically during mitosis, and its depletion in *P. falciparum* reduces spindle formation,

prevents proper karyokinesis, disrupts segmentation and impedes egress. Second, during the

rapid mitoses of male gametogony, *P. berghei* ARK1 dynamics follow spindle poles;

knockdown shortens spindles, block MTOC separation and axoneme biogenesis, and abolishes

transmission. These phenotypes position ARK1 upstream of both chromosome segregation and

gamete formation.

Our findings show that ARK1 is preferentially expressed and targeted during the mitotic stages

marked by spindle formation. Live cell imaging and expansion microscopy revealed that ARK1

locates to the spindle apparatus and the spindle poles, corresponding to the inner core of the

bipartite MTOC[4,21]. Expansion microscopy using centrosome (centrin) and microtubule

(tubulin) markers confirmed that ARK1 associates with spindle poles within the inner MTOC

in both mitotic stages. Some of these features had been observed earlier for *P. falciparum* ARK1

and further it had been shown that its location is limited to mitotically dividing stages[17]. Dual

fluorescence microscopy analysis during male gametogony revealed that it colocalised with

ARK2, another aurora kinase reported recently, and the kinetochore marker NDC80 at the plus

ends of spindle microtubules, suggesting a cooperative role for ARK1 and ARK2 at equatorial

spindles. It also has a role in kinetochore dynamics, as observed previously for ARK2[10].

However, live cell imaging in *P. berghei* clearly showed that it is part of the inner MTOC as it

does not colocalise with the basal body and axoneme marker, kinesin-8B. This contrasts with

T. gondii ARK1, which, when tagged at the N-terminus, exhibited a dynamic localisation to

both nucleus and cytoplasm during division, but remained predominantly nuclear in non-

dividing cells[18,31]. Its colocalisation with kinetochore NDC80 and ARK2 markers further

supports a dual role in spindle organization, kinetochore segregation, and spindle pole

integrity—reminiscent of Aurora B, which localises to centromeres, the central spindle, and the

spindle midzone[5,6,32,33].

Functional genetic studies in *P. falciparum* and *P. berghei* had indicated that ARK1 is likely

essential in asexual blood stages[13,14]. Attempts to generate conditional knockdowns using the

DD/Shield system in *P. falciparum* were unsuccessful, as reported recently[17]. Here using the

DiCre-Rapamycin system for *P. falciparum*[19] and promoter trap strategy for *P. berghei*[20], we

demonstrate that ARK1 depletion leads to defects in spindle formation and karyokinesis. In *P.*

falciparum, ARK1 is required for proper spindle elongation, parasite growth, cytokinesis and

egress. In *Toxoplasma*, overexpression of a kinase dead mutant of TgARK1 disrupts

chromosome segregation and spindle pole body maturation, similar to some of the division

defects observed here in asexual stages, although the mechanism was unexplored[18,31].

During male gametogony, ARK1 knockdown led to defects in the spindle apparatus and failure

of MTOC separation, suggesting its associated function with the outer basal body[4]. This was

accompanied by abnormal axoneme formation and very few flagellated male gametes,

indicating a failure in MTOC-basal body segregation. These results suggest that while ARK1

is part of the inner MTOC, its functional association with the outer MTOC is essential for

successful mitosis and gamete formation, similar to the egress and spindle elongation defects

seen in *P. falciparum*. These features also show some resemblance to those of Aurora B in many

model systems[5,6,32,33].

Our protein interactome analysis reveals that ARK1 is part of a non-canonical chromosomal

passenger complex (CPC), comprising two divergent INCENP homologs and lacking the

canonical chromatin-targeting components Borealin and Survivin. This is consistent with

earlier molecular cell biological studies in *Toxoplasma*[18]. While studies in *Toxoplasma* also

reported two INCENPs, they did not detect the presence or absence of other CPC components.

The lack of chromatin-targeting components might suggest that INCENP1/2 can facilitate such

interactions themselves or simply mean these are lost and replaced by interactions of different

proteins. For instance, INCENPs in apicomplexans may form alternative CPC complexes,

similar to *Trypanosoma*, where INCENP associates with kinesin-1A/B[34]. A potential candidate

kinesin might be kinesin-8X, which shows a similar localisation pattern during different life

stages as ARK1[35]. Interestingly, plants and heterokont algae also harbour divergent Survivin-

like proteins, with a central conserved but highly divergent helix suggesting a possible broader

evolutionary divergence of CPC components across eukaryotes[36]. Future studies will need to

include a further scrutiny of INCENP1/2 localisation and interactomes to determine whether

any new CPC components might be present in apicomplexans.

Comparative genomics provides an evolutionary scenario for this atypical CPC. INCENP

duplication into INCENP1 and INCENP2 likely arose in the apicomplexan ancestor,

concomitant with the loss of Survivin and Borealin (Fig. 6). Notably, INCENP2 has been

secondarily lost in piroplasmids, which also lack ARK2 and ARK3, hinting at functional

coupling between Aurora paralogs and specific INCENP variants. Structural predictions using

AlphaFold3 suggest that apicomplexan INCENPs contain extended IN-box regions that fold

around the ARK1 kinase domain and feature a conserved CxC motif absent from other

eukaryotic homologs. At the same time, ARK1 itself shows a divergent activation loop (GxSH

instead of the canonical RxT), and INCENPs lack the canonical transphosphorylation site

(TSS) site found in other eukaryotes. Such changes might indicate that other kinases are

necessary to activate ARK1. Together, these features indicate that apicomplexans have likely

rewired CPC architecture and regulation, replacing canonical chromatin-targeting logic with

INCENP-centric scaffolds that localise ARK1 at the inner MTOC, spindle and nucleoplasm.

While the function of the other two ARKs-ARK2 and ARK3- is unclear, at least in asexual

stages, our recent studies on PbARK2 have provided insights into its role in sexual transmission

stages. While ARK2 regulates spindle dynamics during male gametogony just like ARK1 in

the present studies, surprisingly, its deletion did not impact exflagellation, unlike that of the

ARK1 mutant (Fig. 4b). Previous studies also revealed that PbARK2 deletion caused an

increase in the expression of ARK1 and ARK3[10]. Therefore, in the light of the present work, it

is reasonable to suggest that ARK1 and ARK2 may have some overlap of function during male

gametogony, particularly in spindle biogenesis. It was also interesting to note that the deletion

of PfARK1 resulted in higher PfARK2 levels in schizonts (Extended Data Fig. 1f), although

this increase was insufficient to complement the PfARK1 mutant. While there may be some
overlap in ARK1 and ARK2 function, each is indispensable for parasite survival.

Overall, this study suggests that *Plasmodium* ARK1 regulates spindle elongation during mitotic
division and associates at the inner nuclear pole of the MTOC. However, its association with
the outer MTOC is essential for karyokinesis and for flagellated gamete formation. ARK1
appears to represent a unique Aurora kinase paralog that, while divergent in structure, shares
functional similarities with Aurora B in humans. This suggests that *Plasmodium* has evolved
an atypical mechanism of cell division, where many of the canonical molecules are either
missing or highly divergent, reflecting the parasite's adaptation to complex life cycle
transitions.

Methods

***P. falciparum* cultures**

Plasmodium falciparum strain 1G5DC was obtained from European Malaria Reagent
Repository (EMRR) and DSM1 was obtained from *BEI* resources, Malaria Research and
Reference Reagents Resource Centre (MR4), American Type Culture Collection (ATCC).
Parasites were maintained in O+ human erythrocytes (5% hematocrit) in RPMI-1640
supplemented with 0.5% Albumax II and 50 μg/mL hypoxanthine[37]. Cultures were maintained
at 37°C under a mixture of 5% CO_2, 3% O_2, and 91.8% N_2 or 5% CO_2. Fresh erythrocytes and
culture medium were used to dilute parasites to maintain 3–5% parasitaemia and 5%
haematocrit. Relevant drugs were used during the culture of various transgenic lines, as
described below. Parasite synchronization was carried out using 5% sorbitol as described
previously[38].

Generation of transgenic parasite lines

All PCR primers used in this study were synthesized by Sigma and are indicated in
Supplementary Table 3.

1G5DC SkipFlox

A dimerisable Cre recombinase (diCre) expressing parasite line was generated by transfecting
pSkipFlox plasmid, a kind gift from Tobias Spielmann (Addgene Plasmid #85797) into the
parent line 1G5DC[19]. Parasites were selected in the presence of 2.5 μg/ml of blasticidin.

pSLI-HA-ARK1- loxP

To generate a conditional knockout transgenic parasite, a selection-linked integration (SLI) approach was used. To generate the construct, a 592 bp homology region corresponding to the 5'-end of the PfARK1 gene was amplified using primers (P1 and P2) derived from 3D7 genomic DNA. The PCR product was cloned into *NotI* and *PmeI* sites upstream of loxP and a Myc-tag coding sequence in the pSLI-N-sandwich-loxP (K13) vector, which was a gift from Tobias Spielmann (Addgene Plasmid #85793). A recodonised sequence version of full-length PfARK1 with a N-terminal 3xHA tag and followed by a loxP site was custom synthesised as a G-block (Genscript). The G-block was cloned in the above construct downstream of yDHODH using *NheI* and *XhoI* restriction sites[19].

P. falciparum **transfection**

The pSLI-HA-PfARK1-loxP construct was transfected in 1G5DC SkipFlox parasites at the ring stage using ~100μg of purified plasmid DNA. Transgenic parasites were selected by using 2nM WR99210 and integrants were then enriched using 1.5μM DSM-1. The parasites that appeared after drug selection were genotyped using PCR primers (Supplementary Table 3 and Extended Data Fig. 1a). The desired integration was confirmed by sequencing of PCR products.

Conditional knockout of PfARK1

HA-PfARK1-loxP parasites were treated with 250 nM rapamycin at ring stage for 24 hours. The genomic DNA was isolated from DMSO/RAP-treated parasites and excision of the PfARK1 locus flanked by loxP sites in the transgenic parasite was confirmed by PCR using primers (P7/P8).

P. falciparum **growth rate assays**

HA-PfARK1-loxP parasites were cultured in the presence of 10 nM WR99210, 2.5 μg/ml blasticidin and 1.5μM DSM-1. Cultures were maintained at 2% hematocrit and 7-8% parasitaemia. To perform the growth rate assay, 1G5DC-SkipFlox and HA-PfARK1-LoxP parasites were synchronised at ring stage for two cycles using 5% sorbitol for 20 min at 37°C, and the rings were seeded at 0.5% parasitaemia and 2% hematocrit. To deplete ARK1, parasites were treated with either DMSO (vehicle control) or 250 nM rapamycin for 24 h. Thin Giemsa-stained blood smears were examined periodically under a microscope to quantify the parasitaemia at different parasite stages. For assessment of parasitaemia by flow cytometry, cells were fixed by incubation in a solution containing 1% paraformaldehyde and 0.0075%

glutaraldehyde at ambient temperature for 15 min on an end-to-end rocker. Post-fixation, samples were either stored at 4°C or stained immediately with Hoechst 33342 dye at 37°C for 10 min. After staining, samples were washed with FACS buffer and then analysed on either BDverse (BD Biosciences) or BD FACSymphony A1 Flow Cytometers, acquiring between 10,000 and 100,000 events per sample. Data analysis was carried out using FlowJo software version 10.10.0.

Ethics statement (Studies on *P. berghei*)

All animal procedures were approved following an ethical review process and conducted in accordance with the United Kingdom Animals (Scientific Procedures) Act 1986, under Home Office Project Licenses (PDD2D5182 and PP3589958). Female CD1 outbred mice (6 to 8 weeks old) were obtained from Charles River Laboratories and used for all experiments carried out in the UK.

Purification of *P. berghei* schizonts and gametocytes

Blood-stage parasites were obtained from infected mice on day 4 post-infection and cultured at 37 °C with gentle rotation (100 rpm) for 11 or 24 hours. Schizonts were purified the following day using a 60% v/v NycoDenz gradient prepared in phosphate-buffered saline (PBS). The NycoDenz stock solution consisted of 27.6% w/v NycoDenz in 5 mM Tris-HCl (pH 7.2), 3 mM KCl, and 0.3 mM EDTA.

Gametocyte purification was performed by first infecting phenylhydrazine-treated mice[39], followed by sulfadiazine treatment two days post-infection to enrich for gametocytes. Blood was collected on day 4 post-infection, and gametocyte-infected cells were purified using a 48% v/v NycoDenz gradient in PBS, prepared from the same NycoDenz stock solution. Gametocytes were collected from the interface and subsequently activated for downstream assays[40].

Generation of *P. berghei* transgenic parasites

GFP-tagging constructs were generated using the p277 plasmid vector and transfected into parasites as previously described[41]. A schematic representation of the wild-type *ark1* locus, the GFP-tagging constructs, and the recombined *ark1* locus is shown in Extended Data Fig. 2a. Correct integration of the *gfp* sequence at the *ark1* locus was confirmed by diagnostic PCR using primer 1 (intArk1tg) and primer 2 (ol492), as indicated in Extended Data Fig. 2b. The

expression of native ARK1-GFP was confirmed by Western blot analysis (Extended Data Fig. 2c).

To investigate the function of ARK1, a promoter trap strategy was employed using double homologous recombination to generate a conditional knockdown line (*ark1PTD*). The knockdown construct was derived from the P*ama1* plasmid (pSS368), in which *ark1* was placed under the control of the *ama1* promoter, as previously described[20]. A schematic representation of the endogenous *ark1* locus, the targeting constructs, and the recombined *ark1* locus is shown in Extended Data Fig. 5a. Diagnostic PCR was performed to confirm successful integration, as outlined in Extended Data Fig. 5a. Integration at the 5′ locus was verified using primers intPTD31_5 (Primer 1) and 5′-intPTD (Primer 2), while integration at the 3′ end was confirmed using primers intPTD31_3 (Primer 3) and 3′-intPTama1 (Primer 4) (Extended Data Fig. 5b). Primer sequences are listed in Supplementary Table 3. Transfections were carried out by electroporation using *P. berghei* ANKA line 2.34 for GFP-tagging experiments and ANKA line 507cl1 expressing GFP for generation of the knockdown line[27].

Generation of dual tagged parasite lines (*P. berghei*)

To generate dual-labelled parasites, ARK1-GFP parasites were mixed in equal proportions with either NDC80-mCherry, ARK2-mCherry, or kinesin-8B-mCherry labelled parasites and co-injected into mice. Four to five days post-infection, when gametocytaemia was high, Anopheles mosquitoes were allowed to feed on the infected mice. Mosquitoes were examined for oocyst development and sporozoite formation at days 14 and 21 post-feeding. Infected mosquitoes were then used to infect naïve mice via mosquito bite-back. Blood-stage infection was assessed 4–5 days later by Giemsa-stained blood smear microscopy. In this way, parasites expressing both ARK1-GFP and either NDC80-mCherry, ARK2-mCherry, or kinesin-8B-mCherry were obtained. Gametocytes from these mixed lines were purified, and fluorescence microscopy was performed as described below to examine protein co-localisation.

P. berghei **phenotype analyses**

Infections were initiated by intraperitoneal injection of approximately 50,000 *P. berghei* parasites from WT-GFP or *ark1PTD* lines into mice. Asexual stages and gametocyte production were monitored on Giemsa-stained thin smears. Four to five days post-infection, exflagellation and ookinete conversion were assessed using a Zeiss AxioImager M2 microscope with an AxioCam ICc1 digital camera. For mosquito transmission assays, 30–50 *Anopheles stephensi* (SD 500) mosquitoes were allowed to feed for 20 min on anaesthetised, infected mice with

~15% asexual parasitaemia and comparable gametocyte levels (as determined by Giemsa-

stained smears). To evaluate midgut infection, ~15 mosquito guts were dissected on day 7, 14

and 21 post-feeding, and oocysts were counted using a 63× oil immersion objective on the

same microscope setup. Sporozoites in oocysts were counted on day 14 and day 21 post

feeding. Salivary glands were isolated on day 21 to count the sporozoites. Bite-back

experiments were performed 21 days post-feeding using naïve mice, and blood smears were

examined after 3–4 days. All experiments were conducted at least three times to assess the

phenotype.

Immunofluorescence Assay (IFAs)

P. falciparum - Immunofluorescence assays (IFA) were conducted on thin blood smears

following a previously described protocol[42]. In brief, air-dried smears were fixed for 2 min

using ice cold 1:1 mixture of methanol and acetone. The smears were dried to remove any

traces of methanol/acetone followed by blocking with 3% BSA for 45 minutes at room

temperature. Samples were incubated with primary antibodies for 12 h at 4°C. After thorough

washing with 1X PBS, Alexa Fluor-conjugated secondary antibodies were added and incubated

for 2 h at room temperature. Smears were then mounted with VectaShield medium (Vector

Laboratories Inc.) containing DAPI for nuclear staining. Microscopy was performed using a

LSM980 AR confocal microscope (Carl Zeiss) or Zeiss Axio Observer microscope and image

processing was carried out using Zeiss ZEN Blue (versions 3.1 or 3.9). Representative z-stack

images were selected for figure preparation unless indicated otherwise. IFA or U-ExM

(described below) with MSP1 or GAP45 antibodies was performed to assess segmentation in

schizonts that had been treated with 10 μM E64 at ~42 h.p.i (cycle 1) for 4-5 hours to prevent

egress and allow complete maturation.

P. berghei - Purified gametocytes were fixed in 4% paraformaldehyde in microtubule-

stabilising buffer on poly-L-lysine–coated slides (Sigma), permeabilised with 0.1% Triton X-

100 in PBS, and blocked with 10% goat serum and 3% BSA in PBS. Primary antibodies—

rabbit anti-GFP (1:250, Thermo Fisher) and mouse anti–α-tubulin (1:1000, Sigma)—were

diluted in 1% BSA and 10% goat serum in PBS and incubated for 1 h. Alexa Fluor 488 anti-

rabbit and Alexa Fluor 568 anti-mouse secondary antibodies (1:1000, Invitrogen) were applied

for 45 min in the dark. Slides were washed with PBS between steps, mounted with Vectashield

containing DAPI (Vector Labs), and sealed. Images were acquired using a Zeiss Axio Imager

M2 microscope with a 63× oil immersion objective and AxioCam ICc1 camera. Fluorescence

was adjusted using Axiovision (Rel. 4.8) software. Experiments were repeated at least three times, and 20–30 cells per stage were analysed. See figure legends for additional details.

P. falciparum live-cell imaging

Live imaging was carried out with minor modifications to previously described protocols[25]. HA-PfARK1-loxP parasites were synchronized at the ring stage and treated with either DMSO or 250 nM rapamycin for 24 hours. Schizonts (cycle 1) were subsequently purified and co-cultured with fresh erythrocytes at 2% hematocrit in RPMI1640 complete medium in the presence of 25 nM PKG inhibitor ML10 [23,24]. ML10 was removed after 4-6 hours and parasites were plated onto μ-slide, 2-well ibiTreat chambers (ibidi, 80286) pre-coated with 0.5 mg/mL concanavalin A. The slides were maintained at 37 °C in a humidified chamber with 5% CO_2. Imaging was performed using a Zeiss Axio Observer microscope, capturing frames every 3 seconds for a minimum duration of 30 min and the acquired time-lapse videos were processed and analyzed using Axio Vision software (version 4.8.2) or ZEN blue 3.1 or 3.9.

P. berghei live cell imaging

To examine PbARK1-GFP expression during erythrocytic development, parasites cultured in schizont medium were imaged at various stages, including ring, trophozoite, schizont, and merozoite. Purified gametocytes were assessed for GFP expression and localisation at multiple time points (0 and 1–15 min) following activation in ookinete medium. For nuclear staining, Hoechst dye was used during live-cell imaging. Zygote and ookinete stages were analysed over a 24-hour culture period. Images were acquired using a Zeiss Axio Imager M2 microscope equipped with a 63× oil immersion objective and an AxioCam ICc1 digital camera (Carl Zeiss) using autoexposure settings. Fluorescence signals were adjusted using Axiovision (Rel. 4.8) software to reduce background fluorescence while maintaining detection sensitivity; representative images are shown in the figures. All imaging experiments were performed in at least three independent replicates, and 30–50 cells were analysed per stage to determine PbARK1-GFP localisation. For additional details, refer to the figure legends.

P. berghei structured illumination microscopy

Formaldehyde-fixed (4%) gametocytes were stained with Hoechst dye, and 2 μl of the cell suspension was placed on a microscope slide and covered with a 50 × 34 mm coverslip to form a thin, immobilised cell monolayer. Imaging was performed using structured illumination microscopy (SIM) on a Zeiss Elyra PS.1 microscope with either a Zeiss Plan-Apochromat

667   63×/1.4 oil immersion or Zeiss C-Apochromat 63×/1.2 W Korr M27 water immersion

668   objective, as previously described[43].

Ultrastructure expansion microscopy

U-ExM was performed largely following previously published protocols[21,44-46]. In brief, to

prepare U-ExM samples of *P. falciparum* schizonts, 12mm coverslips were placed in a 24-well

plate and coated with poly-D-lysine (Gibco, A3890401, 0.1 mg/mL) for 1 h at 37 °C, followed

by three washes with ultrapure water. *P. falciparum* schizonts were seeded onto the coverslips

and incubated for 30 min at 37 °C. Unattached cells were removed gently, and the remaining

cells were fixed using 4% (v/v) formaldehyde in PBS for 15 min at 37 °C. After fixation,

coverslips were rinsed with prewarmed PBS and incubated for 12h at 37 °C in 1.4% (v/v)

formaldehyde and 2% (v/v) acrylamide in PBS. A monomer solution consisting of 19% (w/w)

sodium acrylate (Sigma-Aldrich, 408220), 10% (v/v) acrylamide (Sigma-Aldrich, A3553), and

2% (v/v) N,N′-methylenebisacrylamide (Sigma-Aldrich, M7279) in PBS was prepared 24 h in

advance and stored at −20 °C. For polymerization, 5 μL of 10% (v/v) TEMED (Sigma-Aldrich,

T9281) and 5 μL of 10% (w/v) APS (Sigma-Aldrich, A3678) were mixed with 90 μL of the

thawed monomer solution. The gel mix (35 μL) was pipetted onto parafilm squares pre-cooled

at −20 °C and the coverslip was placed over the drop. Gels were allowed to polymerize for 1 h

at 37 °C. Polymerized gels were transferred to denaturation buffer (200 mM SDS, 200 mM

NaCl, 50 mM Tris, pH 9) and shaken for 15 min at room temperature, then incubated at 95 °C

for 90 min. Denatured gels were expanded by sequential incubations in water (30 min each,

repeated thrice) with gentle shaking. Expanded gels were equilibrated in PBS for 15 min and

blocked in 3% BSA/PBS for 30 min at room temperature. Incubation with primary antibody

(details in Supplementary Table 4) was carried out for 12h at room temperature on a shaker,

followed by three 10 min washes with PBST. Gels were then incubated with secondary

antibodies, 405 NHS Ester (Thermo Fisher, A30000) and SYTOX/Hoechst in PBS for 2.5 h in

the dark with gentle shaking followed by three PBST washes and final equilibration in MilliQ

water (30 min, repeated thrice).

For imaging, μ-slide 2-well ibiTreat chambers (ibidi, 80286) were coated with poly-D-lysine

(0.5 mL, 1 h at 37 °C), rinsed with MilliQ water, and used to mount the gels cell side down.

Imaging was performed using a Zeiss LSM980 confocal microscope with a 63X oil immersion

objective (NA 1.4). Z-stacks (0.5 μm intervals) were captured and processed using ZEN Blue

(versions 3.1 or 3.9). Deconvolution was performed using Zen Deconvolution tool kit. U-ExM

images are displayed as maximum intensity projections of full z-slices unless indicated otherwise.

To prepare U-ExM samples of *P. berghei* male gametocytes, purified and 8 min- and 15 min-activated gametocytes were fixed in 4% formaldehyde in PHEM buffer (60 mM PIPES, 25 mM HEPES, 10 mM EGTA, 2 mM $MgCl_2$, pH6.9) at room temperature for 15 min. Fixed samples were attached to 10 mm round Poly-D-Lysine coated coverslips for 15 min. Coverslips were incubated overnight at 4°C in 1.4% formaldehyde (FA)/ 2% acrylamide (AA). Gelation was performed in ammonium persulfate (APS)/TEMED (10% each)/monomer solution (23% sodium acrylate; 10% AA; 0,1% BIS-AA in PBS) on ice for 5 min and at 37°C for 30 min. Gels were denatured for 15 min at 37°C and for 90 min at 95°C in denaturation buffer (200 mM SDS, 200 mM NaCl, 50 mM Tris, pH 9.0, in water). After denaturation, gels were incubated in distilled water overnight for complete expansion. The following day, circular gel pieces with a diameter of ~13 mm were excised, and the gels were washed in PBS three times for 15 min to remove excess water. The gels were then incubated in blocking buffer (3% BSA in PBS) at room temperature for 30 min, incubated with mouse monoclonal anti-a-tubulin antibody (T9026, Sigma-Aldrich, 1:500 dilution) and/or anti-GFP antibody (A11122, Invitrogen, 1:500 dilution) in blocking buffer (1:500 dilution) at 4°C overnight and washed three times for 15 min in wash buffer (0.5% v/v TWEEN-20 in PBS). The gels were incubated with 8 μg/ml Atto 594 NHS ester (08741, Merck), 10 μg/ml Hoechst 33342 (H1399, Invitrogen), Alexa Fluor 488 goat anti-mouse IgG (A11001, Invitrogen, 1:500 dilution) and Alexa Fluor 568 goat anti-mouse IgG (A11004, Invitrogen, 1:500 dilution) in PBS at 37°C for 3 hours followed by three washes of 15 min each in wash buffer (blocking and all antibody incubation steps were performed with gentle shaking). The gels were then washed three times for 15 min with wash buffer and expanded overnight in ultrapure water. The expanded gel was placed in a 35 mm glass bottom dish (MATTEK) with the 14 mm glass coated with Poly-D-Lysine and mounted with an 18 × 18 mm coverslip to prevent the gel from sliding and to avoid drifting while imaging. High-resolution confocal microscopy images were acquired using a Zeiss Celldiscover 7 with Airyscan using a Plan-Apochromat 50 ×/1.2NA Water objective, with 405, 488 and 561 nm lasers. Confocal z-stacks were acquired using line scanning and the following settings: 55 × 55 nm pixel size, 170 nm z-step, 2.91 μs/pixel dwell time, 850 gain and 3.5% (405 nm), 4.5% (488 nm) and 5.0% (561 nm) laser powers. The z-stack images were processed and analysed using Fiji (Version 1.54f) [47].

Calculation of the expansion factor for the U-ExM of male gametocytes

The expansion factor for the U-ExM of male gametocytes was calculated by comparing

axoneme diameters from two types of images (U-ExM and TEM). The diameters were

measured using Fiji (Version 1.54f) [47]. The average diameter of the axoneme in NHS-stained

U-ExM images of male gametocytes activated for 8 min was 933.83 nm (n = 12), while the

average diameter in TEM images of male gametocytes activated for 8 min was 188.50 nm (n

= 8). We used the calculated expansion factor of 4.954 for the analysis.

Electron microscopy of male gametocytes

P. berghei gametocytes activated for 8 min and 15 min were fixed in 3% glutaraldehyde in 0.1

M cacodylate buffer and processed for electron microscopy[48]. Briefly, samples were post-fixed

in 1% aqueous osmium tetroxide, treated en bloc with 2% aqueous uranyl acetate, dehydrated

in graded ethanol series and embedded in Spurr's epoxy resin. Thin sections were stained with

lead citrate prior to examination in a Tecnai G2 12 BioTwin (FEI UK, UK) or a Jeol JEM-

1400Flash (JEOL, Japan) electron microscope.

Liver stage parasite imaging

For *P. berghei* liver-stage imaging, 100,000 Hepatocelluar carcinoma cells (HepG2) were

seeded onto glass coverslips in 48 well plate. Salivary glands from *Anopheles stephensi*

mosquitoes infected with PbARK1-GFP parasites were dissected and homogenized using a

pestle to release sporozoites. The sporozoites were gently pipetted onto the HepG2 cell

monolayer and incubated at 37 °C with 5% CO_2 in complete minimum Eagle's medium The

culture medium was replaced 2 hours post-infection and subsequently changed daily. For live-

cell imaging, Hoechst 33342 (Molecular Probes) was added to a final concentration of 1 μg/ml

to stain host and parasite nuclei. Infected cells were imaged at, 48-, and 64 -hours post-infection

using a Zeiss Axio Imager M2 microscope equipped with a 63× oil immersion objective and

an AxioCam ICc1 digital camera (Carl Zeiss) using autoexposure settings. and Leica

Application Suite X software. Each experiment was performed in triplicate, and 20–30 infected

cells were analysed per time point to assess PbARK1-GFP localisation.

Immunoblotting

P. falciparum - Immunoblotting was performed as described previously[49] using primary

antibodies (Supplementary Table 4) and HRP-conjugated secondary antibodies and protein

bands were visualized using either SuperSignal™ West Pico or Femto chemiluminescent

substrates (Pierce, USA) using X-ray films.

P. berghei - Purified gametocytes were lysed in buffer containing 10 mM Tris-HCl (pH 7.5),

150 mM NaCl, 0.5 mM EDTA, and 1% NP-40. After lysis, Laemmli buffer was added, and

samples were boiled for 10 min at 95 °C, followed by centrifugation at 13,000 × g for 5 min.

The supernatants were separated on a 4–12% SDS-polyacrylamide gel and transferred onto a

nitrocellulose membrane (Amersham Biosciences). Immunoblotting was performed using the

Western Breeze Chemiluminescence Anti-Rabbit Kit (Invitrogen), following the

manufacturer's protocol. An anti-GFP polyclonal antibody (Invitrogen) was used at a 1:1,250

dilution to detect PbARK1-GFP.

Quantitative Real-Time PCR (qRT-PCR) analyses

P. falciparum - Total RNA was extracted using TRIzol reagent (G Biosciences) followed by

purification with the Qiagen RNeasy Mini Kit (Cat. No. 74104). cDNA synthesis was

performed using RevertAid H Minus Reverse Transcriptase (Thermo scientific, EP0451) and

1 μg of DNase-treated RNA as the template. Quantitative PCR was conducted using the CFX96

Real-Time PCR Detection System (Bio-Rad). Reactions were normalized using 18S rRNA as

the internal control. Relative gene expression levels were calculated using the $2^{-\Delta\Delta Ct}$ method.

Primer sequences used for the qRT-PCR are listed in Supplementary Table 3.

P. berghei - Total RNA was extracted from gametocytes using the RNA Purification Kit

(Stratagene), and cDNA was synthesised using the RNA-to-cDNA Kit (Applied Biosystems).

qPCR was performed with 80 ng of RNA using SYBR Green Fast Master Mix (Applied

Biosystems) on an Applied Biosystems 7500 Fast system. Cycling conditions were: 95 °C for

20 s, followed by 40 cycles of 95 °C for 3 s and 60 °C for 30 s. Primers were designed using

Primer3 (https://primer3.ut.ee/). Each gene was tested in three biological and technical

replicates. *hsp70* (PBANKA_081890) and *arginyl-tRNA synthetase* (PBANKA_143420) were

used as reference genes. Primer sequences are listed in Supplementary Table 3.

Transcriptome analysis using RNA-seq

Total RNA was extracted from activated gametocytes and schizonts of *P. berghei* WT-GFP and

ark1PTD parasites (three biological replicates each) using the RNeasy Kit (Qiagen), vacuum-

concentrated, and transported in RNA-stable tubes (Biomatrica). Strand-specific mRNA

libraries were prepared with the TruSeq Stranded mRNA Sample Prep Kit (Illumina) and

sequenced on an Illumina HiSeq 4000 platform (paired-end, 150 bp reads). Read quality was

assessed with FASTQC, and trimming of low-quality reads and adapter sequences was done

using Trimmomatic[50]. Reads were mapped to the *P. berghei* ANKA genome (PlasmoDB release

40) using HISAT2 v2.1.0[51], and gene counts were generated with FeatureCounts[52]. Low-
expression genes (CPM <1) were excluded. Data were normalized using the TMM method
(EdgeR[53]), transformed with voom (limma[54]), and analysed for differential expression using
DESeq2[55]. Genes with fold-change >2 and FDR <0.05 were considered differentially
expressed.

Immunoprecipitation and mass spectrometry

P. berghei purified schizonts and gametocytes (activated for 1.5 to 2 min) were crosslinked
using formaldehyde (10-min incubation with 1% formaldehyde, followed by 5-min incubation
in 0.125M glycine solution and 3 washes with PBS (pH 7.5). Immunoprecipitation was
performed using crosslinked protein and a GFP-Trap_A Kit (Chromotek) following the
manufacturer's instructions. Proteins bound to the GFP-Trap_A beads were digested using
trypsin and the peptides were analysed by LC-MS/MS and subsequently visualised using both
Principal Component Analysis (PCA) and enrichment visualised using a volcano plot[43,56].

Evolutionary bioinformatics

Our predicted proteome set was based on a version previously used (see Supplementary Table
5). Phyletic profiles for ARK1-3 and INCENP were partially derived from previous studies[10,57].
Sequence searches for missing orthologues were performed with the HHPred webserver from
MPI-toolkit[58,59] and the hmmer package[58], as previously described[60]. For the larger gene
families like Aurora kinases, we used IQ-tree-based maximum-likelihood phylogenetics
(standard setting - model finding, 1000 bootstraps[61]) to check the validity of bidirectional-best-
blast (BBH) hits amongst our genome sets on an ad hoc basis. Off note, we could not reliably
generate a single phylogenetic tree harbouring all three ARKs within the Aurora kinase
orthologous group (OG), suggesting that ARK2 and ARK3 are either extremely divergent or
not Aurora kinases. Similarly, for the INCENP gene family, multiple alignments were generally
too divergent and short to infer any consistent phylograms. All sequences used for this study
can be found in Supplementary Table 5.

AlphaFold3 modelling

822 Co-folds of 3D protein structures for ARK1 and the extended IN-box of either INCENP1 or 2
823 were modelled using the AlphaFold3 webserver (https://alphafoldserver.com/) with standard
824 settings (seed set to 100) [62].

Data availability

The RNA-seq data generated in this study have been deposited in the NCBI Sequence Read

Archive with accession number: PRJNA1309997.

Figure Legends

Figure 1. Expansion of the Aurora kinase family among apicomplexan parasites coincides

with divergent mitotic mechanisms.

a. The schematic shows schizogony in the asexual blood stage and male gametogony in

Plasmodium, highlighting some of the subcellular structures involved (below). **b.** Left: five

different cellular locations of Aurora kinases during mitosis/meiosis[10], based on findings from

model organisms. Right: Overview of the location and putative protein complex/interactors of

three Aurora kinase paralogs ARK1-2-3 amongst the three apicomplexan parasites

Plasmodium, Toxoplasma and Cryptosporidium.

Figure 2. Location of ARK1 during *P. falciparum* asexual division and *P. berghei* sexual

male gametogony.

a. Schematic of asexual division during *P. falciparum* blood stage schizogony. HA-PfARK1-

loxP parasites were synchronized and IFA was performed using anti-HA (green) and anti-

centrin (magenta) antibody on trophozoites and schizonts. DNA was stained with DAPI.

PfARK1 was observed transiently as a punctum only at nuclei with duplicated centrin foci.

Scale bar = 2 μm. **b.** Ultrastructure expansion microscopy (U-ExM) was performed on HA-

PfARK1-loxP schizonts using anti-HA and anti-tubulin antibodies. Hemispindles (white

arrow), mitotic spindles (cyan arrow) and interpolar spindles (yellow arrow) that correspond to

different stages of nuclear division were observed. PfARK1-positive puncta were mainly

associated with mitotic spindles with the puncta flanking the spindle. Scale bar = 2 μm. **c.**

Developing merozoites exhibiting PfARK1 foci (panel b) in association with hemispindles,

mitotic spindles or interpolar spindles were quantified across three independent biological

replicates (SEM ± SE, one way ANOVA, ****$P < 0.0001$, n = 3, number of parasites = 40). **d.**

Schematic of sexual male gametogony. Live cell images show the location of PbARK1-GFP

(green) at different time points (1–8 min) during male gametogony. DNA (blue) was stained

with Hoechst. DIC, differential interference contrast. Representative images of more than 50

cells with more than three biological replicates. Scale bar: 5μm. **e.** U-ExM images showing

location of PbARK1-GFP (green) and α- tubulin (magenta) detected with anti-GFP and anti-

tubulin antibodies, respectively. Hoechst was used to stain DNA (blue). Scale bar = 5 μm.

Representative images of more than 10 cells from two biological replicates. **f.** Live cell imaging

showing the location of PbARK1-GFP (green) in relation to kinetochore marker (NDC80-ch

[mCherry; magenta]) at different time points during gametogony. DNA is stained with Hoechst

dye (blue). Merge: green, magenta, and blue images merged. Representative images of more

than 20 cells with three biological replicates. Scale bar= 5 μm (left panel). 3D-SIM images of

fixed gametocytes at 1.5 min post activation (at the first spindle stage) showing the location of

PbARK1-GFP and NDC80-mCherry. Representative image of more than 10 cells from more

than two biological replicates. Scale bar = 1 μm (right panel). **g.** Left panels: live cell imaging

showing the dynamics of PbARK1-GFP (green) in relation to spindle protein kinase ARK2-

mCherry (magenta) at different time points during gametogony. DNA is stained with Hoechst

dye (blue). Merge: green, magenta, and blue images merged. Representative images of more

than 20 cells with three biological replicates. Scale bar = 5 μm (left panel). Right panels: 3D-

SIM images of fixed gametocytes at 1.5 min post activation (at the first spindle stage) showing

the location of PbARK1-GFP and ARK2-mCherry. Representative image of more than 10 cells

from more than two biological replicates. Scale bar = 1 μm. **h.** Left panels: live cell imaging

showing the dynamics of PbARK1-GFP (green) in relation to basal body/axoneme marker

kinesin-8B-mCherry (magenta) at two different time points during gametogony. DNA is

stained with Hoechst dye (blue). Merge: green, magenta, and blue images merged.

Representative images of more than 20 cells with three biological replicates. Scale bar = 5 μm

Right panels: 3D-SIM images of fixed gametocytes at 1.5 min post activation (at the first

spindle stage) showing the location of PbARK1-GFP and kinesin-8B-mCherry. Representative

image of more than 10 cells from more than two biological replicates. Scale bar = 1 μm.

Figure 3. PfARK1 controls karyokinesis, spindle formation and cytokinesis during

asexual division.

a. Growth rate assays were established in the presence of DMSO or RAP with synchronized

HA-PfARK1-loxP ring stage parasites. After 24h, DMSO or RAP was washed away, and

parasite growth was assessed after each cycle by flow cytometry. *Inset*: fold change in

parasitaemia of RAP-treated parasites relative to control (DMSO) (SEM ± SE, Two-way

ANOVA, n = 3, **$P < 0.01$, ***$P < 0.001$). **b.** HA-PfARK1-loxP parasites treated with either

DMSO or RAP as described in panel **a** were smeared, Giemsa-stained, and examined by

microscopy at the indicated time point. In the DMSO-treated parasites, the intraerythrocytic

developmental cycle progressed as expected, with ring, trophozoite, and subsequent ring stages

typical of the 48-hour parasite cycle. However, in RAP-treated, PfARK1-depleted parasites, a

significant proportion of parasites had pyknotic morphology. **c.** To assess the impact of RAP

on schizont development, HA-PfARK1-loxP parasites treated with DMSO or RAP as described

above were treated with E64 to block merozoite egress. Thin blood smears were made at ~40-

44 h.p.i in cycle 1 and stained with DAPI. RAP treated parasites contained an unsegmented

nuclear mass. Scale bar = 2 μm. **d.** HA-PfARK1-loxP parasites were treated with DMSO or

RAP. U-ExM was performed on schizonts (~40-44 h.p.i in cycle 1) during division using anti-

tubulin and anti-centrin antibodies, SYTOX and NHS-ester, visualised with magenta, green,

blue and inverted greyscale, respectively. In DMSO-treated control parasites, hemispindles

(yellow arrow), mitotic spindles (cyan arrow) and interpolar spindles (white arrow) were

observed in association with single or double centrin foci. In RAP treated parasites, either

speckles of tubulin (grey arrows) or almost complete loss of tubulin staining were observed. In

addition, unsegregated nuclei were observed and clusters of centrin were found associated with

each nuclear mass (enclosed in squares) Scale bar = 2 μm. U-ExM images are displayed as

maximum intensity projection of multiple z slices. Upper Right Panel: parasites with normal

tubulin staining reflected by the presence of spindles were counted from U-ExM images

(SEM ± SE, unpaired t test, **** $P < 0.0001$, n = 3, number of parasites = 55). Lower Right

Panel: % parasites with clustered and/or detached centrin foci (SEM ± SE, unpaired t test, ****

$P < 0.0001$, n = 3, number of parasites = 55). **e and f.** HA-PfARK1-loxP parasites were treated

with DMSO or RAP as above and cycle 1 schizonts were treated then with E64 for 2 h to block

merozoite egress and allow complete maturation and segmentation of the schizonts. IFA (e) or

U-ExM (f) was performed on unexpanded (e) or expanded (f) parasites using anti-MSP1 (e) or

anti-GAP45 and anti-tubulin (f), respectively. In DMSO treated parasites, clear segmentation

was observed whereas RAP-treated parasites either lacked MSP1 and GAP45 staining or

contained unsegregated nuclei which were circumscribed by a large MSP1 contour (cyan

arrow). U-ExM images are displayed as maximum intensity projection of multiple z slices.

Right Panel: Quantification of parasites with normal GAP45 or MSP1 staining was performed

(SEM ± SE, unpaired t test, ** $P < 0.01$ *** $P < 0.001$, n=3, number of parasites = 75 for MSP1

and 40 for GAP45). Scale bar = 2 μm. **g.** Time lapse microscopy of HA-PfARK1-loxP parasites

cultured in the presence or absence of DMSO or RAP as above and cycle 1 schizonts were

anlysed. Selected still images from videos (Supplementary Videos 6-7) are used to illustrate

the egress and dispersal of merozoites. DMSO treated parasites egressed from the erythrocytes and dispersed efficiently. RAP treated parasites egressed from erythrocytes but continued to clump-together even after prolonged duration (magenta arrow). Scale bar = 2 μm. Right Panel: parasites undergoing normal egress as defined above were counted (SEM ± SE, unpaired t test, **** P <0.0001, n=3, number of parasites= 110).

Figure 4. Conditional knockdown of PbARK1 affects male gametogony, disrupting parasite transmission.

a. qRT-PCR showing normalised expression of PbARK1 transcripts in *ark1PTD* and WT-GFP parasites. Shown is mean ± SEM; n = 3 independent experiments. Student t test was used to examine significant difference. ***p<0.01. **b**. Exflagellation centres per field at 15 min post-activation. n = 3 independent experiments (>10 fields per experiment). Error bar ± SEM. Student t test was used to assess differences between control and experimental groups. Statistical significance is indicated as ***p<0.01. **c**. Percentage ookinete conversion from zygote. n = 3 independent experiments (>100 cells). Error bar ± SEM. Student t test was used to assess differences between control and experimental groups. Statistical significance is indicated as ***p<0.01. **d**. Total number of GFP-positive oocysts per infected mosquito in *ark1PTD* compared to WT-GFP parasites at 7-, 14-, and 21 d post-infection. Mean ± SEM. n = 3 independent experiments. Two-way ANOVA test was employed to assess differences between control and experimental groups. Statistical significance is indicated as **P < 0.01, ***P < 0.001. **e.** Mid guts at 10 x and 63 x magnification showing oocysts of *ark1PTD* and WT-GFP lines at 14-, and 21 days post-infection. Scale bar: 50 μm in 10 x and 20 μm in 63 x. **f.** The diameter of GFP-positive oocysts in *ark1PTD* compared to WT-GFP parasites at 7-, 14-, and 21 d post-infection. Mean ± SEM. n=3 independent experiments. Two-way ANOVA test was employed to assess differences between control and experimental groups. Statistical significance is indicated as **P < 0.01, ***P < 0.001. **g.** Expansion microscopy images of wild-type-GFP (WT-GFP) and *ark1* gene-knockdown (*ark1PTD*) male gametocytes at 8 min or 15 min post-activation (mpa). The three images on the left show maximum intensity projections of whole-cell z-stack images labelled with NHS-ester (grey), α-tubulin (green) and Hoechst (blue). The three images on the right show magnified views of the areas enclosed by the magenta squares. BB: basal body, BBs: basal bodies (clumped), IM: inner MTOC, A: axoneme. The scale bars are scaled by an expansion factor of 4.954, calculated from a comparison of axoneme diameters measured using expansion and transmission electron microscopy images.

Figure 5. *Pbark1* gene-knockdown disrupts spindle and axoneme formation.

a. Expansion microscopy images of *P. berghei* wild-type-GFP (WT-GFP) and *ark1* gene-knockdown (*ark1PTD*) male gametocytes at 8 min post-activation (mpa). The maximum intensity projections of entire cells from a z-stack, containing spindles. Cells were labelled with NHS-ester (grey), a-tubulin (green) and Hoechst (blue). Spindles are indicated by magenta lines with a and b. **b.** Magnified views of each spindle of WT-GFP and *ark1PTD* highlighted with magenta lines with a and b in Fig. 5a. **c.** Comparison of spindle lengths measured from expansion microscopy data of WT-GFP and *ark1PTD*. Data represent mean ± SEM. (n = 7 spindles for WT-GFP and 8 spindles for *ark1PTD*, respectively across 3 different cells). Student t test was used to examine significant difference. ****p<0.0001. The scale bars and spindle lengths are scaled by an expansion factor of 4.954, calculated from a comparison of axoneme diameters measured using expansion and transmission electron microscopy images. **d.** Transmission electron microscopy images of WT-GFP and *ark1* gene-knockdown male gametocytes at 8 min or 15 min post-activation. b, f, j and n are magnified views of the areas enclosed by the magenta squares in a, e, i and m, respectively. N: nucleus, IM: inner spindle MTOC, BB: basal body, A: axoneme, K: kinetochore, Sp: spindle microtubule, F: flagellum, BBs: basal bodies (clumped), Mt: microtubule.

Figure 6. PbARK1 is the divergent Aurora kinase of the alternative chromosomal passenger complex (CPC) in *Plasmodium* spp.

a. Volcano plot of GFP-trap pulldown followed by LC-MS/MS analysis for ARK1-GFP from both 8-9 hr schizont and 1.5-min male gametocyte lysates. **b.** Zoom-in on the domain/motif topology of INCENP1/2 in apicomplexans (Pb = *P. berghei*; Tg = *T. gondii*; Cmur = *C. muris)* and model organisms (Hsap = *Homo sapiens*; Ddis = *Dictyostelium discoideum*; Gint = *Giardia intestinalis*), focused on the extended IN-box region (blue-yellow-red) that interacts with the kinase domain of ARK1. Arrows indicate an antiparallel beta sheet formed by the N- and C-terminal extensions of the IN-box. Green: TSS transphosphorylation site (asterisks indicate (putative) phosphorylated sites) in eukaryotic INCENP homologs, but not in INCENP1/2 in *P. berghei* (Pb) and *T. gondii* (Tg). RET is a candidate transphosphorylation site in PbINCENP2 (Pb-2). Protein structures are predicted using AlphaFold3, for aligned error plot, see Extended Data Fig. 8. **c.** Presence/absence of ARK1/2/3, and components of the chromosomal passenger

complex amongst the model eukaryotes, the SAR supergroup with a specific focus on Apicomplexa. Right: consensus sequence logos of the activation and P+1 loop of Aurora kinases in Clustal color scheme: divergent phosphorylation consensus in ARK1 (GxSH instead of RxT). **d.** Summary of findings and discussion points from this study.

Extended Data Figure legend

Extended Data Fig 1. Conditional Knockout of PfARK1 and effect on expression of other PfARKs.

a. Schematic representation of the strategy used to modify the PfARK1 locus using selection-linked integration (SLI). A homology region corresponding to the 5'-end of PfARK1 was cloned in pSLI-N-sandwich-loxP vector. In addition, recodonized PfARK1 was cloned downstream of yDHODH. *-stop codon, arrows- location of PCR primers that were used for genotyping and assessing excision. The above-mentioned construct was transfected in 1G5DC parasites followed by pSkipFlox plasmid, which contains a spilt Cre recombinase insert. **b**. Genotyping of HA-PfArk1-loxP parasites that appeared after drug selection. PCR amplification was carried out using primers targeting the indicated sites in panel a. PCR products of expected size, which is indicated in the table, were amplified and suggested the desired 5'- and 3'- integration whereas the wild-type (WT) locus was absent in the transgenic HA-PfARK1-loxP parasites. The amplicons were sequenced to further confirm the integration at the desired locus. **c**. HA-PfARK1-loxP parasites were treated with DMSO or 250 nM RAP for 24h. Subsequently, genomic DNA was isolated at ~44 h.p.i and PCR was performed using primer sets P7 and P8. A ~1.13kb product was obtained upon RAP treatment indicating successful excision. **d**. Western blot examination of HA-PfARK1-loxP parasite lysates (~44 h.p.i, cycle 0) treated with DMSO or RAP and using anti-HA antibody revealed that the RAP treatment resulted in effective depletion of HA-PfARK1. β-actin antibody was used as a loading control. **e**. HA-PfARK1-loxP parasites were synchronized and were harvested at Ring (R), Trophozoite (T) and Early Schizont (ES) or Late Schizont (LS) stages and Western blotting was performed using anti-HA or BiP antibodies. **f**. HA-PfARK1-loxP parasites were treated with DMSO or 250 nM RAP. Subsequently, RNA was isolated (~44 h.p.i, cycle1) and used for RT-qPCR analysis to assess the expression of ARK1/2/3 (SEM ± SE, two-way ANOVA, * P <0.05, ** P <0.01, n=3)

Extended Data Fig. 2. Generation of PbARK1-GFP parasites and analysis of subcellular location.

a. Schematic representation of the endogenous *Pbark1* locus, the GFP-tagging construct, and the recombined *ark1* locus generated by single homologous recombination. Arrows 1 and 2 indicate the positions of PCR primers used to confirm successful integration of the construct. **b.** Diagnostic PCR of ark1 and WT parasites using primers IntArk1tg (Arrow 1) and ol492 (Arrow 3) to show the correct integration. Integration of the ark1 tagging construct gives a band of 1.1 kb. **c.** Western blot showing the expression of endogenous ARK1-GFP1 detected by anti-GFP antibody. PK2 was used as control. **d.** Live cell images showing the location of ARK1-GFP during asexual blood stages of *P. berghei*. Representative of more than 50 images analysed in 3 different experiments. Scale bar = 5 μm. **e.** Live cell imaging showing the location of PbARK1-GFP (green) in relation to kinetochore marker NDC80-mCherry (magenta) during asexual blood stages of *P. berghei*. DNA is stained with Hoechst dye (blue). Merge: green, magenta, and blue images merged. Representative images of more than 20 cells with three biological replicates. Scale bar = 2 μm. **f.** Indirect immunofluorescence assays with tubulin and GFP-specific antibodies showing location of ARK1–GFP (green) in relation to tubulin (magenta) and DNA (Hoechst, blue). Merge: green, magenta, and blue images merged. Representative of more 20 images analysed in more than 3 different experiments; the scale bar is 5 μm. **g.** Live cell imaging showing the location of PbARK1-GFP (green) in relation to kinetochore marker NDC80-mCherry (magenta) at different time points during gametogony. DNA is stained with Hoechst dye (blue). Merge: green, magenta, and blue images merged. Representative images of more than 20 cells with three biological replicates. Scale bar= 5 μm. **h.** Live cell imaging showing the dynamics of PbARK1-GFP (green) in relation to spindle protein kinase ARK2-mCherry (magenta) at different time points during gametogony. DNA is stained with Hoechst dye (blue). Merge: green, magenta, and blue images merged. Representative images of more than 20 cells with three biological replicates. Scale bar= 5 μm **i.** Live cell imaging showing the dynamics of PbARK1-GFP (green) in relation to basal body/axoneme marker kinesin-8B-mCherry (magenta) at different time points during gametogony. DNA is stained with Hoechst dye (blue). Merge: green, magenta, and blue images merged. Representative images of more than 20 cells with three biological replicates. Scale bar= 5 μm.

Extended Data Fig. 3. Realtime dynamics of PbARK1-GFP during meiosis and other asexual developmental stages.

a. Live-cell imaging showing PbARK1-GFP (green) location in zygote and during ookinete development, relative to the nuclear DNA (blue, Hoechst dye), and cy3-conjugated 13.1 antibody (magenta), which recognises P28 protein on the surface of zygotes and ookinetes. Representative of more than 50 images analysed in more than three different experiments. Scale bar = 5 μm. **b.** Live cell images showing the location of PbARK1-GFP during oocysts and sporozoite stages in mosquito. Representative of more than 50 images analysed in 3 different experiments. Scale bar = 5 μm. **c.** Live cell images showing the location of PbARK1-GFP in liver stages. Representative of more 20 images analysed in 3 different experiments. Scale bar = 5 μm.

Extended Data Fig. 4. Effect of PfARK1 cKO on parasite development and division.

a and **b**. The unmodified 1G5DC SkipFlox (a) or HA-PfARK1-loxP (b) parasites were synchronized and ring stage parasites and were either left untreated or treated with RAP as described for Figure 3a. Parasites were either analyzed at indicated time points by counting parasites from Giemsa-stained thin blood smears (b) and flow cytometry (a) and % parasitaemia was determined (SEM ± SE, two-way ANOVA, ** $P < 0.01$, ****$p < 0.0001$, ns = non-significant, n=3). **c**. HA-PfARK1-loxP parasites were synchronized and ring stage parasites were treated with RAP as described for Figure 3a. The individual ring, trophozoites, and schizont stages were counted at the indicated time points from Giemsa-stained thin blood smears of parasite cultures and are represented as % of total number of parasites. In the case of RAP treated parasites, pyknotic morphology (Fig. 3b) was observed, and there were also quantified separately (SEM ± SE, two-way ANOVA, ** $P < 0.01$, n=3). **d**. This figure is the quantitation for Figure 3c. HA-PfARK1-loxP parasites were treated with DMSO or RAP and parasites were treated with E64 to prevent egress but allow complete maturation and division. Thin blood smears were made ~40-44 h.p.i in cycle 1 and nuclei were stained with DAPI. Percentage parasites with normal segregated nuclei were counted (SEM ± SE, unpaired t test, ** $P < 0.01$, n=3, number of parasites counted = 75). **e**. HA-PfARK1-loxP parasites were treated with DMSO or RAP and U-ExM was performed on schizonts (cycle 1) as described for figure 3d (hemispindle - yellow arrow, mitotic spindles – cyan arrow, interpolar spindles – white arrow). RAP treatment resulted in a loss of spindles (grey arrow) in most parasites and

1084 unsegregated nuclei were observed and several CP clusters were observed which were often

1085 detached (square).

1086

Extended Data Fig. 5. Generation of *Pbark1PTD* parasites and phenotypic analysis.

a. Schematic representation of the promoter swap strategy to construct *Pbark1PTD* parasites

(placing *ark1* under the control of the ama1 promoter) by double homologous recombination.

Arrows 1 and 2 indicate the primer positions used to confirm 5' integration and arrows 3 and 4

indicate the primers used to confirm 3' integration. **b.** Integration PCR of the promotor swap

construct into the *ark1* locus. Primer 1 (IntPTD31_5) and primer 2 (5'-IntPTD) were used to

confirm successful integration of the selectable marker, resulting in a band of 1.8 kb. Primer 3

(3'intPTama1) and primer 4 (IntPTD31_3) were used to determine the successful integration

of the clag promoter, resulting in a band of 1kb. Primer 1 (IntPTD31_5) and primer 4

(IntPTD31_3) were used to confirm a complete knock-in of the construct with a band at 4.9 kb

and the absence of a band at 1.8 kb. Representative image of more than 3 experiments. **c.** Total

number of sporozoites per gut at day 14-, and 21 post-infections; and in salivary gland at day-

21 post-infection in *ark1PTD* compared to WT-GFP parasites. Mean ± SEM. n = 3 independent

experiments**.** Two-way ANOVA test was employed to assess differences between control and

experimental groups. Statistical significance is indicated as **P < 0.01. **d.** Bite back experiment

showing successful transmission of *ark1PTD* and WT-GFP parasites from mosquito to mouse

with two different clones in three different experiments. **e.** Gene expression profile analysis

using a volcano plot to highlight up- and downregulated genes in *ark1-PTD* compared to WT-

GFP gametocytes (non-activated) with an adjusted P-value cutoff of 0.05. **f.** Gene expression

profile analysis using a volcano plot to highlight up- and downregulated genes in *ark1-*

PTD compared to WT-GFP gametocytes (activated for 30 min) with an adjusted P-value cutoff

of 0.05. **g.** Gene ontology enrichment analysis for downregulated genes in *ark1-*

PTD compared to WT-GFP gametocytes, activated for 30 min.

1110

1111 **Extended Data Fig. 6. Additional expansion microscopy and transmission electron**

1112 **microscopy images of *ark1* gene-knockdown male gametocytes.**

1113 **a.** *ark1* gene-knockdown (*ark1PTD*) male gametocytes at 8 min or 15 min post-activation

1114 (mpa). The three images on the left show maximum intensity projections of whole-cell z-stack

1115 images labelled with NHS-ester (grey), a-tubulin (green) and Hoechst (blue). The three images

on the right show magnified views of the areas enclosed by the magenta squares. BBs: basal bodies (clumped), IM: inner MTOC, A: axoneme. The scale bars are scaled by an expansion factor of 4.954, calculated from a comparison of axoneme diameters obtained using expansion and transmission electron microscopy. **b.** Transmission electron microscopy images of *ark1* gene-knockdown (*ark1PTD*) male gametocytes at 8 min or 15 min post-activation. B and F are magnified views of the areas enclosed by the magenta squares in a and e, respectively. N: nucleus, BBs: basal bodies (clumped), A: axoneme, Mt: microtubule. **c.** Additional transmission electron microscopy images revealing short spindle MTs spanning between two basal body MTOCs remained close together. IM: inner spindle MTOC, Sp: spindle microtubule, K: kinetochore.

Extended Data Fig. 7. ARK1 interactomics in schizonts and male gametocytes.
Principal Component Analysis (PCA) of duplicate **a.** schizont 8-9 h and **b.** male gametocytes 1.5 min lysates for GFP-trap of ARK1-GFP vs. WT-GFP. **c.** Table with unique peptide values for ARK1, INCENP1/2 and other associated proteins in both male gametocytes and schizonts. Values can be found in Supplementary Table 2**.**

Extended Data Fig. 8. Aligned Error (AE) plot for AF3 predictions for ARK1-INCENP1 and ARK1-INCENP1 predictions.
a. ARK1-INCENP1 prediction. **b.** ARK1-INCENP2 prediction. (i)pTM scores> 0.7 suggest a robustly predicted interaction. 3D protein structures and protein cartoons are the same as Figure 6b.

Supplementary Table 1
List of genes identified as differentially expressed in *P. berghei ark1PTD* gametocytes compared to wild-type (WT-GFP) controls. Two tabs in the spreadsheet: Tab 1: non-activated gametocytes, Tab 2: 30 min post-activation gametocytes.

Supplementary Table 2
Protein identifications for LC-MS/MS for *P. berghei* ARK1-GFP and WT-GFP. Four tabs in the spreadsheet: Tab 1: table information, Tab 2: volcano plot data, Tab 3: schizonts (8-9 h) WT vs. ARK1-GFP data, Tab 4: male gametocytes (1.5 min) WT vs. ARK1-GFP data.

Supplementary Table 3

PCR primer used in the *P. falciparum* and *P. berghei* study. Two tabs in the spreadsheet: Tab 1: primers used in the *P. falciparum* study, Tab 2: primers used in the *P. berghei* study.

Supplementary Table 4

Primary, secondary antibodies and stains used in the *P. falciparum* and P. *berghei* study. Two tabs in the spreadsheet: Tab 1: primary, secondary antibodies and stains used in the *P. falciparum* study, Tab 2: primary, secondary antibodies and stains used in the *P. berghei* study.

Supplementary Table 5

Source and sequences used during this study. Three tabs in the spreadsheet: Tab 1: table information, Tab 2: sources for predicted proteome database, Tab 3: number and amino acid sequences for ARK1/Aurora, ARK2, ARK3, INCENP1/2, borealin, survivin genes across eukaryotes with a focus on alveolata, apicomplexa.

Supplementary Video 1

Time-lapse video microscopy of gametogony in the Hoechst (blue)-stained PbARK1-GFP (green) male gametocyte after 2-3 min activation. Video is played back at 15× speed. Scale bar = 2 μm.

Supplementary Video 2

Time-lapse video microscopy of gametogony in the Hoechst (blue)-stained PbARK1-GFP (green) male gametocyte after 4-5 min activation. Video is played back at 15× speed. Scale bar = 2 μm.

Supplementary Video 3

Time-lapse video microscopy of gametogony in the PbARK1-GFP (green) male gametocyte after 6-8 min activation. Video is played back at 15× speed. Scale bar = 2 μm.

1178 **Supplementary Video 4**

1179 Time-lapse video microscopy of gametogony in the PbARK1-GFP (green) and PbNDC80-

1180 mCherry (magenta) male gametocyte after 4-5 min activation. Video is played back at 15×

1181 speed. Scale bar = 2 μm.

1182

1183 **Supplementary Video 5**

1184 Time-lapse video microscopy of gametogony in the PbARK1-GFP (green) and PbARK2-

1185 mCherry (magenta) male gametocyte after 2-3 min activation. Video is played back at 15×

1186 speed. Scale bar = 2 μm.

1187

1188 **Supplementary Video 6**

1189 Time-lapse video microscopy to assess egress of DMSO-treated HA-PfARK1-loxP parasites

1190 (related to Fig. 3g(i))

1191

1192 **Supplementary Video 7**

1193 Time-lapse video microscopy assess egress of RAP-treated HA-PfARK1-loxP parasites

1194 (related to Fig. 3g(ii))

1195

1196 **Supplementary Video 8**

1197 Time-lapse video microscopy assess egress of RAP-treated HA-PfARK1-loxP parasites

1198 (related to Fig. 3g(iii))

1199

Contributions

RT, PS, and ECT conceptualised the project. AN and PS performed the *P. falciparum*

experiments. MZ, SLP, AM, ER, and RT performed the *P. berghei* live-cell imaging. RY, DJF,

and SV performed the TEM observation. RY and MZ performed the *P. berghei* U-ExM

observation. MZ, AE, and DB performed the qPCR experiment. SA and KGLR performed the

RNAseq analysis. RT, DB, ARB, and ECT performed the proteomics analysis. ECT performed

the bioinformatics analysis. AN, RY, MZ, DJF, SLP, AM, ARB, SV, KGLR, DSG, AAH, ECT, PS, and RT performed formal analysis. AN, RY, MZ, DJF, SA, SLP, AM, ECT, and RT performed visualization. RT, PS, and ECT acquired funding. RT, PS, and ECT supervised the project. AN, RY, MZ, AAH, ECT, PS, and RT wrote the manuscript draft. All the authors reviewed and edited the manuscript.

Acknowledgement

RT is supported by ERC advance grant funded by UKRI Frontier Science (EP/X024776/1), MRC UK (MR/K011782/1), and BBSRC (BB/L013827/1, BB/X014681/1). MZ, DB, and AM were supported as research fellows by (EP/X024776/1). RY and DSG are supported by BBSRC (BB/X014681/1). SLP is supported by Wellcome DBT India Alliance/Team Science (IA/TSG/21/1/600261). AAH is supported by the Francis Crick Institute (FC001097), which receives core funding from the Cancer Research UK (FC001097), the UK Medical Research Council (FC001097), and the Wellcome Trust (FC001097). KGLR is supported by the NIH/NIAID (R01 AI136511) and the University of California, Riverside NIFA-Hatch-225935. ECT was supported by a personal fellowship from the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), the Netherlands (grant no. VI. Veni.202.223). Studies on *P. falciparum* were also supported by a Team Science grant (IA/TSG/21/1/600261) to PS, RT, and TSKP from DBT-Wellcome Trust India Alliance and funds from NII core. PS is a recipient of J.C. Bose Fellowship. We thank the School of Life Sciences Imaging (SLIM) for providing access to confocal and SIM microscopy and Dr Robert Markus for technical assistance. We thank the Nanoscale and Microscale Research Centre (nmRC) for providing access to instrumentation and Dr Michael W. Fey, Dr Julie Watts and Ms Nicola J. Weston for technical assistance. For Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. We thank Cleidiane Zampronio at Warwick University for mass spectrometry methods and Bio Support Unit, University of Nottingham for maintenance of mice used in this study.

References

Guttery, D. S., Zeeshan, M., Ferguson, D. J. P., Holder, A. A. & Tewari, R. Division and Transmission: Malaria Parasite Development in the Mosquito. *Annu Rev Microbiol* **76**, 113-134 (2022). https://doi.org:10.1146/annurev-micro-041320-010046

Sinden, R. E. Mitosis and meiosis in malarial parasites. *Acta Leiden* **60**, 19-27 (1991).

Sinden, R. E., Canning, E. U., Bray, R. S. & Smalley, M. E. Gametocyte and gamete development in Plasmodium falciparum. *Proc R Soc Lond B Biol Sci* **201**, 375-399 (1978).

Zeeshan, M. *et al.* Plasmodium SAS4: basal body component of male cell which is dispensable for parasite transmission. *Life Sci Alliance* **5** (2022). https://doi.org:10.26508/lsa.202101329

Carmena, M., Earnshaw, W. C. & Glover, D. M. The Dawn of Aurora Kinase Research: From Fly Genetics to the Clinic. *Front Cell Dev Biol* **3**, 73 (2015). https://doi.org:10.3389/fcell.2015.00073

Carmena, M., Ruchaud, S. & Earnshaw, W. C. Making the Auroras glow: regulation of Aurora A and B kinase function by interacting proteins. *Curr Opin Cell Biol* **21**, 796-805 (2009). https://doi.org:10.1016/j.ceb.2009.09.008

Hochegger, H., Hegarat, N. & Pereira-Leal, J. B. Aurora at the pole and equator: overlapping functions of Aurora kinases in the mitotic spindle. *Open Biol* **3**, 120185 (2013). https://doi.org:10.1098/rsob.120185

Zhang, X. Aurora kinases. *Curr Biol* **18**, R146-148 (2008). https://doi.org:10.1016/j.cub.2007.12.004

Li, H. *et al.* Dictyostelium Aurora kinase has properties of both Aurora A and Aurora B kinases. *Eukaryot Cell* **7**, 894-905 (2008). https://doi.org:10.1128/EC.00422-07

Zeeshan, M. *et al.* Plasmodium ARK2 and EB1 drive unconventional spindle dynamics, during chromosome segregation in sexual transmission stages. *Nat Commun* **14**, 5652 (2023). https://doi.org:10.1038/s41467-023-41395-3

Vader, G., Medema, R. H. & Lens, S. M. The chromosomal passenger complex: guiding Aurora-B through mitosis. *J Cell Biol* **173**, 833-837 (2006). https://doi.org:10.1083/jcb.200604032

Hadders, M. A. & Lens, S. M. A. Changing places: Chromosomal Passenger Complex relocation in early anaphase. *Trends Cell Biol* **32**, 165-176 (2022). https://doi.org:10.1016/j.tcb.2021.09.008

Solyakov, L. *et al.* Global kinomic and phospho-proteomic analyses of the human malaria parasite Plasmodium falciparum. *Nat Commun* **2**, 565 (2011). https://doi.org:10.1038/ncomms1558

Tewari, R. *et al.* The systematic functional analysis of Plasmodium protein kinases identifies essential regulators of mosquito transmission. *Cell Host Microbe* **8**, 377-387 (2010). https://doi.org:10.1016/j.chom.2010.09.006

Zhang, M. *et al.* Uncovering the essential genes of the human malaria parasite Plasmodium falciparum by saturation mutagenesis. *Science* **360** (2018). https://doi.org:10.1126/science.aap7847

Reininger, L., Wilkes, J. M., Bourgade, H., Miranda-Saavedra, D. & Doerig, C. An essential Aurora-related kinase transiently associates with spindle pole bodies during Plasmodium falciparum erythrocytic schizogony. *Mol Microbiol* **79**, 205-221 (2011). https://doi.org:10.1111/j.1365-2958.2010.07442.x

Wyss, M. *et al.* The three Plasmodium falciparum Aurora-related kinases display distinct temporal and spatial associations with mitotic structures in asexual blood stage parasites and gametocytes. *mSphere* **9**, e0046524 (2024). https://doi.org:10.1128/msphere.00465-24

Berry, L. *et al.* Toxoplasma gondii chromosomal passenger complex is essential for the organization of a functional mitotic spindle: a prerequisite for productive endodyogeny. *Cell Mol Life Sci* **75**, 4417-4443 (2018). https://doi.org:10.1007/s00018-018-2889-6

Collins, C. R. *et al.* Robust inducible Cre recombinase activity in the human malaria parasite Plasmodium falciparum enables efficient gene deletion within a single asexual

erythrocytic growth cycle. *Mol Microbiol* **88**, 687-701 (2013). https://doi.org:10.1111/mmi.12206

Sebastian, S. *et al.* A Plasmodium calcium-dependent protein kinase controls zygote development and transmission by translationally activating repressed mRNAs. *Cell Host Microbe* **12**, 9-19 (2012). https://doi.org:10.1016/j.chom.2012.05.014

Liffner, B. *et al.* Atlas of Plasmodium falciparum intraerythrocytic development using expansion microscopy. *Elife* **12** (2023). https://doi.org:10.7554/eLife.88088

Li, J. *et al.* Disruption of Plasmodium falciparum kinetochore proteins destabilises the nexus between the centrosome equivalent and the mitotic apparatus. *Nat Commun* **15**, 5794 (2024). https://doi.org:10.1038/s41467-024-50167-6

Ressurreicao, M. *et al.* Use of a highly specific kinase inhibitor for rapid, simple and precise synchronization of Plasmodium falciparum and Plasmodium knowlesi asexual blood-stage parasites. *PLoS One* **15**, e0235798 (2020). https://doi.org:10.1371/journal.pone.0235798

Baker, D. A. *et al.* A potent series targeting the malarial cGMP-dependent protein kinase clears infection and blocks transmission. *Nat Commun* **8**, 430 (2017). https://doi.org:10.1038/s41467-017-00572-x

Ekka, R., Gupta, A., Bhatnagar, S., Malhotra, P. & Sharma, P. Phosphorylation of Rhoptry Protein RhopH3 Is Critical for Host Cell Invasion by the Malaria Parasite. *mBio* **11** (2020). https://doi.org:10.1128/mBio.00166-20

Silvie, O. *et al.* A role for apical membrane antigen 1 during invasion of hepatocytes by Plasmodium falciparum sporozoites. *J Biol Chem* **279**, 9490-9496 (2004). https://doi.org:10.1074/jbc.M311331200

Janse, C. J., Ramesar, J. & Waters, A. P. High-efficiency transfection and drug selection of genetically transformed blood stages of the rodent malaria parasite Plasmodium berghei. *Nat Protoc* **1**, 346-356 (2006). https://doi.org:10.1038/nprot.2006.53

Nair, S. C. *et al.* A Plasmodium yoelii HECT-like E3 ubiquitin ligase regulates parasite growth and virulence. *Nat Commun* **8**, 223 (2017). https://doi.org:10.1038/s41467-017-00267-3

Bushell, E. *et al.* Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes. *Cell* **170**, 260-272 e268 (2017). https://doi.org:10.1016/j.cell.2017.06.030

Segura-Pena, D. *et al.* The structural basis of the multi-step allosteric activation of Aurora B kinase. *Elife* **12** (2023). https://doi.org:10.7554/eLife.85328

Berry, L. *et al.* The conserved apicomplexan Aurora kinase TgArk3 is involved in endodyogeny, duplication rate and parasite virulence. *Cell Microbiol* **18**, 1106-1120 (2016). https://doi.org:10.1111/cmi.12571

Carmena, M. & Earnshaw, W. C. The cellular geography of aurora kinases. *Nat Rev Mol Cell Biol* **4**, 842-854 (2003). https://doi.org:10.1038/nrm1245

Carmena, M., Wheelock, M., Funabiki, H. & Earnshaw, W. C. The chromosomal passenger complex (CPC): from easy rider to the godfather of mitosis. *Nat Rev Mol Cell Biol* **13**, 789-803 (2012). https://doi.org:10.1038/nrm3474

Ballmer, D. & Akiyoshi, B. Dynamic localization of the chromosomal passenger complex in trypanosomes is controlled by the orphan kinesins KIN-A and KIN-B. *Elife* **13** (2024). https://doi.org:10.7554/eLife.93522

Zeeshan, M. *et al.* Plasmodium kinesin-8X associates with mitotic spindles and is essential for oocyst development during parasite proliferation and transmission. *PLoS Pathog* **15**, e1008048 (2019). https://doi.org:10.1371/journal.ppat.1008048

Komaki, S. *et al.* Molecular convergence by differential domain acquisition is a hallmark of chromosomal passenger complex evolution. *Proc Natl Acad Sci U S A* **119**, e2200108119 (2022). https://doi.org:10.1073/pnas.2200108119

Trager, W. & Jensen, J. B. Cultivation of erythrocytic stages. *Bull World Health Organ* **55**, 363-365 (1977).

Lambros, C. & Vanderberg, J. P. Synchronization of Plasmodium falciparum erythrocytic stages in culture. *J Parasitol* **65**, 418-420 (1979).

Beetsma, A. L., van de Wiel, T. J., Sauerwein, R. W. & Eling, W. M. Plasmodium berghei ANKA: purification of large numbers of infectious gametocytes. *Exp Parasitol* **88**, 69-72 (1998). https://doi.org:10.1006/expr.1998.4203

Billker, O. *et al.* Calcium and a calcium-dependent protein kinase regulate gamete formation and mosquito transmission in a malaria parasite. *Cell* **117**, 503-514 (2004). https://doi.org:10.1016/s0092-8674(04)00449-0

Guttery, D. S. *et al.* Genome-wide functional analysis of Plasmodium protein phosphatases reveals key regulators of parasite development and differentiation. *Cell Host Microbe* **16**, 128-140 (2014). https://doi.org:10.1016/j.chom.2014.05.020

Tonkin, C. J. *et al.* Localization of organellar proteins in Plasmodium falciparum using a novel set of transfection vectors and a new immunofluorescence fixation method. *Mol Biochem Parasitol* **137**, 13-21 (2004). https://doi.org:10.1016/j.molbiopara.2004.05.009

Zeeshan, M. *et al.* Plasmodium NEK1 coordinates MTOC organisation and kinetochore attachment during rapid mitosis in male gamete formation. *PLoS Biol* **22**, e3002802 (2024). https://doi.org:10.1371/journal.pbio.3002802

Rashpa, R. & Brochet, M. Expansion microscopy of Plasmodium gametocytes reveals the molecular architecture of a bipartite microtubule organisation centre coordinating mitosis with axoneme assembly. *PLoS Pathog* **18**, e1010223 (2022). https://doi.org:10.1371/journal.ppat.1010223

Bertiaux, E. *et al.* Expansion microscopy provides new insights into the cytoskeleton of malaria parasites including the conservation of a conoid. *PLoS Biol* **19**, e3001020 (2021). https://doi.org:10.1371/journal.pbio.3001020

Yanase, R. *et al.* Divergent Plasmodium kinases drive MTOC, kinetochore and axoneme organisation in male gametogenesis. *Life Sci Alliance* **8** (2025). https://doi.org:10.26508/lsa.202403056

Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676-682 (2012). https://doi.org:10.1038/nmeth.2019

Ferguson, D. J. *et al.* Maternal inheritance and stage-specific variation of the apicoplast in Toxoplasma gondii during development in the intermediate and definitive host. *Eukaryot Cell* **4**, 814-826 (2005). https://doi.org:10.1128/EC.4.4.814-826.2005

Rawat, R. S. *et al.* Protein kinase PfPK2 mediated signalling is critical for host erythrocyte invasion by malaria parasite. *PLoS Pathog* **19**, e1011770 (2023). https://doi.org:10.1371/journal.ppat.1011770

Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014). https://doi.org:10.1093/bioinformatics/btu170

Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements. *Nat Methods* **12**, 357-360 (2015). https://doi.org:10.1038/nmeth.3317

Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014). https://doi.org:10.1093/bioinformatics/btt656

Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010). https://doi.org:10.1093/bioinformatics/btp616

Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015). https://doi.org:10.1093/nar/gkv007

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org:10.1186/s13059-014-0550-8

Zeeshan, M. *et al.* A novel SUN1-ALLAN complex coordinates segregation of the bipartite MTOC across the nuclear envelope during rapid closed mitosis in Plasmodium berghei. *Elife* **14** (2025). https://doi.org:10.7554/eLife.106537

Tromer, E. C., van Hooff, J. J. E., Kops, G. & Snel, B. Mosaic origin of the eukaryotic kinetochore. *Proc Natl Acad Sci U S A* **116**, 12873-12882 (2019). https://doi.org:10.1073/pnas.1821945116

Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**, e1002195 (2011). https://doi.org:10.1371/journal.pcbi.1002195

Soding, J. Protein homology detection by HMM-HMM comparison. *Bioinformatics* **21**, 951-960 (2005). https://doi.org:10.1093/bioinformatics/bti125

Zeeshan, M. *et al.* Real-time dynamics of Plasmodium NDC80 reveals unusual modes of chromosome segregation during parasite proliferation. *J Cell Sci* **134** (2020). https://doi.org:10.1242/jcs.245753

Trifinopoulos, J., Nguyen, L. T., von Haeseler, A. & Minh, B. Q. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* **44**, W232-235 (2016). https://doi.org:10.1093/nar/gkw256

Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org:10.1038/s41586-024-07487-w

Figure 1

a

Plasmodium falciparum **erythrocytic schizogony** *(blood stage - vertebrate host)*

two hemispindles — **Schizont** — **8-32x merozoites**

Plasmodium berghei **(male) gametogony** *(mosquito gut vector 1N>8N)*

1N — **continuous spindle** — 2N — 4N — **octaploid nucleus** 8N — **8x flagellated male gametes**

1 minute post activation — 2-3 mpa — 4-5 mpa — 6-8 mpa — 15 mpa

○ red blood cell   ○ cell membrane   ● nucleus   ● MTOC   ◆ basal body

═ (spindle) microtubules   axoneme   ○ (clustered) centromeres

b

Aurora localisation

MTOC — cytokinesis *(central spindle)* — microtubules — kinetochores — nucleoplasm — chromosome

Scaffolds *(e.g. human, yeast, fly, worm etc.)*

Chromosomal Passenger Complex (CPC)
4 subunits: Aurora-INCENP-Borealin-Survivin

Species-specific scaffolds
e.g. TPX2, Bora & Cep192

Three Aurora-Related Kinases (ARKs) in Apicomplexa

stage	*Plasmodium* schizont	*Plasmodium* male gametocyte	*Toxoplasma* tachyzoite	*Cryptosporidium* -
ARK3	?	*absent?*	?	*absent*
ARK2	?	*EB1-MyoK MISFIT*	?	*absent*
ARK1	?	?	*INCENP1-2*	*untested*

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Extended Data Fig. 1

a

b

Locus	Primer set	Expected size
WT	P3/P6	2.2 Kb
5' Integration	P3/P4	1.11 Kb
3' Integration	P5/P6	1.9 Kb

c

d

e

f

Extended Data Fig. 2

Extended Data Fig. 3

Extended Data Fig. 4

a

(Graph: % Parasitemia vs Time (h), DMSO (black circles) and RAP (gray triangles). Inset: Fold change vs Cycle with "ns" annotations for cycles 1 and 2.)

b

*(Graph: % Parasitemia vs Time (h), DMSO (black circles) and RAP (gray triangles). Inset: Fold change vs Cycle with "**" for cycle 1 and "****" for cycle 2.)*

Time (h): 0   24   48   72   96   120   144
Cycle: 0 | 1 | 2

c

(Bar graph: % Parasitemia with legend — Rings (black), Trophozoites (gray), Schizonts (yellow), Pyknotic/abnormal (orange). DMSO and RAP conditions.)

Time (h): 0 | 24 | 48 | 72 | 96 | 120 | 144
Cycle: 0 | 1 | 2

d

*(Scatter plot: % Normal Parasites for DMSO and RAP, ** significance.)*

e

Tubulin/Centrin/Hoechst

DMSO

RAP

Extended Data Fig. 5

a Endogenous *ark1* locus / Exchange construct / Recombinant locus

b 5'-Int PCR, 3'-Int PCR, Knock in PCR (WT/Mut); ~1.8 kb, ~1 kb, ~1.8 kb, ~4.9 kb

c Sporozoites/gut or sal gland — WT-GFP vs *ark1-PTD*; Mid gut Day 14, Mid gut Day 21, Sal. gland Day 21 (**)

d Bite-back Experiment

Exp	Parasite line	Blood Infection Day	Exp	Parasite line	Blood Infection Day
1	WT-GFP	4	5	WT-GFP	4
	ark1-PTD	No infection		*ark1-PTD*	14
2	WT-GFP	4	6	WT-GFP	4
	ark1-PTD	No infection		*ark1-PTD*	10
3	WT-GFP	4	7	WT-GFP	4
	ark1-PTD	14		*ark1-PTD*	No infection
4	WT-GFP	4	8	WT-GFP	4
	ark1-PTD	No infection		*ark1-PTD*	No infection

e Nonactivated *ark1PTD* vs WT: Volcano plot — Total = 5265 variables

f 30 min post-activation *ark1PTD* vs WT: Volcano plot — Total = 5265 variables

g 10 genes up, 8 genes down / 23 genes up, 229 genes down

Extended Data Fig. 6

Extended Data Fig. 7

a

Schizont (8-9 h)

b

Male gametocyte (1.5 min)

c

Protein IDs	Description	gene_name	category	AA	Schizonts				Male gametocytes			
					WT1	WT2	ARK1-1	ARK1-2	WT1	WT2	ARK1-1	ARK1-2
PBANKA_1343200	**inner centromere protein**	**INCENP2**	**CPC**	**2305**	0	0	60	44	0	0	76	41
PBANKA_1353900	**inner centromere protein**	**INCENP1**	**CPC**	**1126**	0	0	21	21	0	0	42	28
PBANKA_0104100	**aurora-related kinase ARK1**	**ARK1**	**CPC**	**353**	0	0	13	15	0	0	24	13
PBANKA_1011400	DNA topoisomerase 2	TOP2	chromatin architecture	1472	12	14	15	14	4	4	29	14
PBANKA_1312000	condensin-2 complex subunit G2	CAPG2	chromatin architecture	1181	0	1	2	2	0	0	10	2
PBANKA_0913000	condensin-2 complex subunit D3	CAPD3	chromatin architecture	1706	0	0	0	0	0	0	8	3
PBANKA_0301800	condensin-2 complex subunit H2	CAPH2	chromatin architecture	775	0	0	0	0	0	0	6	2
PBANKA_1337500	protein STU2	STU2	kinetochore	3198	3	1	5	2	0	0	14	6
PBANKA_0612300	apicomplexan kinetochore protein 7	AKiT7/MAD1	kinetochore	832	2	1	0	2	0	0	10	2
PBANKA_0612200	apicomplexan kinetochore protein 3	AKiT3	kinetochore	1251	0	0	0	0	0	0	8	0
PBANKA_1243900	apicomplexan kinetochore protein 5	AKiT5	kinetochore	1522	1	1	1	2	0	0	6	0
PBANKA_0621300	apicomplexan kinetochore protein 1	AKiT1	kinetochore	1737	1	1	2	2	0	0	4	3
PBANKA_1358800	kinetochore protein SPC25	SPC25	kinetochore	216	0	0	2	0	0	0	3	0
PBANKA_1115700	kinetochore protein NDC80	NDC80	kinetochore	589	0	0	0	0	0	0	3	0
PBANKA_1442300	kinetochore protein SPC24	SPC24	kinetochore	819	0	0	0	0	0	0	3	0
PBANKA_0522000	apicomplexan kinetochore protein 6	AKiT6	kinetochore	465	0	0	0	0	0	0	2	0
PBANKA_0414300	kinetochore protein NUF2	NUF2	kinetochore	457	0	0	0	0	0	0	0	2
PBANKA_0908500	myosin K	MyoK	spindle	1629	0	0	0	0	0	0	12	0
PBANKA_1038200	nuclear formin-like protein MISFIT	MISFIT	spindle	1594	0	0	0	0	0	0	7	2
PBANKA_0405600	end-binding protein 1	EB1	spindle	435	0	0	0	0	0	0	6	3
PBANKA_1458800	kinesin-15	kinesin-15	spindle	1414	0	0	0	0	0	0	5	6
PBANKA_0407400	aurora-related kinase ARK2	ARK2	spindle	1719	0	0	0	0	0	0	4	0
PBANKA_0805900	kinesin-8X	kinesin-8X	spindle	1409	0	0	0	0	0	0	4	0
PBANKA_0416100	dynein heavy chain	DHC3	MTOC	5834	4	7	1	3	7	12	46	40
PBANKA_0925400	dynein heavy chain	DHC4A	MTOC	5176	2	4	3	3	6	8	36	36
PBANKA_0214000	dynein heavy chain	DHC1	MTOC	5058	8	6	6	9	2	4	35	4
PBANKA_0507300	dynein heavy chain	DHC4B	MTOC	5309	0	4	1	0	4	6	29	31
PBANKA_0801500	conserved Plasmodium protein	CCDC63	MTOC	745	0	0	0	0	0	2	14	9
PBANKA_0202700	kinesin-8B	kinesin-8B	MTOC	1460	0	0	0	0	2	10	13	10
PBANKA_1361500	WD repeat-containing protein 16	CFAP52	MTOC	723	0	2	0	0	0	2	12	10
PBANKA_0102700	conserved Plasmodium protein	CFAP157	MTOC	372	0	0	0	0	0	0	8	4
PBANKA_1430900	SUN domain-containing protein	SUN1	MTOC	872	0	0	0	0	0	0	6	0
PBANKA_1018300	dynein intermediate chain	DNAI2	MTOC	592	2	5	2	1	0	3	5	9
PBANKA_0104200	calcium-binding protein	104200	MTOC	157	0	0	0	0	0	0	4	3
PBANKA_1035700	WD repeat-containing protein 65	CFAP57/WDR65	MTOC	1440	0	0	0	0	0	0	4	0
PBANKA_0507800	conserved protein	CFAP100	MTOC	492	0	0	0	0	0	0	3	2
PBANKA_1125300	centrosomal protein CEP135	CFAP58/CEP135	MTOC	906	0	0	0	0	0	0	3	0
PBANKA_1312800	CLASP domain-containing protein	CLASP	MTOC	859	0	0	0	0	0	0	3	0
PBANKA_1446800	conserved Plasmodium protein	CEP104	MTOC	916	0	0	0	0	0	0	3	0
PBANKA_1109600	conserved Plasmodium protein	RSP9	MTOC	672	2	2	2	2	0	0	3	2
PBANKA_0934700	conserved protein	CFAP73/CCDC42	MTOC	362	0	0	0	0	0	0	2	3
PBANKA_0504300	flagellar outer arm dynein-associated protein	DYNLRB1	MTOC	101	0	0	0	0	0	0	2	2
PBANKA_1133600	dynein light chain	1133600	MTOC	121	0	0	0	0	0	0	2	2
PBANKA_0415800	WD repeat-containing protein 66	CFAP251	MTOC	1111	0	0	0	0	0	0	2	0
PBANKA_1443000	NIMA related kinase 1	NEK1	MTOC	915	0	0	0	0	0	0	2	0
PBANKA_0708100	dynein light chain 1	DLC1	MTOC	198	0	0	0	0	0	0	0	2
PBANKA_1108000	CFA20 domain-containing protein	CFAP20	MTOC	364	0	0	0	0	3	4	0	3
PBANKA_1454600	nucleoporin NUP269	NUP269	MTOC	2260	0	0	0	0	0	0	7	0
PBANKA_0100800	zinc finger protein	100800	other	1382	2	3	5	4	0	0	19	7
PBANKA_0818500	conserved protein	818500	other	665	2	2	3	1	0	0	15	12
PBANKA_1417900	conserved protein	1417900	other	1052	1	3	1	2	0	2	14	5
PBANKA_0314200	calcium-dependent protein kinase 1	CDPK1	other	523	2	1	2	2	3	2	13	9
PBANKA_1426500	cullin-1	CUL1	other	844	1	1	1	2	0	0	12	3
PBANKA_1003400	amino acid decarboxylase AAD	UIS14	other	2262	1	1	0	2	0	0	11	0
PBANKA_1320100	conserved Plasmodium protein	UIS25	other	789	0	0	0	0	0	2	10	5
PBANKA_0933700	mitogen-activated protein kinase 2	MAPK2	other	523	0	0	0	0	0	2	7	7
PBANKA_1129100	DNA polymerase epsilon catalytic subunit A	POLE-A	DNA replication & repair	2826	7	7	9	10	0	0	41	17
PBANKA_1034100	FANCJ-like helicase	FANCJ	DNA replication & repair	1069	3	4	10	7	0	0	31	13
PBANKA_0613200	DNA polymerase alpha catalytic subunit A	POLA-A	DNA replication & repair	1709	4	5	7	6	0	3	30	17
PBANKA_1442200	protein SOC2	SOC2	DNA replication & repair	5372	1	0	5	3	0	0	25	7
PBANKA_0712100	DNA helicase	712100	DNA replication & repair	1018	2	3	5	5	0	0	24	18
PBANKA_0804300	DNA mismatch repair protein MSH2	MSH2-2	DNA replication & repair	853	1	1	7	6	4	8	24	15
PBANKA_1302600	DNA primase small subunit	PRIM1	DNA replication & repair	447	2	2	3	2	0	2	15	7
PBANKA_0307100	conserved Plasmodium protein	Timeless/Tof1	DNA replication & repair	2137	2	2	5	3	0	0	14	10
PBANKA_1358300	conserved Plasmodium protein	MSH2	DNA replication & repair	1419	0	1	1	2	0	0	12	4
PBANKA_1337600	BRCT domain-containing protein	1337600	DNA replication & repair	1265	1	0	3	3	0	0	11	5
PBANKA_0812000	DNA primase large subunit	PRIM2	DNA replication & repair	537	2	0	3	4	0	0	10	6
PBANKA_1230200	cdc2-related protein kinase 5	CRK5	DNA replication & repair	712	0	0	0	0	0	0	8	0
PBANKA_1326900	DNA polymerase alpha subunit B	POLA-B	DNA replication & repair	541	0	3	5	2	0	0	8	8
PBANKA_1006100	flap endonuclease 1	FEN1	DNA replication & repair	479	2	1	1	2	2	3	8	4
PBANKA_1356400	DNA repair protein RAD5	RAD5	DNA replication & repair	1433	0	0	0	0	0	0	7	3
PBANKA_0306500	ARMT1-like domain-containing protein	ARMT1	DNA replication & repair	454	5	3	3	0	0	0	7	2
PBANKA_1133800	DNA helicase	1133800	DNA replication & repair	867	0	0	0	0	0	0	6	3
PBANKA_1349400	DNA replication complex GINS protein	PSF2	DNA replication & repair	502	0	0	0	0	0	0	5	4
PBANKA_1448900	DNA polymerase epsilon subunit B	POLE-B	DNA replication & repair	623	1	0	1	2	0	0	5	2
PBANKA_1344300	chromatin assembly factor 1 subunit B	CAF1B	DNA replication & repair	513	0	0	0	0	0	0	5	0
PBANKA_1102900	cell division control protein 6	CDC6	DNA replication & repair	869	1	2	3	2	0	0	24	15
PBANKA_1356300	CDT1-like protein	CDT1	DNA replication & repair	955	4	5	0	5	0	0	19	8
PBANKA_0312500	origin recognition complex subunit 5	ORC5	DNA replication & repair	815	0	0	3	0	0	0	16	7
PBANKA_1348800	origin recognition complex subunit 4	ORC4	DNA replication & repair	885	2	0	3	4	0	0	12	8
PBANKA_0513900	ORC3 domain-containing protein	ORC3	DNA replication & repair	1346	0	0	2	1	0	0	11	8
PBANKA_1336900	conserved Plasmodium protein	MCM10-like	DNA replication & repair	508	0	2	3	4	0	0	11	7
PBANKA_0602000	origin recognition complex subunit 1	ORC1	DNA replication & repair	1038	0	1	0	0	0	0	11	5
PBANKA_0803000	origin recognition complex subunit 2	ORC2	DNA replication & repair	781	1	1	4	2	0	0	10	4
PBANKA_0202500	nucleoside triphosphate hydrolase	RFC-like	DNA replication & repair	1043	0	0	2	1	0	0	8	2
PBANKA_0303300	conserved Plasmodium protein	CDC45	DNA replication & repair	619	1	1	3	1	0	0	8	8

Extended Data Fig. 8

a ARK1 + INCENP1 (IN-box)

b ARK1 + INCENP2 (IN-box)

pTM=0.7; ipTM=0.777

pTM=0.79; ipTM=0.79

*Pb*INCENP1   642   979   1126

*Pb*INCENP2   1275   1507   2305

- N-terminal extension
- C-terminal extension *with CxC motif*
- coiled-coil
- IN-BOX
- TSS *(transphosphorylation)*