Plasmodium **ARK1 regulates spindle formation during atypical mitosis and forms a**
divergent chromosomal passenger complex

Annu Nagar[1+], Ryuji Yanase[2+], Mohammad Zeeshan[2#+], David J. P. Ferguson[3], Steven Abel[4],
Sarah L. Pashley[2], Akancha Mishra[2], Antonius Eze[2], Edward Rea[2], Declan Brady[2], Andrew
R. Bottrill[5], Sue Vaughan[3], Karine G. Le Roch[4], David S. Guttery[2], Anthony A. Holder[6],
Eelco C. Tromer[7&], Pushkar Sharma[1&], Rita Tewari[2&]

[1]Eukaryotic Gene Expression Laboratory, National Institute of Immunology, New Delhi,
India-10067.
[2]School of Life Sciences, University of Nottingham, Nottingham, UK.
[3]Department of Biological and Medical Sciences, Oxford Brookes University, Oxford, UK.
[4]Department of Molecular, Cell and Systems Biology, University of California Riverside, 900
University Ave., Riverside, USA.
[5]School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry, UK.
[6]Malaria Parasitology Laboratory, The Francis Crick Institute, London, UK.
[7]Cell Biochemistry, Groningen Biomolecular Sciences and Biotechnology Institute, Faculty
of Science and Engineering, University of Groningen, Groningen, The Netherlands.

[#]Present address: Division of Molecular Microbiology and Immunology, CSIR-Central Drug
Research Institute, Lucknow, India.
[+]Contributed equally
[&]joint corresponding author:
Tromer, E.C; e.c.tromer@rug.nl
Pushkar Sharma; pushkar@nii.ac.in
Rita Tewari; rita.tewari@nottingham.ac.uk (Lead)

Abstract

Mitosis in *Plasmodium* spp., the causative agent of malaria, is fundamentally different from model eukaryotes, proceeding via a bipartite microtubule organising centre (MTOC) and lacking canonical regulators such as Polo kinases. During schizogony, asynchronous nuclear replication produces a multinucleate schizont, while rapid male gametogony generates an octaploid nucleus before gamete formation. Here, we identify Aurora-related kinase 1 (ARK1) as a key component of inner MTOC and spindle formation, controlling kinetochore dynamics and driving mitotic progression. Conditional ARK1 depletion disrupts spindle biogenesis, kinetochore segregation, karyokinesis and cytokinesis in both stages, and affects parasite transmission. Interactome analysis shows that ARK1 forms the catalytic core of a non-canonical chromosomal passenger complex (CPC) containing two highly divergent inner centromere proteins (INCENPs), which we term INCENP-A and INCENP-B, and lacking the canonical chromatin-targeting subunits Survivin and Borealin. Comparative genomics suggests that apicomplexan INCENPs arose through recurrent lineage-specific duplications, reflecting an evolutionary rewiring of CPC architecture in this eukaryotic lineage. Together, these findings reveal key adaptations in *Plasmodium* mitosis involving ARK1 and its INCENP scaffolds, and identify the ARK1–INCENP interface as a potential multistage target for antimalarial intervention.

Keywords: *Plasmodium*; Aurora kinase; chromosomal passenger complex (CPC); INCENP; spindle; karyokinesis; male gametogony; MTOC; cytokinesis; transmission

Introduction

Plasmodium spp., the causative agent of malaria, is a haploid unicellular eukaryote of the phylum Apicomplexa, which is evolutionarily distant from most model organisms and has some unusual divergent features of cell division. During its life cycle, mitosis is closed and occurs during two distinct forms of cell division: schizogony within vertebrate host cells, and male gametogony within the female mosquito gut (Fig. 1a). During asexual proliferation within host red blood cells (schizogony), repeated asynchronous nuclear division, without chromosome condensation, produces a multinucleate schizont. Subsequent budding produces invasive merozoites to continue the cycle. In contrast, during male gametogony three rounds of rapid genome replication to octaploid (1N to 8N) occur in 8 minutes within the intact nucleus. Karyokinesis and cytokinesis are synchronised with axoneme dynamics by replication and separation of a bipartite microtubule-organising centre (MTOC), chromosome condensation, and complementary *de novo* basal body formation, leading to the budding of eight flagellated male gametes. These events occur within 12 to 15 min of gametocyte activation, which is probably the most rapid mitosis amongst eukaryotes[1]. Compared to schizogony, mitosis in male gametogony is unusual, with atypical continuous spindles and a bipartite MTOC that spans the nuclear envelope and coordinates the simultaneous assembly of spindle microtubules and the flagellar axoneme. Male gametes are the only flagellated stage in the parasite's life cycle, and therefore the only stage where centriole formation and *de novo* replication of the basal body occur[1, 2, 3, 4].

Aurora-related kinases (ARKs) are multi-functional mitotic regulators that coordinate spindle assembly, chromosome segregation, and late mitotic events across eukaryotes. They act on spindle pole duplication, spindle assembly kinetochore-microtubule attachment, mitotic checkpoints and finally cytokinesis[5, 6, 7, 8]. Budding yeast and the amoeba *Dictyostelium discoideum*[9] have a single ARK to execute all functions, suggesting these to be ancestral to all eukaryotes[7]. However, multiple different lineages (e.g. vertebrates, plants, kinetoplastids and others) have one or more paralogues that resulted from parallel duplications, which subdivided functions over the different paralogues[7, 10]. One paralogue (Aurora B in humans) is the catalytic centre of the chromosomal passenger complex (CPC), a key regulator of mitotic events at the cell equator, which initially hitchhikes on the inner centromere and finally remains at the central spindle to orchestrate cytokinesis. In canonical model systems, the CPC consists of Aurora B, inner centromere protein (INCENP), Survivin and Borealin[11,

[12] that guide its localisation to kinetochores, spindle and the central spindle. It is also critical

for mitotic histone phosphorylation, fixing errors in kinetochore-spindle attachment, and

correct cytokinesis (Fig. 1b).

Apicomplexans, including *Plasmodium* and *Toxoplasma* encode three ARKs, most of which

lack canonical localisation and activation motifs found in model eukaryotes[10, 13, 14]. ARK1, 2

and 3 are refractory to gene disruption and indispensable during asexual blood stage

schizogony in both *P. berghei* and *P. falciparum*[13, 14, 15]. Consistent with this, a recent study

has chemically validated Aurora kinases as promising targets for the development of multi-

stage antimalarials[16, 17]. ARK1 has been observed at the MTOC during schizogony and in *P.*

falciparum gametocytes[18, 19]. However, its precise function in asexual division and its

behaviour during sexual development have remained unclear. Our recent studies on ARK2

revealed its association with spindle microtubules (MTs) forming a divergent protein complex

with EB1, myosin K and MISFIT[10], but little is known of ARK1 and 3 functions during male

gametogony or whether they are essential for parasite transmission. . ARK1 function has also

been explored in another apicomplexan, *Toxoplasma gondii*, where it interacts with two

INCENP paralogues and is important for asexual replication. Whether *Plasmodium* INCENPs

are direct orthologues of *Toxoplasma* INCENPs is not clear. The function of *Toxoplasma*

ARK1 during male gametogony is difficult to study because the sexual cycle of these

parasites only takes place in the gut of cats (Fig. 1b)[20].

Here, we have investigated *Plasmodium* ARK1, integrating studies of location, function and

molecular interactions across two life-cycle stages using *P. falciparum* (Pf, for asexual blood-

stage schizogony) and *P. berghei* (Pb, for male gametogony). High-resolution live-cell

imaging and expansion microscopy revealed ARK1 associated with the mitotic spindle,

kinetochores, and the inner MTOC. Conditional gene knockout using rapamycin-DiCre in *P.*

falciparum[21] and gene knockdown using a promoter-trap approach in *P. berghei*[22] revealed

that ARK1 regulates spindle biogenesis, karyokinesis and cytokinesis during both schizogony

and male gametogony, suggesting that ARK1 is a critical regulator of *Plasmodium* mitosis.

Interactome analysis identifies ARK1 as the catalytic component of a non-canonical

Chromosomal Passenger Complex (CPC) comprising two divergent INCENP paralogues–

here termed INCENP-A and INCENP-B–in the absence of Survivin and Borealin. Together,

these findings reveal how ARK1 and its INCENP scaffolds contribute to the distinctive

mechanisms of apicomplexan nuclear division.

Results

ARK1 is present in asexual stage parasites undergoing mitotic division

We studied the location of PfARK1 in asexual blood stages, using a rapamycin-inducible dimerizable Cre recombinase DiCre-based strategy to tag and knockout *PfARK1* in the 1G5DC parasite line (Supplementary Fig. 1a)[21]. These parasites (HA-PfARK1-loxP) were used to locate PfARK1 during parasite division using an antibody to the HA-tag included at its N-terminal. Indirect immunofluorescence (IFA) and Western blot indicated that ARK1 is expressed only in trophozoite and schizont stages during asexual division, but was not detectable in ring and segmentor (Fig. 2a, Supplementary Fig. 1e). ARK1 was present as a punctum proximal to the MTOC, as identified by staining with an anti-centrin antibody, present only in cells with two associated centrin puncta, indicative of ongoing mitosis (Fig. 2a, c). These observations were largely consistent with previous studies of PfARK1[18, 19].

To gain a deeper insight into ARK1 dynamic organisation, Ultrastructure Expansion Microscopy (U-ExM) was performed, and visualized ARK1 along with spindle microtubules using an anti-tubulin antibody to track spindle biogenesis. As reported previously[23], parasites possessed hemispindles prior to the onset of division, but PfARK1 was not expressed in these parasites or in parasites containing interpolar spindles (Fig. 2b, d). Strikingly, PfARK1 was found specifically associated with mitotic spindles and PfARK1-positive double puncta seemed to be present at the ends of the growing mitotic spindles (Fig. 2b, d). These data strongly indicate that ARK1 is expressed transiently, and only when parasite nuclei are undergoing mitotic division.

These observations were also validated using the rodent malaria parasite, *P. berghei*, a well-established model for studying male gametogony and parasite development within the mosquito host[10]. The subcellular location of PbARK1 in asexual stages was examined by generating an endogenous, C-terminal GFP-tagged PbARK1 (PbARK1-GFP) transgenic parasite line (Supplementary Fig. 2a-c and Supplementary Note 1). Live cell imaging of PbARK1-GFP in asexual blood stages revealed a diffuse location in early ring stages. During trophozoite and early schizont stages, PbARK1-GFP re-localised to discrete focal points in the nucleus (Supplementary Fig. 2d). As schizogony progressed, the focal points of fluorescence divided into two closely associated puncta, which followed subsequent asynchronous nuclear divisions and eventually became diffuse during the later stages

(Supplementary Fig. 2d, e). These observations are consistent with the ARK1 location in *P.*
falciparum as described above (Fig. 2a).

PbARK1 is located in the nucleoplasm, at the spindle poles and along microtubules
during male gametogony

In gametocytes both before and immediately after activation, PbARK1-GFP had a diffuse
distribution predominantly within the nucleoplasm. Within one-min post-activation, part of
this was concentrated to a single discrete focal point adjacent to the DNA (Fig. 3a), and then
the distribution rapidly extended to form an arc-like structure, which within three-min post-
activation divided into two distinct puncta positioned at opposite sides of the nucleus (Fig. 3a
and Supplementary Video 1). The two PbARK1-GFP foci elongated again, forming two arc-
like structures along the periphery of the nucleus (Fig. 3a and Supplementary Video 2). By 6-
to 8-min post-activation, a subsequent round of extension and division had produced eight
discrete PbARK1-GFP foci, coincident with the completion of three successive rounds of
mitosis (Fig. 3a and Supplementary Video 3). 3D-structured illumination microscopy
(SIM) at 1.5–2 min post-activation clearly resolved the arc-like localisation of PbARK1-GFP
across the nucleus, with distinct foci visible at both ends of the arc (Fig. 3b).

The arc-like pattern of PbARK1 observed in live cell imaging and 3D-SIM coincided with
the pattern of anti-tubulin staining in U-ExM, suggesting an association with the mitotic
spindle and spindle poles or inner MTOCs (Fig 3c and Supplementary Fig. 3a). To further
investigate ARK1 location, dynamics, and association with mitotic spindles and kinetochore
segregation, we compared the location of PbARK1 with that of the spindle marker ARK2, the
kinetochore marker NDC80, and cytoplasmic axonemal protein kinesin-8B. We generated
dual fluorescent parasite lines expressing PbARK1-GFP (green) and either PbARK2-
mCherry (magenta), PbNDC80-mCherry (magenta), or Pbkinesin-8B-mCherry (magenta) by
genetic cross as described previously[4]. Live-cell imaging showed that both PbARK1-GFP
and PbARK2-mCherry were co-located at the spindle (Fig. 3d and Supplementary Video 5).
Further 3D-SIM at 1.5–2 min post-activation showed an association of PbARK1-GFP with
PbARK2-mCherry at the spindle, with PbARK1-GFP more outside at the spindle poles (Fig.
3e and Supplementary Fig. 3b). In parasite lines expressing PbARK1-GFP and PbNDC80-
mCherry, these proteins were located next to the Hoechst-stained DNA, partially overlapping
at the spindle but not at the spindle poles (Fig. 3f and Supplementary Video 4). 3D-SIM at
1.5–2 min post-activation showed a clear association of PbNDC80-mCherry with PbARK1-

GFP at the spindle but not at the spindle poles (Fig. 3g and Supplementary Fig. 3c). However,

in a parasite line expressing PbARK1-GFP and Pbkinesin-8B-mCherry (a cytoplasmic

marker) there was no overlap between these two proteins (Fig. 3h), an observation which was

further substantiated by 3D-SIM imaging (Fig. 3i).

During meiotic stages of development post-fertilisation, PbARK1-GFP was diffuse as seen

during schizogony, and male gametogony in the cytoplasm, with transient nuclear foci

forming in zygotes and disappearing in mature ookinetes (Supplementary Fig. 4a and

Supplementary Note 2). A diffuse cytosolic signal with some foci was also observed

throughout oocyst and sporozoite stages, including those from salivary glands

(Supplementary Fig. 4b and Supplementary Note 2). In liver stages (in HepG2-infected cells),

the PbARK1-GFP signal remained diffuse with faint foci even at 48 hours post infection

(hpi), with slight perinuclear enrichment at 64 hpi (Supplementary Fig. 4c and Supplementary

Note 2).

PfARK1 regulates spindle formation and karyokinesis

The functional role of PfARK1 in asexual blood stage development was assessed by

examining the effect of deleting the gene. Rapamycin (RAP) treatment of HA-PfARK1-loxP

parasites efficiently excised the floxed sequence (Supplementary Fig. 1c) and caused

depletion of PfARK1 protein (Supplementary Fig. 1d). Synchronised ring stage cultures of

1G5DC-SkipFlox (control) (Supplementary Fig. 5a) or HA-PfARK1-loxP parasite lines (Fig.

4a and Supplementary Fig. 5b) were treated with RAP or DMSO (control) and parasite

growth was monitored every 24 h. In cycle 0 (RAP-treatment) there was little effect on the

parasitaemia of HA-PfARK1-loxP parasites. However, a significant growth defect was

observed at the beginning of the next cycle (cycle 1) as indicated by a marked reduction in

parasitaemia (Fig. 4a and Supplementary Fig. 5b), whilst there was almost no effect of RAP

treatment on 1G5DC-SkipFlox (Supplementary Fig. 5a). This growth defect was further

accentuated in subsequent cycles. Microscopic examination of Giemsa-stained blood smears

suggested normal ring to trophozoite development in cycle 1 of DMSO-treated parasites

(Supplementary Fig. 5c). However, following RAP treatment and PfARK1 depletion, a

population of parasites were pyknotic (at ~96 and 144 hpi) and fewer rings were found in

cycle 1 and 2 (Fig. 4b and Supplementary Fig. 5c). Schizont development following

treatment with E64 to block late schizont maturation and merozoite egress was analysed: a

significant number PfARK1-depleted cycle 1 schizonts had abnormal morphology, with

unsegregated nuclei (Fig. 4c and Supplementary Fig. 5d), which is consistent with the role of
PfARK1 in nuclear division. This suggestion, that the decrease in parasitaemia resulting from
PfARK1 depletion was due to its role in nuclear division and/or schizont maturation, was
investigated further.

Next, efforts were made to investigate whether PfARK1 regulates MTOC duplication and
spindle formation during asexual division of the parasite. U-ExM revealed the formation of
various forms of spindles as described earlier with one or two centrin-labeled foci on the
cytoplasmic side of the nuclear envelope (Fig. 4d) and associated with NHS-ester labelled
MTOCs or associated with mitotic spindles[23, 24]. In PfARK1-depleted parasites, a very few
had associated spindles indicated by either absent or speckled tubulin staining (Fig. 4d, grey
arrow and Supplementary Fig. 5e) and as mentioned above the nuclei appeared unsegregated.
Also, multiple centrin foci were associated with these unsegregated nuclei, and in some cases
centrin aggregates were seen away from the nuclei and disorganised (Fig. 4d and
Supplementary Fig. 5e, square). Collectively, these data suggested that while the MTOC
duplication was unaltered, it was disorganised and spindle formation and karyokinesis were
impaired, indicating that PfARK1-depletion leads to defective mitosis.

PfARK1 depletion impairs segmentation of daughter merozoites
At the end of schizogony, segmentation of the schizont by invagination of the parasite plasma
membrane leads to formation of merozoites prior to their egress[25]. IFAs were performed
using antibody to PfMSP1, a parasite surface marker, to assess segmentation following
ARK1 depletion. PfMSP1 fluorescence with a bunch of grape-like contours, characterised the
membrane of control parasites, but in contrast, PfARK1-depleted parasites either lacked
PfMSP1 staining, or contained large aggregations that encompassed multiple merozoites or
unsegregated nuclei (Fig. 4e, cyan arrow).

U-ExM of mature (segmented) schizonts with anti-GAP45 antibody-labelling revealed inner
membrane complex (IMC) and tubulin labelled subpellicular MT, and rhoptries labeled with
NHS-ester (Fig. 4f). However, for the PfARK1-depleted parasites, several unsegregated
nuclei were present and these cells either lacked GAP45 staining, or the parasite pellicle (the
plasma membrane/IMC) was disorganised (Fig. 4f, cyan arrow, square, zoom and
Supplementary Fig. 5f). Although NHS-ester labelled rhoptries were found, their organisation
seemed to be disorganised which may be due to aberrant segmentation. In addition,

subpellicular MT were either not observed or were mislocalised (Fig. 4f, cyan arrow, square,

zoom). These defects were possibly due to arrest in division of the parasites as also indicated

by aberrant GAP45 staining. These observations suggested impaired cytokinesis of daughter

merozoites upon PfARK1 depletion, the aberrant karyokinesis or nuclear division possibly

contributes to these defects.

To study the effect of PfARK1 depletion on merozoite release and egress, time-lapse

microscopy was performed[26]. For this purpose, parasites were treated with protein kinase G

(PKG)-inhibitor ML-10 to allow schizont maturation but prevent egress, followed by its

removal which allowed egress and was captured by time-lapse microscopy[27, 28]. Egress and

dispersal of merozoites for control parasites was observed (Fig. 4g and Supplementary Video

6), and while egress from the erythrocyte initially appeared almost normal in PfARK1-

depleted parasites, the merozoites either remained attached to the residual body or clumped

together (Fig. 4g, magenta arrows, and Supplementary Videos 7, 8) even after an extended

period post-egress. These defects likely resulted from the abnormal nuclear segregation and

cytokinesis in PfARK1-depleted parasites.

PbARK1 knockdown reveals an essential role during male gametogony and parasite

transmission

Previous analysis by gene disruption had shown that PbARK1 is probably essential

for asexual blood-stage development[14]. To be able to manipulate its expression in sexual

stages we used a promoter trap strategy, replacing the *ark1* promoter with the promoter from

ama1, resulting in $P_{ama1}ark1$ hereafter referred as (*ark1PTD*), a gene which is not transcribed

in gametocytes but is highly expressed in asexual blood stages (Supplementary Fig. 6a)[29].

Correct genetic integration in the transgenic parasite line was confirmed by PCR

(Supplementary Fig. 6b), and *ark1* transcription was shown to be significantly downregulated

in *ark1PTD* gametocytes by qRT-PCR (Fig. 5a). A phenotypic analysis of these *ark1PTD*

parasites was then performed at different stages of parasite development within the mosquito.

Phenotypic analysis revealed that male gametogony (as scored by exflagellation centres) was

severely affected in the *ark1PTD* line compared to GFP control (wild-type-GFP; WT-GFP)

parasite (Fig. 5b)[30], and therefore significantly fewer ookinetes (differentiated zygotes after

fertilisation) were formed (Fig. 5c). In mosquitoes fed with *ark1PTD* parasites, we observed

significantly fewer oocysts in the midgut at day 7 post infection, in contrast to the hundreds

of oocysts in mosquitoes fed WT-GFP parasites (Fig. 5d). The number of *ark1PTD* oocysts

was further reduced by day 14 and day 21 post infection compared to WT-GFP oocysts (Fig.

5d). The *ark1PTD* oocysts were also significantly smaller than the WT-GFP oocysts 7 days

post infection and diminished further by day 21 post infection (Fig. 5e, f). The number of

salivary glands sporozoites was significantly reduced in *ark1PTD* parasites on day 14

(Supplementary Fig. 6c). However, the ratio of salivary gland to midgut sporozoites on day

21 was comparable to that of WT-GFP parasites (Supplementary Fig. 6c), suggesting that the

surviving *ark1PTD* sporozoites retain their invasive potential and ability to migrate to the

salivary glands. To further confirm the absence of sporogony, mosquitoes that had been fed

with *ark1PTD* or WT-GFP parasites were allowed to feed on susceptible naïve mice (bite-

back experiments). Mosquitoes infected with WT-GFP parasites successfully transmitted the

parasite, with blood-stage infection detected in naïve mice after 4 days. In contrast, *ark1PTD*

parasites failed to transmit in most experiments (five out of eight) (Supplementary Fig. 6d).

However, on three occasions, they were able to establish infection in these bite-back

experiments, although the prepatent period was markedly prolonged, with blood-stage

parasites first detected 14 days post-infection (Supplementary Fig. 6d). Collectively, these

findings indicate that the observed transmission defect is primarily driven by the reduction in

sporozoite numbers rather than a loss of sporozoite invasiveness.

To investigate the genome-wide transcriptional impact of PbARK1 depletion, we performed

RNA-seq analysis. While few changes were observed in non-activated gametocytes, activated

ark1PTD gametocytes (30 mpa) exhibited a broad downregulation of genes involved in

signalling, lipid biosynthesis, and ribosome biogenesis. This transcriptional repression is

consistent with the observed developmental arrest (Supplementary Fig. 6e-g, and

Supplementary Note 3).

ark1PTD parasites have defective spindle formation, MTOC separation, and axoneme

formation during male gametogony

To further investigate the function of PbARK1 during male gametogony, WT-GFP and

ark1PTD male gametocytes were observed during gametogony using U-ExM. In WT-GFP

male gametocytes at 8 mpa, individual MTOCs were distinct and separated, whereas in

ark1PTD male gametocytes multiple MTOCs were clumped and aggregated together (Fig. 5g

and Supplementary Fig. 7a). At 15 mpa, WT-GFP male gametocytes were observed

undergoing exflagellation, whereas in *ark1PTD* male gametocytes the MTOCs were still

clumped together and no exflagellation was observed in the examined cells (Fig. 5g and

Supplementary Fig. 7a).

We examined whether the lack of MTOC separation in the *ark1PTD* line was caused by a

defect in spindle formation. At 8 mpa, long spindles with clearly separated MTOCs were

present in WT-GFP parasites, but in *ark1PTD* lines, short spindles, with the two MTOCs

remaining close together (Fig. 6a, b) were present. Measurement showed that *ark1PTD*

spindles were significantly shorter (~0.8 μm) than WT-GFP spindles (~2.5 μm; Fig. 6c).

These findings corroborated studies on PfARK1 during schizogony as depletion of the kinase

resulted in abnormal spindle formation and accumulation of MTOCs around unsegregated

nuclei (Fig. 4d).

TEM observations further revealed structural defects in *ark1PTD* male gametocytes. At 8

mpa, in the *ark1PTD* line, MTOCs formed a normal bipartite structure consisting of an inner

spindle MTOC and an outer basal body MTOC, similar to those in WT-GFP male

gametocytes, and normal axoneme formation was observed (Fig. 6dA-D, I-L and

Supplementary Fig. 7bA-D), However, we also observed basal body MTOCs that appeared

detached from the nucleus in the *ark1 PTD* line (dBB; Fig. 6dI, J and Supplementary Fig.

7bA, B, D). As revealed by expansion microscopy, multiple basal body MTOCs were often

aggregated and clumped together (Fig. 6dL). Within the cytoplasm, we observed singlet or

doublet MT that may have been axoneme precursors and likely represent failures in axoneme

formation (Fig. 6dK, L).

At 15 mpa, WT-GFP gametocytes had started to form gametes with flagellum and nucleus

(Fig. 6dE, F, G, H), while in the *ark1PTD* line there was incomplete axoneme formation and

exflagellation rarely occurred (Fig. 6dM-P and Supplementary Fig. 7bE-H). In addition, two

basal body MTOCs remained close together without proper separation (Fig. 6dM, N, P), with

short spindle MT visible between them (Fig. 6dP and Supplementary Fig. 7c). These results

suggest that PbARK1 is essential for successful mitosis during male gametogony.

PbARK1 is part of a divergent chromosome passenger complex (CPC) with two

INCENP paralogues

To understand whether ARK1 localisation could be explained by recruitment to specific

regulatory scaffolds, we investigated which interacting proteins might direct ARK1 activity

and positioning during both asexual and sexual stages of the parasite life cycle. We prepared

lysates from *P. berghei* schizonts and male gametocytes expressing PbARK1-GFP, performed

GFP-Trap immunoprecipitation, and identified co-purifying proteins by mass spectrometry.

At both stages two divergent inner centromere protein paralogues — here termed INCENP-A

and INCENP-B — were the predominant ARK1 interactors (Fig. 7a and Supplementary Fig.

8). This physical interaction is further supported by a recent study demonstrating that

PbARK1 co-localises with both INCENP-A and INCENP-B at spindle poles and

kinetochores in *P. berghei*[31]. Although replicate numbers were limited (n = 2), ARK1

pulldowns from male gametocytes also showed coordinated enrichment of several spindle

and kinetochore proteins amongst the pulldowns (Supplementary Data 2) including

components of the ARK2-associated spindle complex (MISFIT-MyoK-EB1, multiple

kinetochore proteins (AKiT1,3,5 and 7) and the condensin-II- subunits (CapH2-G2-D3).

No Survivin- or Borealin-like proteins were detected, supporting earlier predictions that

Plasmodium spp., and more generally apicomplexans, lack these conserved components of

the CPC (Fig. 7c). Given the distinct localisation dynamics of ARK1 suggesting that it may

not act as the classic passenger complex on chromosomes, it is possible that the chromatin-

directing function of Survivin and Borealin has become obsolete for apicomplexans and that

the structural motifs for localisation and activation of ARK1 by INCENP-A and -B might be

revealed upon detailed comparative sequence analyses.

Because *T. gondii* also encodes two INCENP paralogues that interact with TgARK1[20], we

examined INCENP evolution with a focus on alveolates and apicomplexans (Fig. 7b, c).

Phylogenetic reconstructions did not reveal a clear scenario supporting either a single

ancestral duplication or multiple recurrent duplication events. Instead, the observed patterns

are more consistent with parallel, lineage-specific duplications, rather than a conserved

INCENP1/INCENP2 split inherited from the apicomplexan ancestor – hence we named the

paralogue in *Plasmodium* INCENP-A/-B to prevent any misinterpretation of the predicted

functional equivalence of INCENP between the *Toxoplasma* and *Plasmodium* paralogues. We

found parallel but separate duplications in other different alveolate lineages such as

apicomplexpan-like *Squirmida*, *Chrompodellida* and dinoflagellate-related lineages

concomitant with the loss of Borealin/Survivin (Fig. 7c), suggesting INCENP duplications

might compensate for loss of these chromatin-directing subunits. Notably, INCENP-B is

absent from *Piroplasmida*, which also lack ARK2 and ARK3, suggesting functional co-

evolution among specific Aurora and INCENP paralogues.

AlphaFold3—based co-folding of *Pb*ARK1 with INCENP-A or INCENP-B predicted a

conserved IN-box-ARK1 kinase interaction flanked in all apicomplexan proteins by N- and

C-terminal extensions that form a beta-sheet – folding INCENP-A/-B around ARK1 (Fig. 7b

and Supplementary Fig. 9). In addition, the C-terminal extension has a conserved CxC motif,

possibly involved in posttranslational modification (Fig. 7b). We could only predict a coiled-

coil in the N-terminus of INCENP-B, and no further structural features could be discerned for

INCENP-A using AlphaFold3. Such absence of any feature makes it hard to predict the

binding mode of INCENP-A either as a single ARK1-INCENP-A interaction or forming an

INCENP-A-ARK1-INCENP-B trimeric complex.

Finally, we examined whether non-canonical sequence features in ARK1 or its INCENP

partners might point to alternative models of CPC regulation in *Plasmodium* spp., given that

autophosphorylation sites in both Aurora B and INCENP are critical for CPC function in

vertebrates[32]. Both the broadly conserved Aurora-kinase activation-loop motif (RxTxCGT)

and the canonical INCENP activation site (RTSS in vertebrates) are absent from the

Plasmodium proteins: ARK1 instead carries a divergent GxSH motif, and INCENP-A and

INCENP-B lack an RTSS-equivalent position. However, INCENP-A and INCENP-B retain

Aurora consensus motifs in adjacent regions (Fig. 6b, green boxes). For comparison, ARK2

paralogues retain a canonical Aurora activation loop motif, whereas ARK3 possesses a

serine-rich activation loop containing multiple candidate phosphorylation sites (Fig. 7c). The

absence of clear conserved auto/trans-phosphorylation sites between ARK1 and INCENP-A/-

B in *Plasmodium* spp. suggest that CPC kinase activity may rely on a distinct regulatory

mechanism, potentially involving input from an additional kinase.

Discussion

Our cross-stage analysis identified ARK1 as a central organiser of *Plasmodium* mitosis using

P. falciparum and *P. berghei* model systems. First, ARK1 localises to the spindle and inner

MTOC specifically during mitosis, and its depletion in *P. falciparum* reduces spindle formation, prevents proper karyokinesis, disrupts segmentation and impedes egress. Second, during the rapid mitoses of male gametogony, *P. berghei* ARK1 dynamics follow spindle poles; knockdown shortens spindles, block MTOC separation and axoneme biogenesis, and abolishes transmission. These phenotypes demonstrate that ARK1 regulates spindle formation and MTOC separation, thereby ensuring successful sister chromatid segregation and gamete formation.

Our findings show that ARK1 is preferentially expressed and targeted during the mitotic stages marked by spindle formation. Live cell imaging and expansion microscopy revealed that ARK1 locates to the spindle apparatus and the spindle poles, corresponding to the inner core of the bipartite MTOC[4, 23]. Expansion microscopy using MTOC (centrin) and microtubule (tubulin) markers confirmed that ARK1 associates with spindle poles within the inner MTOC in both mitotic stages. Some of these features had been observed earlier for *P. falciparum* ARK1 and further it had been shown that its location is limited to mitotically dividing stages[19]. Dual fluorescence microscopy analysis during male gametogony revealed that it colocalised with ARK2, another aurora kinase reported recently[10], and the kinetochore marker NDC80 at the plus ends of spindle microtubules, suggesting a cooperative role for ARK1 and ARK2 at equatorial spindles. It also has a role in kinetochore dynamics, as observed previously for ARK2[10]. However, live cell imaging in *P. berghei* clearly showed that it is part of the inner MTOC as it does not colocalise with the basal body and axoneme marker, kinesin-8B. This contrasts with *T. gondii* ARK1, which, when tagged at the N-terminus, exhibited a dynamic localisation to both nucleus and cytoplasm during division, but remained predominantly nuclear in non-dividing cells [20, 33]. Its colocalisation with kinetochore NDC80 and ARK2 markers potentially supports a dual role in spindle organisation, kinetochore segregation, and spindle pole integrity—reminiscent of Aurora B, which localises to centromeres, the central spindle, and the spindle midzone[5, 6, 34, 35].

Functional genetic studies in *P. falciparum* and *P. berghei* had indicated that ARK1 is likely essential in asexual blood stages[13, 14]. Attempts to generate conditional knockdowns using the DD/Shield system in *P. falciparum* were unsuccessful, as reported recently[19]. Here, using the DiCre-Rapamycin system for *P. falciparum*[21] and promoter trap strategy for *P. berghei*[22], we demonstrate that ARK1 depletion leads to defects in spindle formation and karyokinesis. In *P. falciparum*, ARK1 is required for proper spindle elongation, which is critical for karyokinesis.

As a result of defects in these processes in PfARK1-depleted parasites there is an impact on
subsequent cytokinesis. As a result, parasite growth and dispersal of individual merozoites
after egress from RBCs is impaired upon PfARK1 depletion. In *Toxoplasma*, overexpression
of a kinase dead mutant of TgARK1 disrupts chromosome segregation and MTOC
maturation, similar to some of the division defects observed here in asexual stages, although
the mechanism was unexplored[20, 33].

During male gametogony, ARK1 knockdown led to defects in the spindle apparatus and the
failure of MTOC separation, suggesting its associated function with the outer basal body[4].
This was accompanied by abnormal axoneme formation and very few flagellated male
gametes, indicating a failure in MTOC-basal body segregation. These results suggest that
while ARK1 is part of the inner MTOC, its functional association with the outer MTOC is
essential for successful mitosis and gamete formation. This phenotype is reminiscent of
defects reported for other inner MTOC proteins, where the physical connection between the
inner and outer plaque components is destabilised[36]. These features also show some
resemblance to those of Aurora B in many model systems[5, 6, 34, 35].

Our protein interactome analysis reveals that ARK1 is part of a non-canonical chromosomal
passenger complex (CPC), comprising two divergent INCENP homologues and lacking the
canonical chromatin-targeting components Borealin and Survivin. Given that the distinct
localisation dynamics of ARK1 suggest that it may not act as the classic passenger complex
on chromosomes, it is possible that the chromatin-directing function of Survivin and Borealin
has become obsolete for apicomplexans and that the structural motifs for localisation and
activation of ARK1 by INCENP-A and -B might be revealed upon detailed comparative
sequence analyses. This is consistent with earlier molecular cell biological studies in
Toxoplasma[20]. While studies in *Toxoplasma* also reported two INCENPs, they did not detect
the presence or absence of other CPC components. The lack of chromatin-targeting
components might suggest that INCENP-A/-B can facilitate such interactions themselves or
simply mean these are lost and replaced by interactions of different proteins. For instance,
INCENPs in apicomplexans may form alternative CPC complexes, similar to *Trypanosoma*,
where INCENP associates with kinesin-1A/B[37]. A potential candidate kinesin might be
kinesin-8X, which shows a similar localisation pattern during different life stages as ARK1[38].
Interestingly, plants and heterokont algae also harbour divergent Survivin-like proteins, with
a central conserved but highly divergent helix suggesting a possible broader evolutionary

divergence of CPC components across eukaryotes[39]. Future studies will need to include a

further scrutiny of INCENP-A/-B localisation and interactomes to determine whether any

new CPC components might be present in apicomplexans.

INCENP duplication into INCENP-A and INCENP-B likely arose in parallel to the

duplication observed for *Toxoplasma*. Our attempted phylogenetic analyses do not provide

evidence to support a clear duplication of INCENPs in the ancestor of both Hematozoa and

Coccidia, suggesting that parallel duplications must have occurred in different lineages.

Remarkably such recurrent duplications can be observed throughout apicomplexans and

consistently coincide with the loss of Survivin (Fig. 7). Notably, INCENP-B has been

secondarily lost in piroplasmids, which also lack ARK2 and ARK3, hinting at functional

coupling between Aurora paralogues and specific INCENP variants. or pointing towards a

loss of meiosis-specific regulators as Roques et al. find that INCENP-B is specific to meiotic

stages[31]. Structural predictions using AlphaFold3 suggest that apicomplexan INCENPs

contain extended IN-box regions that fold around the ARK1 kinase domain and feature a

conserved CxC motif absent from other eukaryotic homologues. At the same time, ARK1

itself shows a divergent activation loop (GxSH instead of the canonical RxT), and INCENPs

lack the canonical transphosphorylation site (TSS) site found in other eukaryotes. Such

changes might indicate that other kinases are necessary to activate ARK1. Together, these

features indicate that apicomplexans have likely rewired CPC architecture and regulation,

replacing canonical chromatin-targeting logic with INCENP-centric scaffolds that localise

ARK1 at the inner MTOC, spindle and nucleoplasm.

While the function of the other two ARKs-ARK2 and ARK3- is unclear, at least in asexual

stages, our recent studies on PbARK2 have provided insights into its role in sexual

transmission stages. While ARK2 regulates spindle dynamics during male gametogony just

like ARK1 in the present studies, surprisingly, its deletion did not impact exflagellation,

unlike that of the ARK1 mutant (Fig. 5b). Previous studies also revealed that PbARK2

deletion caused an increase in the expression of ARK1 and ARK3[10]. Therefore, in the light of

the present work, it is reasonable to suggest that ARK1 and ARK2 may have some overlap of

function during male gametogony, particularly in spindle biogenesis. It was also interesting to

note that the deletion of PfARK1 resulted in higher PfARK2 levels in schizonts

(Supplementary Fig. 1f). While this increase in PfARK2 may be insufficient to fully

complement the PfARK1 mutant, it is possible that it may compensate for some of the defects

in the growth of PfARK1-depleted parasites. While there may be some overlap in ARK1 and

ARK2 function, our results confirm that ARK1 is indispensable for parasite survival, while

ARK2 has been previously reported to be essential for asexual development[13, 14, 40]. Recent

study also substantiates that inhibition of PfARK1 results in mitotic defects and microtubule

disorganisation during schizogony[16].

Overall, this study suggests that *Plasmodium* ARK1 regulates spindle elongation during

mitotic division and associates at the inner nuclear pole of the MTOC. However, its

association with the outer MTOC is essential for karyokinesis and for flagellated gamete

formation. ARK1 appears to represent a unique Aurora kinase paralogue that, while divergent

in structure, shares functional similarities with Aurora B in humans. This suggests that

Plasmodium has evolved an atypical mechanism of mitosis, where many of the canonical

molecules are either missing or highly divergent, reflecting the parasite's adaptation to

complex life cycle transitions.

Methods

Ethics statement

All animal procedures were approved following an ethical review process and conducted in

accordance with the United Kingdom Animals (Scientific Procedures) Act 1986, under Home

Office Project Licenses (PDD2D5182 and PP3589958). Female CD1 outbred mice (6 to 8

weeks old) were obtained from Charles River Laboratories and used for all experiments

carried out in the UK. The conditions of mice kept are a 12 hour light and 12 hour dark (7 till

7) light cycle, the room temperature is kept between 20-24 degrees Celsius and the humidity

is kept between 40-60%.

P. falciparum **cultures**

Plasmodium falciparum strain 1G5DC was obtained from European Malaria Reagent

Repository (EMRR) and DSM1 was obtained from *BEI* resources, Malaria Research and

Reference Reagents Resource Centre (MR4), American Type Culture Collection (ATCC).

Parasites were maintained in O+ human erythrocytes (5% hematocrit) in RPMI-1640

supplemented with 0.5% Albumax II and 50 μg/mL hypoxanthine[41]. Cultures were

maintained at 37°C under a mixture of 5% CO_2, 3% O_2, and 91.8% N_2 or 5% CO_2. Fresh

erythrocytes and culture medium were used to dilute parasites to maintain 3–5% parasitaemia

and 5% haematocrit. Relevant drugs were used during the culture of various transgenic lines,

as described below. Parasite synchronisation was carried out using 5% sorbitol as described

previously[42].

Generation of transgenic parasite lines

556 All PCR primers used in this study were synthesized by Sigma and are indicated in

557 Supplementary Data 3.

1G5DC SkipFlox

559 A dimerisable Cre recombinase (diCre) expressing parasite line was generated by transfecting

560 pSkipFlox plasmid, a kind gift from Tobias Spielmann (Addgene Plasmid #85797) into the

561 parent line 1G5DC[21]. Parasites were selected in the presence of 2.5 µg/ml of blasticidin.

562 For simplicity of representation, this parasite line is referred to as 1G5DC in the text and

563 figures.

HA-PfARK1-loxP To generate a conditional knockout transgenic parasite, a selection-linked

integration (SLI) approach was used. To generate the pSLI-HA-ARK1- loxP construct, a 592

bp homology region corresponding to the 5'-end of the PfARK1 gene was amplified using

primers (P1 and P2) derived from 3D7 genomic DNA. The PCR product was cloned

into *NotI* and *PmeI* sites upstream of loxP and a Myc-tag coding sequence in the pSLI-N-

sandwich-loxP (K13) vector, which was a gift from Tobias Spielmann (Addgene Plasmid

#85793). A recodonised sequence version of full-length PfARK1 with a N-terminal 3xHA tag

and followed by a loxP site was custom synthesised as a G-block (Genscript). The G-block

was cloned in the above construct downstream of yDHODH using *NheI* and *XhoI* restriction

sites.

***P. falciparum* transfection**

The pSLI-HA-PfARK1-loxP construct was transfected in 1G5DC SkipFlox parasites at the

ring stage using ~100µg of purified plasmid DNA. Transgenic parasites were selected by

using 2nM WR99210 and integrants were then enriched using 1.5µM DSM-1. The parasites

that appeared after drug selection were genotyped using PCR primers (Supplementary Data 3

and Supplementary Fig. 1a). The desired integration was confirmed by sequencing of PCR

products.

Conditional knockout of PfARK1

HA-PfARK1-loxP parasites were treated with 250 nM rapamycin at ring stage for 24 hours.
The genomic DNA was isolated from DMSO/RAP-treated parasites and excision of the
PfARK1 locus flanked by loxP sites in the transgenic parasite was confirmed by PCR using
primers (P7/P8).

P. falciparum growth rate assays

HA-PfARK1-loxP parasites were cultured in the presence of 10 nM WR99210, 2.5 µg/ml
blasticidin and 1.5µM DSM-1. Cultures were maintained at 2% hematocrit and 7-8%
parasitaemia. To perform the growth rate assay, 1G5DC-SkipFlox and HA-PfARK1-LoxP
parasites were synchronised at ring stage for two cycles using 5% sorbitol for 20 min at 37°C,
and the rings were seeded at 0.5% parasitaemia and 2% hematocrit. To deplete PfARK1,
parasites were treated with either DMSO (vehicle control) or 250 nM rapamycin for 24 h.
Thin Giemsa-stained blood smears were examined periodically under a microscope to
quantify the parasitaemia at different parasite stages. For assessment of parasitaemia by flow
cytometry, cells were fixed by incubation in a solution containing 1% paraformaldehyde and
0.0075% glutaraldehyde at ambient temperature for 15 min on an end-to-end rocker. Post-
fixation, samples were either stored at 4°C or stained immediately with Hoechst 33342 dye at
37°C for 10 min. After staining, samples were washed with FACS buffer and then analysed
on either BDverse (BD Biosciences) or BD FACSymphony A1 Flow Cytometers, acquiring
between 10,000 and 100,000 events per sample. Data analysis was carried out using FlowJo
software version 10.10.0.

Purification of *P. berghei* schizonts and gametocytes

Blood-stage parasites were obtained from infected mice on day 4 post-infection and cultured
at 37 °C with gentle rotation (100 rpm) for 11 or 24 hours. Schizonts were purified the
following day using a 60% v/v NycoDenz gradient prepared in phosphate-buffered saline
(PBS). The NycoDenz stock solution consisted of 27.6% w/v NycoDenz in 5 mM Tris-HCl
(pH 7.2), 3 mM KCl, and 0.3 mM EDTA.

Gametocyte purification was performed by first infecting phenylhydrazine-treated mice[43],
followed by sulfadiazine treatment two days post-infection to enrich for gametocytes. Blood
was collected on day 4 post-infection, and gametocyte-infected cells were purified using a
48% v/v NycoDenz gradient in PBS, prepared from the same NycoDenz stock solution.
Gametocytes were collected from the interface and subsequently activated for downstream
assays[44].

Generation of *P. berghei* transgenic parasites

GFP-tagging constructs were generated using the p277 plasmid vector and transfected into parasites as previously described[45]. A schematic representation of the wild-type *ark1* locus, the GFP-tagging constructs, and the recombined *ark1* locus is shown in Supplementary Fig. 2a. Correct integration of the *gfp* sequence at the *ark1* locus was confirmed by diagnostic PCR using primer 1 (intArk1tg) and primer 2 (ol492), as indicated in Supplementary Fig. 2b. The expression of native ARK1-GFP was confirmed by Western blot analysis (Supplementary Fig. 2c).

To investigate the function of ARK1, a promoter trap strategy was employed using double homologous recombination to generate a conditional knockdown line (*ark1PTD*). The knockdown construct was derived from the P*ama1* plasmid (pSS368), in which *ark1* was placed under the control of the *ama1* promoter, as previously described[22]. A schematic representation of the endogenous *ark1* locus, the targeting constructs, and the recombined *ark1* locus is shown in Supplementary Fig. 6a. Diagnostic PCR was performed to confirm successful integration, as outlined in Supplementary Fig. 6a. Integration at the 5′ locus was verified using primers intPTD31_5 (Primer 1) and 5′-intPTD (Primer 2), while integration at the 3′ end was confirmed using primers intPTD31_3 (Primer 3) and 3′-intPTama1 (Primer 4) (Supplementary Fig. 6b). Primer sequences are listed in Supplementary Data 3. Transfections were carried out by electroporation using *P. berghei* ANKA line 2.34 for GFP-tagging experiments and ANKA line 507cl1 expressing GFP for generation of the knockdown line[30].

Generation of dual tagged parasite lines (*P. berghei*)

To generate dual-labelled parasites, ARK1-GFP parasites were mixed in equal proportions with either NDC80-mCherry, ARK2-mCherry, or kinesin-8B-mCherry labelled parasites and co-injected into mice. Four to five days post-infection, when gametocytaemia was high, Anopheles mosquitoes were allowed to feed on the infected mice. Mosquitoes were examined for oocyst development and sporozoite formation at days 14 and 21 post-feeding. Infected mosquitoes were then used to infect naïve mice via mosquito bite-back. Blood-stage infection was assessed 4–5 days later by Giemsa-stained blood smear microscopy. In this way, parasites expressing both ARK1-GFP and either NDC80-mCherry, ARK2-mCherry, or kinesin-8B-mCherry were obtained. Gametocytes from these mixed lines were purified, and fluorescence microscopy was performed as described below to examine protein co-localisation.

P. berghei phenotype analyses

Infections were initiated by intraperitoneal injection of approximately 50,000 *P. berghei* parasites from WT-GFP or *ark1PTD* lines into mice. Asexual stages and gametocyte production were monitored on Giemsa-stained thin smears. Four to five days post-infection, exflagellation and ookinete conversion were assessed using a Zeiss AxioImager M2 microscope with an AxioCam ICc1 digital camera. For mosquito transmission assays, 30–50 *Anopheles stephensi* (SD 500) mosquitoes were allowed to feed for 20 min on anaesthetised, infected mice with ~15% asexual parasitaemia and comparable gametocyte levels (as determined by Giemsa-stained smears). To evaluate midgut infection, ~15 mosquito guts were dissected on day 7, 14 and 21 post-feeding, and oocysts were counted using a 63× oil immersion objective on the same microscope setup. Sporozoites in oocysts were counted on day 14 and day 21 post feeding. Salivary glands were isolated on day 21 to count the sporozoites. Bite-back experiments were performed 21 days post-feeding using naïve mice, and blood smears were examined after 3–4 days. All experiments were conducted at least three times to assess the phenotype.

Immunofluorescence Assay (IFAs)

Immunofluorescence assays (IFA) were conducted on thin blood smears following a previously described protocol[46]. In brief, air-dried smears were fixed for 2 min using ice cold 1:1 mixture of methanol and acetone. The smears were dried to remove any traces of methanol/acetone followed by blocking with 3% BSA for 45 minutes at room temperature. Samples were incubated with primary antibodies for 12 h at 4°C. After thorough washing with 1X PBS, Alexa Fluor-conjugated secondary antibodies were added and incubated for 2 h at room temperature. Smears were then mounted with VectaShield medium (Vector Laboratories Inc.) containing DAPI for nuclear staining. Microscopy was performed using a LSM980 AR confocal microscope (Carl Zeiss) or Zeiss Axio Observer microscope and image processing was carried out using Zeiss ZEN Blue (versions 3.1 or 3.9). Representative z-stack images were selected for figure preparation unless indicated otherwise. IFA or U-ExM (described below) with MSP1 or GAP45 antibodies was performed to assess segmentation in schizonts that had been treated with 10 μM E64 at ~42 hpi (cycle 1) for 4-5 hours to prevent egress and allow complete maturation.

P. falciparum live-cell imaging

Live imaging was carried out with minor modifications to previously described protocols[28].

HA-PfARK1-loxP parasites were synchronised at the ring stage and treated with either

DMSO or 250 nM rapamycin for 24 hours. Schizonts (cycle 1) were subsequently purified

and co-cultured with fresh erythrocytes at 2% hematocrit in RPMI1640 complete medium in

the presence of 25 nM PKG inhibitor ML10 [26, 27](a gift from LifeArc). ML10 was removed

after 4-6 hours and parasites were plated onto μ-slide, 2-well ibiTreat chambers (ibidi, 80286)

pre-coated with 0.5 mg/mL concanavalin A. The slides were maintained at 37 °C in a

humidified chamber with 5% CO_2. Imaging was performed using a Zeiss Axio Observer

microscope, capturing frames every 3 seconds for a minimum duration of 30 min and the

acquired time-lapse videos were processed and analysed using Axio Vision software (version

4.8.2) or ZEN blue 3.1 or 3.9.

P. berghei live cell imaging

To examine PbARK1-GFP expression during erythrocytic development, parasites cultured in

schizont medium were imaged at various stages, including ring, trophozoite, schizont, and

merozoite. Purified gametocytes were assessed for GFP expression and localisation at

multiple time points (0 and 1–15 min) following activation in ookinete medium. For nuclear

staining, Hoechst dye was used during live-cell imaging. Zygote and ookinete stages were

analysed over a 24-hour culture period. Images were acquired using a Zeiss Axio Imager M2

microscope equipped with a 63× oil immersion objective and an AxioCam ICc1 digital

camera (Carl Zeiss) using autoexposure settings. Fluorescence signals were adjusted using

Axiovision (Rel. 4.8) software to reduce background fluorescence while maintaining

detection sensitivity.

P. berghei structured illumination microscopy

Formaldehyde-fixed (4%) gametocytes were stained with Hoechst dye, and 2 μl of the cell

suspension was placed on a microscope slide and covered with a 50 × 34 mm coverslip to

form a thin, immobilised cell monolayer. Imaging was performed using structured

illumination microscopy (SIM) on a Zeiss Elyra PS.1 microscope with either a Zeiss Plan-

Apochromat 63×/1.4 oil immersion or Zeiss C-Apochromat 63×/1.2 W Korr M27 water

immersion objective, as previously described[36].

Ultrastructure expansion microscopy

U-ExM was performed largely following previously published protocols[23, 47, 48, 49]. In brief, to

prepare U-ExM samples of *P. falciparum* schizonts, 12mm coverslips were placed in a 24-

well plate and coated with poly-D-lysine (Gibco, A3890401, 0.1 mg/mL) for 1 h at 37 °C,

followed by three washes with ultrapure water. *P. falciparum* schizonts were seeded onto the

coverslips and incubated for 30 min at 37 °C. Unattached cells were removed gently, and

the remaining cells were fixed using 4% (v/v) formaldehyde in PBS for 15 min at 37 °C.

After fixation, coverslips were rinsed with prewarmed PBS and incubated for 12h at 37 °C

in 1.4% (v/v) formaldehyde and 2% (v/v) acrylamide in PBS. A monomer solution consisting

of 19% (w/w) sodium acrylate (Sigma-Aldrich, 408220), 10% (v/v) acrylamide (Sigma-

Aldrich, A3553), and 2% (v/v) N,N'-methylenebisacrylamide (Sigma-Aldrich, M7279) in

PBS was prepared 24 h in advance and stored at −20 °C. For polymerisation, 5 μL of

10% (v/v) TEMED (Sigma-Aldrich, T9281) and 5 μL of 10% (w/v) APS (Sigma-Aldrich,

A3678) were mixed with 90 μL of the thawed monomer solution. The gel mix (35 μL) was

pipetted onto parafilm squares pre-cooled at −20 °C and the coverslip was placed over the

drop. Gels were allowed to polymerise for 1 h at 37 °C. Polymerised gels were transferred

to denaturation buffer (200 mM SDS, 200 mM NaCl, 50 mM Tris, pH 9) and shaken for

15 min at room temperature, then incubated at 95 °C for 90 min. Denatured gels were

expanded by sequential incubations in water (30 min each, repeated thrice) with gentle

shaking. Expanded gels were equilibrated in PBS for 15 min and blocked in 3% BSA/PBS

for 30 min at room temperature. Incubation with primary antibody (details in Supplementary

Data 4) was carried out for 12h at room temperature on a shaker, followed by three 10 min

washes with PBST. Gels were then incubated with secondary antibodies, 405 NHS Ester

(Thermo Fisher, A30000) and SYTOX/Hoechst in PBS for 2.5 h in the dark with gentle

shaking followed by three PBST washes and final equilibration in MilliQ water (30 min,

repeated thrice).

For imaging, μ-slide 2-well ibiTreat chambers (ibidi, 80286) were coated with poly-D-lysine

(0.5 mL, 1 h at 37 °C), rinsed with MilliQ water, and used to mount the gels cell side

down. Imaging was performed using a Zeiss LSM980 confocal microscope with a 63X oil

immersion objective (NA 1.4). Z-stacks (0.5 μm intervals) were captured and processed

using ZEN Blue (versions 3.1 or 3.9). Deconvolution was performed using Zen

Deconvolution tool kit. U-ExM images are displayed as maximum intensity projections of

full z-slices unless indicated otherwise.

To prepare U-ExM samples of *P. berghei* male gametocytes, purified and 8 min- and 15 min-
activated gametocytes were fixed in 4% formaldehyde in PHEM buffer (60 mM PIPES, 25
mM HEPES, 10 mM EGTA, 2 mM MgCl$_2$, pH6.9) at room temperature for 15 min. Fixed
samples were attached to 10 mm round Poly-D-Lysine coated coverslips for 15 min.
Coverslips were incubated overnight at 4°C in 1.4% formaldehyde (FA)/ 2% acrylamide
(AA). Gelation was performed in ammonium persulfate (APS)/TEMED (10%
each)/monomer solution (23% sodium acrylate; 10% AA; 0,1% BIS-AA in PBS) on ice for 5
min and at 37°C for 30 min. Gels were denatured for 15 min at 37°C and for 90 min at 95°C
in denaturation buffer (200 mM SDS, 200 mM NaCl, 50 mM Tris, pH 9.0, in water). After
denaturation, gels were incubated in distilled water overnight for complete expansion. The
following day, circular gel pieces with a diameter of ~13 mm were excised, and the gels were
washed in PBS three times for 15 min to remove excess water. The gels were then incubated
in blocking buffer (3% BSA in PBS) at room temperature for 30 min, incubated with mouse
monoclonal anti-a-tubulin antibody (T9026, Sigma-Aldrich, 1:500 dilution) and/or anti-GFP
antibody (A11122, Invitrogen, 1:500 dilution) in blocking buffer (1:500 dilution) at 4°C
overnight and washed three times for 15 min in wash buffer (0.5% v/v TWEEN-20 in PBS).
The gels were incubated with 8 μg/ml Atto 594 NHS ester (08741, Merck), 10 μg/ml Hoechst
33342 (H1399, Invitrogen), Alexa Fluor 488 goat anti-mouse IgG (A11001, Invitrogen, 1:500
dilution) and Alexa Fluor 568 goat anti-mouse IgG (A11004, Invitrogen, 1:500 dilution) in
PBS at 37°C for 3 hours followed by three washes of 15 min each in wash buffer (blocking
and all antibody incubation steps were performed with gentle shaking). The gels were then
washed three times for 15 min with wash buffer and expanded overnight in ultrapure water.
The expanded gel was placed in a 35 mm glass bottom dish (MATTEK) with the 14 mm glass
coated with Poly-D-Lysine and mounted with an 18 × 18 mm coverslip to prevent the gel
from sliding and to avoid drifting while imaging. High-resolution confocal microscopy
images were acquired using a Zeiss Celldiscoverer 7 with Airyscan using a Plan-Apochromat
50 ×/1.2NA Water objective, with 405, 488 and 561 nm lasers. Confocal z-stacks were
acquired using line scanning and the following settings: 55 × 55 nm pixel size, 170 nm
z-step, 2.91 μs/pixel dwell time, 850 gain and 3.5% (405 nm), 4.5% (488 nm) and 5.0%
(561 nm) laser powers. The z-stack images were processed and analysed using Fiji (Version
1.54f)[50].

Calculation of the expansion factor for the U-ExM of male gametocytes

The expansion factor for the U-ExM of male gametocytes was calculated by comparing
axoneme diameters from two types of images (U-ExM and TEM). The diameters were
measured using Fiji (Version 1.54f)[50]. The average diameter of the axoneme in NHS-stained
U-ExM images of male gametocytes activated for 8 min was 933.83 nm (n = 12), while the
average diameter in TEM images of male gametocytes activated for 8 min was 188.50 nm (n
= 8). We used the calculated expansion factor of 4.954 for the analysis.

Electron microscopy of male gametocytes

P. berghei gametocytes activated for 8 min and 15 min were fixed in 3% glutaraldehyde in
0.1 M cacodylate buffer and processed for electron microscopy[51]. Briefly, samples were post-
fixed in 1% aqueous osmium tetroxide, treated en bloc with 2% aqueous uranyl acetate,
dehydrated in graded ethanol series and embedded in Spurr's epoxy resin. Thin sections were
stained with lead citrate prior to examination in a Tecnai G2 12 BioTwin (FEI UK, UK) or a
Jeol JEM-1400Flash (JEOL, Japan) electron microscope.

Liver stage parasite imaging

For *P. berghei* liver-stage imaging, 100,000 Hepatocelluar carcinoma cells (HepG2) were
seeded onto glass coverslips in 48 well plate. Salivary glands from *Anopheles stephensi*
mosquitoes infected with PbARK1-GFP parasites were dissected and homogenised using a
pestle to release sporozoites. The sporozoites were gently pipetted onto the HepG2 cell
monolayer and incubated at 37 °C with 5% CO_2 in complete minimum Eagle's medium
The culture medium was replaced 2 hours post-infection and subsequently changed daily. For
live-cell imaging, Hoechst 33342 (Molecular Probes) was added to a final concentration of
1 μg/ml to stain host and parasite nuclei. Infected cells were imaged at, 48-, and 64 -hours
post-infection using a Zeiss Axio Imager M2 microscope equipped with a 63× oil immersion
objective and an AxioCam ICc1 digital camera (Carl Zeiss) using autoexposure settings. and
Leica Application Suite X software. Each experiment was performed in triplicate, and 20–30
infected cells were analysed per time point to assess PbARK1-GFP localisation.

Immunoblotting

P. falciparum - Immunoblotting was performed as described previously[52] using primary
antibodies (Supplementary Data 4) and HRP-conjugated secondary antibodies and protein
bands were visualized using either SuperSignal™ West Pico or Femto chemiluminescent
substrates (Pierce, USA) using X-ray films.

P. berghei - Purified gametocytes were lysed in buffer containing 10 mM Tris-HCl (pH 7.5),

150 mM NaCl, 0.5 mM EDTA, and 1% NP-40. After lysis, Laemmli buffer was added, and

samples were boiled for 10 min at 95 °C, followed by centrifugation at 13,000 × g for 5

min. The supernatants were separated on a 4–12% SDS-polyacrylamide gel and transferred

onto a nitrocellulose membrane (Amersham Biosciences). Immunoblotting was performed

using the Western Breeze Chemiluminescence Anti-Rabbit Kit (Invitrogen), following the

manufacturer's protocol. An anti-GFP polyclonal antibody (Invitrogen) was used at a 1:1,250

dilution to detect PbARK1-GFP.

Quantitative Real-Time PCR (qRT-PCR) analyses

P. falciparum - Total RNA was extracted using TRIzol reagent (G Biosciences) followed by

purification with the Qiagen RNeasy Mini Kit (Cat. No. 74104). cDNA synthesis was

performed using RevertAid H Minus Reverse Transcriptase (Thermo scientific, EP0451) and

1 µg of DNase-treated RNA as the template. Quantitative PCR was conducted using the

CFX96 Real-Time PCR Detection System (Bio-Rad). Reactions were normalized using 18S

rRNA as the internal control. Relative gene expression levels were calculated using the $2^{-\Delta\Delta Ct}$

method. Primer sequences used for the qRT-PCR are listed in Supplementary Data 3.

P. berghei - Total RNA was extracted from gametocytes using the RNA Purification Kit

(Stratagene), and cDNA was synthesised using the RNA-to-cDNA Kit (Applied Biosystems).

qRT-PCR was performed with cDNA from 80 ng of RNA using SYBR Green Fast Master

Mix (Applied Biosystems) on an Applied Biosystems 7500 Fast system. Cycling conditions

were: 95 °C for 20 s, followed by 40 cycles of 95 °C for 3 s and 60 °C for 30 s.

Primers were designed using Primer3 (https://primer3.ut.ee/). Each gene was tested in three

biological and technical replicates. *hsp70* (PBANKA_081890) and *arginyl-tRNA synthetase*

(PBANKA_143420) were used as reference genes. Primer sequences are listed in

Supplementary Data 3.

Transcriptome analysis using RNA-seq

Total RNA was extracted from activated gametocytes and schizonts of *P. berghei* WT-GFP

and *ark1PTD* parasites (three biological replicates each) using the RNeasy Kit (Qiagen),

vacuum-concentrated, and transported in RNA-stable tubes (Biomatrica). Strand-specific

mRNA libraries were prepared with the TruSeq Stranded mRNA Sample Prep Kit (Illumina)

and sequenced on an Illumina HiSeq 4000 platform (paired-end, 150 bp reads). Read quality

was assessed with FASTQC, and trimming of low-quality reads and adapter sequences was

done using Trimmomatic[53]. Reads were mapped to the *P. berghei* ANKA genome (PlasmoDB

release 40) using HISAT2 v2.1.0[54], and gene counts were generated with FeatureCounts[55].

Low-expression genes (CPM < 1) were excluded. Data were normalized using the TMM

method (EdgeR[56]), transformed with voom (limma[57]), and analysed for differential

expression using DESeq2[58]. Statistical significance was determined using a two-sided Wald

test, with *P*-values adjusted for multiple comparisons using the Benjamini-Hochberg

procedure. Genes with fold-change > 2 and FDR < 0.05 were considered differentially

expressed.

842 Immunoprecipitation and mass spectrometry

P. berghei purified schizonts and gametocytes (activated for 1.5 to 2 min) were crosslinked

using formaldehyde (10-min incubation with 1% formaldehyde, followed by 5-min

incubation in 0.125M glycine solution and 3 washes with PBS (pH 7.5). Immunoprecipitation

was performed using crosslinked protein and a GFP-Trap_A Kit (Chromotek) following the

manufacturer's instructions. Proteins bound to the GFP-Trap_A beads were digested using

trypsin and the peptides were analysed by LC-MS/MS and subsequently visualised using both

Principal Component Analysis (PCA) and for relative ratios of WT/POI using a volcano plot

as published previously[36, 59]. In short, significance of enrichment was assessed using the two-

tailed Student t-test (implemented in R), and p-values were corrected for multiple testing

using the Benjamini–Hochberg method. iBAQ values were normalized by the sum of all

iBAQ values over one experiment. Only those proteins were considered with more than 2

unique peptides and at least 2 values in ARK1 pulldowns. Volcano plots were visualised

using VolcaNoseR[60].

856 Evolutionary bioinformatics

Our predicted proteome set was based on a version previously used (see Supplementary Data

5)[61]. Phyletic profiles for ARK1-3 and INCENP were partially derived from previous

studies[10, 62]. Sequence searches for missing orthologues were performed with the HHPred

webserver from MPI-toolkit[63, 64] and the hmmer package[63], as previously described[61]. For the

larger gene families like Aurora kinases, we used IQ-tree-based maximum-likelihood

phylogenetics (standard setting - model finding, 1000 bootstraps[65]) to check the validity of

bidirectional-best-blast (BBH) hits amongst our genome sets on an ad hoc basis. Off note, we

could not reliably generate a single phylogenetic tree harbouring all three ARKs within the

Aurora kinase orthologous group (OG), suggesting that ARK2 and ARK3 are either

extremely divergent or not Aurora kinases. Similarly, for the INCENP gene family, multiple

alignments were generally too divergent and phylogenetically informative stretches too short

to infer any consistent phylograms. All sequences used for this study can be found in

Supplementary Data 5.

AlphaFold3 modelling

Co-folds of 3D protein structures for PbARK1 and the extended IN-box of either

PbINCENP-A or -B were modelled using the AlphaFold3 webserver

(https://alphafoldserver.com/) with standard settings (seed set to 100)[66].

Data availability

The RNA-seq data generated in this study have been deposited in the NCBI Sequence Read

Archive under accession number PRJNA1309997

[https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1309997]. The mass spectrometry

proteomics data generated in this study have been deposited in the ProteomeXchange

Consortium via the PRIDE[67] partner repository under the dataset identifier PXD068821 and

10.6019/PXD068821 [https://www.ebi.ac.uk/pride/archive/projects/PXD068821]. All other

source data are provided with this paper.

References

1.　Guttery DS, Zeeshan M, Ferguson DJP, Holder AA, Tewari R. Division and
Transmission: Malaria Parasite Development in the Mosquito. *Annu Rev Microbiol*
76, 113-134 (2022).

888
889　2.　Sinden RE. Mitosis and meiosis in malarial parasites. *Acta Leiden* **60**, 19-27 (1991).

890
891　3.　Sinden RE, Canning EU, Bray RS, Smalley ME. Gametocyte and gamete
892　　　development in Plasmodium falciparum. *Proc R Soc Lond B Biol Sci* **201**, 375-399
893　　　(1978).

894
895　4.　Zeeshan M*, et al.* Plasmodium SAS4: basal body component of male cell which is
896　　　dispensable for parasite transmission. *Life Sci Alliance* **5**, (2022).

897
898　5.　Carmena M, Earnshaw WC, Glover DM. The Dawn of Aurora Kinase Research:
899　　　From Fly Genetics to the Clinic. *Front Cell Dev Biol* **3**, 73 (2015).

901 6.   Carmena M, Ruchaud S, Earnshaw WC. Making the Auroras glow: regulation of
902   Aurora A and B kinase function by interacting proteins. *Curr Opin Cell Biol* **21**, 796-
903   805 (2009).

904
905 7.   Hochegger H, Hegarat N, Pereira-Leal JB. Aurora at the pole and equator:
906   overlapping functions of Aurora kinases in the mitotic spindle. *Open Biol* **3**, 120185
907   (2013).

908
909 8.   Zhang X. Aurora kinases. *Curr Biol* **18**, R146-148 (2008).

910
911 9.   Li H, Chen Q, Kaller M, Nellen W, Graf R, De Lozanne A. Dictyostelium Aurora
912   kinase has properties of both Aurora A and Aurora B kinases. *Eukaryot Cell* **7**, 894-
913   905 (2008).

914
915 10. Zeeshan M*, et al.* Plasmodium ARK2 and EB1 drive unconventional spindle
916   dynamics, during chromosome segregation in sexual transmission stages. *Nat
917   Commun* **14**, 5652 (2023).

918
919 11. Vader G, Medema RH, Lens SM. The chromosomal passenger complex: guiding
920   Aurora-B through mitosis. *J Cell Biol* **173**, 833-837 (2006).

921
922 12. Hadders MA, Lens SMA. Changing places: Chromosomal Passenger Complex
923   relocation in early anaphase. *Trends Cell Biol* **32**, 165-176 (2022).

924
925 13. Solyakov L*, et al.* Global kinomic and phospho-proteomic analyses of the human
926   malaria parasite Plasmodium falciparum. *Nat Commun* **2**, 565 (2011).

927
928 14. Tewari R*, et al.* The systematic functional analysis of Plasmodium protein kinases
929   identifies essential regulators of mosquito transmission. *Cell Host Microbe* **8**, 377-387
930   (2010).

931
932 15. Zhang M*, et al.* Uncovering the essential genes of the human malaria parasite
933   Plasmodium falciparum by saturation mutagenesis. *Science* **360**, (2018).

934
935 16. Langeveld H*, et al.* Targeting Aurora Kinases as Essential Cell-Cycle Regulators to
936   Deliver Multi-Stage Antimalarials Against Plasmodium Falciparum. *Angew Chem Int
937   Ed Engl*, e202518493 (2025).

938
939 17. Wicht KJ*, et al.* The Human Chk1 Inhibitor CHIR-124 Shows Multistage Activity
940   Against Plasmodium falciparum via Dual Inhibition of PfArk1 and Hemozoin
941   Formation. *bioRxiv*, (2025).

943   18.   Reininger L, Wilkes JM, Bourgade H, Miranda-Saavedra D, Doerig C. An essential
944   Aurora-related kinase transiently associates with spindle pole bodies during
945   Plasmodium falciparum erythrocytic schizogony. *Mol Microbiol* **79**, 205-221 (2011).

946
947   19.   Wyss M, Thommen BT, Kofler J, Carrington E, Brancucci NMB, Voss TS. The three
948   Plasmodium falciparum Aurora-related kinases display distinct temporal and spatial
949   associations with mitotic structures in asexual blood stage parasites and gametocytes.
950   *mSphere* **9**, e0046524 (2024).

951
952   20.   Berry L*, et al.* Toxoplasma gondii chromosomal passenger complex is essential for
953   the organization of a functional mitotic spindle: a prerequisite for productive
954   endodyogeny. *Cell Mol Life Sci* **75**, 4417-4443 (2018).

955
956   21.   Collins CR*, et al.* Robust inducible Cre recombinase activity in the human malaria
957   parasite Plasmodium falciparum enables efficient gene deletion within a single
958   asexual erythrocytic growth cycle. *Mol Microbiol* **88**, 687-701 (2013).

959
960   22.   Sebastian S*, et al.* A Plasmodium calcium-dependent protein kinase controls zygote
961   development and transmission by translationally activating repressed mRNAs. *Cell*
962   *Host Microbe* **12**, 9-19 (2012).

963
964   23.   Liffner B*, et al.* Atlas of Plasmodium falciparum intraerythrocytic development using
965   expansion microscopy. *Elife* **12**, (2023).

966
967   24.   Li J*, et al.* Disruption of Plasmodium falciparum kinetochore proteins destabilises the
968   nexus between the centrosome equivalent and the mitotic apparatus. *Nat Commun* **15**,
969   5794 (2024).

970
971   25.   Rudlaff RM, Kraemer S, Marshman J, Dvorin JD. Three-dimensional ultrastructure of
972   Plasmodium falciparum throughout cytokinesis. *PLoS Pathog* **16**, e1008587 (2020).

973
974   26.   Ressurreicao M*, et al.* Use of a highly specific kinase inhibitor for rapid, simple and
975   precise synchronization of Plasmodium falciparum and Plasmodium knowlesi asexual
976   blood-stage parasites. *PLoS One* **15**, e0235798 (2020).

977
978   27.   Baker DA*, et al.* A potent series targeting the malarial cGMP-dependent protein
979   kinase clears infection and blocks transmission. *Nat Commun* **8**, 430 (2017).

980
981   28.   Ekka R, Gupta A, Bhatnagar S, Malhotra P, Sharma P. Phosphorylation of Rhoptry
982   Protein RhopH3 Is Critical for Host Cell Invasion by the Malaria Parasite. *mBio* **11**,
983   (2020).

984
985   29.   Silvie O*, et al.* A role for apical membrane antigen 1 during invasion of hepatocytes
986   by Plasmodium falciparum sporozoites. *J Biol Chem* **279**, 9490-9496 (2004).

30. Janse CJ, Ramesar J, Waters AP. High-efficiency transfection and drug selection of
genetically transformed blood stages of the rodent malaria parasite Plasmodium
berghei. *Nat Protoc* **1**, 346-356 (2006).

31. Roques M, Niu C, Brochet M, Brusini L. A modular chromosomal passenger complex
rewires chromosome segregation in Plasmodium berghei. *bioRxiv*, (2025).

32. Segura-Pena D*, et al.* The structural basis of the multi-step allosteric activation of
Aurora B kinase. *Elife* **12**, (2023).

33. Berry L*, et al.* The conserved apicomplexan Aurora kinase TgArk3 is involved in
endodyogeny, duplication rate and parasite virulence. *Cell Microbiol* **18**, 1106-1120
(2016).

1001

1002 34. Carmena M, Earnshaw WC. The cellular geography of aurora kinases. *Nat Rev Mol*
1003   *Cell Biol* **4**, 842-854 (2003).

1004

1005 35. Carmena M, Wheelock M, Funabiki H, Earnshaw WC. The chromosomal passenger
1006   complex (CPC): from easy rider to the godfather of mitosis. *Nat Rev Mol Cell Biol* **13**,
1007   789-803 (2012).

1008

1009 36. Zeeshan M*, et al.* Plasmodium NEK1 coordinates MTOC organisation and
1010   kinetochore attachment during rapid mitosis in male gamete formation. *PLoS Biol* **22**,
1011   e3002802 (2024).

1012

1013 37. Ballmer D, Akiyoshi B. Dynamic localization of the chromosomal passenger complex
1014   in trypanosomes is controlled by the orphan kinesins KIN-A and KIN-B. *Elife* **13**,
1015   (2024).

1016

1017 38. Zeeshan M*, et al.* Plasmodium kinesin-8X associates with mitotic spindles and is
1018   essential for oocyst development during parasite proliferation and transmission. *PLoS*
1019   *Pathog* **15**, e1008048 (2019).

1020

1021 39. Komaki S, Tromer EC, De Jaeger G, De Winne N, Heese M, Schnittger A. Molecular
1022   convergence by differential domain acquisition is a hallmark of chromosomal
1023   passenger complex evolution. *Proc Natl Acad Sci U S A* **119**, e2200108119 (2022).

1024

1025 40. Bushell E*, et al.* Functional Profiling of a Plasmodium Genome Reveals an
1026   Abundance of Essential Genes. *Cell* **170**, 260-272 e268 (2017).

1027

1028 41. Trager W, Jensen JB. Cultivation of erythrocytic stages. *Bull World Health Organ* **55**,
1029   363-365 (1977).

1030

1031 42. Lambros C, Vanderberg JP. Synchronization of Plasmodium falciparum erythrocytic
1032 stages in culture. *J Parasitol* **65**, 418-420 (1979).

1033

1034 43. Beetsma AL, van de Wiel TJ, Sauerwein RW, Eling WM. Plasmodium berghei
1035 ANKA: purification of large numbers of infectious gametocytes. *Exp Parasitol* **88**,
1036 69-72 (1998).

1037

1038 44. Billker O, Dechamps S, Tewari R, Wenig G, Franke-Fayard B, Brinkmann V.
1039 Calcium and a calcium-dependent protein kinase regulate gamete formation and
1040 mosquito transmission in a malaria parasite. *Cell* **117**, 503-514 (2004).

1041

1042 45. Guttery DS*, et al.* Genome-wide functional analysis of Plasmodium protein
1043 phosphatases reveals key regulators of parasite development and differentiation. *Cell*
1044 *Host Microbe* **16**, 128-140 (2014).

1045

1046 46. Tonkin CJ*, et al.* Localization of organellar proteins in Plasmodium falciparum using
1047 a novel set of transfection vectors and a new immunofluorescence fixation method.
1048 *Mol Biochem Parasitol* **137**, 13-21 (2004).

1049

1050 47. Rashpa R, Brochet M. Expansion microscopy of Plasmodium gametocytes reveals the
1051 molecular architecture of a bipartite microtubule organisation centre coordinating
1052 mitosis with axoneme assembly. *PLoS Pathog* **18**, e1010223 (2022).

1053

1054 48. Bertiaux E*, et al.* Expansion microscopy provides new insights into the cytoskeleton
1055 of malaria parasites including the conservation of a conoid. *PLoS Biol* **19**, e3001020
1056 (2021).

1057

1058 49. Yanase R*, et al.* Divergent Plasmodium kinases drive MTOC, kinetochore and
1059 axoneme organisation in male gametogenesis. *Life Sci Alliance* **8**, (2025).

1060

1061 50. Schindelin J*, et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
1062 *Methods* **9**, 676-682 (2012).

1063

1064 51. Ferguson DJ*, et al.* Maternal inheritance and stage-specific variation of the apicoplast
1065 in Toxoplasma gondii during development in the intermediate and definitive host.
1066 *Eukaryot Cell* **4**, 814-826 (2005).

1067

1068 52. Rawat RS*, et al.* Protein kinase PfPK2 mediated signalling is critical for host
1069 erythrocyte invasion by malaria parasite. *PLoS Pathog* **19**, e1011770 (2023).

1070

1071 53. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
1072 sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

1073

1074   54.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
1075   requirements. *Nat Methods* **12**, 357-360 (2015).

1076

1077   55.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
1078   assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

1079

1080   56.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
1081   differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
1082   139-140 (2010).

1083

1084   57.   Ritchie ME, *et al.* limma powers differential expression analyses for RNA-sequencing
1085   and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

1086

1087   58.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
1088   for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1089

1090   59.   Zeeshan M, *et al.* A novel SUN1-ALLAN complex coordinates segregation of the
1091   bipartite MTOC across the nuclear envelope during rapid closed mitosis in
1092   Plasmodium berghei. *Elife* **14**, (2025).

1093

1094   60.   Goedhart J, Luijsterburg MS. VolcaNoseR is a web app for creating, exploring,
1095   labeling and sharing volcano plots. *Sci Rep* **10**, 20560 (2020).

1096

1097   61.   Zeeshan M, *et al.* Real-time dynamics of Plasmodium NDC80 reveals unusual modes
1098   of chromosome segregation during parasite proliferation. *J Cell Sci* **134**, (2020).

1099

1100   62.   Tromer EC, van Hooff JJE, Kops G, Snel B. Mosaic origin of the eukaryotic
1101   kinetochore. *Proc Natl Acad Sci U S A* **116**, 12873-12882 (2019).

1102

1103   63.   Eddy SR. Accelerated Profile HMM Searches. *PLoS Comput Biol* **7**, e1002195
1104   (2011).

1105

1106   64.   Soding J. Protein homology detection by HMM-HMM comparison. *Bioinformatics*
1107   **21**, 951-960 (2005).

1108

1109   65.   Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ. W-IQ-TREE: a fast online
1110   phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res* **44**, W232-235
1111   (2016).

1112

1113   66.   Abramson J, *et al.* Accurate structure prediction of biomolecular interactions with
1114   AlphaFold 3. *Nature* **630**, 493-500 (2024).

1115

67.   Perez-Riverol Y, *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res* **53**, D543-D553 (2025).

68.   Zeeshan M, *et al.* Kinesin-8B controls basal body function and flagellum formation and is key to malaria transmission. *Life Sci Alliance* **2**, (2019).

Acknowledgement

RT is supported by ERC advance grant funded by UKRI Frontier Science (EP/X024776/1), MRC UK (MR/K011782/1), and BBSRC (BB/L013827/1, BB/X014681/1). MZ, DB, and AM were supported as research fellows by (EP/X024776/1). RY and DSG are supported by BBSRC (BB/X014681/1). SLP is supported by Wellcome DBT India Alliance/Team Science (IA/TSG/21/1/600261). AAH is supported by the Francis Crick Institute (FC001097), which receives core funding from the Cancer Research UK (FC001097), the UK Medical Research Council (FC001097), and the Wellcome Trust (FC001097). KGLR is supported by the NIH/NIAID (R01 AI136511) and the University of California, Riverside NIFA-Hatch-225935. ECT was supported by a personal fellowship from the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), the Netherlands (grant no. VI. Veni.202.223). Studies on *P. falciparum* were also supported by a Team Science grant (IA/TSG/21/1/600261) to PS, RT, and TSKP from DBT-Wellcome Trust India Alliance and funds from NII core. PS is a recipient of J.C. Bose Fellowship. We thank the School of Life Sciences Imaging (SLIM) for providing access to confocal and SIM microscopy and Dr Robert Markus for technical assistance. Most microscopy on *P. falciparum* was done in National Institute of Immunology (NII)-confocal microscopy facility. We thank the Nanoscale and Microscale Research Centre (nmRC) for providing access to instrumentation and Dr Michael W. Fey, Dr Julie Watts and Ms Nicola J. Weston for technical assistance. We are grateful to Dr Sabrina Absalon (Indiana University) for technical advice on U-ExM. For Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. We thank Cleidiane Zampronio at Warwick University for mass spectrometry methods and Bio Support Unit, University of Nottingham for maintenance of mice used in this study.

Author Contributions

RT, PS, and ECT conceptualised the project. AN and PS performed the *P. falciparum* experiments. RY, MZ, SLP, AM, ER, and RT performed the *P. berghei* live-cell imaging. RY, DJF, and SV performed the TEM observation. RY and MZ performed the *P. berghei* U-ExM observation. MZ, AE, and DB performed the qPCR experiment. SA and KGLR performed the RNAseq analysis. RT, DB, ARB, and ECT performed the proteomics analysis. ECT performed the evolutionary bioinformatics analysis. AN, RY, MZ, DJF, SLP, AM, ARB, SV, KGLR, DSG, AAH, ECT, PS, and RT performed formal analysis. AN, RY, MZ, DJF, SA, SLP, AM, ECT, and RT performed visualisation. RT, PS, and ECT acquired funding. RT, PS, and ECT supervised the project. AN, RY, MZ, AAH, ECT, PS, and RT wrote the manuscript draft. All the authors reviewed and edited the manuscript.

Competing Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Expansion of the Aurora kinase family among apicomplexan parasites coincides with divergent mitotic mechanisms.
a. The schematic shows schizogony in the asexual blood stage and male gametogony in *Plasmodium*, highlighting some of the subcellular structures involved (below). For schizogony: based on U-ExM classifications[23] - sequence of intranuclear MT structures: one hemispindle → bipolar mitotic spindle → interpolar microtubules (the outer plaque is closely associated with the IMC/plasma membrane[24]). **b.** Left: five different cellular locations of Aurora kinases during mitosis/meiosis[10], based on findings from model organisms – projected on the nucleus of *Plasmodium* spp. with a closed nucleus during mitosis and nuclear envelope-embedded MTOCs. The cytokinesis-associated region is shown only as a functional analogue based on ARK3 localisation in *Toxoplasma*[33], and does not imply a canonical central spindle or midbody in *Plasmodium*. Right: Overview of the location (pictograms) and putative protein complex/interactors (italicised text) of three Aurora kinase paralogues ARK1-2-3 in the three apicomplexan parasite models *Plasmodium, Toxoplasma and Cryptosporidium*.

Figure 2. Location of ARK1 during *P. falciparum* asexual division.

a. Schematic of asexual division during *P. falciparum* blood stage schizogony. HA-PfARK1-loxP parasites were synchronised and IFA was performed using anti-HA (green) and anti-centrin (magenta) antibodies on schizonts. DNA was stained with DAPI. PfARK1 was observed transiently as a punctum only at nuclei with duplicated centrin foci. Scale bar = 2 μm. **b.** Ultrastructure expansion microscopy (U-ExM) was performed on HA-PfARK1-loxP schizonts using anti-HA (green) and anti-tubulin (magenta) antibodies. Hemispindles (white arrow), mitotic spindles (cyan arrow) and interpolar spindles (yellow arrow) that correspond to different stages of nuclear division were observed. PfARK1-positive puncta were mainly associated with mitotic spindles with the puncta flanking the spindle (square and magnification in the inset). Scale bar = 2 μm. **c.** Percent nuclei that exhibited PfARK1 staining (panel a) were quantified (SEM ± SE, two-sided unpaired t-test, ***P = 0.0005, n = 37 parasites, three independent biological replicates. **d**. Nuclei exhibiting PfARK1 foci (panel b) in association with hemispindles, mitotic spindles or interpolar spindles were quantified and represented as % of total nuclei (SEM ± SE, one way ANOVA, ****P < 0.0001, n > 40 parasites, three independent biological replicates). Source data are provided as a Source Data file.

Figure 3. Location of ARK1 during *P. berghei* male gametogony.

In all images, PbARK1-GFP is shown in green and DNA stained with Hoechst dye in blue. **a.** Top panel: Schematic of male gametogony. Lower panel: Live cell imaging of PbARK1-GFP during male gametogony (1-8 min post activation). DIC: differential interference contrast. Scale bar: 5μm (n > 50 cells, 5 biological replicates). **b.** 3D-SIM of fixed male gametocytes at 1.5–2 min post activation (first spindle stage). Arrows indicate PbARK1-GFP foci. Scale bar = 1 μm (n > 10 cells, 5 biological replicates). **c.** U-ExM of male gametocytes at 1.5–2 min post activation (first spindle stage). PbARK1-GFP (green), α-tubulin (magenta), and NHS-ester (grey). Top panels: whole cell. Bottom panels: magnified views of the spindle (white dashed squares). Arrows: PbARK1-GFP foci at the spindle poles (or inner MTOCs). Scale bars = 5 μm (whole cell) and 2 μm (spindle) (n > 8 cells, 2 biological replicates). **d.** Live cell imaging of PbARK1-GFP and ARK2-mCherry[10] (magenta) during male gametogony (1-8 min post activation). Scale bar = 5 μm (n > 20 cells, 3 biological replicates). **e.** 3D-SIM of fixed male gametocytes at 1.5–2 min post activation (first spindle stage). PbARK1-GFP and ARK2-mCherry. Scale bar = 1 μm (n > 10 cells, 3 biological

replicates). **f.** Live cell imaging of PbARK1-GFP and NDC80-cherry[61] (magenta) during

male gametogony (1-8 min post activation). Scale bar = 5 μm (n > 20 cells, 3 biological

replicates). **g.** 3D-SIM of fixed male gametocytes at 1.5–2 min post activation (first spindle

stage). PbARK1-GFP and NDC80-mCherry. Scale bar = 1 μm (n > 10 cells, 5 biological

replicates). **h.** Live cell imaging of PbARK1-GFP and kinesin-8B-mCherry[68] (magenta)

dynamics during male gametogony (1-8 min post activation). Scale bar = 5 μm (n > 20 cells,

3 biological replicates). **i.** 3D-SIM of male gametocytes at 1.5–2 min post activation (first

spindle stage) of PbARK1-GFP and kinesin-8B-mCherry. Scale bar = 1 μm (n > 10 cells, 2

biological replicates). Note that the Hoechst signal appears weak in the 3D-SIM image

because the focal plane was optimised for the PbARK1-GFP signal located at the apex of the

nucleus.

1224

Figure 4. PfARK1 controls karyokinesis, spindle formation and cytokinesis during

asexual division.

a. Growth assays of synchronised HA-PfARK1-loxP ring stage parasites treated with DMSO

or RAP. *Inset*: fold change in parasitemia of RAP-treated parasites relative to control

(DMSO) (SEM□±□SE, two-sided paired t-test, three independent biological replicates,

*P□=0.0345, **P = 0.0035). **b.** Giemsa stains of HA-PfARK1-loxP parasites treated with

either DMSO or RAP as described in panel **a**. **c.** HA-PfARK1-loxP parasites were treated

with DMSO or RAP and cycle 1 schizonts with E64. Parasites (~40-44 hpi, cycle 1) stained

with DAPI (blue). Scale bar = 2 μm. n >70 parasites, three independent biological replicates

(see Extended Data Fig. 5d). **d.** U-ExM of HA-PfARK1-loxP schizonts (~40-44 hpi, cycle 1)

treated with DMSO or RAP and stained with anti-tubulin (magenta), anti-centrin (green),

SYTOX (blue) and NHS-ester (grey). Hemispindles (yellow arrow), mitotic spindles (cyan

arrow), interpolar spindles (white arrow). Speckles of tubulin (grey arrows) in RAP-treated

parasites are also shown as zoom. Unsegregated nuclei and clusters of centrin (squares) are

also shown in zoom. Scale bar = 2 μm. Upper Right Panel: parasites with normal tubulin

staining of spindles (SEM□±□SE, two-sided unpaired t-test, ****P < 0.0001, n > 55

parasites, three independent biological replicates). Lower Right Panel*:* % parasites with

clustered and/or detached centrin foci (SEM□±□SE, two-sided unpaired t test, ****P

<0.0001, n > 55 parasites, three independent biological replicates). **e and f.** HA-PfARK1-

loxP parasites were treated with DMSO or RAP and cycle 1 schizonts were treated with E64.

IFA (e) or U-ExM (f) was performed on unexpanded (e) or expanded (f) parasites using anti-

MSP1 (e, green) or anti-GAP45 (f, magenta) and anti-tubulin (f, green). RAP-treated

parasites either lacked MSP1 and GAP45 staining or contained unsegregated nuclei (f, cyan

arrow, square, zoom) circumscribed by a large MSP1 contour (e, cyan arrow). Middle Panel:

Quantification of parasites with normal GAP45 (bottom for panel f) or MSP1 staining (top

for panel e) (SEM □±□SE, two-sided unpaired t test, **P = 0.0016, ***P = 0.0006, n > 70

parasites for MSP1 (e) and 40 parasites for GAP45 (f), three independent biological

replicates). Scale bar = 2 µm. U-ExM images are displayed as maximum intensity projection

of multiple z slices (d and f). **g.** Time lapse microscopy of HA-PfARK1-loxP cycle 1

schizonts cultured in the presence of DMSO or RAP. Selected still images from videos

(Supplementary Videos 6-7) showing normal egress. RAP treated parasites that egressed from

erythrocytes but continued to clump-together even after prolonged duration are indicated

(magenta arrow). Scale bar = 2 µm. Right Panel: parasites undergoing normal egress

(SEM □±□SE, two-sided unpaired t test, ***P = 0.0002, n > 100 parasites, three independent

biological replicates). Source data are provided as a Source Data file.

1260

Figure 5. Conditional knockdown of PbARK1 affects male gametogony, disrupting

parasite transmission.

a. qRT-PCR showing normalised expression of PbARK1 transcripts in WT-GFP and

ark1PTD parasites. Mean ± SEM. n = 3 (WT-GFP) and 5 (*ark1PTD*) independent

experiments. **P < 0.01 (P = 0.006), two-sided Welch's t-test. **b**. Exflagellation centres per

field at 15 min post-activation. n = 3 independent experiments (> 7 fields per experiment).

Mean ± SEM. ****P < 0.0001 (P = 1.6 × 10⁻⁸), two-sided Welch's t-test. **c**. Percentage

ookinete conversion. n = 3 independent experiments (> 100 cells). Mean ± SEM. ***P <

0.001 (P = 0.00012), two-sided Welch's t-test. **d**. Total GFP-positive oocysts per infected

mosquito in *ark1PTD* compared to WT-GFP parasites at 7-, 14-, and 21 days post-infection.

Mean ± SEM. n = 3 independent experiments. Statistical significance was determined using

two-sided Welch's t-tests. **P < 0.001, ****P < 0.0001. P = 0.0009 (Day 7), 1.2 × 10⁻⁸ (Day

14), and 1.0 × 10⁻⁵ (Day 21). **e.** Midguts at 10 x and 63 x magnification showing oocysts of

ark1PTD and WT-GFP lines at 14-, and 21 days post-infection (n > 20 oocysts, 3 biological

replicates). Scale bars: 50 µm (10×) and 20 µm (63×). **f.** The diameter of GFP-positive

oocysts in *ark1PTD* compared to WT-GFP parasites at 7-, 14-, and 21 days post-infection.

Mean ± SEM. n = 3 independent experiments. Statistical significance was determined using

two-sided Welch's t-tests. ****P < 0.0001. P = 2.2 × 10⁻⁸ (Day 7), 5.0 × 10⁻⁵ (Day 14), and

1.4 × 10^{-10} (Day 21). **g.** Expansion microscopy images of WT-GFP and *ark1PTD* male
gametocytes at 8 min or 15 min post-activation (mpa). Three images on the left show
maximum intensity projections of whole-cell z-stack images labelled with NHS-ester (grey),
α-tubulin (green) and Hoechst (DNA - blue). The three images on the right show magnified
views of the areas enclosed by the magenta squares. BB: basal body, BBs: basal bodies
(clumped), IM: inner MTOC, A: axoneme. Images are representative of more than 7 cells
analysed across 3 independent experiments. The scale bars are scaled by an expansion factor
of 4.954. Source data are provided as a Source Data file.

1287

Figure 6. *Pbark1* gene-knockdown disrupts spindle and axoneme formation.
a. Expansion microscopy images of *P. berghei* wild-type-GFP (WT-GFP) and *ark1* gene-
knockdown (*ark1PTD*) male gametocytes at 8 min post-activation (mpa). The maximum
intensity projections of entire cells from a z-stack, containing spindles. Cells were labelled
with NHS-ester (grey), a-tubulin (green) and Hoechst (blue). Spindles are indicated by
magenta lines with a and b. Images are representative of more than 7 cells analysed across 3
independent experiments. **b.** Magnified views of each spindle of WT-GFP and *ark1PTD*
highlighted with magenta lines with a and b in Fig. 6a. **c.** Comparison of spindle lengths
measured from expansion microscopy data of WT-GFP and *ark1PTD*. Data represent mean ±
SEM. (n = 7 spindles for WT-GFP and 8 spindles for *ark1PTD*, respectively across 3 different
cells). Statistical significance was determined using a two-sided Welch's t-test. ****$P <$
0.0001 ($P = 0.00009$). The scale bars and spindle lengths are scaled by an expansion factor of
4.954, calculated from a comparison of axoneme diameters measured using expansion and
transmission electron microscopy images. **d.** Transmission electron microscopy images of
WT-GFP and *ark1* gene-knockdown male gametocytes at 8 min or 15 min post-activation. B,
F, J and N are magnified views of the areas enclosed by the magenta squares in A, E, I and M,
respectively. N: nucleus, IM: inner spindle MTOC, BB: basal body, dBB: basal body that
appeared detached from the nucleus, A: axoneme, K: kinetochore, Sp: spindle microtubule, F:
flagellum, BBs: basal bodies (clumped), Mt: microtubule. Images are representative of more
than 50 cells analysed across 2 independent experiments. Source data are provided as a
Source Data file.

1309

Figure 7. PbARK1 is the divergent Aurora kinase of the alternative chromosomal passenger complex (CPC) in *Plasmodium* spp.

a. Fold enrichment (Log 2 transformed ratio) of normalized iBAQ values of proteins found in co-immunoprecipitates of ARK1-GFP versus control GFP-only for both 8-9 hr schizont and 1.5-min male gametocyte lysates (n = 2 for all four pulldowns experiments). *P*-values were derived using a two-tailed Student's *t*-test and corrected for multiple testing using the Benjamini-Hochberg method. False Discovery Rate (FDR) was set at 5%, which corresponds to an unadjusted *P*-value of 0.0067 (see dotted line). The (raw) data underlying this figure can be found in in Supplementary Data 2. **b.** Zoom-in on the domain/motif topology of INCENP-A/-B in apicomplexans (Pb = *P. berghei*; Tg = *T. gondii*; Cmur = *C. muris)* and model organisms (Hsap = *Homo sapiens*; Ddis = *Dictyostelium discoideum*; Gint = *Giardia intestinalis*), focused on the extended IN-box region (blue-yellow-red) that interacts with the kinase domain of ARK1. Arrows indicate an antiparallel beta sheet formed by the N- and C-terminal extensions of the IN-box. Green: TSS transphosphorylation site (asterisks indicate (putative) phosphorylated sites) in eukaryotic INCENP homologs, but not in INCENP-A/-B in *P. berghei* (Pb) and *T. gondii* (Tg). RET is a candidate transphosphorylation site in PbINCENP-A (Pb-A). Protein structures are predicted using AlphaFold3, for Predicted Aligned Error (PAE) plot, see Supplementary Fig. 9. **c.** Presence/absence of ARK1/2/3, and components of the chromosomal passenger complex amongst the model eukaryotes, the SAR supergroup with a specific focus on Apicomplexa. Lack of phylogenetic signal for one ancestral duplication of INCENP amongst apicomplexans implies that multiple different subgroups underwent independent INCENP duplications. In Plasmodium these paralogues are called INCENP-A and -B; in Toxoplasma these are called INCENP-1 and -2. Right: consensus sequence logos of the activation and P+1 loop of Aurora kinases in Clustal color scheme: divergent phosphorylation consensus in ARK1 (GxSH instead of RxT). **d.** Summary of findings and discussion points from this study.

a

Plasmodium falciparum **erythrocytic schizogony** *(blood stage - vertebrate host)*

hemispindle   interpolar spindle   schizont   8-32x merozoites

Plasmodium berghei **(male) gametogony** *(mosquito gut vector 1N>8N)*

1N   continuous spindle   2N   4N   8N octaploid nucleus   8x flagellated male gametes

1 minute post activation   2-3 mpa   4-5 mpa   6-8 mpa   15 mpa

Legend: ○ red blood cell   ○ cell membrane   ○ nucleus   ● MTOC   ◆ basal body   ⫦ (spindle) microtubules   ◣ axoneme   ⦂○ (clustered) centromeres

b

Aurora localisation

MTOC   karyokinesis or cytokinesis   microtubules   kinetochores   chromosomes   nucleoplasm

Scaffolds *(e.g. human, yeast, fly, worm etc.)*

Chromosomal Passenger Complex (CPC)
4 subunits: Aurora-INCENP-Borealin-Survivin

Lineage-specific scaffolds
e.g. TPX2, Bora & Cep192

Three Aurora-Related Kinases (ARKs) in Apicomplexa

	Plasmodium		*Toxoplasma*	*Cryptosporidium*
stage	schizont	male gametocyte	tachyzoite	-
ARK3	?	absent	?	absent
ARK2	?	*EB1-MyoK MISFIT*	?	absent
ARK1	?	?	*INCENP1-2*	untested

a

Plasmodium falciparum erythrocytic schizogony

Trophozoite   Early-Mid schizont   Late schizont

○ red blood cell   ○ cell membrane   ● nucleus   ● MTOC
⫽ (spindle) microtubules   ⁸○ (clustered) centromeres

Schizonts

	Early	Mid	Late	Segmentor

HA-PfARK1
Centrin
Merge DAPI
DIC

b

c

d

a *Plasmodium berghei* male gametogony

1 minute post activation — continuous spindle — 2-3 mpa — 4-5 mpa — octaploid nucleus — 6-8 mpa

1N → 2N → 4N → 8N

red blood cell   cell membrane   nucleus   MTOC   basal body
(spindle) microtubules   axoneme   (clustered) centromeres

1 min | 2-3 min | 4-5 min | 6-8 min

PbARK1-GFP
PbARK1-GFP Hoechst
DIC

b PbARK1-GFP / Hoechst

c PbARK1-GFP / α-tubulin / Hoechst / NHS-ester
Whole cell
Spindle

d PbARK1-GFP
PbARK2-ch
Merge Hoechst

e Merge / Hoechst
PbARK1-GFP
PbARK2-ch

f PbARK1-GFP
PbNDC80-ch
Merge Hoechst

g Merge / Hoechst
PbARK1-GFP
PbNDC80-ch

h PbARK1-GFP
Pbkinesin-8B-ch
Merge Hoechst

i Merge / Hoechst
PbARK1-GFP
Pbkinesin-8B-ch