Evaluation of potential stress marker genes during acclimation of the Patagonian *Lactiplantibacillus plantarum* UNQLp11 strain

Zerbino Antonella[1,2], Lannutti Lucas[1,2], Olguin Nair*[1,2]

[1]Departamento de Ciencia y Tecnología, Universidad de Morón, Machado 914 (B1708EOH), Morón, Buenos Aires, Argentina.
[2]Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Godoy Cruz 2290 (C1425FQB) CABA, Argentina.

*Corresponding author
Correspondence: ntolguin@gmail.com

ABSTRACT

In this study, we evaluated the expression of genes that may serve as markers of temperature- and ethanol-induced stress responses during the acclimation of *Lactiplantibacillus plantarum* strains of enological interest. The transcriptional responses of eight genes potentially related to stress adaptation were analyzed in two strains acclimated at 18 °C and 21 °C. Gene expression was assessed at the start (time zero) and after 48 hours of acclimation. Following acclimation, the cells were inoculated into sterile Pinot noir wine, and L-malic acid consumption was monitored over 19 days to determine whether acclimation temperature influenced cell survival and fermentation performance. Both strains exhibited similar L-malic acid consumption and survival patterns, although transcriptional responses varied depending on the temperature. Notably, *gapB*, *glmS*, and *rfbB* emerged as promising gene markers for future studies. These findings suggest that gene expression profiling of acclimated cells may support the selection of robust *Lpb. plantarum* strains for use as starter cultures in winemaking.

Keywords: gene expression, L-malic acid consumption, Real Time RT-qPCR, acclimation, *Lactiplantibacillus plantarum*

Introduction

In wine, lactic acid bacteria (LAB) are responsible for malolactic fermentation (MLF), a decarboxylation process in which L-malic acid is converted into L-lactic acid. This process is desirable in most red wines and in some white and sparkling wines for three main reasons: it lowers acidity (leading to a softer mouthfeel), confers microbiological stability, and it modifies the wine's flavor (Bartowsky, 2017). Several LAB species can perform MLF, with *Oenococcus oeni* being the main responsible in many countries. As such, it is the most extensively studied and widely used as starter culture (Renouf et al., 2006; Nisiotou et al., 2015; Bartowsky, 2017). Recently, *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*) has emerged as an alternative species. It possesses a more diverse enzymatic profile than *O. oeni,* which may significantly influence flavor development (Iorizzo et al. 2016; Krieger-Weber et al., 2020). Additionally, *Lpb. plantarum* has the genetic potential to produce antimicrobial peptides (plantaricins), which could enhance its dominance during MLF and offer the added benefits of reducing the need of sulphurdioxide (SO_2) commonly used to inhibit undesirable microbial growth (du Toit et al., 2011).

Wine presents a harsh environment due to its high ethanol content, low pH, the presence of SO_2 and other inhibitory compounds. Not all LAB species or strains can survive under such conditions, but *Lpb. plantarum* has demonstrated stress tolerance comparable to that of *O. oeni* (Iorizzo et al., 2016; Krieger-Weber et al., 2020). Thanks to transcriptomic and other omics-bases analyses, we are beginning to understand the gene and protein expression profiles that contribute to LAB survival in wine (Olguin et al., 2010; 2015; 2019; Bordas et al., 2015; Margalef-Català et al., 2016). However, most of this research has focused on *O. oeni*, while studies involving *Lpb.plantarum* remain limited (Fiocco et al., 2007; Miller et al., 2011).

One of the main criteria for selecting LAB strains as starter cultures is their resistance to wine conditions and their metabolic performance. However, although commercial starter cultures are widely available, a limitation is that they are often derived from specific regions and may not adequately capture the microbial diversity representative of other terroirs. The potential contribution of microorganisms to the concept of terroir has long been overlooked (Knight et al., 2015). These microbes not only participate in fermentation but also influence soil chemistry and nutrition, as well as grapevine health, yield, and quality (Belda et al., 2017). Several studies have linked wine characteristics to regional microbial communities (Vitulo et al., 2019), with some specifically highlighting the role of bacteria in shaping wine quality (Portillo et al., 2016; Grangateau et al., 2017; Lorentzen & Lucas, 2019). Therefore, it is important to evaluate whether isolated microbial strains are beneficial within a given terroir. In some cases, the preservation of terroir itselfmay guide the isolation, identification, and characterization of strains for use as starter cultures.

LAB populations in Patagonian wines display high species and strain diversity during spontaneous MLF (Valdés La Hens et al., 2015). In previous work, we applied an acclimation protocol (Bravo-Ferrada et al., 2014), in which cells were cultured in a broth with moderate ethanol concentration prior to wine inoculation. We investigated the influence of two temperatures (18 °C and 21 °C) on gene expression using *O. oeni* strains isolated from Patagonian Pinot noir wines (Olguin et al., 2019). We found that L-malic acid consumption, population survival, andtranscriptional behavior differed depending on acclimation temperature.

The aim of the present study was to evaluate the expression of stress-related genes in *Lpb. plantarum* UNQLp11 induced by acclimation at two temperatures (18 °C and 21 °C). Gene selection was based on previous studies involving both *O. oeni* (Olguin et al., 2010; 2019) and *Lpb. plantarum* (Fiocco et al., 2007; Miller et al., 2011). Gene expression was quantified by reverse transcription real-time quantitative PCR (Real Time RT-qPCR). Population survival and L-malic acid consumption after acclimation were monitored in Pinot noir wine.

Material and methods

Growth conditions

Lpb. plantarum subsp. *plantarum* ATCC 14917 (DSM 20174, Zheng et al., 2020) and the strain UNQLp11 (RGPRN-/-M-/-357-/-Semorile-/-626-/-UNQLp11-/-2024), previously isolated from Patagonian red wines (Bravo-Ferrada et al., 2013; Iglesias et al., 2020) where cultured at 28 °C in tubes containing MRS broth medium (De Man et al., 1960), supplemented with L-malic acid (4 gL^{-1}) and fructose (5 gL^{-1}) at pH 5.0. After 48 h cells were harvested by centrifugation and resuspended in the acclimation medium (50 gL^{-1} MRS, 40 gL^{-1} fructose, 20 gL^{-1} glucose, 4.5 gL^{-1} L-malate, 1 gL^{-1} Tween 80, 0.1 mgL^{-1} pyridoxine, pH 4.6) containing 6% (v/v) of ethanol (Bravo-Ferrada et al., 2014). One of the acclimation tubes was cultured at 18 °C and the other one at 21 °C. We chose these temperatures because both are common during MLF in wineries. Samples were taken at time zero (at the beginning of cell acclimation) and 48 h after acclimation (just before wine inoculation).

Then, cells were harvested by centrifugation and inoculated in tubes containing sterile Pinot noir wine (vintage 2014), at the final stage of alcoholic fermentation (14.3 % ethanol, pH 3.7, 2.3 gL^{-1} L-malic acid). These tubes were kept at 21 °C and cell survival was analysed every 2-3 days until the 19th day. All assays were performed in duplicate at different times using independent precultures and cell survival was monitored by counting colonies on plates of MRS medium supplemented as described above. L-malic acid consumption was measured using the L-Malic Acid Enology Enzymatic kit (BioSystems SA, Barcelona, Spain) according to the manufacturer's instructions.

Sovereign Rights Over Natural Resources: The indigenous LAB strain mentioned above is a genetic resources belonging to the Provinces of Rio Negro (UNQLp11). It was deposited in the strain collection of the Laboratorio de Microbiología Molecular (IMBA–DCyT–UNQ-CIC) and registered with the Environment and Climate Change Secretariat— Province of Rio Negro Government. Their preservation follows the protocols of the Convention on Biological Diversity and the Nagoya Protocol, which safeguard the ownership and use of genetic resources from the regions where they were isolated (RESOL-2024-626-E-GDERNE-SAYCC#SGG, NO-2024-00628737-GDERNE-SAYCC#SGG).

RNA extraction

Lpb. plantarum cells were harvested by centrifugation and frozen until RNA extraction. Total RNA extractions were performed using the SV Total RNA Isolation System #Z3100 (Promega) following the manufacturer's instructions. RNA concentrations were calculated by measuring absorbance at 260 nm using a Nanodrop spectrophotometer (Thermo Fisher Scientific, 1000).

Selection of candidate genes and nucleotide sequences

The gene sequences of *Lpb. plantarum* reference strain SRCM100442 (Genome assembly ASM991365v1) were obtained from the National Center for Biotechnology Information (NCBI). The reference genome assembly strain SRCM100442 was used for sequence comparison against *O. oeni* PSU-1 (NC_008528) (Table 1) and for primers design when applicable (Table 2). We also tested sequence alignment and *in silico* PCR with the strain ATCC 14917 gene sequences. Sequence alignment and primer pairs were tested in silico using the NCBI Needleman-Wunsch Global Align Nucleotide Sequences and Primer-BLAST, respectively (https://blast.ncbi.nlm.nih.gov/Blast.cgi).

133 **Table 1** - Comparison of genes sequence alignment between *Lactiplantibacillus*
134 *plantarum* SRCM100442 and *Oenococcus oeni* PSU-1 (NC_008528)

Target gene *Lpb. plantarum*	Gene name	Gene sequence identity (%)	Protein sequence identity (%)	*O. oeni* gene symbol
C7M33_RS14920	*ldhD*	60	51	OEOE_RS01985
C7M33_RS06325	*gyrA*	66	69	OEOE_RS00030
C7M33_RS14590	*rpoD*	59	61	OEOE_RS04785
C7M33_RS01170	*recA*	65	68	OEOE_RS06650
C7M33_RS09485	*gapB*	72	72	OEOE_RS01940
C7M33_RS05025	*hsp20*	57	41	OEOE_RS01385
C7M33_RS11210	*rfbB (rmlB)*	57	45	OEOE_RS06990
C7M33_RS14780	*dnaK*	70	74	OEOE_RS06310
C7M33_RS09640	*glmS*	63	60	OEOE_RS03035
C7M33_RS01055	*trxA*	60	50	OEOE_RS08215
C7M33_RS10385	*cspA*	79	81	OEOE_RS06620
C7M33_RS04410	*dacA*	51	32	OEOE_RS03435

135 Note: *O. oeni* PSU-1 genes locus tag have been discontinued but are still accessible. Since
136 the new sequences are variable, we decided to use the ones we already had to compare
137 with the *Lpb. plantarum* reference genome assembly (strain SRCM100442).

Reverse transcription and Real Time RT-qPCR

cDNA was synthesized from RNA (15 ng μL^{-1}) using TaqMan Reverse Transcription
Reagents (Applied Bio- systems) as recommended. Ten pairs of primers were designed (Table
2) to be about 18–22 bases long, to contain over 50 % G/C and to have a melting temperature
(Tm) above 60 °C. Two pairs of primers were taken from previous works (Fiocco et al., 2008;
Duary et al., 2012). The length of the PCR products ranged from 95 to 127 bp (Table 2). RT-
qPCR was performed in 12 µL final volume containing 1.5 ng of total cDNA, 1.0 µL of each
primer at 3.15 µmol L^{-1}, 2.5 µL of RNase free water and 6.5 µL of SYBR Green PCR Master Mix
(Applied Biosystems). Amplifications were carried out using a QuantStudio 3 Real-Time PCR
Systems (Applied Biosystems) with an initial step at 95 °C for 5 min followed by 35 cycles of 95
°C for 10 s and 60 °C for 30 s and 72 ºC for 30 s. An additional step to establish a melting curve
was used to verify the specificity of the real-time PCR reaction for each primer pair. The
threshold value used in this study was automatically determined by the instrument.
Results were analysed using the comparative C_T method ($2^{-\Delta\Delta C_T}$) as described by Schmittgen
& Livak (2008). Two statistical algorithms were used to determine the stability of the candidate
reference genes: NormFinder (Andersen et al., 2004) and geNorm (Vandesompele et al., 2002).
The three candidates, *ldhD*, *rpoD* and *gyrA* were chosen as the best combination.
155

156 **Table 2** – Primers used in this work

Target gene	Description	Forward primer (5'→3')	Reverse primer (5'→3')	Amplicon length (bp)	Reference
ldhD	D-2-hydroxyacid dehydrogenase	ACGCCCAAGCTGATGTTATC	AGTGTCCCACGAGCAAAGTT	127	Fiocco et al. (2008)
recA	recombinase A	CCTTAGATGACGCGTTGG	CAGCAACCGCATGTAGTG	108	This work
gyrA	DNA gyrase subunit	GTCAAACGGACACCAGTTCAAG	GTGACCATCCGTGATTGTAACC	111	This work
gapB	type I glyceraldehyde 3-phosphate dehydrogenase	ACTGCCTTGCACCATTAGC	TACAGGGCCGTCAAGTAAC	113	This work
hsp20	Hsp20/alpha crystallin family protein	ATGGGTCGGTCCTAAAGACTG	CGATAATGTGCCATCCCGATAC	119	This work
rfbB	dTDP-glucose 4,6-dehydratase	AACGCTTCCACCATGTTTCC	GTGTAAACTTCTCGCCAACACC	99	This work
dnaK	molecular chaperone DnaK	TTAGTACGGCGGCTGATAAC	GAAGCGACCTAACGTCTTG	95	This work
glmS	glutamine-fructose-6-phosphate transaminase	AATGTGATGCGGAAGCTATCCC	CCAGCAGCCACAATGTATAGTC	116	This work
trxA	thioredoxin	AACGGATACGGCTTCAGG	CACCCATCTCGTCAGCTAATTG	107	This work
cspA	cold shock protein	CTGGTGAAGATGGCAACGATG	CCACGATCGCTTGATTCTTC	115	This work
rpoD	RNA polymerase sigma factor RpoD	GCTTACGGTTCGGACTTGATG	GCCTTGGCTTCGATTTGG	103	Duary et al. (2010)
dacA	D-alanyl-D-alanine carboxypeptidase	AAGCAGCCAGTACACCATC	CGCCATTGGCTTACTACTGTTC	101	This work

158 .

Results

The *Lpb. plantarum* strain used in this study, isolated from a Patagonian Pinot noir wine
undergoing spontaneous MLF, was identified as UNQLp11 (Bravo-Ferrada et al., 2013). It was
selected from a pool of strains obtained from the same wine based on its oenological properties
(Brizuela *et al.*, 2018). *Lpb. plantarum* DSM 20174, isolated from pickled cabbage, was used as
a reference strain. Since this strain originates from a non-wine environment, any observed
metabolic or genetic similarities are likely species-specific rather than environmentally induced.

Acclimation temperatures of 18∟°C and 21∟°C were selected based on prior work showing
temperature-dependent behavior in *O. oeni* strains (Olguin et al., 2019).

During the 48-hour acclimation period, cell density increased by approximately two
logarithmicunits (LOGs) at both temperatures for both strains.

Genes from *Lpb. plantarum* were selected based on their known functions, under the
hypothesis that they may respond to wine-associated stress conditions similarly to *O. oeni* (Table
1). Prior to quantitative analysis, Real Time RT-qPCR amplification of the selected genes was
tested in both strains. The amplified fragment sizes were as expected (Table 2). Amplification
efficiency was calculated for each primer pair; the gene *dacA* was excluded due to poor
efficiency (E = 80.21%, R² = 0.9958). All remaining primers had efficiencies within 10% of each
other, making them suitable for relative quantification using the comparative Ct (ΔΔCt) method
(Schmittgen & Livak, 2008).

Relative gene expression was first analyzed using the sample at time zero under acclimation
conditions (A0) as the calibrator (Fig. 1). At 18∟°C, the genes *rfbB* (also referred to as *rmlB*)
and *glmS* were clearly overexpressed, especially in the Patagonian strain UNQLp11 (Fig. 1A). In
contrast, *recA* was slightly overexpressed in the reference strain ATCC 14917 but
underexpressed in UNQLp11. Genes *dnaK*, *trxA*, *cspA*, and *gapB* were consistently
underexpressed in both strainsat this temperature.

At 21∟°C, gene expression trends differed (Fig. 1B). In ATCC 14917, most genes showed
stable or reduced expression, except for a slight overexpression of *cspA* and a marked
overexpression of *gapB*. In UNQLp11, there was slight overexpression of *glmS* and *trxA*,
with *gapB* again notably overexpressed. All other genes were stably expressed, with the
exception of *rfbB*, which was significantly underexpressed.

Figure 1 - Relative expression of eight genes related to stress response (*hsp20*,
rfbB, *glmS*, *recA*, *dnaK*, *trxA*, *cspA*, *gapB*) after 48 h acclimation at 18 °C (A) and
at 21 °C (B). The calibration condition used was time zero when *Lactiplantibacillus*
plantarum cells were inoculated into the acclimation medium.

To assess the impact of temperature, gene expression levels after 48 hours of acclimation
(A48) were compared between 18∟°C and 21∟°C, using the latter as the calibrator condition
(Fig. 2). Expression patterns were largely consistent between strains. Both strains showed strong
overexpression of *rfbB* and slight overexpression of *glmS*. Genes *dnaK*, *trxA*, *cspA*,

and *gapB* were underexpressed under the lower temperature. Differences between strains were
limited to slight overexpression of *hsp20* and *recA* in the ATCC 14917 strain.
203

204
205
206 **Figure 2** - Relative expression of eight genes related to stress response (*hsp20*,
207 *rfbB*, *glmS*, *recA*, *dnaK*, *trxA*, *cspA*, *gapB*) in *Lactiplantibacillus plantarum* strains
208 after 48 h acclimation at 18 °C. The calibration condition used was time 48 h at 21
209 °C.
210
Cells acclimated at both temperatures (18 °C and 21 °C) were inoculated into sterile Pinot
noir wine and monitored over 19 days. Cell survival and L-malic acid consumption were
evaluated at different time points (Fig. 3).
Viable cell counts for both ATCC 14917 and UNQLp11 declined during the fermentation
period (Fig. 3A). However, both strains were capable of consuming the majority of the L-malic
acid within the first seven days of incubation (Fig. 3B).
217

218
219
Figure 3 - Evolution of cells viability (log CFU mL^{-1}) and L-malic acid consumption
in Pinot noir wine of the *Lactiplantibacillus plantarum* strains acclimated at 18 and
21 °C. (A) ATCC 14917 (circles), UNQLp11 (squares), solid symbols and lines
indicate strains acclimated at 18 °C and empty symbols and dashed lines indicate
strains acclimated at 21 °C. (B) Percentage of L-malic acid consumed by cells
acclimated at 18 °C (black bars) and at 21 °C (grey bars) after 7 days (7d) in sterile
wine.
227
228

Discussion

230 The aim of this study was to expand our understanding of the genetic mechanisms involved
in the acclimation of *Lpb. plantarum*, which may enhance its survival under the stressful
conditions present in wine. Previous studies have demonstrated that wine-associated LAB, such
as *O. oeni* and *Lpb. plantarum*, benefit from an acclimation step prior to inoculation into wine
(Cecconi et al., 2009; Bravo-Ferrada et al., 2013; Costantini et al., 2015). Stress-related genes
in *O. oeni* have been proposed as markers of cellular adaptation (Hamon et al., 2013), but the
genetic response of *Lpb. plantarum* to wine-like conditions remains poorly characterized. Thus,
we evaluated the expression of stress-related genes in two *Lpb. plantarum* strains subjected
toacclimation at two different temperatures.

239 Sequence comparisons between *O. oeni* and *Lpb. plantarum* indicate that while their genes
240 and proteins perform homologous functions, their regulatory mechanisms can differ. It is known
241 that different stress-resistance phenotypes may protect against the same environmental stressor,
242 even when the molecular pathways involved are distinct (Papadimitriou et al., 2016). Therefore, it
243 is not unexpected that some genes in *Lpb. plantarum* respond similarly to those in *O. oeni*, while
244 others do not (Fig. 1).

245 The gene expression profiles of strains UNQLp11 and ATCC 14917 showed broadly similar
responses, which corresponded with their survival and L-malic acid consumption in wine
following acclimation. Among the stress response genes (*hsp20*, *cspA*, *dnaK*, and *trxA*), we
observed general under-expression at 18 ℃. These temperature conditions are moderate, and
more extreme values are typically required to elicit strong responses in heat-shock or cold-shock
genes (Liao et al., 2010; Fiocco et al., 2007). Nevertheless, due to the potential for cross-
protection mechanisms (Papadimitriou et al., 2016; Ma et al., 2021), we included
both *hsp20* and *cspA* in our analysis. The under-expression of *cspA* may align with previous
findings in *O. oeni*, where theprotein was downregulated following acclimation in 8% ethanol
(Cecconi et al., 2009a).

255 While earlier studies reported that *Lpb. plantarum* strains overexpressing heat shock proteins
were more ethanol-tolerant (Fiocco et al., 2007), our results differ from some previous findings.
For example, *hsp* gene expression in *Lpb. plantarum* has been correlated with that in *O.
oeni* (Capozzi et al., 2012), but in our study, *hsp20* did not show significant upregulation,
consistent with previous work on *O. oeni* under similar conditions (Olguin et al., 2019). Similarly,
although *dnaK* expression has been shown to increase under various stress conditions (Liao et
al., 2010; Reverón et al., 2012), we did not observe this in our study, again aligning with earlier
findings in *O. oeni* (Olguin et al., 2019). The gene *trxA*, typically upregulated under oxidative and
ethanol stress in both species (Serrano et al., 2007; Reverón et al., 2012; Margalef-Català et
al.,2017), also did not show the expected increase in our experiment.

265 With respect to carbohydrate metabolism, *gapB* is often considered a housekeeping gene
(Duray et al., 2012), yet in our study, its expression appeared temperature-dependent. This
aligns with previous observations where *gapB* was upregulated under oxidative stress in
specific *Lpb. plantarum* strains (Serrano et al., 2007). Additionally, other studies have reported
changes in protein expression linked to carbohydrate metabolism in response to different culture
conditions in both *Lpb. plantarum* (Cecconi et al., 2009b; Hamon et al., 2013) and *O.
oeni* (Cecconi et al.,2009a; Olguin et al., 2015; Costantini et al., 2015).

272 In the context of amino acid metabolism, we analyzed *glmS*, which showed a notable
273 increase in expression, especially at 18 ℃. Previous work has shown that *glmS* is
274 downregulated in response to antioxidant stress induced by hydroxytyrosol or gallic acid
275 (Reverón et al., 2015; 2020), and its expression also decreases in *O. oeni* following ethanol
276 shock (Olguin et al., 2015). Our results contrast with those findings, suggesting a different
277 regulatory pattern undertemperature-induced stress.

278 The *recA* gene, associated with DNA repair and recombination, has been variously classified
279 as a housekeeping or stress-related gene (Miller & Kokjohn, 1990; Curiel et al., 2011; Reverón
280 etal., 2012, 2015). In our study, its expression remained relatively stable under both conditions.

An especially interesting result was observed for *rfbB*, a gene involved in cell envelope biogenesis. It was strongly overexpressed at 18 °C but downregulated at 21 °C. Previous studies have reported no significant change in *rfbB* protein levels under acidic conditions in *Lpb. plantarum* (Hamon et al., 2013), while in *O. oeni*, both over- and under-expression have beenobserved depending on the acclimation conditions (Olguin et al., 2015; 2019).

The impact of acclimation on both *O. oeni* and *Lpb. plantarum* using the same medium has been studied in previous research (Bravo-Ferrada et al., 2014; Brizuela et al., 2018). Thus, the similar survival and L-malic acid consumption profiles observed in both *Lpb. plantarum* strains in this study were not unexpected. Notably, strain UNQLp11 was isolated from Patagonian Pinot noir wine, and the selection of indigenous strains as starter cultures is gaining global interest for enhancing regional wine identity (Bartowsky et al., 2017). As suggested by Mao et al. (2021), environmental selection pressure can drive genomic changes that improve strain adaptability to specific ecological niches. Therefore, while our results are promising, further research usingadditional wine-isolated *Lpb. plantarum*strains is necessary to validate these findings.

To our knowledge, this is the first study to evaluate stress-related gene expression in *Lpb. plantarum* as potential markers for winemaking applications. As the use of this species in starter culture development continues to grow, further research incorporating gene expression and omics approaches will be valuable. Similar to ongoing work in *O. oeni*, these studies can help correlate genetic behavior with enological performance, ultimately supporting the selection of robust, regionally adapted strains for the wine industry.

In conclusion, this study offers new insights into the genetic mechanisms underlying the acclimation of *Lpb. plantarum* to wine conditions—an area that remains less understood compared to *O. oeni*. By analyzing the gene expression profiles of *Lpb. plantarum* strains at two temperatures, we identified both shared and distinct responses relative to *O. oeni*. While several stress-related genes (*hsp20*, *cspA*, *dnaK*, *trxA*) were under-expressed, genes suchas *glmS* and *rfbB* showed temperature-dependent overexpression.

These findings highlight the potential of *Lpb. plantarum* as a viable candidate for use in starter culture development, particularly when selecting indigenous strains to support regional wine identity. Overall, the results deepen our understanding of *Lpb. plantarum*'s stress adaptation and support its further exploration for enological applications, laying the groundwork for future studies aimed at optimizing strain selection and performance in winemaking.

Acknowledgements

We would like to thank the Environment and Climate Change Secretariat, Government of the Province of Rio Negro, for authorizing access to provincial genetic resources and their use for academic and research purposes. The authors also thank the Laboratorio de Microbiología Molecular (LMM) of the Universidad Nacional de Quilmes (UNQ) for lending them the equipment to perform the Real Time RT-qPCR.

Funding

This work was funded by grants from CONICET [PIBAA 2022 No 28720210100183CO], Agencia Nacional de Promoción Científica y Tecnológica, [PICT 2021 GRF-TI-00350] and, University of Morón [PICT 2024 - 80020230100046UM], Buenos Aires, Argentina. AZ and LL are PhD students from CONICET-UM. NTO is a member of the Research Career of CONICET.

Conflict of interest disclosure

The authors have no conflicts of interest to declare.

Data information availability

327   Data are available online: https://doi.org/10.1101/2025.08.28.672453; Zerbino et al., 2025.
328 Additional documents are available: http://hdl.handle.net/11336/270401; Zerbino et al., 2025.

References

Andersen C, Jensen J, Orntoft T (2004) Normalization of real time quantitative reverse
transcription—PCR data: a model based variance estimation approach to identify genes
suited for normalization, applied to bladder and colon cancer data sets. *Cancer Research*, **64**,
5245. https://doi.org/10.1158/0008

Bartowsky E J (2017) *Oenococcus oeni* and the genomic era. *FEMS Microbiology*
Reviews, **41**(Supp_1), S84-S94. https://doi.org/10.1093/femsre/fux034

Belda I, Zarraonaindia I, Perisin M, Palacios A, Acedo A (2017) From vineyard soil to wine
fermentation: microbiome approximations to explain the "terroir" concept. *Frontiers in*
Microbiology, **8**, 821. https://doi.org/10.3389/fmicb.2017.00821

Bordas M, Araque I, Bordons A, Reguant C (2015) Differential expression of selected
Oenococcus oeni genes for adaptation in wine-like media and red wine. *Annals*
of Microbiology, **65**, 22772285. https://doi.org/10.1007/s1321 3-015-1069-2

Bravo-Ferrada B M, Hollmann A, Delfederico L, Valdes La Hens D, Caballero A, Semorile L
(2013) Patagonian red wines: selection of *Lactobacillus plantarum* isolates as potential starter
cultures for malolactic fermentation. *World Journal of Microbiology and Biotechnology*, **29**,
1537–1549. https://doi.org/10.1007/s11274-013-1337-x

Bravo-Ferrada B M, Tymczyszyn E E, Gómez-Zavaglia A, Semorile L (2014) Effect of
acclimation medium on cell viability, membrane integrity and ability to consume malic acid in
synthetic wine by oenological *Lactobacillus plantarum* strains. *Journal of Applied*
Microbiology, **116**, 360–367.https://doi.org/10.1111/jam.12372

Brizuela N S, Bravo-Ferrada B M, Pozo-Bayón M A, Semorile L, Tymczyszyn E E (2018)
Changes in the volatile profile of Pinot noir wines caused by Patagonian *Lactobacillus*
plantarum and *Oenococcus oeni* strains. *Food Research International*, **106**, 22–8.
https://doi.org/10.1016/j.foodres.2017.12.032.

Capozzi V, Fiocco D, Weidmann S, Guzzo J, Spano G (2012) Increasing membrane protection
in *Lactobacillus plantarum* cells overproducing small heat shock proteins. *Annals of*
*Microbiology,***62**, 517-522. https://doi.org/10.1007/s13213-011-0285-7

Cecconi D, Milli A, Rinalducci S, Zolla L, Zapparoli G (2009a) Proteomic analysis of *Oenococcus*
oeni freeze-dried culture to assess the importance of cell acclimation to conduct
malolacticfermentation in wine. *Electrophoresis*, **30**, 2988–2995.
https://doi.org/10.1002/elps.200900228

Cecconi D, Cristofoletti M, Milli A, Rinalducci S, Zolla L, Zapparoli G (2009b) Effect of tannic
acid on *Lactobacillus plantarum* wine strain during starvation: a proteomic study.
*Electrophoresis,***30**, 957–965. https://doi.org/10.1002/elps.200800310

Costantini A, Rantsiou K, Majumder A, Jacobsen S, Pessione E, Svensson B, Garcia-Moruno E,
Cocolin L (2015) Complementing DIGE proteomics and DNA subarray analyses to shed light
on *Oenococcus oeni* adaptation to ethanol in wine-simulated conditions. *Journal of*
*Proteomics,***123**, 114–127. https://doi.org/10.1016/j.jprot.2015.04.019

Curiel J A, Rodríguez H, de las Nieves B, Anglade P, Baraige F, Zagorec M, Champomier-
Vergès M, Muñoz R, López de Felipe F (2011) Response of *Lactobacillus plantarum* human
isolate to tannic acid challenge assessed by proteomic analyses. *Molecular Nutrition Food*
Research, **55**,1454-1465. https://doi.org/10.1002/mnfr.201000621

De Man J C, Rogosa M, Sharpe M E (1960) A medium for the cultivation of Lactobacilli.
Journalof Applied Bacteriology, **23**, 130–135. https://doi.org/10.1111/j.1365-
2672.1960.tb00188.x

Duary R K, Batish V K, Grover S (2012) Relative gene expression of bile salt hydrolase and
surface proteins in two putative indigenous *Lactobacillus plantarum* strains under in vitro gut

conditions. *Molecular Biology Reports*, **39**, 2541-2552. https://doi.org/10.1007/s11033-011-1006-9

du Toit M, Engelbrecht L, Lerm E, Krieger-Weber S (2011) *Lactobacillus*: the next generation of malolactic fermentation starter cultures – an overview. *Food and Bioprocess Technology*, **4**,876-906. https://doi.org/10.1007/s11947-010-0448-8

Fiocco D, Capozzi V, Goffin P, Hols P, Spano G (2007) Improved adaptation to heat, cold, and solvent tolerance in *Lactobacillus plantarum*. *Applied Microbiology and Biotechnology*, **77**, 909915. https://doi.org/10.1007/s00253-007-1228-x

Fiocco D, Crisetti E, Capozzi V, Spano G (2008) Validation of an internal control gene to apply reverse transcription quantitative PCR to study heat, cold and ethanol stresses in *Lactobacillus plantarum*. *World Journal of Microbiology and Biotechnology*, **24**, 899-902.https://doi.org/10.1007/s11274-007-9556-7

Grangeteau C, Roullier-Gall C, Rousseaux S, Gougeon R D, Schmitt-Kopplin P, Alexandre H, Guilloux-Benatier M (2017) Wine microbiology is driven by vineyard and winery anthropogenicfactors. *Microbial Biotechnology*, **10**, 354–370. https://doi.org/10.1111/1751-7915.12428

Hamon E, Horvatovich P, Marchioni E, Aoudé-Werner D, Ennahar S (2013) Investigation of potential markers of acid resistance in *Lactobacillus plantarum* by comparative proteomics.*Journal of Applied Microbiology*, **116**, 134-144. https://doi.org/10.1111/jam.12339

Iglesias N G, Brizuela N S, Tymczyszyn E E, Hollmann A, Valdés La Hens D, Semorile L, Bravo-Ferrada B M (2020) Complete genome sequencing of *Lactobacillus plantarum* UNQLp11 isolated from a Patagonian Pinot noir wine. *South African Journal of Enology and Viticulture*,**41**(2), 197-209. https://doi.org/10.21548/41-2-4066

Iorizzo M, Testa B, Lombardi S J, García-Ruiz A, Muñoz-González C, Bartolomé B, Moreno-Arribas M V (2016) Selection and technological potential of *Lactobacillus plantarum* bacteria suitable for wine malolactic fermentation and grape aroma release. *LWT Food Science andTechnology*, **73**, 557-566. https://doi.org/10.1016/j.lwt.2016.06.062

Knight S, Klaere S, Fedrizzi B, Goddrard M R (2015) Regional microbial signatures positively correlate with differential wine phenotypes: evidence for a microbial aspect to terroir. *ScientificReports*, **5**, 14233. https://doi.org/10.1038/srep14233

Krieger-Weber S, Heras J M, Suarez C (2020) *Lactobacillus plantarum*, a new biological tool to control malolactic fermentation: a review and an outlook. *Beverages*, **6**(2), 23.https://doi.org/10.3390/beverages6020023

Liao Q, Hang X, Liu X, Pan J, Zhang H, Yang H (2010) The influence of pH on heat stress response by probiotic *Lactobacillus plantarum* LP-Onlly. *Annals of Microbiology*, **60**, 341-348. https://doi.org/10.1007/s13213-010-0048-x

Lorentzen M P G, Lucas P M (2019) Distribution of *Oenococcus oeni* populations in natural habitats. *Applied Microbiology and Biotechnology*, **103**, 2937–2945. https://doi.org/10.1007/s00253-019-09689-z

Ma J, Xu C, Liu F, Hou J, Shao H, Yu W (2021) Stress adaptation and cross-protection of *Lactobacillus plantarum* KLDS 1.0628. *CyTA Journal of Food*, **19**(1), 72–80.https://doi.org/10.1080/19476337.2020.1859619

Mao B, Yin R, Li X, Cui S, Zhang H, Zhao J, Chen W (2021) Comparative genomic analysis of *Lactobacillus plantarum* isolated from different niches. *Genes*, **12**(2), 241. https://doi.org/10.3390/genes12020241

Margalef-Català M, Araque I, Bordons A, Reguant C, Bautista-Gallego J (2016) Transcriptomic and proteomic analysis of *Oenococcus oeni* adaptation to wine stress conditions. *Frontiers inMicrobiology*, **7**, 1554. https://doi.org/10.3389/fmicb.2016.01554

Margalef-Català M, Araque I, Bordons A, Reguant C (2017) Genetic and transcriptional study of glutathione metabolism in *Oenococcus oeni*. *International Journal of Food Microbiology*, **242**,61-69. https://doi.org/10.1016/j.ijfoodmicro.2016.11.013

Miller B J, Franz C M A P, Cho G S, du Toit M (2011) Expression of the malolactic enzyme gene
(*mle*) from *Lactobacillus plantarum* under winemaking conditions. *Current Microbiology*,
62,1682-1688. https://doi.org/10.1007/s00284-011-9914-4
Miller R V, Kokjohn T A (1990) General microbiology of recA: environmental and evolutionary
significance.   *Annual   Reviews*,   **44**,   365-
394.https://doi.org/10.1146/annurev.mi.44.100190.002053
Nisiotou A A, Dourou D, Filippousi M E, Diamantea E, Fragkoulis P, Tassou C, Banilas G (2015)
Genetic and technological characterization of vineyard- and winery-associated lactic
acidbacteria. *BioMed Research International*, **508254**. https://doi.org/10.1155/2015/508254
Olguin N, Bordons A, Reguant C (2010) Multigenic expression analysis as an approach to
understanding the behaviour of *Oenococcus oeni* in wine-like conditions. *International
Journalof Food Microbiology*, **144**, 88–95. https://doi.org/10.1016/j.ijfoodmicr o.2010.08.032
Olguin N, Champomier-Vergès M, Anglade P, Baraige F, Cordero-Otero R, Bordons A, Zagorec
M, Reguant C (2015) Transcriptomic and proteomic analysis of *Oenococcus oeni* PSU-1
response   to   ethanol   shock.   *Food   Microbiology*,   **51**,   87–
95.https://doi.org/10.1016/j.fm.2015.05.005
Olguin N T, Valdés la Hens D, Delfederico L, Semorile L (2019) Relative expression of stress-
related genes during acclimation at low temperatures of psychrotrophic *Oenococcus oeni*
strains from Patagonian wine. *World Journal of Microbiology and Biotechnology*, **35**(5), 1-
9.https://doi.org/10.1007/s11274-018-2577-6
Papadimitriou K, Alegría A, Bron P A, de Angelis M, Gobbetti M, Kleerebezem M, Lemos J A,
Linares D M, Ross P, Stanton C, Turroni F, van Sinderen D, Varmanen P, Ventura M, Zúñiga
M, Tsakalidou E, Kok J (2016) Stress physiology of lactic acid bacteria. *Microbiology
andMolecular Biology Reviews*, **80**, 837-890. https://doi.org/10.1128/mmbr.00076-15
Portillo M, Del C, Franquès J, Araque I, Reguant C, Bordons A (2016) Bacterial diversity of
Grenache and Carignan grape surface from different vineyards at Priorat wine region
(Catalonia,   Spain).   *International   Journal   of   Food   Microbiology*,   **219**,   56–
63.https://doi.org/10.1016/j.ijfoodmicro.2015.12.002
Reverón I, de las Rivas B, Muñoz R, López de Felipe F (2012) Genome-wide transcriptomic
responses of a human isolate of *Lactobacillus plantarum* exposed to p-coumaric acid
stress.*Molecular   Nutrition   Food   Research*,   **56**,   1848-1859.
https://doi.org/10.1002/mnfr.201200384
Reverón I, de las Rivas B, Matesanz R, Muñoz R, López de Felipe F (2015) Molecular
adaptation of *Lactobacillus plantarum* WCFS1 to gallic acid revealed by genome-scale
transcriptomic signature and physiological analysis. *Microbial Cell Factories*, **14**, 160.
https://doi.org/10.1186/s12934-015-0345-y
Reverón I, Plaza-Vinuesa L, Santamaría L, Oliveros J C, de las Rivas B, Muñoz R, López de
Felipe F (2020) Transcriptomic evidence of molecular mechanisms underlying the response of
Lactobacillus   plantarum   WCFS1   to   hydroxytyrosol.   *Antioxidants*,   **9**(5),
422.https://doi.org/10.3390/antiox9050442
Renouf V, Claisse O, Miot-Sertier C, Lonvaud-Funel A (2006) Lactic acid bacteria evolution
during winemaking: Use of *rpoB* gene as a target for PCR-DGGE analysis. *Food
Microbiology*, **23**(2),136-145. https://doi.org/10.1016/j.fm.2005.01.019
Serrano L M, Molenaar D, Wels M, Teusink B, Bron P A, de Vos W M, Smid E J (2007)
Thioredoxin reductase is a key factor in the oxidative stress response of *Lactobacillus
plantarum* WCFS1.*Microbial Cell Factories*, **6**, 29. https://doi.org/10.1186/1475-2859-6-29
Schmittgen T D, Livak K J (2008) Analyzing real-time PCR data by the comparative CT method.
Nature Protocols, **3**, 1101–1108. https://doi.org/10.1038/nprot .2008.73
Valdés La Hens D, Bravo-Ferrada B M, Delfederico L, Caballero A C, Semorile L C (2015)
Prevalence of *Lactobacillus plantarum* and *Oenococcus oeni* during spontaneous malolactic
fermentation in Patagonian red wines revealed by polymerase chain reaction-denaturing
gradient gel electrophoresis with two targeted genes. *Australian Journal of Grape and
WineResearch*, **21**, 49–56. https://doi.org/10.1111/ajgw.12110

Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002)
Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of
multiple internal control genes. *Genome Biology*, **3**, 31–34. https://doi.org/10.1186/gb-2002-
37-research0034
Vitulo N, Fernandes Lemos W J, Calgaro L J M, Confalone M, Felis G E, Zapparoli G, Nardi T
(2019) Bark and Grape Microbiome of *Vitis viniferna*: influence of geographic patterns and
agronomic management on bacterial diversity. *Frontiers in Microbiology*, **9**,
3203.https://doi.org/10.3389/fmicb.2018.03203
Zheng J, Wittouck S, Salvetti E, Franz C M A P, Harris H M B, Mattarelli P, O'Toole P W, Pot B,
Vandamme P, Walter J, Watanabe K, Wuyts S, Felis G E, Ganzle M G, Lebeer S (2020) A
taxonomic note on the genus *Lactobacillus*: description of 23 novel genera, emended
description of the genus *Lactobacillus* Beijerinck 1901, and union of *Lactobacillaceae* and
Leuconostocaceae. *International Journal of Systematic and Evolutionary Microbiology*, **70**,
2782-2858. https://doi.org/10.1099/ijsem.0.004107
495