Evaluation of potential stress marker genes during acclimation of the Patagonian *Lactiplantibacillus plantarum* UNQLp11 strain

Antonella Zerbino[1,2], Lucas Lannutti[1,2], Nair Temis Olguin[1,2*]

[1]Departamento de Ciencia y Tecnología, Universidad de Morón, Machado 914, Motón, Buenos Aires, Argentina.

[2]Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Godoy Cruz 2290 (C1425FQB) CABA, Argentina.

Authors E-mail address: azerbino@unimoron.edu.ar, llannutti@unimoron.edu.ar, ntolguin@gmail.com

*Corresponding author: Nair Temis Olguin

Address: Machado 914, (B1876EOH) Morón, Buenos Aires, Argentina

Phone: + 54-11-56272000 int. 732

E-mail: ntolguin@gmail.com

ORCID: 0000-0001-5742-9236

Abstract

In this study, we evaluated the expression of genes that may serve as markers of temperature- and ethanol-induced stress responses during the acclimation of *Lactiplantibacillus plantarum* strains of enological interest. The transcriptional responses of eight genes potentially related to stress adaptation were analyzed in two strains acclimated at 18 °C and 21 °C. Gene expression was assessed at the start (time zero) and after 48 hours of acclimation. Following acclimation, the cells were inoculated into sterile Pinot noir wine, and L-malic acid consumption was monitored over 19 days to determine whether acclimation temperature influenced cell survival and fermentation performance. Both strains exhibited similar L-malic acid consumption and survival patterns, although transcriptional responses varied depending on the temperature. Notably, *gapB*, *glmS*, and *rfbB* emerged as promising gene markers for future studies. These findings suggest that gene expression profiling of acclimated cells may support the selection of robust *Lpb. plantarum* strains for use as starter cultures in winemaking.

Keywords

gene expression, L-malic acid consumption, Real Time RT-qPCR, acclimation, *Lactiplantibacillus plantarum*

1. Introduction

In wine, lactic acid bacteria (LAB) are responsible for malolactic fermentation (MLF), a decarboxylation process in which L-malic acid is converted into L-lactic acid. This process is desirable in most red wines and in some white and sparkling wines for three main reasons: it lowers acidity (leading to a softer mouthfeel), confers microbiological stability, and it modifies the wine's flavor (Bartowsky, 2017). Several LAB species can perform MLF, with *Oenococcus oeni* being the main responsible in many countries. As such, it is the most extensively studied and widely used as starter culture (Renouf *et al.*, 2006; Nisiotou *et al.*, 2015; Bartowsky, 2017). Recently, *Lactiplantibacillus plantarum* (formerly *Lactobacillus plantarum*) has emerged as an alternative species. It possesses a more diverse enzymatic profile than *O. oeni,* which may significantly influence flavor development (Iorizzo *et al.* 2016; Krieger-Weber *et al.*, 2020). Additionally, *Lpb. plantarum* has the genetic potential to produce antimicrobial peptides (plantaricins), which could enhance its dominance during MLF and offer the added benefits of reducing the need of sulphur dioxide (SO_2) commonly used to inhibit undesirable microbial growth (du Toit *et al.*, 2011).

Wine presents a harsh environment due to its high ethanol content, low pH, the presence of SO_2 and other inhibitory compounds. Not all LAB species or strains can survive under such conditions, but *Lpb. plantarum* has demonstrated stress tolerance comparable to that of *O. oeni* (Iorizzo *et al.*, 2016; Krieger-Weber *et al.*, 2020). Thanks to transcriptomic and other omics-bases analyses, we are beginning to understand the gene and protein expression profiles that contribute to LAB survival in wine (Olguin *et al.*, 2010; 2015; 2019; Bordas *et al.*, 2015; Margalef-Català *et al.*, 2016). However, most of this research has focused on *O. oeni*, while studies involving *Lpb. plantarum* remain limited (Fiocco *et al.*, 2007; Miller *et al.*, 2011).

One of the main criteria for selecting LAB strains as starter cultures is their resistance to wine conditions and their metabolic performance. However, although commercial starter cultures are widely available, a limitation is that they are often derived from specific regions and may not adequately capture the microbial diversity representative of other terroirs. The potential contribution of microorganisms to the concept of terroir has long been overlooked (Knight et al., 2015). These microbes not only participate in fermentation but also influence soil chemistry and nutrition, as well as grapevine health, yield, and quality (Belda et al., 2017). Several studies have linked wine characteristics to regional microbial communities (Vitulo et al., 2019), with some specifically highlighting the role of bacteria in shaping wine quality (Portillo et al., 2016; Grangateau et al., 2017; Lorentzen and Lucas, 2019). Therefore, it is important to evaluate whether isolated microbial strains are beneficial within a given terroir. In some cases, the preservation of terroir itself may guide the isolation, identification, and characterization of strains for use as starter cultures.

LAB populations in Patagonian wines display high species and strain diversity during spontaneous
MLF (Valdés La Hens *et al*., 2015). In previous work, we applied an acclimation protocol (Bravo-
Ferrada *et al*., 2014), in which cells were cultured in a broth with moderate ethanol concentration
prior to wine inoculation. We investigated the influence of two temperatures (18 °C and 21 °C)
on gene expression using *O. oeni* strains isolated from Patagonian Pinot noir wines (Olguin *et al*.,
2019). We found that L-malic acid consumption, population survival, and transcriptional behavior
differed depending on acclimation temperature.

The aim of the present study was to evaluate the expression of stress-related genes in *Lpb.*
plantarum UNQLp11 induced by acclimation at two temperatures (18 °C and 21 °C). Gene
selection was based on previous studies involving both *O. oeni* (Olguin *et al*., 2010; 2019) and *Lpb.*
plantarum (Fiocco *et al*., 2007; Miller *et al*., 2011). Gene expression was quantified by reverse
transcription real-time quantitative PCR (RT-qPCR). Population survival and L-malic acid
consumption after acclimation were monitored in Pinot noir wine.

2. Materials and methods
2.1. Growth conditions
Lpb. plantarum subsp. *plantarum* ATCC 14917 (DSM 20174, Zheng *et al*., 2020) and the strain
UNQLp11 (RGPRN-/-M-/-357-/-Semorile-/-626-/-UNQLp11-/-2024), previously isolated from
Patagonian red wines (Bravo-Ferrada *et al*., 2013; Iglesias *et al*., 2020) where cultured at 28 °C in
tubes containing MRS broth medium (De Man *et al*., 1960), supplemented with L-malic acid (4 gL^{-1})
and fructose (5 gL^{-1}) at pH 5.0. After 48 h cells were harvested by centrifugation and resuspended
in the acclimation medium (50 gL^{-1} MRS, 40 gL^{-1} fructose, 20 gL^{-1} glucose, 4.5 gL^{-1} L-malate, 1
gL^{-1} Tween 80, 0.1 mgL^{-1} pyridoxine, pH 4.6) containing 6% (v/v) of ethanol (Bravo-Ferrada *et al*.,
2014). One of the acclimation tubes was cultured at 18 °C and the other one at 21 °C. We chose
these temperatures because both are common during MLF in wineries. Samples were taken at time
zero (at the beginning of cell acclimation) and 48 h after acclimation (just before wine inoculation).
Then, cells were harvested by centrifugation and inoculated in tubes containing sterile Pinot noir
wine (vintage 2014), at the final stage of alcoholic fermentation (14.3 % ethanol, pH 3.7, 2.3 gL^{-1}
L-malic acid). These tubes were kept at 21 °C and cell survival was analysed every 2-3 days until
the 19th day. All assays were performed in duplicate at different times using independent
precultures and cell survival was monitored by counting colonies on plates of MRS medium
supplemented as described above. L-malic acid consumption was measured using the L-Malic Acid
Enology Enzymatic kit (BioSystems SA, Barcelona, Spain) according to the manufacturer's
instructions.

Sovereign Rights Over Natural Resources: The indigenous LAB strain mentioned above is a genetic
resources belonging to the Provinces of Rio Negro (UNQLp11). It was deposited in the strain
collection of the Laboratorio de Microbiología Molecular (IMBA–DCyT–UNQ-CIC) and registered
with the Environment and Climate Change Secretariat— Province of Rio Negro Government. Their
preservation follows the protocols of the Convention on Biological Diversity and the Nagoya
Protocol, which safeguard the ownership and use of genetic resources from the regions where they
were isolated (RESOL-2024-626-E-GDERNE-SAYCC#SGG, NO-2024-00628737-GDERNE-
SAYCC#SGG).
150

2.2. RNA extraction

152 *Lpb. plantarum* cells were harvested by centrifugation and frozen until RNA extraction. Total RNA
153 extractions were performed using the SV Total RNA Isolation System #Z3100 (Promega) following
154 the manufacturer's instructions. RNA concentrations were calculated by measuring absorbance at
155 260 nm using a Nanodrop spectrophotometer (Thermo Fisher Scientific, 1000).
156

2.3. Selection of candidate genes and nucleotide sequences

The gene sequences of *Lpb. plantarum* reference strain SRCM100442 (Genome
assembly ASM991365v1) were obtained from the National Center for Biotechnology Information
(NCBI). The reference genome assembly strain SRCM100442 was used for sequence comparison
against *O. oeni* PSU-1 (NC_008528) (Table 1) and for primers design when applicable (Table 2).
We also tested sequence alignment and *in silico* PCR with the strain ATCC 14917 gene sequences.
Sequence alignment and primer pairs were tested in silico using the NCBI Needleman-Wunsch
Global Align Nucleotide Sequences and Primer-BLAST, respectively
(https://blast.ncbi.nlm.nih.gov/Blast.cgi).
166

2.4. Reverse transcription and RT-qPCR

cDNA was synthesized from RNA (15 ng μL^{-1}) using TaqMan Reverse Transcription Reagents
(Applied Bio- systems) as recommended. Ten pairs of primers were designed (Table 2) to be about
18–22 bases long, to contain over 50 % G/C and to have a melting temperature (Tm) above 60 °C.
Two pairs of primers were taken from previous works (Fiocco *et al*., 2008; Duary *et al*., 2012). The
length of the PCR products ranged from 95 to 127 bp (Table 2). RT-qPCR was performed in 12 μL
final volume containing 1.5 ng of total cDNA, 1.0 μL of each primer at 3.15 $\mu mol\ L^{-1}$, 2.5 μL of
RNase free water and 6.5 μL of SYBR Green PCR Master Mix (Applied Biosystems).
Amplifications were carried out using a QuantStudio 3 Real-Time PCR Systems (Applied
Biosystems) with an initial step at 95 °C for 5 min followed by 35 cycles of 95 °C for 10 s and 60

°C for 30 s and 72 ℃ for 30 s. An additional step to establish a melting curve was used to verify the

specificity of the real-time PCR reaction for each primer pair. The threshold value used in this study

was automatically determined by the instrument.

Results were analysed using the comparative C_T method ($2^{-\Delta\Delta CT}$) as described by Schmittgen and

Livak (2008). Two statistical algorithms were used to determine the stability of the candidate

reference genes: NormFinder (Andersen *et al*., 2004) and geNorm (Vandesompele *et al*., 2002). The

three candidates, *ldhD*, *rpoD* and *gyrA* were chosen as the best combination.

3. Results

3.1. Cells survival and L-malic acid consumption during acclimation

The *Lpb. plantarum* strain used in this study, isolated from a Patagonian Pinot noir wine undergoing

spontaneous MLF, was identified as UNQLp11 (Bravo-Ferrada *et al*., 2013). It was selected from a

pool of strains obtained from the same wine based on its oenological properties (Brizuela *et al*.,

2018). *Lpb. plantarum* DSM 20174, isolated from pickled cabbage, was used as a reference strain.

Since this strain originates from a non-wine environment, any observed metabolic or genetic

similarities are likely species-specific rather than environmentally induced.

Acclimation temperatures of 18 °C and 21 °C were selected based on prior work showing

temperature-dependent behavior in *O. oeni* strains (Olguin *et al*., 2019).

During the 48-hour acclimation period, cell density increased by approximately two logarithmic

units (LOGs) at both temperatures for both strains.

3.2. Gene expression after 48 hours acclimation

Genes from *Lpb. plantarum* were selected based on their known functions, under the hypothesis that

they may respond to wine-associated stress conditions similarly to *O. oeni* (Table 1). Prior to

quantitative analysis, RT-qPCR amplification of the selected genes was tested in both strains. The

amplified fragment sizes were as expected (Table 2). Amplification efficiency was calculated for

each primer pair; the gene *dacA* was excluded due to poor efficiency (E = 80.21%, R^2 = 0.9958).

All remaining primers had efficiencies within 10% of each other, making them suitable for relative

quantification using the comparative Ct ($\Delta\Delta$Ct) method (Schmittgen and Livak, 2008).

Relative gene expression was first analyzed using the sample at time zero under acclimation

conditions (A0) as the calibrator (Fig. 1). At 18 °C, the genes *rfbB* (also referred to as *rmlB*)

and *glmS* were clearly overexpressed, especially in the Patagonian strain UNQLp11 (Fig. 1A). In

contrast, *recA* was slightly overexpressed in the reference strain ATCC 14917 but underexpressed

in UNQLp11. Genes *dnaK*, *trxA*, *cspA*, and *gapB* were consistently underexpressed in both strains

at this temperature.

At 21 °C, gene expression trends differed (Fig. 1B). In ATCC 14917, most genes showed stable or
reduced expression, except for a slight overexpression of *cspA* and a marked overexpression
of *gapB*. In UNQLp11, there was slight overexpression of *glmS* and *trxA*, with *gapB* again notably
overexpressed. All other genes were stably expressed, with the exception of *rfbB*, which was
significantly underexpressed.
To assess the impact of temperature, gene expression levels after 48 hours of acclimation (A48)
were compared between 18 °C and 21 °C, using the latter as the calibrator condition (Fig. 2).
Expression patterns were largely consistent between strains. Both strains showed strong
overexpression of *rfbB* and slight overexpression of *glmS*. Genes *dnaK*, *trxA*, *cspA*, and *gapB* were
underexpressed under the lower temperature. Differences between strains were limited to slight
overexpression of *hsp20* and *recA* in the ATCC 14917 strain.
223
3.3. Cell survival and L malic acid consumption in wine
Cells acclimated at both temperatures (18 °C and 21 °C) were inoculated into sterile Pinot noir
wine and monitored over 19 days. Cell survival and L-malic acid consumption were evaluated at
different time points (Fig. 3).
Viable cell counts for both ATCC 14917 and UNQLp11 declined during the fermentation period
(Fig. 3A). However, both strains were capable of consuming the majority of the L-malic acid within
the first seven days of incubation (Fig. 3B).
231
4. Discussion
The aim of this study was to expand our understanding of the genetic mechanisms involved in the
acclimation of *Lpb. plantarum*, which may enhance its survival under the stressful conditions
present in wine. Previous studies have demonstrated that wine-associated LAB, such as *O.*
oeni and *Lpb. plantarum*, benefit from an acclimation step prior to inoculation into wine (Cecconi *et*
al., 2009; Bravo-Ferrada *et al*., 2013; Costantini *et al*., 2015). Stress-related genes in *O. oeni* have
been proposed as markers of cellular adaptation (Hamon *et al*., 2013), but the genetic response
of *Lpb. plantarum* to wine-like conditions remains poorly characterized. Thus, we evaluated the
expression of stress-related genes in two *Lpb. plantarum* strains subjected to acclimation at two
different temperatures.
Sequence comparisons between *O. oeni* and *Lpb. plantarum* indicate that while their genes and
proteins perform homologous functions, their regulatory mechanisms can differ. It is known that
different stress-resistance phenotypes may protect against the same environmental stressor, even
when the molecular pathways involved are distinct (Papadimitriou *et al*., 2016). Therefore, it is not

unexpected that some genes in *Lpb. plantarum* respond similarly to those in *O. oeni*, while others

do not (Fig. 1).

The gene expression profiles of strains UNQLp11 and ATCC 14917 showed broadly similar

responses, which corresponded with their survival and L-malic acid consumption in wine following

acclimation. Among the stress response genes (*hsp20*, *cspA*, *dnaK*, and *trxA*), we observed general

under-expression at 18 °C. These temperature conditions are moderate, and more extreme values

are typically required to elicit strong responses in heat-shock or cold-shock genes (Liao *et al*., 2010;

Fiocco *et al*., 2007). Nevertheless, due to the potential for cross-protection mechanisms

(Papadimitriou *et al*., 2016; Ma *et al*., 2021), we included both *hsp20* and *cspA* in our analysis. The

under-expression of *cspA* may align with previous findings in *O. oeni*, where the protein was

downregulated following acclimation in 8% ethanol (Cecconi *et al*., 2009a).

While earlier studies reported that *Lpb. plantarum* strains overexpressing heat shock proteins were

more ethanol-tolerant (Fiocco *et al*., 2007), our results differ from some previous findings. For

example, *hsp* gene expression in *Lpb. plantarum* has been correlated with that in *O. oeni* (Capozzi

et al., 2012), but in our study, *hsp20* did not show significant upregulation, consistent with previous

work on *O. oeni* under similar conditions (Olguin *et al*., 2019). Similarly, although *dnaK* expression

has been shown to increase under various stress conditions (Liao *et al*., 2010; Reverón *et al*., 2012),

we did not observe this in our study, again aligning with earlier findings in *O. oeni* (Olguin *et al*.,

2019). The gene *trxA*, typically upregulated under oxidative and ethanol stress in both species

(Serrano *et al*., 2007; Reverón *et al*., 2012; Margalef-Català *et al*., 2017), also did not show the

expected increase in our experiment.

With respect to carbohydrate metabolism, *gapB* is often considered a housekeeping gene (Duray *et*

al., 2012), yet in our study, its expression appeared temperature-dependent. This aligns with

previous observations where *gapB* was upregulated under oxidative stress in specific *Lpb.*

plantarum strains (Serrano *et al*., 2007). Additionally, other studies have reported changes in

protein expression linked to carbohydrate metabolism in response to different culture conditions in

both *Lpb. plantarum* (Cecconi *et al*., 2009b; Hamon *et al*., 2013) and *O. oeni* (Cecconi *et al*., 2009a;

Olguin *et al*., 2015; Costantini *et al*., 2015).

In the context of amino acid metabolism, we analyzed *glmS*, which showed a notable increase in

expression, especially at 18 °C. Previous work has shown that *glmS* is downregulated in response

to antioxidant stress induced by hydroxytyrosol or gallic acid (Reverón *et al*., 2015; 2020), and its

expression also decreases in *O. oeni* following ethanol shock (Olguin *et al*., 2015). Our results

contrast with those findings, suggesting a different regulatory pattern under temperature-induced

stress.

The *recA* gene, associated with DNA repair and recombination, has been variously classified as a

housekeeping or stress-related gene (Miller and Kokjohn, 1990; Curiel *et al*., 2011; Reverón *et al*.,

2012, 2015). In our study, its expression remained relatively stable under both conditions.

An especially interesting result was observed for *rfbB*, a gene involved in cell envelope biogenesis.

It was strongly overexpressed at 18 °C but downregulated at 21 °C. Previous studies have

reported no significant change in *rfbB* protein levels under acidic conditions in *Lpb.*

plantarum (Hamon *et al*., 2013), while in *O. oeni*, both over- and under-expression have been

observed depending on the acclimation conditions (Olguin *et al*., 2015; 2019).

The impact of acclimation on both *O. oeni* and *Lpb. plantarum* using the same medium has been

studied in previous research (Bravo-Ferrada *et al*., 2014; Brizuela *et al*., 2018). Thus, the similar

survival and L-malic acid consumption profiles observed in both *Lpb. plantarum* strains in this

study were not unexpected. Notably, strain UNQLp11 was isolated from Patagonian Pinot noir

wine, and the selection of indigenous strains as starter cultures is gaining global interest for

enhancing regional wine identity (Bartowsky *et al*., 2017). As suggested by Mao *et al*. (2021),

environmental selection pressure can drive genomic changes that improve strain adaptability to

specific ecological niches. Therefore, while our results are promising, further research using

additional wine-isolated *Lpb. plantarum* strains is necessary to validate these findings.

To our knowledge, this is the first study to evaluate stress-related gene expression in *Lpb.*

plantarum as potential markers for winemaking applications. As the use of this species in starter

culture development continues to grow, further research incorporating gene expression and omics

approaches will be valuable. Similar to ongoing work in *O. oeni*, these studies can help correlate

genetic behavior with enological performance, ultimately supporting the selection of robust,

regionally adapted strains for the wine industry.

5. Conclusions

This study offers new insights into the genetic mechanisms underlying the acclimation of *Lpb.*

plantarum to wine conditions—an area that remains less understood compared to *O. oeni*. By

analyzing the gene expression profiles of *Lpb. plantarum* strains at two temperatures, we identified

both shared and distinct responses relative to *O. oeni*. While several stress-related genes

(*hsp20*, *cspA*, *dnaK*, *trxA*) were under-expressed, genes such as *glmS* and *rfbB* showed temperature-

dependent overexpression.

These findings highlight the potential of *Lpb. plantarum* as a viable candidate for use in starter

culture development, particularly when selecting indigenous strains to support regional wine

identity. Overall, the results deepen our understanding of *Lpb. plantarum*'s stress adaptation and

support its further exploration for enological applications, laying the groundwork for future studies

aimed at optimizing strain selection and performance in winemaking.

Data availability

Data sharing not applicable to this article as no datasets were generated or analysed during the

current study.

Conflicts of interest

The authors have no conflicts of interest to declare.

Funding statement

This work was funded by grants from CONICET [PIBAA 2022 No 28720210100183CO], Agencia

Nacional de Promoción Científica y Tecnológica, [PICT 2021 GRF-TI-00350] and, University of

Morón [PICT 2024 - 80020230100046UM], Buenos Aires, Argentina. AZ and LL are PhD students

from CONICET-UM. NTO is a member of the Research Career of CONICET.

Acknowledgements

We would like to thank the Environment and Climate Change Secretariat, Government of the

Province of Rio Negro, for authorizing access to provincial genetic resources and their use for

academic and research purposes. The authors also thank the Laboratorio de Microbiología

Molecular (LMM) of the Universidad Nacional de Quilmes (UNQ) for lending them the equipment

to perform the Real Time RT-qPCR.

References

Andersen, C, Jensen, J, & Orntoft, T. (2004). Normalization of real time quantitative reverse

transcription—PCR data: a model based variance estimation approach to identify genes suited for

normalization, applied to bladder and colon cancer data sets. *Cancer Research*, 64:5245.

https://doi.org/10.1158/0008

Bartowsky, E. J, (2017). *Oenococcus oeni* and the genomic era. *FEMS Microbiology*

Reviews, 41(Supp_1), S84-S94. https://doi.org/10.1093/femsre/fux034

Belda, I, Zarraonaindia, I, Perisin, M, Palacios, A, & Acedo, A. (2017). From vineyard soil to wine

fermentation: microbiome approximations to explain the "terroir" concept. *Frontiers in*

Microbiology, 8, 821. https://doi.org/10.3389/fmicb.2017.00821

Bordas, M, Araque, I, Bordons, & A, Reguant, C. (2015). Differential expression of selected
Oenococcus oeni genes for adaptation in wine-like media and red wine. *Annals of Microbiology,* 65,
2277–2285. https://doi.org/10.1007/s1321 3-015-1069-2

Bravo-Ferrada, B. M, Hollmann, A, Delfederico, L, Valdes La Hens, D, Caballero, A, & Semorile,
L. (2013). Patagonian red wines: selection of *Lactobacillus plantarum* isolates as potential starter
cultures for malolactic fermentation. *World Journal of Microbiology and Biotechnology*, 29, 1537–
1549.

Bravo-Ferrada, B. M, Tymczyszyn, E. E, Gómez-Zavaglia, A, & Semorile, L. (2014). Effect of
acclimation medium on cell viability, membrane integrity and ability to consume malic acid in
synthetic wine by oenological *Lactobacillus plantarum* strains. *Journal of Applied*
*Microbiology,*116, 360–367. https://doi.org/10.1111/jam.12372

Brizuela, N. S, Bravo-Ferrada, B. M, Pozo-Bayón, M. A, Semorile, L, & Tymczyszyn, E. E. (2018).
Changes in the volatile profile of Pinot noir wines caused by Patagonian *Lactobacillus plantarum*
and *Oenococcus oeni* strains. *Food Research International*, 106, 22–8.
https://doi.org/10.1016/j.foodres.2017.12.032.

Capozzi, V, Fiocco, D, Weidmann, S, Guzzo, J, & Spano, G. (2012). Increasing membrane
protection in *Lactobacillus plantarum* cells overproducing small heat shock proteins. *Annals of*
Microbiology, 62, 517-522. https://doi.org/10.1007/s13213-011-0285-7

Cecconi, D, Milli, A, Rinalducci, S, Zolla, L, & Zapparoli, G. (2009a). Proteomic analysis of
Oenococcus oeni freeze-dried culture to assess the importance of cell acclimation to conduct
malolactic fermentation in wine. *Electrophoresis*, 30, 2988–2995.
https://doi.org/10.1002/elps.200900228

Cecconi, D, Cristofoletti, M, Milli, A, Rinalducci, S, Zolla, L, & Zapparoli, G. (2009b). Effect of
tannic acid on *Lactobacillus plantarum* wine strain during starvation: a proteomic study.
Electrophoresis, 30, 957–965. https://doi.org/10.1002/elps.200800310

Costantini, A, Rantsiou, K, Majumder, A, Jacobsen, S, Pessione, E, Svensson, B, Garcia-Moruno,
E, & Cocolin, L. (2015). Complementing DIGE proteomics and DNA subarray analyses to shed
light on *Oenococcus oeni* adaptation to ethanol in wine-simulated conditions. *Journal of*
Proteomics, 123, 114–127. https://doi.org/10.1016/j.jprot.2015.04.019

Curiel, J. A, Rodríguez, H, de las Nieves, B, Anglade, P, Baraige, F, Zagorec, M, Champomier-
Vergès, M, Muñoz, R, & López de Felipe, F. (2011). Response of *Lactobacillus plantarum* human
isolate to tannic acid challenge assessed by proteomic analyses. *Molecular Nutrition Food*
Research, 55, 1454-1465. https://doi.org/10.1002/mnfr.201000621

De Man, J. C, Rogosa, M, & Sharpe, M. E. (1960). A medium for the cultivation of Lactobacilli.
Journal of Applied Bacteriology, 23, 130–135. https://doi.org/10.1111/j.1365-2672.1960.tb00188.x

Duary, R. K, Batish, V. K, & Grover, S. (2012). Relative gene expression of bile salt hydrolase and

surface proteins in two putative indigenous *Lactobacillus plantarum* strains under in vitro gut

conditions. *Molecular Biology Reports*, 39, 2541-2552. https://doi.org/10.1007/s11033-011-1006-9

du Toit, M, Engelbrecht, L, Lerm, E, & Krieger-Weber, S. (2011). *Lactobacillus*: the next

generation of malolactic fermentation starter cultures – an overview. *Food and Bioprocess*

Technology, 4, 876-906. https://doi.org/10.1007/s11947-010-0448-8

Fiocco, D, Capozzi, V, Goffin, P, Hols, P, & Spano, G. (2007). Improved adaptation to heat, cold,

and solvent tolerance in *Lactobacillus plantarum. Applied Microbiology and Biotechnology*, 77,

909-915. https://doi.org/10.1007/s00253-007-1228-x

Fiocco, D, Crisetti, E, Capozzi, V, & Spano, G. (2008). Validation of an internal control gene to

apply reverse transcription quantitative PCR to study heat, cold and ethanol stresses in

Lactobacillus plantarum. World Journal of Microbiology and Biotechnology, 24, 899-902.

https://doi.org/10.1007/s11274-007-9556-7

Grangeteau, C, Roullier-Gall, C, Rousseaux, S, Gougeon, R. D, Schmitt-Kopplin, P, Alexandre, H,

& Guilloux-Benatier, M. (2017). Wine microbiology is driven by vineyard and winery

anthropogenic factors. *Microbial Biotechnology*, 10, 354–370. https://doi.org/10.1111/1751-

7915.12428

Hamon, E, Horvatovich, P, Marchioni, E, Aoudé-Werner, D, & Ennahar, S. (2013). Investigation of

potential markers of acid resistance in *Lactobacillus plantarum* by comparative proteomics. *Journal*

of Applied Microbiology, 116, 134-144. https://doi.org/10.1111/jam.12339

Iglesias, N. G, Brizuela, N. S, Tymczyszyn, E. E, Hollmann, A, Valdés La Hens, D, Semorile, L, &

Bravo-Ferrada, B. M. (2020). Complete genome sequencing of *Lactobacillus plantarum* UNQLp11

isolated from a Patagonian Pinot noir wine. South African Journal of Enology and Viticulture,

41(2), 197-209. https://doi.org/10.21548/41-2-4066

Iorizzo, M, Testa, B, Lombardi, S. J, García-Ruiz, A, Muñoz-González, C, Bartolomé, B, &

Moreno- Arribas, M. V. (2016). Selection and technological potential of *Lactobacillus plantarum*

bacteria suitable for wine malolactic fermentation and grape aroma release. *LWT Food Science and*

Technology, 73, 557-566. https://doi.org/10.1016/j.lwt.2016.06.062

Knight, S, Klaere, S, Fedrizzi, B, & Goddrard, M. R. (2015). Regional microbial signatures

positively correlate with differential wine phenotypes: evidence for a microbial aspect to terroir.

Scientific Reports, 5, 14233. https://doi.org/10.1038/srep14233

Krieger-Weber, S, Heras, J. M, & Suarez, C. (2020). *Lactobacillus plantarum*, a new biological tool

to control malolactic fermentation: a review and an outlook. *Beverages*, 6(2), 23.

https://doi.org/10.3390/beverages6020023

Liao, Q, Hang, X, Liu, X, Pan, J, Zhang, H, & Yang, H. (2010). The influence of pH on heat stress response by probiotic *Lactobacillus plantarum* LP-Onlly. *Annals of Microbiology*, 60, 341-348. https://doi.org/10.1007/s13213-010-0048-x

Lorentzen, M. P. G. & Lucas, P. M. (2019). Distribution of *Oenococcus oeni* populations in natural habitats. *Applied Microbiology and Biotechnology*, 103, 2937–2945. https://doi.org/10.1007/s00253-019-09689-z

Ma, J, Xu, C, Liu, F, Hou, J, Shao, H, & Yu, W. (2021). Stress adaptation and cross-protection of *Lactobacillus plantarum* KLDS 1.0628. *CyTA Journal of Food*, 19(1), 72–80. https://doi.org/10.1080/19476337.2020.1859619

Mao, B, Yin, R, Li, X, Cui, S, Zhang, H, Zhao, J, & Chen, W. (2021). Comparative genomic analysis of *Lactobacillus plantarum* isolated from different niches. *Genes*, 12(2), 241. https://doi.org/10.3390/genes12020241

Margalef-Català, M, Araque, I, Bordons, A, Reguant, C, & Bautista-Gallego, J. (2016). Transcriptomic and proteomic analysis of *Oenococcus oeni* adaptation to wine stress conditions. *Frontiers in Microbiology*, 7, 1554. https://doi.org/10.3389/fmicb.2016.01554

Margalef-Català, M, Araque, I, Bordons, A, & Reguant, C. (2017). Genetic and transcriptional study of glutathione metabolism in *Oenococcus oeni*. *International Journal of Food Microbiology*, 242, 61-69. https://doi.org/10.1016/j.ijfoodmicro.2016.11.013

Miller, B. J, Franz, C. M. A. P, Cho, G. S, & du Toit, M. (2011). Expression of the malolactic enzyme gene (*mle*) from *Lactobacillus plantarum* under winemaking conditions. *Current Microbiology*, 62, 1682-1688. https://doi.org/10.1007/s00284-011-9914-4

Miller, R. V, & Kokjohn, T. A. (1990). General microbiology of recA: environmental and evolutionary significance. *Annual Reviews*, 44, 365-394. https://doi.org/10.1146/annurev.mi.44.100190.002053

Nisiotou, A. A, Dourou, D, Filippousi, M. E, Diamantea, E, Fragkoulis, P, Tassou, C, & Banilas, G. (2015). Genetic and technological characterization of vineyard- and winery-associated lactic acid bacteria. *BioMed Research International*, 508254. https://doi.org/10.1155/2015/508254

Olguin, N, Bordons, A, & Reguant, C. (2010). Multigenic expression analysis as an approach to understanding the behaviour of *Oenococcus oeni* in wine-like conditions. *International Journal of Food Microbiology*, 144, 88–95. https://doi.org/10.1016/j.ijfoodmicr o.2010.08.032

Olguin, N, Champomier-Vergès, M, Anglade, P, Baraige, F, Cordero-Otero, R, Bordons, A, Zagorec, M, & Reguant, C. (2015). Transcriptomic and proteomic analysis of *Oenococcus oeni* PSU-1 response to ethanol shock. *Food Microbiology*, 51, 87–95. https://doi.org/10.1016/j.fm.2015.05.005

Olguin, N. T, Valdés la Hens, D, Delfederico, , & Semorile, L. (2019). Relative expression of

stress-related genes during acclimation at low temperatures of psychrotrophic *Oenococcus oeni*

strains from Patagonian wine. *World Journal of Microbiology and Biotechnology*, 35(5), 1-9.

https://doi.org/10.1007/s11274-018-2577-6

Papadimitriou, K, Alegría, A, Bron, P. A, de Angelis, M, Gobbetti, M, Kleerebezem, M, Lemos, J.

A, Linares, D. M, Ross, P, Stanton, C, Turroni, F, van Sinderen, D, Varmanen, P, Ventura, M,

Zúñiga, M, Tsakalidou, E, & Kok, J. (2016). Stress physiology of lactic acid bacteria. *Microbiology*

and Molecular Biology Reviews, 80, 837-890. https://doi.org/10.1128/mmbr.00076-15

Portillo, M, Del, C, Franquès, J, Araque, I, Reguant, C, and Bordons, A. (2016). Bacterial diversity

of Grenache and Carignan grape surface from different vineyards at Priorat wine region (Catalonia,

Spain). *International Journal of Food Microbiology*, 219, 56–63.

https://doi.org/10.1016/j.ijfoodmicro.2015.12.002

Reverón, I, de las Rivas, B, Muñoz, R, & López de Felipe, F. (2012). Genome-wide transcriptomic

responses of a human isolate of *Lactobacillus plantarum* exposed to p-coumaric acid stress.

Molecular Nutrition Food Research, 56, 1848-1859. https://doi.org/10.1002/mnfr.201200384

Reverón, I, de las Rivas, B, Matesanz, R, Muñoz, R, & López de Felipe, F. (2015). Molecular

adaptation of *Lactobacillus plantarum* WCFS1 to gallic acid revealed by genome-scale

transcriptomic signature and physiological analysis. *Microbial Cell Factories*, 14, 160.

https://doi.org/10.1186/s12934-015-0345-y

Reverón, I, Plaza-Vinuesa, L, Santamaría, L, Oliveros, J. C, de las Rivas, B, Muñoz, R, & López de

Felipe, F. (2020). Transcriptomic evidence of molecular mechanisms underlying the response of

Lactobacillus plantarum WCFS1 to hydroxytyrosol. *Antioxidants*, 9(5), 422.

https://doi.org/10.3390/antiox9050442

Renouf, V, Claisse, O, Miot-Sertier, C, & Lonvaud-Funel, A. (2006). Lactic acid bacteria evolution

during winemaking: Use of *rpoB* gene as a target for PCR-DGGE analysis. *Food Microbiology*,

23(2), 136-145. https://doi.org/10.1016/j.fm.2005.01.019

Serrano, L. M, Molenaar, D, Wels, M, Teusink, B, Bron, P. A, de Vos, W. M, & Smid, E. J. (2007).

Thioredoxin reductase is a key factor in the oxidative stress response of *Lactobacillus plantarum*

WCFS1. *Microbial Cell Factories*, 6, 29. https://doi.org/10.1186/1475-2859-6-29

Schmittgen, T. D, & Livak, K. J. (2008). Analyzing real-time PCR data by the comparative CT

method. *Nature Protocols*, 3, 1101–1108. https://doi.org/10.1038/nprot .2008.73

Valdés La Hens, D, Bravo-Ferrada, B. M, Delfederico, L, Caballero, A. C, & Semorile, L. C.

(2015). Prevalence of *Lactobacillus plantarum* and *Oenococcus oeni* during spontaneous malolactic

fermentation in Patagonian red wines revealed by polymerase chain reaction-denaturing gradient

gel electrophoresis with two targeted genes. *Australian Journal of Grape and Wine Research*, 21,

49–56. https://doi.org/10.1111/ajgw.12110

Vandesompele, J, De Preter, K, Pattyn, F, Poppe, B, Van Roy, N, De Paepe, A, & Speleman, F.

(2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of

multiple internal control genes. *Genome Biology*, 3, 31–34. https://doi.org/10.1186/gb-2002-3-7-

research0034

Vitulo, N, Fernandes Lemos, W. J, Calgaro, L. J. M, Confalone, M, Felis, G. E, Zapparoli, G, &

Nardi, T. (2019). Bark and Grape Microbiome of *Vitis viniferna*: influence of geographic patterns

and agronomic management on bacterial diversity. *Frontiers in Microbiology*, 9, 3203.

https://doi.org/10.3389/fmicb.2018.03203

Zheng, J, Wittouck, S, Salvetti, E, Franz, C. M. A. P, Harris, H. M. B, Mattarelli, P, O'Toole, P. W,

Pot, B, Vandamme, P, Walter, J, Watanabe, K, Wuyts, S, Felis, G. E, Ganzle, M. G, & Lebeer, S.

(2020). A taxonomic note on the genus *Lactobacillus*: description of 23 novel genera, emended

description of the genus *Lactobacillus* Beijerinck 1901, and union of *Lactobacillaceae* and

Leuconostocaceae. *International Journal of Systematic and Evolutionary Microbiology*, 70, 2782-

2858. https://doi.org/10.1099/ijsem.0.004107

Figure 1. Relative expression of eight genes related to stress response (*hsp20*, *rfbB*, *glmS*, *recA*,
***dnaK*, *trxA*, *cspA*, *gapB*) after 48 h acclimation at 18 °C (A) and at 21 °C (B). The calibration**
condition used was time zero when *Lactiplantibacillus plantarum* cells were inoculated into the
acclimation medium.

Figure 2. Relative expression of eight genes related to stress response (*hsp20*, *rfbB*, *glmS*, *recA*,

***dnaK*, *trxA*, *cspA*, *gapB*) in *Lactiplantibacillus plantarum* strains after 48 h acclimation at 18**

°C. The calibration condition used was time 48 h at 21 °C.

Figure 3. Evolution of cells viability (log CFU mL^{-1}) and L-malic acid consumption in Pinot

noir wine of the *Lactiplantibacillus plantarum* strains acclimated at 18 and 21 ºC. (A) ATCC

14917 (circles), UNQLp11 (squares), solid symbols and lines indicate strains acclimated at 18

ºC and empty symbols and dashed lines indicate strains acclimated at 21 ºC. (B) Percentage of

L-malic acid consumed by cells acclimated at 18 ºC (black bars) and at 21 ºC (grey bars) after

7 days (7d) in sterile wine.

Table 1. Comparison of genes sequence alignment between *Lactiplantibacillus plantarum* SRCM100442 and *Oenococcus oeni* PSU-1 (NC_008528).

Target gene *Lpb. plantarum*	Gene name	Gene sequence identity (%)	Protein sequence identity (%)	*O. oeni* gene symbol
C7M33_RS14920	*ldhD*	60	51	OEOE_RS01985
C7M33_RS06325	*gyrA*	66	69	OEOE_RS00030
C7M33_RS14590	*rpoD*	59	61	OEOE_RS04785
C7M33_RS01170	*recA*	65	68	OEOE_RS06650
C7M33_RS09485	*gapB*	72	72	OEOE_RS01940
C7M33_RS05025	*hsp20*	57	41	OEOE_RS01385
C7M33_RS11210	*rfbB (rmlB)*	57	45	OEOE_RS06990
C7M33_RS14780	*dnaK*	70	74	OEOE_RS06310
C7M33_RS09640	*glmS*	63	60	OEOE_RS03035
C7M33_RS01055	*trxA*	60	50	OEOE_RS08215
C7M33_RS10385	*cspA*	79	81	OEOE_RS06620
C7M33_RS04410	*dacA*	51	32	OEOE_RS03435

*O. oeni PSU-1 genes locus tag have been discontinued but are still accessible. Since the new sequences are variable, we decided to use the ones we already had to compare with the *Lpb. plantarum* reference genome assembly (strain SRCM100442).

550 **Table 2. Primers used in this work.**

Target gene	Description	Forward primer (5'→3')	Reverse primer (5'→3')	Amplicon length (bp)	Reference
ldhD	D-2-hydroxyacid dehydrogenase	ACGCCCAAGCTGATGTTATC	AGTGTCCCACGAGCAAAGTT	127	Fiocco et al. (2008)
recA	recombinase A	CCTTAGATGACGCGTTGG	CAGCAACCGCATGTAGTG	108	This work
gyrA	DNA gyrase subunit	GTCAAACGGACACCAGTTCAAG	GTGACCATCCGTGATTGTAACC	111	This work
gapB	type I glyceraldehyde 3-phosphate dehydrogenase	ACTGCCTTGCACCATTAGC	TACAGGGCCGTCAAGTAAC	113	This work
Hsp20	Hsp20/alpha crystallin family protein	ATGGGTCGGTCCTAAAGACTG	CGATAATGTGCCATCCCGATAC	119	This work
rfbB	dTDP-glucose 4,6-dehydratase	AACGCTTCCACCATGTTTCC	GTGTAAACTTCTCGCCAACACC	99	This work
dnaK	molecular chaperone DnaK	TTAGTACGGCGGCTGATAAC	GAAGCGACCTAACGTCTTG	95	This work
glmS	glutamine-fructose-6-phosphate transaminase	AATGTGATGCGGAAGCTATCCC	CCAGCAGCCACAATGTATAGTC	116	This work
trxA	thioredoxin	AACGGATACGGCTTCAGG	CACCCATCTCGTCAGCTAATTG	107	This work
cspA	cold shock protein	CTGGTGAAGATGGCAACGATG	CCACGATCGCTTGATTCTTC	115	This work
rpoD	RNA polymerase sigma factor RpoD	GCTTACGGTTCGGACTTGATG	GCCTTGGCTTCGATTTGG	103	Duary et al. (2010)
dacA	D-alanyl-D-alanine carboxypeptidase	AAGCAGCCAGTACACCATC	CGCCATTGGCTTACTACTGTTC	101	This work

551 **Table 3. Influence of the acclimation medium on the viability and L-malic acid consumption**

552 **of *Lactiplantibacillus plantarum* plantarum ATCC 14917 and UNQLp11 strains after 48 hours**

553 **incubation at two different temperatures (18 and 21 ºC).**

	ATCC 14917		UNQLp11	
	18 °	21 °	18 °	21 °
*A0 h Log CFU mL^{-1}	8.77 ± 0.72	8.45 ± 0.21	8.71 ± 0.27	8.45 ± 0.21
*A48 h Log CFU mL^{-1}	10.82 ± 1.83	9.53 ± 1.03	10.79 ± 1.79	10.04 ± 0.30
% MLF completed at 48 h	94.47 ± 7.82	100 ± 0.0	94.4 ± 7.92	100 ± 0.0

554 *A0 represent time zero (inoculation of *Lpb. plantarum* into the acclimation medium), and A48

555 represents time after 48 h of inoculation.