Delivery and expression of mRNA therapeutics during ex vivo kidney perfusion: A

feasibility study in porcine and human donor kidneys

Cristina Ballester[1,2†]**,** Ping Song[3†], Katerina B Knudsen[1,2], Chris Jaynes[4,5], Henri Leuvenink[4,5],

Victor R Llorente[1,2], Annick Van Furth[4], Rikke Nørregaard[1], Jan-Luuk Hillebrands[6], Anna

Krarup Keller[1,7], Bente Jespersen[1,2], Jørgen Kjems[3#], Marco Eijken[1,2#]

[1]Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

[2]Department of Renal Medicine, Aarhus University Hospital, Aarhus, Denmark

[3]Interdisciplinary Nanoscience Center (iNANO), Aarhus University, Aarhus, Denmark

[4]Department of Surgery, University Medical Center Groningen, Groningen, The Netherlands

[5]34Lives, Public Benefit Company, West-Lafayette, Indiana, United States

[6]Department of Pathology, University Medical Center Groningen, Groningen, The

Netherlands

[7]Department of Urology, Aarhus University Hospital, Aarhus, Denmark

[†] Shared first authors

[#] Shared last authors

Corresponding author: Cristina Ballester Bergada. Department of Clinical Medicine-Renal

Medicine, Aarhus University, Aarhus, Denmark. Addressee: Palle Juul-Jensens Blvd. 82,

8200 Aarhus, Denmark address. Mail: crisballester@clin.au.dk.

Running title: mRNA delivery during donor kidney perfusion

ABSTRACT

Ischemia–reperfusion injury (IRI) is a major cause of graft dysfunction in donor kidneys. Normothermic machine perfusion (NMP) offers a clinically relevant ex vivo platform for targeted intervention during organ preservation, while minimizing systemic exposure. Here, we investigated lipid nanoparticle (LNP)-mediated mRNA delivery during NMP as a strategy to enable immediate post-transplant therapeutic activity. Porcine and human kidneys were perfused at 37 °C for 6–12 h with an erythrocyte-based perfusate, and LNP-encapsulated mCherry or human erythropoietin (hEPO) mRNA was administered via the renal artery. Intravascular LNP delivery resulted in widespread renal parenchymal uptake of reporter mRNA without evidence of vascular obstruction or histological injury. hEPO protein became detectable in the perfusate approximately 2 h after infusion and continued to increase throughout perfusion, reaching biologically relevant concentrations. LNPs incorporating PEG-lipid TPGS (D-α-tocopherol polyethylene glycol 1000 succinate) achieved significantly higher mRNA delivery efficiency than conventional DSPE-PEG2000 formulations. The findings provide a foundation for the use of LNPs and mRNA therapeutics during organ machine perfusion as a platform for precision medicine in the field of kidney transplantation.

INTRODUCTION

Recent advancements in mRNA-based therapeutics have significantly expanded their potential applications in healthcare. In the field of infectious diseases, the application of SARS-CoV-2 mRNA-based vaccines has been effective[1, 2]. Additionally, mRNA cancer vaccines have shown promising results for melanoma treatment[3]. Beyond vaccines for infectious diseases and cancer, synthetic mRNAs are currently being developed to encode for therapeutic protein replacement therapies, immunotherapies, genome editing and tissue repair[4]. Numerous clinical trials exploring mRNA therapeutics are ongoing across a wide range of diseases, including cystic fibrosis transmembrane conductance regulator (CFTR) mRNA for cystic fibrosis and vascular endothelial growth factor A (VEGF-A) mRNA for myocardial infarction[5, 6], reflecting the growing versatility and therapeutic potential of this platform. Delivery of therapeutic mRNAs requires safe, effective and stable delivery systems to avoid nucleic acid degradation and overcome biological barriers. Lipid nanoparticles (LNPs) have been used as an efficient delivery platform to enhance mRNA stability and cellular uptake[7, 8]. Developing mRNA therapies for non-vaccine applications presents unique challenges. Unlike vaccines, therapeutic mRNAs typically require higher dosing, alternative routes of administration beyond intramuscular injections, and formulation strategies for enhanced targeted delivery and functional expression. In the bloodstream, certain LNPs can bind to apolipoprotein E (APOE), which, in turn, interacts with the abundant low-density lipoprotein receptors (LDLRs) in the liver. This interaction promotes the preferential uptake of LNPs by the liver, thereby limiting the distribution of mRNA to other tissues[9]. *Ex vivo* organ machine perfusion, a technique that

circulates a nutrient and oxygen-rich preservation solution or immune cell-depleted blood

substitute through the organ's vascular network to maintain its function and viability, is

increasingly adopted in clinical practice(10-12). The perfusion can be performed under

hypothermic conditions (typically 4-10°C), which minimizes metabolic activity to reduce

cellular damage, or under normothermic conditions (35–37°C), closely mimicking

physiological conditions and metabolic rates. This dynamic organ preservation technique

offers a unique opportunity for *ex vivo* delivery of mRNA therapeutics. It enables precise

tissue targeting with minimal off-target effects while achieving coverage of a whole organ,

which remains a critical barrier in the mRNA field. Organ machine perfusion is promising

for direct mRNA delivery to organs, potentially eliminating the need for complex LNP

modifications with targeting ligands. This approach allows for localized delivery, enabling

lower doses with improved safety, efficiency and cost-effectiveness. In particular,

normothermic machine perfusion (NMP) supports active metabolism, which can facilitate

LNP uptake and initiate mRNA translation prior to transplantation of the organ. We

hypothesize that mRNAs encoding for therapeutic proteins, delivered during NMP, present

a promising strategy to protect donor kidneys against ischemia-reperfusion injury (IRI). By

reducing the pathological reactivity of the graft triggered upon transplantation into the

recipient, this approach may help reduce fibrosis development, which can eventually

compromise graft function and survival(13). Therefore, the aim of this study is to evaluate

the potential of LNP-based delivery of therapeutic mRNAs to donor kidneys during *ex vivo*

NMP using model mRNAs as proof-of-concept.

RESULTS

Preparation and characterization of LNPs

In the present study, we developed three LNP formulations based on our in-house

synthesized ionizable lipid, C14120, using different coating materials: PEG-lipid TPGS

(TPGS-LNP), PEG-lipid DSPE-PEG2000 (DSPE-LNP), and a PEG-based block copolymer

(F127-LNP). The LNPs ranged between 150 nm to approximately 175 nm in diameter and

exhibited a positive surface charge with zeta potential over 30 mV and generally low

polydispersity (<0.2) (Figure 1a-c). LNPs were loaded with mRNA through electrostatic

interactions between ionizable lipids within the formulations and negatively charged

mRNAs. mRNA encoding mCherry protein was encapsulated within the three LNP

formulations at N/P ratios of 6, 12, and 16. The optimal N/P ratio for mRNA delivery and

protein expression was evaluated across the three LNP formulations in OK cells. For F127-

LNP and DSPE-LNP, transfection efficiency increased with higher N/P ratios, achieving

maximal mCherry expression at N/P ratio of 12 and 16 respectively (Figure1d). In contrast,

TPGS-LNP exhibited consistently high mCherry expression across all tested N/P ratios.

Importantly, no apparent cytotoxicity was observed after 24 hours of incubation with any of

the three LNP formulations at these N/P ratios (Figure 1e). Based on the balance of efficacy

and safety, an N/P ratio of 12 was selected for all subsequent experiments. At this ratio, all

three LNP types showed high mRNA encapsulation efficiency ranging from 80 to 90%

(Figure 1f).

To assess the stability of the encapsulated mRNA, LNPs-encapsulated mCherry mRNA

and free (naked) mCherry mRNA were incubated in the kidney perfusion solution (RBC-

free) at 37°C for up to 24 hours, with samples collected at multiple time points. The

remaining mCherry mRNA levels were quantified by RT-qPCR. Free mRNA exhibited rapid

degradation, with a sharp decline in detectable levels within the first hour of incubation. In

comparison, mRNA encapsulated within all three LNP formulations remained stable over

24 hours (Figure 1g).

The transfection efficiency of F127-, TPGS- and DSPE-LNP for mRNA delivery was assessed

in OK cells, using Cy5-labeled mRNA encoding EGFP (Cy5-EGFP). Both mRNA uptake (Cy5

signal) and protein expression (EGFP signal) were quantified using flow cytometry. No

significant difference in uptake was observed among the three formulations at 4 hours,

while DSPE-LNP exhibited the highest cellular uptake after 24 hours (Figure 2a,b).

Regarding mRNA-mediated protein expression, F127-LNP exhibited rapid EGFP expression

at 4 hours, whereas TPGS-LNP demonstrated the highest EGFP levels at 24 hours (Figure 2

a,c).

To further analyze protein expression efficiency, mRNAs encoding for mCherry and the

secreted protein, human erythropoietin (hEPO) were transfected in HKC-8 and OK cells

(supplemental figure 1). Similar trends were observed using both flow cytometry and ELISA

with these alternative reporter proteins, with TPGS-LNP consistently emerging as the most

efficient candidate showing a comparable pattern.

hEPO mRNA delivery during kidney-NMP

To evaluate mRNA delivery and translation during NMP, human erythropoietin (hEPO)

encoding mRNA was chosen as a reporter, enabling direct measurement of secreted hEPO

in both the perfusate and urine. Porcine kidneys (n=3-4 per treatment) were connected to

an NMP setup and 150 µg of DSPE or TPGS encapsulated hEPO mRNA (DSPE-hEPO and

TPGS-hEPO, respectively) was infused via the renal artery. The contralateral kidneys (n=6)

served as non-mRNA treated controls. After LNP-mRNA infusion, kidneys remained on

NMP for an additional six hours for follow-up. hEPO was measured in the perfusate as early

as two hours following mRNA administration, with levels gradually increasing to about 4

mIU/mL for DSPE-hEPO after 6 hours. TPGS-hEPO treated kidneys showed significantly

higher hEPO levels with up to 360 mIU/mL after 6 hours (Figure 3a). hEPO was also

detected in the urine reaching up to 6 mIU/mL and 420 mIU/mL for DSPE-hEPO and TPGS-

hEPO, respectively, 4-6 hours after administration (Figure 3b).

Perfusion characteristics, including renal blood flow, oxygen consumption and urine

production, were similar between porcine kidneys infused with DSPE-hEPO, TPGS-hEPO

and non-treated controls (Supplemental Figure 4). Hematoxylin and eosin (H&E) as well as

periodic acid-schiff (PAS) staining of the tissue sections showed no signs of LNP-induced

structural damage (Supplemental Figure 5).

mCherry mRNA delivery during kidney-NMP

To asses the possibility for localized expression of an intracellular protein, mRNA encoding

mCherry was administered to porcine kidneys using both DSPE- and TPGS-LNP, following

the same protocol as in the hEPO studies.

In tissue biopsies taken hourly up to 9 hours of NMP and analyzed by western blotting,

mCherry protein expression became apparent after 4-5 hours for TPGS-LNP and 7-8 hours

for DSPE-LNP, both increasing towards the end of the perfusion (9 hours; Figure 4a, b).

Similar to hEPO expression, TPGS-LNP exhibited greater efficacy than DSPE-LNP (Figure

4a, b). Analysis of cortical tissue collected from different kidney poles and the medulla

confirmed that the delivered mRNA reached all regions within the organ. However,

substantial variation in mCherry levels was observed between the tested kidneys.

Notably, in the 9-hour follow-up period, high intrinsic autofluorescence in most kidneys

interfered with the reliable detection of mCherry fluorescence, limiting the accuracy of

direct fluorescence measurements. To overcome this, immunohistochemistry was

employed to visualize mCherry localization at the cellular level. This showed broad

expression across the tissue, including expression within both glomerular and tubular

structures (Figure 5a). BRISH analysis detecting mCherry mRNA confirmed the widespread

tissue distribution of mCherry mRNA, highlighting the effective tissue penetration of LNPs

after the *ex vivo* intra-arterial administration (Figure 5b). The mRNA signal was

predominantly detected intracellularly within the endothelium, although it was also

observed in other renal cell types (Figure 5b).

Translational studies in human kidneys

Next, the feasibility of mRNA therapy in human deceased donor kidneys was investigated

(see Supplemental Table 2 for donor characteristics). DSPE- and TPGS-LNPs carrying

mCherry mRNA were administered to 7 human kidneys during NMP using the same

protocol as the experiments conducted on porcine kidneys.

Within the 12-hour follow-up, mCherry expression was not as abundant compared to the

porcine kidneys. Using immunohistochemistry, however, mCherry protein could be

detected throughout the tissue of 5 of the treated kidneys, whereas the 2 others failed to

induce mCherry expression (Figure 6a). A greater mCherry signal was observed following

mRNA delivery via TPGS-LNPs (Figure 6a).

BRISH analysis confirmed that both DSPE- and TPGS-LNPs achieved intracellular mRNA

delivery throughout the human kidneys (Figure 6b), consistent with the findings in porcine

kidneys.

Similarly to the porcine kidneys, perfusion characteristics including renal blood flow,

oxygen consumption and urine production were not statistically affected by the LNP-

mRNA infusion in the human deceased donor kidneys (supplemental Figure 6).

DISCUSSION

This study establishes a clinically compatible strategy for organ-directed mRNA delivery

during *ex vivo* perfusion. By delivering the mRNA to the isolated organ, off-target effects

commonly associated with systemic mRNA delivery are minimized, including hepatic

interactions (9). Intra-vascular LNP administration via the renal artery achieved effective

tissue penetration and biodistribution of the reporter mRNAs. hEPO expression became

detectable around 2 hours after mRNA administration, reaching biologically relevant levels

during NMP. The rapid appearance of the mRNA ensures the presence of the encoded

proteins during the critical post-transplant period, potentially driving new therapeutic

strategies to mitigate IRI, promote tissue regeneration, and ultimately improve organ

transplantation outcomes.

Building upon previously validated ionizable lipid-based LNP formulations(14), we utilized

three different PEG-lipids: Pluronic F127, TPGS, and DSPE-PEG2000, each contributing

unique properties (15-18). While all formulations showed comparable physical

characteristics, DSPE-PEG2000 and F127 achieved slightly higher mRNA encapsulation

efficiencies than TPGS. This may relate to TPGS's smaller lipid head group and shorter PEG

chain, potentially compromising particle stability.

Our results indicate that when delivering mRNA to whole porcine and human kidneys *ex*

vivo, LNPs containing TPGS as the PEG-lipid achieved markedly higher protein expression

compared to the conventional DSPE-PEG2000. This correlates with the *in vitro* findings,

where TPGS-LNP outperformed the other formulations in terms of expression of the

reporter protein. The enhanced potency of TPGS-containing LNPs under *ex vivo* conditions

may be partly explained by the ability of TPGS to inhibit P-glycoprotein activity(16, 19, 20)

which is highly expressed in proximal tubule cells and plays a central role in efflux

transport. Beyond serving as a passive stabilizer, TPGS can improve LNP stability within the

extracellular milieu, facilitate cellular uptake, and reduce nanoparticle efflux. Moreover, its

surfactant-like properties may promote endosomal escape, thereby increasing the

probability that delivered mRNA reaches the ribosomal machinery for translation(16).

Collectively, these complementary mechanisms could account for the improved

biocompatibility, tissue accumulation, and delivery efficiency observed.

In our previous *in vivo* work in rodents, mRNA was administered systemically at doses

ranging from 0.01 to 1 mg mRNA/kg (22-24). Based on these studies a dose of 0.15 mg

mRNA encapsulated in 1 mg of LNPs was selected per porcine or human kidney(21-23). In

comparison, a clinical trial investigating VEGF mRNA delivery via epicardial injection used

a total dose of 3 mg mRNA(24). Another study investigating an mRNA-based replacement

therapy for cystic fibrosis utilized CO-hCFTR mRNA via nebulizer, with nominal doses

ranging between 4-24 mg mRNA(5). Notably, the relatively low dose of 0.15 mg mRNA per

porcine kidney (weighing approximately 100–150 g) used in our study resulted in high levels

of human EPO reaching up to 400 mIU/mL in the perfusate. For comparison, normal

circulating EPO concentrations in healthy individuals range from 2.8 to 17.1 mIU/mL(25).

In the present study, kidney-NMP was used as a delivery platform and perfusion time was

extended to monitor therapeutic protein expression for up to 12 hours. Using mRNA

therapeutics, *in vitro* systems have demonstrated sustained protein expression for up to a

few days(26, 27), while *in vivo*, systemic peak expression typically occurs at 6 hours and

then decreases(28, 29). A transient expression of therapeutic proteins without long-term

effects is particularly advantageous for targeting the pathogenic pathways following IRI,

observed directly in the period after organ transplantation(30-32).

Ex vivo mRNA delivery was successfully achieved using human-sized porcine kidneys or

human donor kidneys using a clinically compatible NMP setup, identical to those

employed in clinical studies(33, 34). Furthermore, LNPs were administered through a

simple infusion via the arterial port of the NMP setup, confirming the clinical feasibility of

the approach. The lower mCherry protein expression observed in the human kidneys may

reflect their larger organ size and variable donor-related factors, including age and

comorbidity, duration of warm ischemia during procurement surgery and cold storage,

potentially necessitating higher doses.

LNP-mRNA treatment was well tolerated during NMP, with preserved perfusion parameters

and renal architecture. Additionally, no evidence of cytotoxicity was detected *in vitro* or

observed *ex vivo*. Adverse effects seen in systemically delivered LNP-mRNA studies are not

anticipated in our ex vivo approach, as systemic exposure is minimized. Moreover,

following NMP, it is standard clinical procedure to flush the kidney and/or return it to HMP

prior to transplantation, which facilitates the removal of any free LNPs.

Given the complex and multifactorial nature of IRI, it may be necessary to target multiple

pathways(35). Donor kidneys would greatly benefit from a combined and individualized

treatment strategy and mRNA therapy offers a flexible platform for such an approach, as it

can encode both gain-of-function activators and inhibitors, which could potentially be

combined into a single solution. The rapid protein expression following *ex vivo* delivery of

mRNA therapeutics enables the use of these therapeutics to transiently modify the donor

organ during preservation, enhancing graft resilience during the critical period immediately

following transplantation(31, 32, 35).

In summary, we have successfully developed and screened LNPs for efficient mRNA

delivery to the kidney during NMP. TPGS-based LNPs emerged as a promising vector for

mRNA therapeutics to an isolated kidney, demonstrating their efficacy both *in vitro* and *ex*

vivo in porcine and human kidneys. Furthermore, we established a practical strategy for

using mRNA therapeutics in donor organs, providing a foundation for their application in *ex*

vivo normothermic machine perfusion as a precision medicine platform prior to organ

transplantation.

MATERIALS AND METHODS

Cell culture

Opossum kidney cells (OK cells) were cultured in low glucose Dulbecco's Modified Eagle

Medium (DMEM) (Thermo Fisher, Waltham, MA), supplemented with 10% fetal bovine

serum (FBS, Thermo Fisher, Waltham, MA) and 1% penicillin-streptomycin (Thermo Fisher,

Waltham, MA) at 37°C, 5% CO_2 and 100% humidity.

Preparation of nanoparticles

Lipid nanoparticles (LNP) were synthesized through nanoprecipitation. Briefly, Pluronic

(F127, Sigma-Aldrich), C14120(36), cholesterol (Sigma-Aldrich), 1,2-dioleoyl-sn-glycero-3-

phosphocholine (DOPC, Avanti Polar Lipids) and d-α-tocopheryl polyethyleneglycol 1000

succinate (TPGS, Sigma-Aldrich) were dissolved in 100% ethanol. For F127-LNP, F127 and

C14120 were combined at the weight ratio of 2:1. For DSPE-LNP and TPGS-LNP, C14120,

cholesterol, DOPC, and the PEG-Lipid either TPGS or 2-distearoyl-sn-glycero-3-

phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000] (DSPE-PEG2000, Avanti

Polar Lipids) were combined at weight ratio of 4:2:1:1. Subsequently, sodium acetate

(200mM, pH 5.5) was rapidly injected into the organic lipid mixture with vigorous stirring,

leading to be self-assembly of empty nanoparticles. Ethanol was removed by dialysis

against 200 mM sodium acetate pH 5.5 using a 3.5 kDa dialysis tube (Thermo Scientific).

LNPs were stored at 4°C and used within one month.

mRNA encapsulation and delivery

The following mRNAs were used: CleanCap EPO mRNA (5moU, Trilink, L-7209), CleanCap

mCherry mRNA (5moU, Trilink, L-7203) and CleanCap Cy5 eGFP mRNA (5 moU, Trilink, L-

7701). mRNA was encapsulated through electrostatic interaction when thoroughly mixed

with the LNPs.

For in vitro studies, LNPs (1 mg/mL, concentration of C14120) in 200 mM sodium acetate

buffer pH 5.5) were mixed with mRNA (at 200 ng/µL) at different N/P ratios. 100 ng of mRNA

was mixed with 0.75 µL, 1.5 µL, and 2 µL of LNPs to obtain N/P ratios of 6, 12 and 16,

respectively. The LNP/mRNA mixture was incubated for 10 minutes at room temperature

(RT) and added to the culture medium.

For delivery during kidney NMP, 1100 µg LNPs (in 2.5 mL 200 mM acetate buffer, pH 5.5)

were mixed with 150 µg of mRNA (prediluted in 0.5 mL of nuclease-free water) and

incubated for 10 minutes at RT. Prior to the infusion, 2 mL of isotonic saline (0.9 mg/mL

NaCl Fresenius Kabi, B05BB01) was added. This solution was transferred to a 20 mL

syringe and administered via the arterial side of the NMP circuit at an infusion rate of 120

mL/h, starting 30 minutes after the initiation of kidney-NMP. Following the initial infusion,

the line was flushed with 3 mL of isotonic saline.

Characterization of LNPs

The hydrodynamic diameter and surface charge of the LNPs were determined by photon

correlation spectroscopy (PCS) and Laser Doppler Velocimetry (LDV) at 25 °C on a

Zetasizer Nano ZS (Malvern Instruments, Malvern, UK). The encapsulation efficiency of

mRNA was quantified using the RiboGreen assay (Invitrogen, Copenhagen) following the

manufacturer's protocol. Briefly, 50 µL of LNP/mRNA solution was mixed with 50 µL of

RiboGreen working solution (1:100 in TE buffer: 10 mM Tris, 0.1 mM EDTA, pH 7.5). After 5

minutes of incubation at RT, the fluorescence intensity was measured by FLUOstar

OPTIMA (Moritex BioSience) at an excitation/emission wavelength of 480/520 nm.

Cell viability test

Cell viability was evaluated via AlarmaBlue (Molecular Probes, Life Technologies) assay

according to the instructions from the manufacturer. OK cells were seeded in a 96-well

plate at a density of $1x10^4$ cells/well. The following day, the growth medium was refreshed

and each well was transfected with 33 ng LNP-encapsulated mCherry mRNA. After 24

hours, growth medium supplemented with 10% (v/v) of AlarmaBlue reagent was added.

After 2 hours of incubation, 80 µL of the supernatant was transferred to an opaque 96-well

NUNC plate (Sigma-Aldrich, P8616), and the fluorescence intensity was measured using a

FLUOstar OPTIMA plate reader (Moritex BioScience) at excitation and emission

wavelengths of 540 nm and 590 nm, respectively.

Stability of mRNA encapsulated in LNPs

To assess stability, 88 ng of LNP-encapsulated mCherry mRNA was incubated in 45 μL of

NMP perfusate (see supplemental table 1) at 37°C for 0 minutes (right after mixing), 15

minutes, 30 minutes, 1 hour, 2 hours, 4 hours, 6 hours and 24 hours. After incubation, lysis

buffer (0.25% Igepal CA-630, 10mM Tris·HCl pH 7.4, 150 mM NaCl, and 1 U/µL RNase

inhibitor) was added and samples were centrifuged at 10,000 g for 5 minutes. Next, 10 µL

of supernatant was used directly for cDNA preparation using Revert Aid RT Reverse

Transcription Kit (Thermo Fisher). The cDNA was diluted 40 times, and 4 µL was used as

template for quantitative real-time PCR using SYBR Green Master Mix (Thermo Fisher),

running on a Light Cycler 480 Real-Time PCR System (Roche). The primer sequences for

mCherry are: Forward 5'-CGGGTGATGAACTTCGAGGA-3' and Reverse 5'-

TCGAAGTTCATCACCCGCTC-3'.

In vitro performance of LNPs for mRNA delivery

OK cells were seeded in 24-well plate (1x10^5 cells/well). The next day, growth medium was

refreshed and 200 ng LNP-encapsulated mRNA was added per well. After 24 hours, cells

were detached using 0.05% Trypsin-EDTA (Thermo Fisher, Waltham, MA) and fluorescent

intensity was quantified by flow cytometry (NovoCyte, ACEA Biosciences).

For microscopy studies, OK cells were seeded in 8-well µ-Slide (ibiTreat, ibidi; 5x10^4

cells/well). The next day, the growth medium was refreshed and 100 ng LNP-encapsulated

mRNA was added per well. After 24 hours, medium was replaced by FluoroBrite DMEM

(Thermo Fisher Scientific, Waltham, MA), and cells were imaged by confocal laser

scanning microscopy (LSM 710, Zeiss, Germany) and analyzed by Fiji ImageJ software

2.3.0.

Organ procurement and normothermic machine perfusion

A comprehensive account of the ethical approval process, surgical techniques and blood

processing procedures is available in Supplemental Material 1.

Porcine kidneys were procured under general anesthesia following a standardized surgical

protocol, including midline laparotomy and bilateral nephrectomy with a warm ischemia

time as in a living donation procedure. Special care was taken to preserve vascular

integrity and obtain sufficient arterial length for optimal cannulation. The renal artery was

cannulated with a straight t-cannula immediately post-nephrectomy and with minimal

warm ischemic time. Then, the kidney was flushed with heparinized saline followed by cold

preservation solution before being stored at 4 °C. Human kidneys, procured initially with

the intent for transplantation but later declined, were transported on hypothermic machine

perfusion and re-cannulated upon arrival using appropriately sized arterial and ureteral

cannulas. Further procedural details are provided in the Supplemental Material 1.

After a maximum of 32 hours of cold ischemia, all kidneys were connected to an open

clinically compatible NMP system via the arterial cannula and venous outflow drained

freely into the reservoir under gravity, permitted via an uncannulated renal vein.

The NMP circuit was primed with 600 mL human albumin erythrocyte-based solution (see

composition supplemental material 2). The perfusate temperature was kept at 37°C and

oxygenated with 0.5 L/minute carbogen gas (95% O_2, 5% CO2). After the kidney was

connected to the system, a mean arterial pressure of 80 mmHg was maintained. Gas

analysis was carried out every hour (ABL90 FLEX, Radiometer) and hourly perfusate, urine

and biopsies were taken. Urine output was measured, and the rest of the urine was

recirculated. During NMP, glucose and pH were adjusted if glucose dropped under 5 mmol

and pH under 7.35 by adding a bolus of glucose 5% (50 mg/mL, Fresenius Kabi,

7340022100429) or sodium bicarbonate 8.4% (8.4 g/mL, Fresenius Kabi, 03058282)

directly to the organ chamber.

Sampling and storage

During follow-up on NMP, perfusate from the vein and urine samples were collected.

Samples were centrifuged for 12 minutes at 3000 g at 4°C and supernatant was stored at -

80°C. Blood gases from arterial and venous perfusate as well as urine, were measured

hourly (ABL90 FLEX, Radiometer) to monitor the perfusion process. Each 6 mm biopsy was

stored in RNA-later (Merck KGaA) at -80°C. At the end of the perfusion 1x1 cm biopsies with

both cortex and medulla were taken and formalin fixed (Biopsafe, 3178-200-03) for about

24 hours at 4°C, and were then changed to phosphate buffered saline until embedding.

Human EPO ELISA

Human EPO protein expression in kidney NMP perfusate and urine samples was quantified

by a Human EPO ELISA (Invitrogen, BMS2035-2). Human EPO in OK cell cultures was

quantified by an EPO DuoSet ELISA (R&D Systems, DY286-05) according to the

manufacturer's protocols.

mCherry Western Blot

In total, 25 mg of kidney tissue was homogenized in M tubes (Miltenyi Biotec,130093236) in 500 μL of Pierce RIPA buffer (Thermo Scientific, 89901) supplemented with cOmplete Mini protease inhibitor cocktail (Merck KGaA, 11836153001). Lysates were centrifuged (10 minutes, 3000 g, 4°C) and supernatant was stored at -80°C until further use. Total protein was quantified using Pierce BCA Protein Assay Kit (Thermo Fisher, 23225) according to the manufacturer's protocols. In total, 30 μg of protein was run on an Any kD CriterionTM TGX Stain-Free gel (BIO-RAD, 5678124) and transferred onto nitrocellulose Trans-Blot®TurboTM membranes (BIO-RAD, 1704159). Blots were blocked using 5% Skim milk powder (Millipore, 999999-99-4) and washed in 20 mM TBS-T (Tris-buffered saline, pH 7.6, 0.1% TWEEN 20). Membranes were incubated with 0.35 μg/ml DSHB-mCherry-3A11-s antibody (Development Studies Hybridoma Bank, DSHB) in 20 mL of TBS-T overnight at 4°C, followed by 1 hour of incubation at RT. After being washed in TBS-T, blots were incubated with 0.4 μg/ml polyclonal Goat Anti-mouse Immunoglobulins/HRP antibody (Aligent, P0447) in TBS-T for 1 hour at RT. After wash, blots were incubated with 3 mL ECLTM Prime Western Blotting Detection Reagent (Cytvia, RPN2236) for 5 minutes at RT, and imaged using the ChemiDocTM MP Imaging System (BIO-RAD). Quantification of bands in the blot was done with ImageLab (version 6.0.1, 2017) and signal was corrected for total protein of the respective gel.

Immunofluorescence (IF) and microscopy

The formalin-fixed paraffin-embedded (FFPE) samples were sectioned at 2.5 µm with a

microtome (Thermo Scientific, HM3555). Hematoxylin and eosin (H&E) and Periodic Acid–

Schiff (PAS) staining were performed on FFPE kidney sections (3–4 µm thick) using

standard protocols. H&E was used to evaluate general tissue morphology, while PAS

staining was used to assess glomerular basement membranes, tubular brush borders, and

mesangial matrix integrity. Histological scoring was performed by a trained pathologist

blinded to the treatment groups. Kidney sections were evaluated for features including

tubular damage, dilatation, cast formation, vacuolization, glomerular injury, interstitial

inflammation, and edema. Each parameter was semi-quantitatively scored based on

predefined criteria, and overall tissue integrity was assessed to compare damage and

structural preservation between groups.

Before IF staining, kidney FFPE samples were heated at 60°C and deparaffinized with

xylene (BDH Chemicals, 28973-363). Antigen retrieval was performed by immersing the

slides in a preheated antigen retrieval buffer (1.21 g Tris (Sigma T-1378), 0.19 g EGTA

(Triplex VI, Merck 8435) in 1L of MilliQ water, pH 9.0) and heating them at 95-100°C for 10-

20 minutes. After, the sections were allowed to cool down for about 20 minutes at RT.

Then, the sections were washed and permeabilized in TBS-T for 10 minutes at RT. Sections

were placed at RT for 10 minutes then fixed with 4% paraformaldehyde (PFA) for 15

minutes. Following a 5-minute wash with TBS (Tris-NaCl buffered saline, 20 mM Tris, 150

mM NaCl, pH 7.2) the sections were blocked for 1 hour at RT using horse serum (5% for

human kidneys and 1% for porcine kidneys, Thermo Fisher, 26050088) in TBS-T (0.2%

Triton X100 in TBS). Next, the sections were incubated for 2 hours with 2 µg/mL DSHB-

mCherry-3A11-s antibody in TBS-T supplemented with 2.5% horse serum for human

kidneys and 0.5% horse serum for porcine kidneys. After three 10-minute washes with TBS,

the sections were incubated with 5 µg/mL Polyclonal Goat Anti-Mouse

Immunoglobulins/CF647 antibody (Sigma Aldrich, SAB4600182) in TBS-T, for 1 hour at RT.

This was followed by another three 5-minute washes with TBS. Finally, the sections were

mounted with ProLong Diamond Antifade Mountant with 4',6-diamidino-2-phenylindole

(DAPI) mounting media (Thermo Fisher Scientific, P36962) and stored at 4°C until

observation with confocal microscopy (LSM 800 Airyscan, Zeiss) and VS120 Slides Scanner

(Olympus). The images obtained were processed using either Fiji (version 1) or Qu Path

(version 0.5.1-arm64) software.

Brightfield RNA In Situ Hybridization (BRISH)

RNA-ISH was performed by using the RNAscope Probe-mCherry-O10-C1 (Cat. No.

1264111-C1, target region 2-714 bp), which is a probe specific mCherry mRNA. For RNA

quality controls, probes targeting the housekeeping gene peptidylprolyl isomerase B

(cyclophilin B) were used as follows: Ss-PPIB (Cat. No. 428591, target region 16-642 bp) for

pig tissue, and Hs-PPIB (Cat. No. probe. 313901, target region 139-989 bp) for human

tissue and cells.

As specificity control for mCherry detection, about 100.000 HEK293 cells were transfected

with 200 ng mCherry mRNA carried by 1.5 µg of TPGS-LNP for 24 hours. HEK293 cells

incubated for 24 hours without transfection were used as controls. Cells were fixed with

4% paraformaldehyde (PFA) for 1 hour and placed in agar solution (0.18 g of agar in 10 mL

of demineralized water). Followed by a second fixation with 4% PFA and paraffin

embedding.

RNA-BRISH was performed on 5-µm-thick FFPE sections on SuperFrost PLUS microscope

slides. Pretreatment was done following the RNAscope Sample Preparation and

Pretreatment Guide for FFPE Tissue (Cat. No. 322381). This included blocking H_2O_2 by

adding the RNAscope hydrogen peroxide to the slides and incubating them for 10 minutes

at RT. Antigen retrieval was performed by heating the slides for 18 minutes in the

RNAscope antigen retrieval buffer (1:10 dilution and 98-100 °C) using a steamer. Then,

protease treatment was performed by adding the RNAscope Protease Plus to the sections

and incubating them at 40°C for 30 minutes in the HybEZ oven, followed by probe

hybridization overnight. Signals were detected by using the RNAscope 2.0 HD Detection Kit

(Cat. No. 322310). This included amplification by a series of amplification steps and signal

detection by using DAB (3,3'-Diaminobenzidine). The reactivity was visualized as brown

dots. Following BRISH, nuclear counterstaining was performed using hematoxylin.

Sections were coverslipped and digitized using an Olympus SLIDEVIEW VS200 slide

scanner (Olympus Nederland B.V., Hoofddorp, Netherlands).

Data analysis

GraphPad Prism (version 10.2.2) was used to perform statistical analyses and compose

graphs. Data are presented as mean ± SD unless otherwise stated. Statistical tests

included two-way ANOVA, t-tests, and appropriate non-parametric tests based on data

distribution. Statistical significance was set to $p < 0.05$.

DATA AVAILABILITY STATEMENT

Raw numerical data underlying Figures 1–4 and Supplementary Figures S1–S4 (GraphPad

Prism files) are available upon request.

ACKNOWLEDGEMENTS

The authors would like to express their gratitude to the 34Lives team, especially, Kathleen

St Jean, Stephen Eilert, Haley Frost and Ali Smith, for hosting the translational experiments

providing the opportunity to perform studies with human deceased donor kidneys and

arranging the logistics for transport and infrastructure. This study would not have been

possible without the gracious gifts from the generous donor families and through

collaborations with Kentucky Organ Donor Affiliates (Network for Hope), Life Connection of

Ohio, Gift of Hope, Texas Organ Sharing Alliance, LifeGift, Indiana Donor Network, Iowa

Donor Network, and Lifeline of Ohio. We acknowledge Mirjiam Mastik for her valuable

assistance in carrying out the BRISH studies. Fluorescent Microscopy experiments were

supported by the Scientific Service Units (SSU) of Institute of Science and Technology

(ISTA) through resources provided from the Imaging and Optics Facility (IOF), and personal

support of Josephine Pfiefer and Bernhard Hochreiter. Scoring of kidney tissue sections

was supported by pathologist Søren Rasmus Palmelund Krag. And lastly, the authors

would like to acknowledge Lene Mejer Rosenlund, Maria Arge and Birgitte Kildevæd Sahl

for their expert biotechnician assitance. This work was supported by Augustinus Fonden

and Novo Nordisk Foundation grant NNF23OC0081177 (RNA-META).

AUTHOR CONTRIBUTIONS

ME, JK, BJ, AK, RN, HL, CBB and PS conceptualized the research design. CBB, PS, KBK,

VRL, CJ and AVF participated in the performance of the experiments. CBB, PS and JH

participated in the analysis. CBB and PS drafted the manuscript. ME performed critical

revision. ME, JK, BJ and AK supervised the project and reviewed the final version of this

manuscript. All authors approved the final version of the manuscript.

543 **DECLARATION OF INTERESTS**

544 There are no conflicts of interest.

565 **KEY WORDS**

566 mRNA therapeutics, , donor organs, kidney machine perfusion, graft optimization, LNPs

REFERENCES

1. Walsh EE, Frenck RW, Jr., Falsey AR, Kitchin N, Absalon J, Gurtman A, et al. Safety
and Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. N Engl J Med.
2020;383(25):2439-50.
2. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. Efficacy and Safety
of the mRNA-1273 SARS-CoV-2 Vaccine. N Engl J Med. 2021;384(5):403-16.
3. Sahin U, Oehm P, Derhovanessian E, Jabulowsky RA, Vormehr M, Gold M, et al. An
RNA vaccine drives immunity in checkpoint-inhibitor-treated melanoma. Nature.
2020;585(7823):107-12.
4. Parhiz H, Atochina-Vasserman EN, Weissman D. mRNA-based therapeutics:
looking beyond COVID-19 vaccines. Lancet. 2024;403(10432):1192-204.
5. Rowe SM, Zuckerman JB, Dorgan D, Lascano J, McCoy K, Jain M, et al. Inhaled
mRNA therapy for treatment of cystic fibrosis: Interim results of a randomized, double-
blind, placebo-controlled phase 1/2 clinical study. J Cyst Fibros. 2023;22(4):656-64.
6. Anttila V, Saraste A, Knuuti J, Jaakkola P, Hedman M, Svedlund S, et al. Synthetic
mRNA Encoding VEGF-A in Patients Undergoing Coronary Artery Bypass Grafting: Design of
a Phase 2a Clinical Trial. Mol Ther Methods Clin Dev. 2020;18:464-72.
7. Hou X, Zaks T, Langer R, Dong Y. Lipid nanoparticles for mRNA delivery. Nat Rev
Mater. 2021;6(12):1078-94.
8. Whitehead KA, Langer R, Anderson DG. Knocking down barriers: advances in siRNA
delivery. Nat Rev Drug Discov. 2009;8(2):129-38.
9. Sebastiani F, Yanez Arteta M, Lerche M, Porcar L, Lang C, Bragg RA, et al.
Apolipoprotein E Binding Drives Structural and Compositional Rearrangement of mRNA-
Containing Lipid Nanoparticles. ACS Nano. 2021;15(4):6709-22.
10. Moers C, Smits JM, Maathuis MH, Treckmann J, van Gelder F, Napieralski BP, et al.
Machine perfusion or cold storage in deceased-donor kidney transplantation. N Engl J
Med. 2009;360(1):7-19.
11. Ceresa CDL, Nasralla D, Pollok JM, Friend PJ. Machine perfusion of the liver:
applications in transplantation and beyond. Nat Rev Gastroenterol Hepatol.
2022;19(3):199-209.
12. Rega F, Lebreton G, Para M, Michel S, Schramm R, Begot E, et al. Hypothermic
oxygenated perfusion of the donor heart in heart transplantation: the short-term outcome
from a randomised, controlled, open-label, multicentre clinical trial. Lancet.
2024;404(10453):670-82.
13. Lasorsa F, Rutigliano M, Milella M, d'Amati A, Crocetto F, Pandolfo SD, et al.
Ischemia-Reperfusion Injury in Kidney Transplantation: Mechanisms and Potential
Therapeutic Targets. Int J Mol Sci. 2024;25(8).
14. Song P, Yang C, Thomsen JS, Dagnaes-Hansen F, Jakobsen M, Bruel A, et al.
Lipidoid-siRNA Nanoparticle-Mediated IL-1beta Gene Silencing for Systemic Arthritis
Therapy in a Mouse Model. Mol Ther. 2019;27(8):1424-35.
15. Hald Albertsen C, Kulkarni JA, Witzigmann D, Lind M, Petersson K, Simonsen JB. The
role of lipid components in lipid nanoparticles for vaccines and gene therapy. Adv Drug
Deliv Rev. 2022;188:114416.

16.   Yang C, Wu T, Qi Y, Zhang Z. Recent Advances in the Application of Vitamin E TPGS
for Drug Delivery. Theranostics. 2018;8(2):464-85.
17.   Yu H, Dyett BP, Zhai J, Strachan JB, Drummond CJ, Conn CE. Formation of
particulate lipid lyotropic liquid crystalline nanocarriers using a microfluidic platform. J
Colloid Interface Sci. 2023;634:279-89.
18.   Akash MS, Rehman K. Recent progress in biomedical applications of Pluronic
(PF127): Pharmaceutical perspectives. J Control Release. 2015;209:120-38.
19.   Banan Sadeghian R, Ueno R, Takata Y, Kawakami A, Ma C, Araoka T, et al. Cells
sorted off hiPSC-derived kidney organoids coupled with immortalized cells reliably model
the proximal tubule. Commun Biol. 2023;6(1):483.
20.   Collnot EM, Baldes C, Wempe MF, Kappl R, Huttermann J, Hyatt JA, et al.
Mechanism of inhibition of P-glycoprotein mediated efflux by vitamin E TPGS: influence on
ATPase activity and membrane fluidity. Mol Pharm. 2007;4(3):465-74.
21.   Evers MJW, Du W, Yang Q, Kooijmans SAA, Vink A, van Steenbergen M, et al.
Delivery of modified mRNA to damaged myocardium by systemic administration of lipid
nanoparticles. J Control Release. 2022;343:207-16.
22.   Jiang Y, Zhang Y, Liu C, Liu J, Xue W, Wang Z, et al. Tumor-activated IL-2 mRNA
delivered by lipid nanoparticles for cancer immunotherapy. J Control Release.
2024;368:663-75.
23.   Zhao S, Gao K, Han H, Stenzel M, Yin B, Song H, et al. Acid-degradable lipid
nanoparticles enhance the delivery of mRNA. Nat Nanotechnol. 2024;19(11):1702-11.
24.   Anttila V, Saraste A, Knuuti J, Hedman M, Jaakkola P, Laugwitz KL, et al. Direct
intramyocardial injection of VEGF mRNA in patients undergoing coronary artery bypass
grafting. Mol Ther. 2023;31(3):866-74.
25.   Grote Beverborg N, Verweij N, Klip IT, van der Wal HH, Voors AA, van Veldhuisen DJ,
et al. Erythropoietin in the general population: reference ranges and clinical, biochemical
and genetic correlates. PLoS One. 2015;10(4):e0125215.
26.   Liu J, Zhang Y, Liu C, Jiang Y, Wang Z, Guo Z, et al. A single dose of VEGF-A circular
RNA sustains in situ long-term expression of protein to accelerate diabetic wound healing.
J Control Release. 2024;373:319-35.
27.   Chen R, Wang SK, Belk JA, Amaya L, Li Z, Cardenas A, et al. Engineering circular RNA
for enhanced protein production. Nat Biotechnol. 2023;41(2):262-72.
28.   Pardi N, Tuyishime S, Muramatsu H, Kariko K, Mui BL, Tam YK, et al. Expression
kinetics of nucleoside-modified mRNA delivered in lipid nanoparticles to mice by various
routes. J Control Release. 2015;217:345-51.
29.   Chen H, Liu D, Aditham A, Guo J, Huang J, Kostas F, et al. Chemical and topological
design of multicapped mRNA and capped circular RNA to augment translation. Nat
Biotechnol. 2024.
30.   Sharfuddin AA, Molitoris BA. Pathophysiology of ischemic acute kidney injury. Nat
Rev Nephrol. 2011;7(4):189-200.
31.   Legrand M, Mik EG, Johannes T, Payen D, Ince C. Renal hypoxia and dysoxia after
reperfusion of the ischemic kidney. Mol Med. 2008;14(7-8):502-16.
32.   Ponticelli C. Ischaemia-reperfusion injury: a major protagonist in kidney
transplantation. Nephrol Dial Transplant. 2014;29(6):1134-40.

33.   Hosgood SA, Callaghan CJ, Wilson CH, Smith L, Mullings J, Mehew J, et al.
Normothermic machine perfusion versus static cold storage in donation after circulatory
death kidney transplantation: a randomized controlled trial. Nat Med. 2023;29(6):1511-9.
34.   An Observational Study of a Dedicated Preservation and Assessment Service to
Optimize Organ Utilization for Hard-to-Place Kidneys [Internet]. 2024. Available from:
https://clinicaltrials.gov/study/NCT06263023.
35.   Ogurlu B, Hamelink TL, Van Tricht IM, Leuvenink HGD, De Borst MH, Moers C, et al.
Utilizing pathophysiological concepts of ischemia-reperfusion injury to design
renoprotective strategies and therapeutic interventions for normothermic ex vivo kidney
perfusion. Am J Transplant. 2024;24(7):1110-26.
36.   Love KT, Mahon KP, Levins CG, Whitehead KA, Querbes W, Dorkin JR, et al. Lipid-
like materials for low-dose, in vivo gene silencing. Proc Natl Acad Sci U S A.
2010;107(5):1864-9.

 FIGURE LEGENDS

Figure 1. In vitro Characterization of F127-LNP, TPGS-LNP and DSPE-LNP. (a)

Hydrodynamic diameter. (b) Polydispersity index (PDI). (c) Zeta potential. (d) mCherry

protein expression and cell viability (e) in OK cells after transfection with LNP-

encapsulated mCherry mRNA at N/P ratios of 6,12 and 16. (f) The encapsulation efficiency

of F127-LNP, TPGS-LNP and DSPE-LNP at N/P ratio of 12. (g) Stability of free and LNP-

encapsulated mCherry mRNA (F127-LNP, TPGS-LNP and DSPE-LNP) in kidney perfusion

solution over 24 hours at 37 °C, indicated by ct values measured by RT-qPCR and

presented as Mean ± SD (n = 3).

Figure 2. mRNA uptake and protein expression in OK cells transfected with DSPE-LNP,

TPGS-LNP and F127-LNP. (a) Representative confocal images of OK cells transfected with

LNP encapsulated with Cy5-labeled eGFP mRNA. Green: eGFP protein expression; red:

Cy5-labeled eGFP mRNA (mRNA-Cy5); blue: nucleus stained with Hoechst 33342. (b)

Quantitative analysis of cellular uptake of Cy-5 labeled eGFP mRNA and (c) eGFP protein

expression. Data are presented as Mean $\pm$ SD (n = 3). Statistical significance was

determined by multiple mixed effects analysis.* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 3: Levels of human erythropoietin (hEPO) detected in perfusate (kidney perfusion

solution) and urine following arterial infusion of 150 μg hEPO mRNA via DSPE- and TPGS-

LNP during NMP of porcine kidneys. hEPO levels detected by ELISA. (a) Human EPO levels

measured in the perfusate over 6 hours. Significance of TPGS-LNP or DSPE-LNP against

the control is determined by 2-way ANOVA test with Dunn's multiple comparisons. * p <

0.05, ** p < 0.01, *** p < 0.001. (b) Human EPO levels detected in the urine 4-6 hours after

mRNA infusion. Significance is determined by Kruskal-Wallis. * p < 0.05, ** p < 0.01, *** p

< 0.001.

Figure 4: Levels of mCherry detected by western blot throughout several kidney regions

following arterial infusion of 150 µg mCherry mRNA via DSPE- and TPGS-LNP during NMP

of porcine kidneys. (a) Representative western blot showing mCherry expression (28 kDa)

in kidney cortical tissue after delivery with DSPE- and TPGS-LNPs. (b) Bar plot quantifying

mCherry levels in different kidney regions, including the cortex (cranial, lateral, caudal)

and medulla obtained after the end of the perfusion. Significance is determined by two-

way ANOVA and Šídák's multiple comparisons. * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 5: mCherry mRNA and mCherry protein presence in kidney tissue following arterial

infusion of 150 µg mCherry mRNA via DSPE- and TPGS-LNP during NMP of porcine kidneys.

mCherry mRNA presence detected by brightfield microscopy after Brightfield RNA In Situ

Hybridation (BRISH) analysis and mCherry protein presence observed by fluorescence

microscopy after immunofluorescence (IF). (a) Fluorescence microscopy images of

porcine kidney tissue showing mCherry protein expression by IF (red, Alexa 647) and

nuclear staining (blue, DAPI) after delivery with DPSE-, TPGS-LNPs and perfused kidneys

without mRNA-LNP treatment. (b) Brightfield imaging of BRISH analysis showing the

presence of mCherry mRNA (brown precipitate) and hematoxylin counterstaining (blue).

Kidney structures are marked accordingly: glomeruli (▲), tubuli (■) and vascular structures

(●).

Figure 6: mCherry mRNA and mCherry protein presence in kidney tissue following arterial

infusion of 150 µg mCherry mRNA via DSPE- and TPGS-LNP during NMP of human kidneys.

mCherry mRNA presence detected by brightfield microscopy after Brightfield RNA In Situ

Hybridation (BRISH) analysis and mCherry protein presence observed by fluorescence

microscopy after immunofluorescence (IF). (a) Fluorescence microscopy images of

porcine kidney tissue showing mCherry protein expression by IF (red, Alexa 647) and

nuclear staining (blue, DAPI) after delivery with DPSE-, TPGS-LNPs and kidneys with no

protein expression after mRNA-LNP treatment. (b) Brightfield imaging of BRISH analysis

showing the presence of mCherry mRNA (brown precipitate) and hematoxylin

counterstaining (blue). Kidney structures are marked accordingly: glomeruli (▲), tubuli (■)

and vascular structures (●).

a

	Control	F127-LNP	TPGS-LNP	DSPE-LNP	LIPO
eGFP					
mRNA-Cy5					
Hoechst (Nuclei)					
Merge					

b

c

a

0 1 2 3 4 5 6 7 8 9

35kDa ——
25kDa ——   DSPE-LNP

35kDa ——
25kDa ——   TPGS-LNP

b

a

Red: mCherry protein
Blue: DAPI (nuclei)
―――― 100 μm

b

Brown: mCherry mRNA
Blue: Hematoxylin (counterstaining)
―――― 100 μm

a

DSPE-LNP

TPGS-LNP

NEGATIVE

Red: mCherry protein
Blue: DAPI (nuclei)
———— 100 µm

b

DSPE-LNP

DSPE-LNP

TPGS-LNP

Brown: mCherry mRNA
Blue: Hematoxylin (counterstaining)
———— 100 µm