# Strain, procedures, and tools for reproducible genetic transformation and genome editing of *Spirodela polyrhiza* (L.) Schleid.

*Verónica Barragán-Borrero[1*], Amanda de Santana Lopes[2*], Enrico Diniz Rodrigues Batista[2], Martin Höfer[2], Rana Elias[1], Abhisek Chakraborty[2], Arturo Ponce-Mañe[1], Clotilde Descombes[1,3], Laura Diezma-Navas[1], Lydia Petraki[1,4], Meret Huber[2**], Shuqing Xu[2**] and Arturo Marí-Ordóñez[1**].*

[1] Gregor Mendel Institute of Molecular Plant Biology (GMI), Austrian Academy of Sciences, Vienna Biocenter (VBC), Dr. Bohr-Gasse 3, 1030 Vienna, Austria.

[2] Institute of Organismic and Molecular Evolution (IomE), Johannes Gutenberg University, Mainz Biozentrum I, Hanns-Dieter-Hüsch-Weg 15, 55128 Mainz. Germany.

[3] Current address: Institute of Plant Sciences Paris-Saclay (IPS2), Centre National de la Recherche Scientifique, Université Paris-Saclay, Gif Sur Yvette 91190, France.

[4] Current address: Agricultural University of Athens, 75 Iera Odos, 11855 Athens, Greece.

* These authors contributed equally.

** The authors responsible for correspondence and distribution of materials integral to the findings presented in this article are:

Arturo Marí-Ordóñez (arturo.mari-ordonez@gmi.oeaw.ac.at)

Shuqing Xu (shuqing.xu@uni-mainz.de)

Meret Huber (meret.huber@uni-mainz.de)

**Keywords**: Duckweeds, Spirodela, transformation, regeneration, CRISPR/Cas9

# Summary

Duckweeds (*Lemnaceae*) have excellent potential as plant systems for fundamental and applied research due to the ease of cultivation, their small size, and continuous fast clonal growth. However, their usage as model organisms and platforms for biotechnological applications is often limited by the lack of universal genetic manipulation methods necessary for insertional mutagenesis, reporter gene expression, gene editing, and other methods to modify gene expression. Here, we present a robust transformation protocol for a publicly available strain of *Spirodela polyrhiza*, *SP162,* that we identified to be amenable to *Agrobacterium*-mediated stable transformation via tissue culture. The transformation procedure allows stable expression of different reporter genes and selectable markers and enables CRISPR/Cas9-mediated genome editing. Due to the lack of a small RNA-based silencing response, *S. polyrhiza SP162* also sustains prolonged periods of transgene activity in *Agrobacterium*-mediated transient expression assays. Thus, *SP162* is an ideal reference strain for developing improved transformation procedures and applications in *S. polyrhiza*. To promote duckweed research and encourage the adoption of *S. polyrhiza* in research laboratories, we have made *SP162* (ID#: *5676*) and its genome publicly available and provide here detailed procedures for its cultivation and transformation.

## Introduction

Despite the vast diversity of land plants, most mechanistic understanding of plant gene function stems from few model systems with well-established genetic engineering protocols, such as *Arabidopsis thaliana*, *Oryza sativa* (rice), and *Marchantia polymorpha*. Research with these models has advanced plant biology tremendously yet revealed only a narrow slice of the evolutionary, ecological, and physiological diversity across the plant kingdom. Expanding the repertoire of genetically tractable species is therefore essential, not only to uncover lineage-specific adaptations, but also to develop sustainable applications capitalizing on the unique traits of underexplored plants.

Among these are duckweeds, floating aquatic plants in the family *Lemnaceae* comprising 36 species across five genera (Tippery et al. 2021). As the smallest flowering plants, they develop a simple body with 1-10 mm leaf-like fronds and hairless adventitious roots, with some genera having completely lost roots and vasculature (Acosta et al. 2021; Ziegler et al. 2023). Their exceptionally rapid growth, ease of cultivation, and efficient uptake of labeled compounds made duckweeds attractive model organisms for plant physiology and ecology in the mid-20th century (Hillman 1961; Acosta et al. 2021). However, interest declined in the 1990s with the rise of more genetically tractable models, like *A. thaliana*.

Recently, duckweed research has resurged, driven by advances in omics, renewed interest in plant ecological diversity, and potential applications in environmental and agricultural biotechnology (Acosta et al. 2021; Yang et al. 2021). Their small size and clonal propagation make them especially suitable for controlled multigenerational experiments, crucial for studying transgenerational plasticity, epigenetic memory, and stress responses (Huber et al. 2021; Antro et al. 2023; Chávez et al. 2025). Duckweeds are also emerging as key contributors to circular bioeconomy, through applications such as wastewater nutrient recycling (Cheng and Stomp 2009), phytoremediation (Ekperusi et al. 2019), carbon sequestration, and bio-based protein and starch production for food, feed, and fuels (Cui and Cheng 2015; Appenroth et al. 2017; 2021).

Among duckweeds, *Spirodela polyrhiza* stands out as a promising model. It possesses the smallest known genome in its family (140–158 Mb) (Wang et al. 2014; Harkess et al.

2021), and, despite its name "giant duckweed", still offers rapid clonal growth and large fronds for experimental manipulation. Its simple development, high phenotypic plasticity, and stress tolerance, make it valuable for fundamental molecular, developmental, evolutionary and ecological studies as well as applied research (Huber et al. 2021; Baggs et al. 2022; Ware et al. 2023; Sun et al. 2023; Turcotte et al. 2024; Wang et al. 2024; Malacrinò et al. 2024; Smith et al. 2024; Chávez et al. 2025; Böttner et al. 2025).

Despite these desirable characteristics, absence of a reliable, reproducible protocol for stable genetic transformation limits the exploitation of *S. polyrhiza* as a model. While transient gene expression via carbon nanotubes (Islam et al. 2024) and *Agrobacterium*-mediated DNA delivery (Peterson et al. 2021; Dombey et al. 2025) have been reported, only one publication described stable transformation (Yang et al. 2018). However, the study in a not publicly available genotype lacked adequate genotypic and phenotypic characterization, not allowing to reproduce the findings.

To complement the toolbox for duckweed research, we developed a detailed, reproducible *Agrobacterium*- and tissue culture-based transformation protocol for a publicly available *S. polyrhiza* genotype. We generated stable transgenic lines expressing various reporter genes and selectable markers and demonstrated CRISPR/Cas9-mediated genome editing. This know-how will support the growing duckweed research community and help unlock the potential of *S. polyrhiza* for fundamental discoveries and sustainable biotechnological applications.

## Results

### *S. polyrhiza* genotypes vary for callus induction rates.

Success rates in transformation protocols including tissue culture steps vary between strains and genotypes of the same species depending on the ability to form callus (Abe and Futsuhara 1986; Nguyen et al. 2020; Lardon et al. 2020). Therefore, we screened a collection of *S. polyrhiza* genotypes (Wang et al. 2024) for strains displaying high callus induction rates. We first established a suitable callus induction procedure by testing three published protocols for

*S. polyrhiza* (Wang 2016; Huang et al. 2018; Yang et al. 2018), and one for the related species *Lemna aequinoctialis* (Liu et al. 2019). The protocols, differing in the type of explants, composition of phytohormones, and their concentration (**Figure 1a**), were tested on axenic cultures (**Method S1, S2**) of four *S. polyrhiza* genotypes from diverse geographical origin. The methods based on wound-induced callus formation (Wang 2016) and root-derived calli from dorsoventrally laid fronds (Huang et al. 2018) had the highest callus induction rates (**Figure 1b**) and produced calli from frond cuttings or root meristems as expected from either protocol (**Figure 1c**).

The less laborious procedure with root-derived calli was chosen to expand the screen to 35 *S. polyrhiza* genotypes. Callus induction rates varied across genotypes, with some South-East Asian and European genotypes displaying mean callus induction rates above 60% (**Figure 1d**). We also observed variability between genotypes for the tissue-of-origin of the calli: in addition to the calli initiated from root meristems (R), some genotypes formed calli also from vegetative budding pockets (Po), where shoot meristems are located, or from stipes (Sti), the stalk-like structure connecting mother-to-daughter fronds (**Figure 1e**). Hence, *S. polyrhiza* displays variation across genotypes in callus induction responses, both in initiation rate and tissue-of-origin.

**The *S. polyrhiza SP162* genotype can be regenerated into fronds.**

High rates of callus induction does not necessarily result in successful plant regeneration, two independent and often uncorrelated processes (Abe and Futsuhara 1986; Ikeuchi et al. 2013; Motte et al. 2014; Ikeuchi et al. 2019). Therefore, we examined regeneration in two genotypes with high callus induction, the Asian *SP162* and European *9509* (**Figure 1d**). *9509* offers extensive genomic resources, including an assembled reference genome, transcriptomes and epigenomes (Hoang et al. 2018; Harkess et al. 2021; 2024; Dombey et al. 2025). *SP162*, despite lacking such resources, exhibited good formation of turions, sinking dormant

**Figure 1: Identification of compatible procedures and genotypes for *S. polyrhiza* callus induction and frond regeneration. (a)** Types of explants and concentrations of phytohormones (growth regulators) used in previously published protocols for callus induction tested here. **(b)** Mean callus induction rate (as % of explants initiating callus formation) in four different *S. polyrhiza* genotypes, from different origins indicated above, using the protocols in (a). Dots display individual values from independent replicates. Error bars represent standard deviation of the mean. N.D.: not determined. **(c)** Pictures of callus induced from cutting explants or roots according to the indicated published protocols. **(d)** Mean callus induction rate in % of 35 *S. polyrhiza* genotypes from different geographic origin using the Huang *et al.* callus induction protocol. Dots display individual values from three independent replicates. Error bars represent standard deviation of the mean. **(e)** Schematic representation of *S. polyrhiza* frond structure from a ventral position and pictures of different genotypes with ongoing callus formation from different tissues using Huang *et al.* callus induction protocol. R: roots (black arrow), Po: budding pockets (magenta arrow), Sti: stipe, stalk-like connecting structure between mother and daughter fronds (blue arrow). **(f)** *S. polyrhiza* SP162 calli growth rate in % (as increase of biomass) (dots display individual values from three independent replicates. Error bars represent standard deviation of the mean) and representative images during regeneration following Huang *et al.* protocol with increased thidiazuron (TDZ) to 1 mg/L. **(g)** Same as (f) but for *S. polyrhiza* 9509.

vegetative propagules (Landolt 1986; Appenroth 2002; Ziegler et al. 2023; Kim et al. 2024), with near 100% germination after months of cold storage (**Figure S1, Method S3**), providing a convenient storage solution. In contrast, *9509* lacked turion production in our hands.

Despite Huang and colleagues had shown that *S. polyrhiza* regeneration is best induced at low concentrations of the synthetic cytokinin thidiazuron (TDZ) (0.2 mg/L, 0.9 µM), or in absence of any cytokinin (Huang et al. 2018), neither genotype initiated regeneration under these conditions. Others had reported increasing regeneration efficiencies at higher TDZ concentrations (1 mg/L, 4.54 µM) (Wang 2016; Yang et al. 2018). Indeed, both genotypes initiated regeneration in the presence of 1mg/L TDZ (**Figure 1f,g**). However, complete regeneration of individual fronds was only achieved in *SP162* (**Figure 1f**), whereas *9509* calli produced only elongated frond-like structures but not fully regenerated fronds (**Figure 1f**). Therefore, *SP162* was selected as the genotype for further studies.

**Callus induction and maintenance under low light improves *SP162* callus growth and proliferation.**

Although *SP162* calli did regenerate, they were prone to browning despite the presence of anti-browning agents, polyvinylpyrrolidone (PVP) and citric acid, in the callus induction medium (CIM) (**Figure 2a, Method S1**) and grew slowly (**Figure 1f**), two factors negatively affecting *Agrobacterium*-mediated transformation and regeneration (Potrykus 1990; Sangwan et al. 1992; Greenwood and Glaus 2022; Liu et al. 2024). Browning of *S. polyrhiza* calli can be prevented in darkness, although this results in decreased proliferation (Huang et al. 2018). Therefore, we tested whether low light conditions could represent a compromise to reduce browning without limiting growth and proliferation. We performed *SP162* callus induction on CIM at 21ºC and covered the culture plates with two sheets of white paper (80 gr/m$^2$), bringing light intensity during the 16 h light period to 20 µmol/m$^2$/s (**Figure 2b, Method S4**). Swelling of young roots was already apparent after a week (**Figure S2**). By the fourth week, 2-3 mm wide dark-green calli had formed at root tips and budding pockets (**Figure 2c, S2**). Newly

**Figure 2:** *S. polyrhiza SP162* **callus induction and frond regeneration under low light. (a)** Composition of the *SP162* callus induction medium (CIM). **(b)** Schematics of *SP162* callus induction conditions under low light and images of experimental setup and fronds in dorsoventral position on CIM. **(c)** *SP162* calli induced from either roots (R) or budding pockets (Po) after four weeks in CIM under low light. **(d)** Images of initial calli transferred to CIM and 10 weeks after callus induction starting to develop pale sectors cultured on CIM under low light. **(e)** Images of selected and amplified pale friable calli sectors grown under low light. **(f)** Comparative growth rate in % (as increase of biomass) of dark-green calli cultured under high light and pale-green calli cultured under low light. Dots display individual values from three independent replicates. Error bars represent standard deviation of the mean. *p* indicates two-tailed P-value of unpaired t-test between the two groups. **(g)** Composition of the *SP162* frond regeneration medium (FRM) medium. **(h)** Schematic of the transfer procedure from CIM to FRM to initiate regeneration under low light. **(i)** Image of early regenerating fronds (marked with white arrow) after 4 weeks of regeneration (wor) on solid FRM. **(j)** Schematics and images of the two-step acclimation for frond regeneration in liquid media together with pictures of regenerating fronds after removal of cytokinin and sugar after 6 wor. **(k)** Images of a regenerated frond in SH-medium in culture plate and proliferation after individualization.

formed calli under low light showed little or no browning (**Figure 2c, S2**). Hence, reducing light intensity had a beneficial impact on callus induction by producing healthier calli.

After four weeks, calli were detached and subcultured under low light conditions, but temperature was increased to 26ºC to promote faster growth (**Figure 2d**). About six weeks later, pale-green sectors started to appear. These were less compact, more friable (**Figure 2d**), and seemed to proliferate faster. They were isolated, subcultured and amplified (**Figure 2e, Method S4**). Even after several months of culture under low light, pale-green calli did not show browning and grew five to six times faster than dark-green calli cultured without shading (**Figure 2f**).

**_SP162_ calli can regenerate through a two-step acclimatation process under low light.** Given the positive impact of low light on callus health and growth, we tested whether _SP162_ pale calli could regenerate under low light on frond regeneration medium (FRM) containing either 1 mg/L TDZ (**Figure 1f**), or zeatin, another cytokinin shown to induce high levels of regeneration in _S. polyrhiza_ at 2.2 mg/L (10 µM) (Wang 2016) (**Figure 2g, Method S1**). Callus fragments were cultured on solid FRM with either cytokinin under callus induction and maintenance conditions (**Figure 2h, Method S5**). After four weeks, regeneration initiated, with small dark-green frond-like structures appearing, singly or in clusters, in both TDZ or zeatin containing FRM (**Figure 2i**, **Figure S3**).

To progressively acclimate regenerating fronds to autotrophic growth in standard liquid culture conditions, we applied a stepwise process. First, calli with regenerating fronds were accustomed to liquid media and increased light intensity (125 µmol/m$^2$/s) by culturing them in Magenta vessels in 5 cm-deep liquid Schenk and Hildebrant (SH)-medium with sucrose but without growth regulators for seven days. Afterwards, they were moved to 1 cm shallow SH-medium without sugar in culture dishes to improve gas exchange and light exposure at 21ºC (**Figure 2j, Method S1, S5**).

Around six weeks after induction, regenerating fronds visible to the naked eye provided increased buoyancy and calli emerged to the surface (**Figure 2j, S4a**). Two weeks later, individual fronds started to appear on the surface of the culture and were regularly transferred to individual flasks (**Figure 2k, Method S5**). Frond clusters continued to appear up to twelve

weeks after induction (**Figure S4b**). Thus, we established culture conditions enabling healthy

and fast-growing calli of *SP162* to regenerate into fronds.

**Figure 3: Sensitivity of *S. polyrhiza SP162* calli to antibiotics. (a)** Survival rates (%) of calli exposed to different concentrations of the indicated antibiotics along a 20-day period. Dots indicate average survival rates and error bars represent standard deviation of the mean of 5 replicates. Survival was assessed as absence of chlorosis, browning or impact on growth. **(b)** Representative images of calli exposed to different concentrations of indicated antibiotics in CIM media after transfer to media with antibiotics, or without as control, and two weeks later. See Figure S5 for a broader range of concentrations.

**SP162 calli are sensitive to several selective antibiotics.**

Crucial for the successful identification and amplification of transgenic cells is the use of selectable markers to eliminate or arrest the proliferation of non-transgenic cells. To identify suitable selective agents for *SP162* calli, we tested their sensitivity to different concentrations of four broadly used antibiotics in plant selection: kanamycin, hygromycin, G418/geneticin, and spectinomycin. *SP162* calli were sensitive to all four antibiotics, but to different degrees. While calli showed lower sensitivity to kanamycin, they displayed high sensitivity to the other three antibiotics, with spectinomycin and hygromycin being the most efficient at a broad range of concentrations (**Figure 3, S5**). Given that hygromycin had the fastest impact on calli survival at low concentrations (**Figure 3, S5**), hygromycin resistance was chosen as selectable marker for testing transformation in *SP162*.

**SP162 calli are suitable for stable *Agrobacterium*-mediated transformation.**

To assess the feasibility of *Agrobacterium*-mediated transformation in *SP162*, we aimed at transforming calli with green fluorescent protein (*GFP)* driven by the maize *UBIQUITIN* promoter (*pZmUBQ::eGFP*), suitable for strong ubiquitous expression in monocots (Christensen and Quail 1996), and hygromycin resistance (*HptII*) under the *NOPALINE SYNTHASE* promoter *(pNOS)* as a selectable marker. *Agrobacterium tumefaciens EHA105* was chosen as it was shown to deliver DNA to *S. polyrhiza* for transient expression (Dombey et al. 2025).

Small callus fragments were inoculated with *Agrobacterium* carrying the *pZmUBQ::eGFP* T-DNA vector and co-cultured for two days before rinsing off the *Agrobacterium* (**Figure 4a, Method S1, S6**). Calli were transferred to CIM medium containing antibiotics, ticarcillin and cefotaxime, to select against residual *Agrobacterium* for recovery (**Figure 4b, Method S1, S6**). To confirm transformation and assess the impact of *Agrobacterium* on callus health, calli appearance and *GFP* expression were regularly monitored. Already two days post inoculation (dpi*),* bright GFP fluorescence spots were observed under UV light (**Figure 4b**). Calli grew for two weeks without displaying browning or

any side effects from exposure to the bacteria, while GFP fluorescent sectors increased in number and size (**Figure 4b**).

**Figure 4:** *Agrobacterium*-mediated stable transformation and selection of *S. polyrhiza SP162* calli. **(a)** Schematic representation of the calli transformation procedure. **(b)** Representative images under visible and UV light of *SP162* calli in recovery 2, 7 and 14 days after inoculation (dpi) with *Agrobacterium* (*EHA105*) carrying a T-DNA vector for constitutive e*GFP* expression. **(c,d)** Same as in (b) but during weak (5 mg/L hygromycin, c) and high (10 mg/L hygromycin, d) selection of transformed cells. **(e)** Schematics of isolation and amplification of hygromycin-resistant calli and images under visible or UV light of a selected transgenic callus and plates of non-transformed and transgenic calli. In all UV images: green: GFP fluorescence, red: chlorophyll autofluorescence. **(f)** Southern blot analysis of T-DNA integration events in five independent calli. T-DNA integration was assessed by digesting gDNA with *XhoI* and detected with a probe against the *pZmUBQ* as indicated in the T-DNA scheme. Ethidium bromide (EtBr) staining of the gel prior to transfer is shown as loading control. n.t.: non-transformed regenerated wild-type (WT) control; Ld: DNA size ladder.

Hygromycin selection was applied progressively to avoid browning of dying cells, which can have toxic effects on neighboring transformed cells (Sangwan et al. 1992; Zhang et al. 2020; Liu et al. 2024). Weak selection with a low hygromycin concentration (5 mg/L) resulted in chlorosis and loss of chlorophyll autofluorescence (**Figure 4c, Method S1, S6**). After two weeks, selection pressure was increased to 10 mg/L hygromycin (**Method S1, S6**), causing further chlorosis and browning, while hygromycin-resistant, GFP-positive sectors started to enlarge (**Figure 4d**). These healthy sectors were transferred to either CIM medium without antibiotics for amplification or kept on 10 mg/L hygromycin if resistant sectors were small, heterogenous, or difficult to isolate from non-resistant tissue (**Figure 4d,e, Method S6**). Individual hygromycin-resistant sectors were subcultured to obtain larger quantities of fast-growing and uniformly GFP-positive calli, candidates for successful integration of the transgenes (**Figure 4e**).

T-DNA integration into the genome was confirmed by Southern blot analysis of five calli cultures, revealing single to multiple T-DNA insertion events (**Figure 4f**). Thus, *Agrobacterium*-mediated transformation enables stable genetic transformation of *SP162* calli.

**Transformed *SP162* calli regenerate into stable transgenic lines.**

Since T-DNAs can become unstable during tissue culture and plant regeneration (Peerbolte et al. 1987; Phillips et al. 1994; Risseeuw et al. 1997), we investigated T-DNA stability in line #1 (**Figure 4f**) with a single insertion of the *pZmUBQ::eGFP* transgene. As non-transformed *SP162* calli, transgenic calli started to regenerate after four weeks in FRM, with the formation of darker frond-like structures with higher chlorophyll autofluorescence (**Figure 5a**). Transgenic frond regeneration also proceeded normally after acclimation (**Figure 5b**). Fully regenerated GFP-positive fronds started to appear within 7-8 weeks and were able to clonally propagate (**Figure 5c**). Ten regenerated lines were amplified and subjected to Southern blot analysis. All lines carried the single T-DNA insertion (**Figure 5d**) from the initial callus line. Therefore, at least for this single T-DNA insertion, the integrated DNA remained stable during tissue culture and frond regeneration.

**Figure 5: Frond regeneration and molecular characterization of *S. polyrhiza SP162* transgenic plants. (a-c)** Schematics and representative images under visible and UV light of *SP162* single T-DNA insertion transgenic calli expressing e*GFP* during the regeneration initiation (a), at the end of acclimation (b) or at the end of regeneration (c). In all UV images: green: GFP fluorescence, red: chlorophyll autofluorescence. **(d)** Southern blot analysis of T-DNA integration in 10 cultures of independently regenerated transgenic fronds. T-DNA integration was assessed as indicated in Figure 4f. Ethidium bromide (EtBr) staining of the gel prior to transfer is shown as loading control. n.t.: non-transformed regenerated wild-type (WT) control; Ld: DNA size ladder. **(e)** Images of individual fronds from one non-transformed and five transgenic regenerated lines scanned under white or UV light in a biomolecular scanner. In UV images, green: GFP florescence, red: chlorophyll autofluorescence. **(f)** qPCR analysis of e*GFP* expression relative to endogenous *ACTIN* mRNA levels in the same regenerated lines as in (e). Median values (bars) from 3 technical replicates (dots) are represented for each biological sample. **(g)** Protein blot analysis of GFP accumulation in the same lines as in (e,f). Rubisco (Rbcs) bands in Coomassie (coom.) staining of the blot after GFP immuno-detection is shown as loading control. Mw: Molecular weight. **(h)** Images of transgenic callus and regenerated fronds expressing the *RUBY* visual reporter. **(i)** Images of transgenic fronds expressing the *DsRed* fluorescent reporter under visible or UV light.

Furthermore, after continuous culture for six months, GFP fluorescence was homogeneous across single-insertion regenerated lines (**Figure 5e**), with high levels of *GFP* mRNA expression and comparable levels of GFP protein accumulation across lines (**Figure 5f,g**). Thus, transgene expression also remained stable after prolonged periods of clonal propagation.

**_SP162_ transformation can be independently reproduced with different reporter genes and selectable markers.**

To test the reproducibility of *S. polyrhiza SP162* transformation, the process was repeated with different reporter genes and in a different laboratory. First, we tested the betalain-based visual reporter *RUBY, pZmUBQ::RUBY* (He et al. 2020). Both, transgenic calli and regenerated fronds, displayed the bright red color characteristic of betalain accumulation (**Figure 5h**). Next, the transformation and regeneration procedures were independently reproduced in a second laboratory to obtain transgenic *S. polyrhiza* cultures transformed with *pZmUBQ::eGFP* (**Figure S5**), as well as other fluorescent reporters such as *DsRed2* (*pZmUBQ::DsRed2*) (**Figure 5i, S5b**), or using spectinomycin resistance as selectable marker (**Figure S5b, Method S6**). These results showcase the applicability of different reporter genes or selectable markers and further consolidate the reproducibility and practicality of the methodology used here to perform genetic transformation in the *S. polyrhiza SP162* strain.

**CRISPR-Cas9 genome editing can be achieved in _SP162_ to generate mutations.**

Lack of regular flowering and sexual reproduction in *S. polyrhiza* (Fourounjian et al. 2021) hinders classic mutagenesis methods that require selfing or backcrossing to fix desired mutations and clean the background (Koornneef 2002). However, CRISPR/Cas technology allows to precisely introduce mutations through targeted genome editing (Jung and Till 2021; Cardi et al. 2023). Hence, we tested if CRISPR/Cas9 was suitable to generate mutants in *S. polyrhiza SP162*.

**Figure 6: CRISPR/Cas-9-mediated genome editing in *S. polyrhiza SP162*.** **(a)** Vector for simultaneous expression of *zCas9i-2A-eGFP* and multiplexed sgRNAs. **(b)** Schematic representation of the *SpZMET* gDNA region targeted by sgRNAs (orange boxes), the SpZMET protein and its domains. NLS: nuclear localization signal; BAH: Bromo-adjacent homology domain; Chromo: Chromatin organization modifier domain; DNA methylase: C-5 cytosine-specific DNA methylase. **(c)** Image of WT and transgenic *SP162* calli transformed with the indicated plasmids under visible and UV light. Green: GFP fluorescence; red: chlorophyll autofluorescence. **(d)** Protein blot analysis of zCas9i-2A and GFP accumulation in regenerated WT, e*GFP,* and three *zCas9i-2A-eGFP 4x@ZMET* calli. Rubisco (Rbcs) in Coomassie (coom.) staining of blots is shown as loading control. Mw: Molecular weight. **(e)** Images of individual fronds from one non-transformed (WT), one *eGFP* and four *zCas9i-2A-eGFP 4x@ZMET* transgenic regenerated lines scanned under white or UV light in a biomolecular scanner. In UV images, green: GFP fluorescence, red: chlorophyll autofluorescence. **(f)** Protein blot analysis of zCas9i-2A and GFP accumulation in non-transgenic and four *zCas9i-2A-eGFP 4x@ZMET* regenerated lines. Rubisco (Rbcs) in Coomassie (coom.) staining of blots is shown as loading control. **(g)** Agarose-gel electrophoresis of PCR amplification of *SpZMET* gDNA in non-transgenic and four *zCas9i-2A-eGFP 4x@ZMET* regenerated lines. Primers located outside of sgRNA targeted regions were used as indicated by arrows in the diagram above. Bands framed in red were isolated for subsequent cloning and Sanger sequencing analysis to investigate editing events. Ld: Ladder. **(h)** Graphical summary of the analysis of PCR products from (g) displayed in Figure S8. Resulting changes in DNA and predicted resulting protein sequences are shown. **(i)** Schematic representation of Cas9-mediated cleavage relative to the Protospacer Adjacent Motif (PAM) and target sgRNA sequence. **(j)** WT reference sequence, and Sanger sequencing chromatograms of examples of single nucleotide deletion and large nucleotide deletion between sgRNAs found in *SP162* regenerated line #1. PAM sequence is highlighted in blue and target sequence in orange.

The CRISPR/Cas9 genome editing requires designing single-guide RNAs (sgRNAs) that exactly match the DNA target site. Therefore, we sequenced and assembled *S. polyrhiza SP162* using the PromethION (Oxford Nanopore Technologies) (**Figure S7**). We then built a single-vector sgRNA multiplexing CRISPR/Cas9 system based on Liu's laboratory tools (Ma et al. 2015; Ma and Liu 2016). We used the intron-containing and maize codon-optimized *Cas9* (*zCas9i*), shown to improve editing efficiencies in plants (Grützner et al. 2021; Stuttmann et al. 2021), under the *ZmUBQ* promoter (**Figure 6a**). To facilitate selection and monitor *zCas9i* expression, we transcriptionally fused *GFP* to *zCas9i* through the *porcine teschovirus-1 2A* ribosome skipping peptide (2A), allowing independent translation of Cas9 and GFP proteins from the same ORF (*pZmUBQ::zCas9i:P2A:eGFP*; **Figure 6a**). As 2A amino acids remain appended at the C-terminus of the upstream protein, serving as tag for Cas9 detection.

As proof-of-concept, we targeted *ZMET* (SpZMET), a chromomethyltransferase required for DNA methylation at CHG sites (Bewick et al. 2017). Because *S. polyrhiza* has low CHG methylation (Michael et al. 2017; Dombey et al. 2025), we hypothesized that SpZMET loss-of-function would not impair callus survival or regeneration. To test small (1-100 bp) and larger (500-800 bp) deletions, four sgRNAs were designed in pairs flanking the SpZMET nuclear localization signal (NLS) (**Figure 6b**). The resulting plasmid (*pZmUBQ::zCas9i:P2A:eGFP_4x@ZMET*) was transformed into *SP162* calli as described above. Although GFP fluorescence was weaker in *Cas9* transgenic calli than in GFP-only lines (**Figure 6c**), western blotting confirmed Cas9-2A and GFP accumulation (**Figure 6d**). Cas9 expression did not impair frond regeneration and, albeit GFP fluorescence was low, Cas9-2A and GFP continued to be expressed in regenerated lines (**Figure 6e-f**).

Editing was first assessed in four regenerated lines by PCR amplification of the region targeted by the sgRNAs. Two of the lines produced additional shorter amplicons, compatible with large (~600 bp) deletions (**Figure 6g**). Sequencing of PCR products confirmed that in three lines, all sequenced molecules carried mutations ranging from single nucleotide to larger (600 bp) deletions (**Figure 6h, S8**). Mutations were consistent with those expected from Cas9-

mediated cleavage upstream of the third nucleotide after the PAM sequence (Jinek et al. 2012) (**Figure 6i**), resulting in single nucleotide deletions (SND) or elimination of sequences between two cleavages (**Figure 6h,j, S8**). Other known mutations induced by Cas9, such as single nucleotide insertions or larger deletions (2-20 nt), were also observed (**Figure 6h, S8**). Low frequency mutations (e.g. sgRNA4 at band 1.1, 1.2 and 3; **Figure S8**) likely resulted from constitutive Cas9 expression leading to chimeric individuals or cultures.

SND and small deletions generally resulted in out-of-frame mutations, generating putative knockout (ko.) alleles in all lines (**Figure 6h**). In-frame deletions between sgRNA1 and sgRNA3 eliminated much of the N-terminal region, including the NLS, in lines #1 and 2 (**Figure 6h,i**). Thus, our data demonstrate that CRISPR/Cas9 efficiently induced both frameshift and domain-specific deletions in *SP162*, so that the transformation protocol can be applied to create targeted gene editing in *S. polyrhiza*.

**Regenerated *SP162* lines display phenotypic variation.**

Despite the reproducibility and stability of transformation, we noticed that lines of regenerated plants looked and grew differently from the original *WT SP162* and between themselves. The most obvious observable difference was frond size, as regenerated lines were smaller than *WT SP162* (**Figure S9a**). While this difference could be as little as 0.5 mm in length, *pZmUBQ::RUBY* lines were up to 2 mm, about 30%, shorter than *WT* (**Figure S9b**).

In addition, cultures of regenerated plants initiated from similar frond numbers grew to occupy less surface and displayed increased senesce in the same amount of time compared to *WT* (**Figure S10a-c**). While reduced culture surfaces could be explained by smaller frond size, their growth rates were also lower (**Figure S10d**). Although growth patterns of all lines still fit an exponential curve (**Figure S10e,f**), doubling times were delayed by nearly one day for *GFP* lines (3.1-3.5 days) and two for *RUBY* lines (4.2-4.5 days), almost twice the doubling time of *WT* (2.4 days) (**Figure S10f**). Similar phenotypes were observed in the *Cas9*-expressing lines (**Figure S11**).

The reduced growth rates of *RUBY* lines could be consequence of metabolism rerouting and tyrosine depletion for betalain production (He et al. 2020), as high *RUBY*

expression negatively affects growth and development also in other plants (Lee et al. 2023; Tse et al. 2024; Swinnen et al. 2025). Thus, phenotypic variation between *RUBY* lines (**Figure S9, S10**) could be due to different transgene expression levels. Likewise, there was phenotypic variation between the *Cas9* lines (**Figure S11**). It remains to be determined whether such variation is the result of the modified trait in other cases of mutants or transgene-expressing lines. However, similar variation was observed between *GFP* lines (**Figure S9, S10**) that all originate from the same single T-DNA insertion event (**Figure 4f, 5d**). Hence, the underlying cause of the phenotypic differences might be induced rather by tissue culture and/or regeneration.

**Phenotypic variation in regenerated lines is not associated with changes in ploidy.**

Plants obtained through tissue culture usually display phenotypic variation from parental lines due to somaclonal variation generated during callus induction, propagation and/or regeneration (Bairu et al. 2011). Although its causes can be of epigenetic or genetic origin (Evans and Sharp 1986; Neelakandan and Wang 2012; Wibowo et al. 2022), changes in ploidy often correlate with changes in organ and plant size (Kuppu et al. 2015; Corneillie et al. 2018; Šmarda et al. 2023; Dermail et al. 2024). This is of special relevance for genome editing as gene copy number increases through polyploidy impact editing efficiency (Sánchez-Gómez et al. 2023). Therefore, we analyzed DNA content and estimated genome size of several regenerated lines, including those expressing *GFP* and *Cas9*, through flow cytometry (**Figure 2a,b**). No significant variations in genome size were observed between the original *WT SP162* and regenerated lines (**Figure S12c**). Hence, although a more detailed analysis of their genomes and epigenomes will be required to investigate the underlying cause behind the observed phenotypic variation, changes in ploidy during *SP162* tissue culture can be excluded.

**SP162 is also suitable for transient expression.**

In addition to stable transformation, *Agrobacterium*-mediated transient expression enables rapid transgene testing without the arduousness and time requirements of tissue culture. As

**Figure 7: *Agrobacterium*-mediated transient expression in *S. polyrhiza SP162*. (a)** Biomolecular scanning of GFP fluorescence (green) and chlorophyll auto-fluorescence (red) of *SP162* cultures infiltrated with *ZmUBQ::eGFP* at different days post infiltration (dpi). **(b)** Images of non-infiltrated and 21 dpi infiltrated cultures. **(c)** Colony forming units (cfu) in bacteria extracts from individual fronds from samples in (b) plated on *Agrobacterium* EHA105 selective media (magenta; rifampicin, streptomycin) or for bacteria carrying the *ZmUBQ::eGFP* plasmid (green; rifampicin, streptomycin, spectinomycin). Lines link extracts from single fronds plated in the two media **(d)** qPCR analysis of e*GFP* expression relative to endogenous *ACTIN* mRNA levels at the indicated samples and timepoints. Mean values (bars) from 3 biological replicates (dots) are represented for each sample. Error bars represent standard deviation of the mean. *p* indicates two-tailed P-value of unpaired t-test between the indicated groups. **(e)** Two exposure times of protein blot analysis of GFP accumulation during transient expression in three biological replicates (rep#) for each sample. Rubisco (Rbcs) in Coomassie (coom.) staining of blots is shown as loading control. Mw: Molecular weight. **(f)** Visible and UV images of *SP162* fronds infiltrated with *eGFP* under the control of an estradiol-inducible promoter in presence or absence of estradiol. Non-infiltrated and constitutively driven *eGFP* infiltrated samples are shown as control. **(g)** Schematic representation of *ZmUBQ::eGFP* and the *eGFP* derived hairpin (hp) *ZmUBQ::FP:PFhp* plasmids. **(h)** Small RNA blot analysis of samples used in (a-e) and infiltrated with *ZmUBQ::FP:PFhp*. *eGFP*-derived siRNAs were probed against full-length *eGFP*. Detection of *SpTAS3*- tasiRNAs served as control for siRNA biogenesis; miRNA159 and U6 snoRNA detection served as size and loading control, respectively.

the technique has been previously applied and achieved high levels of protein expression in other *S. polyrhiza* strains (Peterson et al. 2021; Dombey et al. 2025), we tested GFP (*pZmUBQ::eGFP*) transient expression in *SP162* using the *Agrobacterium* vacuum infiltration (agro-infiltration) method (Dombey et al. 2025) (**Method S7**).

GFP foci appeared three days post infiltration (dpi) and expanded for two weeks (**Figure 7a, S13**), a pattern independently reproduced in a second laboratory (**Figure S14**). However, after 21 days, infiltrated plants appeared unhealthy, showing chlorosis and growth reduction (**Figure 7b**). Such symptoms resembled those of *S. polyrhiza* infected with pathogenic bacteria (Baggs et al. 2022). Indeed, bacteria extracted from surface-sterilized fronds grew on selective media for *Agrobacterium* (rifampicin and streptomycin) and for plasmid-containing cells (rifampicin, streptomycin and spectinomycin) (**Figure 7c**). Thus, *Agrobacterium* likely persisted within tissues despite the continuous presence of cefotaxime and ticarcillin antibiotics (**Figure 7a, Method S7**), compromising long-term culture viability. To determine transgene expression dynamics, GFP RNA and protein levels were monitored. *GFP* mRNA was already detected at 3 dpi and peaked at 10 dpi and, albeit lower, remained high at 14 dpi (**Figure 7d**). GFP protein levels mirrored this trend, low at 3 dpi (only detectable in one of the biological replicates), rising at 6 dpi and persisting to 14 dpi (**Figure 7e**). We also tested the applicability of the methodology to more complex expression systems with an estradiol-inducible GFP construct. As expected, fluorescence only occurred after estradiol application to the media (**Figure 7f**). Similar induction was obtained in stable transgenic calli and regenerated fronds (**Figure S15**). Thus, *SP162* supports robust transient expression as an experimental system and is suitable for testing constructs before investing into the time and labor for stable transformation.

**Transient transgene expression does not trigger S-PTGS in *S. polyrhiza*.**

The transient expression dynamics seen here resembled previously reported *GFP* expression driven from the viral promoter *p35S* in *S. polyrhiza,* lasting up to 30 dpi (Peterson et al. 2021). Such a long period of transient transgene activity is unusual. Generally, in most tested experimental model plants like *Nicotiana benthamiana*, expression decays within days after

infiltration due to the onset of silencing. Sense post-transcriptional gene silencing (S-PTGS) upon overexpression of transgenes impedes continuous high mRNA levels during transient assays through the production of 21-22-nt-long short interfering (si)RNAs (**Figure S16a**) (Vaucheret et al. 1997; Voinnet et al. 2000; Hamilton et al. 2002; Kościańska et al. 2005; Luo and Chen 2007; Paoli et al. 2009; Parent et al. 2015). To counteract this, transgenes of interest are usually co-infiltrated with viral suppressors of RNA silencing (VSRs), like P19, to reduce S-PTGS (Voinnet et al. 2000; Hamilton et al. 2002; Qu et al. 2003; Mérai et al. 2005; Chen et al. 2013; Jay et al. 2023).However, P19 does not enhance GFP expression in *S. polyrhiza* (Peterson et al. 2021). Furthermore, while endogenous siRNAs (tasiRNAs) whose biogenesis depends on S-PTGS factors (Peragine et al. 2004; Yoshikawa et al. 2005; Gasciolli et al. 2005) are present (**Figure S16b**), no siRNA after *RUBY* agro-infiltration could be detected (Dombey et al. 2025), supporting no or only weak transgene-triggered S-PTGS.

Previous studies lacked either simultaneous siRNA or transgene mRNA measurements, especially relevant for *RUBY* as betalain pigmentation can remain long after transgene expression has faded (Tabara et al. 2024), leaving uncertainty about any correlation between mRNA levels and siRNAs. Therefore, we investigated the presence of *GFP*-derived siRNA in the same samples used to quantify *GFP* transient expression. As control, we infiltrated a *GFP*-derived hairpin construct (*pZmUBQ::FP:PFhp*) to trigger siRNAs independently of S-PTGS (**Figure 7g**, **S16c**) and monitored *SpTAS3*-tasiRNAs (**Figure S16d-g**). *GFP* siRNAs in cultures infiltrated with *FP:PFhp* were already detected at 3 dpi and accumulated strongly at 6 and 10 dpi but were undetectable in plants expressing *GFP*, even at 10 dpi with maximum *GFP* mRNA levels (**Figure 7h**). However, *SpTAS3*-tasiRNAs were present in all samples (**Figure 7h**), further supporting the hypothesis that the transgene-induced branch of S-PTGS might be reduced or compromised in *S. polyrhiza*, enabling prolonged transgene expression.

# Discussion

Despite recent advances (Yamamoto et al. 2001; Cantó-Pastor et al. 2015; Yang et al. 2018; Liu et al. 2019), the deployment of duckweeds as model organisms has been hindered by the lack of publicly available strains accessible for genetic modification and suitable protocols. We identified *S. polyrhiza SP162* as a competent strain for stable transformation and genome editing and provide detailed, versatile, and reproducible protocols, including also transient expression. In addition, and to facilitate the adoption of *S. polyrhiza* in research laboratories, we have also made *SP162* available through the European, North American and Asian duckweed germplasm collections under the ID#: *5676*, along with its genome and other aspects of *SP162* culture. We hope that *SP162* serves as a common platform to stimulate duckweed research and the approach, tools, and procedures described here will help screen and identify other suitable strains for transformation.

The variability in callus induction and response to growth regulators during regeneration observed by us and others (Wang 2016; Huang et al. 2018; Yang et al. 2018), suggests indeed natural diversity and potential to identify additional strains transformable with appropriate media and phytohormone compositions by systematically testing different conditions. Furthermore, novel technologies such as expression of *GRF-GIF* chimeric morphogenic regulators, shown to significantly boost transformation, editing, and regeneration efficiencies in plants (Debernardi et al. 2020; Maher et al. 2020; Vandeputte et al. 2024; Swinnen et al. 2025) could further boost these processes in *SP162* and other genotypes. Future research should address some of the current limitations of the system. Mutations introduced via genome editing might impair regeneration, limiting the recovery of mutant lines in *S. polyrhiza* without the chance to maintain certain alleles in the heterozygous configuration and generate homozygous mutants only upon sexual reproduction. Our successful use of inducible promoters to drive Cas9 expression after regeneration should circumvent this hurdle and reduce the continuous appearance of mutations due to constitutive expression.

Additionally, the origin of somaclonal variation warrants investigation to understand and minimize its confounding effects during phenotypic analysis of transgenic or mutant lines. While genetic causes cannot be excluded, the observed phenotypes - reduced size, slower

growth, and early senescence - are characteristic of *Arabidopsis* regenerated from root- but not from leaf-derived callus due to the inheritance of root-specific epigenetic signatures (Wibowo et al. 2022). Therefore, as most of the calli in our study were derived from roots, it remains possible that the phenotypic variation is of epigenetic origin. Hence, not only the epigenetic profiles of regenerated plants, but also the difference between those regenerated from shoot- (budding pocket) or root-derived calli, deserve future investigation.

Finally, while stable transgenesis and genome editing are powerful tools, transient expression is a versatile technique with multiple applications such as molecular farming or as an experimental system to quickly assess protein localization, function, or interactions (Chen et al. 2013; Krenek et al. 2015). The absence of S-PTGS during transient expression in *S. polyrhiza*, allowing high transgene expression over prolonged periods of time, offers advantages over more traditional systems like *Nicotiana benthamiana*, particularly as a monocot transient expression platform. Given the inability of eudicots to properly splice complex introns, monocot genes cannot be directly expressed from genomic DNA and often require cDNA cloning to function in eudicots (Keith and Chua 1986; Goodall and Filipowicz 1991; Sinibaldi and Mettler 1992; Nitovska et al. 2018; Dombey et al. 2025), which is not always feasible. Moreover, despite recent advances (Li et al. 2025), widespread recalcitrance to *Agrobacterium* in monocots limits its application for transient expression and generally requires the use of viruses or direct DNA delivery to plant cells by complex methods, such as microinjection, biolistics, or electroporation and transfection of protoplasts (Lee et al. 2015; Ren et al. 2021), entailing specialized equipment and skills. Therefore, *S. polyrhiza* also offers a robust, fast, low-cost and technically simple monocot transient expression system.

## Experimental procedures

**Plant material and growth conditions**

*S. polyrhiza* accessions, described in (Wang et al. 2024), were obtained from the Institute of Organismic and Molecular Evolution, Johannes Gutenberg University (Mainz, Germany). *SP162* was deposited at the Landolt-IBBA duckweed collection (Institute of Agriculture Biology and Biotechnology, CNR, Milano, Italy; https://biomemory.cnr.it/collections/CNR-IBBA-MIDW) (Morello et al. 2024), and the Rutgers Duckweed Stock Cooperative (RDSC; Rutgers University, New Jersey, USA; http://www.ruduckweed.org) under ID#: *5676*, and Prof. Huan Weijuan's lab in Shenzhen University of Advanced Technology, China. *Lemna minor 8623* was obtained from the Landolt-IBBA collection.

Plants were grown under sterile conditions on liquid N-medium (Appenroth 2002), SH-medium (*SP162*) [3.2 g/L Schenk and Hildebrandt salts, Duchefa #S0225, pH 5.8] or 0,5X SH-medium (*9509*) (full compositions in **Method S1**). Growth containers included MagentaTM GA-7 boxes (Sigma Aldrich, #V8505), 6-well plates (Greiner Bio-one, #657185), 94 or 145 mm diameter petri dishes (Greiner Bio-one, #633181, #639161), glass beakers covered with a sterile plastic lid or Erlenmeyer flasks, sealed with Parafilm or Leucopore, inside growth chambers under long-day (16-h light / 8-h dark) at 21ºC or 26ºC with a light intensity of 70-125 µmol/m$^2$/s. Plants were sterilized in 1% NaClO (**Method S2**). Turions were induced in N-medium at 21ºC, stored at 4ºC, and germinated at 21/26ºC in SH-medium (**Method S3, S8**).

**Callus induction screening and regeneration test**

Axenic cultures were grown in liquid N-medium with 10 g/L sucrose at 26ºC under long-day (70 µmol/m$^2$/s) for seven days before callus induction. Explants and CIM media were prepared as in (Wang 2016; Huang et al. 2018; Yang et al. 2018; Liu et al. 2019). In most experiments, callus induction rates (number of explants initiating callus formation) were recorded in triplicates. Calli growth rates were expressed as percentage of fresh-weight (FW) biomass gain of 20-50 calli at 7-10-day interval. Frond regeneration was first tested according to (Huang et al. 2018) in the *SP162* and *9509* accessions and later as in (Wang 2016; Yang et al. 2018)

by adding 1 mg/L TDZ or 2.2 mg/L zeatin to the original Frond Regeneration Media by Huang and colleagues and later modified for *SP162* (see below).

***SP162* callus induction and maintenance**

Sterile fronds were transferred dorsoventrally to solid CIM medium [4.4 g/L MS-salts (Duchefa #M0245), 0.5 g/L MES-salts (Duchefa #M1503), 4 g/L Gelrite (Duchefa #G1101), 10 g/L sucrose, 0.1 g/L PVP-40 (Sigma-Aldrich #PVP40-100), 50 mg/L citric acid (Merck #27109-100G-R), 0,221 mg/L 2,4-D (Duchefa #D0911), 0.4 mg/L TDZ (Sigma-Aldrich #P6186-25MG), pH 5.8]. Plates were kept at 21ºC under low light (20 µmol/m$^2$/s) in long-day conditions (see above) and refreshed every two weeks until ~2-3 mm calli were formed at the root tip. Calli were transferred to CIM plates at 26ºC under the same light conditions. Appearing pale-green calli were transferred to separate plates under the same conditions. Sensitivity to antibiotics was performed in 5 to 6 replicates, each composed of a small round plate (60 x 15 mm) containing 3 callus fragments (~3 mm). Sensitivity was tracked by weekly imaging and comparing the developed phenotypes (growth arrest, whitening, and/or browning and death) to calli growing on CIM without antibiotics. Media and step-by-step procedure can be found in **Method S1, S4**.

***SP162* frond regeneration**

Calli fragments (2-3 mm) were placed on FRM [4.4 g/L MS-salts, 0.5 g/L MES-salts, 4 g/L Gelrite, 10 g/L sucrose, 1 mg/L TDZ or 2.2 mg/L zeatin (Sigma-Aldrich #Z0164), pH 5.8] at 26ºC under low light (20 µmol/m$^2$/s) and long-day conditions and refreshed biweekly. Calli with signs of frond regeneration were transferred to liquid SH-medium with 1% w/v sucrose and cultured in Magenta boxes for 7-10 days at 26ºC under high light (125 µM/m$^2$/s) in long-day conditions. Regenerating calli were transferred to 145 mm diameter petri dishes with liquid SH-medium under the same conditions. Emerging fronds were transferred to individual flasks containing SH-media and maintained at 21 or 26ºC. Media and step-by-step procedures are in **Method S1, S5**.

## Callus transformation and selection

*Agrobacterium* (*EHA105*) overnight culture was pelleted (3000 *g*, 20 min), resuspend to $OD_{600}$ 0.7 in 30 ml agroinfection medium [10 mM $MgCl_2$, 50 g/L sucrose, 200 µM acetosyringone (Sigma-Aldrich #D134406)] and incubated 1h in darkness. Fifty 2-3 mm calli were incubated 10 min in the solution under gentle agitation. After 10 min, calli were transferred to solid co-culture medium (solid CIM with 100 µM acetosyringone) and incubated 2 days at 26ºC under low light (20 µmol/m$^2$/s). Excess *Agrobacterium* was rinsed off from calli in liquid SH-medium with 0.3 mg/L cefotaxime (Duchefa #G0124.0025) and 0.25 mg/L ticarcillin (Duchefa #T0190.0010) for 1 min and transferred to agroelimination medium (solid CIM with 0.25 mg/L ticarcillin and 0.3 mg/L cefotaxime). After 2 weeks, calli were transferred to low selection media, solid agroelimination-CIM containing low-dose antibiotic (e.g., 5 mg/L hygromycin, Sigma-Aldrich #400051-1MU). Two weeks later, resistant-looking calli were transferred to media with increased selection pressure. Media were refreshed every two weeks until resistant calli started to proliferate and could be individually transferred to CIM for amplification. Detailed media and step-by-step procedures are in **Method S1, S6**.

## Transient expression in SP162

*Agrobacterium*-mediated transient expression was performed as in (Dombey et al. 2025). 70-80% confluent axenic *SP162* cultures were pretreated in ice-cold 20 mg\L L-glutamine (PanReac Applichem #A1420) for 20 min before being infiltrated twice at -70 kPa twice for 10 min with an induced $OD_{600}$ 0.6 *Agrobacterium* solution in modified agroinfection medium [10 mM $MgCl_2$, 50 g/L sucrose, 150 µM acetosyringone, 0.02% (v/v) Silwet L-77 (Kurt-Obermeier #7060-10)]. Co-cultured was carried in 94 mm diameter petri dishes with SH-medium containing 100 µM acetosyringone for three days in standard growth conditions. After, plants were transferred to SH-medium containing 0.25 mg/L ticarcillin and 0.3 mg/L cefotaxime refreshed weekly. Detailed media and step-by-step procedures are in **Method S1, S7**. Extraction and quantification of bacteria from infiltrated fronds are described in **Method S8**.

**Genome sequencing and annotation.**

DNA sequencing was performed in a PromethION flow cell (v10.4.1 FLO-PRO114M), scaffolded into 20 pseudochromosomes via synteny with the *SP9509* reference genome using Chromosemble in Satsuma_v2 (Grabherr et al. 2010; Ernst et al. 2025). Liftoff v1.6.3 (Shumate and Salzberg 2021), was used to transfer annotations from the *SP9509* genome (Ernst et al. 2025). Further details in **Method S8**.

**Plant imaging and image-based phenotyping**

Images were acquired with a Canon EOS M50 MarkII camera, Andonstar AD409 digital microscope, or Epson V600 scanner. Fluorescent images were taken using a fluorescence stereo zoom microscope Zeiss Axio Zoom V16 or a Sapphire FL Biomolecular scanner (Azure biosystems). Images were analyzed using Ilastik (v1.4.1; https://www.ilastik.org), a supervised machine-learning software (Sommer et al. 2011), suitable for duckweed image analysis (Romano et al. 2022), or Fiji (V2.16.0; https://imagej.net/software/fiji/)(Schindelin et al. 2012). More details about image analysis in **Methods S8**.

**Plasmids generated in this study**

Details on molecular cloning procedures, including sgRNA design, can be found in **Method S8**. All pre-existing used plasmids are publicly available from Addgene (https://www.addgene.org) and listed in **Method S8**. Plasmids newly generated in this study have been deposited in the European Plasmid Repository (https://www.plasmids.eu): *pZmUBQ::2x_BsaI::35St_HygR* (#877); *pZmUBQ::2x_BsaI::35St_KanR* (#878); *pZmUBQ::eGFP* (#869); *pZmUBQ::zCas9i:P2A:GFP* (#870); *pOsU3m::sgRNAscf* (#872); *pOsU6a::sgRNAscf* (#873); *pOsU6aLacZ::sgRNAscf* (#874); *pOsU6b::sgRNAscf* (#875); *pOsU6c::sgRNAscf* (#876); *pZmUBQ::zCas9i:P2A:GFP_4x_sg_ZMET* (#871); *pZmUBQ::FPPFhp* (#879). Plasmids were introduced into *Agrobacterium* through the freeze-thaw method (Weigel and Glazebrook 2006).

**DNA, sRNA, and protein blotting and detection**

DNA, RNA, and protein blotting and detection followed standard procedures described in **Method S8**. Sequences of oligonucleotides used as probes or as PCR primers for Klenow-based DNA probes are listed in **Method S9**. For protein detection, the following antibodies were used. Primary antibodies: monoclonal (rat) anti-GFP [3H9] (Chromotek, #3H9-100), dilution 1:5000; monoclonal (mouse) anti-2A peptide antibody, clone 3H4 (Sigma Aldrich, #MABS2005-25UG), dilution 1:1000. Secondary antibodies: polyclonal (goat) anti-mouse HRP-conjugated (ThermoFisher, #62-6520), dilution 1:10000; polyclonal (goat) anti-rat HRP-conjugated (cell signalling, #7077S), dilution 1:10000.

**qPCR analysis of gene expression**

RNA was extracted from 100 mg frozen ground tissue using TRIzol (#93289, Sigma). One µg of RNA treated with DNase I (Thermo Scientific, #EN0521) was reverse-transcribed with Maxima First-Strand cDNA Synthesis Kit (Thermo Scientific, #K1641). qPCR was performed on a QuantStudio5 (Applied Biosystems) system using KAPA SYBR FAST qPCR Kit (Roche, SFLRAB). Ct values were determined by the 2nd derivative max method of three technical replicates for each biological replicate. Relative expression values were calculated as the ratio of target Ct values relative to reference mRNA (*SpACT*) Ct. Oligonucleotides are listed in **Method S9**.

**Genome size estimation by flow cytometry**

DNA content was estimated by flow cytometry following established methodology for plants (Doležel et al. 2007) and further detailed in **Method S8**. *L. minor 8623* was used as standard with a known DNA content of 1 C = 0,4182 pg (Hoang et al. 2019).

**Statistical analysis**

Statistical tests were performed using GraphPad Prism (V10.2.3; https://www.graphpad.com) or R-dependent ggpubr (v0.6.0) (https://rpkgs.datanovia.com/ggpubr/). Statistical tests performed are indicated accordingly on each figure legend.

# Acknowledgements

We thank present and past members of the Marí-Ordóñez, Hubert and Xu groups for helpful discussions, in particular Daniel Buendía for advice regarding flow cytometry measurements of DNA content and Rodolphe Dombey for help installing and running sgRNA design tools. We are also grateful to the IMP Molecular Biology Service for providing reagents and Sanger sequencing. We thank Dr. Siân Stockton and Dr. Christian Siadjeu for help in isolating HMW DNA of *SP162*. We would also like to thank Andreas Wachter for providing plasmids and sharing the S1 facilities, the nucleic acid core facility from the faculty of biology at JGU Mainz for supporting the genome sequencing. The authors gratefully acknowledge the computing time granted on the supercomputer MOGON 2 at Johannes Gutenberg University Mainz (hpc.uni-mainz.de).

# Conflicts of interest

The authors declare no competing interests.

# Author contributions

AMO, SX, and MHu conceived and designed the study. MHo performed callus induction screens. MHo, RE, and VBB tested regeneration in *9509* and *SP162*. VBB and RE stablished *SP162* culture conditions. ERB, ASL, and VBB tested antibiotic sensitivity. VBB established *SP162* transformation and regeneration after transgenesis and ASL and ERB reproduced it. CD designed sgRNAs and transformed CRISPR/Cas9 vectors. VBB, AMO, and RE characterized regenerated, transgenic, and genome edited lines. AC and SX sequenced and assembled the *SP162* genome. APM performed transient expression experiments. ERB and ASL validated transient expression and tested inducible promoters. LD and LP performed turion experiments. AMO, SX, MHu, VBB, ASL, ERB, and MHo analyzed the data. AMO assembled the figures and wrote the manuscript together with ASL, ERB, VBB, SX, and MHu.

# Funding

This work was supported by core funding of the Gregor Mendel Institute of Plant Molecular Biology of the Austrian Academy of Sciences (GMI-ÖAW) attributed to AMO and from the University of Mainz attributed to SX and MHu. The study was also funded by the Emmy Noether Programme (project#. 512079118), the Stufe-1 funding of University of Mainz, and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) GRK2526 to MHu (Project#. 407023052) and DFG individual grants to SX (Project no. 438887884 and 427577435). LP was supported by the Erasmus+ Student Mobility for Traineeships program (Grant: 14002787004056).

# Data availability statement

All raw source data used in this study can be found in the Zenodo open repository under the following doi: 10.5281/zenodo.16970721. The sequencing data generated have been submitted to NCBI under bio-project number PRJNA1308930. The genome assembly and annotation have been deposited on Figshare: https://figshare.com/articles/dataset/Genome_assembly_and_annotation_of_i_Spirodela_polyrhiza_i_strain_SP162_ID_5676_/29942036.

# References

Abe T, Futsuhara Y. 1986. Genotypic variability for callus formation and plant regeneration in rice (Oryza sativa L.). *Theor Appl Genet* **72**: 3–10.

Acosta K, Appenroth KJ, Borisjuk L, Edelman M, Heinig U, Jansen MAK, Oyama T, Pasaribu B, Schubert I, Sorrels S, et al. 2021. Return of the Lemnaceae: duckweed as a model plant system in the genomics and postgenomics era. *Plant Cell* **33**: 3207–3234.

Antro MV, Prelovsek S, Ivanovic S, Gawehns F, Wagemaker NCAM, Mysara M, Horemans N, Vergeer P, Verhoeven KJF. 2023. DNA methylation in clonal duckweed (Lemna minor L.) lineages reflects current and historical environmental exposures. *Mol Ecol* **32**: 428–443.

Appenroth K -J. 2002. Co-action of temperature and phosphate in inducing turion formation in Spirodela polyrhiza (Great duckweed)*. *Plant, Cell Environ* **25**: 1079–1085.

Appenroth K-J, Sree KS, Böhm V, Hammann S, Vetter W, Leiterer M, Jahreis G. 2017. Nutritional value of duckweeds (Lemnaceae) as human food. *Food Chem* **217**: 266–273.

Appenroth K-J, Ziegler P, Sree KS. 2021. Accumulation of starch in duckweeds (Lemnaceae), potential energy plants. *Physiol Mol Biol Plants* **27**: 2621–2633.

Baggs EL, Tiersma MB, Abramson BW, Michael TP, Krasileva KV. 2022. Characterization of defense responses against bacterial pathogens in duckweeds lacking EDS1. *N Phytol* **236**: 1838–1855.

Bairu MW, Aremu AO, Staden JV. 2011. Somaclonal variation in plants: causes and detection methods. *Plant Growth Regul* **63**: 147–173.

Bewick AJ, Niederhuth CE, Ji L, Rohr NA, Griffin PT, Leebens-Mack J, Schmitz RJ. 2017. The evolution of CHROMOMETHYLASES and gene body DNA methylation in plants. *Genome Biol* **18**: 65.

Böttner L, Dudenhausen F, Nouere S, Malacrinò A, Schäfer M, Koene JM, Huber M, Xu S. 2025. Herbivory can increase plant fitness via reduced interspecific competition—evidence from models and mesocosms. *Proc B* **292**: 20241149.

Cantó-Pastor A, Mollá-Morales A, Ernst E, Dahl W, Zhai J, Yan Y, Meyers BC, Shanklin J, Martienssen R. 2015. Efficient transformation and artificial miRNA gene silencing in Lemna minor. *Plant Biol* **17**: 59–65.

Cardi T, Murovec J, Bakhsh A, Boniecka J, Bruegmann T, Bull SE, Eeckhaut T, Fladung M, Galovic V, Linkiewicz A, et al. 2023. CRISPR/Cas-mediated plant genome editing: outstanding challenges a decade after implementation. *Trends Plant Sci* **28**: 1144–1165.

Chávez A, Schreyer A, Prüsener P, Schäfer M, Xu S, Huber M. 2025. Copper-Induced Transgenerational Plasticity in Plant Defence Boosts Aphid Fitness. *Plant, Cell Environ* **48**: 3997–4010.

Chen Q, Lai H, Hurtado J, Stahnke J, Leuzinger K, Dent M. 2013. Agroinfiltration as an Effective and Scalable Strategy of Gene Delivery for Production of Pharmaceutical Proteins. *Adv Tech Biol Med* **2013**: 103.

Cheng JJ, Stomp A. 2009. Growing Duckweed to Recover Nutrients from Wastewaters and for Production of Fuel Ethanol and Animal Feed. *CLEAN Soil, Air, Water* **37**: 17–26.

Christensen AH, Quail PH. 1996. Ubiquitin promoter-based vectors for high-level expression of selectable and/or screenable marker genes in monocotyledonous plants. *Transgenic Res* **5**: 213–218.

Corneillie S, Storme ND, Acker RV, Fangel JU, Bruyne MD, Rycke RD, Geelen D, Willats WGT, Vanholme B, Boerjan W. 2018. Polyploidy Affects Plant Growth and Alters Cell Wall Composition. *Plant Physiol* **179**: 74–87.

Cui W, Cheng JJ. 2015. Growing duckweed for biofuel production: a review. *Plant Biol* **17**: 16–23.

Debernardi JM, Tricoli DM, Ercoli MF, Hayta S, Ronald P, Palatnik JF, Dubcovsky J. 2020. A GRF–GIF chimeric protein improves the regeneration efficiency of transgenic plants. *Nat Biotechnol* **38**: 1274–1279.

Dermail A, Mitchell M, Foster T, Fakude M, Chen Y-R, Suriharn K, Frei UK, Lübberstedt T. 2024. Haploid identification in maize. *Front Plant Sci* **15**: 1378421.

Doležel J, Greilhuber J, Suda J. 2007. Estimation of nuclear DNA content in plants using flow cytometry. *Nat Protoc* **2**: 2233–2244.

Dombey R, Buendía-Ávila D, Barragán-Borrero V, Diezma-Navas L, Ponce-Mañe A, Vargas-Guerrero JM, Elias R, Marí-Ordóñez A. 2025. Atypical epigenetic and small RNA control of degenerated

transposons and their fragments in clonally reproducing Spirodela polyrhiza. *Genome Res* **35**: 522–544.

Ekperusi AO, Sikoki FD, Nwachukwu EO. 2019. Application of common duckweed (Lemna minor) in phytoremediation of chemicals in the environment: State and future perspective. *Chemosphere* **223**: 285–309.

Ernst E, Abramson B, Acosta K, Hoang PTN, Mateo-Elizalde C, Schubert V, Pasaribu B, Albert PS, Hartwick N, Colt K, et al. 2025. Duckweed genomes and epigenomes underlie triploid hybridization and clonal reproduction. *Curr Biol* **35**: 1828-1847.e9.

Evans DA, Sharp WR. 1986. Applications of Somaclonal Variation. *BioTechnol* **4**: 528–532.

Fourounjian P, Slovin J, Messing J. 2021. Flowering and Seed Production across the Lemnaceae. *Int J Mol Sci* **22**: 2733.

Gasciolli V, Mallory AC, Bartel DP, Vaucheret H. 2005. Partially redundant functions of Arabidopsis DICER-like enzymes and a role for DCL4 in producing trans-acting siRNAs. *CURBIO* **15**: 1494–1500.

Goodall GJ, Filipowicz W. 1991. Different effects of intron nucleotide composition and secondary structure on pre-mRNA splicing in monocot and dicot plants. *EMBO J* **10**: 2635–2644.

Grabherr MG, Russell P, Meyer M, Mauceli E, Alföldi J, Palma FD, Lindblad-Toh K. 2010. Genome-wide synteny through highly sensitive sequence alignment: Satsuma. *Bioinformatics* **26**: 1145–1151.

Greenwood JR, Glaus AN. 2022. Optimized rice transformation protocol for transformation of the blast susceptible Indica rice accession CO39. *CABI Agric Biosci* **3**: 35.

Grützner R, Martin P, Horn C, Mortensen S, Cram EJ, Lee-Parsons CWT, Stuttmann J, Marillonnet S. 2021. High-efficiency genome editing in plants mediated by a Cas9 gene containing multiple introns. *Plant Commun* **2**: 100135.

Hamilton AJ, Voinnet O, Chappell L, Baulcombe DC. 2002. Two classes of short interfering RNA in RNA silencing. *Embo Journal* **21**: 4671–4679.

Harkess A, Bewick AJ, Lu Z, Fourounjian P, Michael TP, Schmitz RJ, Meyers BC. 2024. Unusual predominance of maintenance DNA methylation in Spirodela polyrhiza. *G3: Genes, Genomes, Genet* jkae004. https://doi.org/10.1093/g3journal/jkae004.

Harkess A, McLoughlin F, Bilkey N, Elliott K, Emenecker R, Mattoon E, Miller K, Czymmek K, Vierstra RD, Meyers BC, et al. 2021. Improved Spirodela polyrhiza genome and proteomic analyses reveal a conserved chromosomal structure with high abundance of chloroplastic proteins favoring energy production. *J Exp Bot* **72**: 2491–2500.

He Y, Zhang T, Sun H, Zhan H, Zhao Y. 2020. A reporter for noninvasively monitoring gene expression and plant transformation. *Hortic Res* **7**: 152.

Hillman WS. 1961. The Lemnaceae, or duckweeds. *Bot Rev* **27**: 221–287.

Hoang PNT, Michael TP, Gilbert S, Chu P, Motley ST, Appenroth KJ, Schubert I, Lam E. 2018. Generating a high-confidence reference genome map of the Greater Duckweed by integration of cytogenomic, optical mapping, and Oxford Nanopore technologies. *Plant J* **96**: 670–684.

Hoang PTN, Schubert V, Meister A, Fuchs J, Schubert I. 2019. Variation in genome size, cell and nucleus volume, chromosome number and rDNA loci among duckweeds. *Sci Rep* **9**: 3234.

Huang M, Ma X, Zhong Y, Hu Q, Fu M, Han Y. 2018. Callus induction and plant regeneration of Spirodela polyrhiza. *Plant Cell, Tissue Organ Cult (PCTOC)* **135**: 445–453.

Huber M, Gablenz S, Höfer M. 2021. Transgenerational non-genetic inheritance has fitness costs and benefits under recurring stress in the clonal duckweed Spirodela polyrhiza. *Proc R Soc B* **288**: 20211269.

Ikeuchi M, Favero DS, Sakamoto Y, Iwase A, Coleman D, Rymen B, Sugimoto K. 2019. Molecular Mechanisms of Plant Regeneration. *Annu Rev Plant Biol* **70**: 1–30.

Ikeuchi M, Sugimoto K, Iwase A. 2013. Plant Callus: Mechanisms of Induction and Repression. *Plant Cell* **25**: 3159–3173.

Islam T, Kalkar S, Tinker-Kulberg R, Ignatova T, Josephs EA. 2024. The "Duckweed Dip": Aquatic Spirodela polyrhiza Plants Can Efficiently Uptake Dissolved, DNA-Wrapped Carbon Nanotubes from Their Environment for Transient Gene Expression. *ACS Synth Biol* **13**: 687–691.

Jay F, Brioudes F, Voinnet O. 2023. A contemporary reassessment of the enhanced transient expression system based on the tombusviral silencing suppressor protein P19. *Plant J* **113**: 186–204.

Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. 2012. A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity. *Science* **337**: 816–821.

Jung C, Till B. 2021. Mutagenesis and genome editing in crop improvement: perspectives for the global regulatory landscape. *Trends Plant Sci* **26**: 1258–1269.

Keith B, Chua N-H. 1986. Monocot and dicot pre-mRNAs are processed with different efficiencies in transgenic tobacco. *EMBO J* **5**: 2419–2425.

Kim M, Hyeon DY, Kim K, Hwang D, Lee Y. 2024. Phytohormonal regulation determines the organization pattern of shoot aerenchyma in greater duckweed (Spirodela polyrhiza). *Plant Physiol* **195**: 2694–2711.

Koornneef M. 2002. Classical mutagenesis in higher plants. In *Molecular Plant Biology* (eds. P.M. Gilmartin and C. Bowler), Vol. 1 of, pp. 1–10, Oxford University Press, Oxford, GB.

Kościańska E, Kalantidis K, Wypijewski K, Sadowski J, Tabler M. 2005. Analysis of RNA Silencing in Agroinfiltrated Leaves of Nicotiana Benthamiana and Nicotiana Tabacum. *Plant Mol Biol* **59**: 647–661.

Krenek P, Samajova O, Luptovciak I, Doskocilova A, Komis G, Samaj J. 2015. Transient plant transformation mediated by Agrobacterium tumefaciens: Principles, methods and applications. *Biotechnol Adv* **33**: 1024–1042.

Kuppu S, Tan EH, Nguyen H, Rodgers A, Comai L, Chan SWL, Britt AB. 2015. Point Mutations in Centromeric Histone Induce Post-zygotic Incompatibility and Uniparental Inheritance. *PLoS Genet* **11**: e1005494.

Landolt E. 1986. *Biosystematic investigations in the family of duckweeds (Lemnaceae) (Vol 2). The family of Lemnaceae—a monographic study*. ed. V. des G.I. Der ETH. Veroeffentlichungen des Geobotanischen Instituts der ETH, Stiftung Ruebel, Switzerland.

Lardon R, Wijnker E, Keurentjes J, Geelen D. 2020. The genetic framework of shoot regeneration in Arabidopsis comprises master regulators and conditional fine-tuning factors. *Commun Biol* **3**: 549.

Lee K, Kang M, Ji Q, Grosic S, Wang K. 2023. New T-DNA binary vectors with NptII selection and RUBY reporter for efficient maize transformation and targeted mutagenesis. *Plant Physiol* **192**: 2598–2603.

Lee W-S, Hammond-Kosack KE, Kanyuka K. 2015. Recent Advancements in Gene Expression and Enabling Technologies in Crop Plants. 391–422.

Li E, Geng Y, Khristoforova TR, Wang Y, Jones JW, Demirer GS. 2025. Vacuum and sonication treatment enables efficient transient gene expression in various monocot and eudicot plant seedlings. *bioRxiv* 2025.03.13.642903.

Liu C, Fan H, Zhang J, Wu J, Zhou M, Cao F, Tao G, Zhou X. 2024. Combating browning: mechanisms and management strategies in in vitro culture of economic woody plants. *For Res* **4**: e032.

Liu Y, Wang Y, Xu S, Tang X, Zhao J, Yu C, He G, Xu H, Wang S, Tang Y, et al. 2019. Efficient genetic transformation and CRISPR/Cas9-mediated genome editing in Lemna aequinoctialis. *Plant Biotechnol J* **17**: 2143–2152.

Luo Z, Chen Z. 2007. Improperly terminated, unpolyadenylated mRNA of sense transgenes is targeted by RDR6-mediated RNA silencing in Arabidopsis. *The Plant cell* **19**: 943–958.

Ma X, Liu Y. 2016. CRISPR/Cas9-Based Multiplex Genome Editing in Monocot and Dicot Plants. *Curr Protoc Mol Biol* **115**: 31.6.1-31.6.21.

Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R, Wang B, Yang Z, Li H, Lin Y, et al. 2015. A Robust CRISPR/Cas9 System for Convenient, High-Efficiency Multiplex Genome Editing in Monocot and Dicot Plants. *Mol Plant* **8**: 1274–1284.

Maher MF, Nasti RA, Vollbrecht M, Starker CG, Clark MD, Voytas DF. 2020. Plant gene editing through de novo induction of meristems. *Nat Biotechnol* **38**: 84–89.

Malacrinò A, Böttner L, Nouere S, Huber M, Schäfer M, Xu S. 2024. Induced responses contribute to rapid adaptation of Spirodela polyrhiza to herbivory by Lymnaea stagnalis. *Commun Biol* **7**: 81.

Mérai Z, Kerényi Z, Molnár A, Barta E, Válóczi A, Bisztray G, Havelda Z, Burgyán J, Silhavy D. 2005. Aureusvirus P14 Is an Efficient RNA Silencing Suppressor That Binds Double-Stranded RNAs without Size Specificity. *J Virol* **79**: 7217–7226.

Michael TP, Bryant D, Gutierrez R, Borisjuk N, Chu P, Zhang H, Xia J, Zhou J, Peng H, Baidouri ME, et al. 2017. Comprehensive definition of genome features in Spirodela polyrhiza by high-depth physical mapping and short-read DNA sequencing strategies. *Plant J* **89**: 617–635.

Morello L, Braglia L, A.Iannelli M, Lauria M, Gavazzi F, Gianì S, Perna C, Bucci G, Ianigro M. 2024. IBBA Duckweed Dataset (DUCKWEED-01). *Consiglio Nazionale delle Ricerche, Istituto di Bioscienze e BioRisorse (CNR-IBBR) Occurrence dataset*.

Motte H, Vercauteren A, Depuydt S, Landschoot S, Geelen D, Werbrouck S, Goormachtig S, Vuylsteke M, Vereecke D. 2014. Combining linkage and association mapping identifies RECEPTOR-LIKE PROTEIN KINASE1 as an essential Arabidopsis shoot regeneration gene. *Proc Natl Acad Sci* **111**: 8305–8310.

Neelakandan AK, Wang K. 2012. Recent progress in the understanding of tissue culture-induced genome level changes in plants and potential applications. *Plant Cell Rep* **31**: 597–620.

Nguyen THN, Winkelmann T, Debener T. 2020. Genetic analysis of callus formation in a diversity panel of 96 rose genotypes. *Plant Cell, Tissue Organ Cult (PCTOC)* **142**: 505–517.

Nitovska I, Vasylenko MYu, Morgun B. 2018. The effect of monocot introns on transgene expression in Nicotiana genus plants. *Biotechnol Acta* **11**: 73–83.

Paoli ED, Dorantes-Acosta A, Zhai J, Accerbi M, Jeong D-H, Park S, Meyers BC, Jorgensen RA, Green PJ. 2009. Distinct extremely abundant siRNAs associated with cosuppression in petunia. *RNA* **15**: 1965–1970.

Parent J-S, Jauvion V, Bouché N, Beclin C, Hachet M, Zytnicki M, Vaucheret H. 2015. Post-transcriptional gene silencing triggered by sense transgenes involves uncapped antisense RNA and differs from silencing intentionally triggered by antisense transgenes. *Nucleic Acids Research* **43**: 8464–8475.

Peerbolte R, Ruigrok P, Wullems G, Schilperoort R. 1987. T-DNA rearrangements due to tissue culture: somaclonal variation in crown gall tissues. *Plant Mol Biol* **9**: 51–57.

Peragine A, Yoshikawa M, Wu G, Albrecht HL, Poethig RS. 2004. SGS3 and SGS2/SDE1/RDR6 are required for juvenile development and the production of trans-acting siRNAs in Arabidopsis. *Genes & development* **18**: 2368–2379.

Peterson A, Kishchenko O, Zhou Y, Vasylenko M, Giritch A, Sun J, Borisjuk N, Kuchuk M. 2021. Robust Agrobacterium-Mediated Transient Expression in Two Duckweed Species (Lemnaceae) Directed by Non-replicating, Replicating, and Cell-to-Cell Spreading Vectors. *Front Bioeng Biotechnol* **9**: 5.

Phillips RL, Kaeppler SM, Olhoft P. 1994. Genetic instability of plant tissue cultures: breakdown of normal controls. *Proc Natl Acad Sci* **91**: 5222–5226.

Potrykus I. 1990. Gene Transfer to Cereals: An Assessment. *BioTechnol* **8**: 535–542.

Qu F, Ren T, Morris TJ. 2003. The Coat Protein of Turnip Crinkle Virus Suppresses Posttranscriptional Gene Silencing at an Early Initiation Step. *Journal of Virology* **77**: 511–522.

Ren R, Gao J, Yin D, Li K, Lu C, Ahmad S, Wei Y, Jin J, Zhu G, Yang F. 2021. Highly Efficient Leaf Base Protoplast Isolation and Transient Expression Systems for Orchids and Other Important Monocot Crops. *Front Plant Sci* **12**: 626015.

Risseeuw E, Dijk MEIF, Hooykaas PJJ. 1997. Gene targeting and instability of Agrobacterium T-DNA loci in the plant genome. *Plant J* **11**: 717–728.

Romano LE, Iovane M, Izzo LG, Aronne G. 2022. A Machine-Learning Method to Assess Growth Patterns in Plants of the Family Lemnaceae. *Plants* **11**: 1910.

Sánchez-Gómez C, Posé D, Martín-Pizarro C. 2023. Genome Editing by CRISPR/Cas9 in Polyploids. In *Polyploidy, Methods and Protocols* (ed. Y.V. De Peer), Vol. 2545 of *Methods in Molecular Biology*, pp. 459–473, Humana, New York, NY, USA.

Sangwan RS, Bourgeois Y, Brown S, Vasseur G, Sangwan-Norreel B. 1992. Characterization of competent cells and early events of Agrobacterium-mediated genetic transformation in Arabidopsis thaliana. *Planta* **188**: 439–456.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, et al. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**: 676–682.

Shumate A, Salzberg SL. 2021. Liftoff: accurate mapping of gene annotations. *Bioinformatics* **37**: btaa1016.

Sinibaldi RM, Mettler IJ. 1992. Intron Splicing and Intron-mediated Enhanced Expression in Monocots. *Prog Nucleic Acid Res Mol Biol* **42**: 229–257.

Šmarda P, Klem K, Knápek O, Veselá B, Veselá K, Holub P, Kuchař V, Šilerová A, Horová L, Bureš P. 2023. Growth, physiology, and stomatal parameters of plant polyploids grown under ice age, present-day, and future $CO_2$ concentrations. *N Phytol* **239**: 399–414.

Smith KE, Schäfer M, Lim M, Robles-Zazueta CA, Cowan L, Fisk ID, Xu S, Murchie EH. 2024. Aroma and metabolite profiling in duckweeds: Exploring species and ecotypic variation to enable wider adoption as a food crop. *J Agric Food Res* **18**: 101263.

Sommer C, Straehle C, Kothe U, Hamprecht FA. 2011. ILASTIK: INTERACTIVE LEARNING AND SEGMENTATION TOOLKIT. *2011 IEEE Int Symp Biomed Imaging: Nano Macro* **1**: 230–233.

Stuttmann J, Barthel K, Martin P, Ordon J, Erickson JL, Herr R, Ferik F, Kretschmer C, Berner T, Keilwagen J, et al. 2021. Highly efficient multiplex editing: one-shot generation of 8× Nicotiana benthamiana and 12× Arabidopsis mutants. *Plant J* **106**: 8–22.

Sun Z, Zhao X, Li G, Yang J, Chen Y, Xia M, Hwang I, Hou H. 2023. Metabolic flexibility during a trophic transition reveals the phenotypic plasticity of greater duckweed (Spirodela polyrhiza 7498). *N Phytol* **238**: 1386–1402.

Swinnen G, Lizé E, Sánchez ML, Stolz S, Soyk S. 2025. Application of a GRF-GIF chimera enhances plant regeneration for genome editing in tomato. *Plant Biotechnol J*.

Tabara M, Matsumoto A, Kibayashi Y, Takeda A, Motomura K. 2024. Straightforward and affordable agroinfiltration with RUBY accelerates RNA silencing research. *Plant Mol Biol* **114**: 61.

Tippery NP, Les DH, Appenroth KJ, Sree KS, Crawford DJ, Bog M. 2021. Lemnaceae and Orontiaceae Are Phylogenetically and Morphologically Distinct from Araceae. *Plants* **10**: 2639.

Tse SW, Annese D, Romani F, Guzman-Chavez F, Bonter I, Forestier E, Frangedakis E, Haseloff J. 2024. Optimizing Promoters and Subcellular Localization for Constitutive Transgene Expression in Marchantia polymorpha. *Plant Cell Physiol* **65**: 1298–1309.

Turcotte MM, Kaufmann N, Wagner KL, Zallek TA, Ashman T-L. 2024. Neopolyploidy increases stress tolerance and reduces fitness plasticity across multiple urban pollutants: support for the "general-purpose" genotype hypothesis. *Evol Lett* **8**: 416–426.

Vandeputte W, Coussens G, Aesaert S, Haeghebaert J, Impens L, Karimi M, Debernardi JM, Pauwels L. 2024. Use of GRF-GIF chimeras and a ternary vector system to improve maize (Zea mays L.) transformation frequency. *Plant J* **119**: 2116–2132.

Vaucheret H, Nussaume L, Palauqui JC, Quillere I, Elmayan T. 1997. A Transcriptionally Active State Is Required for Post-Transcriptional Silencing (Cosuppression) of Nitrate Reductase Host Genes and Transgenes. *Plant cell* **9**: 1495–1504.

Voinnet O, Lederer C, Baulcombe DC. 2000. A Viral Movement Protein Prevents Spread of the Gene Silencing Signal in Nicotiana benthamiana. *Cell* **103**: 157–167.

Wang W, Haberer G, Gundlach H, Gläßer C, Nussbaumer T, Luo MC, Lomsadze A, Borodovsky M, Kerstetter RA, Shanklin J, et al. 2014. The Spirodela polyrhiza genome reveals insights into its neotenous reduction fast growth and aquatic lifestyle. *Nat Commun* **5**: 3311.

Wang Y. 2016. Callus induction and frond regeneration in Spirodela polyrhiza. *Czech J Genet Plant Breed* **52**: 114–119.

Wang Y, Duchen P, Chávez A, Sree KS, Appenroth KJ, Zhao H, Höfer M, Huber M, Xu S. 2024. Population genomics and epigenomics of Spirodela polyrhiza provide insights into the evolution of facultative asexuality. *Commun Biol* **7**: 581.

Ware A, Jones DH, Flis P, Chrysanthou E, Smith KE, Kümpers BMC, Yant L, Atkinson JA, Wells DM, Bhosale R, et al. 2023. Loss of ancestral function in duckweed roots is accompanied by progressive anatomical reduction and a re-distribution of nutrient transporters. *Curr Biol* **33**: 1795-1802.e4.

Weigel D, Glazebrook J. 2006. Transformation of Agrobacterium Using the Freeze-Thaw Method. *Cold Spring Harb Protoc* **2006**: pdb.prot4666.

Wibowo AT, Antunez-Sanchez J, Dawson A, Price J, Meehan C, Wrightsman T, Collenberg M, Bezrukov I, Becker C, Benhamed M, et al. 2022. Predictable and stable epimutations induced during clonal plant propagation with embryonic transcription factor. *PLOS Genet* **18**: e1010479.

Yamamoto YT, Rajbhandari N, Lin X, Bergmann BA, Nishimura Y, Stomp A-M. 2001. Genetic transformation of duckweed Lemna gibba and Lemna minor. *Vitr Cell Dev Biol - Plant* **37**: 349–353.

Yang G-L, Feng D, Liu Y-T, Lv S-M, Zheng M-M, Tan A-J. 2021. Research Progress of a Potential Bioreactor: Duckweed. *Biomolecules* **11**: 93.

Yang J, Li G, Hu S, Bishopp A, Heenatigala PPM, Kumar S, Duan P, Yao L, Hou H. 2018. A protocol for efficient callus induction and stable transformation of Spirodela polyrhiza (L.) Schleiden using Agrobacterium tumefaciens. *Aquat Bot* **151**: 80–86.

Yoshikawa M, Peragine A, Park MY, Poethig RS. 2005. A pathway for the biogenesis of trans-acting siRNAs in Arabidopsis. *Genes Development* **19**: 2164–2175.

Zhang K, Su J, Xu M, Zhou Z, Zhu X, Ma X, Hou J, Tan L, Zhu Z, Cai H, et al. 2020. A common wild rice-derived BOC1 allele reduces callus browning in indica rice transformation. *Nat Commun* **11**: 443.

Ziegler P, Appenroth KJ, Sree KS. 2023. Survival Strategies of Duckweeds, the World's Smallest Angiosperms. *Plants* **12**: 2215.