1

**Earliest Evidence of Elephant Butchery at Olduvai Gorge (Tanzania) Reveals the**
**Evolutionary Impact of Early Human Megafaunal Exploitation**

Manuel Domínguez-Rodrigo[1,2,3], Enrique Baquedano[1,4], Abel Moclán[5,1], David Uribelarrea[6],
Alejandro Velázquez-Tello[1,4], José Ángel Correa Cano[1,4], Fernando Diez-Martín[1,7], Elia
Organista[8], Eduardo Mendez-Quintas[1,9], Marina Vegara-Riquelme[1,2], Agness Gidna[11], Audax
Mabulla[12].

[1]Institute of Evolution in Africa (IDEA), University of Alcalá and Archaeological and Paleontological
Museum of Madrid, Madrid, Spain

[2]Area of Prehistory, Department of History and Philosophy, University of Alcalá de Henares, Alcalá de
Henares, Spain.

[3]Department of Anthropology, Rice University, 6100 Main St., Houston, TX 77005-1827, USA

[4]Archaeological and Paleontological Museum of Madrid, Plaza de las Bernardas s/n, Alcalá de Henares,
Madrid, Spain.

[5]PALEVOPRIM laboratory, UMR CNRS 7262, and Université de Poitiers, Poitiers, France.

[6]Department of Geodynamics, Stratigraphy and Paleontology, Complutense University, Madrid, Spain.

[7]Department of Prehistory and Archaeology, University of Valladolid, Valladolid, Spain.

[8]Osteoarkeologiska forskningslaboratoriet och antikens kultur Stockholm Universitet, Stockholm, Sweden.

[9]Universidade de Vigo, MAPAS Lab, 36310, Vigo, Spain.

[10]UNIARQ – Centre for Archaeology. School of Arts and Humanities. University of Lisbon. Alameda da
Universidade, 1600-214, Lisbon, Portugal.

[11]Ngorongoro Conservation Area Authority, Department of Cultural Heritage, 23602 Karatu, Tanzania.

[12]Department of Archaeology, University of Dar es Salaam, 35091 Dar Es Salaam, Tanzania

*The role of megafaunal exploitation in early human evolution remains debated. Occasional use*

*of large carcasses by early hominins has been considered by some as opportunistic, possibly a*

*fallback dietary strategy, and for others a more important survival strategy. At Olduvai Gorge,*

*evidence for megafaunal butchery is scarce in the Oldowan of Bed I, but becomes more*

*frequent and widespread after 1.8 Ma in Bed II, coinciding with the emergence of Acheulean*

*technologies. Here, we present the earliest direct evidence of proboscidean butchery, including*

*a newly documented elephant butchery site (EAK). This shift in behavior is accompanied by*

*larger, more complex occupation sites, signaling a profound ecological and technological*

*transformation. Rather than opportunistic scavenging, these findings suggest a strategic*

*adaptation to megafaunal resources, with implications for early human subsistence and social*

*organization. The ability to systematically exploit large prey represents a unique evolutionary*

*trajectory, with no direct modern analogue, since modern foragers do so only episodically.*

**Introduction**

Spatially concurrent lithic tools and proboscidean bones have a long history in human evolution.
Such concurrency (sometimes in conjunction with other variables) has been frequently
interpreted as the result of hominin butchery (Adrien Hannus, 2018; Agam and Barkai, 2018,
2016; Altamura et al., 2020; Barkai, 2019; Ben-Dor and Barkai, 2020; Berthelet and Chavaillon,
2001; Boschian and Saccà, 2015; Chavaillon et al., 1987; Chavaillon and Berthelet, n.d.;
Delagnes et al., 2006; Gaudzinski et al., 2005; Gaudzinski-Windheuser et al., 2023a, 2023b;
Gomez et al., 1978; Goren-Inbar et al., 1994; Haynes, 2022; Konidaris et al., 2021; Leakey,
1971; Lemorini et al., 2023; Lev and Barkai, 2016; Mussi and Villa, 2008; Reshef and Barkai,
2015; Rocca et al., 2021; Saccà, 2012; Santucci et al., 2016; Solodenko et al., 2015; Villa et al.,
2005; Yravedra et al., 2019, 2010). Spatial concurrency is insufficient to establish a causal
relationship between tools and megafaunal remains. Given the time-averaged nature of a large
part of the Paleolithic record, the deposition of tools and elephants could have occurred
sequentially and unrelatedly in several sites; especially taking into account that most of these
sites were in paleo-floodplains where elephants and hippos were intensively active and formed
natural taphocoenoses. Examples of this abound in the paleobiological record. For example, at

BK (upper Bed II, Olduvai Gorge, Tanzania), some *Pelorovis* carcasses were naturally

deposited on the same alluvial bar and among the fauna processed by hominins, thereby

creating an artificial spatial concurrence between independent depositional events and agencies

(Organista et al., 2016). In the Middle Pleistocene deposits of Torralba and Ambrona (Spain),

artifact-littered areas occur in spatial association with naturally-deposited elephant carcasses,

only a few of which were also exploited to a limited degree by hominins, as demonstrated by a

few proboscidean remains bearing cut marks (Villa, 1990; Villa et al., 2005). Post-

depositional/sedimentation processes also might have artificially created associations of faunal

remains and unrelated lithic artifacts (Villa, 1990; Villa et al., 2005).

Direct taphonomic evidence of hominin exploitation of megafauna (in the form of cutmarked or

dynamically-percussed bone) has been so far the main argument to establish a causal link

between hominin agency and megafaunal remains (Tappen et al., 2022). If restricted to the

Early Pleistocene, most sites containing partial carcasses of megafauna (e.g., elephant,

hippopotamus) either show poor preservation, do not exhibit any of these anthropogenic traces,

or these modifications cannot be securely identified (Delagnes et al., 2006; Domínguez-Rodrigo

et al., 2007; Haynes, 2022, 1991; Isaac and Isaac, 2023; Leakey, 1971). In contrast, for the

Middle and Upper Pleistocene, there is more abundant evidence of hominin butchery of

megafaunal remains, especially of elephants (Gaudzinski-Windheuser et al., 2023a, 2023b;

Haynes, 2022; Yravedra et al., 2010); see review of taphonomic evidence in Agam and Barkai,

(2018), Haynes (2022) and Konidaris et al. (2021). The question that we address here is if such

a behavior has its roots in the early Pleistocene.

Once hominin agency is empirically demonstrated at taphonomically-supported butchery sites, a

second inferential step is attributing carcass butchery to hunting or scavenging. Occasionally,

the occurrence of embedded flint in proboscidean remains has been taken as direct evidence of

predatory behaviors (Agam and Barkai, 2018). This evidence is marginal and chronologically

restricted to the late Upper Pleistocene with composite hafted tools. Recently, a supportive

argument used to detect proboscidean hunting lies on the intensity of cut marks on carcasses

and mortality profiles dominated by male prime adults (Gaudzinski-Windheuser et al., 2023a,

2023b). Prime adult mortality is documented also in other natural palimpsests (Haynes, 1991),

in some cases probably caused by fights leading also to broken tusks (Villa and d'Errico, 2001).

Ethnoarchaeological observation of modern foragers butchering elephants and hippos shows

that they leave very few cut marks after bulk defleshing (Crader, 1983). Modern elephant

butchery after culling events (based on a sample of >500 carcasses) also shows that bulk

defleshing virtually leaves no traces on bones (Haynes and Krasinski, 2021). Only butchery

aiming at the extraction of meat scraps after bulk defleshing imparts most of the cut marks

documented experimentally (Haynes and Krasinski, 2021). This casts doubts on whether

cutmarking intensity relates to hunting, scavenging or a maximization strategy feasible in all

scenarios. To complicate things more, there is a clear mismatch between the butchery intensity

documented at several sites and the resulting tool kit associated with it. Some Middle

Pleistocene elephant carcasses display abundant cut marks, but there are very few or no stone

tools associated with them (Gaudzinski-Windheuser et al., 2023a); whereas at other sites,

elephant carcasses exhibit not a single cut mark but they occur with hundreds or thousands of

associated tools (Chavaillon et al., 1987; Delagnes et al., 2006; Yravedra et al., 2010) (Table 1).

In the latter case, it is probable that in some cases the bulk of tools may have been used to

process other non-megafaunal fauna. In the particular case of the Nadung´a elephant (Kenya,

1-0.7 Ma), that explanation for the presence of lithics is improbable since the only other

mammal carcass remains found are from one bovid and one suid, most likely a background

scatter associated with the bird, fish and reptile remains with which they are spatially

associated. The question remains as to why such a divergent range of tool sets (both in quantity

and typological composition) accompany similar partial megafaunal skeletal remains if they are

truly functionally related to them.

To make interpretations even more challenging, many butchery sites exhibit fairly complete and

(semi)articulated carcasses with limb bones intact. In some of these instances, elements were

dismembered but not demarrowed or degreased (Gaudzinski-Windheuser et al., 2023a).

Defleshing or demarrowing/degreasing do not require disarticulating bones, and the energy

costs involved in cutting through tendons and ligaments of a proboscidean carcass are better

explained as a maximizing strategy aiming at reducing the costs of element transportation

(neither defleshing or demarrowing/degreasing requires disarticulation), and/or further

exploitation of within-bone fat resources. None of this is clearly documented in those sites. Fat

was a crucial resource during human evolution (Ben-Dor and Barkai, 2024 ; Ben-Dor et al.,

2011; Reshef and Barkai, 2015; Solodenko et al., 2015), and the large fat deposits stored in

proboscidean carcasses rationally would have been a target of prehistoric butchery, especially

in strategies of carcass maximization (as documented in the abundant cut mark record at some

megafaunal butchery sites). The question of why our ancestors chose to maximize flesh

extraction from proboscideans, and yet abandoned their fat deposits untouched remains

unanswered. Modern Bisa people (Zambia) butcher elephants and do not exploit their fat, but

they follow a "satisficing" strategy (Haynes and Krasinski, 2021), through bulk defleshing (barely

impacting bone surfaces) (Crader, 1983), and not a "maximizing" strategy, as suggested for

some highly anthropogenically-impacted archaeological proboscidean sites (Gaudzinski-

Windheuser et al., 2023a). It could be argued that the meat of healthy elephants contains

already a high amount of fat and that other anatomical areas (like the podal pads) contain

extremely nutritious fat, but if the strategy was to maximize food resources, long bone medullary

contents should also have been consumed (Ben-Dor et al., 2011; Haynes et al., 2021), as

documented in sites like Castel di Guido (Italy), Preresa (Spain) (Boschian and Saccà, 2015;

Yravedra et al., 2012), or BK (see below). Megafaunal long bone green breakage (linked to

continuous spiral fractures on thick cortical bone) is probably a less ambiguous trace of

butchery than "cut marks", since many of the latter could be equifinal and harder to identify,

especially in contexts of high abrasion and trampling (Haynes et al., 2021, 2020). For example,

the purported cut-marked pelvis from Fuentenueva 3 (Yravedra el al. 2024) exhibits intense

biostratinomic -in addition to diagenetic- modifications impacted by abrasive processes, which

renders the pristine experimental reference collection inadequate to confidently classify it as

human-made. This is further supported by the fact that the experimental trampling collection

used depicts trampling marks as extremely shallow and with wide divergent walls. This does not

include the large diversity of morphologies generated by trampling marks, whose cross-sections

frequently are indistinguishable from stone-tool cut marks (Domínguez-Rodrigo et al., 2010b).

Given the abrasive sedimentary context, the abundance of abrasion marks on the affected

specimens, and the lack of description of microscopic features that could unambiguously be

used to identify those traces as human-made, we remain skeptical of their anthropogenic

nature.

Despite the occurrence of green fractures on naturally-broken bones, such as those trampled by

elephants (Haynes et al., 2020), those occurring through traumatic fracturing or gnawed by

carnivores (Haynes and Hutson, 2020), these fail to reproduce the elongated, extensive, or

helicoidal spiral fractures (uninterrupted by stepped sections), the overlapping conchoidal

cortical scars, the reflected scarring, or the impact hackled flakes typical of dynamic percussive

breakage. Evidence of this type of green breakage had not been documented earlier for the

Early Pleistocene proboscidean or hippopotamid carcasses, beyond the documentation of

flaked bone with the purpose of elaboration of bone tools (Backwell and d'Errico, 2004; Pante et

al., 2020; Sano et al., 2020). Such type of dynamic breaking, in contrast, has been documented

in *Pelorovis* and *Sivatherium* carcasses at the 1.3 Ma site of BK (Olduvai Gorge)(Domínguez-
Rodrigo et al., 2014a; Labrado, 2017; Organista et al., 2016).
173
Regardless of the type of butchery evidence -and with the taphonomic caveat that no
unambiguous evidence exists to confirm that megafaunal carcasses were hunted or scavenged
other than hominins accessed them in different taphonomically-defined stages- stages – the
principal reasons for exploring megafaunal consumption in early human evolution is its origin, its
episodic or temporally-patterned occurrence, its impact on hominin adaptation to certain
landscapes, and its reflection on hominin group size and site functionality. If hominins actively
sought the exploitation of megafauna, especially if targeting early stages of carcass
consumption, the recovery of an apparent surplus of resources reflects a substantially different
behavior from the small-group/small-site pattern documented at several earlier Oldowan
anthropogenic sites (Domínguez-Rodrigo et al., 2019) -or some modern foragers, like the
Hadza, who only exploit megafaunal carcasses very sporadically, mostly upon opportunistic
encounters (Marlowe, 2010; O'Connell et al., 1992; Wood, 2010; Wood and Marlowe, 2013).
When the process started of becoming megafaunal commensal started has major implications
for human evolution.
188
The multiple taphonomic biases intervening in the palimpsestic nature of most of these butchery
sites often prevent the detection of the causal traces linking megafaunal carcasses and
hominins. Functional links have commonly been assumed through the spatial concurrence of
tools and carcass remains; however, this perception may be utterly unjustified as we argued
above. Functional association of both archaeological elements can more securely be detected
through objective spatial statistical methods. This has been argued to be foundational for
heuristic interpretations of proboscidean butchery sites (Giusti, 2021). Such an approach
removes ambiguity and solidifies spatial functional association, as demonstrated at sites like
Marathousa 1 (Konidaris et al., 2018) or TK *Sivatherium (Panera et al., 2019)*. This method will
play a major role in the present study.
199
Here, we present the discovery of a new elephant butchery site (Emiliano Aguirre Korongo,
EAK), dated to 1.78 Ma, from the base of Bed II at Olduvai Gorge. It is the oldest unambiguous
proboscidean butchery site at Olduvai. In combination with its study, we have carried out
extensive survey at a landscape scale on one stratigraphic section of Bed I (Oldowan) and three
stratigraphic sections of Bed II: lower Bed II-Lower Augitic Sands (1.8.1.69 Ma), upper middle

Bed II (from tuff IIB to IID, 1.5 Ma), and upper Bed II (overlying Tuff IID, 1.3 Ma), which are

known to contain Acheulian materials. The goal was to find megafaunal fossils bearing traces of

hominin modification across the landscape. Starting at 1.78 Ma, Olduvai Gorge records

extensive evidence of hominin involvement with megafaunal carcasses (Domínguez-Rodrigo et

al., 2014b, 2009b), namely Proboscidea and Hippopotamidae, followed closely by Giraffidae

and big Bovidae. The nature of this interaction is elaborated and analyzed in this paper. We

have used a spatial approach, combined with a more general taphonomic analysis, and an

analysis of hammerstone-broken bone implements occurring widespread on the three targeted

Bed II stratigraphic intervals. Also provided is a technological study of the lithic artifacts.

**Results**

*The Emiliano Aguirre Korongo (EAK) site*

EAK is located in the area of the confluence of the two gorges, next to geolocality 45a, between

the FLK-N and FLK-NN sites (Hay, 2023) (Fig. 1). Stratigraphically, the archaeological remains

rest on Tuff IF (1.78 M.a), i.e. at the beginning of Bed II (Deino, 2012). As in the whole area of

the gorge, the lowermost Bed II is similar to the uppermost Bed I, which is composed of clays

deposited in the margins of the lake. The drying that characterizes Bed II had not yet taken

place during this moment.  The sedimentary context of the Olduvai basin in this area is

dominated by low energy processes, with predominantly clay deposition, controlled by small

seasonal rises and falls in lake level. The lake has a very high pH and high salinity in the central

zone (Deocampo et al., 2017), but this value is lowered at the margins by the arrival of runoff

water. EAK is located roughly coinciding with the FLK fault zone (Fig. 1). The bone and lithic

remains lie on a thin layer of clay (<5 cm) on top of tuff IF. They are also covered by the

lacustrine clay of the lowermost Bed II. It is therefore the oldest documented archaeological site

formed in Bed II in Olduvai.

During the recent Holocene, the EAK site was affected by a small landslide which displaced the

site 6.25 m vertically and approximately 12 m downslope. Although the stratigraphic sequence

remains clearly recognizable, Tuff I-F, which serves as the paleosurface of the site, is

fragmented into numerous blocks up to 1 m in diameter (Fig. 2). The archaeological deposit has

moved together with the blocks and their original spatial distribution has been altered by this

post-sedimentary deformation (Supplementary Information).

Olduvai Gorge is located in a sedimentary basin bounded to the south by several volcanoes,
and to the north by large outcrops of metamorphic rocks. These two groups are the sources of
raw material (basalts and quartzites respectively) during practically the entire Pleistocene at
Olduvai. Both, the aforementioned basement rocks and the Plio-Quaternary and Quaternary
sedimentary fill are affected by these major east-west faults known as 1$^{st}$, 2$^{nd}$, 3$^{rd}$, 4$^{th}$, KK, FLK
and 5$^{th}$ faults. It should be noted that the FLK fault can only be identified in the landscape
through the deformation of the southern escarpment of the gorge, at the beginning of the gully
that gives its name to this deposit (EAK). It is a well-known fault because it defines the FLK-5
fault block, described stratigraphically by Hay(Hay, 2023). Although it is an escarpment
degraded by the current geodynamics, it affects the most recent aeolian deposits of Olduvai
(Naisiusiu). This geomorphological feature indicates that the FLK fault remained active at least
until the Upper Pleistocene to Holocene. The exact direction of the fault is unknown, although
the stratigraphy suggests that it must be SW-NE (Fig. 1), and its vertical drop is several tens of
meters. The movement of the fault has probably triggered local earthquakes of varying
magnitude. It is well known that earthquakes are one of the main causes of landslides (Keefer,
1984; Ugai et al., 2012; Wasowski et al., 2011), and in this particular case they may have
triggered several landslides affecting deposits from the base of the present gully, i.e. from the
upper part of Bed I to the uppermost Bed (Fig. 2). The fluvial erosion of the stream at the base
of the slopes also destabilized the base of the slopes in this gully, favoring this type of
gravitational process. In the case of EAK, this is a very small landslide, although much larger
landslides have been recorded in the surrounding area, affecting Bed II units such as the Lower
Augitic Sandstone (LAS), which slides over the waxy clay of the lowermost Bed II. Interestingly,
there is a similar archaeological site on the Colorado Plateau in northern New Mexico (USA), in
this case a mammoth, presenting a similar post-depositional history (Rowe et al., 2022).
263
In June 2022, a partial elephant carcass was found at EAK on a fragmented stratigraphic block
situated topographically below FLK North, containing most of the sequence from Tuff ID to the
carbonated root clays overlying Tuff IF that constitute the base of Bed II. The carcass must have
been eroding over the years, since proboscidean bone fragments and lithics had been found
over more than 100 m repeatedly along the seasonally-active stream at the bottom of the gulley.
In January of the same year, we had not documented anything eroding on the same block, but
after the rainy season, several bones emerged, making it obvious that the site contained
remains of a proboscidean carcass (Fig. 3). Excavation started right after the discovery. We
initially set up a 4x2 m trench, followed by an extension consisting of two additional trenches

(Figs. 3). The northern side presented an erosive front, probably responsible for several of the

scattered elephant bones found along the bottom gully. The remaining excavated partial

carcass contained the complete pelvis associated with both hindlimbs (Table 2). These included

both femora and tibiae, plus one set of tarsal elements (including calcaneus, astragalus) and a

fully articulated foot (Figure S1). In addition, several rib fragments belonging to a minimum of 8

ribs, as well as a portion of an ulna were also discovered (Fig. S2). It is interesting to note that

no complete vertebrae were discovered (only one fragment was identified), despite the

presence of a portion of the axial skeleton. The skull was positioned upside down and the

projecting tusks were partially preserved (Fig. 3). In addition to the elephant carcass, a

background scatter of few bones, including a Parmularius $M^2$, a vertebra, rib, tibia and

metatarsal of a small-sized artiodactyl were found.

A total of 80 stone tools, mostly flakes and flake fragments, were spatially distributed in tight

connection with 153 bone remains, pertaining to 46 elements of a proboscidean carcass (Fig. 3;

Table S1). Their preservation is exceptional and the edges remain sharp. The close spatial

association must be functionally linked. Although the cortical bone in several elements was

affected by diagenesis, preventing the preservation of any potential bone surface modifications,

the good preservation in the remaining assemblage did not yield any marks created either by

hominins or other carnivores. On-going restoration work will provide an accurate estimate of

well-preserved and modified fractions of the assemblage. Two elephant bone specimens

showed attributes of green breakage, inferred to be of anthropic origin (See Supplementary

Information). The tibiae were well preserved, but the epiphyseal portions of the femora were

missing, probably removed by carnivores, which would also explain why a large portion of the

rib cage and almost all vertebrae are missing. Further support for this likelihood can be found in

the deletion of bones from three feet, which are one of the richest sources of fat and affected

early in the process of carnivore ravaging (Haynes, 1991, 1988; Haynes and Hutson, 2020).

The absence of the forelimbs could be also explained if carnivores had dragged them away

from the core axial area; however, the fact that the carcass was detached from its original

depositional locus and gravitationally moved (with its overlying sediment) after faulting might

also explain that the missing carcass parts could still be in their original depositional spot. Our

testing of the area where the block was supposed to have been originally formed yielded no

results, mainly because of the thick overburden of Ndutu sediments in the area. The fact that

the carcass was moved while encased in its sedimentary context, along with the close

association of stone tools with the elephant bones, is in agreement with the inference that the

animal was butchered by hominins. A more objective way to assess this association is through
spatial statistical analysis.

*The spatial statistical analyses of EAK*

The Spatial Point Pattern (SPP) of the elephant bones has an intensity of 7.74 specimens per
square meter. However, this intensity varies significantly across the analysed surface, with a
very high-intensity area in the centre of the excavated area and very low intensity surrounding
this zone. This pattern is evident both when analysing the intensity itself and when examining
the hot spot maps (Fig. S24 a-c). The use of MAD and DCLF tests has rejected the null
hypothesis of spatial randomness with the inhomogeneous $K$ and $L$ functions, as well as with
the $K$scaled and $L$scaled functions (p-value = 0.025), while it could not be rejected with the $F$, $G$,
and $J$ functions (p-value = 1). These results align with the use of spatial functions (Fig. S24 d-k),
as the $F$, $G$, and $J$ functions do not reject the random model, whereas the $K$, $L$, $K$scaled, and
$L$scaled functions reject randomness and suggest a clustered SPP. Additionally, the
inhomogeneous $K$ and $L$ functions indicate that this clustering occurs only at short distances
and that at distances greater than 1 meter, the SPP follows a regular pattern. The pair
correlation function also appears to indicate the presence of a clustered pattern at very short
distances. According to the nearest-neighbor cleaning test, most of the faunal assemblage is
highly likely to be part of the cluster (Fig. S24 l-n).

When performing different regression models, we found that the linear regression had an AIC of
-861.53, the quadratic regression -1990.689, the cubic regression -2076.40, and the 4th-degree
polynomial regression -2100.39. However, we used the quadratic regression for the analysis
because the cubic and 4th-degree polynomial regressions were found to be unreliable due to
issues related to sample size and collinearity. Therefore, we performed modelling using the
quadratic regression (Fig. S24 o-r), reinforcing the interpretation of a very high-intensity cluster
in the central part of the excavated area. Furthermore, it is proposed that the accumulation
should not extend significantly beyond the excavated area. This supports that site erosion may
have affected the preserved concentration less than initially suggested.

The analysis of the SPP for lithic remains shows an intensity of 1.67. Although the sample size
is much smaller, it exhibits a pattern similar to that of the bone remains. It also presents intensity
and hot spot maps (Fig. S25 a-c), with very high intensity in the central area of the spatial
window and very low intensity in the surrounding areas. The DCLF and MAD tests yield the

same results as those for the elephant remains. However, the functions (Fig. S25 d-k) show
some differences. The *F, G, J*, and pair correlation functions do not reject the null hypothesis,
while the *K, L, K*scaled, and *L*scaled functions do reject it. In this case, the inhomogeneous
functions suggest the presence of a regular pattern, while the scaled functions indicate a
clustered pattern. Once again, the nearest-neighbor cleaning test (Fig. S25 l-n) suggests that
the lithic remains in the high-intensity area are highly likely to be part of a cluster.

The regression analyses in this case show much higher AIC values compared to the faunal
remains. However, as before, the $4^{th}$-degree polynomial regression has the lowest AIC (-87.99),
followed by the cubic regression (-83.02), the quadratic regression (-71.73), and finally the linear
regression (79.5). However, once again, the cubic and $4^{th}$-degree polynomial regressions
cannot be used due to collinearity and sample size issues, making the quadratic regression the
most appropriate choice. Modelling with quadratic regression (Fig. S25 o-r) further reinforces
the presence of a large cluster and the possible absence of remains outside the analysed
window. Interestingly, the area of maximum intensity does not exactly coincide with that of the
elephant remains but is slightly shifted to the left (west) within the spatial window.

The analysis of the SPP with marks (elephant bones vs. lithic tools) shows that, as expected
due to the imbalance in sample size, there is always a higher probability of finding elephant
remains than lithic remains (Fig. 4 a), with the only exception being at the far left (north) of the
spatial window. The analysis of spatial marks reveals a significant segregative correlation
between fauna and lithics when using the *K*cross and *L*cross functions (Fig. 4 b-c). The pattern
is of regular segregation, meaning that although lithics and bones form part of the same cluster,
they tend to occupy different positions, with lithics clustered in the periphery, as would
correspond to their use and discard upon the processing of the elephant carcass, which
occupies the central part of the cluster.

Finally, analyzing the bones as a covariate has further reinforced the presence of high lithic tool
intensity in the center of the spatial window and the relation between fauna and lithic artifacts
(Fig. 5). First, modelling intensity based on the covariate and the estimated intensity ($\rho$) shows
that intensity is very high when lithic remains are close to elephant remains, but it drops sharply
when moving just 50 cm away. This finding is further supported by the ANOVA test on the
model (p-value = $<2.2\times10^{-16}$) and the results of the Berman $Z_1$ (p-value = $<8.456\times10^{-9}$) and $Z_2$
(p-value = $<1.585\times10^{-30}$) tests. This also shows that the functional connection between the
elephant bones and the tools has been maintained despite the block post-sedimentary
movement. This is further supported by the technological analysis (Supplementary Information).

*Traces of megafaunal exploitation along the paleolandscape at Olduvai Gorge*

<u>*Bed I*</u>

During our survey of the Bed I targeted strata in 2024, no significant traces of megafaunal green bone breakage were observed, with the exception of one locality (Fig. S3). In previous years, we had observed few megafaunal fossils, mostly restricted to hippopotamus scattered remains. They were not particularly abundant, despite the clearly lacustrine and alluvial depositional settings sampled at the junction.

<u>*Bed II*</u>

Out of the three targeted units (lower, intermediate and upper), the lower unit yielded the highest concentration of elephant and hippopotamus remains, followed by the upper unit (Figure S3). The distribution shows a clustered pattern, despite the virtual continuity of most of the sequence in larger areas. The junction area contains the highest density or megafaunal remains for the lower unit. Figure S3 shows the thorough overlap of those elephant and hippopotamus carcasses with areas containing high density of stone artifacts. All the carcasses documented in the lower unit but one (EAK) occur within the Lower Augitic Sands (LAS). This unit contains a landscape-scale concentration of lithic artifacts spreading from HWKE to FLK West (Domínguez-Rodrigo et al., 2023; Uribelarrea et al., 2019, 2017).

The distribution of carcasses in the intermediate unit is less dense and mostly restricted to a small portion of the junction. This small area also shows concentration of lithic remains, but its extent is still unknown since only two localities have been excavated [one to the west of FLK West (Fujioka et al., 2022) and BBS] and no much surface material is found that could be securely attributed to this unit in between both sites. BBS (Bob Brain Site) is a recent site discovered in middle Bed II on the same stratigraphic unit as the site named by Fujioka et al. (2022) as FLK West, recently renamed T69 Complex.  In the side gorge, SHK represents this unit, with well documented presence of megafauna and cut-marked hippopotamus remains (Diez-Martín et al., 2017; Domínguez-Rodrigo et al., 2014b).

The other large cluster of elephant and hippopotamus carcasses is documented in the upper section of the side gorge (Fig S3). There, most of them concentrate around two sites (BK and SC). These sites are also relevant because they contain other smaller megafauna, such as

407 *Sivatherium* and *Pelorovis* in larger numbers (Domínguez-Rodrigo et al., 2014a; Organista et
408 al., 2019, 2017, 2016). It is at these two sites and the landscape surrounding them that the
409 highest densities of lithic artifacts are also found in the upper section of Bed II (Diez-Martín et
410 al., 2009).
411
The clustering of the elephant (and hippopotamus) carcasses in the areas containing the
highest densities of landscape surface artifacts is suggestive of a hominin agency in at least
part of their consumption and modification. The presence of green broken elephant long bone
elements in the area surveyed is only documented within such clusters, both for lower and
upper Bed II. This constitutes inverse negative evidence for natural breaks occurring on those
carcasses through natural (i.e., non-hominin) pre- and peri-mortem limb breaking (Haynes et al.,
2021, 2020; Haynes and Hutson, 2020). In this latter case, it would be expected for green-
broken bones to show a more random landscape distribution, and occur in similar frequencies in
areas with intense hominin landscape use (as documented in high density artifact deposition)
and those with marginal o non-hominin intervention (mostly devoid of anthropogenic lithic
remains).
423
This indicates that from lower Bed II (1.78 Ma) onwards, there is ample documented evidence of
anthropogenic agency in the modification of proboscidean bones across the Olduvai
paleolandscapes. The discovery of EAK constitutes in this respect the oldest evidence thereof
at the gorge. The taphonomic evidence of dynamic proboscidean bone breaking across time
and space supports, therefore, the inferences made by the spatial statistical analyses of bones
and lithics at the site.
430
431 **Discussion**
432
A recent discovery of a couple of hippopotamus partial carcasses at the 3.0-2.6 Ma site of
Nyayanga (Kenya), spatially concurrent with stone artifacts, has been argued to be causally
linked by the presence of cut marks on some bones (Plummer et al., 2023). The only evidence
published  thereof is a series of bone surface modifications on a hippo rib and a tibial crest,
which we suggest may be the result of byproduct of abiotic abrasive processes; the marks
contrast noticeably with the well-defined cut marks found on smaller mammal bones (Plummer
et al.´s 2023: fig. 3C,D) associated with the hippo remains (Plummer et al., 2023). The two
trenches from which both hippo remains derive are located in the stratigraphic unit NY1,

consisting of a channel containing cobble conglomerates and coarse sands in addition to silts.
Gravel and coarse sand can generate the same types of marks seen in the figures of the
Nyayanga hippo bones (Domínguez-Rodrigo et al., 2009a; Manuel Domínguez-Rodrigo et al.,
2010b). Hippopotamidae are the most abundant taxa in the Nyayanga excavations, indicating a
productive taphocoenotic environment, in which hippopotamus remains co-occur with a large
diversity of mammal taxa, most of which could have been naturally deposited and mixed. No
green fractures have been reported on the hippopotamus bones as found in some other
mammal fauna. Although Nyayanga could potentially be one of the earliest examples of
megafaunal butchery, the taphonomic evidence remains ambiguous.
450
The 1.8 Ma *Elephas recki* of FLK North (level 6, Olduvai Bed I) has the potential of being the
oldest proboscidean butchery site (Leakey, 1971). It was argued that cut marks existed on its
complete bones (Bunn, 1982; Delagnes et al., 2006), but a subsequent taphonomic study of
those marks suggested that they might have been made by trampling (Domínguez-Rodrigo et
al., 2007). Additionally, it is unclear if the spatial association of stone tools and elephant bones
is functional or artificial. Leakey (1971) did not report on the third dimension of her excavations
and this is crucial. In our excavations of FLK N6, we documented that the level was a deposit
spanning more than half a meter depth and fossils and stone tools were vertically scattered
throughout (Domínguez-Rodrigo et al., 2010a). It would have been essential to document that
the FLK N6 tools associated with the elephant were either on the same depositional surface as
the elephant bones and/or on the same vertical position.
462
The ambiguity about the FLK N6 elephant renders EAK the oldest secure proboscidean
butchery evidence at Olduvai, and also probably one of the oldest in the early Pleistocene
elsewhere in Africa. FLK North Deinotherium occupied a position above the EAK stratigraphic
level, and evidence of hominin intervention there is missing (Leakey, 1971). A cut-marked bone
fragment of a large animal (probably a hippopotamus) was documented at El-Kherba (Algeria)
dated to 1.78 Ma (Sahnouni et al., 2013). These findings are suggestive of the beginning of
megafaunal exploitation by humans in similar times both in East and North Africa.
470
Our landscape study of the megafaunal remains at Olduvai show that elephants and
hippopotamus overlap on the same clustering areas. In all of them, the geology is indicative of
extensive alluvial zones. At the base of Bed II, the gorge junction is characterized by extensive
wetlands (Ashley, 2003; Ashley et al., 2009; Deocampo et al., 2002; Hay, 2023; Liutkus and

Ashley, 2003). In the overlying LAS, wetlands are crossed by important fluvial systems

(Domínguez-Rodrigo et al., 2023; Uribelarrea et al., 2017). In the middle and upper Bed II fluvial

and wetland systems intertwined as the lake was reduced and moved away from its former

shoreline in the Bed I-Bed II transition (Arráiz et al., 2017; Garrett, 2017; Uribelarrea del Val and

Domínguez-Rodrigo, 2017). In all cases, these wet environments must have been preferred

places for water-dependent megafauna, like elephants and hippos, and their overlapping

ecological niches are reflected in the spatial co-occurrence of their carcasses. Both types of

megafauna show traces of hominin use through either cutmarked or percussed bones, green-

broken bones or both (Supplementary Information).

Green-broken elephant bones are documented at Olduvai at a landscape scale during the LAS

deposition (1.7 Ma)(Diez-Martín et al., 2015; Domínguez-Rodrigo et al., 2023; Uribelarrea et al.,

2017). This record is now added to the newly discovered EAK site, showing tight spatial

association (both vertically and horizontally) between bones and stone tools. This trend of

spatial association of stone artifacts and proboscidean bones continues through the deposition

of Bed II. Although no cut marks have been documented on the proboscidean bones that we

identified (part of which had cortical surfaces impacted by biostratinomic and diagenetic

modification), the presence of green-broken bones at the site, the butchery of carcasses as

indicated by spatial statistical analyses at sites like EAK and TK (Domínguez-Rodrigo et al.,

2014a; Organista et al., 2016; Panera et al., 2019), and the presence of hammerstone-broken

bones on the surrounding paleo-landscapes of these and other sites, support the occasional

exploitation of megafaunal remains by humans. In most green-broken limb bones, we document

the presence of a medullary cavity, despite the continuous presence of trabecular bone tissue

on its walls. We do not know if the bone breakage was intended to obtain marrow or to create

raw material for making artifacts or tools with functional purposes or both. We have been able to

identify only one potential bone tool in our sample (Figures 6, S4-S23); the scarcity of bone

implements supports the former option. The exploitation of megafauna is not anecdotal, since at

sites like BK or TK (upper Bed II), the taphonomic analysis of other megafaunal remains

(namely, *Sivatherium* and *Pelorovis*) clearly shows intensive exploitation of complete carcasses

in both of these taxa, especially at BK (Domínguez-Rodrigo et al., 2014a; Organista et al., 2016;

Panera et al., 2019). The food surplus generated by the exploitation of some of these animals

would have been a significant advantage for the survival and adaptation of the hominin groups.

These results  of widespread conspicuous evidence of hammerstone-breaking behaviors during

Bed II partially confirm the recent report on bone artefacts discovered in Bed II by another

research team (de la Torre et al., 2025). Excavations at T69 Complex area unearthed a wealth

of megafaunal (proboscidean and hippopotamus) remains displaying hammerstone green

breakage and hominin modifications. These remains seem to be spatially connected with a

highly productive paleolandscape, since at its northern end it connects with BBS, a recently

discovered site with megafaunal (Hippopotamus) remains. Green breakage at BBS also reflects

hominin intervention in the palustrine environment congregating so many megafaunal

carcasses. This unit would be equivalent to our second unit in Bed II.

We are not confident that the artefacts reported by dela Torre et al are indeed tools. They are

similar to the green-broken specimens that we have reported here. They are characterized by

repeated number of flake scars along the edges; however, this is not a diagnostic feature, since

multiple scars along edges also indirectly result from the repeated battering until the final

breakage of the bone during butchery (Haynes et al., 2021, 2020). In large bovids, it can also be

derived from the gnawing of durophagous carnivores (Villa and Bartram, 1996) (Fig. S31).

Carnivores capable of breaking elephant bones and producing the same break shapes go back

to at least the early Middle Miocene (Fig S32). Another argument used by the authors is that the

tools are longer and bear more flake scars compared to other faunal fragments (approximate

average 13 versus 4). Again, this is documented among durophagous carnivores (Villa and

Bartram, 1996). A higher scar number in the T69 artifact collection could result from a

correlation of different variables (i.e.,  longer dimension of the compared specimens, or different

carcass sizes) and be, therefore, unrelated to anthropic intentionality. A third argument provided

is that the items share consistent technological features, such as the production of

morphologically similar, elongated, pointed, and notched bone tools, suggesting a patterned

behavior. Again, this is equifinal with unintentional breaking of megafaunal remains during

butchery (Haynes et al., 2021, 2020). There are several arguments that could be used to

support this alternative:

1.  The scars exhibit an asymmetric distribution, characterized by stepped or reflected,

incomplete fractures predominantly on the medullary surface, with minimal invasiveness.

This contrasts with the expected conchoidal and complete scarring distributed equally

across both cortical (i.e., periosteal) and medullary surfaces, as would be anticipated in

cases of intentional flaking. Given that bone is more easily flaked than metamorphic

rock, and that bifacial knapping of stone typically produces symmetrical, conchoidal

scars on both surfaces, this would also be expected in the T69 bone specimens.

2.  This bilateral symmetry principle is observed in Middle Pleistocene bone handaxes,

where intentional modification is inferred based on the equal treatment of both sides and

evidence of artificial shaping (Zutovski and Barkai, 2016). It is also documented on the

similarly aged 1.4 Ma bone handaxe from Konso (Sano et al., 2020), and a purported

handaxe from FC (Bed II, Olduvai) (Backwell and d'Errico, 2004), but it is absent from

the reported items in T69, as well as those of the present study. Why, otherwise, would

scars on the medullary surface be mostly limited to the wall of the breakage plane?

3.  The repeated presence of incomplete, stepped scars on the medullary surfaces of the

artifacts found at T69 aligns more closely with damage resulting from deliberate

battering associated with breakage rather than subsequent edge retouching. The latter

process would be expected to generate more controlled and complete conchoidal scars

indicative of intentional shaping (Backwell and d'Errico, 2004; Haynes et al., 2021,

2020). This reflected scarring is also documented in Bos/Bison bones broken by hyenas,

caused by redundancy in gnawing and the resulting overlapping micro-cracking leading

to final collapse(Villa and Bartram, 1996) (Fig. S31).

4.  Scar frequency is not a taphonomically valid criterion to attribute agency, since elephant

bones can be broken in multiple ways, thus impacting experimentally-derived scar

patterns. Sequential battering of initial crack lines with handheld hammerstones can

generate overlapping notches and scars. This is not necessarily reproduced when

experimental reproductions use alternative methods (such as throwing or bashing

against immobile anvils) (Backwell and d'Errico, 2004).

5.  In several of the artifacts from T69, most of the overlapping scarring occurs in

association to notches (and associated percussion marks) derived from dynamic

loading. This reinforces that they may have been resulted from repeated battering during

bone breaking.

6.  The T69 Complex bone artifacts and those reported here are large-sized. If their

potential function was for heavy-duty activities, those should have left any visible traces

(at the macro and microscales) on the scarred edges. These are absent. A faunal

assemblage that has good preservation of cut and percussion marks should certainly

have good preservation conditions for microwear use and macro-damage scars.

7.  If applying the same criteria defining bone tools at T69 Complex to Bos/Bison modified-

bones, hyenas would also create similar artifacts (Fig. S31).

A final argument used by the authors to justify the intentional artifactual nature of their bone
implements is that the  bone tools were found *in situ* within a single stratigraphic horizon
securely dated to 1.5 million years ago, indicating systematic production rather than episodic
use. This is taphonomically unjustified. In a productive paleosurface where megafauna is
abundant and hominin intervention is intense, the presence of this type of highly-scarred green-
broken bones is fairly common, since they result from butchery and carnivore post-depositional
ravaging (Domínguez-Rodrigo et al., 2014a, 2014b, 2009b; Villa and Bartram, 1996).Here, we
have shown that most green-broken elephant bones cluster in three main areas of the Olduvai
gorge.
585
Leakey collected more than a hundred purported bone tools from the same locations of Bed II
where we collected our hammerstone-broken bones (Leakey, 1971) (Fig S33). All of them
derive from Bed II (Backwell and d'Errico, 2004). A recent taphonomic review cautiously
identified six potential bone tools in that collection (Pante et al., 2020). Some of these authors,
using the much smaller T69 Complex artifact sample, now argue about systematic bone tool
production. Bone artifacts (i.e., hominin-made items) that are used as tools should display
evidence thereof through the macroscopic scarring, the polishing of the actively used surface,
and the presence of microabrasive modifications on it (Backwell and d'Errico, 2008, 2001;
d'Errico and Backwell, 2009; LeMoine, 1994). None of that has been documented in the T69
assemblage. We do not deny that the items reported by de la Torre et al. (2025) have the
potential of having been used as tools, but proof is currently missing. The same applies to our
reported sample (Fig 6; S4-S25). Only one artifact shows modifications that are potentially
derived from its use as a tool. If confirmed, it is interesting to note the convergence in the point
shape targeted in such an implement and those intentionally shaped in the earliest Acheulian
implements documented at Olduvai Gorge (Figs 6, S6-S8). If such a bone implement was a tool,
it would be the oldest bone tool documented to date (>1.7 Ma).
602
603
604    **Conclusions**
605
Evidence documents that the oldest systematic anthropogenic exploitation of proboscidean
carcasses are documented (at several paleolandscape scales) in the Middle Pleistocene sites of
Neumark-Nord (Germany)(Gaudzinski-Windheuser et al., 2023a, 2023b). Evidence of at least
episodic access to proboscidean remains goes back in time (see review in Agam and Barkai,

2018; Ben-Dor et al., 2011; Haynes, 2022). Redundant megafaunal exploitation is well

documented at some early Pleistocene sites from Olduvai Gorge (Domínguez-Rodrigo et al.,

2014a, 2014b; Organista et al., 2019, 2017, 2016). At the very same sites, the stone artifactual

assemblages, as well as the site dimensions, are substantially larger than those documented in

the Bed I Oldowan sites (Diez-Martín et al., 2024, 2017, 2014, 2009). Whether this reflects

larger hominin groups or more prolonged and time-averaged intervention in more productive

environments remains to be determined (it will likely vary according to site). Here, we have

reported a significant change in hominin foraging behaviors during Bed I and Bed II times,

roughly coinciding with the replacement of Oldowan industries by Acheulian tool kits -although

during Bed II, both industries co-existed for a substantial amount of time (Domínguez-Rodrigo et

al., 2023; Uribelarrea et al., 2019, 2017). The evidence presented here, together with that

documented by de la Torre et al. (2025), represents the most geographically extensive

documentation of repeated access to proboscidean and other megafaunal remains at a single

fossil locality. The transition from Oldowan sites, where lithic and archaeofaunal assemblages

are typically concentrated within 30–40 m² clusters, to Acheulean sites that span hundreds or

even over 1000 m² (as in BK), with distinct internal spatial organization and redundancy in

space use across multiple archaeological layers spanning meters of stratigraphic sequence

(Domínguez-Rodrigo et al., 2014a, 2009b; Organista et al., 2017), reflects significant behavioral

and technological shifts. This pattern likely signifies critical innovations in human evolution,

coinciding with major anatomical and physiological transformations in early hominins (Dembitzer

et al., 2022; Domínguez-Rodrigo et al., 2021, 2012).

**Methods**

*Excavation of EAK*

The systematic excavation of the stratigraphic layers involved a small crew. Materials exceeding

2 cm in size were meticulously plotted using laser theodolites (TOPCON total stations). Each

trench was subjected to stereo-photography upon exposure, enabling the creation of

photogrammetric 3D reconstructions of the paleosurface and deposited materials prior to

recovery. The orientation and inclination of the artifacts were recorded using a compass and an

inclinometer, respectively. Sediments removed from the trenches were sieved using 5 mm and

3 mm meshes.

The photogrammetric reconstructions of each trench were subsequently georeferenced and the

resulting orthophotos were imported into AutoCAD, facilitating the detailed mapping of

archaeological materials. The decision to conduct photogrammetry by individual trenches was

made to minimize the exposure time of the excavated materials. Prolonged exposure could

accelerate desiccation, potentially causing the most fragile specimens to become brittle and

fragmentary. To mitigate this risk, excavation was limited to small, manageable areas, with

small trenches proving ideal for efficient excavation due to the limited overburden. Each trench

was excavated until the entire archaeological horizon was exposed, followed by comprehensive

photogrammetric documentation prior to 3D plotting. Throughout the excavation, professional

restoration experts provided necessary treatment for the bones. This was crucial for the

recovery and identification of many bone fragments.

*Restoration of the EAK elephant bones*

The main interventions were carried out in two locations: in situ, and in the field laboratory at the

Emiliano Aguirre Research Station. For the consolidation and extraction of remains at the site,

interventions were performed using Paraloid B-72 diluted in acetone at different concentrations,

applied exclusively to fossils exhibiting powdering. This methodology strengthened the bones

prior to extraction, ensuring better handling and reducing the risk of fractures. The use of

Paraloid B-72 is a standard practice in the conservation of osseous remains due to its stability

and compatibility with the material, though its limited penetration in some bone types has been

noted(Daura et al., 2017). During this process, bandaging with sterile gauze impregnated with

Paraloid B-72 was selectively applied to necessary areas to further protect fragile sections.

Additionally, superficial bandaging was conducted on larger fossils that visibly presented

potential fractures and retained their cortical surface, aiming to prevent deterioration during the

extraction process.

The greatest challenge was extracting long bones due to their size, weight, and fragility. These

bones exhibited a greater presence of cortical surface, although they were not completely

preserved, as they were fragmented but still retained significant areas suitable for study.

Another significant challenge was the extraction of tusk remains, as the ivory exhibited a high

degree of degradation and fragmentation. To recover these, multiple applications of consolidant

were performed, allowing certain tusk blocks to be extracted in a controlled manner.

Larger fossils were extracted in block to minimize the risk of structural damage. The extraction

process involved:

1. Initial consolidation by applying Paraloid B-72 dropwise to the surface, along with
bandaging with sterile gauze in necessary areas to provide additional protection.
2. Surface protection with a sacrificial layer, avoiding direct contact with other reinforcing
materials.
3. Creation of a counter-mold using plaster bandages, reinforced with interwoven steel rods
to provide rigidity and stability.
4. Insertion of metal spatulas beneath the remains to lift and safely transport them.

In the field laboratory, cleaning of the bone remains was carried out, along with adhesion of

fragments and their consolidation when necessary. The primary priority was preparing the

remains for study by the rest of the team, opting for a purely conservative intervention,

prioritizing detailed cleaning of some fragments for identification. Additionally, reassembly of

larger fragments was performed, using Paraloid B-72 diluted in acetone at a higher

concentration to ensure effective adhesion. For bone remains, mechanical and chemical

cleaning procedures were applied depending on the nature of the adhered deposits.

This restoration procedure enabled the preservation of many bone fragments. Cleaning them

properly will require much more time. In their current status, several specimens still have some

adhering sedimentary matrix. This has prevented us from thoroughly examining all bone

surfaces looking for BSM or conducting comprehensive identification. Some elements and

fragments remain unidentified given also the lack of a proper comparative proboscidean

collection. Despite this, most of the assemblage has been successfully studied.

*Lithic analysis of EAK*

A total of 80 lithic specimens have been recovered from EAK. The technological analysis of

these materials follows established classification criteria and methodological frameworks(Diez-

Martín et al., 2009), categorizing artifacts into the following groups:

(a) Unmodified Specimens – Naturally occurring cobblestones lacking evidence of
anthropogenic modification.

(b) Percussion Tools – Includes hammerstones or cobbles exhibiting diagnostic battering,
pitting, and/or impact scars consistent with percussive activities.

(c) Cores – Handheld cores are classified based on key technological attributes: (i) the number
of reduction surfaces (unifacial, bifacial, multifacial), (ii) the number of striking platforms
(unipolar, bipolar, multipolar), and (iii) the spatial organization of knapping sequences (linear,
opposed, orthogonal, centripetal). Bipolar cores are identified based on diagnostic features
associated with anvil-supported percussion.

(d) Detached Products – This category encompasses both complete and fragmented flakes,
derived from either bipolar or freehand reduction. Technological attributes recorded for
detached products include core reduction stage (assessed through the presence and extent of
cortical coverage on dorsal surfaces and striking platforms), platform morphology, and dorsal
scar patterning.

(e) Knapping By-Products – Includes all non-diagnostic lithic debris resulting from core reduction
and flake detachment. This category incorporates non-diagnostic flake fragments with evidence
of direct handheld percussion, shatter, blocky/angular undetermined detached fragments, core-
like fragments, and debris ≤20 mm in size.

727
728　*The spatial statistical analyses of EAK*
729
The spatial point patterns (SPP) of faunal and lithic remains have been analyzed using the
'spatstat' library(Baddeley et al., 2015) in R (www.r-project.org). This methodological approach
is of great interest for paleontological and archaeological contexts, as it allows for the analysis
of aspects such as the degree of randomness of the SPP, potential relationships between
different types of materials, or enabling predictions about unexcavated areas. We conducted the
analysis in three different ways after selecting the spatial window, i.e. the analysed excavated
area (52.56 m$^2$).

First, we analysed the SPPs without marks (i.e., raw SPPs, without considering any specific
spatial variable) for the elephant remains and lithic remains independently. In these initial cases,
we created intensity maps, hot spot maps, and hot spot maps with a 99% Confidence Interval
(C.I.). Next, we applied a chi-squared test with four different grid configurations (5×5, 8×8,
10×10, 12×12)(Moclán et al., 2023) to determine whether the SPPs are homogeneous or
inhomogeneous. These analyses showed that, in all cases, the SPPs can be considered
inhomogeneous (p-value <0.05). Therefore, we proceeded to use tests to analyse the type of
SPP. We employed the Pair Correlation, *K, L, F, G*, and *J* functions in their inhomogeneous
versions, as well as the *Kscaled* and *Lscaled* functions to analyse the typology of the SPPs (i.e.,
random, cluster, or regular). Both versions of the functions were used because the sample size
does not allow us to analyse the type of inhomogeneity through a studentized permutation test.
Similarly, we used the DCLF (Diggle-Cressie-Loosmore-Ford) and MAD (Maximum Absolute
Deviation) tests with the *K, L, F, G, J, Kscaled*, and *Lscaled* functions to check the typology of
the point patterns (i.e., random vs. non-random). Finally, we used the 'nnclean'
function(Baddeley et al., 2015) to detect possible clustering of materials through nearest
neighbour clutter removal. Lastly, we performed spatial modelling using linear, quadratic, cubic,
and $4^{th}$-degree polynomial regressions to delve deeper into the characteristics of the SPPs. The
models were evaluated using the Akaike Information Criterion (AIC) index, and subsequently,
we compared their validity through ANOVA. Finally, we selected the best-performing model (i.e.,
the one with the lowest AIC values and that does not show issues of collinearity or suffer from
the small sample size) and generated four maps: 1) estimated intensity of the point process
based on the fitted model, 2) simulation of a new point pattern using the fitted model with the
Metropolis-Hastings algorithm, 3) a difference map subtracting the intensity of the simulated
new point pattern from the "real" intensity of the spatial point pattern(Domínguez-Rodrigo et al.,
2024), and 4) an extrapolated estimation of potential points within a larger spatial window
(308.23 $m^2$) than the original one.

The second analysis consists of examining the degree of statistical correlation between two
marks, i.e., spatial variables, in this case, fauna vs. lithic tools. To do this, we first calculated the
relative risk to assess the likelihood of each type of material appearing in the analysed window.
Then, we applied a chi-square test to verify once again, using the same grid configuration,
whether the samples are homogeneous or inhomogeneous. The chi-square test again indicates
that the spatial distribution is inhomogeneous. Therefore, we used the *K·*, *L·*, *K*cross, and
*L*cross inhomogeneous functions to analyse whether there is spatial correlation between the two
material types and, if so, what kind it is.

Finally, given the size disparity between the elephant remains and the lithics, we analysed the
relationship between both types of materials by considering the location of the elephant remains
as a covariate. First, a distance map was generated to the centroids of the bone remains.
ANOVA was then conducted to assess the fitted effect of the covariate in the SPPs. This was
followed by the application of the ρ function ('rhohat' function) to investigate the influence of the
distance to the bones. This test is similar to the previous one but provides a more detailed view
of the potential variations in the intensity of remains with respect to distance from the bones.
Finally, a $Z_1$ and $Z_2$ Berman–Lawson–Waller test was performed.

*Megafaunal exploitation along the paleolandscape sequence of Olduvai Gorge*

An intensive survey seeking stratigraphically-associated megafaunal bones was carried out in
the months of June 2023 and 2024. We targeted the Bed I and Bed II exposures of the main
Gorge (From FLK NNN to the KK fault), and the side gorge, from the junction until BK. Linearly,
this comprises over 6 km of outcrops. We targeted this area because it contains the highest
density of fossils of all the gorge. We focused on proboscidean bones and used hippopotamus
bones, some of the most abundant in the megafaunal fossils, as a spatial control. Bones were
counted as separate carcasses when enough distance existed in between findings (i.e.,
localities). Stratigraphic association was carried out by direct observation of the geological
context and with the presence of a Quaternary geologist during the whole survey. When fossils
found were ambiguously associated with specific strata, these were excluded from the present
analysis.

For the sake of analysis, and given that we were not focusing on time-resolution landscapes, we
used four time intervals: Bed I, Bed II lower (from Tuff IF to LAS), Bed II intermediate (from Tuff
IIB to Tuff IIC, and Bed II upper (from Tuff IID to the interface with Bed III). Only in one case,
Bed II lower, were most of the remains circumscribed to a single stratigraphic unit (Lower
Augitic Sands [LAS], which mark the beginning of Hay´s middle Bed II). The goals of this survey
were: a) collect a spatial sample of proboscidean and megafaunal bones enabling us to
understand if carcasses on the Olduvai paleolandscapes were randomly deposited or
associated to specific habitats; b) given the water dependence of elephants, we wanted to
understand if they showed the same type of ecological association to water-dependent
megafauna, like hippopotamus, as their modern counterparts (using taphocoenoses as
indicators of past megafaunal biocoenosis); c) approach potential anthropogenic agency in

proboscidean exploitation by comparing carcass dispersal and accumulation and productivity of
anthropogenic landscapes (i.e., concentrations and dispersals of lithic artifacts); d) seek
anthropogenic traces of proboscidean exploitation in the form of cut and percussion marks and
green bone breakage; e) associate this latter point with the hominin use of space, by
determining if these taphonomic evidences occurred randomly or clustered and spatially
associated with stone tools.

**Acknowledgements**

We thank the Spanish Ministry of Science and Innovation for funding this research (PID2023-
146260NB-C2), and the Spanish Ministry of Culture for their funding through the program of
Archaeology Abroad. We thank the Commission for Science and Technology (COSTECH), the
Ngorongoro Conservation Area Authorities (NCAA), the Division of Antiquities, and the
Tanzanian Ministry of Natural Resources and Tourism for their permission to conduct research
in Tanzania. AM is funded by a postdoctoral contract grant from the Fyssen Foundation. We are
grateful to the Tanzanian co-workers at Olduvai Gorge.

**References**

Adrien Hannus L. 2018. Clovis Mammoth Butchery: The Lange/Ferguson Site and Associated
    Bone Tool Technology. Texas A&M University Press.
Agam A, Barkai R. 2018. Elephant and Mammoth Hunting during the Paleolithic: A Review of
    the Relevant Archaeological, Ethnographic and Ethno-Historical Records. *Quaternary* **1**:3.
Agam A, Barkai R. 2016. Not the brain alone: The nutritional potential of elephant heads in
    Paleolithic sites. *Quat Int* **406**:218–226.
Altamura F, Gaudzinski-Windheuser S, Melis RT, Mussi M. 2020. Reassessing Hominin Skills at
    an Early Middle Pleistocene Hippo Butchery Site: Gombore II-2 (Melka Kunture, Upper
    Awash valley, Ethiopia). *Journal of Paleolithic Archaeology* **3**:1–32.
A matter of fat: Hunting preferences affected Pleistocene megafaunal extinctions and human
    evolution. n.d.
Arráiz H, Barboni D, Uribelarrea D, Mabulla A, Baquedano E, Domínguez-Rodrigo M. 2017.
    Paleovegetation changes accompanying the evolution of a riverine system at the BK
    paleoanthropological site (Upper Bed II, Olduvai Gorge, Tanzania). *Palaeogeogr
    Palaeoclimatol Palaeoecol* **488**:84–92.
Ashley G. 2003. Tracks, Trails and Trampling by Large Vertebrates in a Rift Valley Paleo-
    Wetland, Lowermost Bed II, Olduvai Gorge, Tanzania. *Ichnos* **9**:23–32.
Ashley GM, Tactikos JC, Owen RB. 2009. Hominin use of springs and wetlands: Paleoclimate
    and archaeological records from Olduvai Gorge (~1.79–1.74 Ma). *Palaeogeogr
    Palaeoclimatol Palaeoecol* **272**:1–16.
Backwell L, d'Errico F. 2008. Early hominid bone tools from Drimolen, South Africa. *J Archaeol*

*Sci* **35**:2880–2894.

Backwell L, d'Errico F. 2004. The first use of bone tools: a reappraisal of the evidence from
Olduvai Gorge, Tanzania. *Palaeontologia africana* **40**:95–158.

Backwell LR, d'Errico F. 2001. Evidence of termite foraging by Swartkrans early hominids. *Proc*
*Natl Acad Sci U S A* **98**:1358–1363.

Baddeley A, Rubak E, Turner R. 2015. Spatial Point Patterns, Chapman & Hall/CRC
Interdisciplinary Statistics. Oakville, MO: Apple Academic Press.

Barkai R. 2019. When Elephants Roamed Asia: The Significance of Proboscideans in Diet,
Culture and Cosmology in Paleolithic Asia In: Kowner R, Bar-Oz G, Biran M, Shahar M,
Shelach-Lavi G, editors. Animals and Human Society in Asia: Historical, Cultural and
Ethical Perspectives. Cham: Springer International Publishing. pp. 33–62.

Ben-Dor, M., Barkai, R. 2024. A matter of fact: hunting preferences affected Pleistocene
megafaunal extinctions and human evolution. Quaternary Science Reviews, 331.
https://doi.org/10.1016/j.quascirev.2024.108660.

Ben-Dor M, Barkai R. 2020. The importance of large prey animals during the Pleistocene and
the implications of their extinction on the use of dietary ethnographic analogies. *Journal of*
*Anthropological Archaeology* **59**:101192.

Ben-Dor M, Gopher A, Hershkovitz I, Barkai R. 2011. Man the fat hunter: the demise of Homo
erectus and the emergence of a new hominin lineage in the Middle Pleistocene (ca. 400
kyr) Levant. *PLoS One* **6**:e28689.

Berthelet A, Chavaillon J. 2001. The early Palaeolithic butchery site of Barogali (Republic of
Djibouti)The World of Elephants. International Congress Proceedings, Consiglio Nazionale
Delle Ricerche, Rome. pp. 176–179.

Boschian G, Saccà D. 2015. In the elephant, everything is good: Carcass use and re-use at
Castel di Guido (Italy). *Quat Int* **361**:288–296.

Bunn HT. 1982. Meat-eating and Human Evolution: Studies on the Diet and Subsistence
Patterns of Plio-Pleistocene Hominids in East Africa. University of California, Berkeley.

Chavaillon, J., Berthelet, A., 2004. The archaeological sites of Melka Kunture. In: Chavaillon, J.,
Piperno, M. (Eds.), Studies on the Early Paleolithic Site of Melka Kunture, Ethiopia. Istituto
Italiano di Preistoria e Protostoria, Roma, pp. 25–80.

Chavaillon J, Boisaubert J-L, Faure M, Guerin C, Ma J-L, Nickel B. 1987. Le site de dépeçage
pléistocène à Elephas recki de Barogali (République de Djibouti): nouveaux résultats et
datation. *C R Acad Sci II* **305**:1259–1266.

Crader DC. 1983. Recent single-carcass bone scatters and the problem of "butchery" sites in
the archaeological record. *Animal and Archaeology 1 Hunters and their Prey, BAR*
*International Series* **163**:107–141.

Daura J, Sanz M, Arsuaga JL, Hoffmann DL, Quam RM, Ortega MC, Santos E, Gómez S,
Rubio A, Villaescusa L, Souto P, Mauricio J, Rodrigues F, Ferreira A, Godinho P, Trinkaus
E, Zilhão J. 2017. New Middle Pleistocene hominin cranium from Gruta da Aroeira
(Portugal). *Proc Natl Acad Sci U S A* **114**:3397–3402.

Deino AL. 2012. (40)Ar/(39)Ar dating of Bed I, Olduvai Gorge, Tanzania, and the chronology of
early Pleistocene climate change. *J Hum Evol* **63**:251–273.

Delagnes A, Lenoble A, Harmand S, Brugal J-P, Prat S, Tiercelin J-J, Roche H. 2006.
Interpreting pachyderm single carcass sites in the African Lower and Early Middle
Pleistocene record: A multidisciplinary approach to the site of Nadung'a 4 (Kenya). *J*
*Anthropol Archaeol* **25**:448–465.

de la Torre I, Doyon L, Benito-Calvo A, Mora R, Mwakyoma I, Njau JK, Peters RF,
Theodoropoulou A, d'Errico F. 2025. Systematic bone tool production at 1.5 million years
ago. *Nature*. doi:10.1038/s41586-025-08652-5

Dembitzer J, Barkai R, Ben-Dor M, Meiri S. 2022. Levantine overkill: 1.5 million years of hunting
down the body size distribution. *Quat Sci Rev* **276**:107316.

Deocampo DM, Berry PA, Beverly EJ, Ashley GM, Jarrett RE. 2017. Whole-rock geochemistry
tracks precessional control of Pleistocene lake salinity at Olduvai Gorge, Tanzania: A
record of authigenic clays. *Geology* **45**:683–686.
Deocampo DM, Blumenschine RJ, Ashley GM. 2002. Wetland Diagenesis and Traces of Early
Hominids, Olduvai Gorge, Tanzania. *Quat Res* **57**:271–281.
d'Errico F, Backwell L. 2009. Assessing the function of early hominin bone tools. *J Archaeol Sci*
**36**:1764–1773.
Diez-Martín F, Fraile C, Uribelarrea D, Sánchez-Yustos P, Domínguez-Rodrigo M, Duque J,
Díaz I, De Francisco S, Yravedra J, Mabulla A, Baquedano E. 2017. SHKExtension: a new
archaeological window in theSHKfluvial landscape of Middle BedII(Olduvai Gorge,
Tanzania). *Boreas* **46**:831–859.
Diez-Martín F, Fraile-Márquez C, Duque-Martínez J, Sánchez-Yustos P, de Francisco S,
Baquedano E, Mabulla A, Domínguez-Rodrigo M. 2024. On time scales and "synchronic"
variability in the archaeology of human origins: short-term technological variations at SHK
(Olduvai Gorge, Tanzania). *Archaeol Anthropol Sci* **16**. doi:10.1007/s12520-024-02092-4
Diez-Martín F, Sánchez P, Domínguez-Rodrigo M, Mabulla A, Barba R. 2009. Were Olduvai
Hominins making butchering tools or battering tools? Analysis of a recently excavated lithic
assemblage from BK (Bed II, Olduvai Gorge, Tanzania). *Journal of Anthropological
Archaeology* **28**:274–289.
Diez-Martín F, Sánchez Yustos P, Uribelarrea D, Baquedano E, Mark DF, Mabulla A, Fraile C,
Duque J, Díaz I, Pérez-González A, Yravedra J, Egeland CP, Organista E, Domínguez-
Rodrigo M. 2015. The origin of the Acheulean: The 1.7 million-year-old site of FLK West,
Olduvai Gorge (Tanzania). *Sci Rep* **5**:17839.
Diez-Martín F, Sánchez-Yustos P, Uribelarrea D, Domínguez-Rodrigo M, Fraile-Márquez C,
Obregón R-A, Díaz-Muñoz I, Mabulla A, Baquedano E, Pérez-González A, Bunn HT. 2014.
New archaeological and geological research at SHK main site (Bed II, Olduvai Gorge,
Tanzania). *Quat Int* **322-323**:107–128.
Domínguez-Rodrigo M, Baquedano E, Organista E, Cobo-Sánchez L, Mabulla A, Maskara V,
Gidna A, Pizarro-Monzo M, Aramendi J, Galán AB, Cifuentes-Alcobendas G, Vegara-
Riquelme M, Jiménez-García B, Abellán N, Barba R, Uribelarrea D, Martín-Perea D, Diez-
Martin F, Maíllo-Fernández JM, Rodríguez-Hidalgo A, Courtenay L, Mora R, Maté-
González MA, González-Aguilera D. 2021. Early Pleistocene faunivorous hominins were
not kleptoparasitic, and this impacted the evolution of human anatomy and socio-ecology.
*Sci Rep* **11**:16135.
Domínguez-Rodrigo M, Barba R, Egeland CP. 2007. Deconstructing Olduvai: A Taphonomic
Study of the Bed I Sites. Springer Science & Business Media.
Domínguez-Rodrigo M, Bunn HT, Mabulla AZP, Baquedano E, Uribelarrea D, Pérez-González
A, Gidna A, Yravedra J, Díez-Martín F, Egeland CP, Others. 2014a. On meat eating and
human evolution: A taphonomic analysis of BK4b (Upper Bed II, Olduvai Gorge, Tanzania),
and its bearing on hominin megafaunal consumption. *Quat Int* **322**:129–152.
Domínguez-Rodrigo M, Cobo-Sánchez L, Aramendi J, Gidna A. 2019. The meta-group social
network of early humans: A temporal–spatial assessment of group size at FLK Zinj (Olduvai
Gorge, Tanzania). *J Hum Evol* **127**:54–66.
Domínguez-Rodrigo M, Cobo-Sánchez L, Baquedano E, Mabulla A, Gidna A, Diez-Martin F.
2024. Reconstructing Olduvai: The Behavior of Early Humans at David's Site. Elsevier
Science.
Domínguez-Rodrigo M, de Juana S, Galán AB, Rodríguez M. 2009a. A new protocol to
differentiate trampling marks from butchery cut marks. *J Archaeol Sci* **36**:2643–2654.
Domínguez-Rodrigo M, Diez-Martín F, Yravedra J, Barba R, Mabulla A, Baquedano E,
Uribelarrea D, Sánchez P, Eren MI. 2014b. Study of the SHK Main Site faunal assemblage,
Olduvai Gorge, Tanzania: Implications for Bed II taphonomy, paleoecology, and hominin

utilization of megafauna. *Quat Int* **322-323**:153–166.

Domínguez-Rodrigo M, Mabulla A, Bunn HT, Barba R, Diez-Martín F, Egeland CP, Espílez E,
Egeland A, Yravedra J, Sánchez P. 2009b. Unraveling hominin behavior at another
anthropogenic site from Olduvai Gorge (Tanzania): new archaeological and taphonomic
research at BK, Upper Bed II. *J Hum Evol* **57**:260–283.

Domínguez-Rodrigo M, Mabulla AZP, Bunn HT, Diez-Martin F, Baquedano E, Barboni D, Barba
R, Domínguez-Solera S, Sánchez P, Ashley GM, Yravedra J. 2010. Disentangling hominin
and carnivore activities near a spring at FLK North (Olduvai Gorge, Tanzania). *Quat Res*
**74**:363–375.

Domínguez-Rodrigo M, Pickering TR, Bunn HT. 2010b. Configurational approach to identifying
the earliest hominin butchers. *Proceedings of the National Academy of Sciences*
**107**:20929–20934.

Domínguez-Rodrigo M, Pickering TR, Diez-Martín F, Mabulla A, Musiba C, Trancho G,
Baquedano E, Bunn HT, Barboni D, Santonja M, Uribelarrea D, Ashley GM, Martínez-Ávila
M del S, Barba R, Gidna A, Yravedra J, Arriaza C. 2012. Earliest porotic hyperostosis on a
1.5-million-year-old hominin, olduvai gorge, Tanzania. *PLoS One* **7**:e46414.

Domínguez-Rodrigo M, Uribelarrea D, Diez-Martín F, A, Mabulla, Gidna A, Cobo-Sánchez L,
Martín-Perea DM, Organista E, Barba R, Baquedano E. 2023. Earliest Acheulian
paleolandscape reveals a 1.7 million-year-old megasite at Olduvai Gorge (Tanzania). *Quat*
*Sci Rev* **316**:108262.

Fujioka T, Benito-Calvo A, Mora R, McHenry L, Njau JK, de la Torre I. 2022. Direct cosmogenic
nuclide isochron burial dating of early Acheulian stone tools at the T69 Complex (FLK
West, Olduvai Bed II, Tanzania). *J Hum Evol* **165**:103155.

Garrett K. 2017. Multi-proxy reconstuction of a ~1.3 Ma freshwater wetland, Olduvai Gorge,
Tanzania. doi:10.7282/T3T43WJ0

Gaudzinski S, Turner E, Anzidei AP, Àlvarez-Fernández E, Arroyo-Cabrales J, Cinq-Mars J,
Dobosi VT, Hannus A, Johnson E, Münzel SC, Scheer A, Villa P. 2005. The use of
Proboscidean remains in every-day Palaeolithic life. *Quat Int* **126-128**:179–194.

Gaudzinski-Windheuser S, Kindler L, MacDonald K, Roebroeks W. 2023a. Hunting and
processing of straight-tusked elephants 125.000 years ago: Implications for Neanderthal
behavior. *Sci Adv* **9**:eadd8186.

Gaudzinski-Windheuser S, Kindler L, Roebroeks W. 2023b. Widespread evidence for elephant
exploitation by Last Interglacial Neanderthals on the North European plain. *Proc Natl Acad*
*Sci U S A* **120**:e2309427120.

Giusti D. 2021. Investigating the spatio-temporal dimension of past human-elephant
interactions: a spatial taphonomic approach.

Gomez MS, Lopez N, Gonzalez AP. 1978. Acheulean occupation sites in the Jarama valley
(Madrid, Spain). *Curr Anthropol* **19**:394–395.

Goren-Inbar N, Lister A, Werker E, Chech M. 1994. A Butchered elephant skull and associated
artifacts from the Acheulian site of Gesher Benot Ya'Aqov, Israel. *Paléorient* **20**:99–112.

Haynes G. 2022. Late Quaternary Proboscidean Sites in Africa and Eurasia with Possible or
Probable Evidence for Hominin Involvement. *Quaternary* **5**:18.

Haynes G. 1991. Mammoths, Mastodonts, and Elephants: Biology, Behavior and the Fossil
Record. Cambridge University Press.

Haynes G. 1988. Longitudinal studies of african elephant death and bone deposits. *J Archaeol*
*Sci* **15**:131–157.

Haynes G, Hutson J. 2020. African elephant bones modified by carnivores: Implications for
interpreting fossil proboscidean assemblages. *Journal of Archaeological Science: Reports*
**34**:102596.

Haynes G, Krasinski K. 2021. Butchering marks on bones of Loxodonta africana (African
savanna elephant): Implications for interpreting marks on fossil proboscidean bones.

*Journal of Archaeological Science: Reports* **37**:102957.
Haynes G, Krasinski K, Wojtal P. 2021. A Study of Fractured Proboscidean Bones in Recent
and Fossil Assemblages. *Journal of Archaeological Method and Theory* **28**:956–1025.
Haynes G, Krasinski K, Wojtal P. 2020. Elephant bone breakage and surface marks made by
trampling elephants: Implications for interpretations of marked and broken Mammuthus
spp. bones. *Journal of Archaeological Science: Reports* **33**:102491.
Hay R. 2023. Geology of the Olduvai Gorge. Univ of California Press.
Isaac GL, Isaac B. 2023. Koobi Fora Research Project: Plio-Pleistocene archaeology.
Clarendon.
Keefer DK. 1984. Landslides caused by earthquakes. *GSA Bulletin* **95**:406–421.
Konidaris GE, Athanassiou A, Tourloukis V, Thompson N, Giusti D, Panagopoulou E, Harvati K.
2018. The skeleton of a straight-tusked elephant (Palaeoloxodon antiquus) and other large
mammals from the Middle Pleistocene butchering locality Marathousa 1 (Megalopolis
Basin, Greece): preliminary results. *Quat Int* **497**:65–84.
Konidaris GE, Barkai R, Turlukēs B, Harvati K. 2021. Human-elephant Interactions: From Past
to Present. Tübingen University Press.
Labrado EO. 2017. Estudio tafonómico de los niveles arqueológicos de BK (Bell Korongo),
garganta de Olduvai, Tanzania.
Leakey MD. 1971. Olduvai Gorge: Volume 3, Excavations in Beds I and II, 1960-1963.
Cambridge University Press.
LeMoine GM. 1994. Use wear on bone and antler tools from the Mackenzie Delta, Northwest
Territories. *Am Antiq* **59**:316–334.
Lemorini C, Santucci E, Caricola I, Nucara A, Nunziante-Cesaro S. 2023. Life around the
elephant in space and time: An integrated approach to study the human-elephant
interactions at the Late Lower Paleolithic site of La Polledrara di Cecanibbio (Rome, Italy). *J*
*Archaeol Method Theory* **30**:1233–1281.
Lev M 'ayan, Barkai R. 2016. Elephants are people, people are elephants: Human–
proboscideans similarities as a case for cross cultural animal humanization in recent and
Paleolithic times. *Quat Int* **406**:239–245.
Liutkus CM, Ashley GM. 2003. Facies Model of a Semiarid Freshwater Wetland, Olduvai Gorge,
Tanzania. *J Sediment Res* **73**:691–705.
Marlowe F. 2010. The Hadza: Hunter-gatherers of Tanzania. University of California Press.
Moclán A, Cobo-Sánchez L, Domínguez-Rodrigo M, Méndez-Quintas E, Rubio-Jara S, Panera
J, Pérez-González A, Santonja M. 2023. Spatial analysis of an Early Middle Palaeolithic
kill/butchering site: the case of the Cuesta de la Bajada (Teruel, Spain). *Archaeol Anthropol*
*Sci* **15**. doi:10.1007/s12520-023-01792-7
Mussi M, Villa P. 2008. Single carcass of Mammuthus primigenius with lithic artifacts in the
Upper Pleistocene of northern Italy. *J Archaeol Sci* **35**:2606–2613.
O'Connell JF, Hawkes K, Blurton-Jones NG. 1992. Patterns in the distribution, site structure and
assemblage composition of Hadza kill-butchering sites. *J Archaeol Sci* **19**:319–345.
Organista E, Arriaza MC, Barba R, Gidna A, Ortega MC, Uribelarrea D, Mabulla A, Baquedano
E, Domínguez-Rodrigo M. 2019. Taphonomic analysis of the level 3b fauna at BK, Olduvai
Gorge. *Quat Int* **526**:116–128.
Organista E, Domínguez-Rodrigo M, Egeland CP, Uribelarrea D, Mabulla A, Baquedano E.
2016. Did Homo erectus kill a Pelorovis herd at BK (Olduvai Gorge)? A taphonomic study
of BK5. *Archaeol Anthropol Sci* **8**:601–624.
Organista E, Domínguez-Rodrigo M, Yravedra J, Uribelarrea D, Arriaza MC, Ortega MC,
Mabulla A, Gidna A, Baquedano E. 2017. Biotic and abiotic processes affecting the
formation of BK Level 4c (Bed II, Olduvai Gorge) and their bearing on hominin behavior at
the site. *Palaeogeogr Palaeoclimatol Palaeoecol* **488**:59–75.
Panera J, Rubio-Jara S, Domínguez-Rodrigo M, Yravedra J, Méndez-Quintas E, Pérez-

González A, Bello-Alonso P, Moclán A, Baquedano E, Santonja M. 2019. Assessing functionality during the early Acheulean in level TKSF at Thiongo Korongo site (Olduvai Gorge, Tanzania). *Quat Int* **526**:77–98.

Pante M, Torre I de la, d'Errico F, Njau J, Blumenschine R. 2020. Bone tools from Beds II–IV, Olduvai Gorge, Tanzania, and implications for the origins and evolution of bone technology. *J Hum Evol* **148**:102885.

Plummer TW, Oliver JS, Finestone EM, Ditchfield PW, Bishop LC, Blumenthal SA, Lemorini C, Caricola I, Bailey SE, Herries AIR, Parkinson JA, Whitfield E, Hertel F, Kinyanjui RN, Vincent TH, Li Y, Louys J, Frost SR, Braun DR, Reeves JS, Early EDG, Onyango B, Lamela-Lopez R, Forrest FL, He H, Lane TP, Frouin M, Nomade S, Wilson EP, Bartilol SK, Rotich NK, Potts R. 2023. Expanded geographic distribution and dietary strategies of the earliest Oldowan hominins and Paranthropus. *Science* **379**:561–566.

Reshef H, Barkai R. 2015. A taste of an elephant: The probable role of elephant meat in Paleolithic diet preferences. *Quat Int* **379**:28–34.

Rocca R, Boschin F, Aureli D. 2021. Around an elephant carcass: Cimitero di Atella and Ficoncella in the behavioural variability during the Early Middle Pleistocene in italy.

Rowe TB, Stafford TW, Fisher DC, Enghild JJ, Quigg JM, Ketcham RA, Sagebiel JC, Hanna R, Colbert MW. 2022. Human Occupation of the North American Colorado Plateau ~37,000 Years Ago. *Frontiers in Ecology and Evolution* **10**. doi:10.3389/fevo.2022.903795

Saccà D. 2012. Taphonomy of Palaeloxodon antiquus at Castel di Guido (Rome, Italy): Proboscidean carcass exploitation in the Lower Palaeolithic. *Quat Int* **276-277**:27–41.

Sahnouni M, Rosell J, van der Made J, Vergès JM, Ollé A, Kandi N, Harichane Z, Derradji A, Medig M. 2013. The first evidence of cut marks and usewear traces from the Plio-Pleistocene locality of El-Kherba (Ain Hanech), Algeria: implications for early hominin subsistence activities circa 1.8 Ma. *J Hum Evol* **64**:137–150.

Sano K, Beyene Y, Katoh S, Koyabu D, Endo H, Sasaki T, Asfaw B, Suwa G. 2020. A 1.4-million-year-old bone handaxe from Konso, Ethiopia, shows advanced tool technology in the early Acheulean. *Proc Natl Acad Sci U S A* **117**:18393–18400.

Santucci E, Marano F, Cerilli E, Fiore I, Lemorini C, Palombo MR, Anzidei AP, Bulgarelli GM. 2016. Palaeoloxodon exploitation at the Middle Pleistocene site of La Polledrara di Cecanibbio (Rome, Italy). *Quat Int* **406**:169–182.

Solodenko N, Zupancich A, Cesaro SN, Marder O, Lemorini C, Barkai R. 2015. Fat residue and use-wear found on Acheulian biface and scraper associated with butchered elephant remains at the site of Revadim, Israel. *PLoS One* **10**:e0118572.

Tappen M, Bukhsianidze M, Ferring R, Coil R, Lordkipanidze D. 2022. Life and death at Dmanisi, Georgia: Taphonomic signals from the fossil mammals. *J Hum Evol* **171**:103249.

Ugai K, Yagi H, Wakai A. 2012. Earthquake-Induced Landslides: Proceedings of the International Symposium on Earthquake-Induced Landslides, Kiryu, Japan, 2012. Springer Science & Business Media.

Uribelarrea del Val D, Domínguez-Rodrigo M. 2017. Geoarchaeology in a meandering river: A study of the BK site (1.35 Ma), Upper Bed II, Olduvai Gorge (Tanzania). *Palaeogeogr Palaeoclimatol Palaeoecol* **488**:76–83.

Uribelarrea D, Martín-Perea D, Díez-Martín F, Sánchez-Yustos P, Dominguez-Rodrigo M, Baquedano E, Mabulla A. 2017. A reconstruction of the paleolandscape during the earliest Acheulian of FLK West: The co-existence of Oldowan and Acheulian industries during lowermost Bed II (Olduvai Gorge, Tanzania). *Palaeogeogr Palaeoclimatol Palaeoecol* **488**:50–58.

Uribelarrea D, Perea DM, Díez-Martín F. 2019. A geoarchaeological reassessment of the co-occurrence of the oldest Acheulean and Oldowan in a fluvial ecotone from lower middle Bed II (1.7 ma) at …. *Quaternary*.

Villa P. 1990. Torralba and Aridos: elephant exploitation in Middle Pleistocene Spain. *J Hum*

*Evol* **19**:299–309.

Villa P, Bartram L. 1996. Flaked bone from a hyena den. *Paléo* **8**:143–159.

Villa P, d'Errico F. 2001. Bone and ivory points in the Lower and Middle Paleolithic of Europe. *J Hum Evol* **41**:69–112.

Villa P, Soto E, Santonja M, Pérez-González A, Mora R, Parcerisas J, Sesé C. 2005. New data from Ambrona: closing the hunting versus scavenging debate. *Quat Int* **126-128**:223–250.

Wasowski J, Keefer DK, Lee C-T. 2011. Toward the next generation of research on earthquake-induced landslides: Current issues and future challenges. *Eng Geol* **122**:1–8.

Wood BM. 2010. Household and Kin Provisioning by Hadza Males. Harvard University.

Wood BM, Marlowe FW. 2013. Household and kin provisioning by Hadza men. *Hum Nat* **24**:280–317.

Yravedra J, Domínguez-Rodrigo M, Santonja M, Pérez-González A, Panera J, Rubio-Jara S, Baquedano E. 2010. Cut marks on the Middle Pleistocene elephant carcass of Áridos 2 (Madrid, Spain). *J Archaeol Sci* **37**:2469–2476.

Yravedra J, Rubio-Jara S, Panera J, Martos JA. 2019. Hominins and Proboscideans in the Lower and Middle Palaeolithic in the Central Iberian Peninsula. *Quat Int* **520**:140–156.

Yravedra J, Rubio-Jara S, Panera J, Uribelarrea D, Pérez-González A. 2012. Elephants and subsistence. Evidence of the human exploitation of extremely large mammal bones from the Middle Palaeolithic site of PRERESA (Madrid, Spain). *J Archaeol Sci* **39**:1063–1071.

Yravedra, J., Courtenay, L., Gutiérrez-Rodríguez, M., Reinoso-Gordo, M., Saarinen, J., Égüez, N., Luzón, C., Rodríguez-Alba, J., Solano, J., Titton, S., Montilla-Jiménez, E., Cámara-Donoso, E., Herranz-Rodrigo, D., Estaca, V., Serrano-Ramos, A., Amorós, G., Azanza, B., Bocherens,H., DeMiguel,D., Fagoaga, A., García-Alix,A.,González-Quiñones,J., Jiménez-Espejo, F., Kaakinen, H., Munuera,M., Ochando, J., Piñero, P., Sánchez-Bandera, C., Viranta, S., Fortelius, M.,Agustí,J., Hugues-Alexandre Blain, Carrión, J., Barsky, D., Oms, O., Mallol,C., Jiménez-Arenas,J.M. 2024. Not seen before. Unveiling depositional context and Mammuthus meridionalis exploitation at Fuente Nueva 3 (Orce, southern Iberia) through taphonomy and microstratigraphy. Quaternary Science Reviews,Volume 329, https://doi.org/10.1016/j.quascirev.2024.108561.

Zutovski K, Barkai R. 2016. The use of elephant bones for making Acheulian handaxes: A fresh look at old bones. *Quat Int* **406**:227–238.

**FIGURE CAPTIONS**

Figure 1. Location of EAK in the junction of the main and secondary branches of Olduvai Gorge, main faults and Naibor Soit quartzites (A), within the area where the Bed I sites cluster at the junction (B), and the specific EAK locus with 1 m contour lines (C).

Figure 2. A, Location of EAK on the south side of the Main Gorge (view to the east). B, Stratigraphic section of the gully (Korongo) showing the correlation between the different marker

tuffs of Bed I on either side. The landslide displaced the boundary between Bed I and Bed II,

including the following sequence: waxy claystone, tuff 1F, waxy claystone. Vertical displacement

is 6.25 m. C, detail of the stratigraphy of the site, showing how the archaeological level rests

directly on Tuff 1F and was covered by the clay of the lowermost Bed II. The landslide has

created vertical cracks that separate both the Tuff 1F and the archaeological level itself. The

archaeological remains have moved along with the tuff blocks. The original clay of the

lowermost layer II protected the fossils from erosion after the landslide. An incipient soil formed

on this surface and was subsequently buried by colluvium. At present, erosion is acting on the

part closest to the stream, affecting some of the fossils.

1153

1154 Figure 3. Photogrammetry (left) and planimetry (right) of the EAK site in 2022 and 2023. The

1155 site-wide distribution of materials, lithics (blue) and bones (red), topographic profiles, and the 3D

1156 terrain model are shown.

1157

1158 Figure 4. Relative risk (i.e., probability of occurrence) of bones and lithic artifacts at EAK, with

1159 second-order functions showing regular correlation between both types of items.

1160

1161 Figure 5. Correlation between lithics and bones at EAK (using the elephant bones as a

1162 covariate)  with the ρ ('rhohat') function and a Z 1 and Z 2Berman–Lawson–Waller.

1163

Figure 6. Left: Intentional shaping of points in a quartzite LCT from FLK West, and on a

proboscidean femur shaft. Arrows indicate conchoidal scars probably caused by use (given their

location away from the edge) and their reflection and stepped morphology by the pointed area.

Notice the polishing at the point, which contrasts with the unpolished state of the remainder of

the artifact. More complete cortical and medullary views of this artifact can be seen in Figures

S6-S8. Right: Comparison of the configuration strategies identified on the quartz LCT and the

EAK proboscidean femur shaft. A. LCT on a quartz slab from FLK West, Level 6; B. The

proboscidean femoral shaft from EAK; C. Diacritical diagram showing point conceptualization

sequencing identified in the previous specimens: 1. First thinning work (oblique to the distal end

in the quartz LCT and vertical in the bone shaft); 2. Right-sided retouch; 3. Left-sided retouch.

Series 2 and 3 determine the final shaping of the distal tip.

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187    Table 1. African Early Pleistocene archaeological sites containing potential evidence of hominin

1188    involvement.

1189

| | Area excavated (m2) | Nº bones | Nº lithics | Megafaunal taxon | Other non megafauna | Age (Ma) | Green breaks | Cut marks |
|---|---|---|---|---|---|---|---|---|
| Nyayanga 3 (Kenya) | 49 | 1580* (241 Hippo) | 42** | Hippo | Present | 3-2.6 | Unreported | Ambiguous |
| Nyayanga 5 (Kenya) | 24 | 196* | 14** | Hippo | Present | 3-2.6 | Unreported | Ambiguous |
| FLK North 6 (Tanzania) | 37 | 400 | 123*** | Elephant | Present | 1.8 | Absent | Ambiguous |
| FLK North Deinotherium site (Tanzania) | - | | 23 | Elephant | Present | 1.7 | Absent | Absent |

| | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| Barogali (Djibuti) | 37 | - | 569 | Elephant | Absent | 1.6-1.3 | Unreported | Absent |
| Nadung´a (Kenya) | 53 | 65 | 6797 | Elephant | Present | 1-0.7 | Unreported | Absent |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |

- \* Numbers in main text and supplementary tables do not coincide.
- \*\*Only artifacts reported as associated to the skeletal remains (total in both excavations is 135)
- \*\*\*Excluding purported manuports

**A**

Museum

FLK-W

FLK-NN

EAK

**B**

NE

SO

**Emiliano Aguirre Korongo**

Waxy claystone

Bed II

1F

Bed I

6.25 m

Waxy claystone

EAK

Ng´eju

1E

1D

1C

CHT

**C**

Waxy claystone

Bed II

Bed I

1F

Waxy claystone

Edaphic

m

2.0

1.5

1.0

0.5

A    Waxy claystone          10YR 4/1 (Dark grey)
B    (Colluvium)             5Y 4/3 (Olive)
Ab                           2.5Y 5/3
Cb   Waxy claystone
     (Lowermost Bed I)       10YR 3/2

EAK

1F

**a) Distance to elephant bones**

**b) Fitted model**

ANOVA p-value = <2.2e-16

**c) Estimated intensity (ρ)**

**d) Berman Z1 test**

z1 statistic = -5.759
p-value = 8.456e-09

**e) Berman Z2 test**

z2 statistic = -11.48
p-value = 1.585e-30

A

1 Thinning
2 Lateral retouch
3 Lateral retouch

B

C