Early oligodendrocyte dysfunction signature in Alzheimer's disease: Insights from DNA methylomics and transcriptomics

Katherine Fodder[1], Hannah M.G Smith[1], Umran Yaman[2], Ignazio S. Piras[3], Megha Murthy[4], John Hardy[1,2], Tammaryn Lashley[1], Rohan de Silva[4,5], Dervis A Salih[1,2], Conceição Bettencourt[1*]

1. Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, London, UK

2. UK Dementia Research Institute, London, UK

3. Neurogenomics Division, Translational Genomics Research Institute, Phoenix, AZ, USA

4. Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, London, UK

5. Reta Lila Weston Institute, UCL Queen Square Institute of Neurology, London, UK

*Corresponding Author:

Prof. Conceição Bettencourt, PhD

Department of Neurodegenerative Disease

UCL Queen Square Institute of Neurology

1 Wakefield Street

London WC1N 1PJ

United Kingdom

Email c.bettencourt@ucl.ac.uk

ABSTRACT

INTRODUCTION

Alzheimer's disease (AD) research has largely focused on neurons, with glial contributions, particularly from oligodendrocytes (OLGs), often overlooked. DNA methylation alterations are well documented in AD, but cell-type-specific insights remain limited. Here, we investigate OLG-associated DNA methylation across brain regions to further understand its role in AD.

METHODS

We applied weighted-gene correlation network analysis (WGCNA) to human DNA methylation data from four brain regions, human brain single-nuclei RNA sequencing, and mouse brain RNA sequencing. Networks were examined to identify disease-associated modules enriched for OLG genes and compared across datasets.

RESULTS

We identified an AD-associated DNA methylation signature enriched for OLGs, which was preserved across brain regions, and linked to altered OLG gene expression. Dysregulated signatures spanned multiple disease stages, suggesting early OLG alterations are consistent features of AD.

DISCUSSION

Our findings implicate early OLG-specific DNA methylation changes as a potential initiating factor of AD pathogenesis.

Keywords: Alzheimer's disease, neurodegeneration, neurodegenerative disease, early pathogenesis, oligodendrocyte dysfunction, brain tissue, DNA methylation, gene expression, WGCNA, multiomics, gene networks

1 INTRODUCTION

Alzheimer's disease (AD) is the leading cause of dementia worldwide[1]. Although neurodegenerative diseases such as AD have been associated with grey matter and neuronal damage, there is evidence for decline and involvement of white matter and glial cells during disease progression[2–6]. Oligodendrocytes (OLGs), which are not well studied in neurodegenerative diseases, are glial cells in the CNS, responsible for the production, stability, and maintenance of myelin, the lipid-rich, multilamellar membrane which wraps around axons and enables fast transmission of electrical signals[7]. Early evidence of the disruption of myelin in AD suggests those regions of the cortex, such as the temporal and frontal lobes, that are myelinated later in development are more likely to present with AD pathology earlier[2,4,8]. This suggests that those regions that myelinate later are more vulnerable to pathogenic mechanisms which result in neurodegeneration. Moreover, brain imaging data has indicated that Aβ deposition may change white matter microstructure in early disease stages[9]. Although the precise role of OLGs in AD pathology is still unclear, there is evidence from human post-mortem studies showing alterations in the numbers and morphology of OLG lineage cells[3] as well as decreases in Olig2￭+￭cells[10], and an increased number of oligodendrocyte precursor cells (OPCs) in white matter lesions[11]. Morphological changes are also seen in OLGs derived from AD post-mortem brains, specifically decreased nuclear diameter in the parahippocampal white matter[12]. Genome-wide association studies (GWAS) have also implicated specific myelin/OLG-related genes in neurodegeneration, including the bridging integrator 1 (*BIN1*) gene, which is the second strongest genetic risk factor for late onset AD[13–15] and known to be largely expressed by mature OLGs in white matter tracts[16]. As well as observed changes in OLGs in AD, recent research has provided evidence OLG derived Aβ contributes to AD pathology, and that suppression of such production by OLGs in mice reduces AD associated pathology[17]. Transcriptomic analyses reveal gene expression changes in additional myelin-related/OLG genes in AD[18], further supporting the importance of myelination changes in AD. Interestingly, there are also genetic links between AD and some leukodystrophies, a rare group of inherited disorders characterised by loss of myelin[19]. For example, genetic variants in the AD associated gene *TREM2* are thought to be causative for an adult onset leukodystrophy[20,21]. Several other Mendelian leukodystrophy genes (e.g. *CSF1R* and *NOTCH3*) are also linked to AD[22]. This genetic overlap underscores the need to elucidate mechanisms of dysfunction of myelin in AD. Although OLG and myelin dysfunction in AD have historically been considered as consequences of neuronal pathology rather than causative factors, emerging evidence suggests

that changes in myelin are an early feature of AD[3,23–26], indicating that oligodendrocyte dysfunction may play a contributory or even initiating role in disease development.

DNA methylation, the most widely studied epigenetic modification, is crucial in the spatio-temporal control of gene expression, but has also been implicated in neurodegenerative disease[27–29]. In AD, epigenome-wide association studies (EWAS) utilising bulk brain tissue have identified multiple genes with DNA methylation changes associated with the disease and its pathological burden[30–33], and meta-analyses have identified significant changes across multiple brain regions[29,34–36]. More recently, various studies have investigated DNA methylation changes in glia and neurons separately, and have identified many AD-associated DNA methylation changes that had previously been detected in 'bulk' tissue but were indeed driven by changes in non-neuronal cells, including in OLGs[34,37]. However, detailed characterization of glial cell-type specific effects of dysregulated DNA methylation, particularly in OLGs, is limited.

In this study, we aimed to investigate dysregulation of OLG DNA methylation and their contribution towards AD. We utilised co-methylation network analysis, an agnostic systems biology approach to identify higher order relationships between biological entities, to uncover DNA methylation signatures derived from human post-mortem brain tissue that are associated with AD and relevant to OLG lineage cells. Following the identification of these AD-associated co-methylation OLG signatures, we investigated whether they were present in brain regions differently affected by AD pathology as well as in additional neurodegenerative diseases, such as primary tauopathies. Additionally, we integrated human brain single-nucleus RNA sequencing data and mouse brain gene co-expression networks to further investigate these OLG dysregulated signatures and whether they might be occurring early in the disease process. By bringing together multiple data modalities, we provide a multi-layered view of epigenetic and transcriptional changes supporting an important role for molecular dysregulation of OLGs early in the neurodegenerative process, an area which has so far been overlooked.

2 METHODS

2.1 DNA methylation datasets and pre-processing

Data used to investigate DNA methylation signatures in AD was generated as described by Semick et al.[27] and retrieved from GEO under the accession code GSE125895. This dataset comprised samples across multiple brain regions, including the cerebellum (CRB) (N = 67), which is affected last by AD pathology, while the hippocampus (HIPPO) (N = 65), entorhinal cortex (ERC) (N = 69), and dorsolateral prefrontal cortex (DLPFC) (N = 67) are all affected to varying degrees in AD, with the ERC and HIPPO among the earliest sites of pathology and the DLPFC becoming involved at later disease stages (**Table 1**).

Beta-values ranging from 0 to 1 (approximately 0% to 100% methylated, respectively), were used to estimate the methylation levels of each CpG site using the ratio of intensities between methylated and unmethylated alleles. Data analysis was conducted using several R Bioconductor packages as previously described[38]. After importing raw data in the form of idat files, all datasets were subjected to pre-processing and quality control checks using minfi[39], wateRmelon[40], and ChAMP[41] packages. Probes were removed if they: (1) showed poor quality, (2) were cross reactive, (3) included common genetic variants, and (4) mapped to X or Y chromosome. Samples were also dropped if: (1) they presented with a high failure rate (≥ 2% of probes), (2) the predicted sex did not match the phenotypic sex, and (3) they clustered inappropriately on multidimensional scaling analysis. Beta-values were normalised with ChAMP using the Beta-Mixture Quantile (BMIQ) normalisation method. M-values, computed as the logit transformation of beta-values, were used for all statistical analysis, as recommended by Du et al. [42], due to their reduced heteroscedasticity and improved statistical validity in comparison to beta-values for differential methylation analysis.

An overview of the methodological framework of this study is presented in **Figure 1**. Following the approach described by Fodder et al[43], and further detailed below, we conducted weighted gene correlation network analysis (WGCNA). Prior to input into WGCNA, we carried out covariate adjustment using linear regression modelling to account for unwanted technical and biological effects (**Table 1**). Brain cell-type proportions were estimated using a cell-type deconvolution method and reference panel (NeuNP, NeuN-positive nuclei; DoubleN, NeuN–/SOX10– nuclei; Sox10P, SOX10-positive nuclei) described by Shireby et al.[34].

FIGURE 1

Overview of the methodological framework with a multi-omic network analysis approach to identify oligodendrocyte (OLG)-associated gene dysregulation in AD. 1) Human AD DNA methylation co-methylation networks were constructed using adjusted M-values from multiple human brain regions (e.g. DLPFC, ERC, and HIPPO. **2)** Human AD single-nucleus RNA sequencing (snRNA-seq) data was then used to identify OLG clusters, where differentially expressed genes were identified. HdWGCNA was applied to build OLG-specific co-expression networks. **3)** Mouse hippocampal gene expression data of AD models was then used to construct co-expression networks that were enriched for OLG signatures. **4)** Modules identified from the three network types were compared to uncover genes present across multiple modules. DLPFC, dorsolateral prefrontal cortex; ERC, entorhinal cortex; HIPPO, hippocampus; AD, Alzheimer's disease; OLG, oligodendrocyte; WGCNA, weighted gene-correlation network analysis; hdWGCNA, high-dimensional weighted gene correlation analysis; snRNAseq, single-nucleus RNA sequencing.

TABLE 1. Characterization of the four brain region AD datasets and selected regression models for input to WGCNA

Dataset	AD cases and controls included after quality control	Mean age ± SD (years)	Sex	Regression models used for covariate adjustment	Number (%) of methylation sites included for WGCNA	Soft-thresholding power used for WGCNA
AD-HIPPO	AD (*N* = 17)	81.5 ± 9.16	8M/9F	*~0+Sample_group+plate +DoubleN+sex+NeuNP+ age+array*	44,609 (20%)	12
	Controls (*N* = 48)	61.7 ± 7.66	29M/19F			
AD-DLPFC	AD (*N* = 21)	79.9 ± 9.46	11M/10F	*~0+Sample_group+Neu NP+age+sex+plate*	53,826 (20%)	14
	Controls (*N* = 46)	61.3 ± 7.56	27M/19F			
AD-ERC	AD (*N* = 20)	79.7 ± 9.64	9M/11F	*~0+Sample_group+slide +DoubleN+Sox10P+age + 1/1 surrogate variables detected*	52,296 (20%)	12
	Controls (*N* = 49)	61.6 ± 7.61	29M/20F			
AD-CRB	AD (*N* = 24)	79.5 ±10.0	11M/13F	*~0+Sample_group+slide +sex+age+NeuNP+Dou bleN*	52,712 (20%)	16
	Controls (*N* = 43)	60.6 ± 7.04	24M/19F			

AD, Alzheimer's disease; HIPPO, hippocampus; DLPFC, dorsolateral prefrontal cortex; ERC, entorhinal cortex; CRB, cerebellum; SD, standard deviation; F, females; M, males; NeuNP, estimated NeuN-positive nuclei proportion; DoubleN, estimated NeuN–/SOX10– nuclei proportion; Sox10P, estimated SOX10-positive nuclei proportion; WGCNA, weighted gene correlation network analysis.

2.2 Human brain gene expression

We used human brain single-nucleus RNA sequencing (snRNAseq) data from the previously published and publicly available ROSMAP study[44] obtained through Synapse (synapse ID [syn18485175](syn18485175)). We used filtered cell counts from 24 individuals showing little or no pathology and 24 showing mild to severe AD pathology (*N* = 48 total). Pathology stage was also assigned to each individual as 'no-pathology', 'early-pathology' or 'late-pathology' as described by Mathys et al.[44]. Filtered data consisted of 70,634 droplet-based single-nucleus covering a total of 17,926 genes. We used Seurat to normalise the data with the function "NormalizeData" with the option "LogNormalize", using a scale factor of 10,000. We then assigned cell cluster identities as described by Mathys et al.[44], which contained 8 cell-type clusters: Astrocytes (Ast), Endothelial

cells (End), Excitatory neurons (Ext), Inhibitory neurons (In), Microglia (Mic), Oligodendrocytes (Oli), Oligodendrocyte precursor cells (Opc) and pericytes (Per). Subclusters of these cells were also defined, which for OLGs were Oli_0,1,3,4 and 5.

2.3 Mouse brain gene expression

As a way to investigate early disease changes, we also examined mouse brain gene expression to complement the human post-mortem end stage disease findings. Such mouse models represent the early-stage of amyloid pathology but no tau pathology. The gene expression dataset was generated using AD mouse models as described by Salih *et al.*[45], Matarin *et al.*[46]. In brief, four transgenic mouse lines which develop Aβ pathology at varying rates were included: *APP, PSEN1,* homozygous *APP/PSEN1* and heterozygous *APP/PSEN1* alongside age-matched wild-type mice (**Table S1**). For each transgenic line, at least three mice were used, except for *APP* mice at 4 months, where only two mice were available (**Table S1**). A minimum of seven littermate controls (*WT*) from the original parental lines were also included. The hippocampi of male mice at the age of 2, 4, 8 and 18 months were used for transcriptomic analysis. Procedures involving mice were conducted in accordance with local ethical agreement, and in compliance with the UK Animals Act (1986).

As described by Salih *et al.*[45], RNA-seq library preparation and sequencing were performed by Eurofins Genomics (Ebersberg, Germany). Briefly, RNA strand-specific libraries were created using TruSeq Stranded mRNA Library Preparation Kit, Illumina according to the manufacturer's instructions. Raw sequencing data were processed using RTA version 1.18.64, and FASTQ files were generated using bcl2fastq-1.8.4. Low-quality base pairs and adaptors were removed using Trim Galore (Babraham Bioinformatics). Transcripts were quantified using Salmon[47] and gene level quantification, expressed as transcripts per million (TPM), was obtained using the tximport package in R[48]. Gene expression values were transformed by applying log2 (normalised counts+1), and lowly expressed genes, i.e. those with an average detection level of below 1.5 TPM were excluded, reducing the dataset from 48,525 to 15,453 genes.

2.4 Weighted gene correlation network analysis (WGCNA)

We used a systems biology approach to identify co-methylation and co-expression modules, i.e. clusters of highly correlated DNA methylation sites (CpGs) or genes. For this we have used weighted gene correlation network analysis (WGCNA[49]) for the four human bulk brain tissue

DNA methylation datasets (**Table 1**) and for the mouse gene expression dataset (**Table S1**). For all DNA methylation datasets, we excluded those CpGs not mapping to genes, and selected only those showing the highest variance (top 20%) across individuals within each dataset. The percentage of total methylation sites and number of methylation sites included for each analysis is given in **Table 1**, as is the soft-thresholding power used for network construction. We used covariate adjusted M-values for the selected CpGs as input for this analysis. For the mouse expression data, following sample clustering, three outliers were removed leaving 83 samples for the network analysis. A soft-thresholding power of 6 was used. To construct all networks, pairwise DNA methylation/expression correlations were calculated to create a similarity matrix, as previously described[49].

Module membership (MM) was then reassigned for each network using the applyKMeans function of the CoExpNets package[50]. CpGs/genes with high MM are considered hub genes, meaning they are highly connected and likely key biological drivers within that module. Gene significance was also considered, which here represents the correlation between the methylation levels at a specific site and the disease status[49].

After co-methylation/expression modules were identified, we carried out module-trait correlation analysis (as part of the WGCNA workflow[49]) to understand the relationship between co-methylation modules and disease traits. For the human co-methylation modules, this association was carried out with AD status (i.e. AD or control sample). Whilst for the mouse co-expression modules association was based on a semi-quantitative score based on Aβ pathology severity, ranging from 0-5 obtained as previously described[45]. We applied Bonferroni multiple testing correction, i.e. corrected correlation p-value over the number of modules detected, to minimize the finding of false-positives.

2.5 Cell-type enrichment analyses for co-regulation modules

Cell-type enrichment analysis for the human co-methylation and mouse co-expression modules was conducted using the EWCE package and the associated single-cell mouse transcriptomic dataset[51,52]. This analysis aimed to determine whether the genes within each co-methylation/co-expression module were enriched for markers of a specific cell type. Overrepresented cell types were identified using p-values derived from 1000 iterations. For the human co-methylation modules we additionally checked this enrichment using an OLG gene list from human snRNA-

sequencing data generated by Mathys et al.[44] and curated by Piras et al.[53]. We used Fisher's exact test to confirm whether OLG modules identified through the EWCE package were also statistically significantly enriched for genes within this OLG list.

2.6 High dimensional WGCNA (hdWGCNA)

To confirm the cell-type specificity and the OLG relevance of the identified co-methylation signatures, we also utilised hdWGCNA to identify co-expression networks within the high-dimensional snRNA-seq dataset. Using cells belonging to the five OLG clusters (Oli_0-5 described above), we restricted our analysis to genes that were expressed in at least 1% of cells (total genes = 9,254). Following the pipeline described by Morabito et al.[54], we generated metacells, which aggregate similar cells within each group (OLG subclusters) and averaged their expression profiles, thereby reducing noise and improving module detection. The resulting metacell expression matrix was then log-normalized, before soft-thresholding was carried out to identify a power of 8 for network construction.

2.7 Preservation Analysis

To identify which co-methylation modules in each of AD networks were preserved (i.e. shared) or perturbed (i.e. unique) in the other brain regions, and against additional AD and other neurodegenerative disease datasets, we employed module preservation analysis, as described by Langfelder et al.[55]. For each network (taken as the "reference dataset"), module preservation in the other datasets (the "test data") was calculated using the modulePreservation function implemented in WGCNA, carried out with 200 permutations for each analysis. As a measurement of module preservation, we used the Z-summary statistic (a composite measure to summarise multiple preservation statistics). A Z-summary greater than 10 indicates a strong preservation of a module in the "test data", a Z-summary of between 2 and 10 indicates moderate preservation, and a Z-summary less than 2 indicates no preservation. To investigate preservation, we used five additional complementary DNA methylation datasets derived from AD, frontotemporal lobar degeneration (FTLD) and movement disorders (MD), which are summarised in **Table 2**.

TABLE 2. Pathological and demographic characteristics of DNA methylation datasets used for preservation

Dataset	Disease subtypes and controls (N after quality control)	Mean age ± SD (years)	Sex	Regression models used for adjustment
FTLD1 (Frontal Cortex) Fodder et al.[43]	FTLD-TDP (*N* = 15)	70.07 ± 5.59	7M/8F	*~ 0 + disease + age + sex + SOX10P proportions + DoubleN proportions + array* (0 surrogate variables detected)
	Controls (*N* = 8)	75.75 ± 5.63	3M/5F	
FTLD2 (Frontal Cortex) Menden et al.[56]	FTLD-TDP/FTLD-tau (*N* = 34)	63.18 ± 7.92	14M/20F	*~ 0 + disease + age + sex + SOX10P proportions + DoubleN proportions + array + slide* (0 surrogate variables detected)
	Controls (*N* = 14)	78.43 ± 11.76	5M/9F	
FTLD3 (Frontal Cortex) Weber et al.[57]	FTLD-Tau (PSP) (*N* = 93)	71.16 ± 5.32	54M/39F	*~ 0 + disease + age + sex + SOX10P proportions + DoubleN proportions + array + slide + surrogate variable* (1/1 surrogate variables detected)
	Controls (*N* = 70)	76.17 ± 7.93	45M/25F	
AD (Dorsolateral Prefrontal Cortex) De Jager et al.[30]	AD (*N* = 201)	85.4 ± 5.39	120M/81F	*~0 + disease + slide + bs_conversion + NeuNP proportion + DoubleN proportion + array + PC1+ PC2*
	Controls (*N* = 329)	87.3 ± 3.91	212M/117F	
MD (Frontal lobe white matter) Murthy et al.[58]	Controls (*N* = 17)	72.35 ± 4.47	8M/9F	*~0 + disease + age + sex + PMI + NeuNP proportion + DoublN proportion + slide, + array*
	MSA (*N* = 17)	67.71 ± 6.47	9M/8M	
	PD (*N* = 17)	68.06 ± 3.67	8M/9F	
	PSP (*N* = 17)	65.35 ± 3.75	9M/8F	

analysis

FTLD, Frontotemporal lobar degeneration; FTLD-TDP, FTLD with 43 kDa transactive response DNA-binding protein (TDP-43) positive inclusions; FTLD-Tau, FTLD with tau-positive inclusions; MD, movement disorders; PSP, progressive supranuclear palsy; AD, Alzheimer's Disease; PD, Parkinson's Disease; MSA, Multiple System Atrophy, SD, Standard deviation; F, Females; M, Males; NeuNP, estimated NeuN-positive nuclei proportion; DoubleN, estimated NeuN–/SOX10– nuclei proportion; Sox10P, estimated SOX10-positive nuclei proportion; PC1, 1st principal component; PC2, 2nd principal component, PMI; post-mortem interval.

2.8 Genetic enrichment assessment

To investigate whether any of the co-methylation modules we produced during the analysis described above were enriched for genes known to confer genetic risk of AD, we utilised MAGMA-Generalized Gene-Set Analysis of GWAS Data[59]. MAGMA uses a multiple regression framework to enable powerful and efficient gene- and gene-set level analysis.

3 RESULTS

3.1 Human brain co-methylation network analysis reveals significant oligodendrocyte dysregulation associated with AD

We generated co-methylation networks for four brain regions affected by different degrees of AD pathology (**Figure S1**), with the HIPPO representing the brain region displaying pathology earlier in AD, followed by the ERC and DLPFC, and the CRB, which is usually spared from typical AD pathology. For the DLPFC, ERC, and HIPPO and CRB networks, 6/16 ($p < 0.003$, 0.05/16 modules), 17/25 ($p < 0.002$, 0.05/25 modules), 7/18 ($p < 0.003$, 0.05/18 modules) and 4/6 ($p < 0.008$, 0.05/6 modules) co-methylation modules, respectively, were found to be associated with the disease status (i.e. AD vs control) (**Figure S1**). We next analysed which of these modules were enriched for OLG/OPC relevant genes, and identified one module in each of the DLPFC, ERC and HIPPO networks, the DLPFC-greenyellow, ERC-tan and the HIPPO-grey60 (**Figure 2, Figures S1-S2, Tables S2.1-2.3**). For the CRB networks, we did not find significant enrichment for OLGs/OPCs (**Figure S1B**). All these modules, the DLPFC-greenyellow, ERC-tan and the HIPPO-grey60 also showed significant positive correlation between module membership and gene-level significance for AD status (r = 0.69, r = 0.91, r = 0.53; p < 1x10^{-138}; respectively), and a positive correlation with disease status, i.e. increased DNA methylation levels in AD compared to controls (**Figure S1**).

Hub gene analysis, which identifies in this case the most strongly connected CpGs and related genes for each OLG-enriched module, reinforces the relevance for OLG lineage cells in AD-associated modules. The top hub gene (highest module membership (MM)) for the DLPFC-greenyellow module was *MOG* (myelin oligodendrocyte glycoprotein) (**Figure 2A, Table S2.1**), a well described marker of mature OLGs[60], which has also been found differentially methylated

in chronic demyelinated lesions in multiple sclerosis[61]. The CpG with the second highest MM mapped to *ATP11A* (ATPase phospholipid transporting 11A), a gene which is more highly expressed in OLGs compared to other brain cell types ([ATP11A Tissue Expression](#)) (**Table S2.1),** and is mutated in hypomyelinating leukodystrophy[62]. We noted that 26 CpGs within this greenyellow module mapped to this gene *ATP11A* (**Table S2.1**).

The top hub gene for the ERC-Tan module, with the two CpGs with highest MM, was *C11ORF9* (also known as *MYRF*) (**Figure 2A, Table S2.2**), which codes for myelin regulatory factor, an important driver of post-OLG differentiation myelination and myelin maintenance[63]. In a previous study using gene expression and proteomic networks, *MYRF* was found to be a key driver of AD-associated disrupted OLG signatures[64]. The *ATP11A* gene, which was also highlighted in the DLPFC-greenyellow module, has 12 CpGs within this tan module (**Table S2.2**). A CpG mapping to *MBP* (a well described marker of mature OLGs) also presents with high MM within the module (**Table S2.2**).

The hub gene for the HIPPO-grey60 module was *MGC14436,* according to the Illumina DNA methylation array manifest, which maps to chromosome 12 and is not well characterised. However, further inspection of the genomic location of this CpG revealed that it lies within the region of the *FAM222A* gene*,* which encodes a protein previously reported to build up in Aβ plaques in AD[65], and has been shown to contain genetic variants associated with AD risk[26]. The gene has also been reported to be co-expressed with several other AD related genes[66]. This disease-associated OLG module also contained 27 CpGs mapping to the gene *ATP11A* (**Table S2.3)**, as we had seen with the ERC-tan and the DLPFC-greenyellow module. Full details of all CpGs within each module and corresponding MM information are given in **Tables S2.1-3.**

We also examined similarities across brain regions for the OLG-enriched AD co-methylation modules. Pairwise comparisons of module membership (MM) values revealed moderate and highly significant positive correlations between regions (**Figure 2B**). We also utilised module preservation statistics to evaluate how well the co-methylation modules identified in one region were preserved in the others. The DLPFC-greenyellow module showed strong preservation in both ERC and HIPPO datasets (Z.summary > 10), as did the ERC-tan module in the DLPFC and the HIPPO networks. The HIPPO-grey60 module was preserved in ERC and DLPFC, albeit only moderately in DLPFC (**Figure 2C**). Looking in more detail at which CpGs were present across all three, we found that 74 CpGs belonged to all of the DLPFC-greenyellow, ERC-tan

and HIPPO-grey60, mapping to 54 distinct genes (**Table S3**). Notably, 8/74 probes mapped to the gene *ATP11A* (highlighted above). Other genes containing multiple probes within this set included *INPP5A* (to which 5/74 CpGs mapped), *RASA3* (3/74) and *FAM107B* (3/74).

To assess the reproducibility of the co-methylation modules, we evaluated their preservation in an independent AD DNA methylation dataset from the ROSMAP study (derived from prefrontal cortex tissue). All three modules showed strong preservation, suggesting that they represent robust biological signals (**Figure 2C**). We also investigated whether this preservation extended to other neurodegenerative diseases, namely several frontotemporal lobar degeneration (FTLD) subtypes we previously investigated ranging between FTLD-tau and FTLD-TDP pathology[43] and a DNA methylation dataset derived from the white matter (highly enriched for OLGs) of multiple system atrophy (MSA), progressive supranuclear palsy (PSP) and Parkinson's disease (PD) samples[58]. Both the DLPFC-greenyellow and the ERC-tan AD-associated modules (OLG enriched), were highly preserved (the most preserved of all co-methylation modules) in the FTLD3 dataset (composed of sporadic primary tauopathy cases - PSP) and moderately preserved in the FTLD1 and FTLD2 dataset (mostly TDP-43 proteinopathies) (**Figure 2C**). All three modules were also highly preserved in the white matter MSA/PSP/PD dataset (**Figure 2C**), indicating that in addition to AD these dysregulated OLG signatures are also highly relevant to other neurodegenerative diseases with multiple underlying pathologies.

We also checked to see if there was any enrichment of a specific CpG feature (i.e. where within the gene the CpG was located) within these modules. We found that there was a significant overrepresentation of CpGs located within the gene body in all three modules (**Figure 2D**). Although the role of DNA methylation within the gene body is less well characterised compared to that in the promoter regions (where increased DNA methylation is often associated with a decrease in gene expression), our data points towards the importance of gene body DNA methylation dysregulation in AD.

FIGURE 2

Human brain co-methylation network analysis shows significant dysregulation of OLG signatures across brain regions in AD.
A) Top 25 genes with highest module membership across the DLPFC-greenyellow, ERC-tan and HIPPO-grey60 modules (top to bottom). Edge surrounding the genes indicates the feature of the CpG mapping to each gene as given by the legend. B) Scatterplots show pairwise relationships between CpG-level MM values from the grey60, tan, and greenyellow modules with Pearson correlation coefficients (r) and significance indicated (***$p < 0.001$). C) Module preservation scores between co-methylation modules (DLPFC, ERC and HIPPO), ROSMAP (AD), FTLD, and MD test data. D) Enrichment of methylation site features within 3 AD-associated modules of interest. AD, Alzheimer's Disease; FTLD, Frontotemporal lobar degeneration; HIPPO, hippocampus; DLPFC, dorsolateral prefrontal cortex; ERC, entorhinal cortex; MD, movement disorder; MM, module membership; TSS200, transcription start site -200; TSS1500, transcription start site -1500-200; 3'UTR, 3 prime untranslated region; 5'UTR, 5 prime untranslated region.

We checked whether any of the modules were linked to AD risk genes using MAGMA and associated GWAS summary statistics[14], and found no enrichment within the three modules (data not shown), suggesting that known AD-associated genetic effects are unlikely to be driving the OLG dysregulation we report above.

3.2 Genes in AD-associated OLG co-methylation signatures show upregulation in AD OLGs

Next, we investigated whether expression of genes present across the co-methylation modules was distinct between AD and controls using a human OLG snRNAseq dataset. First, we checked for genome-wide differential expression of all genes within the three co-methylation modules (**Figure 3A**). We found that most of these genes showed upregulation in AD compared to control OLGs, and also that the majority of those genes were differentially expressed in a specific single-cell OLG cluster - Oli3. The finding that many genes were upregulated was interesting given that we had seen an enrichment of methylation in gene body features within our three OLG AD DNA methylation modules. As described previously, increased DNA methylation in the body of the gene is often associated with an increase in gene expression[67]. Therefore, our data is in accordance with this direction of effect (increased gene body methylation and consequent increased gene expression).

We also investigated which of these genes were differentially expressed between early-pathology/late-pathology and controls. We saw that genes within these modules were differentially expressed at both stages of disease (**Figure 3B**), indicating that these changes represented are not solely occurring at end-disease stages.

Of the genes that were most highly differentially expressed in the Oli3 cluster, several were included in at least one of our co-methylation modules, including *QDPR* and *CRYAB* (**Figure 3C**)*,* which Mathys et al.[44] have previously described as also being upregulated at the protein level in white matter of individuals with AD[44].

FIGURE 3

Expression levels of genes present in disease associated co-methylation modules in OLG snRNA-sequencing clusters.
A) Bar plots showing the number of genes from each co-methylation module (DLPFC-greenyellow, ERC-tan, and HIPPO-grey60) that are upregulated (red) or downregulated (blue) in each snRNAseq OLG subcluster (Oli0–Oli3). **B)** Bar plots showing the number of genes from each co-methylation module (DLPFC-greenyellow, ERC-tan, and HIPPO-grey60) that are upregulated (red) or downregulated (blue) in each OLG subcluster (Oli0–Oli3) and stratified by early (light shades) and late (dark shades) pathology stages. **C)** Volcano plots showing differential gene expression across Oli0, Oli1, and Oli3. Significantly differentially expressed genes (adjusted p-value < 0.05) from the co-methylation modules are coloured by their module membership. Genes present in multiple modules are shown in combined colours. Labelled genes are the top 10 most significantly dysregulated within each subcluster which are also module members. AD, Alzheimer's disease; DLPFC, dorsolateral prefrontal cortex; ERC, entorhinal cortex; HIPPO, hippocampus; OLG, oligodendrocyte.

In order to understand how patterns of co-methylation might relate to gene co-expression changes, we used the snRNA-sequencing data and generated co-expression modules using the OLG clusters to identify OLG genes that are co-regulated. Using hdWGCNA, we identified 5 modules within the OLG clusters: hd-green, hd-brown, hd-turquoise, hd-blue and hd-yellow. All modules showed significant association with at least one pathological trait (i.e. amyloid, Braak stage and/or cognitive decline) within one subcluster of OLGs (**Figure 4A**). As we did between co-methylation modules across brain regions, we looked at correlation between module membership within co-methylation OLG modules and snRNAseq gene expression modules. As co-methylation modules are built at the probe level, for each gene within the co-methylation modules, we selected the CpG with the highest module membership (MM) for this comparison. The snRNAseq hd-turquoise module showed significant positive correlations with the DLPFC-greenyellow and ERC-tan modules (**Figure 4B**). While the snRNAseq hd-blue module was significantly correlated with the HIPPO-grey60 module. We next looked at which genes showed high MM across co-methylation and co-expression modules (**Table S4.1-4.2**). The gene with the second highest module membership in the hd-turquoise module was *CRYAB,* which was also part of the DLPFC-greenyellow module (**Table S2.1**). Other genes with high MM in the hd-turquoise module (top 5%) included *QDPR* (**Figure 4C**)*,* which was included in all three co-methylation modules (**Tables S2.1-2.3**).

FIGURE 4

Co-expression network analysis of human oligodendrocyte snRNAseq clusters. A) Heatmap showing correlations between module eigengenes of hdWGCNA co-expression modules and neuropathological traits (pathology stage, amyloid burden, Braak stage, and cognitive diagnosis) for all oligodendrocyte clusters and for individual subclusters (Oli0–Oli5). Red indicates positive and blue negative correlations. Significance is indicated: $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***). **B)** Heatmap of Pearson correlations between module membership (MM) values of three disease-associated co-methylation modules (ERC-tan, HIPPO-grey60 and DLPFC-greenyellow) and MM in each hdWGCNA module. **C)** Network plots of two hdWGCNA modules (blue and turquoise), showing connections amongst top 15 hub genes. hdWGCNA; high-dimensional weighted gene correlation network analysis, MM; module membership.

3.3 AD-associated OLG co-methylation networks show dysregulation of gene expression in mouse models of early AD stages

To complement human post-mortem brain tissue derived datasets and be able to infer the relevance of dysregulated OLG signatures at an early stage of AD pathology, we have used gene co-expression networks using a dataset derived from early-stage AD mouse models. A total of 10 mouse brain co-expression modules were identified and then related to sample traits: Aβ pathology, age and genotype (**Figure S3A**). Three modules were significantly positively associated with Aβ pathology ($p \leq 0.005$ (0.05/10)), indicating that the expression of genes within these modules significantly increases with increasing pathology. We identified the pink mouse co-expression module as significantly associated with AD-related pathology and enriched for OLGs (**Figure S3B**).

As expected, well-known OLG marker genes were present across the pink co-expression module, including *Mobp, Plp1, Fa2h, Sox10* and *Mag*, and hub genes included *Myo1d, Clic4, Ttyh2, Pacs2, Arhgef10, Tmem88b, Carns1, Cntn2,* and *Scd1* (**Figure 5A, Table S5**). There was a significant overlap of the mouse pink module genes with all three human OLG-relevant AD co-methylation modules (**Figure S3C**), and the hub genes of the ERC-tan and DLPFC-greenyellow modules, *Mog* and *Myrf* were also present within the pink module. The gene *Fam222A* (possible HIPPO-grey60 hub gene) was also present in this module. Furthermore, many of the overlapping genes present with high MM in both human and mice networks (**Figure 5B**), strongly supporting those genes as important drivers within these networks across species. Additionally, because these mice do not have tau pathology, this data suggests that dysregulation of these OLG signatures happens at an early stage of AD pathology before tau pathology develops. Interestingly, because these OLG signatures are highly preserved in primary tauopathies and alpha-synucleinopathies (preservation analysis above), this suggests that their dysregulation may also be independent of amyloid pathology. A summary outline of these findings is shown in **Figure 6**.

Figure 5

Brain co-expression network analysis of early-stage AD mouse models. A) WGCNA 'pink' oligodendrocyte enriched co-expression module. Co-expression network for the 'pink' module constructed using Cytoscape software. Nodes represent genes and edge lines represent co-expression connections. The top 25 most connected genes out of the 442 total genes in the module are shown. **B)** Scatter plots showing the relationship between human co-

methylation and mouse co-expression module membership (MM) for genes in the DLPFC-greenyellow, ERC-tan, and HIPPO-grey60 co-methylation modules. The x-axis represents MM in the human co-methylation modules; the y-axis represents MM in the corresponding mouse co-expression module. Genes with high MM in both species are highlighted in blue and labelled. Dashed lines indicate the MM threshold of 0.5.

FIGURE 6

Schematic outlining the datasets and disease contexts in which a preserved oligodendrocyte (OLG) signature across neurodegeneration was identified. AD, Alzheimer's disease; DLPFC, dorsolateral prefrontal cortex; ERC, entorhinal cortex; FTLD, frontotemporal lobar degeneration; HIPPO, hippocampus; MSA, multiple system atrophy; OLG, oligodendrocyte; PSP, progressive supranuclear palsy; PD, Parkinson's disease.

4 DISCUSSION

Here, we have employed network analysis across several DNA methylation and gene expression datasets representing different stages of AD pathology in order to investigate the effect of DNA methylation on dysregulation of OLG relevant genes in AD. Through these networks, constructed on different modalities of omics data derived from human and mouse brain tissue, we have identified AD-associated dysregulated OLG signatures shared between more mildly and severely affected human brain regions as well as in models of early stages of AD pathology. Our findings underpin the importance of studying OLGs as an early contributor to neurodegenerative processes, rather than as passive bystanders, a concept which is in its infancy of being explored[68].

From the human brain co-methylation networks, both the DLPFC-greenyellow and ERC-tan modules had hub genes which are known as important OLG genes and key regulators of myelination: *MOG* and *MYRF* respectively. *MOG* is a gene expressed by mature OLGs. *MYRF* is a transcription factor that is essential for myelin maintenance[69] and a target of *SOX10*, which is another crucial activator of OLG differentiation-related genes[70]. We noticed several methylation sites with high module membership (MM) across all three co-methylation modules mapped to the gene *ATP11A*, a gene implicated in a hypomyelinating leukodystrophy[62]. Mechanisms behind effects of mutations in this gene were suggested to involve disruption of phospholipids in the cell membrane which led to, amongst other effects, disrupted cell cholesterol homeostasis. Given the known importance of cholesterol functioning in OLGs mediated through the *APOE-ε4* allele in AD[71], this was of interest. As described, a link between AD and leukodystrophies has been suggested, with several genetic risk factors found to be shared between the two conditions[22]. In a previous transcriptomic study of AD, *ATP11A* was shown to be upregulated in AD compared to controls[72], and we also observed a nominally significant upregulation across multiple OLG clusters in AD compared to controls.

Additional genes with known relevance to AD pathology and white matter perturbations were also present in these dysregulated OLG modules/signatures. The gene *QDPR,* which codes for quinoid dihydropteridine (an enzyme important in the regeneration of tetrahydrobiopterin (BH4), important in nitric oxide production) was found to present across all three modules, and also within disease-associated human snRNAseq and the mouse OLG co-expression modules.

There are some limitations to the use of bulk DNA methylation data in investigating cell-type specific dysregulation, given that cell-type specific methylation changes may be masked by noise from other cell types. However, by using a complementary analysis based on human snRNA sequencing data, we found a significant overlap between genes within the co-methylation OLG modules and those differentially expressed within a cluster of OLGs that had been previously identified as AD associated by Mathys et al[44]. This suggests that DNA methylation acts on the dysregulation of the identified AD-associated OLG signature. We further investigated co-expression modules within the scRNA-seq data and identified that the hd-turquoise module was associated with several pathology measures in the disease associated Oli3 cluster, and contained a significant overrepresentation of genes that we had identified as showing dysregulated DNA methylation. This finding at the single nucleus level reinforces OLG

specific observations in the bulk human DNA methylation. To address additional limitations of using bulk human brain tissue (e.g. temporality of events), we further complemented our analyses with co-expression networks in early-stage mouse model data, and identified an amyloid pathology associated module that was enriched for genes showing OLG methylation changes in human AD brain tissue. Another limitation of the analysis here is the inability to distinguish between 5-methylcytosine (5mC) and 5-hydroxymethylcytosine (5hmC), an oxidative derivative of DNA methylation that is thought to play a role in OLG fate determination[73]. Understanding the role of this specific epigenetic mark could contribute to further insights into the role of DNA methylation on OPC to OLG maturation, which is suggested to be perturbed in AD pathology[74], and warrants further investigation in future studies.

In summary, we have found consistent OLG AD-associated DNA methylation signatures across brain regions affected at different stages of disease progression; the entorhinal cortex and hippocampus are known to exhibit changes early in AD, whilst the dorsolateral prefrontal cortex is typically affected later on in disease[75]. The finding of aberrant DNA methylation across both early- and late stages of pathology suggests alterations contributing to disease early on rather than a secondary effect of neurodegeneration. This interpretation is supported through our complementary analysis of the mouse co-expression modules, which represent an early-stage of amyloid pathology, where the mice have no tau pathology, suggesting molecular myelin-related changes are present at the time of amyloid deposition or even before. Furthermore, we saw that there was very high preservation of the same OLG co-methylation AD modules within a primary tauopathy (PSP) dataset and to some extent within datasets consisting of TDP-43 proteinopathies and synucleinopathies as well (**Figure 6**). Altogether, our findings suggest that OLG dysregulation may precede both amyloid and tau pathology in AD, possibly representing a more generic early event independent of downstream protein accumulation across neurodegenerative diseases. These findings together highlight the need for further research into the importance of OLGs as important in early neurodegenerative pathology.

Our findings reveal that oligodendrocytes, often overlooked in neurodegeneration, show early and disease-relevant DNA methylation changes in AD, highlighting DNA methylation as a potential driver of pathology and offering novel insights into mechanisms of dysfunction that may be common across neurodegenerative disorders.

AUTHOR CONTRIBUTIONS

KF contributed to data analysis, prepared the figures, and drafted the manuscript; HGMS contributed to data analysis, figure preparation and interpretation; UY, IP, MM, JH, TL, RdS and DAS, contributed with datasets and/or interpretation; CB conceptualised and supervised the study; All authors critically reviewed and approved the manuscript.

ACKNOWLEDGEMENTS

KF is supported by the Medical Research Council (MR/N013867/1) and the Multiple System Atrophy Trust. HGMS is supported by Alzheimer's Research UK. MM is supported by the Multiple System Atrophy Trust. RdS is supported by Reta Lila Weston Trust for Medical Research and CurePSP. TL is supported by Alzheimer's Society, Alzheimer's Research UK and the Association of Frontotemporal Dementia. JH and DS are supported by the UK Dementia Research Institute [award number UK DRI-1210] through UK DRI Ltd, principally funded by the Medical Research Council (MRC). D.A.S. also received funding from MRC award MC_PC_23036 to establish an MRC National Mouse Genetics Network Ageing Cluster, and from the ARUK pump priming scheme via the UCL network. JH is supported by the Dolby Foundation and by the NIHR University College London Hospitals Biomedical Research Centre. CB is supported by Alzheimer's Research UK and the Multiple System Atrophy Trust. The funding bodies had no role in the design of the study or collection, analysis, and interpretation of data nor in writing the manuscript.

CONFLICT OF INTEREST STATEMENT

The authors report no conflicts of interest.

REFERENCES

1.   2021 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2021;17(3):327-406.

2.   Papuć E, Rejdak K. The role of myelin damage in Alzheimer's disease pathology. *Archives of Medical Science : AMS*. 2020;16(2):345-345.

3. Nasrabady SE, Rizvi B, Goldman JE, Brickman AM. White matter changes in Alzheimer's disease: a focus on myelin and oligodendrocytes. *Acta Neuropathologica Communications*. 2018;6(1):22-22.

4. Fodder K, de Silva R, Warner TT, Bettencourt C. The contribution of DNA methylation to the (dys)function of oligodendroglia in neurodegeneration. *acta neuropathol commun*. 2023;11(1). doi:10.1186/s40478-023-01607-9

5. Molina-Gonzalez I, Miron VE, Antel JP. Chronic oligodendrocyte injury in central nervous system pathologies. *Commun Biol*. 2022;5(1):1274.

6. Huang JK, Káradóttir RT. Oligodendrocytes in health and disease. *Neuropharmacology*. 2016;110(Pt B):537-538.

7. Bradl M, Lassmann H. Oligodendrocytes: biology and pathology. doi:10.1007/s00401-009-0601-5

8. Braak H, Braak E. Development of Alzheimer-related neurofibrillary changes in the neocortex inversely recapitulates cortical myelogenesis. *Acta Neuropathologica 1996 92:2*. 1996;92(2):197-201.

9. Dc D, Sa H, Sr K, et al. Association of Amyloid Pathology With Myelin Alteration in Preclinical Alzheimer Disease. *JAMA Neurol*. 2017;74(1):41-49.

10. G, Behrendt, K, Baer, A, Buffo, et al. Dynamic changes in myelin aberrations and oligodendrocyte generation in chronic amyloidosis in mice and men. *Glia*. 2013;61(2):273-286.

11. Je S, Ms F, L, Clark, et al. White matter lesions in an unselected cohort of the elderly: astrocytic, microglial and oligodendrocyte precursor cell responses. *Neuropathology and applied neurobiology*. 2007;33(4):410-419.

12. Gagyi E, Kormos B, Castellanos KJ, et al. Decreased oligodendrocyte nuclear diameter in Alzheimer's disease and Lewy body dementia. *Brain Pathol*. 2012;22(6):803-810.

13. Hu X, Pickering E, Liu YC, et al. Meta-analysis for genome-wide association study identifies multiple variants at the BIN1 locus associated with late-onset Alzheimer's disease. *PLoS One*. 2011;6(2):e16616.

14. Kunkle BW, Grenier-Boley B, Sims R, et al. Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing. *Nat Genet.* 2019;51(3):414-430.

15. Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet.* 2013;45(12):1452-1458.

16. De Rossi P, Buggia-Prévot V, Clayton BLL, et al. Predominant expression of Alzheimer's disease-associated BIN1 in mature oligodendrocytes and localization to white matter tracts. *Molecular Neurodegeneration.* 2016;11(1). doi:10.1186/S13024-016-0124-1

17. Rajani RM, Ellingford R, Hellmuth M, et al. Selective suppression of oligodendrocyte-derived amyloid beta rescues neuronal dysfunction in Alzheimer's disease. *PLoS Biol.* 2024;22(7):e3002727.

18. Allen M, Wang X, Burgess JD, et al. Conserved brain myelination networks are altered in Alzheimer's and other neurodegenerative diseases. *Alzheimer's & Dementia.* 2018;14(3):352-366.

19. Nowacki JC, Fields AM, Fu MM. Emerging cellular themes in leukodystrophies. *Front Cell Dev Biol.* 2022;10:902261.

20. Paloneva J, Manninen T, Christman G, et al. Mutations in two genes encoding different subunits of a receptor signaling complex result in an identical disease phenotype. *Am J Hum Genet.* 2002;71(3):656-662.

21. Guerreiro R, Wojtas A, Bras J, et al. TREM2 variants in Alzheimer's disease. *N Engl J Med.* 2013;368(2):117-127.

22. Sassi C, Nalls MA, Ridge PG, et al. Mendelian adult-onset leukodystrophy genes in Alzheimer's disease: critical influence of CSF1R and NOTCH3. *Neurobiol Aging.* 2018;66:179.e17-e179.e29.

23. Brickman AM. Contemplating Alzheimer's disease and the contribution of white matter hyperintensities. *Curr Neurol Neurosci Rep.* 2013;13(12):415.

24. Dean DC III, Sojkova J, Hurley S, et al. P4-264: Amyloid pathology is associated with

extensive myelin alteration in preclinical Alzheimer's disease: New insights into disease process with novel brain imaging. *Alzheimers Dement.* 2015;11(7S_Part_19):P885-P885.

25. Depp C, Sun T, Sasmita AO, et al. Myelin dysfunction drives amyloid-β deposition in models of Alzheimer's disease. *Nature.* 2023;618(7964):349-357.

26. Graham AC, Bellou E, Harwood JC, et al. Human longevity and Alzheimer's disease variants act via microglia and oligodendrocyte gene networks. *Brain.* 2025;148(3):969-984.

27. Semick SA, Bharadwaj RA, Collado-Torres L, et al. Integrated DNA methylation and gene expression profiling across multiple brain regions implicate novel genes in Alzheimer's disease. *Acta Neuropathol.* 2019;137(4):557-569.

28. Murthy M, Cheng YY, Holton JL, Bettencourt C. Neurodegenerative movement disorders: An epigenetics perspective and promise for the future. *Neuropathol Appl Neurobiol.* 2021;47(7):897-909.

29. Smith RG, Pishva E, Shireby G, et al. A meta-analysis of epigenome-wide association studies in Alzheimer's disease highlights novel differentially methylated loci across cortex. *Nature Communications 2021 12:1.* 2021;12(1):1-13.

30. De Jager PL, Srivastava G, Lunnon K, et al. Alzheimer's disease: early alterations in brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. *Nat Neurosci.* 2014;17(9):1156-1163.

31. Lunnon K, Smith R, Hannon E, et al. Methylomic profiling implicates cortical deregulation of ANK1 in Alzheimer's disease. *Nat Neurosci.* 2014;17(9):1164-1170.

32. Smith RG, Hannon E, De Jager PL, et al. Elevated DNA methylation across a 48-kb region spanning the HOXA gene cluster is associated with Alzheimer's disease neuropathology. *Alzheimers Dement.* 2018;14(12):1580-1588.

33. Watson CT, Roussos P, Garg P, et al. Genome-wide DNA methylation profiling in the superior temporal gyrus reveals epigenetic signatures associated with Alzheimer's disease. *Genome Med.* 2016;8(1). doi:10.1186/S13073-015-0258-8

34. Shireby G, Dempster E, Policicchio S, et al. DNA methylation signatures of Alzheimer's disease neuropathology in the cortex are primarily driven by variation in non-neuronal cell-

types. *bioRxivorg*. Published online 2022:2022.03.15.484508-2022.03.15.484508.

35. Smith AR, Wheildon G, Lunnon K. Invited Review - A 5-year update on epigenome-wide association studies of DNA modifications in Alzheimer's disease: progress, practicalities and promise. *Neuropathology and applied neurobiology*. 2020;46(7):641-653.

36. Zhang L, Silva TC, Young JI, et al. Epigenome-wide meta-analysis of DNA methylation differences in prefrontal cortex implicates the immune processes in Alzheimer's disease. *Nature Communications 2020 11:1*. 2020;11(1):1-13.

37. Gasparoni G, Bultmann S, Lutsik P, et al. DNA methylation analysis on purified neurons and glia dissects age and Alzheimer's disease-specific changes in the human cortex. *Epigenetics Chromatin*. 2018;11(1). doi:10.1186/S13072-018-0211-3

38. Bettencourt C, Foti SC, Miki Y, et al. White matter DNA methylation profiling reveals deregulation of HIP1, LMAN2, MOBP, and other loci in multiple system atrophy. *Acta Neuropathologica*. 2020;139(1):135-135.

39. Aryee MJ, Jaffe AE, Corrada-Bravo H, et al. Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics*. Published online 2014. doi:10.1093/bioinformatics/btu049

40. Schalkwyk LC, Pidsley R, Wong CCY. wateRmelon: Illumina 450 methylation array normalization and metrics. *R package version 122*. Published online 2013.

41. Tian Y, Morris TJ, Webster AP, et al. ChAMP: Updated methylation analysis pipeline for Illumina BeadChips. *Bioinformatics*. Published online 2017. doi:10.1093/bioinformatics/btx513

42. Du P, Zhang X, Huang CC, et al. Comparison of Beta-value and M-value methods for quantifying methylation levels by microarray analysis. *BMC Bioinformatics*. 2010;11. doi:10.1186/1471-2105-11-587

43. Fodder K, Murthy M, Rizzu P, et al. Brain DNA methylomic analysis of frontotemporal lobar degeneration reveals OTUD4 in shared dysregulated signatures across pathological subtypes. *Acta Neuropathologica 2023*. 2023;1:1-19.

44. Mathys H, Davila-Velderrain J, Peng Z, et al. Single-cell transcriptomic analysis of

Alzheimer's disease. *Nature.* doi:10.1038/s41586-019-1195-2

45. Salih DA, Bayram S, Guelfi S, et al. Genetic variability in response to amyloid beta deposition influences Alzheimer's disease risk. *Brain Commun.* 2019;1(1):fcz022.

46. Matarin M, Salih DA, Yasvoina M, et al. A genome-wide gene-expression analysis and database in transgenic mice during development of amyloid or tau pathology. *Cell Rep.* 2015;10(4):633-644.

47. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods.* 2017;14(4):417-419.

48. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000Res.* 2015;4:1521.

49. Langfelder P, Horvath S. WGCNA: An R package for weighted correlation network analysis. *BMC Bioinformatics.* Published online 2008. doi:10.1186/1471-2105-9-559

50. Botía JA, Vandrovcova J, Forabosco P, et al. An additional k-means clustering step improves the biological features of WGCNA gene co-expression networks. *BMC Syst Biol.* Published online 2017. doi:10.1186/s12918-017-0420-6

51. Zeisel A, Muñoz-Manchado AB, Codeluppi S, et al. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science.* 2015;347(6226):1138-1142.

52. Skene NG, Grant SGN. Identification of Vulnerable Cell Types in Major Brain Disorders Using Single Cell Transcriptomes and Expression Weighted Cell Type Enrichment. *Front Neurosci.* 2016;0(JAN):16-16.

53. Piras IS, Huentelman MJ, Pinna F, et al. A review and meta-analysis of gene expression profiles in suicide. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology.* 2022;56. doi:10.1016/j.euroneuro.2021.12.003

54. Morabito S, Reese F, Rahimzadeh N, Miyoshi E, Swarup V. hdWGCNA identifies co-expression networks in high-dimensional transcriptomics data. *Cell Rep Methods.* 2023;3(6):100498.

55. Langfelder P, Luo R, Oldham MC, Horvath S. Is my network module preserved and reproducible? *PLoS Comput Biol.* 2011;7(1):e1001057.

56. Menden K, Francescatto M, Nyima T, et al. A multi-omics dataset for the analysis of Frontotemporal Dementia genetic subtypes. *bioRxiv*. Published online December 2, 2020. doi:10.1101/2020.12.01.405894

57. Weber A, Schwarz SC, Tost J, et al. Epigenome-wide DNA methylation profiling in Progressive Supranuclear Palsy reveals major changes at DLX1. *Nat Commun*. Published online 2018. doi:10.1038/s41467-018-05325-y

58. Murthy M, Fodder K, Miki Y, et al. DNA methylation patterns in the frontal lobe white matter of multiple system atrophy, Parkinson's disease, and progressive supranuclear palsy: a cross-comparative investigation. *Acta Neuropathol*. 2024;148(1):4.

59. de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput Biol*. 2015;11(4):e1004219.

60. Peschl P, Bradl M, Höftberger R, Berger T, Reindl M. Myelin oligodendrocyte glycoprotein: Deciphering a target in inflammatory demyelinating diseases. *Front Immunol*. 2017;8:529.

61. Tiane A, Schepers M, Reijnders RA, et al. From methylation to myelination: epigenomic and transcriptomic profiling of chronic inactive demyelinated multiple sclerosis lesions. *Acta Neuropathol*. 2023;146(2):283-299.

62. Segawa K, Kikuchi A, Noji T, et al. A sublethal ATP11A mutation associated with neurological deterioration causes aberrant phosphatidylcholine flipping in plasma membranes. *J Clin Invest*. 2021;131(18). doi:10.1172/JCI148005

63. Koenning M, Jackson S, Hay CM, et al. Myelin gene regulatory factor is required for maintenance of myelin and mature oligodendrocyte identity in the adult CNS. *J Neurosci*. 2012;32(36):12528-12542.

64. McKenzie AT, Moyon S, Wang M, et al. Multiscale network modeling of oligodendrocytes reveals molecular components of myelin dysregulation in Alzheimer's disease. *Mol Neurodegener*. 2017;12(1):82.

65. Yan T, Liang J, Gao J, et al. FAM222A encodes a protein which accumulates in plaques in Alzheimer's disease. *Nat Commun*. 2020;11(1):411.

66. Liang J, LaFleur B, Hussainy S, Perry G. Gene co-expression analysis of multiple brain

tissues reveals correlation of FAM222A expression with multiple Alzheimer's disease-related genes. *J Alzheimers Dis*. 2024;99(s2):S249-S263.

67. Yang X, Han H, De Carvalho DD, Lay FD, Jones PA, Liang G. Gene body methylation can alter gene expression and is a therapeutic target in cancer. *Cancer Cell*. 2014;26(4):577-590.

68. Bright A, Akay LA, Blanchard J, Tsai LH. Emerging connections between myelin and Alzheimer's disease. *Nat Cell Biol*. Published online July 1, 2025:1-3.

69. Huang H, Zhou F, Zhou S, Qiu M. MYRF: A mysterious membrane-bound transcription factor involved in myelin development and human diseases. *Neurosci Bull*. 2021;37(6):881-884.

70. Hornig J, Fröb F, Vogl MR, Hermans-Borgmeyer I, Tamm ER, Wegner M. The transcription factors Sox10 and Myrf define an essential regulatory network module in differentiating oligodendrocytes. *PLoS Genet*. 2013;9(10):e1003907.

71. Blanchard JW, Akay LA, Davila-Velderrain J, et al. APOE4 impairs myelination via cholesterol dysregulation in oligodendrocytes. *Nature 2022*. Published online November 2022:1-11.

72. Cioffi F, Adam RHI, Bansal R, Broersen K. A review of oxidative stress products and related genes in early Alzheimer's disease. *J Alzheimers Dis*. 2021;83(3):977-1001.

73. Zhang M, Wang J, Zhang K, et al. Ten-eleven translocation 1 mediated-DNA hydroxymethylation is required for myelination and remyelination in the mouse brain. *Nat Commun*. 2021;12(1):5091.

74. Vanzulli I, Papanikolaou M, De-La-Rocha IC, et al. Disruption of oligodendrocyte progenitor cells is an early sign of pathology in the triple transgenic mouse model of Alzheimer's disease. *Neurobiol Aging*. 2020;94:130-139.

75. Planche V, Manjon JV, Mansencal B, et al. Structural progression of Alzheimer's disease over decades: the MRI staging scheme. *Brain Commun*. 2022;4(3):fcac109.