Brain morphology network alterations in adolescents with autism spectrum disorder: a sex-stratified study

Nooshin Safari[1], Isaac Levy[1], Aiying Zhang[1], Chirag Agarwal[1],
Jesus M. Cortes[2,3,4], Kevin A. Pelphrey[5],
John Darrell Van Horn[1,6], Javier Rasero[1]

[1]School of Data Science, University of Virginia, Charlottesville, VA, USA.
[2]Computational Neuroimaging Laboratory, BioBizkaia Health Research Institute, Barakaldo, Bizkaia, Spain.
[3]Ikerbasque, Basque Foundation for Science, VA, USA.
[4]Department of Cell Biology and Histology, University of the Basque Country (UPV/EHU), Leioa, Bizkaia, Spain.
[5]Department of Neurology, University of Virginia, Charlottesville, VA, USA.
[6]Department of Psychology, University of Virginia, Charlottesville, VA, USA.

Contributing authors: rvz7wf@virginia.edu; gbz6qn@virginia.edu; xsa5db@virginia.edu; qze3wn@virginia.edu; jesus.m.cortes@gmail.com; KAP3N@uvahealth.org; jdv7g@virginia.edu; javier.rasero@virginia.edu;

Background: One of the most persistent questions in autism research is why males are more consistently diagnosed than females. Neuroimaging studies have sought to understand this disparity by examining sex differences, primarily through functional and structural connectivity. However, much less is known about how brain networks are organized in autism from a morphological perspective, and how this organization may help explain its sex-related characteristics. This study aimed to elucidate sex-specific alterations in morphology-based connectivity among adolescents with autism spectrum disorder (ASD).

Methods: T1-weighted MRI scans from 337 individuals (181 with ASD; 162 females in total), aged 8–18 years, were preprocessed using FreeSurfer to extract five morphological features—thickness, surface area, sulcal depth, mean curvature, and gray matter volume—across 68 cortical regions. Connectivity was estimated via Kullback–Leibler divergence between region-wise multivariate feature distributions. Network-Based Statistics was applied within sex-stratified models to identify clusters of significantly altered connections in ASD relative to controls (individual edge threshold p = 0.005; FWE-corrected p = 0.05; 5,000 permutations). An exploratory post hoc analysis using partial correlations further associated identified sex-specific network profiles with behavior and comorbidity.

Results: ASD males showed significantly increased connectivity in a cluster centered on the fusiform gyrus, along with the medial orbitofrontal, entorhinal, and parahippocampal cortices. This pattern was positively correlated with restricted and repetitive behaviors (r = 0.22, 95% CI [0.01, 0.41]). In females with ASD, increased connectivity was found in a sub-network implicating mainly the entorhinal cortex, followed by the inferior parietal lobule, and lateral occipital cortex. Notably, the fusiform gyrus was disrupted yet in the opposite direction, exhibiting decreased connectivity with the superior temporal sulcus. These alterations were negatively associated with communication skills (r = –0.25, 95% CI [–0.45, –0.03]). No overlap between male- and female-specific profiles was found, highlighting their distinctiveness.

Limitations: Our cross-sectional sample does not allow for neurodevelopmental inference. Sex-stratified modeling precluded sex-by-diagnosis testing.

Conclusions: Males and females with ASD show non-overlapping profiles of altered morphology-based connectivity, each associated with different behavioral traits, which may contribute to better characterize sex differences in autism.

Keywords: autism spectrum disorder; sex differences; morphological networks; adolescent neuroimaging; cortical features; comorbidity; multivariate analysis

Introduction

Autism Spectrum Disorder (ASD) is a neurodevelopmental condition defined by persistent impairments in social communication alongside restricted and repetitive behaviors, currently affecting approximately one in 36 children in the United States, with a male-to-female diagnosis ratio ranging from 3:1 to 4:1 [1, 2]. The basis of this sex bias is still under debate. Some theories suggest that ASD is characterized by a cognitive trait dominated by systemizing rather than empathizing. Systemizing is more commonly observed in males, which may explain the higher prevalence of autism in this sex. Under this view, females are at the opposite end of the cognitive profile due to their generally stronger ability for empathizing [3]. Another compelling theory is the Female Protective Effect, which posits that females are more resilient to developing autism and require a greater etiological load to reach diagnosis [4]. If these theories are true, they should be at least partially reflected in the brain through sex-specific neural correlates. To examine this, sex should be treated as a stratifying variable rather than a nuisance covariate [5, 6].

Much of our understanding of the neurobiology underlying sex-specificity in autism has emerged from neuroimaging efforts evaluating disruptions in brain architecture. From a functional perspective, females with ASD typically show a trend toward network-wide hyperconnectivity relative to sex-matched typically developing controls (TDC), while males tend toward hypoconnectivity [7]. At a more local level, these patterns can be more nuanced or even reversed [8]. In particular, functional disruptions in females with ASD are predominantly found in somatomotor regions, while males exhibit alterations in both the default mode and somatomotor networks—supporting the so-called Gender Incoherence model [9]. Additional differences in visual, language, and basal ganglia connectivity have also been reported among autistic females [10]. Similar sex disparities have been found in structural connectivity. In preschool-aged children, males with ASD show reduced callosal projections to the orbitofrontal cortex, whereas females exhibit reductions targeting anterior frontal areas, along with elevated diffusivity metrics [11]. During adolescence, females with ASD—but not males—show reduced fiber density and cross-section in callosal pathways [12]. Region-specific reductions in connectivity involving temporal, parietal, and medial posterior cortices have also been reported in females, with no corresponding effects in males [13].

Beyond function and structure, brain morphology has also contributed to further characterizing sex-specific differences in autism. Recent findings indicate that autistic girls exhibit thicker cortices in early development, followed by accelerated thinning relative to autistic boys, suggesting a sex-linked protective mechanism [14]. Overall, morphological features (e.g. cortical thickness, surface area, sulcal depth, mean curvature, and volume) are thought to reflect the cumulative outcome of developmental and genetic processes, positioning them as important markers for disentangling heterogeneity in ASD [15]. Yet, most of this work has focused on region-by-region analyses, typically examining features independently across predefined brain areas, which may be insufficient to capture the spatial interdependencies in brain morphology that sustain neurodevelopment [16].

One promising approach to overcome this limitation is to also adopt a connectivity perspective on brain morphology and examine the similarity in structural features (e.g., cortical thickness, surface area) between brain regions. Most efforts in this direction have employed structural covariance networks, which model connectivity as inter-regional covariation in a single morphometric property across individuals [17, 18]. In the context of autism, studies have reported widespread alterations in connectivity based on region-wise volumetric covariation, with some of these patterns linked to socio-cognitive differences [19, 20]. A recent study further identified significant sex-by-diagnosis interactions in this type of networks within major brain systems, showing that ASD males had increased covariance relative to typically developing males, while ASD females displayed reduced covariance compared to female controls [21]. Despite these findings, whole-brain, sex-specific alterations in morphology-based connectivity in autism remain largely underexplored. Moreover, covariation network approaches are typically limited to a single morphological property and may overlook the benefits of integrating multiple features into a unified framework to provide a more comprehensive description of group and individual differences in cortical organization [22, 23].

To fill these gaps, the present study set out to characterize sex-specific connectivity patterns based on multivariate morphological information in individuals with autism during typical developmental ages. Following similar observations in functional and structural connectivity, we hypothesized that males and females with ASD would show distinct morphological network profiles. To this end, we leveraged a large-scale, multi-site dataset comprising 337 youth participants (aged 8–18 years), balanced by sex (162 females) and diagnosis (181 with ASD). Adopting a recently developed framework for robustly estimating similarity networks from T1-weighted images [24], morphology-based connectivity was computed from multivariate patterns of cortical features, including thickness, mean curvature, surface area, sulcal depth, and gray matter volume. Clusters of statistically disrupted connections in ASD relative to TDC were then identified separately in males and females, and their associations with behavioral and comorbidity traits were further evaluated.

Materials and Methods

Participants

337 individuals from Wave 1 of NIH Autism Centers of Excellence (ACE) Network were selected for this study after quality control assurance, involving visual inspection of segmentation output and exclusion of scans with structural preprocessing failures (see [25] for more details on the ACE network). The study sample included 181 subjects with ASD (Age: 12.73 ± 2.92 years, 84 female) and 156 with TDC (Age: 13.20 ± 3.01 years, 78 female), who were collected across four sites: Harvard University, Seattle Children's/University of Washington, University of California Los Angeles, and Yale University. Additionally, several scores were used to assess the association of findings with behavioral and comorbidity traits within individuals with autism. Specifically, we extracted scores for Communication, Social Interaction, and Restricted, Repetitive, and Stereotyped Behaviors from the Autism Diagnostic Interview–Revised (ADI-R); Social Affect and Restricted and Repetitive Behaviors from the Autism Diagnostic Observation Schedule—2nd Edition (ADOS-2); and subscale scores for Anxiety, Affective Problems, Attention Deficit/Hyperactivity, Oppositional Defiant, Conduct, and Internalizing Problems from the Child Behavior Checklist (CBCL). Descriptive statistics for our sample can be found in Table 1.

MRI Acquisition and Preprocessing

Structural images were acquired using a high-resolution T1-weighted MPRAGE sequence (FOV: 176×256×256; voxel size: $1 \times 1 \times 1 \, \mathrm{mm}^3$). Preprocessing of these images were performed with FreeSurfer's `recon-all` pipeline (v6.0), which involves cortical surface reconstruction and subcortical segmentation. Additionally, gray matter volume maps were generated using a standard voxel-based morphometry pipeline implemented in CAT12 [26], but adapted for FreeSurfer outputs. This included spatial normalization, modulation, and smoothing of gray matter probability maps. All images were manually reviewed to ensure accurate surface reconstruction, parcellation, and segmentation.

Variable	Male		Female	
	ASD	TDC	ASD	TDC
N	97	78	84	78
Age (months)	152.36 [145.02–159.69]	161.02 [153.47–168.57]	153.34 [146.03–160.65]	155.82 [147.10–164.55]
Site (HAR/SCR/UCL/YAL)	24/28/30/15	11/23/26/18	11/25/25/23	13/25/23/17
ICV ($10^6 \times$ mm^3)	1.63 [1.60–1.66]	1.64 [1.61–1.67]	1.47 [1.44–1.50]	1.48 [1.45–1.51]
ADI-R Social	19.38 [18.28–20.49]	—	18.11 [16.74–19.48]	
ADI-R Communication	16.54 [15.67–17.42]	—	15.38 [14.37–16.40]	—
ADI-R Behavioral	6.63 [6.12–7.14]	—	5.70 [5.13–6.28]	—
ADOS-2 Social Affect	9.52 [8.75–10.28]	—	8.06 [7.40–8.72]	—
ADOS-2 Communication	2.63 [2.29–2.97]	—	2.54 [2.15–2.93]	—
CBCL Affective	61.07 [59.28–62.85]	51.60 [50.86–52.33]	64.19 [61.93–66.46]	52.95 [51.87–54.02]
CBCL Anxiety	61.74 [60.13–63.36]	51.90 [51.06–52.73]	62.49 [60.56–64.43]	52.04 [51.14–52.94]
CBCL ADHD	61.92 [60.37–63.48]	50.51 [49.00–52.01]	63.38 [61.47–65.28]	51.18 [50.59–51.77]
CBCL Oppositional	58.65 [57.04–60.26]	51.62 [50.96–52.28]	58.38 [56.67–60.09]	51.53 [50.94–52.13]
CBCL Conduct	56.17 [54.70–57.63]	50.81 [50.32–51.29]	57.33 [55.56–59.10]	50.86 [50.38–51.35]
CBCL Internalizing	60.87 [59.15–62.58]	43.81 [41.66–45.95]	63.04 [60.48–65.60]	46.47 [44.31–48.63]

Table 1: **Demographic, behavioral and comorbidity data stratified by sex and diagnostic group.** Values represent means and 95% confidence intervals. Dashes (—) indicate unavailable scores for TDC participants. HAR: Harvard University; SCR: Seattle Children's/University of Washington; ICV: intracranial volume

Morphology-based Connectivity

Morphology-based connectivity matrices were constructed for all subjects using the Morphometric INverse Divergence framework [24], which estimates pairwise similarity between brain regions based on the symmetrized Kullback–Leibler (KL) divergence of vertex-wise morphometric feature distributions. Main analyses concentrated on cortical networks, where multiple morphological properties could be defined. Additional analyses were performed on whole-brain networks, but these were based solely on volumetric information, since this was the only morphological measure available for subcortical regions.

Specifically, in order to build the connectivity matrices at the cortical level, thickness, mean curvature, surface area, sulcal depth, and gray matter volume were first extracted from the preprocessed structural images and then sampled at the vertex level

within each region of the Desikan–Killiany atlas (68 regions; 34 per hemisphere). Since each region contains many vertices, this procedure yielded a distribution of values for each morphological property within each ROI. These distributions were then modeled with kernel density estimation and jointly used to compute the similarity between pairs of regions via a symmetrized multivariate KL divergence. This resulted in a 68×68 connectivity matrix for each individual, representing the similarity between regions based on their multivariate morphological signatures (See Supplementary Fig. S1 for an overview of the whole connectivity calculation procedure).

For the whole-brain connectivity matrices, 14 subcortical structures obtained from FreeSurfer's *aseg.mgz* segmentation file [27] were added to the aforementioned cortical regions from the the Desikan–Killiany atlas. These subcortical regions consisted of the bilateral thalamus, caudate, putamen, pallidum, hippocampus, amygdala, and nucleus accumbens. An 82×82 connectivity matrix was then built for each individual as the univariate KL divergence between region-wise gray matter volume distributions, extracted from the voxel values within each ROI after the voxel-based morphometry pipeline.

Sex-specific Network Alterations

To identify statistically significant differences in morphology-based connectivity, we applied the Network-Based Statistics (NBS) approach [28], which is analogous to cluster-wise thresholding of statistical brain maps but adapted for network data. For each edge in the connectivity matrix, a test statistic was first computed, followed by an individual threshold to identify sub-networks of suprathreshold connected edges. The significance of each sub-network was then assessed using permutation testing while controlling the family-wise error rate. A two-sided linear regression model was employed to calculate the test statistic at each edge, therefore allowing the existence of significant clusters of both increased and decreased connectivity in ASD relative to TDC. For interpretability, the resulting edges were subsequently separated based on the sign of the observed test statistic. This analysis was implemented using the *lm_nbr* function from the NBR R package [29].

The current study concentrated on **sex-specific** network alterations. As a result, NBS was applied to the connectivity matrices within each sex sub-population (Male and Female) with the diagnosis variable (ASD vs TDC) as the main effect, while controlling for covariation in age, intracranial volume and collection site. Of note, by stratifying our study by sex multi-collinearity issues due to the marked differences in intracranial volume between males and females (see Table 1) were avoided. Main results were reported for NBS with a 5000 total permutations, an individual edge threshold $p = 0.005$ and family-wise error correction ($p_{\text{FWE}} = 0.05$). In NBS, the individual edge threshold is a hyperparameter that determines the potential size of clusters. We chose $p = 0.005$ since it appeared to provide a compromise in cluster size, sufficient for discovery while preserving statistical power [28]. To assess robustness, we also tested the sensitivity of the findings to a more liberal ($p = 0.01$) and stringent threshold ($p = 0.001$).

Importance of Morphological Properties

The relative importance of each morphological property in distinguishing individuals with autism from typically developing controls was assessed by combining the strength of each sub-network identified by NBS with a leave-one-feature-out procedure within a ROC curve analysis framework. Strength was calculated for each individual's connectivity matrix as the sum of values corresponding to the altered sub-networks and was selected because it is the main statistic in NBS. This metric reflects the overall magnitude of morphological similarity within the significant sub-networks and provides a compact summary of network-level disruption. The leave-one-feature-out procedure removed a given morphological property prior to constructing the connectivity matrices via multivariate KL divergence, thereby changing the strength of the sub-networks too. The impact on classification performance was then quantified by examining the change in the area under the ROC curve (AUC) using strength as the discriminating variable. This procedure was repeated for each morphological property individually.

Association with Behavior and Comorbidity

In a post-hoc analysis, the association of the obtained network findings with behavioral and comorbidity symptoms within the autistic group was also tested. Specifically, for each significantly altered pattern and for each contrast identified by the NBS analysis (i.e., ASD > TDC and TDC > ASD), sub-network strength was correlated with the extracted scores from ADI-R, ADOS-2, and CBCL. To maintain consistency with the covariates used in the primary group-level network analysis, a partial correlation approach was adopted, in which the effects of age, site, and intracranial volume were regressed out from both variables involved in the correlation. Correlation coefficients, 95% confidence intervals, and corresponding p-values were reported. Given the exploratory nature of this analysis, p-values were not corrected for multiple comparisons. As such, these results should be interpreted with caution and considered hypothesis-generating rather than conclusive.

Results

ASD Males vs. TDC Males

Group-level network analysis identified a single cluster of 35 significantly altered connections ($p_{\text{FWE}} = 0.037$). As shown in Fig 1A, most of these corresponded to increased morphology-based connectivity in ASD relative to TDC, with a pattern prominently involving the left hemisphere fusiform and medial orbital frontal gyri, as well as the right hemisphere entorhinal cortex and lingual gyrus. This cluster also included a single connection showing decreased connectivity in ASD compared to TDC, linking the left middle temporal gyrus and the right inferior temporal gyrus. This analysis was performed using the individual edge threshold of $p = 0.005$ and 5000 permutations for null-modeling. Additional profiles with other individual edge thresholds in NBS were also reported (see Supplementary Fig. S2). Remarkably, the fusiform continued to exhibit significantly decreased connectivity even under the most stringent threshold

Fig. 1: **Main results in the male cohort**. **A)** Circular plots visualizing clusters of significant inter-regional connections. Nodes represent cortical regions, with olive and teal indicating the left and right hemispheres, respectively. On the left, increased connectivity in TDC relative to ASD; on the right, the opposite. Darker colors denote stronger group differences. **B)** ROC curves and area under the curve (AUC) values showing performance in distinguishing ASD from TDC using the strength of network profiles in **A**. Each curve represents the scenario in which one morphological property is omitted in the computation of the connectivity matrices (CT: cortical thickness, SA: surface area, SD: sulcus depth, Vol: volume, MC: mean curvature). **C)** Forest plot displaying partial correlation coefficients and 95% confidence intervals between the network strength of pattern in **A** and behavioral and comorbidity measures.

($p = 0.001$). No significantly altered networks at the whole-brain level using volumetric information only was found at any of the thresholds.

The most influential morphological property for distinguishing between controls and individuals with autism within this cluster appeared to be cortical thickness, with the AUC dropping from 0.72 (when all features were included) to 0.54 when this feature was removed (see Fig. 1B). This was followed by sulcal depth (AUC = 0.68)

and mean curvature, although in the latter case the reduction in group separability was minimal (AUC = 0.71). In contrast, surface area and volumetric properties did not appear to be individually necessary, as removing either of them resulted in slightly improved group separability (AUC = 0.72 in both cases).

With respect to behavior and comorbidity (see Fig. 1C), the strongest correlations with the single connection of increased connectivity in TDC relative to ASD were found for Anxiety (r = 0.17, 95% CI = [–0.05, 0.37], p = 0.13) and Affective problems (r = 0.16, 95% CI = [–0.06, 0.36], p = 0.16), based on CBCL scores. Interestingly, the major sub-network comprising significantly increased connections in the ASD group compared to TDC showed a moderate positive correlation with restricted, repetitive, and stereotyped behaviors measured by the ADI-R (r = 0.22, 95% CI = [0.01, 0.41], p = 0.037), although this result would not survive correction for multiple comparisons and should therefore be interpreted with caution. Other behavioral measures exhibited weaker correlations, with confidence intervals that included zero.

ASD Females vs. TDC Females

Within the female subgroup, a single cluster comprising 41 significantly altered morphology-based connections (p_{FWE} = 0.028) was identified. Compared to TDC, patterns of increased connectivity in the ASD group were centered around a prominent hub in the bilateral entorhinal cortex, which then extended to the bilateral inferior parietal lobule, lateral occipital cortex, and pre- and postcentral gyri (see Figure 2A, right side), all regions implicated in visuospatial integration, somatosensory encoding, and mnemonic retrieval. In contrast, decreased connectivity, although more scarce, prominently involved the bank of the superior temporal sulcus engaging with a few other regions, among which the fusiform gyrus was found (see Figure 2A, left side). All these alterations remained consistent—yet spatially circumscribed, more notoriously to the left hemisphere enthorinal cortex—at the more stringent individual edge threshold in NBS (p = 0.001). At the more liberal p = 0.01 level, the pattern expanded to include bilateral precentral extensions and deeper parietal-occipital couplings (see Supplementary Fig. S3), suggesting a core-periphery organization sensitive to statistical stringency. Similar to males, no significantly altered networks were found at any of the thresholds at the whole-brain level using only volumetric information.

In terms of group separability, a similar pattern to that observed in the male cohort was found, to a large extent. As shown in Fig. 2B, using all morphological features together yielded an AUC of 0.66. Removing cortical thickness led to the largest reduction in group separability (AUC = 0.58), confirming its key role. In contrast, removing surface area had no effect (AUC = 0.66), while sulcal depth had only a minimal impact in this case (AUC = 0.65). Volumetric properties and mean curvature appeared to be non-essential individually, as their removal slightly increased the AUC to 0.67 in both cases.

Fig. 2C displays a moderate negative correlation between the sub-network of decreased connectivity in ASD relative to TDC, which primarily involved connections to the right hemisphere banks of the superior temporal sulcus, and communication skills, as measured by the ADI-R (r = –0.25, 95% CI = [–0.45, –0.03], p = 0.027). Even

Fig. 2: **Main results in the female cohort**. **A)** Circular plots visualizing clusters of significant inter-regional connections. Nodes represent cortical regions, with olive and teal indicating the left and right hemispheres, respectively. On the left, increased connectivity in TDC relative to ASD; on the right, the opposite. Darker colors denote stronger group differences. **B)** ROC curves and area under the curve (AUC) values showing performance in distinguishing ASD from TDC using the strength of network profiles in **A**. Each curve represents the scenario in which one morphological property is omitted in the computation of the connectivity matrices (CT: cortical thickness, SA: surface area, SD: sulcus depth, Vol: volume, MC: mean curvature). **C)** Forest plot displaying partial correlation coefficients and 95% confidence intervals between the network strength of pattern in **A** and behavioral and comorbidity measures.

though other CBCL subscales exhibited small-to-moderate correlations (e.g. oppositional defiant problems: $r = 0.15$, 95% CI $= [-0.09, 0.37]$, $p = 0.21$), these were not statistically significant. Likewise, none of the correlations with the sub-network of increased connectivity in ASD relative to TDC reached significance, with the strongest

being anxiety problems from CBCL ($r = 0.18$, 95% CI = [–0.06, 0.39], $p = 0.14$). Similar to the male cohort, these findings should be interpreted with caution as not being corrected for multiple comparisons.

Comparing Network Profiles

We conducted an overlap analysis to assess the distinctiveness of the network profiles identified across sexes. For each group comparison (i.e., ASD males vs. TDC males, ASD females vs. TDC females), we first constructed a binary adjacency matrix in which an entry was set to 1 if the corresponding edge was found to be significantly altered by NBS. We then computed the Jaccard index (J) defined as the number of intersecting positive cases divided by the size of their union, i.e., $J = \frac{|A \cap B|}{|A \cup B|}$. At the individual edge threshold of $p = 0.005$ and cluster-level family-wise error correction ($p_{\mathrm{FWE}} = 0.05$), this yielded a Jaccard index of $J = 0$ between the male and female network profiles, indicating no overlapping edges across the sex-specific contrasts.

To further evaluate whether these profiles reflected true diagnostic group effects rather than normative sex differences, we applied the same NBS setup of our main analysis (individual edge threshold $p = 0.005$, cluster-level family-wise error correction $p_{\mathrm{FWE}} = 0.05$, 5000 permutations, and identical covariates) to compare TDC males with TDC females. This analysis revealed a cluster of 48 significantly altered connections, which are depicted by directionality in Supplementary Fig. S4. When overlapping this normative sex-difference network with the diagnostic-related profiles, we found a Jaccard index of $J = 0$ for ASD males vs TDC males, and $J = 0.01$ for ASD females vs TDC females. In the latter case, only a single overlapping connection was found, linking the left rostral middle frontal gyrus and the right superior temporal gyrus. These minimal overlaps suggest that the reported sex-specific morphological network profiles arise from diagnosis-related alterations rather than normative sex differences.

Discussion

This study aimed to delineate sex-specific network profiles based on altered brain morphology-based connectivity in adolescents with autism. To accomplish this, we leveraged a large-scale, multi-study dataset comprising 337 individuals, balanced by sex and diagnostic group. For each participant's T1-weighted image, we extracted vertex-level distributions of cortical morphometric properties across 68 brain regions, including cortical thickness, mean curvature, surface area, sulcal depth, and gray matter volume. We then estimated inter-regional connectivity by quantifying the similarity between these morphometric distributions using a recently proposed approach based on multivariate KL divergence. By means of a network-based statistical framework, we assessed sex-specific alterations in connectivity between ASD and TDC individuals, and found distinct profiles across males and females. Specifically, males with autism showed significantly heightened levels of connectivity in regions prominently involving the fusiform and lingual gyri, anterior cingulate cortex, orbitofrontal cortex, and parietal association cortices, whereas for females, statistical results centered on elevated

connectivity in the entorhinal cortex, inferior parietal lobule, and lateral occipital cortex, as well as a decreased connectivity between the fusiform gyrus and the bank of superior temporal sulcus. Moreover, a post-hoc exploratory analysis showed that these profiles may be linked with a diverse behavioral response, namely, a positive correlation with repetitive and restrictive behavior in males and a negative association with communication skills in females. Our results suggest that examining morphological information from a connectivity perspective can further characterize autism and, importantly, provide novel insights into the sex-specific behavioral differences commonly observed in individuals with this neurodiverse condition.

Our results add to the growing body of evidence reinforcing a network-level perspective for better characterizing sex differences in ASD. In males, we found increased connectivity in regions belonging to the default mode, salience, and frontoparietal networks. One of the central hubs we identified is the fusiform gyrus, a region implicated in social cognition that has been consistently linked to autism through findings of reduced activation during face processing and structural abnormalities associated with social perceptual deficits [30–32]. Other key regions included the parahippocampal gyrus (linked to memory) and the postcentral gyrus (involved in sensorimotor processing). These findings align with theories of network-level developmental plasticity, in which increased long-range integration may reflect delayed synaptic pruning or compensatory adaptations to altered sensory or cognitive demands, as supported by studies using graph-theoretical metrics [33, 34]. Importantly, these areas have also been found to be hyperconnected in fMRI studies using male-predominant cohorts [35–37] and, as we previously demonstrated, contribute to the functional connectivity–based heterogeneity observed in autism [38]. Furthermore, they have also been identified as part of a disrupted structural architecture in ASD, exhibiting reduced fractional anisotropy and aberrant streamline coherence in white matter tracts traversing the cingulum bundle, forceps minor, and superior longitudinal fasciculus [39–41]. Taken together, our findings point to a coherent pattern of multimodal dysconnectivity in males with ASD, centered on frontal, temporal, and occipital integration hubs as key components of ASD neuropathophysiology.

On the other hand, in females with ASD, network disruptions across both functional and structural modalities also appeared to overlap with key regions identified in our morphology-based connectivity profiles. Specifically, resting-state fMRI studies have revealed disruptions within the default-mode network (heightened entorhinal–posterior cingulate; [42]), altered top-down central executive influence—specifically dorsolateral prefrontal to postcentral pathways [43]- and increased coupling between the default-mode and central executive networks [42]. From the perspective of the canonical triple-network model [44], these changes suggest a reconfigured network topology in ASD females, which may reflect compensatory upregulation of executive–memory loops to offset local cortical variability. Additionally, sex-specific salience network alterations—such as elevated connectivity with sensorimotor areas reported in early ASD trajectories (infants at familial risk) [45]—may reflect dynamic sensory reweighting and flexible attention gating across internal–external stimulus axes. Further, reduced network flexibility in executive and salience systems (as quantified via multilayer network analyses) correlates with social–communication symptom

severity in ASD females [46], raising interpretative nuance around whether increased 'integration' denotes an adaptive scaffold or constrained network rigidity. In parallel, DTI studies in ASD females aged 12–30 years have revealed significant reductions in fractional anisotropy within white matter tracts underlying our morphological-based network profiles—including the internal capsule, inferior parietal lobule–lateral occipital cortex association fibers, and posterior corona radiata [47, 48]. In sum, these convergent findings appear to support the presence of a multimodal (morphology, function and structure) hub integration profile that may be specific to the female ASD phenotype.

What, then, do these sex-specific morphological network alterations reveal about the broader neurobiology of autism in light of sex differences? Even though our study did not directly contrast sexes, the involvement of largely distinct regions—and in some cases, with opposite directionality—suggests divergent neural substrates underlying core autistic traits. For example, we found hyperconnectivity of the fusiform gyrus in males but hypoconnectivity in females. Since this region plays a established role in face perception and social cognition [49–51], its increased connectivity in males with autism could mirror a compensatory mechanism or atypical social attention pathways typically found in male-biased ASD studies but at the morphological architecture level. In contrast, the reduced connectivity of the fusiform in females, particularly involving connections with the superior temporal sulcus —another social cognitive hub— may signal a qualitatively different neurodevelopmental trajectory. Interestingly, the enthorinal cortex appears to be more disrupted in females than males. As a central node in memory and spatial navigation [52, 53] this finding may hint at alternative circuitries supporting adaptive behavior. Critically, these highlighted regions are unlikely to be attributed to normative sex differences, as our overlap analysis demonstrated.

We assessed the sensitivity in discriminating utility of each morphological property underlying the obtained sex-specific connectivity patterns using a one-leave-feature-out procedure. Across sexes, the largest drops in diagnostic performance occurred when cortical thickness and sulcus depth were excluded, suggesting that these properties play a particularly sensitive role in the discriminative capacity of morphological networks—or at the very least, offer complementary information relative to the other morphological properties. Cortical thickness is a well-established marker known to be altered in autism and associated with symptom severity [15, 54–56]. Sulcal depth, by contrast, has been less extensively studied in autism, with findings ranging from no statistical differences [57], to specific localized abnormalities [58], and to associations with early cortical folding processes during neurodevelopment [59]. Our study builds on this body of work by highlighting the relevance of both of these morphological properties from a network-level perspective and across sexes. Conversely, removing either surface area or volumetric information from the connectivity calculations led to slight improvements in discriminating utility, which seems to indicate a potentially redundant or less informative contribution from these properties. This can be partially explained by the strong correlation typically observed between surface area and volume at the cortex-level, so very little or none is gained from their combination. Nevertheless, in our analysis based solely on volumetric information, performed to allow the inclusion of the subcortex, we did not find any significant clusters of altered

edges across sexes, reinforcing the limited utility of volumetric features to identify ASD individuals, at least from a network perspective. This appears to contradict prior studies suggesting that brain networks based on volumetric features have high predictive power for classifying ASD individuals [20]. However, this may simply reflect the need for a more distributed morphometric signature, whereas our network-based statistical framework focuses on detecting clusters of significantly altered connections, which might be less efficient for diagnostic classification purposes.

To further contextualize our findings, we also examined the association between the altered networks and core autism-related behavioral scales in autism, such as ADOS-2 and ADI-R. In males with autism, this analysis yielded a small-to-moderate positive correlation with restrictive and repetitive behaviors, and a small-to-moderate negative correlation with communication issues in ASD females. Together, these findings hint at potentially distinct behavioral phenotypes underlying the sex-specific networks. Interestingly, these results appear consistent with commonly hypothesized patterns in autism: Greater alterations in males corresponding to increased symptom severity, while in females, this inverse relationship may reflect a camouflaging effect in social cognition and communication domains [60, 61]. Notwithstanding, these results warrants further exploration as they were not corrected for multiple comparisons due to the exploratory nature of the analysis. No significant associations were found across the rest of behavioral subdomains for either sex. We speculate that this overall null effect may stem from the multifaceted nature of ASD behavior, which may not map onto discrete and localized clusters of altered connectivity but rather arise from more widespread or complex network disruptions across the brain [36, 62, 63]. Likewise, no statistical associations of comorbidity were found, suggesting that the observed morphology-based connectivity alterations are unlikely to manifest general traits of other psychiatric conditions.

Limitations

Several limitations of our study should be acknowledged. First, as consistently stated throughout this document, estimated associations of sex-specific network profiles with behavior and comorbidity were not corrected for multiple comparisons due to the exploratory nature of this analysis. Therefore, future studies should validate these findings. Second, our analysis was stratified by sex to reduce potential multicollinearity arising from sex-specific differences in global morphology. As such, any comparisons across sexes should be interpreted with caution. Future studies with more balanced designs and carefully controlled variables will be needed to directly test for sex-by-diagnosis interaction effects. Third, our data, drawn from Wave 1 of the ACE network, is cross-sectional and therefore precludes any conclusions about the developmental trajectory of our findings. However, subsequent waves (Wave 2 and 3) of the ACE network include follow-up data, which will enable us to test explicitly for longitudinal effects in future avenues. Fourth, our findings were derived using a purely statistical inference approach which, although useful for gaining insight into the morphology-based network architecture characteristic of autism and its sex-specific heterogeneity, does not quantify the utility of these features as biomarkers for diagnostic purposes. Future studies

should adopt a thorough machine learning framework with out-of-sample testing to specifically assess this. Finally, our network results restrict to a single neuroimaging modality, specifically, brain morphology. Hence, their combination with biomarkers from functional and structural connectivity signals may lead to a more comprehensive characterization of the ASD heteroegeneity and improved diagnostic accuracy rates [64]. Future studies should also examine the robustness and generalizability of sex-specific network effects across modalities [65, 66].

Conclusions

Despite the aforementioned limitations, the present study provides evidence for a brain connectivity profile, derived from multivariate morphological signatures, that is expressed differently across sexes in adolescents with autism. This could contribute to improving the identification of autism in female individuals at younger ages, who are typically underdiagnosed. Furthermore, the distinct patterns of observed associations with behavior in males and females, although small, hold promise for informing sex-sensitive models of psychopathology and guiding intervention in autism.

Abbreviations

ASD	Autism Spectrum Disorder
TDC	Typticall Developing Control
ACE	Autism Centers of Excellence
ADOS-2	Autism Diagnostic Observation Schedule—2nd Edition
ADI-R	Autism Diagnostic Interview–Revise
CBCL	Child Behavior Checklist
NBS	Network-Based Statistics
KL	Kullback–Leibler

Ethics approval and consent to participate

The recruitment of participants for this dataset was approved by the Yale Institutional Review Board, the UCLA Office of the Human Research Protection Program, the Boston Children's Hospital Institutional Review Board, the USC Office for the Protection of Research Subjects, and the University of Virginia Institutional Review Board for Health Sciences Research. All procedures were conducted in accordance with the Declaration of Helsinki. Minor participants provided verbal assent and were informed they could withdraw at any point during the experiment. Written informed consent was obtained from a parent or legal guardian of each participating child.

Consent for publication

Not applicable.

Availability of data and materials

Raw neuroimaging data used in this study can be downloaded from the National Institute of Mental Health Data Archive Data (collection #2021). Source files, analysis scripts and figures are available at https://github.com/nooshin23/Autism-sex-stratified-morphology-connectivity.

Competing interests

The authors declare that they have no competing interests.

Funding

NS's work was supported by the UVA School of Data Science. KAP and JDVH were supported by the National Institute of Mental Health (NIMH) Autism Center of Excellence Network Award (R01 MH100028; PI: KAP).

Authors' contributions

All authors contributed to the conceptualization of the project, interpretation of findings, manuscript editing, and approved the final version. KAP led data collection. NS, IL, and JR performed the analysis. NS and JR drafted the manuscript and prepared the figures.

Acknowledgements

Not applicable.

References

[1] Loomes, R., Hull, L., Mandy, W.P.L.: What is the male-to-female ratio in autism spectrum disorder? a systematic review and meta-analysis. Journal of the American Academy of Child & Adolescent Psychiatry **56**(6), 466–474 (2017) https://doi.org/10.1016/j.jaac.2017.03.013

[2] Baio, J., Wiggins, L., Christensen, D.L., Maenner, M.J., Daniels, J., Warren, Z., Kurzius-Spencer, M., Zahorodny, W., Robinson, C., Rosenberg, White, T., Durkin, M.S., Imm, P., Nikolaou, L., Yeargin-Allsopp, M., Lee, L.-C., Harrington, R., Lopez, M., Fitzgerald, R.T., Hewitt, A., Pettygrove, S., Constantino, J.N., Vehorn, A., Shenouda, J., Hall-Lande, J., Van, K., Naarden, Braun, Dowling, N.F.: Prevalence of autism spectrum disorder among children aged 8 years — autism and developmental disabilities monitoring network, 11 sites, united states, 2014. MMWR. Surveillance Summaries **67**(6), 1–23 (2018) https://doi.org/10.15585/mmwr.ss6706a1

[3] Baron-Cohen, S.: The extreme male brain theory of autism. Trends in Cognitive Sciences **6**(6), 248–254 (2002) https://doi.org/10.1016/s1364-6613(02)01904-6

[4] Robinson, E.B., Lichtenstein, P., Anckarsäter, H., Happé, F., Ronald, A.: Examining and interpreting the female protective effect against autistic behavior. Proceedings of the National Academy of Sciences **110**(13), 5258–5262 (2013) https://doi.org/10.1073/pnas.1211070110

[5] Halladay, A.K., Bishop, S., Constantino, J.N., Daniels, A.M., Koenig, K., Palmer, K., Messinger, D., Pelphrey, K., Sanders, S.J., Singer, A.T., Taylor, J.L., Szatmari, P.: Sex and gender differences in autism spectrum disorder: summarizing evidence gaps and identifying emerging areas of priority. Molecular Autism **6**(1) (2015) https://doi.org/10.1186/s13229-015-0019-y

[6] Lai, M.-C., Lombardo, M.V., Auyeung, B., Chakrabarti, B., Baron-Cohen, S.: Sex/gender differences and autism: Setting the scene for future research. Journal of the American Academy of Child & Adolescent Psychiatry **54**(1), 11–24 (2015) https://doi.org/10.1016/j.jaac.2014.10.003

[7] Alaerts, K., Swinnen, S.P., Wenderoth, N.: Sex differences in autism: a resting-state fmri investigation of functional brain connectivity in males and females. Social Cognitive and Affective Neuroscience **11**(6), 1002–1016 (2016) https://doi.org/10.1093/scan/nsw027

[8] Di, X., Biswal, B.B.: Sex-dependent and sex-independent brain resting-state functional connectivity in children with autism spectrum disorder (2016) https://doi.org/10.1101/038026

[9] Floris, D.L., Lai, M.-C., Nath, T., Milham, M.P., Di Martino, A.: Network-specific sex differentiation of intrinsic brain function in males with autism. Molecular Autism **9**(1) (2018) https://doi.org/10.1186/s13229-018-0192-x

[10] Tavares, V., Fernandes, L.A., Antunes, M., Ferreira, H., Prata, D.: Sex differences in functional connectivity between resting state brain networks in autism spectrum disorder. Journal of Autism and Developmental Disorders **52**(7), 3088–3101 (2021) https://doi.org/10.1007/s10803-021-05191-6

[11] Nordahl, C.W., Iosif, A.-M., Young, G.S., Perry, L.M., Dougherty, R., Lee, A., Li, D., Buonocore, M.H., Simon, T., Rogers, S., Wandell, B., Amaral, D.G.: Sex differences in the corpus callosum in preschool-aged children with autism spectrum disorder. Molecular Autism **6**(1) (2015) https://doi.org/10.1186/s13229-015-0005-4

[12] Kirkovski, M., Fuelscher, I., Hyde, C., Donaldson, P.H., Ford, T.C., Rossell, S.L., Fitzgerald, P.B., Enticott, P.G.: Fixel based analysis reveals atypical white matter micro- and macrostructure in adults with autism spectrum disorder: An investigation of the role of biological sex. Frontiers in Integrative Neuroscience **14** (2020) https://doi.org/10.3389/fnint.2020.00040

[13] Irimia, A., Torgerson, C.M., Jacokes, Z.J., Van Horn, J.D.: The connectomes

of males and females with autism spectrum disorder have significantly different white matter connectivity densities. Scientific Reports **7**(1) (2017) https://doi.org/10.1038/srep46401

[14] Andrews, D.S., Diers, K., Lee, J.K., Harvey, D.J., Heath, B., Cordero, D., Rogers, S.J., Reuter, M., Solomon, M., Amaral, D.G., Nordahl, C.W.: Sex differences in trajectories of cortical development in autistic children from 2–13 years of age. Molecular Psychiatry **29**(11), 3440–3451 (2024) https://doi.org/10.1038/s41380-024-02592-8

[15] Yang, D.Y.-J., Beam, D., Pelphrey, K.A., Abdullahi, S., Jou, R.J.: Cortical morphological markers in children with autism: a structural magnetic resonance imaging study of thickness, area, volume, and gyrification. Molecular Autism **7**(1) (2016) https://doi.org/10.1186/s13229-016-0076-x

[16] Bedford, S.A., Park, M.T.M., Devenyi, G.A., Tullo, S., Germann, J., Patel, R., Anagnostou, E., Baron-Cohen, S., Bullmore, E.T., Chura, L.R., Craig, M.C., Ecker, C., Floris, D.L., Holt, R.J., Lenroot, R., Lerch, J.P., Lombardo, M.V., Murphy, D.G.M., Raznahan, A., Ruigrok, A.N.V., Smith, E., Spencer, M.D., Suckling, J., Taylor, M.J., Thurm, A., Lai, M.-C., Chakravarty, M.M.: Large-scale analyses of the relationship between sex, age and intelligence quotient heterogeneity and cortical morphometry in autism spectrum disorder. Molecular Psychiatry **25**(3), 614–628 (2019) https://doi.org/10.1038/s41380-019-0420-6

[17] Tijms, B.M., Series, P., Willshaw, D.J., Lawrie, S.M.: Similarity-based extraction of individual networks from gray matter mri scans. Cerebral Cortex **22**(7), 1530–1541 (2011) https://doi.org/10.1093/cercor/bhr221

[18] Kong, X.-z., Liu, Z., Huang, L., Wang, X., Yang, Z., Zhou, G., Zhen, Z., Liu, J.: Mapping individual brain networks using statistical similarity in regional morphology from mri. PLOS ONE **10**(11), 0141840 (2015) https://doi.org/10.1371/journal.pone.0141840

[19] Cardon, G.J., Hepburn, S., Rojas, D.C.: Structural covariance of sensory networks, the cerebellum, and amygdala in autism spectrum disorder. Frontiers in Neurology **8** (2017) https://doi.org/10.3389/fneur.2017.00615

[20] He, C., Cortes, J.M., Kang, X., Cao, J., Chen, H., Guo, X., Wang, R., Kong, L., Huang, X., Xiao, J., Shan, X., Feng, R., Chen, H., Duan, X.: Individual-based morphological brain network organization and its association with autistic symptoms in young children with autism spectrum disorder. Human Brain Mapping **42**(10), 3282–3294 (2021) https://doi.org/10.1002/hbm.25434

[21] Zielinski, B.A., Andrews, D.S., Lee, J.K., Solomon, M., Rogers, S.J., Heath, B., Nordahl, C.W., Amaral, D.G.: Sex-dependent structure of socioemotional salience, executive control, and default mode networks in preschool-aged children with autism. NeuroImage **257**, 119252 (2022) https://doi.org/10.1016/j.

neuroimage.2022.119252

[22] Li, W., Yang, C., Shi, F., Wu, S., Wang, Q., Nie, Y., Zhang, X.: Construction of individual morphological brain networks with multiple morphometric features. Frontiers in Neuroanatomy **11** (2017) https://doi.org/10.3389/fnana.2017.00034

[23] Seidlitz, J., Váša, F., Shinn, M., *et al.*: Morphometric similarity networks detect microscale cortical organization and predict inter-individual cognitive variation. Neuron **97**(1), 231–247 (2018) https://doi.org/10.1016/j.neuron.2017.11.039

[24] Sebenius, I., Seidlitz, J., Warrier, V., Bethlehem, R.A.I., Alexander-Bloch, A., Mallard, T.T., Garcia, R.R., Bullmore, E.T., Morgan, S.E.: Robust estimation of cortical similarity networks from brain mri. Nature Neuroscience **26**(8), 1461–1471 (2023) https://doi.org/10.1038/s41593-023-01376-7

[25] Jacokes, Z., Adoremos, I., Hussain, A.R., Newman, B.T., Pelphrey, K.A., and, J.D.V.H.: Unsupervised Dimensionality Reduction Techniques for the Assessment of ASD Biomarkers, pp. 614–630. https://doi.org/10.1142/9789819807024_0044

[26] Gaser, C., Dahnke, R., Thompson, P.M., Kurth, F., Luders, E., the Alzheimer's Disease Neuroimaging Initiative: Cat: a computational anatomy toolbox for the analysis of structural mri data. GigaScience **13** (2024) https://doi.org/10.1093/gigascience/giae049

[27] Fischl, B.: Freesurfer. Neuroimage **62**(2), 774–781 (2012)

[28] Zalesky, A., Fornito, A., Bullmore, E.T.: Network-based statistic: Identifying differences in brain networks. NeuroImage **53**(4), 1197–1207 (2010) https://doi.org/10.1016/j.neuroimage.2010.06.041

[29] Gracia-Tabuenca, Z., Alcauter, S.: Nbr: network-based r-statistics for (unbalanced) longitudinal samples. bioRxiv (2020)

[30] Hadjikhani, N., Joseph, R.M., Snyder, J., Chabris, C.F., Clark, J., Steele, S., McGrath, L., Vangel, M., Aharon, I., Feczko, E., Harris, G.J., Tager-Flusberg, H.: Activation of the fusiform gyrus when individuals with autism spectrum disorder view faces. NeuroImage **22**(3), 1141–1150 (2004) https://doi.org/10.1016/j.neuroimage.2004.03.025

[31] Pierce, K., Redcay, E.: Fusiform function in children with an autism spectrum disorder is a matter of "who". Biological Psychiatry **64**(7), 552–560 (2008) https://doi.org/10.1016/j.biopsych.2008.05.013

[32] Floris, D.L., Llera, A., Zabihi, M., Moessnang, C., Jones, E.J.H., Mason, L., Haartsen, R., Holz, N.E., Mei, T., Elleaume, C., Vieira, B.H., Pretzsch, C.M., Forde, N.J., Baumeister, S., Dell'Acqua, F., Durston, S., Banaschewski, T., Ecker, C., Holt, R.J., Baron-Cohen, S., Bourgeron, T., Charman, T., Loth, E., Murphy,

D.G.M., Buitelaar, J.K., Beckmann, C.F., Holz, N.E., Forde, N.J., Banaschewski, T., Langer, N.: A multimodal neural signature of face processing in autism within the fusiform gyrus. Nature Mental Health **3**(1), 31–45 (2025) https://doi.org/10.1038/s44220-024-00349-4

[33] Zielinski, B.A., Gennatas, E.D., Zhou, J., Seeley, W.W.: Network-level structural covariance in the developing brain. Proceedings of the National Academy of Sciences **107**(42), 18191–18196 (2010) https://doi.org/10.1073/pnas.1003109107

[34] Rudie, J.D., Shehzad, Z., Hernandez, L.M., Colich, N.L., Bookheimer, S.Y., Dapretto, M.: Reduced functional integration and segregation of distributed neural systems underlying social and emotional information processing in autism spectrum disorders. Cerebral Cortex **22**(5), 1025–1037 (2013) https://doi.org/10.1093/cercor/bhr171

[35] Keown, C.L., Datko, M.C., Chen, C.P., Maximo, J.O., Jahedi, A., Müller, R.-A.: Network organization is globally atypical in autism: A graph theory study of intrinsic functional connectivity. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging **2**(1), 66–75 (2017) https://doi.org/10.1016/j.bpsc.2016.07.008

[36] Abbott, A.E., Nair, A., Keown, C.L., Datko, M., Jahedi, A., Fishman, I., Müller, R.-A.: Patterns of atypical functional connectivity and behavioral links in autism differ between default, salience, and executive networks. Cerebral Cortex **26**(10), 4034–4045 (2015) https://doi.org/10.1093/cercor/bhv191

[37] Abbott, A.E., Linke, A.C., Nair, A., Jahedi, A., Alba, L.A., Keown, C.L., Fishman, I., Müller, R.-A.: Repetitive behaviors in autism are linked to imbalance of corticostriatal connectivity: a functional connectivity mri study. Social Cognitive and Affective Neuroscience **13**(1), 32–42 (2017) https://doi.org/10.1093/scan/nsx129

[38] Rasero, J., Jimenez-Marin, A., Diez, I., Toro, R., Hasan, M.T., Cortes, J.M.: The neurogenetics of functional connectivity alterations in autism: Insights from subtyping in 657 individuals. Biological Psychiatry **94**(10), 804–813 (2023) https://doi.org/10.1016/j.biopsych.2023.04.014

[39] Shukla, D.K., Keehn, B., Smylie, D.M., Müller, R.-A.: Microstructural abnormalities of short-distance white matter tracts in autism spectrum disorder. Neuropsychologia **49**(5), 1378–1382 (2011) https://doi.org/10.1016/j.neuropsychologia.2011.02.022

[40] Travers, B.G., Adluru, N., Ennis, C., Tromp, D.P.M., Destiche, D., Doran, S., Bigler, E.D., Lange, N., Lainhart, J.E., Alexander, A.L.: Diffusion tensor imaging in autism spectrum disorder: A review. Autism Research **5**(5), 289–313 (2012) https://doi.org/10.1002/aur.1243

[41] Aoki, Y., Abe, O., Nippashi, Y., Yamasue, H.: Comparison of white matter

integrity between autism spectrum disorder subjects and typically developing individuals: a meta-analysis of diffusion tensor imaging tractography studies. Molecular Autism **4**(1), 25 (2017) https://doi.org/10.1186/2040-2392-4-25

[42] Hong, S.-J., Wael, R., Bethlehem, R.A.I., Lariviere, S., Paquola, C., Valk, S.L., Milham, M.P., Di Martino, A., Margulies, D.S., Smallwood, J., Bernhardt, B.C.: Atypical functional connectome hierarchy in autism. Nature Communications **10**(1) (2019) https://doi.org/10.1038/s41467-019-08944-1

[43] Just, M.A., Keller, T.A., Malave, V.L., Kana, R.K., Varma, S.: Autism as a neural systems disorder: A theory of frontal–posterior underconnectivity. Neuroscience & Biobehavioral Reviews **36**(4), 1292–1313 (2012) https://doi.org/10.1016/j.neubiorev.2012.02.007

[44] Menon, V.: Large-scale brain networks and psychopathology: a unifying triple network model. Trends in Cognitive Sciences **15**(10), 483–506 (2011) https://doi.org/10.1016/j.tics.2011.08.003

[45] Tsang, T., Green, S.A., Liu, J., Lawrence, K., Jeste, S., Bookheimer, S.Y., Dapretto, M.: Salience network connectivity is altered in 6-week-old infants at heightened likelihood for developing autism. Communications Biology **7**(1) (2024) https://doi.org/10.1038/s42003-024-06016-9

[46] Gao, L., Cao, Y., Zhang, Y., Liu, J., Zhang, T., Zhou, R., Guo, X.: Sex differences in the flexibility of dynamic network reconfiguration of autism spectrum disorder based on multilayer network. Brain Imaging and Behavior **18**(5), 1172–1185 (2024) https://doi.org/10.1007/s11682-024-00907-5

[47] Cheng, Y., Chou, K.-H., Chen, I.-Y., Fan, Y.-T., Decety, J., Lin, C.-P.: Atypical development of white matter microstructure in adolescents with autism spectrum disorders. NeuroImage **50**(3), 873–882 (2010) https://doi.org/10.1016/j.neuroimage.2010.01.011

[48] Barnea-Goraly, N., Lotspeich, L.J., Reiss, A.L.: Similar white matter aberrations in children with autism and their unaffected siblings: A diffusion tensor imaging study using tract-based spatial statistics. Archives of General Psychiatry **67**(10), 1052 (2010) https://doi.org/10.1001/archgenpsychiatry.2010.123

[49] Kanwisher, N., McDermott, J., Chun, M.M.: The fusiform face area: A module in human extrastriate cortex specialized for face perception. The Journal of Neuroscience **17**(11), 4302–4311 (1997) https://doi.org/10.1523/jneurosci.17-11-04302.1997

[50] Schultz, R.T., Grelotti, D.J., Klin, A., Kleinman, J., Gaag, C., Marois, R., Skudlarski, P.: The role of the fusiform face area in social cognition: implications for the pathobiology of autism. Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences **358**(1430), 415–427 (2003) https:

//doi.org/10.1098/rstb.2002.1208

[51] Schultz, R.T.: Developmental deficits in social perception in autism: the role of the amygdala and fusiform face area. International Journal of Developmental Neuroscience **23**(2–3), 125–141 (2005) https://doi.org/10.1016/j.ijdevneu.2004.12.012

[52] Schultz, H., Sommer, T., Peters, J.: The role of the human entorhinal cortex in a representational account of memory. Frontiers in Human Neuroscience **9** (2015) https://doi.org/10.3389/fnhum.2015.00628

[53] Coutureau, E., Di Scala, G.: Entorhinal cortex and cognition. Progress in Neuro-Psychopharmacology and Biological Psychiatry **33**(5), 753–761 (2009) https://doi.org/10.1016/j.pnpbp.2009.03.038

[54] Ecker, C., Ginestet, C., Feng, Y., Johnston, P., Lombardo, M.V., Lai, M.-C., Suckling, J., Palaniyappan, L., Daly, E., Murphy, C.M., Williams, S.C., Bullmore, E.T., Baron-Cohen, S., Brammer, M., Murphy, D.G.M., MRC AIMS Consortium, f.t.: Brain surface anatomy in adults with autism: The relationship between surface area, cortical thickness, and autistic symptoms. JAMA Psychiatry **70**(1), 59 (2013) https://doi.org/10.1001/jamapsychiatry.2013.265

[55] Khundrakpam, B.S., Lewis, J.D., Kostopoulos, P., Carbonell, F., Evans, A.C.: Cortical thickness abnormalities in autism spectrum disorders through late childhood, adolescence, and adulthood: A large-scale mri study. Cerebral Cortex **27**(3), 1721–1731 (2017) https://doi.org/10.1093/cercor/bhx038

[56] Lucibello, S., Bertè, G., Verdolotti, T., Lucignani, M., Napolitano, A., D'Abronzo, R., Cicala, M.G., Pede, E., Chieffo, D., Mariotti, P., Colosimo, C., Mercuri, E., Battini, R.: Cortical thickness and clinical findings in prescholar children with autism spectrum disorder. Frontiers in Neuroscience **15** (2022) https://doi.org/10.3389/fnins.2021.776860

[57] Arutiunian, V., Gomozova, M., Minnigulova, A., Davydova, E., Pereverzeva, D., Sorokin, A., Tyushkevich, S., Mamokhina, U., Danilina, K., Dragoy, O.: Structural brain abnormalities and their association with language impairment in school-aged children with autism spectrum disorder. Scientific Reports **13**(1) (2023) https://doi.org/10.1038/s41598-023-28463-w

[58] Auzias, G., Viellard, M., Takerkart, S., Villeneuve, N., Poinso, F., Fonséca, D.D., Girard, N., Deruelle, C.: Atypical sulcal anatomy in young children with autism spectrum disorder. NeuroImage: Clinical **4**, 593–603 (2014) https://doi.org/10.1016/j.nicl.2014.03.008

[59] Nordahl, C.W., Dierker, D., Mostafavi, I., Schumann, C.M., Rivera, S.M., Amaral, D.G., Van Essen, D.C.: Cortical folding abnormalities in autism revealed by surface-based morphometry. The Journal of Neuroscience **27**(43), 11725–11735

(2007) https://doi.org/10.1523/jneurosci.0777-07.2007

[60] Lai, M.-C., Lombardo, M.V., Ruigrok, A.N., Chakrabarti, B., Auyeung, B., Szatmari, P., Happé, F., Baron-Cohen, S.: Quantifying and exploring camouflaging in men and women with autism. Autism **21**(6), 690–702 (2016) https://doi.org/10.1177/1362361316671012

[61] Corbett, B.A., Schwartzman, J.M., Libsack, E.J., Muscatello, R.A., Lerner, M.D., Simmons, G.L., White, S.W.: Camouflaging in autism: Examining ¡scp¿sex-based¡/scp¿ and compensatory models in social cognition and communication. Autism Research **14**(1), 127–142 (2020) https://doi.org/10.1002/aur.2440

[62] Benkarim, O., Paquola, C., Park, B.-y., Hong, S.-J., Royer, J., Wael, R., Lariviere, S., Valk, S., Bzdok, D., Mottron, L., C. Bernhardt, B.: Connectivity alterations in autism reflect functional idiosyncrasy. Communications Biology **4**(1) (2021) https://doi.org/10.1038/s42003-021-02572-6

[63] Buch, A.M., Vértes, P.E., Seidlitz, J., Kim, S.H., Grosenick, L., Liston, C.: Molecular and network-level mechanisms explaining individual differences in autism spectrum disorder. Nature Neuroscience **26**(4), 650–663 (2023) https://doi.org/10.1038/s41593-023-01259-x

[64] He, C., Cortes, J.M., Ding, Y., Shan, X., Zou, M., Chen, H., Chen, H., Wang, X., Duan, X.: Combining functional, structural, and morphological networks for multimodal classification of developing autistic brains. Brain Imaging and Behavior (2025) https://doi.org/10.1007/s11682-025-01026-5

[65] Uddin, L.Q., Dajani, D.R., Voorhies, W., Bednarz, H., Kana, R.K.: Progress and roadblocks in the search for brain-based biomarkers of autism and attention-deficit/hyperactivity disorder. Translational Psychiatry **7**, 1218 (2017) https://doi.org/10.1038/tp.2017.164

[66] Hull, J.V., Dokovna, L.B., Jacokes, Z.J., *et al.*: Resting-state functional connectivity in autism spectrum disorders: A review. Frontiers in Psychiatry **7**, 205 (2016) https://doi.org/10.3389/fpsyt.2016.00205

Supplementary Information

Fig. S1: **Methodological Workflow**. We analyzed a sex- and diagnosis-balanced sample of 337 individuals aged 8–18 years. T1w images were preprocessed using FreeSurfer and CAT12 for voxel-based morphometry. From the preprocessed data, we extracted regional distributions of multivariate morphological features (mean curvature, cortical thickness, sulcus depth, surface area, and volume) defined by the Desikan–Killiany cortical atlas. Inter-regional connectivity was then estimated using the Kullback–Leibler divergence between these multivariate distributions. Finally, connectivity matrices were regressed onto group labels while controlling for covariates (age, intracranial volume, and collection site), adopting a network-based statistic approach. Analyses were performed separately by sex.

Fig. S2: **Network profiles with different individual edge thresholds (male cohort)**. Circular plots visualizing clusters of significant inter-regional connections using a more stringent individual edge threshold ($p = 0.001$, first row) and a more lenient threshold ($p = 0.01$, second row), than in the main analysis ($p = 0.005$). Nodes represent cortical regions, with olive and teal indicating the left and right hemispheres, respectively. Blue colors indicate increased connectivity in the TDC group relative to ASD (left panels), and blue colors the opposite (right panels).

Fig. S3: **Network profiles with different individual edge thresholds (female cohort)**. Circular plots visualizing clusters of significant inter-regional connections using a more stringent individual edge threshold ($p = 0.001$, first row) and a more lenient threshold ($p = 0.01$, second row), than in the main analysis ($p = 0.005$). Nodes represent cortical regions, with olive and teal indicating the left and right hemispheres, respectively. Blue colors indicate increased connectivity in the TDC group relative to ASD (left panels), and blue colors the opposite (right panels).

Fig. S4: **Sex-normative morphological network profiles**. Circular plots visualizing clusters of significant inter-regional connections comparing males and females within the TDC group. The same setup as in the main analysis was adopted (individual edge threshold p = 0.005; FWE-corrected p = 0.05; 5,000 permutations). Nodes represent cortical regions, with olive and teal indicating the left and right hemispheres, respectively. Indigo colors indicate increased connectivity in males relative to females, and green the opposite.