1   **Ventral Pallidum GABA Neuron Inhibition Augments Context-Appropriate Defensive**
2   **Responses to Learned Threat Cues**
3
4

5   Erica M. Ramirez*[1], Maricela X. Martinez*[1], Ryan K. Rokerya[1], Vanessa Alizo Vera[1], Christina
6   M. Ruiz[1], Mitchell R. Farrell[1], Shreeya A. Walawalkar[1], Hazael Ramirez-Ramirez[1], Grace J.
7   Kollman[2], and Stephen V. Mahler[1]
8
9   *These authors contributed equally to the manuscript
10
11 [1]Department of Neurobiology & Behavior, University of California, Irvine. 1203 McGaugh Hall,
12 Irvine, CA 92697, USA
13 949-824-4727
14
15 [2]Department of Neurosciences, University of California, San Diego. 9375 Gilman Dr, San Diego,
16 CA 92161, USA
17
18 **Corresponding Author:** Erica M. Ramirez ericamr1@uci.edu
19
20 **Other Authors:** maricexm@uci.edu, ryanrokerya@gmail.com, valizove@uci.edu,
21 cruiz1@uci.edu, Mitchell.Farrell.phd@gmail.com, swalawal@uci.edu, hazaelr@uci.edu,
22 gracekollman@gmail.com, mahlers@uci.edu

23
24
25 **Keywords**
26
27
28 *Aversive Motivation, Fear Conditioning, Defensive Treading, Lateral Habenula, Mediodorsal*
29 *Thalamus*
30
31

Abstract

The ventral pallidum (VP) is embedded within the brain circuits controlling motivated behavior, which are heavily implicated in addiction and other psychiatric disorders. Prior work showed that VP GABAergic neurons (VP[GABA]) promote reward approach and seeking, while the intermixed population of VP glutamate neurons instead promote avoidance and aversion. Some have thus suggested a functional dichotomy between these VP subpopulations in reward versus threat. We test this hypothesis by asking how inhibiting VP[GABA] impacts active and passive defensive responses to learned threat cues in the absence of rewards. We taught GAD1:Cre rats with inhibitory VP[GABA] DREADDs (or control rats) that a metal probe delivers shock, or that a 20sec auditory cue precedes footshocks. These stimuli thereafter elicit active defensive burying, or passive freezing responses, respectively. We found that VP[GABA] inhibition with CNO markedly increased stimulus-appropriate defensive responses to both types of learned threats, but failed to consistently alter new learning about them—suggesting VP[GABA] mediates aversive motivation but not memory formation. VP[GABA] inhibition also altered threat-related c-Fos expression within VP cell populations, and in their efferent target lateral habenula, but not mediodorsal thalamus—pointing to potential underlying circuit mechanisms of defensive responses. Results indicate that VP[GABA] neurons not only promote reward seeking as previously reported, but that they may also actively inhibit defensive responses to threats that might otherwise compete with reward seeking. This refines our understanding of subcortical valanced motivation circuits, and may suggest new targets for intervening in disorders like addiction and depression.

52   *Graphical abstract:*

VP: ventral pallidum, LHb: Lateral Habenula, MDT: Mediodorsal Thalamus

53
54
55
56   ***Highlights:***

57
58   *-Inhibiting VP^{GABA} increases defenses against threatening stimuli*
59   *-Both active (burying) and passive (freezing) responses were enhanced*
60   *-Inhibiting VP^{GABA} excites other VP cells, and neurons in downstream LHb*
61   *-VP subpopulations may interact to bidirectionally modulate circuits and adaptive behaviors*
62

1. Introduction

The ventral pallidum (VP) is best known for its role in reward, especially via its strong inputs from nucleus accumbens, and prominent projections to ventral tegmental area (VTA) (Heimer et al., 1991; Kalivas et al., 1993; Mahler et al., 2014; Ottenheimer et al., 2018; Pardo-Garcia et al., 2019; Smith & Berridge, 2007; Zahm, 1989). However, like other nearby hypothalamic and extended amygdala regions (Barker et al., 2017; Jennings et al., 2013a; Lammel et al., 2012; Nieh et al., 2016), VP also contains glutamatergic populations that instead mediate aversion, in part via their direct projections to lateral habenula (LHb) (Haber et al. 1985; Zahm et al. 1996; Faget et al., 2018, 2024; Stephenson-Jones et al., 2020; Tooley et al., 2018; Wang et al., 2020). Yet VP sends both GABA and glutamate projections to LHb as well as reward-related regions (Soares-Cunha & Heinsbroek, 2023)—suggesting unresolved complexity in the specific behavioral roles of these motivation circuits. Since adaptive decision making depends upon proper balance between appetitive and aversive motivations, understanding how their circuit substrates function normally, as well as how they become imbalanced in psychiatric disorders, is essential.

Cell type-specific manipulation and observation studies in mice suggest that VP^{GABA} neurons selectively mediate reward and approach, while VP^{glutamate} neurons instead mediate aversion and avoidance (Heinsbroek et al., 2020; Stephenson-Jones et al., 2020; Tooley et al., 2018; Wulff et al., 2019, Faget et al., 2018, 2024). In rats, we and others find that VP^{GABA} neurons also mediate pursuit of highly salient rewards, especially when triggered by learned reward cues (Farrell et al., 2021, 2022; Scott et al., 2023, 2025; Prasad et. al. 2020). In contrast, we found that inhibiting VP^{GABA} neurons did not alter unconditioned affective responses to unpleasant stimuli like foot shocks in rats, and caused only minor effects on operant avoidance of shock (Farrell et al., 2021, 2022). These data largely support the idea of a highly specific role for VP^{GABA} in appetitive motivation.

This said, we still wondered if VP^{GABA} neurons might play a role in aversive motivation under certain circumstances, especially in response to aversive conditioned stimuli. Might VP^{GABA} neurons be capable of constraining responses to perceived threats in addition to promoting reward seeking, allowing the organism suppress threat responses and to fully commit to capitalizing on opportunities? Since reward-related behavioral effects of VP^{GABA} manipulations are often clearest when motivation is elicited by conditioned cues (Scott et al., 2023; Farrell et. al., 2022), and since few studies had interrogated VP^{GABA} involvement in conditioned *aversive* motivation, we hypothesized that manipulations of VP^{GABA} might alter motivated responses to cues signifying threat, as they do cues paired with reward.

We therefore asked how chemogenetically inhibiting VP^{GABA} neurons affects defensive responses to perceived threats when they are encountered in the absence of reward-related stimuli. We tested both a spatially-localized threat (a probe that can deliver a shock), which normally elicits active defensive burying of the probe, and a temporally-predictive threat (a 20sec auditory cue predicting foot shock), which normally elicits passive defensive freezing. Inhibiting VP^{GABA} neurons augmented context-appropriate defensive responses to both threat cues (be it active defense or passive freezing), without impacting other behaviors, or initial learning about threats. In addition, inhibiting VP^{GABA} neurons increased threat-induced Fos expression in neighboring, likely non-GABAergic (and possibly glutamatergic) VP cells, as well as in the downstream LHb, but not mediodorsal thalamus (MDT)—consistent with selective disinhibition of aversion-related neural circuits.

This report is thus the first to show a clear behavioral role for VP^{GABA} neurons in conditioned aversive motivation, and it supports the idea that VP^{GABA} circuits may not only promote reward

seeking directly, but also concurrently suppress threat responses that might otherwise constrain or compete with reward pursuit. We thus provide novel insights into the context-dependent neural mechanisms underlying adaptive responses to salient, affectively-valanced stimuli, and how they may go awry in stress-related disorders.

2. Materials and Methods

2.1 Subjects
Long Evans male and female GAD1:Cre rats (n=68; n=24 female) and their Cre-negative wildtype (WT) littermates (n=17; n=9 female) were bred in-house, and housed in pairs or trios in ventilated tub cages with corncob bedding and *ad libitum* chow and water. Experiments were conducted during the dark phase of a reversed 12:12hr light cycle. Rats were 75+ days old when experiments commenced. Procedures were approved by the UCI Institutional Animal Care and Use Committee, and followed NIH Guide for the Care and Use of Laboratory Animals guidelines.

2.2 Stereotaxic Surgery
GAD1:Cre and WT rats were anesthetized with ketamine (56.5mg/kg), xylazine (8.7mg/kg), and meloxicam (1mg/kg), and an AAV2 vector containing a Cre-dependent, mCherry-tagged inhibitory DREADD (hSyn-DIO-hM4Di-mCherry; titer: 5×10^{12} vg/mL; Addgene Cat. #:44362-AAV2) was injected bilaterally with a glass pipette into VP (~300nl/hemisphere), at bregma-relative coordinates (mm): AP: 0.0, ML: ±2.0, DV: −8.0, as previously described (Martinez et al., 2023). Rats were given at least 21 days to recover from surgery and allow for viral expression to occur before commencing behavioral testing.

2.3 CNO Preparation and Injection
Clozapine-N-oxide (CNO) was obtained from the NIDA Drug Supply Program, stored at 4°C in powder aliquots with desiccant, and protected from light. CNO (5mg/kg/ml) was dissolved daily for IP injection in 5% dimethyl sulfoxide, then diluted in 0.9% saline to dose. Vehicle (VEH) or CNO was injected IP 30min prior to commencement of all behavioral tests (Farrell et al., 2019, 2021, 2022; Lawson et al., 2021, 2023; Mahler et al., 2014; Martinez et al., 2024).

2.4 Behavioral Methods
2.4.1 Experimental Timeline: At least 4 weeks after surgery, rats were handled for 5min on 5 consecutive days prior to behavioral training. Male and female GAD1:Cre (n=24) and WT rats (n=11) first underwent testing for shock probe fear expression (2 expression tests, counterbalanced VEH and CNO), then they were tested for expression of freezing elicited by 0.75mA foot shock-paired cues in either VP^{GABA}-Inhibited (GAD1:Cre+CNO; n=16) or Control-treated (GAD1:Cre+VEH: n=8; or WT+CNO: n=11) rats in a between-subjects design. Other male and female rats first underwent testing for expression of auditory cue fear elicited by 0.3mA foot shock-paired cues after VP^{GABA} inhibition (GAD1:Cre+CNO: n=16) or Control treatment (GAD1:Cre+VEH: n=11; WT+CNO: n=6), then they were subsequently used to test effects of Inhibition (GAD1:Cre+CNO; n=14) or Control treatment (GAD1:Cre+VEH: n=8; or WT+CNO: n=6) on acquisition of shock probe fear learning. Another group of male GAD1:Cre rats expressing VP^{GABA} DREADDs that previously underwent auditory cue fear conditioning training and expression testing (0.5mA shock during training; behavioral data from this pilot experiment not shown) were given 2 weeks off, then they were used to examine neural activity (Fos) elicited by a threatening or non-threatening probe after injection of either VEH or CNO (Threatening Probe + VEH: n=5; Threatening Probe + CNO: n=4; Non-Threatening Probe + VEH: n=5; Non-Threatening Probe + CNO: n=3).

165
2.4.2 Shock Probe Test General Methods: All shock probe sessions were videorecorded for
later quantification of behaviors, as described in **Supplemental Methods**. Rats were first
habituated for 30min on two days to a specific plexiglass tub cage filled with corncob bedding
(5cm height from cage bottom). They were then returned to the chamber on the following day
for a <u>shock probe fear acquisition session</u>. The chamber now had a copper-wrapped plastic
probe inserted through a 1cm diameter hole in the wall located 8cm above the cage floor. The
probe delivered a 1.5mA shock when touched (Coulbourn Precision Animal Shocker). Rats
were placed in the opposite end of the cage from the shock probe, and the shock probe
acquisition session continued for 20min after the first rat-initiated shock (Fucich & Morilak, 2018,
Berridge et al., 1988; Craft et al., 1988; Pinel & Treit, 1978). The following day, rats underwent a
<u>shock probe fear expression session</u>. The chamber was configured identically to the shock
probe fear acquisition session, except the probe was inactive and no longer delivered shock
when touched.
179
2.4.3 VP^{GABA} Inhibition Effects on Shock Probe Fear Expression: To test effects of VP^{GABA}
inhibition on expression of defensive responses directed at a threatening probe which previously
delivered shock, rats received VEH or CNO injection, then were placed into the chamber with
the inactive probe for 20min. 48h later, these rats underwent a short (5min) shock probe fear
"re-acquisition" session, where the probe was re-electrified, and on which all rats were shocked
again at least once. The next day, they underwent a second 20min shock probe fear expression
test, conducted identically to the first, other than that they received the alternative injection
(VEH/CNO) prior to the session. Rats received VEH and CNO treatments in counterbalanced
order in these two shock probe fear expression tests, and defensive burying behaviors were
statistically equivalent in rats that received VEH on the first versus the second shock probe fear
expression test (Mann-Whitney U=162, p=0.6072). Therefore, data from both shock probe fear
expression tests were combined for repeated measures statistical analysis. One rat's data was
excluded from this experiment because it never received a shock during the shock probe fear
acquisition session, and 3 rats' data were excluded because they never emitting
treading/digging behavior on either of the two shock probe fear expression tests.
195
196 *2.4.4 VP^{GABA} Inhibition Effects on Shock Probe Fear Acquisition:* To test effects of VP^{GABA}
197 inhibition on initial learning about the shock-delivering probe, rats received either VEH or CNO
198 injections prior to a shock probe fear acquisition session, following prior testing of auditory cue
199 fear expression. 48hrs later, these rats were returned to the chamber with the now-inactive
200 probe, without further injections, for a 20min shock probe fear expression test.
201
2.4.5 VP^{GABA} Inhibition Effects on Auditory Cue Fear Expression: All sessions were
videorecorded for later quantification of behaviors, as described in **Supplemental Methods**.
Auditory cue fear conditioning training and testing took place in Med Associates rat operant
chambers (30.5x24x21cm; St. Albans, VT), housed within sound-attenuating boxes. Chambers
had a peppermint scent (McCormick) during cue/shock training, and standard metal bars that
could deliver foot shock composed the chamber floor. On these 8.5min training sessions, rats
were placed into the novel chamber, and a house light turned on for a 3min baseline period.
Then, a 20sec auditory cue (continuous white noise: 80dB) was played, which was followed
immediately by either a low (0.3mA) or high (0.75mA) intensity foot shock, delivered for 2sec.
Three cue/shock pairings were made during this training session, with 90sec elapsing between
each. 48hrs later, rats received either VEH or CNO injection, then they were placed into a
different Med Associates chamber with a distinct metal grid floor, and an orange scent
(McCormick) for a 47min auditory cue fear expression session. After a 3min baseline period, the

20sec shock-associated white noise cue was played a total of 24 times, with 90sec elapsing
between each. No shocks were delivered during this session.
217
2.4.6 VP^{GABA} Inhibition Effects on Probe Threat-Elicited Neural Activity: Male GAD1:Cre rats
with VP^{GABA} DREADDs were habituated and trained as described in Section 2.4.2 on the shock
probe fear acquisition procedure, except that about half the rats underwent the 20min training
session with a probe that did not deliver shock. The following day, rats were injected with either
VEH or CNO, then they underwent a shock probe fear expression session as described in
section 2.4.2 (probes did not deliver shock on this session). Rats were returned to their home
cages for 90min after the session, then they were perfused and brains were prepared for Fos
staining and quantification. One rat in the No-Threat+VEH group did not express sufficient
mCherry in VP^{GABA} cells for quantification, so it was excluded from mCherry-based VP Fos
analyses, but this rat's data was included in analyses of Fos in VP^{ChAT} neurons, LHb, and MDT,
since these analyses do not require identification of VP^{GABA} versus putative VP^{nonGABA}
populations.
230
2.6 Histology, Immunohistochemistry, and Microscopy Methods
232 Details of staining protocols and Fos quantification methods are provided in **Supplemental**
233 **Methods**.
234
235 *2.6.1 Preparation of Slices for Staining:* All rats were perfused transcardially with chilled 0.9%
236 saline and 4% paraformaldehyde, brains were postfixed in 4% paraformaldehyde for ~16hrs,
237 cryoprotected in 20% sucrose-azide for 3+ days, then sectioned coronally at 40µm into PBS-
238 azide.
239
240 *2.6.2 Visualizing & Quantifying VP Virus Expression:* Cre-dependent virus expression localized
241 predominantly within VP was verified in each GAD1:Cre rat used in these studies, by analyzing
242 the zone of immunohistochemically-amplified mCherry expression around viral injection sites in
243 relation to substance P-defined borders of the VP (Haber & Nauta, 1983; Zahm & Heimer, 1988,
244 1990). Lack of mCherry expression was also verified in each WT Control rat.
245
2.6.3 Visualizing & Quantifying Neural Activity in VP Cell Populations, LHb, and MDT: We used
dual color immunofluorescence to stain for Fos protein, and to amplify Cre-dependent mCherry
expression in VP^{GABA} cells in the immediate vicinity of the center of virus injection sites. We
examined the percentage of VP^{GABA} neurons (i.e. mCherry+ cells, as these were previously
confirmed to be ~91% GABAergic (Farrell et al., 2021)) that also expressed Fos, and we also
quantified the number of Fos+ VP cells that did not also express mCherry in the same zone.
Although we did not verify the identity of these mCherry-negative VP cells directly, they were
well within the zone of Cre-dependent expression of mCherry in VP^{GABA} cells using this highly-
penetrant and specific transfection system (Farrell et al., 2021), so we hereafter refer to them as
putative VP^{nonGABA} cells. We note that we were unable to directly identify whether these
unlabeled cells were VP^{glutamate} using the IHC methods available to us, since vGlut2 protein does
not persist long in soma before it is trafficked axonally. Therefore, we further explored the
potential identity of these Fos+, putatively VP^{nonGABA} cells by staining additional adjacent
sections for Fos and choline acetyltransferase (ChAT), which allowed us to determine whether
this cholinergic population was affected by Threat and/or VP^{GABA} Inhibition. Finally, we also Fos-
stained sections that contained LHb and MDT from the same animals, allowing us to determine
how neural activity in both these VP efferent targets was affected by Threat and/or VP^{GABA}
Inhibition, and how this activity relates to activity in VP subpopulations in the same individuals.
264
2.7 Data Analysis Approach

2.7.1 *Group Comparisons:* To determine effects of VP^{GABA} inhibition on shock probe fear expression, male and female GAD1:Cre and WT rats received VEH and CNO in counterbalanced order before each of two probe fear expression tests, and within-subjects comparisons were used. To determine effects of VP^{GABA} inhibition on shock probe fear acquisition and auditory cue fear expression, GAD1:Cre rats with DREADDs that were injected with CNO (Inhibited Group) were compared to Controls that consisted of a mixture of GAD1:Cre rats with DREADDs that were given VEH (n=8 high-shock intensity experiment, 11 low-shock intensity experiment, 8 shock probe fear acquisition experiment), and WT rats without DREADDs that were given CNO (n=11 high-shock, 6 low-shock, 6 shock probe fear acquisition). No behavioral differences were seen between these subgroups of Control rats in 1st cue freezing, or in defensive burying (ts<1.522, ps>0.05), so they were combined for primary comparisons of Control and Inhibited rats.

2.7.2 *Statistical Analyses:* Statistical analyses were conducted in Graphpad Prism. ANOVAs and t-tests were used to determine main effects and interactions in normally distributed datasets. When data violated normality assumptions of parametric tests (assessed by D'Agostino & Pearson tests), data were either log-normalized (shock probe fear expression experiment), or non-parametric tests (Mann-Whitney U) were used for independent samples. For auditory cue fear expression tests, a repeated measures Trial (24 Cue Deliveries) factor was included in ANOVAs to examine extinction of cue responses within the session. Secondary analyses for each experiment examined main effects of Sex, and its interactions with other variables. Fos data were analyzed with two-way ANOVA incorporating Drug (VEH, CNO) and Probe Type (Threat, No-Threat), and Pearson correlations examined how Fos in LHb or MDT relates to Fos in VP^{GABA}, putative VP^{nonGABA}, or VP^{ChAT} subpopulations. All tests were 2-tailed, with significance criterion set at $p<0.05$.

3. Results

3.1 Analysis of VP DREADD Expression

The previously-validated (Farrell et al., 2021, 2022), Cre-dependent hM4Di DREADD/mCherry vector caused robust expression in VP^{GABA} neurons of GAD1:Cre rats**,** but not in their Cre- WT littermates (**Fig. 1A,D**). Somatic mCherry expression in included rats was localized predominantly within Substance P-defined VP borders of GAD1:Cre rats (**Fig. 1B**), as well as in these neurons' axons, which were present as expected in VP efferent targets including LHb and MDT (Groenewegen & Berendse, 1990; Haber et al., 1993; Root et al., 2015; Zahm et al., 1987) (**Fig. 1C**). Some GAD1:Cre rats were excluded from behavioral analyses for significant DREADD expression observed outside VP borders (n=5), or lack of expected VP mCherry expression (n=9).

3.2 VP^{GABA} Inhibition Enhances Active Defensive Responses to a Localized Threat

We first asked whether inhibiting VP^{GABA} neurons impacts defensive burying behaviors directed at a localized, previously-learned threat—a probe that delivered an electric shock during training, but which was inactive during counterbalanced VEH and CNO fear expression tests **(Fig. 2A)**.

3.2.1 *Effects of VP^{GABA} Inhibition on Probe Fear Expression:* Inhibiting VP^{GABA} neurons markedly increased expression of defensive responses to the previously-learned shock probe. Relative to VEH day, CNO in GAD1:Cre rats (n=11M,13F) increased time animals spent actively burying the shock probe, compared to Controls (n=6F,5M) (DREADD Group x Drug Interaction: $F_{1,28}=4.196$, p=0.050; Holm-Sidak posthoc for VEH versus CNO in GAD1:Cre rats: $t_{28}=3.226$,

317   p=0.007; **Fig. 2B**). Inhibited rats also displayed significantly more bouts of burying behavior
318   ($F_{1,28}$=6.687, p=0.015; Holm-Sidak: t_{28}=3.291, p=0.005; **Fig. 2C**) and non-significant trends
319   toward increased height (cm) of bedding piles formed ($F_{1,32}$=2.068, p=0.156, **Fig. 2D**), and
320   decreased latency to begin burying ($F_{1,32}$=2.585, p=0.117, **Fig. 2E**) were observed.
321

Figure 1: Targeting VP^GABA Neurons with Inhibitory DREADDs in GAD1:Cre Rats
(A) Diagram of hM4Di-vector injection to the VP. **(B)** Representative image of mCherry (red) expression in the Substance P (green)-defined VP from an included GAD1:Cre rat. DAPI is blue. **(C)** mCherry-expressing VP^GABA axons (red) in LHb and MDT are shown, DAPI is blue. **(D)** Location of virus expression in each included GAD1:Cre rat is shown in red, with Substance P-defined VP borders represented in green. Coordinates indicate distance anterior or posterior of bregma, in mm. AP=anterior-posterior coordinate relative to bregma. 3V=3^rd ventricle.

322   In contrast to these active defensive responses to the probe, inhibiting VP^GABA did not alter other
323   behaviors less linked to active defense, such as the number of probe investigations
324   ($F_{1,32}$=0.506, p=0.479, **Fig. 2F**) or escape attempts ($F_{1,32}$=0.27, p=.606). Freezing was
325   uncommon in this active fear expression task and was unaffected by VP^GABA inhibition (data not
326   shown).
327
Effects of CNO in GAD1:Cre rats are attributable to the chemogenetic manipulation of VP^GABA
neurons, as no significant main effects of Drug were observed in the DREADD Group x Drug
analysis for burying bouts (Main effect of Drug: $F_{1,28}$=0.0112, p=0.917) or duration ($F_{1,28}$=3.056,
p=0.09). During initial drug-free learning of the shock probe threat, GAD1:Cre and Control rats
acquired the task similarly, indicating that observed VP^GABA inhibition effects did not result from
pre-existing group differences in training history or efficacy. All rats contacted the probe at least
once during initial shock training, and the duration and number of bouts of defensive burying did
not differ between GAD1:Cre and Control rats (DREADD Group x Drug Interaction: Fs<1.734,
ps>0.199; **Fig. S1A-B**). Results thus indicate that inhibiting VP^GABA neurons robustly and
selectively increased active defensive responses directed at a previously-learned localized
threat.
339
3.2.2 Similar Effects of VP^GABA Inhibition on Probe Fear Expression in Both Sexes: Only minor
sex differences were found in this probe fear expression test. Overall, males spent slightly more
time burying than females (Main effect of Sex, log-normalized data: $F_{1,31}$=5.128, p=0.031), and
built taller piles of bedding ($F_{1,31}$=8.620, p=0.006), while females spent more time attempting to
escape than males (Main effect of Sex: $F_{1,37}$=32.630, p<0.0001). There were no other sex
differences in probe-directed behaviors (burying bouts, latency, pile height, probe investigations)
or freezing (Fs<3.636, ps>0.05; **Fig. S2A-D**).
347
348   No sex differences were seen in effects of VP^GABA inhibition on defensive responses to the
349   probe, since no significant Sex x Drug interactions were observed for any behavior in Inhibited

rats, including burying, pile height, latency to bury, freezing, escape attempts, or probe investigations (all Fs<2.573, ps>0.120; **Fig. S2A-D**). No sex-specific effects of CNO versus VEH were observed in Control animal behaviors either (no Sex x Drug interactions, log-normalized data; Fs<2.572, ps>0.05).

Figure 2: VP^{GABA} Inhibition Enhanced Active Defensive Burying Responses to a Localized Threat (A) Behavioral protocol for probe fear expression tests. **(B-E)** Enhanced defensive burying behaviors in GAD1:Cre rats on CNO day relative to VEH day, with no CNO effect in Control rats. **(F)** In contrast, inhibition caused no change in probe investigations. *p<0.05, Holm-Sidak multiple comparison posthoc. Data were log-normalized to meet ANOVA assumptions, and for display.

3.3 VP^{GABA} Inhibition Does Not Consistently Alter Learning About the Shock Probe

3.3.1 Effects of VP^{GABA} Inhibition on Active Fear Acquisition: Next we asked whether inhibiting VP^{GABA} impacts learning about the shock-delivering probe, by training a cohort of Inhibited (n=15;6F,9M) and Control rats (n=15;6F,9M) to acquire a fear memory during VP^{GABA} manipulation (**Fig. 3A**). VP^{GABA} inhibited rats did not differ from Controls in defensive burying during this acquisition session (Mann-Whitney test; duration: U=85, p=0.563, **Fig. 3B**; bouts: U=65.5, p=0.140, **Fig. 3C**; pile height: U=89, p=0.689, **Fig. 3D**; latency: U=65, p=0.137, **Fig. 3E**). Probe investigations (U=95, p=0.834, **Fig. 3F**) and freezing (U=88, p=0.667) were similarly unaffected during acquisition. Rats were then tested 48hrs later with the non-electrified probe, drug-free, thus querying retention of the threat learning that occurred in the presence or absence of VP^{GABA}. Again, no significant differences between Inhibited and Control rats were observed overall in defensive responses (duration: U=87.5, p=0.643, **Fig. 3G**; bouts: U=84.5, p=0.765, **Fig. 3H**; pile height: U=88.5, 0.673, **Fig. 3I**; latency: U=95.5, p=0.918, **Fig. 3J**) or probe investigations (U=62, p=0.104, **Fig. 3K**), and freezing was not observed in any rats. These data suggest that VP^{GABA} inhibition does not consistently alter unconditioned responses to an electrified shock probe itself, nor learning or consolidation of memories about this threat.

3.3.2 Sex Differences in Active Fear Acquisition Tests: Female rats displayed slightly less burying behavior than males during shock probe acquisition (main effect of Sex; burying duration: $F_{1,22}$=3.378, p=0.080; bouts: $F_{1,22}$=5.864, p=0.024; pile height: $F_{1,24}$=7.225, p=0.013; **Fig. S3A-C**), and also an increased latency to first bury after being shocked ($F_{1,24}$=6.907, p=0.015; **Fig. S3D**). Accordingly, during the subsequent retention test females were also slower to initiate defensive responses ($F_{1,24}$=8.519, p=0.008, **Fig. S3H**), and they created shorter piles than males ($F_{1,24}$=5.226, p=0.032, **Fig. S3G**).

Figure 3: VP^GABA Inhibition During Training Does Not Consistently Affect Acquisition of Defensive Burying **(A)** Diagram shows behavioral protocol, where CNO or VEH injections were given during shock probe training in Control or Inhibited groups, then a drug-free memory test was conducted with a dormant probe. Training day behaviors are shown for defensive burying duration **(B)**, bouts **(C)**, pile height **(D)**, and latency **(E)**, and also probe investigation duration **(F)**. The same behaviors on the memory test are shown in **(G-K).** No statistically significant results were found.

VP^GABA inhibition did not sex-dependently influence defensive behaviors emitted during the
acquisition session, as no Sex × DREADD interactions were observed (all Fs<0.600, ps>0.446).
However, during the drug-free memory retention test conducted 24hrs later, there was a
significant Sex × Inhibition Group interaction, such that prior VP^GABA inhibition appeared to sex-
dependently affect most measures of defensive behavior (duration: $F_{1,21}$=11.76, p=0.003; bouts:
$F_{1,21}$=9.446, p=0.006; pile height: $F_{1,24}$=7.832, p=0.010; latency: $F_{1,24}$=2.789, p=0.108; **Fig. S3E-**
G). Posthoc analysis of behavior on this memory retention test in each Sex showed that males
which were VP^GABA-Inhibited during training spent less time burying on the retention test than
Control males (Holm-Sidak Posthoc: t_{21}=2.471, p=0.044), while females inhibited during training
spent more time burying than Control females (t_{21}=2.450, p=0.044). Males inhibited during
training also had fewer bouts of burying during the retention test (t_{21}=2.559, p=0.036), but no
effect was seen in females on this measure. No significant differences in pile height during the
retention test were seen in females (t_{21}= 1.783, p=0.167), while males showed a trend towards a
decrease in pile height (t_{21}=2.250, p=0.067). Notably, unlike the previous experiment where
effects VP^GABA inhibition on expression of probe fear were specific to active defensive behaviors,
a trend toward a similar sex difference in probe investigations during the memory retention test
was also seen (Sex x Inhibition interaction: $F_{1,24}$=3.369, p=0.079). The significance of this
potential sex difference in responses to a probe initially encountered during VP^GABA inhibition is
unclear, so future studies with larger cohorts and complementary behavioral tasks will be
important to clarify whether sex mediates VP^GABA roles in memory formation or consolidation.
400
401   ***3.4 VP^GABA Inhibition Enhances Passive Defensive Freezing to a Shock-Predictive***
402   ***Auditory Cue***
403   Next we asked whether VP^GABA neurons are also involved in responses to cues that are
404   temporally- but not spatially-predictive of a shock, which elicit passive (freezing) rather than
405   active (probe burying) defensive responses.
406
Rats first learned to associate a 20sec auditory cue that terminated coincident with a foot shock
3 times during a fear acquisition session (low-shock: 0.3mA, or high-shock: 0.75mA). 48hrs later

they underwent a fear expression test in a novel context with VP^GABA Inhibition or Control
treatment. The shock-paired cue (without shock) was played 24 times, allowing examination of
VP^GABA inhibition effects on cued fear expression (response to 1st delivered cue) and extinction
(subsequent cues; **Fig. 4A)**.
413

Figure 4: VP^GABA Inhibition Enhances Freezing Responses to a Previously-Learned Auditory Fear Cue
(A) Diagram of the passive auditory fear expression protocol. During cue/shock Acquisition training, 3 cue/shock (0.3 or 0.75 mA) pairings were made, and 48hrs later, a fear expression test occurred in VP^GABA- Inhibited and Control rats, where the cue was played 24 times without additional shocks. **(B)** The low-shock intensity (0.3mA) cue yielded enhanced freezing in Inhibited rats relative to Controls during its first presentation (indicated by shaded area), and also across all 24 cue presentations. **(C)** Inhibited rats also froze more than Controls during the 90sec periods following each cue presentation. **(D)** The high-shock intensity (0.75mA) cue elicited similar freezing in Inhibited and Control rats during cue presentations, though **(E)** Inhibited rats froze more during post-cue periods. BL: 3min baseline prior to the first cue. Red asterix = main effect of VP^GABA Inhibition (Inhibited versus Control; p<0.05 in a 2-way ANOVA). Black asterix = cue presentations during or after which Inhibited rats froze more than Controls (p<0.05, Holm-Sidak posthoc).

415
3.4.1 Effects of VP^GABA Inhibition on Passive Defensive Responses to Auditory Threat Cues: In
the low-shock intensity cohort, VP^GABA inhibition (n=8M,9F) did not alter behaviors relative to
Control treatment (n=9M,5F) during a baseline period in the novel context (Inhibited versus
Control, Mann-Whitney test; U=110.5, p=0.103, **Fig. 4B**), suggesting no change in
generalization of fear to the novel chamber as a result of VP^GABA inhibition. When the low-
intensity shock-paired cue was played for the first time on this expression test, when rats
maximally expected to be imminently shocked, Inhibited rats froze more than Controls (U=56.5,
p=0.005. **Fig. 4B**, shaded area). Cues were played 23 more times without shocks in the
session, and Inhibited rats froze more than Controls during cues throughout this extinction
training (Main effect of Inhibition: $F_{1,31}$=6.472, p=0.016; including during the 1st, 17th, 22nd, and
23rd cue presentations; Holm-Šídák postdoc, 1st: t_{775}=2.976, p=0.003; 17th: t_{775}=2.486, p=0.013;
22nd: t_{775}=2.194, p=0.028; 23rd: t_{775}=3.390, p=0.001). Inhibited and Control groups otherwise
showed similar extinction rates (Main effect of Cue#: $F_{24,744}$=23.240, p<0.0001; No Inhibition x
Cue# interaction: $F_{24,744}$=1.048, p=0.400).

430
431 In addition to measuring behavior during 20sec cues, we also examined the 90sec periods
432 immediately after each cue, and found that VP^{GABA} inhibited rats froze more than Controls
433 across all cues (Main effect of Inhibition: $F_{1,31}=5.128$, p=0.031; **Fig. 4C**)**,** especially late in the
434 session (Inhibition x Cue# interaction: $F_{24,744}=1.906$, p=0.006; Holm-Sidak posthoc significance
435 for cues 16, and 19-24 (ts>2.106; ps<0.036).
436
For the cohort receiving high-intensity shocks during training, VP^{GABA} inhibition (n=9M,7F) during
the expression test did not notably alter behavior relative to Controls (n=9M,10F) during
baseline (U=122, p=0.293; **Fig. 4D**), during the first cue presentation (U=196.5, p=0.930;
shaded area), or across the 24 cue presentations (no main effect of Inhibition: $F_{1,33}=1.701$,
p=0.201, nor Inhibition x Cue# interaction: $F_{24,792}=0.629$, p=0.916, **Fig. 4D**), potentially due to a
ceiling effect caused by high rates of freezing in both groups. However, during post-cue periods,
increased freezing in Inhibited rats was seen in early, but not late extinction trials (Inhibition x
Cue# interaction: $F_{23,759}=2.316$, p=0.001, **Fig. 4E**), potentially suggesting that VP^{GABA} inhibition
caused "spillover" of fear into these otherwise relatively "safe" post-cue periods. Holm-Sidak
posthocs confirmed elevated post-cue freezing during cues 3-5 and 19 (ts>1.981; ps<0.048).
447
448 Effects of VP^{GABA} inhibition on expression of auditory cue-elicited freezing did not appear to
449 result from group differences in Pavlovian training, prior to VP^{GABA} inhibition. Inhibited and
450 Control rats behaved similarly during these training sessions, with equivalent freezing during
451 baseline (Us>119, ps>0.05), during each of the three 20sec cue presentations (Us>82.5,
452 ps>0.05), and during 90sec periods following each shock (Us>97.5, ps>0.05; **Fig. S1C,D**).
453
VP^{GABA} inhibition enhancement of passive defensive responses to previously-learned auditory
threat cues was specific to freezing, as other measured behaviors in the low-shock cohort were
either unaffected by VP^{GABA} inhibition (rearing, head movements, escape attempts), or they
were decreased at the expense of increased freezing. Sniffing was decreased during the 1st cue
(U=64, p=0.009), and grooming was decreased during the 24 cues (main effect of Inhibition:
$F_{1,31}=10.06$, p=0.004) and the 24 post-cue periods ($F_{1,31}=7.648$, p=0.010). In the high-shock
cohort, decreases in nonfreezing behaviors were only seen in the post-cue interval and
dependent on both a VP^{GABA} inhibition and time of the post-cue interval during the extinction
session (Inhibition x Cue# interaction, rearing: $F_{23,943}=1.780$, p=0.013; head movement:
$F_{23,943}=1.697$, p=0.022). We note that defensive burying was never seen in this experiment,
since a grid floor was placed above bedding in test chambers.
465
466 *3.4.2 Sex-Specific Effects of Passive Fear Expression and VP^{GABA} Inhibition:* Females froze
467 more than males in general during auditory fear cue experiments, with increased freezing during
468 cues (**Fig S4,** main effect of Sex in low-shock cohort: $F_{1,28}=7.945$, p=0.009; high-shock cohort:
469 $F_{1,31}=40.820$, p<0.0001), and in post-cue periods (low-shock cohort: $F_{1,28}=9.943$, p=0.004; high-
470 shock cohort: $F_{1,31}=0.597$, p=0.446).
471
VP^{GABA} inhibition similarly enhanced freezing to the 1st cue presentation during recall in the low-
shock cohort (Main Effect of Inhibition: $F_{1,29}=9.003$, p=0.006) in both sexes (Sex x Inhibition
interaction: $F_{1,29}=0.224$, p=0.639 **Fig. S4A**). However, there was some evidence at VP^{GABA}
inhibition had greater ability to augment conditioned freezing to the low intensity shock-paired
cue in females, relative to males over the entire extinction session (**Fig. S4**; Sex x Inhibition
interaction; low-shock cohort: $F_{1,28}=4.479$, p=0.043; high-shock cohort: $F_{1,31}=0.754$, p=0.392).
Similar effects were observed in post-cue periods (low-shock cohort: $F_{1,29}=5.883$, p=0.022; high-
shock cohort: $F_{1,31}=0.962$, p=0.334). This intriguing finding is a rare example of a potentially sex-
dependent effect of VP^{GABA} inhibition which should be replicated.

3.5 VP^{GABA} Inhibition Effects on Neural Activity in LHb, and in Neighboring VP Neurons

3.5.1 Effects of Probe Threat and VP^{GABA} Inhibition on Fos in VP Efferent Targets: We next sought to investigate the neural circuits that may mediate augmented threat responses when VP^{GABA} neurons are inhibited. To do so, we examined Fos expression that was elicited by a threatening probe relative to a probe that when previously encountered did not deliver shock (Threat versus No Threat), and/or by VP^{GABA} inhibition (GAD1:Cre+CNO versus GAD1:Cre+VEH). We quantified Fos in major targets of VP^{GABA} efferent projections (LHb and MDT), and in subpopulations of VP neurons (**Fig. 5A; Fig. S5**).

We first examined Fos in the LHb, an aversion-related brain region to which VP neurons (including both VP^{GABA} and VP^{glutamate} subpopulations) strongly project (Faget et al., 2018; Groenewegen et al., 1993; Haber et al., 1993; Liu et al., 2020; Root et al., 2014; Tooley et al., 2018), and compared it to another VP target linked more to memory and cognition rather than to aversion, the MDT (Corbit et al., 2003; Floresco et al., 1999b; Leung et al., 2024; Mitchell & Chakraborty, 2013; Ostlund & Balleine, 2008; Parnaudeau et al., 2018; Root et al., 2015; Wolff & Halassa, 2024). Threat (i.e. exposure to a probe that previously delivered shock, or which did not) strongly impacted LHb Fos (2-way ANOVA; main effect of Threat: $F_{1,13}=33.80$, $p<0.0001$), with VP^{GABA} inhibition significantly enhancing LHb Fos for the shock probe only (Drug x Threat Interaction: $F_{1,13}=15.01$, $p=0.002$; Holm-Šídák postdoc for VP^{GABA} inhibition effect on response to Threat: $t_{13}=3.729$, $p=0.005$; or No Threat: $t_{13}=1.835$, $p=0.090$; **Fig. 5B,C**). These effects appeared to be specific to LHb, as in MDT there were no main effects of Threat ($F_{1,12}=1.475$, $p=0.2479$), VP^{GABA} inhibition ($F_{1,12}=2.665$, $p=0.1285$), or interactions of these variables ($F_{1,12}=0.9897$, $p=0.3394$; **Fig. 5D**).

3.5.2 Effects of Probe Threat and VP^{GABA} Inhibition on Fos in VP Subpopulations: We used our ability to Cre-dependently label VP^{GABA} neurons with mCherry to identify GAD1+ cells within VP (**Fig. 5E**), and to distinguish them from unlabeled cells in the immediate vicinity of virus injections, which are presumably largely non-GABAergic. Non-GABA neurons in the central VP region quantified here are 2-3x more likely to be glutamatergic than cholinergic (Faget et al., 2018; Tooley et al., 2018). Although we did not confirm their identity here, we refer to these unlabeled cells as putative VP^{nonGABA} neurons.

Relative to VEH, CNO significantly decreased activity in VP^{GABA} (mCherry+) cells regardless of Threat (main effect of Drug: $F_{1,13}=9.128$, $p=0.0001$; no Drug x Threat interaction: $F_{1,13}=0.071$, $p=0.7935$; **Fig. 5F**), which is consistent with our prior verifications that chemogenetic inhibition in this transgenic system is capable of inhibiting VP^{GABA} neuron activity (Farrell et. al. 2021). We also found that Threat itself elevated activity in VP^{GABA} cells (main effect of Threat: $F_{1,13}=9.005$, $p=0.010$).

Fos expression in non mCherry-labeled, putative VP^{nonGABA} neurons was also significantly increased by Threat (main effect of Threat: $F_{1,13}=15.32$, $p=0.002$). However, in the presence of the threatening probe, putative VP^{nonGABA} neurons were further activated by inhibition of their VP^{GABA} neighbors—an effect that was not seen in these neurons in the absence of the shock-associated probe (Drug x Threat Interaction: $F_{1,12}=10.04$, $p=0.008$; Holm-Šídák postdoc for CNO versus VEH in Threat condition: $t_{12}=3.278$, $p=0.013$; **Fig. 5G**).

Although we were unable to examine Fos in confirmed VP^{glutamate} neurons, we were able to quantify Fos in VP acetylcholine cells (VP^{ChAT}), which comprise the remainder of non-GABA, non-glutamate neurons in VP (Faget et al., 2018; Tooley et al., 2018). VP^{GABA} inhibition (VEH/CNO) did not alter responses of VP^{ChAT} neurons to Threat ($t_9=1.342$, $p=0.380$), nor did

532 VP^{ChAT} Fos differ in either Threat group compared to No Threat rats that received VEH (one way
533 ANOVA: $F_{2,9}$=0.390, p=0.688; Threat+CNO group was not included in this control experiment;
534 **Fig. S5A**). This finding suggests that neither Threat nor VP^{GABA} inhibition appears to alter
535 activity in VP^{ChAT} neurons, suggesting that they are unlikely to account for the clear group
536 differences seen in putative VP^{nonGABA} neurons.
537
3.5.3 Relationship Between Activity in VP Efferent Targets and VP Subpopulations: We also
examined how Fos in VP^{GABA}, putative VP^{nonGABA}, and VP^{ChAT} populations relates to Fos in VP's
efferent targets LHb and MDT within the same animals, thus pointing toward potential circuit
mechanisms that could underlie the observed behavioral effects of VP^{GABA} inhibition in
responses to Threat. There was a clear correlation between Fos in putative VP^{nonGABA} cells and
LHb (R^2=0.468, p=0.004, **Fig. 5H, right**), but no such relationship between LHb and VP^{GABA}
(R^2=0.007, p=0.764, **Fig. 5H, left**), or between LHb and VP^{ChAT} (R^2=0.107, p=0.326, **Fig. S5B**).
No such correlation was seen between Fos in MDT and any of these VP subpopulations either
(R^2s<0.031; Ps>0.517; **Fig. 5I, S5C**). These findings are consistent with activity in LHb (but not
MDT) being related to activity in non-GABA, non-acetylcholine VP neurons in VP, which could
be glutamatergic. However, it is not clear from these data whether VP^{GABA} inhibition yielded an
augmented Threat response in LHb via removal of direct LHb inhibitory inputs, via disinhibition
of excitatory VP projections to LHb, or via another circuit mechanism.

Figure 5: VP^GABA Inhibition Increases Threat-induced Fos in the LHb, and in Putative VP^nonGABA Cells: (A) Schematic of Fos experiment protocol. **(B)** Representative images of LHb Fos staining in Threat+VEH and Threat+CNO conditions, indicated with yellow box. **(C)** Both Threat and VP^GABA inhibition increase Fos in LHb, and Inhibition also augmented Threat-induced LHb Fos without affecting LHb Fos in the absence of Threat. **(D)** VP^GABA Inhibition and Threat do not significantly impact Fos in MDT. **(E)** Representative images of VP cell populations quantified in the immediate vicinity of virus injection sites in central VP, which were stained for Fos (green, identifying active neurons), mCherry (red, identifying VP^GABA neurons), and DAPI (blue, cell nuclei). Yellow arrows indicate active VP^GABA cells; green arrows indicate active, putatively VP^nonGABA cells. **(F)** CNO inhibited mCherry-labeled VP^GABA cells, and Threat activated them, but no Threat specific changes were caused by CNO. **(G)** Putative VP^nonGABA cells were activated by Threat, and inhibiting VP^GABA cells with CNO selectively augmented this Threat response. **(H)** Correlation between Fos in LHb and VP^GABA (*left*) or putative VP^nonGABA (*right*) populations from the same rat are shown. **(I)** Correlation between Fos in MDT and VP^GABA (*left*) or putative VP^nonGABA (*right*) populations from the same rats are shown. 3V=3rd ventricle. Two-Way ANOVA and Holm-Sidak postdoc, or Pearson correlation significance: *p<0.05, **p<0.0, ***p<0.001; two-tailed. Significant main effect of Threat is indicated on X-axis label, and main effect of Drug is shown in legend inset.

4. Discussion

Here we show for the first time that in addition to promoting reward seeking, VP[GABA] neurons are also capable of regulating aversively motivated behaviors. Our findings suggest that chemogenetically inhibiting VP[GABA] neurons "releases the brakes" on context-appropriate defensive responses to learned threats, likely via direct or indirect interactions with LHb. This suggests that VP[GABA] cells may promote reward pursuit both by imbuing reward-associated stimuli with incentive salience (Berridge & Robinson, 2003; Farrell et al., 2018; Smith et al., 2009), but also by concurrently suppressing conditioned fear and aversion that would otherwise limit reward seeking. We found that VP[GABA] inhibition consistently enhanced either active or passive defensive responses to shock threat cues, depending on which response was most appropriate for the perceived threat encountered. In contrast, inhibiting VP[GABA] did not consistently alter acquisition of threat learning, and non-defensive behaviors were unaffected, suggesting a specific role for VP[GABA] in suppressing fear and aversion elicited by threat cues that were spatially or temporally associated with a noxious shock. These findings lend insight into VP modulation of motivation, and may also reflect broader principles by which intermeshed subcortical circuits of valanced motivation govern affective decision making (Carlezon & Thomas, 2009; Gordon-Fennell & Stuber, 2021; Waraczynski, 2016), and thus inform future strategies for intervening in disorders like addiction and depression.

4.1 VP[GABA] Inhibition Augments Defensive Responses to Learned Threats

VP[GABA] neurons have been described as primarily helping drive appetitive behavior like highly-motivated reward seeking (Faget et al., 2018, 2024; Prasad et al., 2020; Scott et al., 2023; Soares-Cunha & Heinsbroek, 2023; Stephenson-Jones et al., 2020; Wulff et al., 2019), while intermingled VP[glutamate] neurons instead seem to mediate aversive and avoidance processes that suppress reward pursuit (Levi et al., 2020; Liu et al., 2020; Tooley et al., 2018; Wang et al., 2020). Other reports have previously found a role for undefined VP cells in fear or aversion (Hernández-Jaramillo et al., 2024; Moaddab et al., 2021; Russell & McDannald, 2024), and one previously showed VP[GABA] cells respond to noxious stimuli (Faget et al., 2024), but their role in conditioned aversive motivation or defensive behaviors was unknown.

We previously showed that chemogenetically inhibiting VP[GABA] in rats failed to alter either primary unconditioned responses to shocks, or to markedly alter instrumental avoidance or escape from foot shocks (Farrell et al., 2021), and here we recapitulated the finding that primary behavioral reactions to shock do not require VP[GABA]. Instead, we found an unappreciated role for these neurons in conditioned threat responses, as inactivating them markedly *enhanced* defensive responses to shock-associated cues. This implies that VP[GABA] neuron activity, in addition to directly promoting appetitive responses, may also engage safety signaling mechanisms that dampen aversive motivation elicited by anticipated dangers (Duvarci, 2024; Luo et al., 2018; Sangha et al., 2020).

In the natural world of rats where opportunities and threats often co-occur, this type of push-pull mechanism would be adaptive in situations in which reward pursuit must be prioritized in the face of danger. Consistent with this idea, when we previously asked how inhibiting VP[GABA] influences risky decision making, we found a shift to a more conservative strategy—promoting choice of a small but safe reward (one food pellet) over a larger, but potentially dangerous option (two pellets, plus a chance of foot shock)(Farrell et al., 2021). We interpreted this as primarily reflecting suppression of motivation to receive reward when VP[GABA] was inhibited, but the present data suggest that such decision making may also have been influenced by an augmented perception of threat. We therefore propose that VP[GABA], and potentially other pallidal, hypothalamic, and extended amygdala GABA circuits with a seemingly similar specific

role in promoting reward seeking and consumption (Gordon-Fennell & Stuber, 2021; Jennings et al., 2013; Nieh et al., 2015; Nieh, Vander Weele, et al., 2016; Waraczynski, 2016; Zhang et al., 2021), may in fact perform a more nuanced function than previously appreciated, by balancing conflicting motivational influences to guide high-stakes decision making.

Given the VP's long-known role in motor activation (Austin & Kalivas, 1990; Churchill et al., 1998; Heimer et al., 1982), it is particularly striking that inhibiting VP^{GABA} led to either active, threat-directed defensive treading, or instead to passive immobility in response to threat— depending on the nature of the shock cue presented. The shock probe used here was a localized stimulus that was spatially paired with shock—analogous to naturalistic threats like a predator chasing a rat down an underground tunnel (Berridge et al., 1988; Craft et al., 1988; Pinel & Treit, 1978). Such proximate, localized threats must either be escaped, or if this is impossible, opposed—for example by pushing and throwing any available material toward the threat, as occurs here when rats attempt to bury the probe in bedding material. VP^{GABA} inhibition selectively augmented this defensive behavior in response to the learned probe threat. In contrast, when a diffuse auditory cue is played just prior to delivery of a foot shock, most rats exhibit freezing behavior when the cue is re-encountered. When VP^{GABA} was inhibited under these circumstances, passive freezing was significantly augmented. Non-defensive behaviors were not altered by VP^{GABA} inhibition in either threat task, nor were defensive behaviors inappropriate to the threat at hand (freezing to the probe, treading to the auditory cue). Therefore, VP^{GABA} inhibition seems to augment aversive motivation in a manner that is upstream of the action selection and generation circuits which determine the best behavioral manifestation of fear under the circumstances. A similar position in threat-processing circuits has been proposed for hypothalamic corticotropin releasing factor cells (Bains et al., 2015; Füzesi et al., 2016), but how exactly these interact with VP, or how either population interacts with downstream mechanisms generating specific defensive actions remains to be determined.

Interestingly, inhibiting VP^{GABA} did not increase freezing in a novel context before cues were presented, suggesting no unusual generalization of fear to this environment, or altered perception of mere novelty as threatening. However, we did observe that VP^{GABA} inhibition caused rats to continue to freeze even after termination of shock cues that were played without delivery of the predicted shock, and to partially oppose extinction of these cues. This suggests that VP^{GABA} inhibition may prolong or intensify cue-elicited fear, thereby interfering with the normal resolution or extinction of the fear response. VP^{GABA} neurons are thus involved in shaping not only the selection of appropriate defensive motor strategies, but also the ability of rats to appropriately terminate defensive states once immediate danger has passed, and to extinguish conditioned fear responses.

Furthermore, we interpret these results to indicate that VP^{GABA} inhibition does not elicit *de novo* fear states in the absence of actual environmental threats. In prior reports, we and others showed that VP^{GABA} inhibition reduces highly motivated, but not less-motivated reward seeking (Farrell et al., 2021, 2022)—if this decreased reward pursuit was secondary to a fear-like state, one would expect all types of reward seeking to be similarly affected. Likewise, inhibiting VP^{GABA} in the absence of threat cues, here or in our prior work, does not elicit either spontaneous freezing in operant boxes, or spontaneous burying in bedding-filled tub cages (Farrell et al., 2021, 2022). We also failed to see activation of LHb by VP^{GABA} inhibition when a non-threatening probe was encountered, but only augmentation of threat-elicited LHb activity. Therefore, we conclude that VP^{GABA} inhibition effects are context- and cue-dependent, and that their behavioral manifestation depends critically upon the affective valence of the circumstances at hand.

4.2 VP^GABA Inhibition Does Not Alter Defenses Against, or Learning About Unconditioned Threats

We found that VP^GABA inhibition did not alter defensive responses directed at an unconditioned and recently experienced threat, as avoidance and burying of a novel, still-electrified probe was unaffected. This suggests that VP^GABA does not mediate affective or sensory aspects of the unconditioned experience of punishment as previously reported (Farrell et al., 2021), and it demonstrates that rats are capable of mounting acute, active defenses in the absence of VP^GABA. In addition, subsequent conditioned responses to the shock probe which was encountered while VP^GABA was inactivated were also largely unaltered. This suggests that disrupting VP^GABA does not consistently affect the ability to form or consolidate aversive Pavlovian memories. There is some evidence that VP plays a role in other types of learning (Dusa et al., 2022; Floresco et al., 1999a; Kaplan et al., 2020; Leung & Balleine, 2013; Macpherson et al., 2019; Péczely et al., 2014; Roman-Ortiz et al., 2021; Stephenson-Jones, 2019; Stephenson-Jones et al., 2020), though most of these studies did not disambiguate motivational from learning processes, or VP cell types. One report showed that optogenetically stimulating VP^GABA suppressed auditory cue-induced freezing without affecting acquisition of new fear learning (Roman-Ortiz et al., 2021), consistent with the present result. However, another showed that stimulating predominantly GABAergic VP enkephalin neurons disrupts formation of an inhibitory avoidance memory (Macpherson et al., 2019). Clearly, further work is required to fully understand the conditions in which VP^GABA is implicated in the complex processes involved in forming, consolidating, storing, and retrieving the many different types of memory (McGaugh, 2003; Sherry & Schacter, 1987; Squire, 2004; White & McDonald, 2002).

4.3 VP^GABA Inhibition "Releases Brakes" on LHb, and on Nearby "Non-GABA" Neurons?

The LHb, which helps encode negative affect and learning, and plays a role in aversion, avoidance, and anhedonia (Bromberg-Martin & Hikosaka, 2011; Li et al., 2013; Liu et al., 2020; Matsumoto & Hikosaka, 2009; Tooley et al., 2018; Y. Yang et al., 2018), is a major efferent target of VP^GABA as well as VP^glutamate neurons (Faget et al., 2018; Liu et al., 2020; Stephenson-Jones et al., 2020; Tooley et al., 2018). We therefore determined how neural activity in LHb, the MDT (another target of VP efferents), and in VP neuronal subpopulations is affected by exposure to a threatening shock prod, and how inhibiting VP^GABA cells affects this activity.

As expected, encountering a threatening probe increased activity in LHb relative to an inert probe, and we found that this activation was further potentiated by inhibiting VP^GABA. This is consistent with a disinhibition mechanism, whereby disrupting VP^GABA removes this direct inhibitory input to LHb, amplifying its response to threat. Accordingly, globally inhibiting VP^GABA neurons induces place avoidance (Faget et al., 2018), and activating other LHb GABA inputs induces place preference (Barker et al., 2017; Stamatakis et al., 2013). However, optogenetic activation of VP^GABA terminals in LHb failed to promote appetitive behavior (Faget et al., 2018), and optogenetic inhibition of this pathway strongly suppresses conditioned reward-seeking, and modestly reduces conditioned avoidance of a noxious stimulus (Stephenson-Jones et al., 2020). Here, we found that VP^GABA inhibition in the absence of a threat was not sufficient to activate LHb, suggesting that VP^GABA inhibition alone does not recruit LHb, but merely disinhibits its activity when it is recruited by relevant environmental threats. Interestingly, no clear effects of either threat or of VP^GABA inhibition were seen in MDT, another major target of VP thought to play a role in memory and cognition, rather than aversion and defensive responding.

Since we did not use pathway-specific approaches to examine functions of VP^GABA projections to LHb, we cannot establish whether VP^GABA inhibition impacts LHb via its direct projections, or via indirect network mechanisms. Indeed, our data raise the intriguing, though untested, possibility that one way in which VP^GABA inhibition could indirectly impact LHb is via disinhibition

of nearby populations of VP^{glutamate} neurons that also project there. VP^{glutamate} projections to LHb
mediate aversion- and avoidance-related processes (Faget et al., 2018; Heinsbroek et al., 2020;
Stephenson-Jones et al., 2020; Tooley et al., 2018; Wulff et al., 2019), and VP^{GABA} cells directly
inhibit their glutamatergic neighbors (Faget et al., 2018; Levi et al., 2020; Tripathi et al., 2013). It
is therefore possible that the activation of putative VP^{nonGABA} cells that we showed occurred after
VP^{GABA} inhibition, and which we found were unlikely to be cholinergic, reflects disinhibition of
local VP^{glutamate}, cells that project to LHb. This is also consistent with the correlation we saw
between Fos in putative VP^{nonGABA} and LHb, but not MDT. However, we were unable to identify
VP^{glutamate} cells here directly, and we cannot exclude the possibility that some of these cells are
unlabeled VP^{GABA} cells, so this hypothesis must be tested in future using appropriate methods
like RNA-based cell identification, vGlut2-based cell- and pathway-targeting approaches, or
physiological techniques.
717
718 *4.4 Sex Differences in VP^{GABA} Functions*
VP (DiBenedictis et al., 2020; Lee et al., 2024; Scott et al., 2025; L. Yang et al., 2024), and the
circuits in which it is embedded (Becker & Chartoff, 2019; Lim et al., 2004; Trainor, 2011;
Zachry et al., 2021), appear to differ based on sex; as does the prevalence of symptoms
defining many psychiatric disorders these circuits contribute to (Bangasser & Cuarenta, 2021;
Cosgrove et al., 2007; Green et al., 2018; Kessler, 2003; Trainor, 2011; Young & Pfaff, 2014).
We noticed VP-independent sex differences in threat response, such that females overall
exhibited more passive freezing to the associated auditory cue presentation, while males
exhibited more active burying of the probe cue. We found only subtle sex differences in the
effects of VP^{GABA} inhibition on threat responding, in that VP^{GABA} inhibited females froze more
than males to auditory fear cues and post-cue intervals throughout the extinction session, while
initial enhanced freezing to the 1st cue under inhibition was independent of sex. In this context it
is surprising that our lab's work in VP has rarely observed sex differences in the effects of VP
manipulations on drug and natural reward seeking (Farrell et al., 2019, 2021, 2022; Mahler et
al., 2014).
733
Another sex difference we did observe in the effects of VP^{GABA} inhibition stems from our
experiment in which VP^{GABA} was inhibited during initial learning about a shock probe, and
subsequent testing of learned responses to that probe in the absence of further VP
manipulation. VP^{GABA} inhibition did not alter responses in either sex to the shock-delivering
probe during their initial encounters with it, and defensive and avoidance behaviors in the
minutes following shock were unaffected. However, when rats' ability to remember this threat
learned under VP^{GABA} inhibition was queried, a possible sex difference emerged: enhanced
defensive burying in females, and decreased burying in males, consistent with a sex difference
in VP^{GABA} participation in memory consolidation. This is an exciting possibility, but given the low
sample sizes involved, and lack of converging evidence from other behaviors in these
experiments, we suggest caution in interpreting this finding. Future studies with larger sample
sizes, complementary behavioral tasks, and designs capable of dissociating memory
consolidation from performance (McGaugh, 1973) will be required to resolve the conditions in
which sex modulates VP roles in negative emotion, learning, and/or memory (Blanchard et al.,
1991).
749
750 *4.5 Caveats and Limitations*
These studies have additional limitations that should be noted, and potentially resolved with
further research. We limited the aversive stimuli tested here to shock delivered two different
ways—it would be valuable to learn whether VP^{GABA} inhibition also augments other responses to
learned (e.g. context fear conditioning) or innate threats (e.g predator odors or looming stimuli).
Likewise, naturalistic threat responding also depends critically on factors like threat imminence

or proximity which could be manipulated in follow-up studies, and other potential responses to
threats like escape or defensive aggression were not possible here, so it remains to be seen
how VP^{GABA} helps prioritize such strategies. In addition, VP^{GABA} inhibition here enhanced
freezing only to a low-intensity shock cue, which likely reflects a ceiling effect for freezing with
the high-shock intensity cue, but could also suggest more complex interactions. Regarding
neural activity data, we cannot exclude the possibility that CNO had DREADD-independent
effects on LHb Fos, since we did not include a DREADD-free control group in this experiment.
However, nonspecific CNO effects are unlikely to account at least for enhanced LHb Fos in
VP^{GABA} inhibited rats exposed to threat, since CNO did not augment Fos in unthreatened rats. In
addition, Fos analyses of rats exposed to threat were performed in males only, and should be
followed up with females, especially since as noted above the nature of sex differences in VP
need further clarification more generally. Although these experiments targeted DREADDs to a
region of VP spanning its rostrocaudal axis, prior reports have shown functional distinctions
between these subregions (Farrell et al., 2022; Ho & Berridge, 2014; Johnson et al., 1993;
Kupchik & Kalivas, 2013; Mahler et al., 2014; Panagis et al., 1995; Root et al., 2013; Smith &
Berridge, 2005; Stout et al., 2016; Zahm & Heimer, 1993) so future work should test whether
such dissociations may also relate to VP^{GABA} roles in aversive motivation.

773
4.6 Conclusions
Here we report a novel role for VP^{GABA} neurons in suppressing responses to threat cues,
complementing their established role in promoting reward seeking. These cells may therefore
facilitate commitment to reward pursuit even in the face of danger, thus prevent vacillation
between behavioral strategies once a decision to pursue reward is made. In addition to
clarifying roles for VP circuits in motivation, these findings might also be relevant to the
organization of wider subcortical circuits involving seemingly analogous intermixed populations
of GABA and glutamate cells that mediate reward and aversion, respectively (Gordon-Fennell &
Stuber, 2021; Hu, 2016; Root et al., 2015, 2020; Soares-Cunha & Heinsbroek, 2023; Wulff et
al., 2019). These data also open new avenues of investigation for understanding how VP^{GABA}
interacts with other VP cell populations involved in stress (Chang & Grace, 2014; Eckenwiler et
al., 2025; Ji et al., 2022), and with wider brain circuits of motivation, emotion, and learning.
Given that imbalance in motivational drives is likely involved in numerous psychiatric disorders
including addiction, depression, schizophrenia, and PTSD, understanding the functions of these
brain networks will be essential for developing next-generation strategies for treating these
devastating conditions.

5. Authorship Contribution Statement
EMR and MXM equally contributed to conceptualization and implementation of the project, and
with SVM analyzed data and prepared the manuscript. CMR, RKR, MRF, VAV, SAW, HRR, and
GJK contributed data to the project.

6. Acknowledgements
This work was supported by the following grants: NIDA R01DA055849, NIMH R01MH132680,
NIDA U01DA053826, NIDA P50DA044118-05A1, TRDRP T31IR1767; EMR Ford Foundation
Predoctoral Fellowship; MXM NSF GRFP DGE-1839285 and NINDS F99NS141399. We thank
Yiyan Xie and Erik Castillo for technical support, Tom Hnasko for clarifying conversations and
technical assistance, and NIDA/NIMH drug distribution programs for supplying CNO.

802
803
7. References

Austin, M. C., & Kalivas, P. W. (1990). Enkephalinergic and GABAergic modulation of motor
activity in the ventral pallidum. *The Journal of Pharmacology and Experimental*
Therapeutics, *252*(3), 1370–1377. https://doi.org/10.1016/S0022-3565(25)20204-X
Bains, J. S., Cusulin, J. I. W., & Inoue, W. (2015). Stress-related synaptic plasticity in the
hypothalamus. *Nature Reviews Neuroscience 2015 16:7*, *16*(7), 377–388.
https://doi.org/10.1038/nrn3881
Bangasser, D. A., & Cuarenta, A. (2021). Sex differences in anxiety and depression: circuits and
mechanisms. *Nature Reviews Neuroscience 2021 22:11*, *22*(11), 674–684.
https://doi.org/10.1038/s41583-021-00513-0
Barker, D. J., Miranda-Barrientos, J., Zhang, S., Root, D. H., Wang, H.-L., Liu, B., Calipari, E.
S., & Morales, M. (2017). Lateral Preoptic Control of the Lateral Habenula through
Convergent Glutamate and GABA Transmission. *Cell Reports*, *21*(7), 1757–1769.
https://doi.org/10.1016/j.celrep.2017.10.066
Becker, J. B., & Chartoff, E. (2019). Sex differences in neural mechanisms mediating reward
and addiction. *Neuropsychopharmacology*, *44*(1), 166–183.
https://doi.org/10.1038/S41386-018-0125-
6;SUBJMETA=1662,1689,378,5,631;KWRD=ADDICTION,MOTIVATION
Berridge, K. C., Fentress, J. C., & Treit, D. (1988). A triggered hyperkinesia induced in rats by
lesions of the corpus striatum. *Experimental Neurology*, *99*(2), 259–268.
https://doi.org/10.1016/0014-4886(88)90146-X
Berridge, K. C., & Robinson, T. E. (2003). Parsing reward. *Trends in Neurosciences*, *26*(9),
507–513. https://doi.org/10.1016/S0166-2236(03)00233-9
Blanchard, D. C., Shepherd, J. K., Carobrez, A. D. P., & Blanchard, R. J. (1991). Sex effects in
defensive behavior: Baseline differences and drug interactions. *Neuroscience &*
Biobehavioral Reviews, *15*(4), 461–468. https://doi.org/10.1016/S0149-7634(05)80132-0
Bromberg-Martin, E. S., & Hikosaka, O. (2011). Lateral habenula neurons signal errors in the
prediction of reward information. *Nature Neuroscience*, *14*(9), 1209–1218.
https://doi.org/10.1038/NN.2902
Carlezon, W. A., & Thomas, M. J. (2009). Biological substrates of reward and aversion: a
nucleus accumbens activity hypothesis. *Neuropharmacology*, *56 Suppl 1*(Suppl 1), 122–
132. https://doi.org/10.1016/J.NEUROPHARM.2008.06.075
Chang, C. H., & Grace, A. A. (2014). Amygdala-Ventral Pallidum Pathway Decreases Dopamine
Activity After Chronic Mild Stress in Rats. *Biological Psychiatry*, *76*(3), 223–230.
https://doi.org/10.1016/J.BIOPSYCH.2013.09.020
Churchill, L., Klitenick, M. A., & Kalivas, P. W. (1998). Dopamine depletion reorganizes
projections from the nucleus accumbens and ventral pallidum that mediate opioid-induced
motor activity. *Journal of Neuroscience*, *18*(19), 8074–8085.
https://doi.org/10.1523/JNEUROSCI.18-19-08074.1998
Corbit, L. H., Muir, J. L., & Balleine, B. W. (2003). Lesions of mediodorsal thalamus and anterior
thalamic nuclei produce dissociable effects on instrumental conditioning in rats. *The*
European Journal of Neuroscience, *18*(5), 1286–1294. https://doi.org/10.1046/J.1460-
9568.2003.02833.X

Cosgrove, K. P., Mazure, C. M., & Staley, J. K. (2007). Evolving Knowledge of Sex Differences
in Brain Structure, Function, and Chemistry. *Biological Psychiatry*, *62*(8), 847–855.
https://doi.org/10.1016/J.BIOPSYCH.2007.03.001

Craft, R. M., Howard, J. L., & Pollard, G. T. (1988). Conditioned defensive burying as a model
for identifying anxiolytics. *Pharmacology Biochemistry and Behavior*, *30*(3), 775–780.
https://doi.org/10.1016/0091-3057(88)90098-6

DiBenedictis, B. T., Cheung, H. K., Nussbaum, E. R., & Veenema, A. H. (2020). Involvement of
ventral pallidal vasopressin in the sex-specific regulation of sociosexual motivation in rats.
Psychoneuroendocrinology, *111*, 104462.
https://doi.org/10.1016/J.PSYNEUEN.2019.104462

Dusa, D., Ollmann, T., Kállai, V., Lénárd, L., Kertes, E., Berta, B., Szabó, Á., László, K., Gálosi,
R., Zagoracz, O., Karádi, Z., & Péczely, L. (2022). The antipsychotic drug sulpiride in the
ventral pallidum paradoxically impairs learning and induces place preference. *Scientific*
Reports, *12*(1), 19247. https://doi.org/10.1038/s41598-022-23450-z

Duvarci, S. (2024). Dopaminergic circuits controlling threat and safety learning. *Trends in*
Neurosciences, *47*(12), 1014–1027.
https://doi.org/10.1016/J.TINS.2024.10.001/ASSET/54F2EAC0-FB9C-4738-8BA3-
E5CA2FB60E71/MAIN.ASSETS/GR2.JPG

Eckenwiler, E. A., Ingebretson, A. E., Stolley, J. J., Fusaro, M. A., Romportl, A. M., Ross, J. M.,
Petersen, C. L., Kale, E. M., Clark, M. S., Schattauer, S. S., Zweifel, L. S., & Lemos, J. C.
(2025). Corticotropin-Releasing Factor Release From a Unique Subpopulation of Accumbal
Neurons Constrains Action-Outcome Acquisition in Reward Learning. *Biological Psychiatry*,
97(6), 637–650. https://doi.org/10.1016/J.BIOPSYCH.2024.08.006

Faget, L., Oriol, L., Lee, W.-C., Zell, V., Sargent, C., Flores, A., Hollon, N. G., Ramanathan, D.,
& Hnasko, T. S. (2024). Ventral pallidum GABA and glutamate neurons drive approach and
avoidance through distinct modulation of VTA cell types. *Nature Communications*, *15*(1),
4233. https://doi.org/10.1038/S41467-024-48340-Y

Faget, L., Zell, V., Souter, E., McPherson, A., Ressler, R., Gutierrez-Reed, N., Yoo, J. H.,
Dulcis, D., & Hnasko, T. S. (2018). Opponent control of behavioral reinforcement by
inhibitory and excitatory projections from the ventral pallidum. *Nat Commun*, *9*(1), 849.
https://doi.org/10.1038/s41467-018-03125-y

Fanselow, M. S. (1979). Naloxone attenuates rat's preference for signaled shock. *Physiological*
Psychology, *7*(1), 70–74.

Fanselow, M. S. (1980). Conditional and unconditional components of post-shock freezing. *The*
Pavlovian Journal of Biological Science: Official Journal of the Pavlovian, *15*(4), 177–182.

Farrell, M. R., Esteban, J. S. D., Faget, L., Floresco, S. B., Hnasko, T. S., & Mahler, S. V.
(2021). Ventral Pallidum GABA Neurons Mediate Motivation Underlying Risky Choice.
Journal of Neuroscience, *41*(20), 4500–4513. https://doi.org/10.1523/JNEUROSCI.2039-
20.2021

Farrell, M. R., Ruiz, C. M., Castillo, E., Faget, L., Khanbijian, C., Liu, S., Schoch, H., Rojas, G.,
Huerta, M. Y., Hnasko, T. S., & Mahler, S. V. (2019). Ventral pallidum is essential for
cocaine relapse after voluntary abstinence in rats. *Neuropsychopharmacology*, *44*(13),
2174–2113. https://doi.org/10.1038/s41386-019-0507-4

Farrell, M. R., Schoch, H., & Mahler, S. V. (2018). Modeling cocaine relapse in rodents:
Behavioral considerations and circuit mechanisms. *Progress in Neuro-*
Psychopharmacology and Biological Psychiatry, *87*(Pt A), 33–47.
https://doi.org/10.1016/j.pnpbp.2018.01.002

Farrell, M. R., Ye, Q., Xie, Y., Esteban, J. S. D., & Mahler, S. V. (2022). Ventral pallidum GABA
neurons bidirectionally control opioid relapse across rat behavioral models. *Addiction*
Neuroscience, *3*. https://doi.org/10.1016/J.ADDICN.2022.100026

Floresco, S. B., Braaksma, D. N., & Phillips, A. G. (1999a). Involvement of the Ventral Pallidum
in Working Memory Tasks With or Without a Delay. *Annals of the New York Academy of*
Sciences, *877*(1), 711–716. https://doi.org/10.1111/J.1749-6632.1999.TB09308.X

Floresco, S. B., Braaksma, D. N., & Phillips, A. G. (1999b). Thalamic-cortical-striatal circuitry
subserves working memory during delayed responding on a radial arm maze. *The Journal*
of Neuroscience : The Official Journal of the Society for Neuroscience, *19*(24), 11061–
11071. https://doi.org/10.1523/JNEUROSCI.19-24-11061.1999

Fucich, E. A., & Morilak, D. A. (2018). Shock-probe Defensive Burying Test to Measure Active
versus Passive Coping Style in Response to an Aversive Stimulus in Rats. *Bio-Protocol*,
8(17). https://doi.org/10.21769/BIOPROTOC.2998

Füzesi, T., Daviu, N., Wamsteeker Cusulin, J. I., Bonin, R. P., & Bains, J. S. (2016).
Hypothalamic CRH neurons orchestrate complex behaviours after stress. *Nature*
Communications 2016 7:1, *7*(1), 1–14. https://doi.org/10.1038/ncomms11937

Gordon-Fennell, A., & Stuber, G. D. (2021). Illuminating Subcortical GABAergic and
Glutamatergic Circuits for Reward and Aversion. *Neuropharmacology*, *198*, 108725.
https://doi.org/10.1016/J.NEUROPHARM.2021.108725

Green, T., Flash, S., & Reiss, A. L. (2018). Sex differences in psychiatric disorders: what we can
learn from sex chromosome aneuploidies. *Neuropsychopharmacology 2018 44:1*, *44*(1), 9–
21. https://doi.org/10.1038/s41386-018-0153-2

Groenewegen, H. J., & Berendse, H. W. (1990). Connections of the subthalamic nucleus with
ventral striatopallidal parts of the basal ganglia in the rat. *Journal of Comparative*
Neurology, *294*(4), 607–622. https://doi.org/10.1002/CNE.902940408

Groenewegen, H. J., Berendse, H. W., & Haber, S. N. (1993). Organization of the output of the
ventral striatopallidal system in the rat: ventral pallidal efferents. *Neuroscience*, *57*(1), 113–
142.

Haber, S. N., Groenewegen, H. J., Grove, E. A., & Nauta, W. J. H. (1985). Efferent connections
of the ventral pallidum: Evidence of a dual striato pallidofugal pathway. *Journal of*
Comparative Neurology, *235*(3), 322–335.
https://doi.org/https://doi.org/10.1002/cne.902350304

Haber, S. N., Lynd-Balta, E., & Mitchell, S. J. (1993). The organization of the descending ventral
pallidal projections in the monkey. *Journal of Comparative Neurology*, *329*(1), 111–128.
https://doi.org/https://doi.org/10.1002/cne.903290108

Haber, S. N., & Nauta, W. J. H. (1983). Ramifications of the globus pallidus in the rat as
indicated by patterns of immunohistochemistry. *Neuroscience*, *9*(2), 245–260.
https://doi.org/https://doi.org/10.1016/0306-4522(83)90291-9

Heimer, L., Switzer, R. D., & Van Hoesen, G. W. (1982). Ventral striatum and ventral pallidum:
Components of the motor system? *Trends in Neurosciences*, *5*, 83–87.

Heimer, L., Zahm, D. S., Churchill, L., Kalivas, P. W., & Wohltmann, C. (1991). Specificity in the projection patterns of accumbal core and shell in the rat. *Neuroscience*, *41*(1), 89–125. https://doi.org/10.1016/0306-4522(91)90202-Y

Heinsbroek, J. A., Bobadilla, A.-C., Dereschewitz, E., Assali, A., Chalhoub, R. M., Cowan, C. W., & Kalivas, P. W. (2020). Opposing Regulation of Cocaine Seeking by Glutamate and GABA Neurons in the Ventral Pallidum. *Cell Reports*, *30*(6), 2018-2027.e3. https://doi.org/https://doi.org/10.1016/j.celrep.2020.01.023

Hernández-Jaramillo, A., Illescas-Huerta, E., & Sotres-Bayon, F. (2024). Ventral Pallidum and Amygdala Cooperate to Restrain Reward Approach under Threat. *Journal of Neuroscience*, *44*(23). https://doi.org/10.1523/JNEUROSCI.2327-23.2024

Ho, C. Y., & Berridge, K. C. (2014). Excessive disgust caused by brain lesions or temporary inactivations: mapping hotspots of the nucleus accumbens and ventral pallidum. *The European Journal of Neuroscience*, *40*(10), 3556–3572. https://doi.org/10.1111/EJN.12720

Hu, H. (2016). Reward and Aversion. *Annual Review of Neuroscience*, *39*(Volume 39, 2016), 297–324. https://doi.org/10.1146/ANNUREV-NEURO-070815-014106/CITE/REFWORKS

Jennings, J. H., Sparta, D. R., Stamatakis, A. M., Ung, R. L., Pleil, K. E., Kash, T. L., & Stuber, G. D. (2013). Distinct extended amygdala circuits for divergent motivational states. *Nature*, *496*(7444), 224–228. https://doi.org/10.1038/nature12041

Ji, M. J., Gao, Z. Q., Yang, J., Cai, J. H., Li, K. X., Wang, J., Zhang, H., Zhou, C. H., Cao, J. L., & Liu, C. (2022). Dynorphin promotes stress-induced depressive behaviors by inhibiting ventral pallidal neurons in rats. *Acta Physiologica*, *236*(3), e13882. https://doi.org/10.1111/APHA.13882;JOURNAL:JOURNAL:17481716A;REQUESTEDJOURNAL:JOURNAL:17481716;PAGE:STRING:ARTICLE/CHAPTER

Johnson, P. I., Stellar, J. R., & Paul, A. D. (1993). Regional reward differences within the ventral pallidum are revealed by microinjections of a mu opiate receptor agonist. *Neuropharmacology*, *32*(12), 1305–1314. https://doi.org/10.1016/0028-3908(93)90025-X

Kalivas, P. W., Churchill, L., & Klitenick, M. A. (1993). GABA and enkephalin projection from the nucleus accumbens and ventral pallidum to the ventral tegmental area. *Neuroscience*, *57*(4), 1047–1060.

Kaplan, A., Mizrahi-Kliger, A. D., Israel, Z., Adler, A., & Bergman, H. (2020). Dissociable roles of ventral pallidum neurons in the basal ganglia reinforcement learning network. *Nature Neuroscience*, *23*(4), 556–564.

Kessler, R. C. (2003). Epidemiology of women and depression. *Journal of Affective Disorders*, *74*(1), 5–13. https://doi.org/10.1016/S0165-0327(02)00426-3

Kupchik, Y. M., & Kalivas, P. W. (2013). The rostral subcommissural ventral pallidum is a mix of ventral pallidal neurons and neurons from adjacent areas: an electrophysiological study. *Brain Structure & Function*, *218*(6), 1487–1500. https://doi.org/10.1007/S00429-012-0471-9

Lammel, S., Lim, B. K., Ran, C., Huang, K. W., Betley, M. J., Tye, K. M., Deisseroth, K., & Malenka, R. C. (2012). Input-specific control of reward and aversion in the ventral tegmental area. *Nature*, *491*(7423), 212–217. https://doi.org/10.1038/nature11527

Lawson, K. A., Flores, A. Y., Hokenson, R. E., Ruiz, C. M., & Mahler, S. V. (2021). Nucleus Accumbens Chemogenetic Inhibition Suppresses Amphetamine-Induced Ultrasonic

Vocalizations in Male and Female Rats. *Brain Sciences 2021, Vol. 11, Page 1255*, *11*(10), 1255. https://doi.org/10.3390/BRAINSCI11101255

Lawson, K. A., Ruiz, C. M., & Mahler, S. V. (2023). A head-to-head comparison of two DREADD agonists for suppressing operant behavior in rats via VTA dopamine neuron inhibition. *Psychopharmacology*, *240*(10), 2101–2110. https://doi.org/10.1007/S00213-023-06429-0/FIGURES/3

Lee, J. D. A., Reppucci, C. J., Huez, E. D. M., Bredewold, R., & Veenema, A. H. (2024). Sex differences in the structure and function of the vasopressin system in the ventral pallidum are associated with the sex-specific regulation of social play behavior in juvenile rats. *Hormones and Behavior*, *163*, 105563. https://doi.org/10.1016/J.YHBEH.2024.105563

Leung, B. K., & Balleine, B. W. (2013). The Ventral Striato-Pallidal Pathway Mediates the Effect of Predictive Learning on Choice between Goal-Directed Actions. *The Journal of Neuroscience*, *33*(34), 13848. https://doi.org/10.1523/JNEUROSCI.1697-13.2013

Leung, B. K., Chieng, B., Becchi, S., & Balleine, B. W. (2024). A ventral pallidal-thalamocortical circuit mediates the cognitive control of instrumental action. *Current Biology*, *34*(15), 3315-3326.e6. https://doi.org/10.1016/j.cub.2024.06.011

Levi, L. A., Inbar, K., Nachshon, N., Bernat, N., Gatterer, A., Inbar, D., & Kupchik, Y. M. (2020). Projection-specific potentiation of ventral pallidal glutamatergic outputs after abstinence from cocaine. *Journal of Neuroscience*, *40*(20), 1276–1285. https://doi.org/10.1523/JNEUROSCI.0612-20.2020

Li, K., Zhou, T., Liao, L., Yang, Z., Wong, C., Henn, F., Malinow, R., Yates, J. R., & Hu, H. (2013). βCaMKII in lateral habenula mediates core symptoms of depression. *Science*, *341*(6149), 1016–1020. https://doi.org/10.1126/SCIENCE.1240729/SUPPL_FILE/1240729.LI.SM.PDF

Lim, M. M., Murphy, A. Z., & Young, L. J. (2004). Ventral striatopallidal oxytocin and vasopressin V1a receptors in the monogamous prairie vole (Microtus ochrogaster). *Journal of Comparative Neurology*, *468*(4), 555–570. https://doi.org/10.1002/CNE.10973

Liu, B., Cao, Y., Wang, J., & Dong, J. (2020). Excitatory transmission from ventral pallidum to lateral habenula mediates depression. *The World Journal of Biological Psychiatry*, *21*(8), 627–633. https://doi.org/10.1080/15622975.2020.1725117

Luo, R., Uematsu, A., Weitemier, A., Aquili, L., Koivumaa, J., McHugh, T. J., & Johansen, J. P. (2018). A dopaminergic switch for fear to safety transitions. *Nature Communications 2018 9:1*, *9*(1), 1–11. https://doi.org/10.1038/s41467-018-04784-7

Macpherson, T., Mizoguchi, H., Yamanaka, A., & Hikida, T. (2019). Preproenkephalin-expressing ventral pallidal neurons control inhibitory avoidance learning. *Neurochemistry International*, *126*, 11–18.

Mahler, S. V, Vazey, E. M, Beckley, J. T., Keistler, C. R., McGlinchey, E. M., Kaufling, J., Wilson, S. P., Deisseroth, K., Woodward, J. J., & Aston-Jones, G. (2014). Designer receptors show role for ventral pallidum input to ventral tegmental area in cocaine seeking. *Nature Neuroscience*, *17*(4), 577–585.

Martinez, M. X., Alizo Vera, V., Ruiz, C. M., Floresco, S. B., & Mahler, S. V. (2024). Adolescent THC impacts on mPFC dopamine-mediated cognitive processes in male and female rats. *Psychopharmacology 2024 242:2*, *242*(2), 309–326. https://doi.org/10.1007/S00213-024-06676-9

Martinez, M. X., Farrell, M. R., & Mahler, S. V. (2023). Pathway-specific chemogenetic
manipulation by applying ligand to axonally expressed DREADDs. In *Vectorology for*
Optogenetics and Chemogenetics (pp. 207–220). Springer.
Matsumoto, M., & Hikosaka, O. (2009). Representation of negative motivational value in the
primate lateral habenula. *Nature Neuroscience*, *12*(1), 77–84.
https://doi.org/10.1038/NN.2233
McGaugh, J. L. (1989). Modulation of Memory Storage Processes. *Memory: Interdisciplinary*
Approaches, 33–64. https://doi.org/10.1007/978-1-4612-3500-2_3
Mitchell, A. S., & Chakraborty, S. (2013). What does the mediodorsal thalamus do? *Frontiers in*
Systems Neuroscience, *7*(JUL). https://doi.org/10.3389/FNSYS.2013.00037
Moaddab, M., Ray, M. H., & McDannald, M. A. (2021). Ventral pallidum neurons dynamically
signal relative threat. *Communications Biology 2021 4:1*, *4*(1), 1–14.
https://doi.org/10.1038/s42003-020-01554-4
Nieh, E. H., Matthews, G. A., Allsop, S. A., Presbrey, K. N., Leppla, C. A., Wichmann, R., Neve,
R., Wildes, C. P., & Tye, K. M. (2015). Decoding neural circuits that control compulsive
sucrose seeking. *Cell*, *160*(3), 528–541.
https://doi.org/10.1016/J.CELL.2015.01.003/ATTACHMENT/A7A8641E-9760-4CF4-8113-
D209EC66DF99/MMC5.PDF
Nieh, E. H., Vander Weele, C. M., Matthews, G. A., Presbrey, K. N., Wichmann, R., Leppla, C.
A., Izadmehr, E. M., & Tye, K. M. (2016). Inhibitory Input from the Lateral Hypothalamus to
the Ventral Tegmental Area Disinhibits Dopamine Neurons and Promotes Behavioral
Activation. *Neuron*, *90*(6), 1286–1298. https://doi.org/10.1016/J.NEURON.2016.04.035
Nieh, E. H., Vander Weele, C. M., Matthews, G. A., Presbrey, K. N., Wichmann, R., Leppla, C.
A., Izadmehr, E. M., & Tye, K. M. (2016). Inhibitory Input from the Lateral Hypothalamus to
the Ventral Tegmental Area Disinhibits Dopamine Neurons and Promotes Behavioral
Activation. *Neuron*, *90*(6), 1286–1298. https://doi.org/10.1016/j.neuron.2016.04.035
Ostlund, S. B., & Balleine, B. W. (2008). Differential involvement of the basolateral amygdala
and mediodorsal thalamus in instrumental action selection. *The Journal of Neuroscience :*
The Official Journal of the Society for Neuroscience, *28*(17), 4398–4405.
https://doi.org/10.1523/JNEUROSCI.5472-07.2008
Ottenheimer, D., Richard, J. M., & Janak, P. H. (2018). Ventral pallidum encodes relative reward
value earlier and more robustly than nucleus accumbens. *Nat Commun*, *9*(1), 4350.
https://doi.org/10.1038/s41467-018-06849-z
Packard, M. G., & Cahill, L. (2001). Affective modulation of multiple memory systems. *Current*
Opinion in Neurobiology, *11*(6), 752–756. https://doi.org/10.1016/S0959-4388(01)00280-X
Panagis, G., Miliaressis, E., Anagnostakis, Y., & Spyraki, C. (1995). Ventral pallidum self-
stimulation: a moveable electrode mapping study. *Behavioural Brain Research*, *68*(2), 165–
172. https://doi.org/10.1016/0166-4328(94)00169-G
Pardo-Garcia, T. R., Garcia-Keller, C., Penaloza, T., Richie, C. T., Pickel, J., Hope, B. T.,
Harvey, B. K., Kalivas, P. W., & Heinsbroek, J. A. (2019). Ventral Pallidum Is the Primary
Target for Accumbens D1 Projections Driving Cocaine Seeking. *The Journal of*
Neuroscience, *39*(11), 2041. https://doi.org/10.1523/JNEUROSCI.2822-18.2018

Parnaudeau, S., Bolkan, S. S., & Kellendonk, C. (2018). The Mediodorsal Thalamus: An
Essential Partner of the Prefrontal Cortex for Cognition. *Biological Psychiatry*, *83*(8), 648–
656. https://doi.org/10.1016/j.biopsych.2017.11.008
Paxinos, G., & Watson, C. (2006). *The rat brain in stereotaxic coordinates: hard cover edition*.
Elsevier.
Péczely, L., Ollmann, T., László, K., Kovács, A., Gálosi, R., Szabó, Á., Karádi, Z., & Lénárd, L.
(2014). Role of D1 dopamine receptors of the ventral pallidum in inhibitory avoidance
learning. *Behavioural Brain Research*, *270*, 131–136.
https://doi.org/https://doi.org/10.1016/j.bbr.2014.04.054
Pinel, J. P., & Treit, D. (1978). Burying as a defensive response in rats. *Journal of Comparative*
and Physiological Psychology, *92*(4), 708–712. https://doi.org/10.1037/H0077494
Prasad, A. A., Xie, C., Chaichim, C., Nguyen, J. H., McClusky, H. E., Killcross, S., Power, J. M.,
& McNally, G. P. (2020). Complementary Roles for Ventral Pallidum Cell Types and Their
Projections in Relapse. *Journal of Neuroscience*, *40*(4), 880–893.
https://doi.org/10.1523/JNEUROSCI.0262-19.2019
Roman-Ortiz, C., Guevara, J. A., & Clem, R. L. (2021). GABAergic basal forebrain projections to
the periaqueductal gray promote food consumption, reward and predation. *Scientific*
Reports, *11*(1), 22638. https://doi.org/10.1038/s41598-021-02157-7
Root, D. H., Barker, D. J., Estrin, D. J., Miranda-Barrientos, J. A., Liu, B., Zhang, S., Wang, H.
L., Vautier, F., Ramakrishnan, C., Kim, Y. S., Fenno, L., Deisseroth, K., & Morales, M.
(2020). Distinct Signaling by Ventral Tegmental Area Glutamate, GABA, and Combinatorial
Glutamate-GABA Neurons in Motivated Behavior. *Cell Reports*, *32*(9), 108094.
https://doi.org/10.1016/j.celrep.2020.108094
Root, D. H., Ma, S., Barker, D. J., Megehee, L., Striano, B. M., Ralston, C. M., Fabbricatore, A.
T., & West, M. O. (2013). Differential roles of ventral pallidum subregions during cocaine
self-administration behaviors. *Journal of Comparative Neurology*, *521*(3), 558–588.
https://doi.org/10.1002/CNE.23191
Root, D. H., Mejias-Aponte, C. A., Qi, J., & Morales, M. (2014). Role of Glutamatergic
Projections from Ventral Tegmental Area to Lateral Habenula in Aversive Conditioning. *The*
Journal of Neuroscience, *34*(42), 13906. https://doi.org/10.1523/JNEUROSCI.2029-
14.2014
Root, D. H., Melendez, R. I., Zaborszky, L., & Napier, T. C. (2015). The ventral pallidum:
Subregion-specific functional anatomy and roles in motivated behaviors. *Progress in*
Neurobiology, *130*, 29–70. https://doi.org/10.1016/J.PNEUROBIO.2015.03.005
Russell, E. L., & McDannald, M. A. (2024). Ventral Pallidum Neurons Are Necessary to
Generalize and Express Fear-Related Responding in a Minimal Threat Setting. *ENeuro*,
11(11). https://doi.org/10.1523/ENEURO.0124-24.2024
Sangha, S., Diehl, M. M., Bergstrom, H. C., & Drew, M. R. (2020). Know safety, no fear.
Neuroscience & Biobehavioral Reviews, *108*, 218–230.
https://doi.org/10.1016/J.NEUBIOREV.2019.11.006
Scott, A., Palmer, D., Newell, B., Lin, I., Cayton, C. A., Paulson, A., Remde, P., & Richard, J. M.
(2023). Ventral Pallidal GABAergic Neuron Calcium Activity Encodes Cue-Driven Reward
Seeking and Persists in the Absence of Reward Delivery. *Journal of Neuroscience*, *43*(28),
51915203. https://doi.org/10.1523/JNEUROSCI.0013-23.2023

Scott, A., Paulson, A., Prill, C., Kermoade, K., Newell, B., Eckenwiler, E. A., Lemos, J. C., & Richard, J. M. (2025). Ventral Pallidal GABAergic Neurons Drive Consumption in Male, But Not Female, Rats. *ENeuro*, *12*(2). https://doi.org/10.1523/ENEURO.0245-24.2025,

Smith, K. S., & Berridge, K. C. (2005). The ventral pallidum and hedonic reward: neurochemical maps of sucrose "liking" and food intake. *Journal of Neuroscience*, *25*(38), 8637–8649.

Smith, K. S., & Berridge, K. C. (2007). Opioid Limbic Circuit for Reward: Interaction between Hedonic Hotspots of Nucleus Accumbens and Ventral Pallidum. *Journal of Neuroscience*, *27*(7), 1594–1605. https://doi.org/10.1523/jneurosci.4205-06.2007

Smith, K. S., Tindell, A. J., Aldridge, J. W., & Berridge, K. C. (2009). Ventral Pallidum Roles in Reward and Motivation. *Behavioural Brain Research*, *196*(2), 155. https://doi.org/10.1016/J.BBR.2008.09.038

Soares-Cunha, C., & Heinsbroek, J. A. (2023). Ventral pallidal regulation of motivated behaviors and reinforcement. *Frontiers in Neural Circuits*, *17*. https://doi.org/10.3389/FNCIR.2023.1086053/FULL

Stamatakis, A. M., Jennings, J. H., Ung, R. L., Blair, G. A., Weinberg, R. J., Neve, R. L., Boyce, F., Mattis, J., Ramakrishnan, C., & Deisseroth, K. (2013). A unique population of ventral tegmental area neurons inhibits the lateral habenula to promote reward. *Neuron*, *80*(4), 1039–1053.

Stephenson-Jones, M. (2019). Pallidal circuits for aversive motivation and learning. *Current Opinion in Behavioral Sciences*, *26*, 82–89. https://doi.org/https://doi.org/10.1016/j.cobeha.2018.09.015

Stephenson-Jones, M., Bravo-Rivera, C., Ahrens, S., Furlan, A., Xiao, X., Fernandes-Henriques, C., & Li, B. (2020). Opposing contributions of GABAergic and glutamatergic ventral pallidal neurons to motivational behaviors. *Neuron*, *105*(5), 921–933. https://doi.org/10.1016/j.neuron.2019.12.006

Stout, K. A., Dunn, A. R., Lohr, K. M., Alter, S. P., Cliburn, R. A., Guillot, T. S., & Miller, G. W. (2016). Selective Enhancement of Dopamine Release in the Ventral Pallidum of Methamphetamine-Sensitized Mice. *ACS Chemical Neuroscience*, *7*(10), 1364–1373. https://doi.org/10.1021/ACSCHEMNEURO.6B00131

Tooley, J., Marconi, L., Alipio, J. B., Matikainen-Ankney, B., Georgiou, P., Kravitz, A. V., & Creed, M. C. (2018). Glutamatergic Ventral Pallidal Neurons Modulate Activity of the Habenula–Tegmental Circuitry and Constrain Reward Seeking. *Biological Psychiatry*, *83*(12), 1012–1023. https://doi.org/10.1016/J.BIOPSYCH.2018.01.003

Trainor, B. C. (2011). Stress responses and the mesolimbic dopamine system: Social contexts and sex differences. *Hormones and Behavior*, *60*(5), 457–469. https://doi.org/10.1016/J.YHBEH.2011.08.013

Tripathi, A., Prensa, L., & Mengual, E. (2013). Axonal branching patterns of ventral pallidal neurons in the rat. *Brain Structure and Function*, *218*, 1133–1157.

Wang, F., Zhang, J., Yuan, Y., Chen, M., Gao, Z., Zhan, S., Fan, C., Sun, W., & Hu, J. (2020). Salience processing by glutamatergic neurons in the ventral pallidum. *Science Bulletin*, *65*(5), 389–401. https://doi.org/10.1016/J.SCIB.2019.11.029

Waraczynski, M. (2016). Toward a systems-oriented approach to the role of the extended amygdala in adaptive responding. *Neuroscience & Biobehavioral Reviews*, *68*, 177–194. https://doi.org/https://doi.org/10.1016/j.neubiorev.2016.05.015

Wolff, M., & Halassa, M. M. (2024). The mediodorsal thalamus in executive control. *Neuron*, *112*(6), 893–908. https://doi.org/10.1016/j.neuron.2024.01.002

Wulff, A. B., Tooley, J., Marconi, L. J., & Creed, M. C. (2019). Ventral pallidal modulation of aversion processing. *Brain Research*, *1713*, 62–69. https://doi.org/10.1016/J.BRAINRES.2018.10.010

Yang, L., Fang, L. Z., Lynch, M. R., Xu, C. S., Hahm, H. J., Zhang, Y., Heitmeier, M. R., Costa, V. D., Samineni, V. K., & Creed, M. C. (2024). Transcriptomic landscape of mammalian ventral pallidum at single-cell resolution. *Science Advances*, *10*(50), 6017. https://doi.org/10.1126/SCIADV.ADQ6017

Yang, Y., Cui, Y., Sang, K., Dong, Y., Ni, Z., Ma, S., & Hu, H. (2018). Ketamine blocks bursting in the lateral habenula to rapidly relieve depression. *Nature 2018 554:7692*, *554*(7692), 317–322. https://doi.org/10.1038/nature25509

Young, L. J., & Pfaff, D. W. (2014). Sex differences in neurological and psychiatric disorders. *Frontiers in Neuroendocrinology*, *35*(3), 253–254. https://doi.org/10.1016/j.yfrne.2014.05.005

Zachry, J. E., Nolan, S. O., Brady, L. J., Kelly, S. J., Siciliano, C. A., & Calipari, E. S. (2021). Sex differences in dopamine release regulation in the striatum. *Neuropsychopharmacology*, *46*(3), 491–499. https://doi.org/10.1038/S41386-020-00915-1;SUBJMETA=1788,1964,378,548,631;KWRD=NEUROTRANSMITTERS,REWARD

Zahm, D. S. (1989). The ventral striatopallidal parts of the basal ganglia in the rat—II. Compartmentation of ventral pallidal efferents. *Neuroscience, 30*(1), 33–50. https://doi.org/10.1016/0306-4522(89)90351-5

Zahm, D. S., & Heimer, L. (1988). Ventral striatopallidal parts of the basal ganglia in the rat: I. Neurochemical compartmentation as reflected by the distributions of neurotensin and substance P immunoreactivity. *Journal of Comparative Neurology*, *272*(4), 516–535. https://doi.org/https://doi.org/10.1002/cne.902720406

Zahm, D. S., & Heimer, L. (1990). Two transpallidal pathways originating in the rat nucleus accumbens. *Journal of Comparative Neurology*, *302*(3), 437–446.

Zahm, D. S., & Heimer, L. (1993). Specificity in the efferent projections of the nucleus accumbens in the rat: Comparison of the rostral pole projection patterns with those of the core and shell. *Journal of Comparative Neurology*, *327*(2), 220–232. https://doi.org/10.1002/cne.903270205

Zahm, D. S., Williams, E., & Wohltmann, C. (1996). Ventral striatopallidothalamic projection: IV. Relative involvements of neurochemically distinct subterritories in the ventral pallidum and adjacent parts of the rostroventral forebrain. *Journal of Comparative Neurology*, *364*(2), 340–362.

Zahm, D. S., Zaborszky, L., Alheid, G. F., & Heimer, L. (1987). The ventral striatopallidothalamic projection: II. The ventral pallidothalamic link. *Journal of Comparative Neurology*, *255*(4), 592–605. https://doi.org/https://doi.org/10.1002/cne.902550410

Zhang, G. W., Shen, L., Tao, C., Jung, A. H., Peng, B., Li, Z., Zhang, L. I., & Tao, H. W. (2021). Medial preoptic area antagonistically mediates stress-induced anxiety and parental behavior. *Nature Neuroscience 2021 24:4*, *24*(4), 516–528. https://doi.org/10.1038/s41593-020-00784-3

1196