Nanoscopic tau aggregates in Parkinson's disease

Authors

Florence Layburn[1,2], Dorothea Boeken[1,2], Yu P. Zhang[1,2], Kristy Halliday[4], Dan Rodgers[4],
Lakmini Kahanawita[3], Bina Patel[3], Annelies Quaegebeur[3,4], Caroline H. Williams-Gray[3], David
Klenerman[1,2]

Author Affiliations

[1]Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW,
U.K.

[2]UK Dementia Research Institute at Cambridge, Cambridge CB2 0XY, U.K.

[3]Department of Clinical Neurosciences, University of Cambridge, Cambridge CB2 0PY, U.K.

[4]Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK.

Abstract:

Post-mortem tau pathology is frequently observed in Parkinson's disease (PD) using
immunohistochemistry (IHC) to measure large inclusions, however, small protein aggregates
that precede inclusions are considered a major driver of toxicity in neurodegenerative disease.
We aimed to uncover the distribution of nanoscopic aggregates across six brain regions in
post-mortem tissue from 14 PD and 15 controls using the single-molecule pull-down assay
(SiMPull). In the hippocampus and amygdala, tau IHC and SiMPull were associated with
advanced age in controls and dementia status in PD. Despite negligible tau IHC-labelled
aggregates in the putamen, we identified a unique population of high-intensity nanoscopic tau
aggregates for a subset of PD cases using SiMPull, ranging from 10–1,000 epitopes per
aggregate and 30–1,000 nm in length. Previous evidence linking nigrostriatal tau pathology
and motor deficits indicates that the nanoscopic tau aggregates identified in this study may
contribute to striatal dysfunction in PD.

Main:

Parkinson's disease (PD) manifests as a movement disorder during life, however, post-
mortem neuropathology is defined by two key features. First is the loss of dopaminergic
neurons from the substantia nigra in the midbrain, a component of the basal ganglia
movement-control circuitry (1, 2). Cell death in the substantia nigra directly translates to the
cardinal motor symptoms of PD, such as slowness of movement and rigidity (3). The second
criterion for PD diagnosis is the presence of Lewy Bodies; large, insoluble protein aggregates
primarily composed of alpha-synuclein (αSyn) and distributed throughout the brain according
to the Lewy Body pathology Braak stages (4-6). Post-mortem neuropathology has informed
the development of biomarkers to help support the diagnosis of PD during life, for example
DaTScans to assess the loss of dopaminergic nerve terminals in the striatum, and the
explosion of αSyn-based fluid biomarkers currently in preclinical development (7-11). Our
understanding of PD neuropathology is overwhelmingly αSyn-centric, however, this is not the
only protein to aggregate in the PD brain.

For many years, post-mortem tau pathology has also been observed in PD. Tau is a
microtubule-associated protein that becomes hyperphosphorylated and forms inclusions such
as neurofibrillary tangles in the brains of people with Alzheimer's disease and other
tauopathies (12, 13). There are surprisingly few studies on the significance of tau in PD

considering that approximately 50% of PD-dementia cases display post-mortem tau pathology
(14); there is an association between genetic variation in MAPT, the gene encoding tau, and
increased risk of parkinsonism (15); and there are several parkinsonian disorders affecting the
nigrostriatal pathway, such as progressive supranuclear palsy, where the primary pathology is
tau (16, 17). Furthermore, a recently published study identified an association between
nigrostriatal tau pathology and motor deficits in a cohort of PD and non-PD individuals,
suggesting that tau may contribute to dopaminergic degeneration (18). Despite the overlap of
pathology in neurodegenerative diseases, we still do not know if tau aggregation is merely co-
incident and independent of PD disease progression, or co-pathological and actively
contributing to cell death in PD.

Currently, neuropathological analysis depends on detection of large insoluble inclusions which
are visible by immunohistochemistry (IHC). However, Lewy Bodies are poorly associated with
neurological dysfunction and mounting evidence suggests that small, intermediate protein
aggregates— which are undetectable using IHC due to their nanoscopic size range— are the
true drivers of disease for a range of neurological disorders (19-21). For example, small
aggregates are more potent at inducing neurotoxicity, and more efficient at spreading between
cells to seed further aggregation when compared to fibrils and inclusion bodies (22, 23). To
overcome the barriers to studying nanoscopic aggregates, our group uses the single-molecule
pull-down assay (SiMPull) and fluorescence imaging to characterise protein aggregates from
human samples in terms of size, concentration, and composition without the need for
excessive purification or denaturing conditions that would disrupt endogenous aggregate
structures. We have used SiMPull to measure protein aggregates from blood, cerebrospinal
fluid, and brain samples; for example, identifying diffusible αSyn aggregates in PD brain tissue
capable of inducing inflammation and membrane permeabilisation (20, 24, 25).

In this study, we aimed to improve our understanding of tau aggregate distributions for the full
range of aggregate sizes in a cohort of 14 PD and 15 controls (CON). To capture the breadth
of pathology in the PD brain, we studied six brain regions; from earliest to latest affected
according to the Lewy Body Braak stages, including the substantia nigra (SN), amygdala
(AMYG), hippocampus (HIP), putamen (PUT), frontal cortex (FC) and occipital cortex (OC).
We measured tau aggregates extracted from fractionated post-mortem brain tissue—
henceforth referred to as nanoscopic aggregates— using SiMPull, and measured tau
inclusions in formalin-fixed, paraffin-embedded tissue sections using IHC. We identified a
disease-specific trend of higher intensity nanoscopic tau aggregates in the basal ganglia using
SiMPull, implicating tau in striatal dysfunction in PD.

Results:

Single-molecule imaging can be used to measure protein aggregates' abundance, intensity, and size

We used immunohistochemistry (IHC) and single-molecule fluorescence imaging to quantify protein aggregates from post-mortem brain tissue from 14 PD and 15 controls across six brain regions (Figure 1a). The groups were similar in age (PD = 80.9 ± 6.5 years; control = 74.7 ± 12.7 years; two-sided T-test p-value = 0.13) and post-mortem interval (PD = 53.2 ± 32.1 hours; control = 53.8 ± 27.1 hours; two-sided T-test p-value = 0.96), and there were slightly more females in the control group (6F/9M) compared to PD (3F/11M) (Methods Table 1). Formalin-fixed, paraffin-embedded (FFPE) brain tissue sections were stained using 3, 3'-diaminobenzidine (DAB) IHC to quantify large insoluble aggregates, and fractionated fresh-frozen brain tissue was used for single-molecule pull-down (SiMPull) to measure nanoscopic aggregates (Figure 1b–c). SiMPull is an aggregate-specific assay, due to the use of identical capture and detection antibodies; if a monomer is captured on the surface, it does not have a second epitope to accommodate the detection antibody and therefore does not generate a fluorescent signal (Figure 1c).

The output of SiMPull depends on the imaging probe; a dye-conjugated detection antibody was used for diffraction-limited imaging, and a DNA-conjugated antibody together with a dye-labelled imaging DNA strand was used for DNA points accumulation for imaging in nanoscale topography (DNA-PAINT) super-resolution imaging. We used diffraction-limited imaging to quantify the density of aggregates per field of view as a measure of abundance (Figure 1d) and aggregate intensity as a measure of size (Figure 1e). To demonstrate the utility of diffraction-limited SiMPull, we used silica nanoparticles (SiNaPs) as aggregate standards, either conjugated with the tau antibody HT7 binding sequence or αSyn monomers. There was a strong linear relationship between SiNaP concentration and SiMPull density for a dilution series of tau and αSyn SiNaPs respectively (Figure 1d). The tau SiNaP intensity distribution was higher than the HT7 antibody binding sequence dipeptide, which can only bind a single detection antibody, demonstrating that larger aggregates with higher numbers of detection antibody binding lead to higher intensity values (Figure 1e).

We used DNA-PAINT to expand on the size information from diffraction-limited SiMPull and measure aggregate lengths on the nanometre scale. DNA-PAINT overcomes the diffraction-limit of light by separating the fluorescent signal over time across thousands of frames, which were subsequently reconstructed to generate the super-resolved image (Figure 1f).

Figure 1. SiMPull workflow and output for characterising protein aggregates. (a) Post-mortem brain tissue cohort overview. The SN was highlighted due to the reduced sample size for frozen tissue. **(b)** IHC was used to study large, insoluble inclusion bodies. **(c)** Schematic representation of SiMPull used to detect nanoscopic aggregates. SiMPull excludes monomers from detection by using identical capture and detection antibodies. **(d)** Examples of diffraction-limited imaging using tau SiNaP aggregate standards. Aggregate density was significantly correlated with aggregate concentration for tau and αSyn SiNaP aggregates (Pearson correlation). Each point represents the mean of 12 FOVs and three well replicates, error bars represent mean ± standard deviation. **(e)** "Auto" contrast-adjusted example images of HT7 dipeptide, tau SiNaPs, and corresponding aggregate intensity distributions. **(f)** Example DNA-PAINT images from one FOV are shown; first the individual image frames, then the super-resolved image. Insets **(i)** and **(ii)** feature close-ups of super-resolved phosphorylated tau aggregates from a PD human brain sample. CON = Control; HIP = Hippocampus; AMYG = Amygdala; PUT = Putamen; SN = Substantia nigra; FC = Frontal cortex; OC = Occipital cortex. SiNaP = silica nanoparticle; FOV = Field of view.

PD cases display typical neuropathology in FFPE sections, but αSyn SiMPull does not detect PD-specific nanoscopic aggregates

All PD cases in this study met the international criteria for post-mortem diagnosis of PD, displaying neurodegeneration in the substantia nigra (Figure 2a–b) and Lewy Body pathology (Figure 2c–e). As expected, αSyn immunolabel coverage was significantly higher in PD compared to controls in all brain regions except the occipital cortex, which had low staining in both groups (Figure 2c). Typical Lewy Body staining and elongated, neuritic αSyn staining was observed by IHC across the PD samples (Figure 2d–e).

We used the αSyn C-terminus binding antibody Syn211 that detects most forms of αSyn for SiMPull to measure nanoscopic αSyn aggregates and compare with Lewy Body deposition (26). SiMPull aggregate densities were within the lower range of detection as compared to αSyn SiNaPs in Figure 1b, mostly below 2 spots/µm^2 for all brain regions, and there was a significant increase in the PD occipital cortex compared to controls (Figure 2f). Aggregate intensities were also low, although there was a significant increase in mean aggregate intensity in the control frontal cortex compared to PD (Figure 2g). In light of the low signal from SiMPull signal, we confirmed that Syn211 binds αSyn in the fractionated brain samples (Supplementary Figure 1a), and obtained similar SiMPull results with five alternative αSyn antibodies (Supplementary Figure 1b–d).

We probed the intensity values further by calculating the approximate number of epitopes per aggregate. To do so, we used another total αSyn antibody SOY1 to capture αSyn monomers on the surface and detected using Syn211 (Methods Figure 1). This analysis revealed that 88.8% of aggregates for controls and 87.8% of aggregates for PD were composed of fewer than five epitopes, and there were no clear differences between the size distributions of PD and controls (Figure 2i). Due to the low abundance and small size of objects detected by αSyn SiMPull, we did not consider these species disease-specific, and continued to investigate tau protein aggregates in the PD brain.

Figure 2. PD cases displayed typical neuropathology in FFPE brain sections, but αSyn SiMPull does not detect PD-specific nanoscopic aggregates. (a–b) Representative images of neuronal loss in the SN in PD compared to CON. **(c)** αSyn IHC coverage was significantly higher in PD (n =14) compared to CON (n = 15) for all brain regions except the OC, which had low staining in both groups (Two-sided Mann-Whitney test, SN $p < 0.0001$; AMYG $p < 0.0001$; PUT $p < 0.0001$; HIP $p < 0.0001$; FC $p = 0.0023$; OC $p = 0.1583$). Each point represents an individual case. **(d–e)** Representative images of αSyn staining in PD brain tissue. **(e)** Depicts different staining morphologies within close proximity in a PUT section, with red arrows highlighting Lewy Bodies as round inclusion-like staining and black arrows highlighting Lewy Neurites as elongated, neuritic staining. **(f)** αSyn SiMPull mean aggregate densities were significantly higher in the PD OC compared to CON (Two-sided Mann-Whitney test, SN $p = 0.9048$; AMYG $p = 0.1658$; PUT $p = 0.1583$; HIP $p = 0.2115$; FC $p = 0.4253$; OC $p = 0.0079$). **(g)** αSyn SiMPull mean aggregate intensities were significantly higher in the CON FC compared to PD (Two-sided Mann-Whitney test, SN $p = 0.4127$; AMYG $p = 0.4776$; PUT

p = 0.7148; HIP *p* = 0.3164; FC *p* = 0.01375; OC *p* = 0.9145). **(f–g)** Each point represents the
mean of 9–12 FOVs and 2–3 independent repeats per case. **(h)** Example αSyn SiMPull
images from one PD and one CON case in the AMYG, PUT, and FC. All images were contrast-
adjusted to the same range. **(i)** Distribution of epitopes per aggregate for CON and PD
compiled for all brain regions, plotted on a log scale. CON = Control; SN = Substantia nigra;
AMYG = Amygdala; HIP = Hippocampus; PUT = Putamen; FC = Frontal cortex; OC = Occipital
cortex; FOV = Field of view.

Tau SiMPull reveals two sub-populations of nanoscopic aggregates, and higher intensity aggregates in the PD putamen

To investigate the distribution of tau pathology in PD, we first used IHC to quantify tau
immunolabel coverage (Figure 3a). There was a wide range of tau staining levels in the
hippocampus and amygdala across the whole tissue cohort, although no significant
differences between PD and controls (Figure 3a). For future analysis, control cases were
subclassified into Path-Positive and Path-Negative based on tau staining levels in the
hippocampus (Figure 3a). There was a significant increase in tau staining in the PD substantia
nigra, although absolute coverage levels were low compared to the hippocampus and
amygdala, which had visibly higher staining coverage compared to the other regions (Figure
3b).

We followed up IHC analysis with tau SiMPull using the total-tau antibody HT7 and observed
similar trends in the hippocampus and amygdala, with high intensity aggregates for both PD
and controls in these regions (Figure 3c–d). Interestingly, there was a significant increase in
aggregate intensities in PD compared to controls in the putamen that was not previously
observed using IHC (Figure 3d). There were no significant differences in aggregate density
between controls and PD in any of the brain regions (Supplementary Figure 2a).

We used the HT7 dipeptide to calculate the approximate number of epitopes per aggregate
and observed two distributions in both PD and controls, reflecting two aggregates populations
of 1–10 and 10–1000 epitopes respectively (Figure 3e). This prompted us to inspect the
underlying intensity distributions for PD, Path-Positive controls and Path-Negative controls per
brain region (Figure 3f). The intensity distributions revealed there were both low and high-
intensity aggregates in PD and Path-Positive controls in the hippocampus and amygdala;
there were predominantly low-intensity aggregates in the frontal and occipital cortices for all
groups; and there were predominantly low-intensity aggregates in Path-Negative controls in
all brain regions. In the putamen, the high-intensity aggregate population was only present in
PD.

Figure 3. Tau SiMPull intensities are higher in the PD putamen compared to CON. (a) Tau IHC coverage in PD (n = 14) compared to CON (n = 15) shows a significant increase in PD in the SN (Two-sided Mann-Whitney test, HIP p = 0.4773; AMYG p = 0.2116; SN p = 0.00044; PUT p = 0.1456; FC p = 0.5045; OC p = 0.3314). Each point represents an individual case. **(b)** Representative images of tau staining. The red arrow highlights tau staining in the SN as opposed to dark pigmented cells also visible. **(c)** Example tau SiMPull images from one PD and one CON case across the HIP, PUT, and OC. Each pair of images was contrast-adjusted to the same range. **(d)** Tau SiMPull mean aggregate intensity, plotted on a log scale.

Aggregate intensities were significantly higher in the PD PUT compared to CON (Two-sided Mann-Whitney test, HIP $p = 0.6075$; AMYG $p = 0.8428$; SN $p = 0.4127$; PUT $p = 0.0013$; FC $p = 0.9358$; OC $p = 0.3761$). Each point represents the mean of 9–12 FOVs and 2–3 independent repeats per case. **(e)** Distribution of epitopes per aggregate for CON and PD respectively compiled for all brain regions, plotted on a log scale, revealing two populations of aggregate sizes. **(f)** Tau aggregate intensity distributions for Path-Negative CON, Path-Positive CON, and PD across all brain regions. The PUT is highlighted to demonstrate the higher intensity distribution in PD. For each region, "n" refers to the number of cases, and "n aggregates" refers to the number of aggregates detected by SiMPull per group. SN sample numbers are italicised to highlight the lower n compared to other brain regions. CON = Control; HIP = Hippocampus; AMYG = Amygdala; PUT = Putamen; SN = Substantia nigra; FC = Frontal cortex; OC = Occipital cortex; FOV = Field of view.

Phosphorylated tau aggregates are higher intensity in PD putamen compared to CON

We repeated SiMPull using the phosphorylated tau (pTau) antibody AT8, which is commonly used for neuropathological diagnosis of tauopathies. Again, high-intensity aggregates were observed in the hippocampus and amygdala for both PD and controls (Figure 4a–b). Furthermore, mean aggregate intensities were correlated with tau SiMPull and pTau SiMPull (Supplementary Figure 2b). Mean aggregate intensities were significantly higher in PD in the putamen and substantia nigra, although the latter had a reduced sample size and did not reach statistical significance, and there was a modest increase in intensity in the PD occipital cortex (Figure 4b). Although aggregate levels were lower in the occipital cortex compared to other brain regions, we also observed a significant increase in aggregate density in PD in this region (Figure 4c). Unlike tau SiMPull, pTau aggregate intensities displayed a single distribution for each group (Figure 4d). There were overlapping distributions for PD and Path-Positive controls in the hippocampus and amygdala, and a distinct shift towards higher pTau aggregate intensities for PD in the putamen and substantia nigra (Figure 4d). In the occipital cortex, the difference in intensities observed in Figure 4c was explained by the predominantly low-intensity aggregates in Path-Negative controls, however pTau aggregate intensities were similar between Path-Positive controls and PD (Figure 4d).

We identified pTau aggregates in the putamen ranging from 30–1,000 nm using super-resolution imaging (Figure 4e–f). There was a significant increase in aggregate length in PD, driven by a subset of PD cases with longer aggregates (Figure 4g). Examples of long pTau aggregates in the six highest PD cases are demonstrated in Figure 4h, accompanied by representative images of smaller aggregates that were present in both PD and controls.

Figure 4. pTau aggregates are higher intensity in PD PUT compared to CON, corresponding to lengths of 30–1,000 nm. (a) Example pTau SiMPull images from one PD and one CON case across the HIP, PUT, and OC. All images were contrast-adjusted to the same range. **(b)** pTau SiMPull mean aggregate intensity for PD (n = 14) and CON (n = 15), plotted on a log scale. Aggregate intensities were significantly higher in the PD PUT and OC (Two-sided Mann-Whitney test, HIP *p* = 0.2115; AMYG *p* = 0.5899; SN *p* = 0.1111; PUT *p* =

0.00022; FC *p* = 0.2703; OC *p* = 0.04817). **(c)** pTau SiMPull mean aggregate density, plotted
on a log scale. Aggregate densities were significantly higher in the PD OC compared to CON
(Two-sided Mann-Whitney test, HIP *p* = 0.9759; AMYG *p* = 1; SN *p* = 0.1905; PUT *p* = 0.1121;
FC *p* = 0.9486; OC *p* = 0.01448). **(b–c)** Each point represents the mean of 9–12 FOVs and
2–3 independent repeats per case. **(d)** Tau aggregate intensity distributions for Path-Negative
CON, Path-Positive CON, and PD across all brain regions. The PUT is highlighted to
demonstrate the significant shift towards higher intensity aggregates in PD. For each region,
"n" refers to the number of cases, and "n aggregates" refers to the number of aggregates
detected by SiMPull per group. SN sample numbers are italicised to highlight the lower n
compared to other brain regions. **(e)** Example super-resolution image of PD13 from a subset
of the FOV. **(f)** Length distributions of super-resolved pTau aggregates in the PUT for PD and
CON, ranging from 30–1,000 nm. **(g)** Aggregate lengths were significantly higher in PD
compared to CON (Two-sided Mann-Whitney test, *p* = 0.0099). Each point represents the
mean of 4 FOVs per case. **(h)** Example images of long aggregates from the highest six PD
cases and representative images of short aggregates from CON cases. A subset of the FOV
was used for demonstration purposes and the fire colour preset was used to represent the
density of localisations. CON = Control; FOV = Field of view; HIP = Hippocampus; AMYG =
Amygdala; PUT = Putamen; SN = Substantia nigra; FC = Frontal cortex; OC = Occipital cortex;

272 SiMPull tau aggregate intensities are correlated with IHC tau coverage the
273 hippocampus and amygdala, but not PD putamen

Next, we investigated the relationship between tau IHC coverage and tau SiMPull aggregate
intensities. We started by classifying the nanoscopic aggregates from SiMPull as "dim" or
"bright" and calculated the percentage of bright aggregates per case, revealing that a
combination the two aggregate populations could be detected within an individual (Figure 5a–
b). The brightness trends were consistent with tau SiMPull mean intensities and there was a
significant increase in PD putamen compared to controls (Figure 5b).

There was a significant correlation between tau SiMPull and tau IHC in the hippocampus for
both controls and PD (Figure 5c). There was also a significant correlation between tau SiMPull
and tau IHC in the amygdala for controls. However, there was no significant correlation
between tau SiMPull and tau IHC for PD in the amygdala, or for PD and controls in the
putamen. The latter can be explained by the low signal for controls in the putamen, however,
this analysis highlighted the presence of bright nanoscopic tau aggregates in PD that were
undetectable by IHC.

In PD, there was a positive correlation between tau SiMPull and αSyn IHC coverage in the
hippocampus, and a negative correlation in the amygdala, although neither relationship
reached statistical significance (Figure 5d). Due to low signal, there was no significant
correlation between SiMPull and IHC in substantia nigra, frontal cortex, and occipital cortex
brain regions (Supplementary Figure 3).

Figure 5. Tau SiMPull is correlated with tau IHC in CON, but not in PD PUT. (a) Pooled tau SiMPull intensity distribution for all regions in PD (n = 14) and CON (n = 15). Aggregates were classified as "dim" or bright", and the percentage of bright aggregates for each case was calculated. A representative image of dim and bright aggregates from tau SiMPull is shown. **(b)** There was a significant increase in bright tau aggregates in PD in the PUT (Two-sided Mann-Whitney test, HIP *p* = 0.6075; AMYG *p* = 0.5899; SN *p* = 0.5556; PUT *p* = 0.0023; FC *p* = 0.3173; OC *p* = 0.4544). Each point represents the mean of 9–12 FOVs and 2–3 independent repeats per case. **(c)** Tau SiMPull bright aggregates were positively correlated with IHC tau coverage in CON HIP and AMYG, but not PUT (Spearman correlation). Tau SiMPull bright aggregates were positively correlated with IHC tau coverage in PD HIP, but not AMYG or PUT (Spearman correlation). **(d)** Tau SiMPull bright aggregates and αSyn IHC coverage were correlated in PD HIP and AMYG, although these did not reach significance (Spearman correlation). **(c–d)** Each point represents the mean per case. CON = Control; HIP = Hippocampus; AMYG = Amygdala; PUT = Putamen; SN = Substantia nigra; FC = Frontal cortex; OC = Occipital cortex; FOV = Field of view.

Tau pathology is associated with age in controls and dementia status in PD

We used principal component analysis (PCA) and k-means cluster analysis to summarise the different pathology trends within this cohort (Figure 6a). The PCA input variables included αSyn IHC coverage (%), tau IHC coverage (%), tau SiMPull bright aggregates (%) and pTau SiMPull mean intensity (photons) from the hippocampus, amygdala, and putamen. αSyn IHC coverage and tau IHC from the substantia nigra were also included, however SiMPull results

from the substantia nigra were excluded due to low sample size. Results from the frontal and
occipital cortices were excluded due to low signal across all markers. We identified four
clusters within the cohort, comprised of two control and two PD sub-groups (Figure 6a). We
plotted the pathology markers for each PCA group to reveal: Group 1 controls had low
pathology in all brain regions; Group 2 controls had high tau pathology predominantly in the
medial temporal lobe (including the hippocampus and amygdala); Group 3 PD cases had αSyn
pathology and moderate tau pathology in the medial temporal lobe; Group 4 PD cases had
αSyn pathology, as well as high tau pathology in the medial temporal lobe and putamen
(Figure 6b). The pathology trends for the individual cases in each group are represented in
Supplementary Figure 4.

We explored whether the different PCA groups were associated with clinical phenotypes such
as age, dementia status, or MAPT genotype. Group 1 controls without pathology were younger
compared to the other groups (Figure 6c) and tau pathology was associated with age in
controls (Supplementary Table 1, Supplementary Figure 5). Most of the PD cases with
dementia were in Group 4 (Figure 6d), and there was no clear association between MAPT
genotype and PCA group (Figure 6e).

Figure 6. Tau pathology is associated with age in CON and dementia status in PD. (a) Principal component analysis (PCA) shows the separation of the cohort into four clusters along PC1 (32.0% variance) and PC2 (18.7% variance). Ellipses indicate confidence interval = 0.95 of groups. MTL = Medial temporal lobe, referring to the HIP and AMYG. **(b)** Summary of pathology for each PCA group, each point represents the group mean and the shaded area represents the standard deviation. **(c)** Age of death was significantly lower in Group 1 compared to the other groups (Dunn's multiple comparison test, Group 1 vs Group 2 p = 0.0099; Group 1 vs Group 3 p = 0.0120; Group 1 vs Group 4 p = 0.0393). **(d)** Group 4 had the highest proportion of cases with a dementia diagnosis. **(e)** There was an even distribution of H1/H1 and H1/H2 MAPT genotype for Groups 1, 2 and 3. The majority of cases in Group 4 had an H1/H2 MAPT genotype. CON = Control; HIP = Hippocampus; AMYG = Amygdala; PUT = Putamen; SN = Substantia nigra; FC = Frontal cortex; OC = Occipital cortex.

Discussion:

In this study, we used single-molecule imaging to compare the distribution of nanoscopic tau aggregates with IHC-labelled tau inclusions in PD post-mortem brain. We identified a unique population of high-intensity nanoscopic tau aggregates in the PD putamen that were not detectable using IHC, ranging from 10–1,000 epitopes per aggregate and up to 1,000 nm in length. Our results align with recently-published results of nigrostriatal tau deposition associated with motor symptom development, as well as implicating soluble tau aggregates in PD-dementia (18).

Although we observed consistently higher αSyn IHC in PD, αSyn SiMPull did not identify a PD-specific nanoscopic αSyn aggregate population. Instead, we observed low numbers of low-intensity αSyn aggregates, predominantly composed of fewer than five epitopes. The lack of association between αSyn SiMPull and IHC forced us to question whether these were true αSyn aggregates, functional low-order assemblies such as tetramers, or alternatively, αSyn monomers sandwiched by other protein binding partners (27, 28). If the latter is true, this may explain the increased αSyn SiMPull intensities in the frontal cortex and occipital cortex compared to other brain regions, as the cortical regions were largely free of Lewy Bodies and may contain more functional αSyn engaged in protein-protein interactions. At this stage we were unable to determine which assemblies were true nanoscopic αSyn aggregates, and future studies should consider screening αSyn binding partners as exclusion criteria for aggregate analysis.

Unlike αSyn, we identified a robust nanoscopic tau aggregate population that aligned with IHC staining in the hippocampus and amygdala, which are highly and early affected in Alzherimer's disease pathology (13). Surprisingly, we observed a striking increase in SiMPull intensity in the PD putamen in the absence of tau IHC staining. From SiMPull, the number of epitopes per aggregate had a bimodal distribution, with one population of aggregates between 2–10 epitopes and a second between 10–1,000 epitopes. The first tau aggregate population had a similar number of epitopes as αSyn SiMPull, and could similarly be interpreted as true tau aggregates, or tau complexes with binding partners. Further analysis of the underlying intensity distributions for each brain region revealed that Path-Negative controls only had low-intensity (or 2–10 epitope) aggregates, reinforcing that protein assemblies in this size range were not associated with IHC, whilst high-intensity aggregates were observed in the hippocampus and amygdala for Path-Positive controls and PD. In the putamen, high-intensity aggregates were uniquely present in PD, and the substantia nigra displayed a similar trend towards higher intensities in a reduced sample size. The putamen is perturbed in PD due to its functional and anatomical proximity to the substantia nigra, although the trigger for cell death in this circuit is not known (29, 30). The recent study by Chu et al., (2024) proposed that dopaminergic neurodegeneration may be tau-mediated, after observing IHC-positive nigrostriatal tau pathology as a common feature of PD and non-PD individuals with motor deficits. The average age of non-PD motor-deficit individuals in this study was 10 years older than our PD cohort, implying that the nanoscopic tau aggregates observed by SiMPull may eventually form large tau inclusions. Furthermore, our results support the conclusion that neuronal cell death in PD may be tau-mediated, but raise the question of which aggregate species are more neurotoxic.

Our findings in the putamen were strengthened by the results of phosphorylated tau (pTau) SiMPull, which revealed a consistent increase in pTau aggregate intensity in PD. This time, there was a single aggregate intensity distribution for each group and lower numbers of aggregates in Path-Negative controls compared to the other groups, indicating that the low-intensity aggregate population from tau SiMPull was not phosphorylated. Although the

occipital cortex had significantly higher pTau aggregate levels in PD, Path-Positive controls
and PD had similar intensity distributions, therefore occipital cortex tau pathology did not
appear PD-specific. Again, the putamen had significantly higher intensity aggregates in PD
compared to both control groups, and we used super-resolution imaging to determine that
their lengths ranged from 30–1,000 nm. The differences in putamen pTau aggregate sizes
were influenced by a subset of PD cases with mean length greater than 120 nm, and the
longest aggregates appeared fibrillar in shape from reconstructed super-resolution images.
One limitation of this study is that we were unable to use Thioflavin T with SiMPull to determine
if nanoscopic aggregates were beta-sheet positive due to high background fluorescence, and
greater structural analysis of nanoscopic tau aggregates would be of interest in the future.

Tau SiMPull and tau IHC aggregates were closely linked in the control group, however, that
relationship was disrupted in PD. In PD, tau SiMPull and IHC were only correlated in the
hippocampus, but there was no relationship in the amygdala or putamen, despite high levels
of bright nanoscopic tau aggregates. Comparing tau SiMPull with αSyn IHC coverage in PD
may provide insights into regional differences; there was a positive correlation between tau
SiMPull and αSyn IHC in the hippocampus, indicating a universal response by aggregation-
prone proteins in this region. In the amygdala, tau SiMPull was negatively correlated with αSyn
IHC, indicating a possible interaction between αSyn and tau; for example tau may be
incorporated into Lewy Bodies, resulting in the absence of nanoscopic tau aggregates (31).
An inherent limitation of working with post-mortem brain tissue is the inability to determine the
sequence of pathological events, so it is not possible to determine from this data whether Lewy
Bodies affect tau aggregate levels or vice versa. Nevertheless, these results indicate a distinct
distribution pattern for tau aggregates in PD compared to controls.

Combining aggregate measurements from SiMPull and IHC prompted two different
interpretations for the role of tau pathology in PD. We used principal component analysis to
divide the cohort into four clusters: Group 1 controls had low pathology in all brain regions;
Group 2 controls had high tau pathology that was associated with advanced age; Group 3 PD
had αSyn pathology and moderate tau pathology in the hippocampus and amygdala; Group 4
PD had αSyn pathology and high tau pathology, with a unique high-intensity nanoscopic tau
aggregate population in the putamen, and the highest rates of dementia. It should be noted
that PD-dementia was not a selection criterion for this cohort, so a follow-up study will
determine the strength of this relationship in a larger cohort of PD versus PD-dementia. The
different patterns of tau pathology in Groups 3 and 4 can be interpreted as: (1) Tau pathology
is independent of PD pathogenesis, beginning in the hippocampus and progressing along the
tau Braak stages, therefore Group 4 is more advanced in the process compared to Group 3;
(2) Tau aggregation initiates in the basal ganglia in a subset of PD cases like Group 4, and
progresses in parallel to age-related tau aggregation that begins in the hippocampus. In the
latter scenario, tau aggregates could have a central role in causing dopaminergic cell loss in
PD, as previously discussed. Alternatively, tau aggregation may be triggered by cell death in
the substantia nigra, for example by reduced dopamination of tau which was recently shown
to protect against aggregation (32). If tau is truly driving cell death in PD, it could be important
to stratify PD patients based on nigrostriatal tau pathology to identify a subpopulation who
would benefit from tau-targeting therapies in the future.

In conclusion, we leveraged single-molecule imaging to measure nanoscopic tau aggregates
in PD that may have been previously underestimated by pathological studies relying on IHC.
Our data provides evidence that nanoscopic tau aggregate formation in the putamen may
contribute to pathology in a subset of cases, and is linked to more advanced PD with dementia.
Future studies should aim to establish the sequence of pathological events and the prevalence

of nanoscopic putamen tau aggregates in the PD brain, which may represent a targetable

aggregate species for disease-modifying therapies.

Methods

Case Information

Post-mortem human brain tissue from 14 idiopathic PD cases and 15 control (CON) cases was acquired from the Cambridge Brain Bank. All research procedures utilising human brain tissue were conducted with approval by the London-Bloomsbury Research Ethics Committee (16/LO/0508) and with informed consent from the families of the donors. Idiopathic PD cases received a clinical diagnosis of PD during life and displayed post-mortem Lewy Body neuropathology. Clinical data pertaining to PD progression was available for a subset of PD cases who attended Cambridge University Hospital clinics during life or from death certificates, and Braak staging for αSyn and tau was assessed by neuropathologist Dr Annelies Quaegebeur (Table 1).

Control cases had no diagnosis of PD during life or PD-associated post-mortem neuropathology. However, some controls were documented as having dementia or other neurological disease on their death certifications (Table 1). Additionally, control cases CON5, CON6, CON7, CON8, CON9, CON11, CON12, and CON14 had tau neurofibrillary tangle deposition, although it was limited to the hippocampus and amygdala.

From the available information in the PD group, there was a mean disease duration of (10.28 ± 5.7 years), 5/14 cases had dementia, a range of Lewy Body Braak stages from IV–VI, and a range of tau Braak stages from I–III.

Upon arrival at the Brain Bank, brains were bisected and one half was flash-frozen and stored at −80 °C while the other half was fixed in 10% neutral buffered formalin for 14 days. The formalin-fixed tissue was embedded in paraffin blocks and cut into 10 μm sections for immunohistochemical analysis. Tissue sections from six brain regions were acquired for this study including hippocampus (PD n = 14; control n = 15), amygdala (PD n = 14; control n = 12), substantia nigra (PD n = 13; control n = 15), putamen (PD n = 14; control n = 15), occipital cortex (PD n = 14; control n = 15), and frontal cortex (PD n = 14; control n = 15).

Frozen tissue was acquired for the same cases and brain regions and processed for single-molecule imaging. It should be noted that frozen tissue availability varied for some regions, and the SN in particular was only available for a subset of cases: hippocampus (PD n = 12; control n = 11); amygdala (PD n = 12; control n = 12), substantia nigra (PD n = 5; control n = 4), putamen (PD n = 14; control n = 15), occipital cortex (PD n = 14; control n = 15), and frontal cortex (PD n = 14; control n = 15).

Table 1. Clinical information for all PD and Control (CON) brain tissue samples. N/A = Not available; AOD = Age of Death; PMI = Post-mortem interval; FTD = Frontotemporal dementia; MND = Motor neuron disease; GI = Gastrointestinal.

Case	Sex	AOD (years)	PMI (hours)	PD Duration (years)	Lewy Body Braak Stage	Tau Braak & Braak stage	Dementia status	Cause of Death
PD1	M	71	64	4.4	VI	II	N	N/A
PD2	M	85	8	18.1	VI	III	Y	PD Dementia
PD3	M	73	57	N/A	V	II	N	PD
PD4	F	78	68	5.2	V	II	N	Pneumonia
PD5	M	76	60	9.7	V	II	Y	Aspiration Pneumonia
PD6	M	77	101	8.5	V	III	N	Pneumonia
PD7	F	86	52	N/A	VI	II	N	PD
PD8	M	86	32	4.8	V	III	N	Myocardial Infarction
PD9	M	84	21	6.3	IV	II	N	Pneumonia
PD10	M	89	N/A	11.9	V	I	Y	Hypertensive heart disease
PD11	M	88	44	22.4	VI	III	N	Pneumonia
PD12	M	73	124	N/A	VI	II	N	Massive GI bleed
PD13	F	90	52	N/A	V	III	Y	Vascular dementia
PD14	M	75	8	11.5	VI	III	Y	PD Dementia
CON1	M	72	39	-	0	II	N	Malignant tumor of spinal cord
CON2	F	65	5	-	0	II	N	Idiopathic Pulmonary Fibrosis
CON3	M	39	59	-	0	0	N	Metastatic cancer of the colon
CON4	M	73	51	-	0	II	N	Myocardial infarction
CON5	M	62	63	-	0	I	N	Hepatocellular carcinoma
CON6	M	74	74	-	0	II	N	N/A
CON7	M	88	18	-	0	III	N	Acute haemopericardium
CON8	F	88	28	-	0	III	N	Multiorgan failure
CON9	F	77	59	-	0	III	N	Burkitts Lymphoma
CON10	F	75	92	-	0	II	N	Metastatic ovarian Cancer
CON11	M	87	49	-	0	III	Y	Dementia
CON12	F	86	67	-	0	III	N	Idiopathic Pulmonary Fibrosis
CON13	F	70	32	-	0	I	N	N/A
CON14	M	90	56	-	0	II	Y	FTD/MND
CON15	M	74	115	-	0	II	Y	Dementia

Immunohistochemistry (IHC)

Table 2. Primary antibodies used for IHC.

Target	Type/Species	Source	Ref	Dilution
αSyn	Polyclonal/Rabbit	Enzo Life Sciences	BML-SA3400-0100	1:250
Tau	Monoclonal/Mouse	Thermo Fisher Scientific	MN1020	1:500

IHC was performed by the Cambridge Brain Bank as previously described (33). In brief, tissue sections were deparaffined and hydrated by sequential incubation in xylene, ethanol, and water. The sections were pre-treated with formic acid for 5 minutes for additional antigen retrieval. Sections were incubated with hydrogen peroxide for 5 mins, followed by washing with water and phosphate-buffered saline (PBS). Sections were blocked using 2% Marvel in PBS for 20 mins, then incubated with the primary antibody (Table 2) in 2% Marvel for 1 hour. The sections were washed with PBS for 5 mins, then incubated with the secondary antibody (1% Rabbit horse radish peroxidase, diluted in 10% serum in PBS) for 1 hour. The sections were washed with PBS, then incubated with 3,3′-diaminobenzidine (DAB) (Vector, SK-4100). Finally, the sections were counterstained with hematoxylin to visualize cell nuclei, and sealed for imaging.

IHC imaging and analysis

Slide scanning at 20 x magnification was performed at the Cambridge Brain Bank or Cancer Research UK Cambridge Institute using the Aperio Scanscope AT2 (Leica Biosystems). Image pre-processing and quantitative analysis was performed using QuPath software (version 0.4.3) as described by a previously published protocol (34). Briefly, colour deconvolution was applied to the scanned brightfield whole slide images to separate the stains into different channels. Slides were then manually inspected to remove pigmented cells (in the substantia nigra), DAB artefacts, or folded tissue, and the region of interest (ROI) was outlined using the wand tool to exclude white matter. A thresholder tool in QuPath was applied to detect DAB-positive pixels (Resolution = Full (µm/pixel), Pre-filter = Gaussian, Smoothing sigma = 0.5, Threshold = 1.3). IHC coverage for each marker was calculated by:

$$100 \text{ x (area of positive pixels)/area of the ROI}$$

Frozen post-mortem human brain tissue fractionation

The steps for frozen brain tissue fractionation were adapted from a previously published (20). The case order of brain tissue processing was randomised to avoid technical bias and different dounce homogenisers were used for PD and CON cases to avoid cross-contamination. Briefly, 200-300 mg of fresh-frozen brain tissue was homogenised manually (Sigma-Aldrich, Cat. D8938) in five volumes (w/v) of ice cold low-salt buffer (124 mM NaCl, 2.8 mM KCl, 1.2 5mM NaH$_2$PO$_4$, 26 mM NaHCO$_3$; pH 7.32, supplemented with 5 mM EDTA, 1 mM EGTA, 5 µg/mL leupeptin, 5 µg/mL aprotinin, 2 µg/mL pepstatin, 20 µg/mL Pefabloc, 5 mM NaF), with 10 strokes of a high-pass pestle followed by 15 strokes of a low-pass pestle. The homogenate was centrifuged for 30 min at 21,000 xg at 4 ˚C and the supernatant was collected. The total protein concentration of the supernatant was measured immediately by Pierce bicinchoninic acid (BCA) protein assay kit (Thermo Scientific, Cat. 23227) according to the manufacturer's instructions, then aliquoted and stored at −80 ˚C.

Antibodies for single molecule pull-down (SiMPull)

Where possible, antibodies were acquired pre-labelled with biotin and Alexa Fluor (AF) dyes. The remaining antibodies were home-labelled as described below.

Table 3. Monoclonal antibodies used for SiMPull. Epitope refers to the amino acid sequence of the target protein. AF= Alexa fluor; N/A = Not available.

Target	ID	Species	Epitope	Source	Ref
αSyn	Syn211	Mouse	121-125	Santa Cruz	SC-12767
αSyn	Syn211-AF647	Mouse	121-125	Santa Cruz	SC-12767 AF647
αSyn	SOY1	Mouse	C-terminus	Merck	MABN1818
Tau	Biotin HT7	Mouse	159-163	Invitrogen	MN1000B
Tau	HT7	Mouse	159-163	Invitrogen	MN1000
pTau	Biotin AT8	Mouse	pSer202, Thr205	Invitrogen	MN1020B
pTau	AT8	Mouse	pSer202, Thr205	Invitrogen	MN1020

The pTau antibody AT8 was also used for super-resolution imaging with DNA points accumulation for imaging in nanoscale topography super-resolution imaging (DNA-PAINT). The antibody was conjugated with the following docking oligonucleotide sequence:

DBCO TEG-AAACCACCACCACCACCACCACCACCACCACCACCA

The complimentary imaging oligonucleotide was conjugated with Cy3b dye, with the sequence: TGGTGGT-AminoC7 - Cy3B

Antibody labelling

Capture antibodies were biotinylated, and detection antibodies were conjugated with a docking oligonucleotide for DNA-PAINT, using the SiteClick™ Antibody Azido Modification Kit (Invitrogen, Cat. S20026) according to the manufacturer's instructions. In brief, at least 150 µg of antibody without preservatives or salts was concentrated and buffer exchanged with Buffer A using a 50 kDa MWCO Amicon spin filter (Merck Millipore, Cat. UFC505024). The antibody was recovered and incubated with beta galatosidose overnight at 37 ˚C. The next day, components E, F, G, and H were combined, and the antibody was added to the mixture. The mixture was incubated overnight at 30 ˚C. The next day, the mixture was concentrated using a 50 kDa MWCO Amicon spin filter and 10 molar equivalents of biotin were added, followed by another overnight incubation at 37 ˚C. Finally, excess biotin was removed using a 40 kDa MWCO Zeba Spin Desalting Column (Thermo Scientific, Cat. A57766), the antibody was concentrated using a 50 kDa MWCO Amicon Spin filter, and the concentration was determined by Nanodrop A280.

The Zip Alexa Fluor™ 647 Rapid Antibody Labeling Kit (Invitrogen, Cat. Z11233) was used to label detection antibodies. The antibodies were first concentrated and buffer exchanged with

PBS using a 50 kDa MWCO Amicon spin filter. The antibody was recovered and incubated
with 1/10 volume 1 M sodium bicarbonate and 10 molar equivalents of the appropriate dye for
1 hour at 37 ˚C. The reaction was quenched by diluting 1:1 in 1 M Tris pH 8, then excess dye
was removed using a 40 kDa MWCO kDa Zeba Spin Desalting Column. The antibody was
concentrated using a 50 kDa MWCO Amicon spin filter, then the final concentration and the
dye-labelling efficiency were determined by Nanodrop.

HT7-binding peptide sequence

548 A dimer of the HT7-binding peptide sequence (KK(PEG)8GAAPPGQKGQ, Cat. No.
549 U9131MEFG0-9, Genscript) was used for tau SiMPull, as previously described (25). The dimer
550 contains two binding sites for the total-tau antibody HT7; when used for SiMPull, the dipeptide
551 may bind the capture antibody and a single detection antibody.

Silica nanoparticle (SiNaP) aggregate standards

553 Tau SiNaP aggregate standards were prepared according to the previously published protocol
554 by Böken et al., (2024), by conjugating the HT7 antibody binding sequence to the SiNaPs, and
555 αSyn SiNaP aggregate standards were prepared according to the previously published
556 protocol by Herrmann et al., (2017) (36, 37).

Coverslip passivation for single-molecule pull-down

Glass coverslips (26 x 76 mm, Epredia) were cleaned using an argon plasma cleaner (PDC-
002, Harrick Plasma) for 30 minutes, then a 50-well PDMS gasket (Grace Bio-Labs, Ref.
103250) was attached to each coverslip. The wells were loaded with 6 µL of coating solution
(1:1 mixture of Rain-X and isopropanol, filtered using a 0.22 µm PVDF filter (Millex, Ref.
SLGV004SL)), and left to evaporate at room temperature as previously described (38). The
surfaces were always prepared the day before a SiMPull experiment.

Single-molecule pull-down (SiMPull)

Tris-buffered saline (TBS, Fisher Bioreagents, Cat. BP2471-500) and TBS-Tween (TBS-T,
Rockland antibodies and assays, Cat. MB-013) buffers were filtered using a 0.2 µm PVDF
filter (Millex, Ref. SLGP033RS) immediately prior to use. A 0.22 µm PVDF filter (Millex, Ref.
SLGV004SL) was used for all other reagents that are described as filtered. All reagents or
wash buffers were added to the wells in 10 µL volumes, and all incubations were carried out
at room temperature.

Pre-prepared Rain-X coated coverslips were placed in a humidity chamber to prevent the
surface from drying out. The wells were washed twice with TBS (Gibco, Ref. 10010-015), then
incubated with NeutrAvdin (Thermo Scientific, Ref. 31050) at 0.1 mg/mL in TBS for 15 minutes.
The wells were washed three times with TBS, then incubated for 1 hour with 1% pluronic F-
127 polymer (Biotium, Cat. 59005, 10% in water) diluted 1:10 in TBS and filtered. The wells
were washed three times with TBS-T (1x PBS with 0.05% Tween 20; Rockland, Code MB-
075-1000), then incubated with 10 nM biotinylated Capture antibody for 15 minutes. The wells
were washed twice with TBS-T, then incubated with filtered blocking solution (500 µL TBS-T,
50 µL BSA (R&D Systems, Cat. 841380), 20 µL normal goat serum (Abcam, Cat. Ab7481), 20
µL recombinant BSA (New England BioLabs, Cat. B9200S)), for 30 minutes to prevent non-

specific protein adsorption to the surface. The wells were subsequently washed twice with
TBS-T before sample addition.

For tau and pTau SiMPull: brain samples were diluted 2-fold in TBS and incubated for 2 hours.
For αSyn SiMPull: brain samples were added to the surface without dilution and incubated for
3 hours. Samples were run in triplicate for each brain region and antibody pair, with one well
per case used on each surface. For every surface, one well was incubated with TBS (negative
control), and another was incubated with a single-detection antibody control (HT7-dipeptide
for tau SiMPull, capture and detect with HT7; αSyn monomer for αSyn SiMPull, capture with
SOY1 and detect with Syn211). Following sample incubation, the wells were washed four
times with TBS-T.

For diffraction-limited imaging, the wells were incubated with 5 nM detection antibody labelled
with Alexa-Fluor 647 in filtered 1% BSA in TBS-T for 15 minutes. For multichannel imaging,
both detection antibodies were added at a final concentration of 5 nM. Finally, the wells were
washed three times with TBS-T, then the buffer was exchanged with TBS for image acquisition.

For super-resolution imaging, another blocking step was performed with filtered 1% BSA in
TBS-T for 30 minutes. Next, the DNA-conjugated detection antibody diluted in TBS was
incubated for 15 minutes. The wells were washed twice with TBS-T, then 1 nM Cy3b-
conjugated imaging strand was added to the wells, and a coverslip was added on top to
prevent evaporation throughout imaging.

Diffraction-limited image acquisition and analysis

Diffraction-limited imaging was performed using a home-built total internal reflection
fluorescence (TIRF) microscope using an inverted Ti-E Eclipse body (Nikon Corporation) fitted
with a 1.49 N.A, 100x objective (Apo TIRF, Nikon) and in-built Perfect Focus System. The
microscope was fitted with 488 nm (iBeam – SMART, Toptica) and 638 nm lasers (MLD, 638,
Cobalt), and the fluorescent emissions was collected by an EMCCD camera (Evolve 512,
Photometrics) through a 1.5x beam expander, with an electron-multiplying gain of 214
ADU/photon, sensitivity of 3.019 electrons/ADU, and offset of 604.2 ADU. Each pixel
corresponded to 109.2 nm and image acquisition (50 frames, 50 ms exposure time, 12 fields
of view) was controlled with MicroManager (39).

Automated analysis of single-molecule localisation was performed in ImageJ (40). The code
was adapted

from https://github.com/YPZ858/DF-single-molecule-couting-/blob/main/DF21.ijm. Briefly, the
image frames 10–50 were combined into a z-stack, then the ThunderSTORM plugin was used
to apply a wavelet filter of 2*std(Wave.F1) and detect single molecules (41). The results were
exported as an Excel spreadsheet and further filtering was performed to retain data that met
the following criteria: higher intensity than the mean intensity of the blank (TBS only);
replicates with positive signal in at least 9 out of 12 fields of view; cases with positive signal in
at least 2 out of 3 replicates.

Example images were prepared in ImageJ by first performing a z-stack of frames 10–50. The
greyscale values were calibrated to photons using the ThunderSTORM plugin, and images
were contrast-adjusted with a minimum value of 0 and a maximum value corresponding to the
range of spot intensities in the image. The calibration bar was used to demonstrate the contrast

ranges of different images, and a higher maximum value corresponded to higher intensity
spots. It should be noted that the maximum value of the calibration bar does not necessarily
indicate the maximum photon value in the image, as contrast ranges were selected as a
compromise between multiple images being compared. The field of view was cropped to half
of the original size to enlarge the image sufficiently for demonstration purposes.

Intensity calibration to calculate number of epitopes per aggregate

The intensity of a single detection antibody was used to calculate the approximate number of
epitopes per aggregate. To obtain this intensity value, HT7 dipeptide and αSyn monomers
(Abcam, cat. AB218818) were used for tau and αSyn SiMPull respectively (Methods Figure
1). To account for the epitope bound by the capture antibody, +1 was added.

# of epitopes/aggregate = (Intensity(Aggregate) / Intensity(Single detection antibody)) +1

Methods Figure 1. Intensity calibrants to calculate the approximate number of epitopes
per aggregate. Schematic representation of **(a)** αSyn and **(b)** tau intensity calibrants used to
measure the intensity of a single detection antibody for calculating the approximate number of
epitopes per aggregate.

Super-resolution image acquisition and analysis

Super-resolution imaging was performed with DNA-PAINT using the same optical setup as
diffraction-limited imaging. Images were acquired using the 561 nm laser (Jive, Cobalt), with
8,000 frames, 100 ms exposure time, and 4 fields of view.

DNA-PAINT image reconstruction and analysis was performed using the Picasso package, as
previously described (25, 42). The custom scripts are available at
https://zenodo.org/records/8027256. In brief, the first 300 frames were discarded, localisations
were identified and fitted, using redundant cross-correlation to correct for microscope drift. The
localisations were filtered using a 30 nm precision threshold, then assigned to clusters with
DBSCAN using the scitkit-learn package (radius of 2 and minimum density of 1, minimum of
3 localisations per cluster) corresponding to individual aggregates (43). Each aggregate was
measured in terms of skeletonised length using SKAN, calculated as the summed branch
distance (44). The results were exported as an Excel spreadsheet, and were further filtered to
retain clusters that had a higher number of localisations compared to the blank (TBS only).

DNA extraction and MAPT genotyping

DNA was isolated from fresh-frozen brain tissue using the QIAamp DNAMini and Blood Mini
kit according to the manufacturer's instructions. H1 and H2 MAPT haplotype was determined

by Taqman assay (Assay ID: C___7563692_10; SnP ID: rs1800547) using the QuantStudio 5
detection system (Applied Biosystems).

Statistical analysis in R

All statistical analysis was performed in R (Version 4.2.2) and all graphs were plotted using the packages ggplot2, tidyverse, plyr, ggpubr, scales, ggh4x, ggrepel (45-52). The appropriate statistical test was determined after checking the data met the normality assumptions, and p-values of less than 0.05 were considered significant. The sample size for all whole-cohort comparisons was PD n = 14 and CON n = 15. For comparisons of IHC coverage, mean SiMPull density or mean SiMPull intensity between PD and CON, non-parametric two-sided Mann-Whitney test was used. For linear correlation analysis of human samples, data was not normally distributed therefore the non-parametric Spearman correlation was performed using stat_cor in the R package ggpubr, and the correlation coefficient rho (ρ) was displayed. For linear correlation analysis of SiNaP SiMPull, data was normally distributed, therefore the Pearson correlation was performed using stat_cor in the R package ggpubr, and the correlation coefficient R^2 was displayed. Box plots show the first quartile to third quartile (box) with the median marked, and whiskers extend to 1.5 times the interquartile range. For intensity distribution histograms, intensity was plotted on a $\log_{10}$ x-axis scale, and the y-axis was plotted as percentage per group to account for the different sample sizes. For multiple comparisons of age of death, non-parametric Dunn's multiple comparison test with Benjamini-Hochberg p-value adjustment was used.

The following input variables were used for principal component analysis (PCA): αSyn IHC coverage (%) for hippocampus, amygdala, putamen, substantia nigra; Tau IHC coverage (%) for hippocampus, amygdala, putamen, substantia nigra; Tau SiMPull mean bright aggregates (%) for hippocampus, amygdala, putamen; pTau SiMPull mean intensity (photons) for hippocampus, amygdala, putamen. Results from the frontal cortex and occipital cortex were excluded due to low signal for all markers, and SiMPull results from the substantia nigra were excluded due to low sample size. For each variable, missing were imputed with the median, and data was centred and scaled prior to PCA. Subsequently, k-means cluster analysis was performed using principal components 1 and 2. For comparisons between PCA groups for age of death, dementia status, or MAPT genotype: Group 1 n = 7; Group 2 n =8; Group 3 n = 9; Group 4 n = 5.

Acknowledgements: We gratefully acknowledge the contributions of our brain tissue donors and their families and the Cambridge Brain and Tissue Bank. We acknowledge the histology core at the Cancer Research UK Cambridge institute for their technical support. The work was supported by a grant from the UK Medical Research Council.

Author Contributions: F.L.: Tissue processing, single-molecule assays, immunohistochemistry image analysis, data analysis, manuscript preparation, conception and design. D.B.: Development of tau assays, super-resolution data analysis. Y.Z.: Assay development, technical support. K.H.: Tissue preparation, immunohistochemistry. D.R.: Tissue preparation, immunohistochemistry. L.K.: Tissue preparation, technical support. B.P.: Data collection. A.Q.: Neuropathological assessment, manuscript preparation. C.H.W.G.: Manuscript preparation, conception and design. D.K.: Manuscript preparation, conception and design, supervision. The final manuscript was edited and approved by all authors.

References

1. Greenfield J, Bosanquet FD. The brain-stem lesions in Parkinsonism. Journal of neurology, neurosurgery, and psychiatry. 1953;16(4):213.

2. Damier P, Hirsch E, Agid Y, Graybiel A. The substantia nigra of the human brain: II. Patterns of loss of dopamine-containing neurons in Parkinson's disease. Brain. 1999;122(8):1437-48.

3. Greffard S, Verny M, Bonnet A-M, Beinis J-Y, Gallinari C, Meaume S, et al. Motor score of the unified Parkinson disease rating scale as a good predictor of Lewy body–associated neuronal loss in the substantia nigra. Archives of neurology. 2006;63(4):584-8.

4. Spillantini MG, Crowther RA, Jakes R, Hasegawa M, Goedert M. α-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. Proceedings of the National Academy of Sciences. 1998;95(11):6469-73.

5. Braak H, Del Tredici K, Rüb U, De Vos RA, Steur ENJ, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. Neurobiology of aging. 2003;24(2):197-211.

6. Braak H, Del Tredici K. Neuropathological staging of brain pathology in sporadic Parkinson's disease: separating the wheat from the chaff. Journal of Parkinson's disease. 2017;7(s1):S71-S85.

7. Bega D, Kuo PH, Chalkidou A, Grzeda MT, Macmillan T, Brand C, et al. Clinical utility of DaTscan in patients with suspected Parkinsonian syndrome: a systematic review and meta-analysis. npj Parkinson's Disease. 2021;7(1):43.

8. Gomes BF, Farris CM, Ma Y, Concha-Marambio L, Lebovitz R, Nellgård B, et al. α-Synuclein seed amplification assay as a diagnostic tool for parkinsonian disorders. Parkinsonism & Related Disorders. 2023;117:105807.

9. Gibbons CH, Levine T, Adler C, Bellaire B, Wang N, Stohl J, et al. Skin biopsy detection of phosphorylated α-synuclein in patients with synucleinopathies. Jama. 2024;331(15):1298-306.

10. Schaffrath A, Schleyken S, Seger A, Jergas H, Özdüzenciler P, Pils M, et al. Patients with isolated REM-sleep behavior disorder have elevated levels of alpha-synuclein aggregates in stool. npj Parkinson's Disease. 2023;9(1):14.

11. Smith R, Capotosti F, Schain M, Ohlsson T, Vokali E, Molette J, et al. The α-synuclein PET tracer [18F] ACI-12589 distinguishes multiple system atrophy from other neurodegenerative diseases. Nature communications. 2023;14(1):6750.

12. Binder LI, Guillozet-Bongaarts AL, Garcia-Sierra F, Berry RW. Tau, tangles, and Alzheimer's disease. Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease. 2005;1739(2-3):216-23.

13. Braak H, Alafuzoff I, Arzberger T, Kretzschmar H, Del Tredici K. Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. Acta neuropathologica. 2006;112(4):389-404.

14. Chin KS, Yassi N, Churilov L, Masters CL, Watson R. Prevalence and clinical associations of tau in Lewy body dementias: A systematic review and meta-analysis. Parkinsonism & related disorders. 2020;80:184-93.

15. Pascale E, Di Battista ME, Rubino A, Purcaro C, Valente M, Fattapposta F, et al. Genetic architecture of MAPT gene region in Parkinson disease subtypes. Frontiers in cellular neuroscience. 2016;10:96.

16. Dickson DW, Rademakers R, Hutton ML. Progressive supranuclear palsy: pathology and genetics. Brain Pathology. 2007;17(1):74-82.

17. Schneider JA, Li JL, Li Y, Wilson RS, Kordower JH, Bennett DA. Substantia nigra tangles are related to gait impairment in older persons. Annals of neurology. 2006;59(1):166-73.

18. Chu Y, Hirst WD, Federoff HJ, Harms AS, Stoessl AJ, Kordower JH. Nigrostriatal tau pathology in parkinsonism and Parkinson's disease. Brain. 2024;147(2):444-57.

19.   Parkkinen L, Kauppinen T, Pirttilä T, Autere JM, Alafuzoff I. α-Synuclein pathology does
not predict extrapyramidal symptoms or dementia. Annals of Neurology: Official Journal of the
American Neurological Association and the Child Neurology Society. 2005;57(1):82-91.
20.   Emin D, Zhang YP, Lobanova E, Miller A, Li X, Xia Z, et al. Small soluble α-synuclein
aggregates are the toxic species in Parkinson's disease. Nature communications.
2022;13(1):5512.
21.   Hong W, Wang Z, Liu W, O'Malley TT, Jin M, Willem M, et al. Diffusible, highly bioactive
oligomers represent a critical minority of soluble Aβ in Alzheimer's disease brain. Acta
neuropathologica. 2018;136:19-40.
22.   Winner B, Jappelli R, Maji SK, Desplats PA, Boyer L, Aigner S, et al. In vivo
demonstration that α-synuclein oligomers are toxic. Proceedings of the National Academy of
Sciences. 2011;108(10):4194-9.
23.   Jiang L, Zhao J, Cheng J-X, Wolozin B. Tau oligomers and fibrils exhibit differential
patterns of seeding and association with RNA binding proteins. Frontiers in Neurology.
2020;11:579434.
24.   Lobanova E, Whiten D, Ruggeri FS, Taylor CG, Kouli A, Xia Z, et al. Imaging protein
aggregates in the serum and cerebrospinal fluid in Parkinson's disease. Brain.
2022;145(2):632-43.
25.   Böken D, Cox D, Burke M, Lam JY, Katsinelos T, Danial JS, et al. Single-Molecule
Characterization and Super-Resolution Imaging of Alzheimer's Disease-Relevant Tau
Aggregates in Human Samples. Angewandte Chemie. 2024;136(21):e202317756.
26.   Kumar ST, Jagannath S, Francois C, Vanderstichele H, Stoops E, Lashuel HA. How
specific are the conformation-specific α-synuclein antibodies? Characterization and validation
of 16 α-synuclein conformation-specific antibodies using well-characterized preparations of α-
synuclein monomers, fibrils and oligomers with distinct structures and morphology.
Neurobiology of disease. 2020;146:105086.
27.   Bartels T, Choi JG, Selkoe DJ. α-Synuclein occurs physiologically as a helically folded
tetramer that resists aggregation. Nature. 2011;477(7362):107-10.
28.   Sanderson JB, De S, Jiang H, Rovere M, Jin M, Zaccagnini L, et al. Analysis of α-
synuclein species enriched from cerebral cortex of humans with sporadic dementia with Lewy
bodies. Brain communications. 2020;2(1):fcaa010.
29.   Zhai S, Shen W, Graves SM, Surmeier DJ. Dopaminergic modulation of striatal
function and Parkinson's disease. Journal of Neural Transmission. 2019;126(4):411-22.
30.   Owens-Walton C, Jakabek D, Li X, Wilkes FA, Walterfang M, Velakoulis D, et al. Striatal
changes in Parkinson disease: An investigation of morphology, functional connectivity and
their relationship to clinical symptoms. Psychiatry Research: Neuroimaging. 2018;275:5-13.
31.   Ishizawa T, Mattila P, Davies P, Wang D, Dickson DW. Colocalization of tau and alpha-
synuclein epitopes in Lewy bodies. Journal of Neuropathology & Experimental Neurology.
2003;62(4):389-97.
32.   Wang Q, Liu Z, Wang Y, Liu Y, Chen Y, Zhang S, et al. Quantitative chemoproteomics
reveals dopamine's protective modification of Tau. Nature Chemical Biology. 2025:1-10.
33.   Pansuwan T, Quaegebeur A, Kaalund SS, Hidari E, Briggs M, Rowe JB, et al. Accurate
digital quantification of tau pathology in progressive supranuclear palsy. Acta
neuropathologica communications. 2023;11(1):178.
34.   Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD, et
al. QuPath: Open source software for digital pathology image analysis. Scientific reports.
2017;7(1):1-7.
35.   Böken D, Xia Z, Lam JY, Fertan E, Wu Y, English EA, et al. Ultrasensitive Protein
Aggregate Quantification Assays for Neurodegenerative Diseases on the Simoa Platform.
Analytical Chemistry. 2024;97(1):290-9.
36.   Herrmann Y, Bujnicki T, Zafiu C, Kulawik A, Kühbach K, Peters L, et al. Nanoparticle
standards for immuno-based quantitation of α-synuclein oligomers in diagnostics of
Parkinson's disease and other synucleinopathies. Clinica chimica acta. 2017;466:152-9.

37.   Boken D, Xia Z, Lam JY, Fertan E, Wu Y, English EA, et al. Ultrasensitive protein
aggregate quantification assays for neurodegenerative diseases on the simoa platform.
Analytical Chemistry. 2024;97(1):290-9.

38.   Zhang YP, Lobanova E, Dworkin A, Furlepa M, Yang WS, Burke M, et al. Improved
imaging surface for quantitative single-molecule microscopy. ACS Applied Materials &
Interfaces. 2024;16(28):37255-64.

39.   Edelstein AD, Tsuchida MA, Amodaj N, Pinkard H, Vale RD, Stuurman N. Advanced
methods of microscope control using µManager software. Journal of biological methods.
2014;1(2).

40.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji:
an open-source platform for biological-image analysis. Nature methods. 2012;9(7):676-82.

41.   Ovesný M, Křížek P, Borkovec J, Švindrych Z, Hagen GM. ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution
imaging. Bioinformatics. 2014;30(16):2389-90.

42.   Schnitzbauer J, Strauss MT, Schlichthaerle T, Schueder F, Jungmann R. Super-
resolution microscopy with DNA-PAINT. Nature protocols. 2017;12(6):1198-228.

43.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-
learn: Machine learning in Python. the Journal of machine Learning research. 2011;12:2825-
30.

44.   Nunez-Iglesias J, Blanch AJ, Looker O, Dixon MW, Tilley L. A new Python library to
analyse skeleton images confirms malaria parasite remodelling of the red blood cell
membrane skeleton. PeerJ. 2018;6:e4312.

45.   Wickham H. Data analysis. ggplot2: elegant graphics for data analysis: Springer;
2016. p. 189-201.

46.   Slowikowski K. Ggrepel: automatically position non-overlapping text labels
with'ggplot2'. R package version 0.9. 6.9999. 2025.

47.   Kassambara A. ggpubr:'ggplot2'based publication ready plots. R package version.
2018:2.

48.   Wickham H, Wickham MH, RColorBrewer I. Package 'scales'. R package version.
2016;1(0).

49.   Wickham H, Averick M, Bryan J, Chang W, McGowan LDA, François R, et al. Welcome
to the Tidyverse. Journal of open source software. 2019;4(43):1686.

50.   Wickham H. The split-apply-combine strategy for data analysis. Journal of statistical
software. 2011;40:1-29.

51.   Brand Tvd. ggh4x: Hacks for 'ggplot2' Github2025 [Available from:
https://github.com/teunbrand/ggh4x.

52.   Team RS. RStudio: Integrated development environment for R. (No Title). 2021.

842