1 **Title:** Population genomics highlight the vulnerability of coral-dwelling gobies to ecological
2 losses due to climatic disturbances
3

Authors: Catheline Y.M. Froehlich[123], Renae L. Kirby[1], Samuel Greaves[4], Siobhan J.
Heatwole[1], O. Selma Klanten[5], Martin Hing[1], Marian Y.L. Wong[1]
Affiliations:
[1]School of Sciences
Faculty of Science, Medicine and Health
University of Wollongong
Wollongong, New South Wales, 2500 Australia
11
12 [2]School of Biological Sciences
13 University of Alabama
14 Tuscaloosa, Alabama, 35487, USA
15
16 [3]Dauphin Island Sea Lab
17 Dauphin Island, Alabama, 36528, USA
18
19 [4]Department of Biology
20 University of Central Florida
21 Orlando, FL, 32816, USA
22
23 [5]School of Life Sciences
24 University of Technology Sydney
25 Sydney, New South Wales, 2007, Australia
26

27 **Statement of authorship:** CF OK MW designed the study; CF MH RK SH OK MW
28 collected the data; CF RK analyzed the data; SG provided and supervised methods; CF wrote
29 the draft; all authors contributed to revisions.
30
31

32 **Keywords (max 10):** genetic bottlenecks; cyclones; heatwaves; coral-dwelling fishes;
33 compounding effects; population structure; climate change; ecology;

34
35 **Data Statement:** The data is available on NCBI project: PRJNA1309028.
36
37 **Abstract**
38

Predicting species resilience to climate change generally focuses on species life history and
ecology, overlooking genomic changes following disturbances. After consecutive cyclones
and heatwaves, coral-dwelling *Gobiodon* gobies have experienced extreme population and
group size reductions, and slower recovery rates than their coral hosts. To assess genomic
contributions to this vulnerability, we analyzed multiple *Gobiodon* species for population
structure (1) among three locations in western Oceania and (2) pre- and post-disturbances at
the central location that underwent the same cyclones and heatwaves that reduced populations
and group sizes. Species compared among locations all exhibited population structure. Post-
disturbances, two species (*G. fuscoruber, G. rivulatus*) were extirpated, two species (*G.
histrio, G. quinquestrigatus*) exhibited genetic bottlenecks, one species had no change to
genetic diversity (*G. brochus*), and another species had consistently low diversity (*G.
erythrospilus*). Our results highlight that coral-dwelling gobies face compounding ecological
and genomic losses from climatic disturbances, warranting targeted conservation.

MAIN TEXT

Introduction

Climate change is driving more frequent and intense disturbances, reshaping ecosystems at an alarming pace (AghaKouchak *et al.* 2018; Hughes *et al.* 2018a; Seidl *et al.* 2017). Most ecosystems face one or more disturbances, such as drought, floods, fires, storms, heatwaves, ocean acidification, increasing salinity, species invasions, and pollution (Bergstrom *et al.* 2021). This can lead to a variety of changes, like disease outbreaks, habitat degradation, behavioral modifications, altered species interactions, population declines, extirpations, and even ecosystem collapse (Sergio *et al.* 2018; Thakur *et al.* 2022; Turner 2010). To understand to what extent organisms are facing climate-related threats, it is crucial to study the impacts of climate change across multiple levels of biological organization (Attrill & Depledge 1997; Borges *et al.* 2021; Hourigan *et al.* 1988; Krishnan *et al.* 2013; Pinna *et al.* 2023). While ecological and physiological studies show that most species face mounting pressures due to climate change (Hill *et al.* 2021; Hughes *et al.* 2018b; Johnstone *et al.* 2016; Pinsky *et al.* 2022), understanding the extent of gene flow among populations has also been central to predicting their resilience and refining management strategies (Baptista *et al.* 2025; Cooke *et al.* 2016; Pauls *et al.* 2013; Woodall *et al.* 2011). By investigating the link between population structure and the potential for genetic bottlenecks, we can provide greater insights into species persistence with continued disturbances and guide effective management strategies (Jangjoo *et al.* 2016; Munguía-Vega *et al.* 2015; Ngeve *et al.* 2023). By combining insights from ecology, population connectivity, and genetic bottlenecks, we can better forecast species' resilience and refine conservation strategies that aim to mitigate the impacts of climate change (Teng *et al.* 2025).

Reef fishes face major setbacks immediately following climatic disturbances, with coral-dwelling fishes being particularly vulnerable (Bellwood *et al.* 2012; Froehlich *et al.* 2021; Richardson *et al.* 2018). Our recent studies have shown that coral-dwelling gobies (genus *Gobiodon*) declined considerably after consecutive cyclones and bleaching events, as they suffered extreme population losses, and recovered slower than their coral hosts (Froehlich *et al.* 2021, 2023). Nine months after these consecutive disturbances, only a few adult and juvenile gobies were found. Although the community of coral hosts differed three years post-disturbances, only a few goby species switched hosts and the majority of corals were left uninhabited (Froehlich *et al.* 2023). When forcibly displaced—as can occur during cyclones—gobies overwhelmingly risked predation to return to their home coral rather than accepting a new host (Froehlich *et al.* 2022). Post-disturbance, these gobies no longer lived in groups (Froehlich *et al.* 2024), despite the well-documented benefits of group living for their survival and reproduction (Wong & Buston 2013). Together, these findings paint a troubling picture of severe and multifaceted changes to goby ecology and population dynamics following extreme climatic events. Such losses raise crucial questions about whether gobies possess long-term resilience to ongoing climate change.

Here, we combined two complementary approaches to explore the interplay between genetic connectivity and diversity of *Gobiodon* populations and declines in their populations from climatic disturbances. We asked two questions: 1) from a spatial perspective, is there consistent gene flow among different areas within their ranges that could facilitate recovery? And 2) from a disturbance perspective, can genetic bottlenecks be detected only three years after consecutive disturbances that triggered ecological and population declines? To address these questions, we sampled goby populations at three locations when in relatively healthy states: Papua New Guinea (Kimbe Bay), northern Great Barrier Reef (Lizard Island), and southern Great Barrier Reef (One Tree Island). We then revisited Lizard Island after the

disturbances that resulted in extreme population and ecological (i.e. host types and sociality)
losses to determine whether there were also losses in genetic diversity and signals of genetic
bottlenecks. Together, these approaches revealed how ecological changes and mass mortality
following extreme disturbances were coupled with population structure to result in
extirpations and genetic bottlenecks for several *Gobiodon* species. Our study underscores the
high vulnerability of coral-dwelling gobies to climate disturbances and need for additional
protection.
111
112 **Methods**
113
114 Study Locations
115
Sampling was completed at three different locations in western Oceania (Fig. 1). The
northern most location was Kimbe Bay, Papua New Guinea (PNG: -5.42896°, 150.09695°).
We visited PNG from 25 Sep-3 Nov 2018 when the reef was relatively undisturbed. The
central location was Lizard Island, northern Great Barrier Reef in Australia (LI; -14.687264°,
145.447039°). We visited LI from 30 Oct-5 Nov 2013 and 4 Feb-26 Feb 2014 when the reef
was relatively undisturbed, and then from 15 Jan-5 Mar 2020 after the reef had been exposed
to four consecutive years of extreme disturbances (category 4 cyclones Ita in 2014 and
Nathan in 2015, and two mass bleaching events from extended marine heatwaves in 2016 and
2017). The southernmost location was One Tree Island, southern Great Barrier Reef in
Australia (OTI; -23.506565°, 152.090954°). We visited OTI from 28 Jan-19 Feb 2019 when
the reef was relatively undisturbed. The direct distance (point to point) between PNG and LI
was ~1140 km, between LI and OTI was ~1200 km, and between PNG and OTI was ~2000
km.
129

131 Fig. 1. Three sampling locations with timelines of sampling events (dashed lines) and
132 climatic disturbances (downward arrows; black spiral symbol for cyclones and white coral
133 symbol for marine heatwaves that caused mass bleaching) for all species studied in western
134 Oceania.
135
136 Sample Collection
137
Tissue collection was completed on SCUBA. At each location, we searched for gobies within
their *Acropora* coral hosts with the aid of an underwater flashlight. We selected goby species
based on their abundance at all locations. When encountered, gobies were temporarily

removed from the coral using an anesthetic solution (1:1:1 clove oil: seawater:70% ethanol)
for anesthesia and a hand wafting motion around the coral to direct them into hand nets.
Gobies were identified to species level and a small fin clip was collected from their caudal
fin, i.e. tail. Individuals were then returned to their coral hosts once they had recovered from
the anesthesia. Collected tissues were stored in 80% ethanol.
146
147　DNA Extraction and Sequencing
148
From Papua New Guinea, we sequenced 36 *G. fuscoruber*, 37 *G. histrio*, 43 *G.*
quinquestrigatus, and 31 *G. rivulatus*. From Lizard Island pre-disturbances, we sequenced 47
G. brochus, 43 *G. erythrospilus,* 38 *G. fuscoruber*, 50 *G. histrio*, 53 *G. quinquestrigatus*, and
47 *G. rivulatus*. From Lizard Island post-disturbances, we sequenced 33 *G. brochus*, 33 *G.*
erythrospilus, 40 *G. histrio*, and 41 *G. quinquestrigatus.* From One Tree Island, we sequenced
31 *G. fuscoruber*, 29 *G. quinquestrigatus*, and 29 *G. rivulatus.* We extracted DNA from the
tissue at the University of Wollongong and Sydney Institute for Marine Science using the SV
Genomic DNA Purification System (Promega) following the manufacturer's protocol. We
concentrated the DNA at 60°C in a vacuum centrifuge. We sent the samples to the Australian
Genome Research Facility (AGRF) for library preparation and sequencing. Samples were
prepared following the Peterson et al. (2012) protocol for double digest restriction-site
associated digest (ddRAD) with restriction enzymes EcoRI and MseI, and size selected for a
60 bp region (350-410 bp fragments). The samples were sequenced on an Illumina NextSeq
500 (1x150 cycles mid-output) at AGRF.
163
164　Data Analysis
165
Initial quality control and demultiplexing was performed by AGRF using STACKS v.2.3d
software (Catchen *et al.* 2013, 2011). Sequences that had exact barcodes were retained and
were then trimmed to 150bp. We then performed genotyping and Single Nucleotide
Polymorphism (SNP) filtering using STACKS v2.54 (Rochette *et al.* 2019). Parameters used
for denovo pipeline were as follows: six for the distance between stacks (-M), six for the
distance between catalogue loci (-n) and three for the minimum stack depth of coverage (-m).
These values were optimized following Rochette & Catchen (2017). Samples with less than
50% of loci called were excluded from analysis.
174
The data was divided into datasets for analyzing (1) differences among locations in relatively
healthy states (PNG 2018, LI 2014, OTI 2019) for each species and (2) differences among
pre- (2014) and post-disturbances (2020) (LI only) for each species. By using the
"populations" component of STACKS, loci were exported by calling more than 80% of
samples (-r) that were present in all populations (-p; 3 for among locations, except 2 for *G.*
histrio which was only found at two locations; 2 for pre/post disturbances), had less than 0.7
observed heterozygosity (--max-obs-het), and had a minimum of 2 minor allele count (--min-
mac). If loci were significantly out of Hardy-Weinberg Equilibrium (HWE) in two or more
sampling locations, then they were removed. To properly calculate HWE, the Hardy-
Weinberg (--hwe) option was used within populations and the p-values were then adjusted
with the "BY" correction of the "p.adjust" function in R v3.6.3 (Rstudio V.1.4.1106) (R Core
Team 2020; RStudio Team 2020) to account for False Discovery Rate. To avoid closely
linked SNPs, the --write-single-snp option was used to export the final set of loci.
188
Population structure among the different locations (PNG, LI, OTI;) and pre and post-
disturbances (LI only) were quantified for analysis of molecular variance (AMOVA) using
Arlequin v3.5.2.2(Excoffier & Lischer 2010). For the AMOVA, 30,000 permutations were
used and pairwise F_{ST} values were generated to assess hierarchical population structure

among populations if the AMOVA was significant. The observed heterozygosity (H_0),
expected heterozygozity (H_e), and inbreeding coefficient (F_{IS}) were calculated as genetic
diversity statistics using the "basic.stats" function in the R package hierfstat (Goudet 2005).
To determine whether there was asymmetric genetic migration, we used the divMigrate
function (Sundqvist *et al.* 2016) from the diveRsity package (Keenan *et al.* 2013) in R, using
the "Nm" statistic to estimate number of migrants, and 1000 bootstrap samples when
calculating significance.
200
To identify patterns of admixture in the populations with each analysis, STRUCTURE v.2.3.4
(Pritchard *et al.* 2000) was used. In each analysis, we included 100,000 iterations for burn-in
and 500,000 iterations of Markov-chain Monte Carlo (MCMC), and for each K value (1-9)
we used 9-25 independent runs per K depending on the species to limit the number of minor
modes created. By applying the ΔK Evanno method (Evanno *et al.* 2005), the most likely K
value was determined through STRUCTURE HARVESTER (Earl & vonHoldt 2012). The
Cluster Markov Packager Across Ks (CLUMPAK) online tool (Kopelman *et al.* 2015) was
used to summarize the results, and we produced the STRUCTURE graphs in R. The R
package adegenet v1.7-16 (Jombart 2008; Jombart & Ahmed 2011) was used to conduct a
Discriminant Analysis of Principal Components (DAPC). Loci were removed if there were
more than 20% missing values, and individuals were removed if there were more than 25%
missing values. For the remaining data, the missing values were imputed using the mean
allele frequency for each locus calculated by the "tab" function. The "find.clusters" function
was used to find the number of *K* values within each analysis regardless of a priori population
groups sampled by using a combination of Bayesian Information Criterion (BIC) to identify
the most likely K value, as well as a k-means approach to identify genetic clusters. The
"xvalDapc" function was used to cross-validate results and identify the number of
discriminant functions and principal components to retain for DAPC for each species per
analysis.
220
221 **Results**
222
223 Population Structure Among Locations
224
After quality control and SNP calling, 101 individuals from *Gobiodon fuscoruber,* 103 *G.*
rivulatus, 99 *G. quinquestrigatus*, and 75 *G. histrio* were retained with 5751, 6345, 6527, and
10006 loci respectively passing filtering parameters for the analysis among locations. Each
species examined had distinct population structure across the locations where it was
observed. However, the degree of clustering, heterozygosity and direction of migration
asymmetry among locations varied by species.
231
Gobiodon fuscoruber exhibited the most distinct population structure among locations
(AMOVA: $F_{ST} = 0.14$, P < 0.0001, see Suppl. Tabs. for all statistical outputs) compared to all
other species (see results below). There was some shared ancestry between LI (central
population) and PNG (northern population; pairwise $F_{ST} = 0.12$) as well as LI and OTI
(southern population; pairwise $F_{ST} = 0.11$; Fig. 2A&B). However, PNG and OTI were highly
distinct from each other with little shared ancestry (pairwise $F_{ST} = 0.22$). Regardless of
population designation, three clusters were identified ($\Delta K = 3$, 9 STRUCTURE
HARVESTER repetitions per K, no minor modes; Suppl. Fig. 1). The smallest cluster
included all PNG individuals whereas OTI and LI each had individuals with ancestry
dispersed among the three clusters (Fig. 2B). The inbreeding coefficient (F_{IS}) was high at LI,
moderate at OTI, and low at PNG (Table 1). Heterozygosity values for both observed and
expected were between 0.05 and 0.09 and observed (H_o) was only higher than expected (H_e)

at PNG (Table 1). There was asymmetry in the migration rates from PNG to LI (100%) and
PNG to OTI (61%; Fig. 2C), yet there was no asymmetry in migration between LI and OTI.
246

247
248 Figure 2. (A) Bayesian clustering algorithm STRUCTURE v2.3.4 with color representing
249 different clusters following (Pritchard *et al.* 2000) and (B) Principal Components Analysis of
250 population differentiation and (C) migration rates filtered for asymmetries for 3 sampled
251 locations (purple = PNG, orange = LI, green = OTI) of each *Gobiodon* species. PNG = Papua
252 New Guinea, LI = Lizard Island, OTI = One Tree Island. ** means p < 0.0001.
253

254 Table 1. Genetic diversity measures for each population of *Gobiodon* species studied. To
255 investigate inbreeding coefficients, we used the range suggested by (Hartl 1999) in which 0-
256 0.05 is low, 0.05-0.15 is moderate, 0.16-0.24 is high, and ≥0.25 is very high.

Species	Population	Inbreeding Coefficient F_{IS}	Observed Heterozygosity H_o	Expected Heterozygosity H_e
G. fuscoruber	Papua New Guinea	0.048	0.06	0.05
	Lizard Island	0.174	0.07	0.09
	One Tree Island	0.106	0.06	0.07
G. histrio	Papua New Guinea	0.125	0.08	0.09
	Lizard Island	0.134	0.09	0.11
	One Tree Island	NA	NA	NA
G. quinquestrigatus	Papua New Guinea	0.195	0.06	0.07
	Lizard Island	0.156	0.05	0.06
	One Tree Island	0.088	0.06	0.06
G. rivulatus	Papua New Guinea	0.137	0.07	0.08
	Lizard Island	0.128	0.06	0.07
	One Tree Island	0.061	0.04	0.04

We found *Gobiodon histrio* at only two of the three sampling locations (PNG and LI). *G.*
histrio had distinct population structure among the two locations (AMOVA: F_{ST} = 0.03, P <
0.0001), yet it had the least structure compared to other species. There were several
individuals with shared ancestry among locations (Fig. 2A&B). Regardless of population
designation, three clusters were identified depending on the analysis used (ΔK = 3, 25
STRUCTURE HARVESTER repetitions per K, no minor modes; Suppl. Fig. 1). However, no
cluster fully encapsulated the population from a single location (Fig. 2B). The F_{IS} was
moderate for both locations and slightly higher at LI (Table 1). Heterozygosity values for
both H_o and H_e were between 0.08 and 0.11 with H_o being lower than H_e at both locations
(Table 1). There was no asymmetry in the migration rates among locations (Fig. 2C).
268
Gobiodon quinquestrigatus had distinct population structure among the three locations
(AMOVA: F_{ST} = 0.05, P < 0.0001), albeit with mixed ancestry present in all locations
(pairwise F_{ST} values: 0.06 for PNG-LI, 0.03 for LI-OTI, 0.05 for PNG-OTI; Fig. 2A&B).
Regardless of population designation, three clusters were identified (ΔK = 3, 25
STRUCTURE HARVESTER repetitions per K, 2 minor modes; Suppl. Fig. 1&2). However,
no cluster fully encapsulated the population from a single location (Fig. 2B). The F_{IS} was
high at PNG, high at LI, and moderate at OTI (Table 1). Heterozygosity values for both H_o
and H_e were between 0.05 and 0.07 and H_o was lower or the same as H_e at all locations (Table
1). There was no asymmetry in the migration rates among locations (Fig. 2C).
278
Gobiodon rivulatus had distinct population structure among the three locations (AMOVA: F_{ST}
= 0.09, P < 0.0001). There was shared ancestry between LI and OTI (pairwise F_{ST} = 0.04) and
some between LI and PNG (pairwise F_{ST} = 0.10), but less between PNG and OTI (pairwise
F_{ST} = 0.13; Fig. 2A&B). Regardless of population designation, three clusters were identified
(ΔK = 3, 25 STRUCTURE HARVESTER repetitions per K, 2 minor modes; Suppl. Fig.
1&3). A single cluster encapsulated the PNG population and some individuals from LI and
OTI, which were both dispersed among several clusters (Fig. 2B). The F_{IS} was moderate at all
sites although it was two times higher at PNG and LI than OTI (Table 1). Heterozygosity
values for both H_o and H_e were between 0.05 and 0.08 and H_o was lower or the same as H_e at
all locations (Table 1). There is asymmetry in the migration rates from OTI to PNG (42%)
and OTI to LI (100%, Fig. 2C), but no asymmetry between PNG and LI.
290
291 Pre- versus Post-Disturbances at Lizard Island
292
Gobiodon fuscoruber and *G. rivulatus* were locally extirpated from Lizard Island following
the climatic disturbances. For the pre- and post-disturbances subsets from Lizard Island, we
retained 62 *G. brochus,* 70 *G. histrio*, 59 *G. quinquestrigatus*, and 62 *G. erythrospilus*
individuals with 9724, 9391, 7656, and 10760 loci, respectively. Two out of the four species
had distinct structure between pre- and post-disturbances and the degree of heterozygosity
varied among species.
299
Gobiodon brochus had no distinct structure pre- versus post-disturbances (AMOVA: F_{ST} =
0.0007, P = 0.87). Regardless of population designation, four clusters were identified (ΔK =
4, 10 STRUCTURE HARVESTER repetitions per K, 0 minor modes; Suppl. Fig. 4). There
was a loss of a single cluster made up of 2 individuals from pre- to post-disturbances (Fig.
3A&B). The F_{IS} was moderate at both time points (Table 2). Heterozygosity values for both
H_o and H_e were between 0.14 and 0.17 and H_o was lower than H_e at both time points (Table
2). There was no asymmetry in the migration rates among sampling periods (Fig. 3C).
307

308
309 Figure 3. (A) Bayesian clustering algorithm STRUCTURE v2.3.4 with color representing
310 different clusters (Pritchard *et al.* 2000) and (B) Principal Components Analysis showing
311 population differentiation and (C) asymmetric gene flow migration for pre- (PRE = orange)
312 and post-disturbances (POST = dark red) at Lizard Island of each *Gobiodon* species. **
313 means p < 0.0001, * means p < 0.02.
314
315 Table 2. Genetic diversity measures for each *Gobiodon* species studied pre- and post-
316 disturbances at Lizard Island, northern Great Barrier Reef.

Species	Population	Inbreeding Coefficient F_{is}	Observed Heterozygosity H_o	Expected Heterozygosity H_e
G. brochus	Pre-disturbances	0.134	0.14	0.17
	Post-disturbances	0.127	0.14	0.17
G. erythrospilus	Pre-disturbances	0.130	0.13	0.16
	Post-disturbances	0.131	0.13	0.15
G. fuscoruber	Pre-disturbances	0.174	0.07	0.09
	Post-disturbances	extirpated	extirpated	extirpated
G. histrio	Pre-disturbances	0.075	0.12	0.13
	Post-disturbances	0.108	0.08	0.10

G. quinquestrigatus	Pre-disturbances	0.183	0.08	0.10
	Post-disturbances	0.262	0.06	0.10
G. rivulatus	Pre-disturbances	0.128	0.06	0.07
	Post-disturbances	extirpated	extirpated	extirpated

Gobiodon erythrospilus had barely significant population structure pre- and post-disturbances (AMOVA: F_{ST} = 0.005, P = 0.018). However, only one cluster was identified regardless of population designation (Suppl. Fig. 4). There were two clusters that disappeared from pre- to post-disturbances, although those clusters had only been found in two individuals pre-disturbances. The rest of the individuals exhibited the same ancestry at both time periods (Fig. 3A&B). The F_{IS} was moderate at both time points (Table 2). Heterozygosity values for both H_o and H_e were between 0.13 and 0.16 and H_o was lower than H_e at both time points (Table 2). There was no asymmetry in the migration rates between either time points (Fig. 3C).

Gobiodon histrio had distinct population structure pre- and post-disturbances (AMOVA: F_{ST} = 0.08, P < 0.0001). The population structure homogenized post-disturbances, with the loss of individuals with partial ancestry from another population (Fig. 3A&B). Regardless of population designation, three clusters were identified (⍰K = 3, 10 STRUCTURE HARVESTER repetitions per K, 0 minor modes; Suppl. Fig. 4). The F_{IS} was moderate at both time points but was 44% higher post-disturbances than pre-disturbances (Table 2). Heterozygosity values for both H_o and H_e were between 0.08 and 0.13 and H_o was lower than H_e at both time points (Table 2). Although there was asymmetric migration identified from post-disturbances to pre-disturbances, the context instead implies the opposite direction (Sundqvist *et al.* 2016) with a founder effect due to the disturbances (Fig. 3C).

Gobiodon quinquestrigatus had distinct population structure pre- and post-disturbances (AMOVA: F_{ST} = 0.08, P < 0.0001). This species similarly homogenized post-disturbance, with almost all individuals exhibiting full ancestry from only 2 populations (Fig. 3A&B). Regardless of population designation, four clusters were identified (⍰K = 4, 10 STRUCTURE HARVESTER repetitions per K, 0 minor modes; Suppl. Fig. 4). Individuals pre-disturbances were found in each of the four clusters whereas one cluster was lost post-disturbances. Most individuals post-disturbances were from one of two clusters (dark blue or cian) with a select few exhibiting also third cluster (red with dark blue) (Fig. 3B). The F_{IS} was high pre-disturbances and 43% higher post-disturbances sitting at a very high value (0.26), which was the highest in the whole study (Table 2). Heterozygosity values for both H_o and H_e were between 0.06 and 0.10 and H_o was lower than H_e at both time points (Table 2). Although there was asymmetric migration identified from post-disturbances to pre-disturbances, the context instead implies the opposite direction (Sundqvist *et al.* 2016) with a founder effect due to the disturbances (Fig. 3C).

Discussion

We used a genomic approach to quantify population structure and the potential for genetic bottlenecks resulting from the severe declines in *Gobiodon* populations and changes in their social structures and host-use patterns following repeated cyclones and heatwave-induced bleaching events (Froehlich *et al.* 2021, 2023, 2024). While all species showed some degree of population structure across the study sites, they differed in genetic diversity and directional gene flow. For example, *G. fuscoruber* exhibited migration patterns from north to south, the opposite was found for *G. rivulatus*, and *G. histrio* and *G. quinquestrigatus* had no clear pattern. Following disturbances at the central location, *G. fuscoruber* and *G. rivulatus* were extirpated. We detected genetic bottlenecks in *G. quinquestrigatus* and *G. histrio*. Our study

365   highlights that population structure and genetic bottlenecks increases the vulnerability of
366   these fishes to population collapse during climatic disturbances, alongside other ecological
367   losses observed for theses fishes. Given these results, we recommend expanding conservation
368   efforts to include more *Gobiodon* species, which are facing multiple climate-driven threats.
369
Population structure was evident for each *Gobiodon* species across all locations in relatively
healthy states, suggesting these populations may be especially vulnerable to climatic
disturbances. Most species exhibited low to moderate levels of genetic isolation, as indicated
by F_{ST} values (0.03–0.13), while *G. fuscoruber* showed moderate to high F_{ST} values (0.11–
0.22) (Hall 2022). Although the distances between study sites (1000–2000 km) are typically
below thresholds for genetic isolation at sea (Cooke *et al.* 2016), *G. fuscoruber*'s minimum
F_{ST} of 0.15 (Frankham *et al.* 2002) indicates that its populations are indeed genetically
isolated. Such populations may be either recruitment-limited or post-recruitment regulated
(Jones *et al.* 2009). High self-recruitment rates are observed in many reef fishes with short
larval durations (D'Aloia *et al.* 2018; Jones *et al.* 1999, 2005; Saenz-Agudelo *et al.* 2012;
Selwyn *et al.* 2016; Swearer *et al.* 2002), like the coral-dwelling goby *Paragobiodon*
xanthosoma (Rueger *et al.* 2021b). For *Gobiodon*, the pelagic larval duration averages around
19.6 days (Pereira *et al.* 2015), which suggests that they are regulated by post-recruitment
and are not recruitment-limited under normal conditions, which reduces the risks of
extirpation but increases recovery time (Jones *et al.* 2009). Nonetheless, two species (*G.*
fuscoruber and *G. rivulatus*) experienced local extirpation in the disturbed location.
Following these severe events, which dramatically reduced adult populations, *Gobiodon* may
have become recruitment-limited as well. Inbreeding levels further highlight species-specific
vulnerabilities. Inbreeding ranged from low to high (Hartl 1999) at different locations
depending on the species (e.g. *G. fuscoruber, G. quinquestrigatus*), or moderate throughout
(*G. rivulatus*). At the central location (Lizard Island, inbreeding was moderate in *G. histrio*
and *G. rivulatus* and high in *G. fuscoruber* and *G. quinquestrigatus*. Such inbreeding patterns
are unsurprising since *Gobiodon*, like other habitat specialists, tend to settle near natal reefs
with siblings (Buston *et al.* 2009; Jones *et al.* 2005). However, limited genetic diversity may
occur from such settlement behavior, therefore increasing their susceptible to extreme
disturbances (Jones *et al.* 2009). Accordingly, low to high population structure coupled with
moderate to high inbreeding patterns for all species likely contribute to their slow recovery
rates.
398
Ocean currents play crucial roles in larval dispersal and the recovery potential of marine
species (Gouvêa *et al.* 2023; Snead *et al.* 2023; Wilson *et al.* 2016), but they do not fully
explain the patterns of directional genetic migration in *Gobiodon* species. Based on
prevailing ocean currents in western Oceania (Ridgway & Hill 2009), the central populations
in our study (Lizard Island) would be expected to act as a source for the northern populations
(Papua New Guinea). Additionally, the presence of competing currents between the southern
and central populations (East Australia and Hirri Currents; Ridgway and Hill 2009) further
complicates the predictions of source-sink dynamics between these locations. Of the four
species studied, only two species (*G. fuscoruber* and *G. rivulatus*) exhibited directional
genetic migration, and both were locally extirpated from the disturbed location. Interestingly,
their migration patterns were in opposite directions: *G. fuscoruber* showed directional genetic
migration from the northern population to the central and southern populations, whereas *G.*
rivulatus exhibited the opposite. It is possible that we did not sample intermediate
populations found between the study locations, which may help explain the contrasting
patterns among species. However, ocean currents alone may not explain the overall pattern of
gene flow in all *Gobiodon* species, especially given that the central population would be
expected to have sourced both the northern and southern locations. However, ocean currents
are affected by climate change, which may alter patterns of larval dispersal (Wilson *et al.*

2016). Although the dispersal and genetic structure of many marine fishes is often driven by
ocean currents (Baltazar-Soares *et al.* 2014; Munguia-Vega *et al.* 2017; Snead *et al.* 2023),
not all tropical species exhibit such patterns (Shulman & Bermingham 1995). Since two
Gobiodon species exhibited contrasting patterns despite their overlapping ranges, their
genetic migration is likely influenced by other ecological factors, like their sociality
(Froehlich *et al.* 2024; Rueger *et al.* 2021a) and the distribution of their preferred hosts
(Froehlich *et al.* 2023). Overall, our findings indicate that ocean currents do not always
influence population structure of all species within a single genus, which has flow-on
implications for their resilience to climatic disturbances.
426
When investigating whether climatic disturbances at the central location affected the genetic
diversity, not all species responded in the same manner. There were few changes to the
population structure of *Gobiodon brochus* and *G. erythrospilus* following the disturbances.
The latter species had little genetic diversity at both time points, which aligns with its
vulnerable status (Larson 2020). Other aquatic taxa that underwent mass mortality following
storms and heatwaves had little changes to their genetic diversity, such as the submerged
aquatic plant *Vallisneria americana* and the red gorgonian *Paramuricea clavata* (Ngeve *et al.*
2023; Pilczynska *et al.* 2016). For the alpine butterfly *Parnassius smintheus*, high
connectivity pre-disturbances resulted in little change to their genetic diversity after mass
mortality (Jangjoo *et al.* 2016). However, not all species are resilient. Two species in our
study, *G. fuscoruber* and *G. rivulatus*, were extirpated. The other two species, *G. histrio* and
G. quinquestrigatus, showed a clear loss of genetic diversity, increased inbreeding, and
marked directional genetic migration post-disturbance. In both species, post-disturbance
populations were more genetically uniform (i.e. less diverse), and at least one ancestral
cluster that was initially present pre-disturbances disappeared entirely. There were
characteristics similar to a founder effects (Sundqvist *et al.* 2016) whereby a few individuals
pre-disturbances gave rise to the post-disturbance populations. This was evident from the
reduction in the number of ancestral clusters, the increase in assignment of individuals to a
single cluster, and the directional genetic migration observed. Broader patterns in marine
ecosystems show similar results in which population structure leads to lower genetic diversity
post-disturbances and the potential for genetic bottlenecks (Noreen *et al.* 2009; Selkoe *et al.*
2016). Similarly, seaweed populations underwent up to 80-100% loss in genetic diversity
following marine heatwaves (Gurgel *et al.* 2020). Taken together, our results suggest that
genetic bottlenecks pose a real threat to several *Gobiodon* species. Conservation strategies
aimed at protecting coral reef ecosystems should include coral-dwelling fishes and
incorporate species-specific management plans at a population level. As climatic disturbances
continue to increase in frequency and severity, genetic bottlenecks may become even more
common, placing many more reef-associated species at risk.
455
When we look at the genetic, life history, and ecological outcomes of climate change on
Gobiodon species, a clearer picture emerges as to why some species recover more slowly
than others. For example, the two species that were extirpated post-disturbances (*G.*
fuscoruber and *G. rivulatus*) had some of the highest F_{ST} values among populations. Pre-
disturbances, both species lived primarily in groups rather than pairs, unlike other goby
species in this study (Froehlich *et al.* 2024; Hing *et al.* 2019). Post-disturbances, the absence
of established social groups for *G. fuscoruber* and *G. rivulatus* likely reduced successful
settlement, compounded by high inbreeding rates and population structure. All *Gobiodon*
species—regardless of whether they typically lived in pairs or groups before the
disturbances—were mostly found living alone or in pairs post-disturbances (Froehlich *et al.*
2024), likely reducing their reproductive success. Additionally, a species' growth potential
also shapes resilience. For example, *G. rivulatus* is one of the smallest species in the genus
(Hing *et al.* 2018, 2019), making it more vulnerable to predation, especially since smaller

individuals are more frequently targeted post-disturbance (Brandl *et al.* 2018; Goatley &
Bellwood 2016). Since *Gobiodon* species fully depend on live coral, how well their preferred
coral hosts recover can directly affect their own recovery. *Gobiodon* species that were
extirpated post-disturbances (*G. fuscoruber* and *G. rivulatus*) primarily inhabited corals that
became rare post-disturbances (*Acropora millepora* and *A. gemmifera*, respectively)
(Froehlich *et al.* 2021, 2023). However, *Gobiodon* species that persisted either were able to
change the coral host they inhabited or benefited from their preferred host recovering quickly
(Froehlich *et al.* 2023). Yet even species that survived were slower to recover than their coral
hosts and several exhibited genetic bottlenecks, which further reduces their ability to respond
to future disturbances. These compounding pressures highlight the urgent need for
conservation efforts that specifically target coral-dwelling gobies as they play important roles
for coral health and resilience to bleaching.
481

482   Animal Ethics and Permits
483   The study was approved under animal ethics protocols AE1404 and AE1725 at the University
484   of Wollongong. This research was conducted under research permits issued by the Great
485   Barrier Reef Marine Park Authority (G13/36197.1, G15/37533.1, and G18/41020.1) and
486   Papua New Guinea Research Visa Permit AA654347.
487
Funding
Field and laboratory work were funded by the Hermon Slade Foundation to MW and M.
Dowton, Sea World Research and Rescue Foundation to MW and T. Rueger, a target grant
from the Faculty of Science, Medicine and Health at the University of Wollongong (UOW) to
MW, funding initiatives from the Centre for Sustainable Ecosystem Solutions to MW, CF and
SH, the Zoltan Florian Marine Biology Fellowship as part of the Lizard Island Doctoral
Fellowship program of the Australian Museum to CF, and an Australian Coral Reef Society
Fellowship to CF. For student support, CF was supported under a University Postgraduate
Award, RK and SH by an Australian Government Research Training Program Scholarship,
both administered through UOW. SG is supported by the National Science Foundation
Graduate Research Fellowship Program under Grant No. DGE-2035702.
499
Acknowledgments
We pay respect to the traditional owners past and present of the Kilu and Tamare
communities of Kimbe Bay in Papua New Guinea, of the Dingaal Aboriginal Traditional
Owners of Lizard Island in Australia, and Gurang, Gooreng Gooreng, Bailai and Bunda
Traditional Owners of One Tree Island in Australia. We greatly appreciate our field helpers:
Kylie Brown, Karen Hing, Theresa Rueger, Rituraj Sharma, Jemma Okapie Smith, Nelson
and Jerry Sikatura. We are thankful to Mark Dowton and Tracey Gibson for guidance
regarding laboratory practices. We are grateful to the facilities and management of field
stations at the Mahonia Na Dari Research and Conservation Centre of Kimbe Bay, Lizard
Island Research Station, and One Tree Island Research Station, with particular appreciation to
Ainsley and Paul Carin, Anne Hoggett, Somei Jonda, Peter and Jane Miller, and Lyle Vail for
their assistance with our field trips. We would like to thank Courtney Hildebrandt for her
Gobiodon artwork and staff at the Sydney Institute of Marine Science for help while carrying
out laboratory work in the facility.

514
515

References

AghaKouchak, A., Huning, L.S., Chiang, F., Sadegh, M., Vahedifard, F., Mazdiyasni, O., *et al.* (2018). How do natural hazards cascade to cause disasters? *Nature*, 561, 458–460.

Attrill, M.J. & Depledge, M.H. (1997). Community and population indicators of ecosystem health: targeting links between levels of biological organisation. *Aquat. Toxicol.*, 38, 183–197.

Baltazar-Soares, M., Biastoch, A., Harrod, C., Hanel, R., Marohn, L., Prigge, E., *et al.* (2014). Recruitment Collapse and Population Structure of the European Eel Shaped by Local Ocean Current Dynamics. *Curr. Biol.*, 24, 104–108.

Baptista, M.S.P., Keuroghlian, A., Tambosi, L.R., Côrtes, M.C., de Góes Maciel, F., Cirino, D.W., *et al.* (2025). Agriculturally developed areas reduce genetic connectivity for a keystone Neotropical ungulate. *Biotropica*, 57, e13418.

Bellwood, D.R., Baird, A.H., Depczynski, M., González-Cabello, A., Hoey, A.S., Lefèvre, C.D., *et al.* (2012). Coral recovery may not herald the return of fishes on damaged coral reefs. *Oecologia*, 170, 567–573.

Bergstrom, D.M., Wienecke, B.C., Hoff, J., Hughes, L., Lindenmayer, D.B., Ainsworth, T.D., *et al.* (2021). Combating ecosystem collapse from the tropics to the Antarctic. *Glob. Change Biol.*, gcb.15539.

Borges, F.L.G., Oliveira, M.D.R., De Almeida, T.C., Majer, J.D. & Garcia, L.C. (2021). Terrestrial invertebrates as bioindicators in restoration ecology: A global bibliometric survey. *Ecol. Indic.*, 125, 107458.

Brandl, S.J., Goatley, C.H.R., Bellwood, D.R. & Tornabene, L. (2018). The hidden half: ecology and evolution of cryptobenthic fishes on coral reefs. *Biol. Rev.*, 93, 1846–1873.

Buston, P.M., Fauvelot, C., Wong, M.Y.L. & Planes, S. (2009). Genetic relatedness in groups of the humbug damselfish Dascyllus aruanus: small, similar-sized individuals may be close kin. *Mol. Ecol.*, 18, 4707–4715.

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A. & Cresko, W.A. (2013). Stacks: an analysis tool set for population genomics. *Mol. Ecol.*, 22, 3124–3140.

Catchen, J.M., Amores, A., Hohenlohe, P., Cresko, W. & Postlethwait, J.H. (2011). Stacks: Building and Genotyping Loci De Novo From Short-Read Sequences. *G3 GenesGenomesGenetics*, 1, 171–182.

Cooke, G.M., Schlub, T.E., Sherwin, W.B. & Ord, T.J. (2016). Understanding the spatial scale of genetic connectivity at sea: unique insights from a land fish and a meta-analysis. *PLoS ONE*, 11, e0150991.

D'Aloia, C.C., Xuereb, A., Fortin, M.-J., Bogdanowicz, S.M. & Buston, P.M. (2018). Limited dispersal explains the spatial distribution of siblings in a reef fish population. *Mar. Ecol. Prog. Ser.*, 607, 143–154.

Earl, D.A. & vonHoldt, B.M. (2012). STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv. Genet. Resour.*, 4, 359–361.

Evanno, G., Regnaut, S. & Goudet, J. (2005). Detecting the number of clusters of individuals using the software structure: a simulation study. *Mol. Ecol.*, 14, 2611–2620.

Excoffier, L. & Lischer, H.E.L. (2010). Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.*, 10, 564–567.

Frankham, R., Ballou, J.D., Briscoe, D.A. & McInnes, K.H. (2002). *Introduction to Conservation Genetics*. Cambridge University Press, Cambridge, UNITED KINGDOM.

Froehlich, C.Y.M., Heatwole, S.J., Klanten, O.S. & Wong, M.Y.L. (2022). Habitat health, size
and saturation do not alter movement decisions in a social coral reef fish. *Anim.*
Behav., 191, 125–133.
Froehlich, C.Y.M., Heatwole, S.J., Selma Klanten, O., Hing, M.L., Hildebrandt, C.A., Smith,
J.O., *et al.* (2024). Multi-level framework to assess social variation in response to
ecological and social factors: modeled with coral gobies. *Oikos*, e10669.
Froehlich, C.Y.M., Klanten, O.S., Hing, M.L., Dowton, M. & Wong, M.Y.L. (2021). Uneven
declines between corals and cryptobenthic fish symbionts from multiple disturbances.
Sci. Rep., 11, 16420.
Froehlich, C.Y.M., Klanten, O.S., Hing, M.L., Dowton, M. & Wong, M.Y.L. (2023). Delayed
recovery and host specialization may spell disaster for coral-fish mutualism. *Ecol.*
Evol., 13, e10209.
Goatley, C.H.R. & Bellwood, D.R. (2016). Body size and mortality rates in coral reef fishes:
a three-phase relationship. *Proc. R. Soc. B Biol. Sci.*, 283, 20161858.
Goudet, J. (2005). hierfstat, a package for r to compute and test hierarchical F-statistics. *Mol.*
Ecol. Notes, 5, 184–186.
Gouvêa, L.P., Fragkopoulou, E., Cavanaugh, K., Serrão, E.A., Araújo, M.B., Costello, M.J.,
et al. (2023). Oceanographic connectivity explains the intra-specific diversity of
mangrove forests at global scales. *Proc. Natl. Acad. Sci.*, 120, e2209637120.
Gurgel, C.F.D., Camacho, O., Minne, A.J.P., Wernberg, T. & Coleman, M.A. (2020). Marine
Heatwave Drives Cryptic Loss of Genetic Diversity in Underwater Forests. *Curr.*
Biol., 30, 1199-1206.e2.
Hall, S.J.G. (2022). Genetic Differentiation among Livestock Breeds—Values for Fst. *Anim.*
Open Access J. MDPI, 12, 1115.
Hartl, D.L. (1999). *A primer of population genetics, 3rd edition*. Sinauer Associates,
Sunderland, MA, USA.
Hill, G.M., Kawahara, A.Y., Daniels, J.C., Bateman, C.C. & Scheffers, B.R. (2021). Climate
change effects on animal ecology: butterflies and moths as a case study. *Biol. Rev.*, 96,
2113–2126.
Hing, M.L., Klanten, O.S., Dowton, M., Brown, K.R. & Wong, M.Y.L. (2018). Repeated
cyclone events reveal potential causes of sociality in coral-dwelling *Gobiodon* fishes.
PLoS ONE, 13, e0202407.
Hing, M.L., Klanten, O.S., Wong, M.Y.L. & Dowton, M. (2019). Drivers of sociality in
Gobiodon fishes: an assessment of phylogeny, ecology and life-history. *Mol.*
Phylogenet. Evol., 137, 263–273.
Hourigan, T.F., Timothy, C.T. & Reese, E.S. (1988). Coral Reef Fishes as Indicators of
Environmental Stress in Coral Reefs. In: *Marine Organisms as Indicators* (eds. Soule,
D.F. & Kleppel, G.S.). Springer, New York, NY, pp. 107–135.
Hughes, T.P., Anderson, K.D., Connolly, S.R., Heron, S.F., Kerry, J.T., Lough, J.M., *et al.*
(2018a). Spatial and temporal patterns of mass bleaching of corals in the
Anthropocene. *Science*, 359, 80–83.
Hughes, T.P., Kerry, J.T., Baird, A.H., Connolly, S.R., Dietzel, A., Eakin, C.M., *et al.* (2018b).
Global warming transforms coral reef assemblages. *Nature*, 556, 492–496.
Jangjoo, M., Matter, S.F., Roland, J. & Keyghobadi, N. (2016). Connectivity rescues genetic
diversity after a demographic bottleneck in a butterfly population network. *Proc. Natl.*
Acad. Sci., 113, 10914–10919.
Johnstone, J.F., Allen, C.D., Franklin, J.F., Frelich, L.E., Harvey, B.J., Higuera, P.E., *et al.*
(2016). Changing disturbance regimes, ecological memory, and forest resilience.
Front. Ecol. Environ., 14, 369–378.
Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinforma. Oxf. Engl., 24, 1403–1405.

Jombart, T. & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide
618   SNP data. *Bioinforma. Oxf. Engl.*, 27, 3070–3071.

619   Jones, G.P., Almany, G.R., Russ, G.R., Sale, P.F., Steneck, R.S., van Oppen, M.J.H., *et al.*
620   (2009). Larval retention and connectivity among populations of corals and reef fishes:
621   history, advances and challenges. *Coral Reefs*, 28, 307–325.

622   Jones, G.P., Milicich, M.J., Emslie, M.J. & Lunow, C. (1999). Self-recruitment in a coral reef
623   fish population. *Nature*, 402, 802–804.

624   Jones, G.P., Planes, S. & Thorrold, S.R. (2005). Coral reef fish larvae settle close to home.
625   *Curr. Biol.*, 15, 1314–1318.

626   Keenan, K., McGinnity, P., Cross, T.F., Crozier, W.W. & Prodöhl, P.A. (2013). diveRsity: An
627   R package for the estimation and exploration of population genetics parameters and
628   their associated errors. *Methods Ecol. Evol.*, 4, 782–788.

629   Kopelman, N.M., Mayzel, J., Jakobsson, M., Rosenberg, N.A. & Mayrose, I. (2015).
630   Clumpak: a program for identifying clustering modes and packaging population
631   structure inferences across K. *Mol. Ecol. Resour.*, 15, 1179–1191.

632   Krishnan, P., George, G., Immanuel, T., Bitopan-Malakar, M. & Anand, A. (2013). Studies on
633   the Recovery of Bleached Corals in Andaman: Fishes as Indicators of Reef Health. In:
634   *Ecology and Conservation of Tropical Marine Faunal Communities* (eds.
635   Venkataraman, K., Sivaperuman, C. & Raghunathan, C.). Springer, Berlin,
636   Heidelberg, pp. 395–407.

637   Larson, H.K. (2020). *Gobiodon erythrospilus. IUCN Red List Threat. Species*,
638   e.T68330883A68333739.

639   Munguia-Vega, A., Marinone, S.G., Paz-Garcia, D.A., Giron-Nava, A., Plomozo-Lugo, T.,
640   Gonzalez-Cuellar, O., *et al.* (2017). Anisotropic larval connectivity and
641   metapopulation structure driven by directional oceanic currents in a marine fish
642   targeted by small-scale fisheries. *Mar. Biol.*, 165, 16.

643   Munguía-Vega, A., Sáenz-Arroyo, A., Greenley, A.P., Espinoza-Montes, J.A., Palumbi, S.R.,
644   Rossetto, M., *et al.* (2015). Marine reserves help preserve genetic diversity after
645   impacts derived from climate variability: Lessons from the pink abalone in Baja
646   California. *Glob. Ecol. Conserv.*, 4, 264–276.

647   Ngeve, M.N., Engelhardt, K.A.M., Gray, M. & Neel, M.C. (2023). Calm after the storm?
648   Similar patterns of genetic variation in a riverine foundation species before and after
649   severe disturbance. *Ecol. Evol.*, 13, e10670.

650   Noreen, A.M.E., Harrison, P.L. & Van Oppen, M.J.H. (2009). Genetic diversity and
651   connectivity in a brooding reef coral at the limit of its distribution. *Proc. R. Soc. B
652   Biol. Sci.*, 276, 3927–3935.

653   Pauls, S.U., Nowak, C., Bálint, M. & Pfenninger, M. (2013). The impact of global climate
654   change on genetic diversity within populations and species. *Mol. Ecol.*, 22, 925–946.

655   Pereira, P.H.C., Munday, P.L. & Jones, G.P. (2015). Competitive mechanisms change with
656   ontogeny in coral-dwelling gobies. *Ecology*, 96, 3090–3101.

657   Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H.S. & Hoekstra, H.E. (2012). Double Digest
658   RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in
659   Model and Non-Model Species. *PLOS ONE*, 7, e37135.

660   Pilczynska, J., Cocito, S., Boavida, J., Serrão, E. & Queiroga, H. (2016). Genetic Diversity
661   and Local Connectivity in the Mediterranean Red Gorgonian Coral after Mass
662   Mortality Events. *PLOS ONE*, 11, e0150590.

663   Pinna, M., Zangaro, F., Saccomanno, B., Scalone, C., Bozzeda, F., Fanini, L., *et al.* (2023).
664   An Overview of Ecological Indicators of Fish to Evaluate the Anthropogenic
665   Pressures in Aquatic Ecosystems: From Traditional to Innovative DNA-Based
666   Approaches. *Water*, 15, 949.

667   Pinsky, M.L., Comte, L. & Sax, D.F. (2022). Unifying climate change biology across realms
668   and taxa. *Trends Ecol. Evol.*, 37, 672–682.

Pritchard, J.K., Stephens, M. & Donnelly, P. (2000). Inference of Population Structure Using
Multilocus Genotype Data. *Genetics*, 155, 945–959.
R Core Team. (2020). R: a language and environment for statistical computing. *R Found.*
Stat. Comput., Vienna, Austria. https://www.R-project.org.
Richardson, L.E., Graham, N.A.J., Pratchett, M.S., Eurich, J.G. & Hoey, A.S. (2018). Mass
coral bleaching causes biotic homogenization of reef fish assemblages. *Glob. Change*
Biol., 24, 3117–3129.
Ridgway, K. & Hill, K. (2009). The East Australian Current (EAC). *NCCARF Publ.*, 1–16.
Rochette, N.C. & Catchen, J.M. (2017). Deriving genotypes from RAD-seq short-read data
using Stacks. *Nat. Protoc.*, 12, 2640–2659.
Rochette, N.C., Rivera-Colón, A.G. & Catchen, J.M. (2019). Stacks 2: Analytical methods for
paired-end sequencing improve RADseq-based population genomics. *Mol. Ecol.*, 28,
4737–4754.
RStudio Team. (2020). RStudio: integrated development environment for R. *RStudio PBC*,
Boston, MA. http://www.rstudio.com/.
Rueger, T., Branconi, R., Froehlich, C.Y.M., Heatwole, S.J., Wong, M.Y.L. & Buston, P.M.
(2021a). The next frontier in understanding the evolution of coral reef fish societies.
Front. Mar. Sci., 8, 665780.
Rueger, T., Buston, P.M., Bogdanowicz, S.M. & Wong, M.Y. (2021b). Genetic relatedness in
social groups of the emerald coral goby *Paragobiodon xanthosoma* creates potential
for weak kin selection. *Mol. Ecol.*, 30, 1311–1321.
Saenz-Agudelo, P., Jones, G.P., Thorrold, S.R. & Planes, S. (2012). Patterns and persistence
of larval retention and connectivity in a marine fish metapopulation. *Mol. Ecol.*, 21,
4695–4705.
Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., *et al.*
(2017). Forest disturbances under climate change. *Nat. Clim. Change*, 7, 395–402.
Selkoe, K.A., Gaggiotti, O.E., Treml, E.A., Wren, J.L.K., Donovan, M.K. & Toonen, R.J.
(2016). The DNA of coral reef biodiversity: predicting and protecting genetic
diversity of reef assemblages. *Proc. R. Soc. B Biol. Sci.*, 283, 20160354.
Selwyn, J.D., Hogan, J.D., Downey-Wall, A.M., Gurski, L.M., Portnoy, D.S. & Heath, D.D.
(2016). Kin-aggregations explain chaotic genetic patchiness, a commonly observed
genetic pattern, in a marine fish. *PLoS ONE*, 11, e0153381.
Sergio, F., Blas, J. & Hiraldo, F. (2018). Animal responses to natural disturbance and climate
extremes: a review. *Glob. Planet. Change*, 161, 28–40.
Shulman, M.J. & Bermingham, E. (1995). Early Life Histories, Ocean Currents, and the
Population Genetics of Caribbean Reef Fishes. *Evolution*, 49, 897–910.
Snead, A.A., Tatarenkov, A., Avise, J.C., Taylor, D.S., Turner, B.J., Marson, K., *et al.* (2023).
Out to sea: ocean currents and patterns of asymmetric gene flow in an intertidal fish
species. *Front. Genet.*, 14.
Sundqvist, L., Keenan, K., Zackrisson, M., Prodöhl, P. & Kleinhans, D. (2016). Directional
genetic differentiation and relative migration. *Ecol. Evol.*, 6, 3461–3475.
Swearer, S.E., Shima, J.S., Hellberg, M.E., Thorrold, S.R., Jones, G.P., Robertson, D.R., *et al.*
(2002). Evidence of self-recruitment in demersal marine populations. *Bull. Mar. Sci.*
Teng, Y., Li, W., Wang, X., Zhang, R., Shen, Y., Wu, R., *et al.* (2025). Integrating population
genomics and environmental data to predict adaptation to climate change in post-
bottleneck Tibetan macaques. *Sci. Adv.*, 11, eadw0562.
Thakur, M.P., Risch, A.C. & Putten, W.H. van der. (2022). Biotic responses to climate
extremes in terrestrial ecosystems. *iScience*, 25.
Turner, M.G. (2010). Disturbance and landscape dynamics in a changing world. *Ecology*, 91,
2833–2849.

Wilson, L.J., Fulton, C.J., Hogg, A.M., Joyce, K.E., Radford, B.T.M. & Fraser, C.I. (2016).
Climate-driven changes to ocean circulation and their inferred impacts on marine
dispersal patterns. *Glob. Ecol. Biogeogr.*, 25, 923–939.
Wong, M.Y.L. & Buston, P.M. (2013). Social systems in habitat-specialist reef fishes: key
concepts in evolutionary ecology. *BioScience*, 63, 453–463.
Woodall, L.C., Koldewey, H.J. & Shaw, P.W. (2011). Historical and contemporary population
genetic connectivity of the European short-snouted seahorse Hippocampus
hippocampus and implications for management. *J. Fish Biol.*, 78, 1738–1756.
727