The chromatin remodeller CHD4 regulates transcription factor binding to both prevent activation of silent enhancers and maintain active regulatory elements

Andria Koulle[1,2,4], Oluwaseun Ogundele[1,2], Devina Shah[1,2,5], India-May Baker[1], Maya Lopez[1], Nicola Reynolds[1,3], Ramy Ragheb[1,3], Ernest Laue[1,2] and Brian Hendrich[1,2,3]

1. Cambridge Stem Cell Institute, Jeffrey Cheah Biomedical Centre, University of Cambridge, Cambridge CB2 0AW, UK
2. Department of Biochemistry, University of Cambridge, Cambridge CB2 1GA, UK
3. Living Systems Institute, University of Exeter, Exeter EX4 4QD, UK

4. Current Address: Department of Life Sciences, Imperial College London, London, SW7 2AZ, UK
5. Current Address: University of Copenhagen, Novo Nordisk Foundation Center for Protein Research, Blegdamsvej 3B, 2200 Copenhagen, Denmark

All correspondence to Brian Hendrich: B.D.Hendrich@exeter.ac.uk

Abstract

Chromatin organisation and transcriptional regulation are tightly coordinated processes that are essential for maintaining cellular identity and function. ATP-dependent chromatin remodelling proteins play critical roles in control of genome structure and in regulating transcription across eukaryotes. Their essential nature, however, has made it difficult to define exactly how these functions are mediated. The chromatin remodeller CHD4 has been shown to be capable of sliding nucleosomes in vitro, and to regulate chromatin accessibility and gene expression in vivo. Using an inducible depletion system, here we identify a second mechanism of action for CHD4 in actively restricting the residence time of transcription factors on chromatin. Together these activities result in distinct, context-dependent outcomes: at highly accessible regulatory elements, CHD4 limits transcription factor binding to maintain regulatory function, while at low-accessibility euchromatic regions, it prevents transcription factor engagement and sustains chromatin compaction, thereby silencing cryptic enhancers. Collectively, these mechanisms enable CHD4 to reduce transcriptional noise while preserving the responsiveness of active regulatory networks.

Introduction

Cell state transitions are driven by the activity of transcription factors. Many transcription factors bind to specific DNA motifs within chromatin, but the frequency of these motifs within mammalian genomes far outnumbers sites at which protein binding is detectable. The ability of a transcription factor to recognise its cognate motif is influenced by how the DNA encoding that motif is packaged in chromatin. Accessible sites, i.e. those associated with a low density of intact nucleosomes, are more likely to be identified and bound by transcription factors than are inaccessible sites, i.e. those associated with higher nucleosome density. Controlling chromatin accessibility is therefore crucial for controlling transcription factor binding to cognate sites in regulatory regions. Transcription factor binding patterns will then determine which gene regulatory regions are used to drive gene expression and thereby define cell identity.

Vertebrate cells contain multiple proteins capable of using energy derived from ATP hydrolysis to remodel nucleosomes. These chromatin remodellers share a conserved sucrose non-fermentable 2 (SNF2) helicase-like ATPase domain but otherwise have various additional functional domains which impact how they organise chromatin, transcription and DNA repair (Hota and Bruneau, 2016; Narlikar et al., 2013). Chromatin remodellers play important roles

in mammalian development, and heterozygous mutations in the genes encoding them

underlie a variety of developmental disorders in humans (Gourisankar et al., 2024; Hota and

Bruneau, 2016; Pierson et al., 2019). Similarly, somatic mutations in chromatin remodeller

subunit genes are increasingly being implicated in cancer initiation and/or progression (Lai

and Wade, 2011; Martincorena et al., 2017; Wilson and Roberts, 2011).

CHD4 is an abundant ATP-dependent chromatin remodelling protein which plays

important roles in chromatin organisation and cell fate decisions in many different aspects of

metazoan development. Depletion of CHD4 in mouse or Drosophila cells, or overexpression

of a dominant negative form of the protein, leads to increased chromatin accessibility at

regulatory elements and DNase hypersensitive sites (Dieuleveult et al., 2016; Morris et al.,

2014; Moshkin et al., 2012), indicating that it predominantly functions to compact chromatin.

Its impact on gene expression is less clear-cut, however. One study found that while CHD4

acted to repress genes with bivalent promoters, its activity at active promoters (marked with

H3K4Me3) mainly facilitated transcription in mouse embryonic stem cells (Dieuleveult et al.,

2016). Another study found CHD4 controlled the probability of gene expression, rather than

levels, during the first cell fate transition in mammalian embryogenesis (O'Shaughnessy-

Kirwan et al., 2015). Several studies in somatic lineages have shown that CHD4 prevents

lineage inappropriate gene expression during cell fate decisions in both mice and Drosophila

(Arends et al., 2019; Aughey et al., 2023; Gómez-del Arco et al., 2016; Sreenivasan et al.,

2021; Wilczewski et al., 2018; Yoshida et al., 2019). Exactly how CHD4 has this varied impact

on chromatin and transcription has not been defined.

CHD4 is the predominant chromatin remodelling subunit of the Nucleosome Remodelling

and Deacetylation (NuRD) complex (Wade et al., 1998; Xue et al., 1998; Zhang et al., 1998) .

NuRD is a highly abundant chromatin remodeller present at active enhancers and promoters

in many cell types. Its activity is important not only to regulate transcription but also the

movement of enhancers in 3D space (Basu et al., 2023) , and to maintain genome integrity

(Polo et al., 2010; Smeenk et al., 2010) . NuRD has been shown to exert two functions in

mammalian cells: one is to control the nucleosome density at active enhancers, thereby

regulating transcription factor binding, enhancer activity and transcriptional output (Bornelöv

et al., 2018; Pundhir et al., 2023), while the other is to prevent low-level, inappropriate

transcription across the genome (Burgold et al., 2019; Montibus et al., 2023; Ragheb et al.,

2020; Saotome et al., 2024).

Although mouse ES cells can survive with a complete loss of the histone deacetylase

subunit of NuRD (Burgold et al., 2019), loss of the chromatin remodelling component, CHD4,

leads to cell death (Stevens et al., 2017). While we have used genetics to create NuRD-low or

NuRD-null ES cells in the past to define NuRD function, in both cases, the remodelling

subcomplex (CHD4, GATAD2A/B, CDK2AP1) remained on chromatin and the extent to which it

could continue to remodel chromatin is not known (Bornelöv et al., 2018). As with any genetic

change, it is also very difficult to know to what extent constitutive loss of protein activity has

resulted in selection of cells using some compensatory mechanism to remain viable. CHD4 is

also known to function outside of the NuRD complex (O'Shaughnessy-Kirwan et al., 2015;

Ostapcuk et al., 2018; Williams et al., 2004) . Assessing the direct function of NuRD's

remodelling activity is therefore difficult as knockout of the remodeller results in a non-viable

cell, while knockdown or exogenous overexpression of a dominant negative version of the

remodeller (Bornelöv et al., 2018; Feng and Zhang, 2001) will likely produce a heterogeneous

mix of cells with varying levels of remodeller activity and displaying increasing degrees of cell

cycle arrest and apoptosis over time.

The CHD4/NuRD function described at enhancers has been defined both in genetic

mutants and over a time course of NuRD reintroduction to null cells (Bornelöv et al., 2018;

Reynolds et al., 2012) . In contrast, the noise reduction function has been described in

genetically NuRD-deficient or NuRD-null cells or after NuRD component knockdown (Burgold

et al., 2019; Montibus et al., 2023; Ragheb et al., 2020; Saotome et al., 2024) , meaning that

this could either be a primary function of NuRD or a downstream consequence and/or cell

adaptation of NuRD deletion/depletion. For these reasons we have employed a degron

system, which allows us to acutely deplete CHD4 protein in mouse ES cells and assess the

consequences to chromatin and gene expression within minutes to hours of protein

depletion, long before the cells begin to exhibit cell cycle defects.

Results

CHD4 has an immediate and widespread impact on chromatin accessibility

NuRD in mouse ES cells can exist in various forms and it can contain different paralogs of

key components such as MBD2 or MBD3, GATAD2A or GATAD2B, HDAC1 or HDAC2, and

MTA1, 2 and/or MTA3 (Reid et al., 2023). In embryonic stem cells NuRD has only one

chromatin remodelling subunit: CHD4. We therefore created ES cell lines in which the

endogenous *Chd4* alleles were homozygously tagged with a mini-Auxin inducible degron

(mAID) (Nishimura et al., 2009). CHD4-mAID was largely depleted from both the nucleoplasm

and chromatin within 60 minutes of Auxin addition (Fig. 1A). CHD4 depleted cells showed a

normal cell cycle profile until 24 hours of depletion, when they began to undergo cell cycle

arrest at the G1/S checkpoint (Fig. 1B). By 30 hours CHD4-depleted cells were undergoing

apoptosis, which increased through 48 hours. We therefore focused our analyses on the first

few hours after CHD4 depletion, well before cells started to undergo cell cycle arrest.

CHD4 depletion had an immediate and widespread impact on chromatin accessibility, as

measured by calibrated ATAC-seq. After 60 minutes of CHD4 depletion there were more than

thirteen thousand sites showing a significant change in accessibility, which increased to over

fifty thousand sites showing increased accessibility and eight thousand showing decreased

accessibility by 4 hours (Fig. 1C). These data are consistent with previous reports of overall

increased accessibility upon CHD4/dMi2 depletion by shRNA in mouse ES cells or Drosophila

S2 cells (Dieuleveult et al., 2016; Moshkin et al., 2012).

Differentially accessible regions increasing in accessibility upon CHD4 depletion exhibited

low accessibility and low overall enrichment for specific histone modifications associated

with active chromatin in undepleted cells (H3K27Ac, H3K4Me1, H3K4Me3), but not

H3K27Me3 (Fig. 1D,E). As a class, these sites predominantly mapped to inactive enhancers

and repetitive elements (Fig. 1F). They had very low average enrichment for CHD4 or MBD3

(Fig. 1G) and would likely not pass the threshold to be counted as "Peaks" in many ChIP-seq

or Cut&Run datasets. Low enrichment for active chromatin marks and NuRD components as

well as low but detectable accessibility indicates that these sites predominantly represent

silent and cryptic regulatory sequences. Upon CHD4 loss accessibility at these sites

increased two- to three-fold at 60 and 240 minutes of depletion (Fig. 1D) indicating that the

low level CHD4 enrichment at these sites may nevertheless be functionally relevant.

Sites showing decreased accessibility without CHD4 were highly accessible in undepleted

cells, enriched for marks of active chromatin such as H3K4Me1 and H3K27Ac, and to a lesser

extent H3K4Me3 (Fig. 1D,E), and often overlapped with known enhancers and promoters

(Fig.1F). These sites also showed high enrichment for CHD4 and MBD3 in the undepleted

state (Fig. 1G). Upon CHD4 depletion accessibility at these sites was reduced by less than

two-fold on average (Fig. 1D). We and others have previously shown that NuRD acts at active

enhancers (Basu et al., 2023; Bornelöv et al., 2018; Dieuleveult et al., 2016) and, consistent

with these findings, CHD4 depletion caused an average decrease in accessibility at active

enhancers (Fig. 1H). We therefore conclude that CHD4 activity maintains closed chromatin

generally at inactive regulatory sequences but also contributes to the maintenance of highly

accessible chromatin at active sites.

Figure 1

Figure 1. The impact of CHD4 depletion on chromatin accessibility. A. Western blots of nuclear soluble (Nucleoplasm) and chromatin fractions across CHD4 depletion probed with antibodies directed against indicated proteins. Times in hours of Auxin addition are indicated across the top. Lamin B1 and Histone H3 act as loading controls. **B**. Cell cycle analysis across CHD4 depletion time course. Hours post Auxin addition are indicated at the bottom, "CTRL" indicates DMSO control. Data represent an average of three replicates. Asterisks indicate significant differences from CTRL using a mixed-effects model with Dunnett's multiple comparisons correction. *p < 0.05; **p < 0.01. **C**. Volcano plots of differentially accessible ATAC-seq peaks between 0 and 1 hour of Auxin addition (left) or 0 and 4 hours of Auxin addition (right). Magenta spots indicate statistically significant differences (FDR > 0.05). Numbers of significant peaks decreased or increased are indicated on the plots. **D**. Heatmaps of ATAC-seq signal for all regions displaying increased accessibility (N=52041) or decreased accessibility (N=8384) at any time across the CHD4 depletion time course are displayed for each time point. **E**. Heatmaps of Cut&Run data for indicated histone modifications at sites increasing or decreasing in accessibility (as in panel **D**) (H3K27Ac and H3K4Me1 from this study; H3K27Me3 and H3K4Me3 taken from (Lando et al., 2024)) at indicated times after CHD4 depletion. **F**. Percentages of sites increasing in accessibility (top, blue) or decreasing in accessibility upon CHD4 depletion (bottom, red) which localise to indicated genomic features. Active enhancers are defined as having H3K4Me1 and K3K27Ac but not K3K4Me3, and inactive enhancers as having H3K4Me1 but not H3K4Me3 or H3K27Ac. **G**. Heatmaps of CHD4 and MBD3 Cut&Run data at upDARs and downDARs in 2iL conditions. **H**. Heatmaps of ATAC-seq signal at active enhancers (N=4707) across the CHD4 depletion time course. Median curves in graphs in **D**, **E**, **G** and **H** are plotted with standard error of the mean in lighter shading.

CHD4 activity maintains expression of active genes while reducing transcriptional noise

Significant changes in gene expression were first detected in nascent and total mRNA 1-2 hours after CHD4 depletion (Fig. 2A,B). Loss of CHD4 resulted in an approximately 2:1 ratio of increased to decreased gene expression from 1 hour onwards (Fig. 2C), consistent with CHD4 activity able to both facilitate and limit transcription, with most genes changing by two-fold or less (Fig. 2D). GO terms associated with activated genes indicate various tissue-specific functions, consistent with the general noise reduction activity described for NuRD (Burgold et al., 2019; Montibus et al., 2023; Ragheb et al., 2020). Downregulated genes, in contrast, are associated with cellular maintenance and early development, consistent with these genes being normally highly expressed in ES cells (Fig. 2E).

To assess whether the observed changes in chromatin accessibility were linked to changes in gene expression, we plotted the distance between differentially accessible regions and the annotated TSS of genes found to be misexpressed up to 4 hours after CHD4 depletion. Sites increasing in accessibility upon CHD4 depletion tended to be located far from genes repressed by CHD4 (Fig. 3A). This makes it unlikely that CHD4 directly silences gene expression by maintaining condensed chromatin at inactive promoters widely, although it

Figure 2. CHD4 acutely regulates gene expression. A and B. Heatmaps from nascent RNA-seq (A) or bulk RNA-seq (B) of genes showing significant (P_adj <0.05) changes in expression at any point during the CHD4 depletion time course. Heatmaps displays z-scores, meaning

expression for each gene has been centred and scaled across the entire time course. **C.** Volcano plots showing significant gene expression changes at indicated time points in nascent RNAseq (top) and bulk RNAseq (bottom). Genes increasing upon CHD4 depletion are shown in red, and those decreasing are shown in blue. The number of significantly misexpressed genes at each time point is indicated in the figure. **D**. Violin plots showing the average $Log_2FoldChange$ of significant upregulated (red) and downregulated (blue) genes during the CHD4 depletion timecourse. **E**. Gene Ontology (GO) enrichment analysis of genes increased or decreased after four hours or 24 hours of CHD4 depletion. The top 10 biological processes are shown for each category, based on smallest adjusted p-value.

could be acting to prevent activation of distal enhancers. In contrast, a notable number of

sites losing accessibility are located within 1Kb of the TSS of misregulated genes (Fig. 3A). This

indicates that CHD4 acts at some highly active promoters to maintain both chromatin

accessibility and transcriptional fidelity.

If loss of CHD4 resulted in activation of normally silent or cryptic enhancers, we would

expect that they should show increases in enhancer-associated histone modifications and

binding of transcription factors (TFs). As expected, Cut&Tag for H3K4Me1 and H3K27Ac

showed increasing enrichment at CHD4-condensed sites 4 hours after CHD4 depletion,

which did not increase further by 24 hours of depletion (Fig. 3B). Although these sites were

largely unbound by NANOG or SOX2 in the presence of CHD4, within 1 hour of CHD4

depletion, they became extensively bound by both transcription factors (Fig. 3C).

Consistently, thousands of new NANOG and SOX2 peaks were detected across the genome

after CHD4 depletion (Fig. 3D). For example, the 25 kb region upstream of the *Eomes* gene

contains multiple CHD4-condensed sites (Fig. 3E). Those labelled 1, 2 and 3 show very little, if

any, accessibility or TF binding, and very low H3K27 acetylation and H3K4 monomethylation

in undepleted cells. All three of these show a gain in accessibility, TF binding and both histone

modifications upon CHD4 depletion. Those sites labelled 4 and 5 both show some

accessibility, TF binding and enrichment for active histone modifications in undepleted cells.

Nevertheless, these sites also show an increase in all of these features upon CHD4 depletion.

In contrast, the nearby *Eomes* promoter ("P") shows little or no change in accessibility, TF

binding or enrichment for H3K27Ac or H3K4Me1 across the CHD4 depletion time course.

We propose that low-level association of CHD4 and NuRD across euchromatin restricts

chromatin accessibility, preventing transcription factors from binding to consensus sequence

motifs. It is also possible, however, that accessibility increases are a consequence of more

stable transcription factor binding. Failure to maintain this level of chromatin inaccessibility

216 results in activation of inactive and/or cryptic enhancers, which can stimulate inappropriate

217 gene expression.

Figure 3. Chromatin opening upon CHD4 depletion. A. Frequency density distribution of the distance of increasing Differentially Accessible Regions ("↑ DARs," red lines) and decreasing regions ("↓ DAR," blue lines) to the TSS of genes showing increased (solid lines) or decreased (dotted lines) expression within four hours of CHD4 depletion. **B**. Heatmaps of Cut&Tag signal for H3K27Ac, and H3K4Me1 at sites increasing in accessibility (N = 52041) at indicated times of CHD4 depletion. **C**. Heatmaps of NANOG and SOX2 Cut&Run signal at increasing accessibility sites at indicated times of CHD4 depletion. Median curves in **B** and **C** are plotted with standard error of the mean in lighter shading. **D**. Pairwise comparisons of called peaks of binding for NANOG (top) and SOX2 (bottom) between undepleted cells (0h) and 30 minutes, 1 hour or 4 hours of CHD4 depletion. Significantly changed (FDR>0.05) binding sites are shown in blue when $\log_2FC>0$ and red when $\log_2FC<0$. **E**. IGV screenshot of the upstream region of the mouse *Eomes* locus displaying ATAC-seq, Cut&Run and Cut&Tag data as indicated at left. Boxed regions labelled 1-5 are CHD4 condensed sites, while the box labelled P corresponds to the *Eomes* promoter.

CHD4 and SALL4 maintain chromatin inaccessibility largely independently

The SALL4 protein is a very abundant transcription factor in mouse ES cells, which has long been known to interact with the NuRD complex (Kloet et al., 2015; Lauberth and Rauchman, 2006; Lu et al., 2009; Miller et al., 2016). SALL4 preferentially binds to A/T-rich DNA genome-wide, where it is proposed to exert a general repression function through NuRD recruitment (Kong et al., 2021; Pantier et al., 2021; Ru et al., 2022; Watson et al., 2023). If this model is correct, we would expect some proportion of the CHD4 condensed sites to be dependent upon both CHD4 and SALL4 to remain inaccessible. To test this model, we assessed chromatin accessibility after dTAG-mediated SALL4 depletion. SALL4 is partially redundant with SALL1 in ES cells, but only SALL4 is required for early mammalian development (Miller et al., 2016; Nishinakamura et al., 2001). We therefore created $Sall1^{-/-}Sall4^{FKBP/-}$ ES cells in which one endogenous $Sall4$ allele was mutated and the remaining allele was targeted to express a SALL4-FKBP fusion protein.

Acute depletion of SALL4-FKBP resulted in rapid and extensive changes in chromatin accessibility as measured using ATAC-seq. Over 18,000 sites showed increased accessibility within 1 hour of SALL4 depletion in $Sall1^{(-/-)}$ ES cells (Fig. 4A,B). These sites showed moderate enrichment for SALL4 in undepleted cells, though less than that seen at active enhancers (Fig. 4C). When we compared these sites to those showing CHD4-dependent chromatin condensation, we find that only 4559 sites of the more than 52,000 CHD4 -dependent sites also show dependency upon SALL4 to maintain chromatin inaccessibility (Fig. 4D-F). This means that only 17.2% of SALL4-dependent sites also rely on CHD4 to prevent chromatin opening. Moreover, over 47,000 CHD4-dependent sites show neither SALL4 binding nor SALL4 dependence to remain inaccessible (Fig4. D-G). Similarly, over 20,000 SALL4-dependent sites show no change in accessibility upon CHD4 depletion, despite showing similar levels of enrichment for NuRD components as those dependent upon both SALL4 and CHD4 (Fig. 4E,H). Consistent with the proposed preference of SALL4 for binding to A/T-rich DNA, SALL4-unique regions showed enrichment for SALL4 binding and a high A/T-content (53.9%) (Fig. 4D). CHD4-unique regions showed little if any SALL4 enrichment and a much lower A/T content (44.2%), while those sites requiring both SALL4 and CHD4 to remain inaccessible showed enrichment for NuRD components as well as SALL4 and displayed intermediate A/T content (48.2%). Together, these data show that both CHD4 and SALL4 broadly restrict chromatin accessibility across the genome, but that their activities overlap at only a relatively small proportion of sites.

Figure 4. CHD4 and SALL4 both restrict chromatin accessibility. A. Volcano plots of differentially accessible ATAC-seq peaks when comparing 1 hour, 4 hours or 24 hours of SALL4 depletion with those seen in undepleted cells (0). Magenta spots indicate statistically significant differences (FDR > 0.05). Numbers of significant peaks decreased or increased are indicated on the plots. **B**. Heatmaps of ATAC-seq signal for all regions displaying increased accessibility (N=26510) across the SALL4 depletion time course are displayed for each time

point. **C**. Heatmap of SALL4 Cut&Run signal in undepleted ES cells (taken from (Ru et al.,
2022)) at all regions displaying increased accessibility across the SALL4 depletion time
course (left, N=26510) or at active enhancers (right, N=4339). **D**. Overlap of sites showing
increased accessibility upon SALL4 depletion with those increasing upon CHD4 depletion
(upDARs). The % A/T base composition of the different categories of sites is indicated. **E, F**.
Heatmaps of ATAC-seq signal at sites increasing upon either SALL4 or CHD4 depletion
(SALL4+CHD4), sites increasing upon CHD4 depletion but not upon SALL4 depletion (CHD4
Only), or sites increasing upon SALL4 depletion but not upon CHD4 depletion (SALL4 Only)
plotted at indicated time points of CHD4 depletion (**E**) and SALL4 depletion (**F**). **G, H**.
Cut&Run signal for SALL4 (**G**) or for CHD4 and MBD3 (**H**) in undepleted ES cells at the three
different classes of sites. Median curves in **B, C,** and **E-H** are plotted with standard error of the
mean in lighter shading.

NuRD limits transcription factor binding to chromatin

Sites decreasing in accessibility upon CHD4 depletion showed an increase in enrichment

for both NANOG and SOX2 after 30 minutes, even though these sites were, on average, losing

accessibility and losing enrichment for the active chromatin marks H3K4Me1 and H3K27Ac at

that time (Fig. 5A,B). Active enhancers similarly showed an increase in NANOG and SOX2

enrichment from 30 minutes of CHD4 depletion (Fig. 5C). An example locus is shown in Figure

5D, where a cluster of enhancers located 50-70Kb downstream of the *Klf4* gene contains

several high accessibility sites. Those located at 68, 57 and 55 Kb downstream of *Klf4*

(labelled 68, 57 and 55, respectively in Figure 5D) all show a loss of ATAC-seq signal but a gain

in NANOG and, to a lesser extent, SOX2 enrichment between 0 and 30 minutes of CHD4

depletion. NANOG enrichment then decreases again at 1 and 4 hours while Sox2 enrichment

remains relatively constant, even though accessibility continues to decrease.

The simultaneous decrease in chromatin accessibility and increase in transcription factor

enrichment after 30 minutes of CHD4 depletion could indicate that CHD4/NuRD plays a

direct role in controlling transcription factor binding to chromatin. To independently assess

the impact CHD4 has on transcription factor binding, we used single molecule imaging in live

ES cells to directly measure transcription factor binding kinetics upon CHD4 depletion. A

HALO-Tag cassette was knocked-in to the endogenous *Nanog*, *Klf4* or *Sox2* loci in the CHD4-

mAID ES cell line to create strains expressing the different C-terminal protein fusions. Single

HALO-tagged protein molecules were then labelled with a photoactivatable dye (PA-JF646,

(Grimm et al., 2016)) and tracked at two distinct temporal regimes where we collected images

either every 20 ms or 500 ms, using double-helix point spread function (DH PSF) microscopy

as they moved within a 4-μm slice of the nucleus. Recording at a 20 ms time resolution allows

the segmentation of trajectories into freely diffusing and chromatin-bound states, and this

data can be used to extract the chromatin binding kinetics of proteins and complexes (Basu et

al., 2023). Here, we used these data to determine residence times – how long the molecules

remain bound to chromatin. The proportion of molecules that remain bound to chromatin

after increasing lengths of time was then plotted before and after CHD4 depletion (for 1 hour)

and the results were fitted with a single exponential decay to determine the apparent

dissociation rate, k_{off} (Fig. 5E). NANOG-HALO, SOX2-HALO and KLF4-HALO all showed a

pronounced decrease in apparent k_{off} after CHD4 depletion (Fig. 5F-H), consistent with CHD4

actively limiting transcription factor residence times on chromatin.

To determine whether it is CHD4 by itself or the intact NuRD complex that can remodel

transcription factor binding, we similarly imaged NANOG-HALO and KLF4-HALO in MBD3-

depletable ES cells after disruption of the NuRD complex (Fig. 5I). These cells were also null

for *Mbd2*, which encodes a protein displaying partial functional redundancy with MBD3 but is

dispensable in ES cells (Hendrich et al., 2001). As CHD4 is largely responsible for the binding

of NuRD to its target enhancers, removal of MBD3 enables disassembly of NuRD, but does not

prevent CHD4 binding to chromatin (Basu et al., 2023; Bornelöv et al., 2018; Zhang et al.,

2016). The nucleosome remodelling activity of CHD4 is, however, greatly reduced outside of

NuRD (Bornelöv et al., 2018). We therefore reasoned that if it is intact NuRD that is required

(as opposed to just CHD4), depletion of MBD3 should also limit TF residence times. Indeed,

we detected a significant decrease in the apparent k_{off} of stably bound NANOG-Halo and

KLF4-Halo after 1 hour of MBD3 depletion (Fig. 5J,K). We therefore conclude that intact NuRD

(and not CHD4 by itself) actively limits transcription factor residence times on chromatin,

while at the same time contributing to the maintenance of chromatin accessibility at active

regulatory regions.

Figure 5. NuRD regulates NANOG and SOX2 binding to active sites. A. Heatmaps of
NANOG and SOX2 Cut&Run signal at indicated times of CHD4 depletion across sites
decreasing in accessibility (N = 8384). **B**. Heatmaps of H3K27Ac and K3H4Me1 Cut&Tag signal

at indicated times of CHD4 depletion across decreasing accessibility sites. **C**. Heatmaps of
Cut&Run signal for NANOG and SOX2 across active enhancers (N = 4707) at indicated times
of CHD4 depletion. Median curves in **A-C** are plotted with standard error of the mean in lighter
shading.**D.** IGV screenshot of the enhancer cluster downstream of the *Klf4* gene displaying
ATAC-seq, Cut&Run and Cut&Tag data as indicated at the left. Boxed regions are labelled with
the distance in Kb from the annotated *Klf4* transcription start site. **E**. Fluorescence survival
curves of chromatin-bound NANOG-HALO molecules in 2iL (blue line) or after 1 hour of CHD4
depletion (purple line). The grey dotted line represents the fluorescence survival curve for
molecules in a fixed-cell control imaged under identical conditions. **F**. Apparent dissociation
rates (k_{off}) of chromatin-bound NANOG molecules calculated through fitting a single
exponential decay model to the survival curves in **panel E**. Error bars represent 95%
confidence intervals for each fit applied to data taken from three independent experiments.
The horizontal dashed line represents the upper 95% confidence limit for a fixed-cell control.
** indicates that 99% confidence intervals do not overlap, i.e. p < 0.01. **G.H**. As in **F** but for
SOX2 (**G**) and KLF4 (**H**). The fixed cell control was not imaged in the SOX2 experiments in
panel **F. I**. Western blots of nuclear soluble (Nucleoplasm) and chromatin fractions across
MBD3 depletion probed with antibodies directed against the indicated proteins. Times in
hours of Auxin addition are indicated across the top. Lamin B1 and Histone H3 act as loading
controls. **J, K.** Apparent dissociation rates (k_{off}) for NANOG-HALO (**I**) and KLF4-HALO (**J**) before
and after 60 minutes of MBD3 depletion. For calculation of k_{off} the trajectories were pooled
from 4 replicates of each time point obtained over 2 days.
364

NuRD activity influences active and inactive regulatory elements differently

We next asked how CHD4 activity could have different impacts on accessibility at different

kinds of sequences. Information about the structure of accessible regions can be obtained

from ATAC-seq data by quantitating recovered fragments of all sizes and determining the

frequency and location of recovered Tn5 integration sites using VPlots, (Henikoff et al., 2011;

Schep et al., 2015; Serizay and Ahringer, 2021)(Supplementary Figure 2). Applying this

analysis at CHD4 -ondensed sites shows an increase in both Tn5 integrations and in reads

corresponding to the nucleosome-free region (NFR) after 30-minutes of CHD4 depletion (Fig.

6A,B). After 60 minutes, when CHD4 is almost completely depleted, there is an increase in

accessibility generally across the entire site, both of short reads within the NFR but also

longer reads (200-300bp) extending from the NFR outwards (Fig. 6B). We conclude that NuRD

is acting to restrict accessibility of both the NFR and the flanking nucleosomal DNA (see

Supplementary Figure 2) at these sites, consistent with the demonstrated role of CHD4/NuRD

in maintaining the density of intact nucleosomes (Bornelöv et al., 2018; Dieuleveult et al.,

2016; Moshkin et al., 2012).

The example of the *Eomes* locus (Fig. 3F) illustrates that some CHD4-condensed sites

show a degree of accessibility in the presence of CHD4, while others show little if any

accessibility. To better understand how CHD4 prevents chromatin opening at these two

different kinds of sites, we constructed Tn5 integration plots and Vplots from the 1000 CHD4-condensed sites showing the least accessibility in undepleted conditions and from the 1000 showing the most accessibility (Fig. 6C-F). Sites showing little or no accessibility in undepleted cells initially show very few reads less than 100bp. The V-plots also show reads of 200-300 bp distributed across the regions, again indicative of generally inaccessible chromatin. Loss of CHD4 results in the formation and progressive expansion of an NFR, as well as an increase in longer reads with one end located within the NFR (Fig. 6C,D). The Tn5 integration plot does not show a general widening of the NFR over time, indicating that CHD4 function is largely focussed on maintaining nucleosome density within a 100-200bp region (Fig. 6C). Sites at which CHD4 activity restricts accessibility at already accessible sites showed a general increase in both small and longer reads as CHD4 was depleted, and a broadening of the existing NFR (Fig. 6 E,F). Together these data are consistent with CHD4/NuRD maintaining chromatin compaction at inactive or low activity enhancers by maintaining nucleosome density.

The sites at which CHD4 activity maintains accessibility displayed a prominent NFR as well as high density of longer reads in undepleted cells (Fig. 6G,H). CHD4 depletion induced a general decrease in Tn5 integrations and VPlot signal intensity across these sites, consistent with CHD4 acting to generally maintain accessibility.

We next asked why CHD4 promoted accessibility at highly accessible active sites (e.g. Fig. 6G,H) but reduced accessibility at slightly less accessible, inactive sites (e.g. Fig. 6E,F). We took advantage of ES cells in which a tamoxifen-inducible MBD3b is expressed in an otherwise *Mbd3*-null ES cell line, allowing us to restore NuRD activity to cells upon tamoxifen addition (Bornelöv et al., 2018; Reynolds et al., 2012). We had previously subjected these cells to micrococcal nuclease (MNase) sequencing at different time points after MBD3 induction to show that restoration of NuRD activity caused increased density of intact nucleosomes across sites of active transcription (Bornelöv et al., 2018). Unlike ATAC-seq, sequencing of MNase-treated DNA results in increased reads at sites protected from digestion by bound proteins, i.e. less accessible regions, and a decrease in signal at locations where the DNA is easily accessible and therefore digested by the MNase. In that study, we used an MNase concentration at which DNA associated with intact nucleosomes is protected from MNase digestion. Lower MNase concentrations, however, will recover DNA protected by other chromatin proteins as well as by partial (fragile) nucleosomes (Kubik et al., 2015; Xi et al., 2011).

We therefore subjected cells undergoing the time course of NuRD reformation to lower MNase concentrations prior to sequencing. Traditional MNase sequencing shows that NuRD increases the density of intact nucleosomes at both classes of accessible sites, as indicated by an increase in signal from the red line (NuRD-deficient) to the black line (24h NuRD restored) in the top panels of Figure 6I. This effect is more pronounced at the most accessible inactive sites ("UP Most Accessible", black line, Fig. 6I), occurring across a broader area than at fully active sites ("All Decreasing", black line). The lower MNase concentration, by contrast (Fig. 6I, lower panels) produces a peak of MNase resistance in the absence of NuRD activity at both classes of sites (red lines), indicating an accumulation of structures which are digested by the higher MNase concentration, such as fragile nucleosomes. Restoration of NuRD activity clears the majority of this signal at the most accessible inactive sites, but only moderately reduces their abundance at fully active sites (Fig. 6I, lower panels). These data indicate that at highly active regions, NuRD acts to limit the abundance of fragile nucleosomes and other proteins, while marginally increasing the density of intact nucleosomes, which has an overall result of maintaining these sites in a fully active state. It does not completely remove the fragile nucleosomes from these sites, which are presumably being created by the activity of other chromatin remodellers (Klein et al., 2023; Nocente et al., 2024). In contrast, at accessible but inactive sites it prevents the accumulation of fragile nucleosomes which would otherwise cause further opening and enable activation of inactive regulatory elements. This means that the activity of CHD4 does not differ at the two classes of sites, but it is the presence of other remodellers producing fragile nucleosomes which dictates whether CHD4 activity acts to maintain or prevent accessibility (Fig. 7).

Together, these data show that CHD4 shows two modes of binding on chromatin: in addition to the high enrichment seen at active enhancers and promoters, it also shows low-level binding at silent and/or cryptic regulatory elements across euchromatin. At the former class of sites, it acts to limit the residence times of chromatin-associated proteins to maintain highly accessible chromatin. This maintenance activity of CHD4/NuRD is necessary for the full activity of these elements to direct appropriate transcription of target genes. At the latter class of sites we propose it maintains chromatin in an inaccessible state, preventing accumulation of fragile nucleosomes and preventing transcription factors from recognising and stably binding their cognate motifs and thereby prevents spurious activation of these elements.

Figure 6. CHD4 controls accessibility differently at different classes of sites. A, **C, E, G**
Tn5 integration frequency was determined from ATAC-seq data and plotted across indicated
sites across the CHD4 depletion time course. The number of sites is shown in parentheses. **B,**
D, F, H Vplots across sites indicated above (corresponding to the Tn5 integration plots) at

indicated times of CHD4 depletion. See **Supplementary Figure 2** for a Vplot schematic. **I.**
MNase-seq data collected 0 (red), 30 minutes (dark red) or 24 hours (black) after of NuRD
reformation in *Mbd3⁻/⁻* ES cells are plotted across indicated sites. Top graphs show results
from "Hard" MNase treatment while bottom graphs show "Soft" MNase treatment (see text).
The y-axis shows normalised read coverages, while the X-axis shows distance in base pairs
from the centre of the feature. Curves show mean and standard error from 3 biological
replicates.
460

Discussion

Assessing CHD4 function by traditional knockout experiments is difficult as ES cells

without CHD4 undergo cell cycle arrest and begin to apoptose after 24 hours (Fig. 1B). Even

with siRNA or inducible deletion, cells in a population will lose CHD4 at different times and, in

the case of siRNA, to different extents, making primary function difficult to ascertain. Genetic

deletion in mouse ES cells has been used extensively to define protein function in cells and in

intact mice. In such an experiment the genetic lesion will often be induced in a single cell, and

a clone of cells will be grown out from that single cell. While this is an extremely powerful

method for defining gene function, there is usually a very large number of cell divisions

occurring and considerable time passing between loss of the gene and assessment of

phenotype, making it of limited use when studying proteins required for the viability of ES

cells. Resulting phenotypes may be due directly to the absence of protein function but may

also reflect adaptations made by cells selected to proliferate in the absence of that protein.

The advent of inducible depletion methods (summarised in (Wit and Nora, 2023)) has meant

we can now infer protein function by assessing the immediate molecular consequences of

loss of a given protein activity in the very short term. Combined with genome-wide profiling,

this has allowed us to identify both a novel transcription factor remodelling activity for CHD4

and to define how this combines with the protein's nucleosome remodelling activity to exert

different functions at different classes of sites.

CHD4 is an abundant chromatin remodeller with a well-defined function in moving intact

nucleosomes. Here we show that CHD4 can also actively limit the residence times of

transcription factors on chromatin. The combination of these two activities has different

consequences at different kinds of sequences. At highly accessible, active sites (Fig. 7A)

CHD4 enrichment is high and it acts to limit the binding of transcription factors and fragile

nucleosomes to regulatory regions, allowing for full accessibility and functionality of these

regulatory elements. At inactive, less accessible euchromatic sites (Fig. 7B) CHD4 associates

more infrequently, but here it's TF removal activity, combined with maintenance of

nucleosome structure, is sufficient to keep accessibility low and to prevent transcription factors from stably binding recognition motifs. This activity is sufficient to prevent the binding of NANOG and SOX2, the latter being capable of binding nucleosomal templates (Dodonova et al., 2020; Zhu et al., 2018). By combining these two activities, CHD4 functions to prevent spurious activation of cryptic and/or silent regulatory regions, thereby reducing transcriptional noise, but also acts to facilitate the activity of highly accessible regulatory regions, allowing cells to accurately and rapidly respond to differentiation cues (Fig. 7).

Many transcription factors recognise and bind to specific sequence motifs in DNA. A transcription factor recognising a 7 base pair sequence should, on average, have more than 150,000 different potential binding sites in a mammalian genome. Yet ChIP-seq and Cut&Run studies have found most transcription factors associate with orders of magnitude fewer sites in any one cell type (Lambert et al., 2018; Srivastava and Mahony, 2020). Most transcription factors bind preferentially or exclusively in accessible chromatin, so it is believed that differences in chromatin accessibility, or nucleosome positioning across sites, determines which subset of the potential consensus DNA binding motifs are available for TF binding at any given time or cell type (Workman and Buchman, 1993; Zhu et al., 2018). Here we identify CHD4 as one factor which acts to prevent transcription factors, even so-called Pioneer Factors, from binding thousands of sites in inaccessible chromatin. Consistently with this model, CHD4 was found to prevent binding of the GATA3 transcription factor to inappropriate sites in breast cancer cells undergoing mesenchemal-to-epithelial transition (Saotome et al., 2024). We propose that CHD4's chromatin remodelling activity keeps the sites inaccessible, so TF binding is not favoured (i.e. k_{on} is low). The ability of CHD4 to remove TFs from chromatin means that even if they are capable of binding to a nucleosomal substrate, CHD4 promotes their dissociation from chromatin so they are quickly removed (Fig. 7B). At highly accessible chromatin, CHD4 similarly limits TF residence times by increasing k_{off}, however accessibility is high so the k_{on} is also high. We propose that at these sites CHD4 activity balances the high k_{on} by maintaining a high k_{off}, thus facilitating the turnover of bound molecules (Fig. 7A). Without CHD4, k_{off} decreases while k_{on} remains high, resulting in more stable binding of TFs to chromatin. CHD4 and NuRD have been shown to be important for cells to properly respond to differentiation cues in a wide variety of organisms and contexts (Ahringer, 2000; Bracken et al., 2019; Lenz and Brehm, 2023; McDonel et al., 2009; Signolet and Hendrich, 2015) and we propose that by facilitating TF turnover and accessibility at promoters and enhancers, CHD4

520 and NuRD facilitate the ability of regulatory elements to respond to differentiation cues (Fig.

521 7).

A. Active sites   Figure 7

B. Inactive sites

Figure 7. Model of CHD4 function. **A. Highly accessible sites.** In undepleted cells these
sites are extensively bound by CHD4/NuRD, where it acts to promote the off rate of
transcription factors to promote accessibility. Tn5 is able to access the central NFR but also
can integrate into the flanking nucleosomal DNA. After CHD4 depletion, the on rate for TFs
does not change, but the off rate is now much reduced, resulting in increased TF binding. The
sites become less accessible to Tn5, such that although it can still access the hypersensitive
site within the NFR, there are fewer integrations extending outwards. These regulatory regions
cannot quickly respond to receipt of external signals. **B**. Model of CHD4 function at
inaccessible, silent enhancers. In undepleted conditions, there is low CHD4 enrichment at
these sites. Here, CHD4 acts to prevent binding of transcription factors and maintain low
accessibility, such that Tn5 cannot frequently access the DNA. After CHD4 depletion, the
locus becomes more accessible, and transcription factors can stably bind. This leads to
spurious activation of distal promoters and an increase in transcriptional noise.

CHD4's activity to prevent spurious TF binding and chromatin opening occurs at many sites

where enrichment of CHD4 by ChIP or Cut&Run is low, but not absent. Despite the low

amount of CHD4 detected at these sites, this activity was sufficient to prevent activation of

silent and cryptic regulatory regions (Fig. 2). This low-level enrichment of CHD4 at sites where

it prevents activation could explain an observation made during Drosophila spermatogenesis,

where the CHD4 orthologue dMi2 was found to prevent transcription from cryptic promoters,

despite not being detected at these promoters by ChIP-seq assays (Kim et al., 2017). A recent

study in *S. cerevisiae* found many transcription factors appear to exert influences over

expression of genes without detectable association at that gene and bind many sites where

they exert no detectable regulatory function (Mahendrawada et al., 2025). While some have

argued that one should focus on high confidence binding targets to assess protein function

(Mindel and Barkai, 2025), in our case this would have resulted in us ignoring most of the

50,000 sites at which we detect CHD4-dependent chromatin compaction (Fig. 1C). While we

have argued that chromatin changes detected within 30 minutes are likely to be direct

consequences of NuRD manipulations (Fig. 6), we do acknowledge that it is possible these

are indirect consequences of loss or gain of NuRD activity. We agree that the definition of

"targets" can be highly variable between studies, which is why we preferred to assess CHD4

enrichment at subsets of sites where function was identified, irrespective of whether it

passed the threshold of being a "peak."

SALL4 presents another example of a protein exerting function away from its predominant

sites of chromatin enrichment: initial ChIP-seq results indicated that SALL4 was

predominantly located at enhancers, whereas more recent Cut&Run data indicates a wider

spectrum of sites bound by SALL4 (Kong et al., 2021; Miller et al., 2016; Pantier et al., 2021; Ru

et al., 2022). We find that SALL4 shows lower enrichment at A/T-rich genomic sites than it

does at enhancers, but that it nevertheless has a major impact on chromatin compaction at

these genomic A/T-rich sites (Fig. 4). While many groups, including ours, have generally

assumed protein function would be focussed at ChIP-seq "peaks," we argue here that

focusing on protein enrichment levels on chromatin is not necessarily the best way to identify

important sites of protein activity.

Together, our data show that CHD4 exerts functions on chromatin beyond its well-

described ability to slide intact nucleosomes along the DNA (Zhang et al., 2016; Zhong et al.,

2020). Consistent with this conclusion, a recent paper found that acute depletion of CHD4

resulted in an increase in fragile nucleosomes at enhancers (Nocente et al., 2024), while we

and others have found that NuRD acts to limit the amount of paused RNA Polymerase II at

active promoters (Bornelöv et al., 2018; Pundhir et al., 2023). It is possible that CHD4

translocates along the DNA within accessible sequences and displaces any bound proteins,

such as transcription factors or fragile nucleosomes, that it encounters. Both transcription

factors and fragile nucleosomes are less tightly bound to DNA than intact nucleosomes, so it

is possible CHD4's nucleosome remodelling activity results in their displacement, while still

sliding intact nucleosomes. CHD4 could also displace transcription factors from preferred

binding sites, thus making binding less favourable. Such activity might not be necessary in

inaccessible chromatin, but at inactive enhancers showing a small amount of accessibility,

infrequent binding and translocation of NuRD across the accessible region could remove any

proteins which have managed to bind. Whether this is achieved through a similar ATP-

dependent mechanism as is used to slide nucleosomes remains to be determined (Reid et

al., 2024; Zhong et al., 2020). Notably, the yeast remodeller INO80 was found to bind

differently to fragile nucleosomes vs intact nucleosomes, possibly indicating different

mechanisms of remodelling the two different substrates (Wu et al., 2023; Zhang et al., 2023).

A recent study showed that the nucleosome remodelling activity of a different remodeller,

SMARCA5, is dictated by the density of nucleosomes on the DNA that it encounters: at high

density sites it maintains that density, while at low density sites it slides nucleosomes across

the DNA to facilitate TF access (Abdulhay et al., 2023). We find that CHD4 increases density

of intact nucleosomes at sites where it prevents or maintains accessibility (Fig. 6I, top panels)

and also limits TF residence times at both classes of sites. Therefore, unlike SMARCA5, we

see no evidence that the activity of CHD4 differs between condensed and accessible

chromatin, but rather the consequences of that activity are different at different kinds of sites

(Fig. 7).

The residence times of transcription factors vary but are generally on the order of a few

seconds (Lu and Lionnet, 2021). Why might it be important to limit the residence times of

transcription factors? We propose that this could fix an enhancer into a specific state, when

one important job of enhancers is to be responsive to changes in stimuli. If no change in

status quo is required, then after CHD4 promotes eviction of a particular transcription factor

from the enhancer it can continually re-bind and exert its influence over that enhancer.

Should signals change, the enhancer needs to be able to remove TFs corresponding to the old

signal to make space for determinants of the new signal. In this model, CHD4/NuRD

maintains this fluidity of transcription factor interactions, ensuring enhancers are rapidly able

to respond immediately upon receipt of differentiation cues.

Methods

Mouse embryonic stem (ES) cells

Mouse ES cells were cultured in 2i+LIF media on gelatin-coated plates (Montibus et al.,

2023; Mulas et al., 2019). All cell lines were genotyped and tested for mycoplasma regularly.

The CHD4-mAID ES cell line was made in BC8, an F1 hybrid from a C57Black/6 and Mus

castaneus cross (40, XY), obtained from Anne Ferguson-Smith (Cambridge). MBD3-AID ES

cells were created in 23AF, a primary ES cell line derived from $Mbd2^{-/-}$, $Mbd3^{Flox/Flox}$ mice (40,

XX) in a mixed C57Black/6 and 129/Ola background. Sall4-FKBP cells were created in $Sall1^{-/-}$

$Sall4^{+/-}$ ES cells (Miller et al., 2016).

Targeting constructs were made using the AID sequence (Nishimura et al., 2009) or mini-

AID (Kubota et al., 2013) amplified from an Oct4-AID plasmid; a gift from José Silva (Bates et

al., 2021) . Targeting plasmids using the FKBP protein (Nabet et al., 2018) were made by

amplifying FKBP from pLEX_305-C-dTAG; a gift from James Bradner & Behnam Nabet

(Addgene plasmid # 91798). Plasmids created as part of this study are available from

Addgene: https://www.addgene.org/Brian_Hendrich/.

To create Auxin-depletable cell lines, parent lines were first transfected with a PiggyBac

construct to constitutively express OsTir1 (created using pMGS56 (GFP-ARF16-PB1-P2A-

OsTIR1); a gift from Michael Guertin (Addgene plasmid # 129668)) linked to either G418 or

Hygromycin resistance, and cells were cultured under selection to maintain OsTir1

expression. To create degron knock-ins, cells were lipofected with a targeting vector and

appropriate gRNA (Table 1) in a Cas9-expression vector (pSpCas9(BB)-2A-GFP (PX458); a gift

from Feng Zhang; Addgene plasmid # 48138). Drug resistant colonies were genotyped and

correctly targeted clones were subsequently transiently transfected with an expression

plasmid for Dre recombinase to remove ROXed drug selection cassettes. *Chd4* targeting

required two rounds of transfection, selection, and drug removal, while *Mbd3* and *Sall4*

targeting required one round. For protein depletion, cells were treated with 500 µM Auxin or

500 µM dTAG-13 in standard culture media. All HALOTag-fusion ES cell lines were

heterozygous for the HALOTag fusion, i.e. HALO/+, and verified by western blotting.

Gene	gRNA	Sequence (5' -- 3')
Chd4	gRNA1	GGTGGAGGTGGATATCACTC
Mbd3 A3xF	gRNA1	TTCTCACGCGTCACTCGCTC
Mbd3	gRNA2	CAGCCATTCCCTGGAAGTAC
Sall4	gRNA1	AATAAGATTGCTGTCAGCTA
Sall4	gRNA2	AAGATTGCTGTCAGCTAAGG
Nanog	gRNA1	AACTACTCTGTGACTCCACC
Sox2	gRNA1	TGCCCCTGTCGCACATGTGA
Klf4	gRNA2	GTGGGTCACATCCACTACGT

Table 1. gRNA sequences used for gene targeting

Nuclear extract fractionation and western blotting

Nuclear fractionation was carried out as described (Gillotin, 2018). Briefly, cells were collected in ice-cold PBS and pelleted in a refrigerated centrifuge. The cell pellet was lysed by gentle up and down pipetting five times in ice-cold buffer E1 (50 mM HEPES, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% NP40, 0.25% Triton X-100, 1 mM DTT, protease inhibitors). After pelleting and washing in E1 buffer, the pellet was resuspended in ice-cold E2 buffer (10 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, protease inhibitors) and shaken for 45 minutes at 1400rpm at 4°C. The supernatant, representing the nuclear fraction, was collected into a fresh tube. After washing in E2 buffer, the pellet was resuspended in ice-cold E3 buffer (50 mM Tris-HCl, 20 mM NaCl, 1 mM $MgCl_2$, 1% NP-40, protease inhibitors). The resuspended pellet was sonicated at 4°C in a Bioruptor Plus (Diagenode) for 5 minutes, using 30-second ON/30-second OFF cycles at high power. Following sonication, nuclear and chromatin fractions were centrifuged at 16,000xg at 4°C for 10 minutes. 10 µg of extract per lane was used for western blots. Antibodies are listed in Table 2.

Antibody target	Raised in	Supplier	Cat #	Dilution (WB)
CHD4	Mouse	Abcam	ab70469	1/5000
CHD4	Rabbit	Abcam	ab72418	1/10000
FLAG	Mouse	Sigma Aldrich	F3165/F1804	1/2000

HA	Mouse	Invitrogen	26183	1/2000
ADNP	Goat	R&D Systems	AF5919	1/2000
Histone H3	Rabbit	Abcam	ab1791	1/5000
MTA2	Mouse	Abcam	ab50209	1/5000
NANOG	Rabbit	Bethyl Labs	A300-387A	1/2000
SOX2	Rat	E-bioscience	14-9811-82	1/2000
Lamin B1	Rabbit	Abcam	ab133741	1/10000
Mouse IgG	Rabbit	Invitrogen	A16162	1/100
Rabbit IgG	Guinea Pig	Antibodies Online	ABIN101961	1/100
Goat IgG	Rabbit	Abcam	ab6697	1/100
Rat IgG	Rabbit	Abcam	ab6703	1/100
Mouse IgG	Sheep	VWR	NA931V	1/2000
Rabbit IgG	Donkey	VWR	NA934V	1/2000

Table 2. Antibodies used in this study with dilutions used for western blot.

Cell cycle analysis

The cell cycle distributions of auxin (500 µM) or dTAG-13 (500 nM) treated ES cell lines were

assessed using propidium iodide (PI) staining coupled with flow cytometry. Approximately

200,000 single cells were fixed using 70% ethanol for at least 24 hours before staining. Fixed

samples were washed twice in PBS and resuspended in 200 µL of PI solution [50 µg/mL

propidium iodide (Invitrogen, P3566) and 50 µg/mL DNase and protease-free RNase A

(ThermoScientific, EN0531), diluted in sterile PBS and incubated overnight at 4°C in the dark.

The fluorescent intensity of stained samples was determined using the Attune NxT

(Thermofisher) equipped with a 561 nm laser line and a minimum of 10,000 single cell events

were recorded. The resulting FCS files were analysed using FlowJo (v10.10) and tested for

significance using a mixed-effects model with Dunnett's multiple comparisons correction.

Single molecule imaging

ES cells were passaged 24 hours before imaging onto either: No 1.0 35 mm glass bottom

dishes (MatTek Corporation P35G-1.0-14-C) or No 1.5 35 mm glass bottom dishes (MatTek

Corporation P35G-1.5-14-C) with their surfaces pre-coated in poly-L-ornithine (Sigma Aldrich

P4957) for ≥30 minutes at 37°C, followed by three PBS rinses at room temperature, followed

by 100μg/ml Laminin (Sigma Aldrich L2020) coating in PBS for >4 hours at room temperature.

Cells were labelled on the day of imaging with either 250 nM HaloTag®-PA-JF646 (a gift from L.

Lavis, Janelia) for 15 minutes, rinsed twice in PBS and incubated for 20 minutes at 37°C in

fresh media. PBS rinsing and 20-minute incubation steps were repeated five more times, and

the cells were then imaged in fresh media.

Single-molecule tracking was carried out using oblique illumination (Tokunaga et al., 2008)

on a custom-built double helix point spread function microscope with a Nikon Eclipse Ti-U

inverted microscope body and a box incubator set to 37°C (Carr et al., 2017). Beams were

expanded and collimated using Galilean beam expanders, then combined using dichroic

mirrors. For 2D imaging, a Nikon 1.49 NA 60× oil immersion objective (CFI Apochromat TIRF

60XC Oil) was used to focus excitation beams onto the sample, and the microscope was also

set to an internal magnification of 1.5x. For 3D DHPSF imaging, a Nikon 1.27 NA 60x water

immersion objective lens (CFI SR Plan Apo IR 60XC WI) was used without internal 1.5x

magnification. The emission path of the microscope was also modified to include a fixed

double helix phase mask (DoubleHelix, Boulder, CO) in the Fourier domain of the emission

path of the microscope (Carr et al., 2017; Pavani et al., 2009) . For long-exposure imaging,

control samples were fixed in 4% formaldehyde (v/w) in PBS for 10 minutes at room

temperature, rinsed twice with PBS and stored in PBS. Samples were imaged in identical pre-

warmed culture media and under identical conditions to live-cell samples. Across all

experiments, three separate samples per condition were imaged in a day and all experiments

were repeated on at least two separate days for biological replications.

Background subtraction with a five-pixel rolling ball radius was carried out on image stacks

using open-source software Fiji (Schindelin et al., 2012). Single molecules imaged in 2D were

localised using the PeakFit tool within the GDSC Single Molecule Light Microscopy (SMLM)

plugin (https://github.com/aherbert/gdsc-smlm (Etheridge et al., 2022)) for Fiji. Localisations

were filtered for precision better than 25 nm. For single molecules imaged in 3D, PeakFit was

used as above but with an initial precision threshold of 40 nm and an additional analysis step

after this. 2D localisations of DHPSF lobes were paired to generate 3D localisations using

DHPSFU (https://github.com/TheLaueLab/DHPSFU). Localisations were then tracked across

frames to generate trajectories using custom Python code

(https://github.com/wb104/trajectory-analysis). Localisations were connected between two

successive frames if they were located within 400 nm of each other.

Short-exposure trajectories were segmented into confined and unconfined sub-trajectories

using the 4P-algorithm from (Basu et al., 2023). Bound fractions were calculated for a sample

based on the number of trajectory frames assigned as being confined, divided by the total

number of trajectory frames. Tracks with a residence time shorter than 1.5 seconds were

filtered out to reduce the impact of noise on further analysis. The decay curve of residence

times for each sample was fitted to a single exponential fit using MATLAB (2018) to yield

apparent dissociation rates. The measured apparent dissociation rate (k_{off}^{app}) was

calculated by fitting a single exponential decay function to the survival (Kaplan-Meier)

function (S) of TF dwell times measured from the point of first observation (t), as below:

$$S = e^{-k_{off}^{app}t}$$

RNAseq

Cells were harvested in Trizol Reagent and RNA purified using Zymo Direct-zol columns

(Zymo Research) according to manufacturer's instructions. 60-100 ng of Ribosomal RNA-

depleted mRNA was used for library preparation with the the NEXTflex RapidDirectional RNA-

seq kit (Illumina). 100 bp paired end (2 x 50 bp) sequencing was performed on a NovaSeq S1

flowcell at the CRUK Cambridge Institute sequencing facility.

Raw data were trimmed using TrimGalore v0.6.4. Reads were indexed and aligned to the

unmasked GRC38/mm10 reference genome using the Burrows-Wheeler Aligner (BWA) (Li and

Durbin, 2009) . Successful removal of adapter and low-quality bases was assessed using

FastQC after trimming. File conversion, sorting, removal of duplicates and mitochondrial

reads were performed using samtools (Li et al., 2009) .

Read counts for genomic features were summarised using featureCounts (Liao et al.,

2014) and a list of the GRC38/mm10 genomic features in a GTF format downloaded from

https://www.ensembl.org/ and https://genome.ucsc.edu/cgi-bin/hgTables. Differential

expression was analysed using DESeq2 (Love et al., 2014) . Raw expression counts were

transformed and normalised to FPKM (fragments per kilobase of transcript per million

mapped reads). Simple linear models were used for pairwise comparative analyses between

the 0-timepoint and depletion timepoints. An adjusted p-value threshold of 0.05 was used to

identify significantly differentially expressed genes. For principal component analysis (PCA)

and the generation of correlation matrices vst-transformed data were used within DESeq2.

Genes were labelled with their Ensembl and their common gene symbols using the

Annotation Hub and ensembldb packages (Rainer et al., 2019) . The ComplexHeatmap

package was used for the generation of expression heatmaps (Gu et al., 2016) . Volcano plots

were generated using the EnhancedVolcano package. To find enriched gene ontology terms

among differentially expressed genes, the enrichGO function of the clusterProfiler R package

and the genome-wide annotations were retrieved using the biomaRt package (Yu et al., 2012) .

ATAC-seq

100,000 nuclei were used for each reaction using the OMNI-ATAC protocol (Corces et al.,

2017) with 10,000 rat nuclei added as a spike-in control. Tagmentation was achieved using

Tn5 made in-house (Picelli et al., 2014) for 30 minutes at 37°C on a shaking block.

Immediately after incubation DNA was purified using a Zymo DNA Clean and Concentrator Kit

(Zymo Research). DNA was then eluted in 21 µL of DNAse/RNAse-free H2O. Barcoding for

library preparation was performed by PCR amplification using NEBNext® High-Fidelity 2X PCR

Master Mix (New England Biolabs) and NEBNext® index primers for Illumina sequencing.

Amplified samples were purified using AMPure SPRI beads (VWR International Ltd) and

resuspended in 25 µL of 10mM Tris-HCl pH 8.0. The Agilent 4200 TapeStation System with

D1000 ScreenTape and D1000 Reagents (Agilent) was used for quantification and

fragmentation check of the samples. Equimolar ratios of all samples (7 conditions x 2

biological replicates x 2 technical replicates each) were pooled for 300 bp paired end (2 x150

bp) sequencing performed on a NovaSeq S2 flowcell with the CRUK Cancer Institute

sequencing facility.

Raw data were trimmed to remove adapter contamination using TrimGalore v0.6.4. Reads

were aligned to the GRC38/mm10 reference genome and the Rattus norvegicus/Rnor_6.0

genome (spike-in) using the BWA aligner. File conversion, sorting, removal of duplicates and

mitochondrial reads were performed using SAMtools (Li et al., 2009) . Fragments up to 120 bp

in length, which represent nucleosome free regions, were used for further analysis. For

visualisation purposes bigwig files were generated using bamCoverage from deepTools

(Ramírez et al., 2014) . A scale factor was applied calculated as the number of uniquely

mapped rat spike-in reads for each sample divided by the number of uniquely mapped reads

in the sample with the lowest count. All replicates were pooled together into a single track

using bigWigMerge from UCSC Tools (Kent et al., 2002) . Mean profiles of the signal and

heatmaps of these regions were plotted using the ComputeMatrix, plotHeatmap and

plotProfile functions from deepTools.

Peak calling was performed for ≤120bp fragments using macs2 (Zhang et al., 2008) and

applying the -f BAMPE -q 0.05 –nolambda –keep-dup auto parameters. Peak annotation and

motif analysis was conducted using the annotatePeaks and findMotifsGenome functions

from HOMER (Heinz et al., 2010). Integration of these annotated peaks with RNA-seq gene

expression data was conducted using custom R scripts. Differential accessibility analysis was

carried out using DiffBind (https://bioconductor.org/packages/DiffBind). Differentially

accessible regions were identified in pairwise comparisons to timepoint-0 using the rat

aligned reads as a spike-in and an adjusted p-value threshold of 0.05. Chromatin state

enrichment analysis for both ADNP and CHD4 differentially accessible regions was

performed using ChromHMM (Ernst and Kellis, 2017) , using a predefined chromatin state

model generated from E14 mouse ESCs ChIP-seq data (Pintacuda et al., 2017) .

To analyse the Tn5 integration sites of the ATAC-seq data the plotFootprint() function in

VplotR (Serizay and Ahringer, 2021) was used with the addition of a code in a loop to calculate

the normalised version of the cuts by manually dividing by the library size of each merged bam

file. To analyse the fragment sizes of the ATAC-seq data that correspond to different

nucleosome structure sizes, the plotVmat() function in VplotR was used. The merged reads of

different time points of CHD4 depletion were normalised using the native libdepth+nloci

option, which normalises the reads using the library depth and the number of loci.

Cut&Run and Cut&Tag

Cut&Run and Cut&Tag were carried out as described (Janssens et al., 2022; Meers et al.,

2019). For Cut&Run 100,000 live ES cells were used per reaction, with 10,000 rat nuclei added

as a spike-in control. Cut&Tag was performed on 100,000 nuclei per reaction. Antibodies

(Table 2) were used at 1/100 dilution. pAG-MNAse and pA-Tn5 were made and purified in-

house as described (Kaya-Okur et al., 2019; Meers et al., 2019). Plasmids 3XFlag-pA-Tn5-Fl

and pAG/MNase were a gift from Steven Henikoff (Addgene plasmids #124601 and #123461,

respectively). Library preparation for sequencing was carried out in the CSCI Genomics

facility. 300 bp paired-end sequencing was performed on a NovaSeqX 25B flowcell at the

CRUK Cancer Institute sequencing facility.

Raw data were trimmed to remove adapter contamination using TrimGalore v0.6.4. Reads were aligned to the unmasked GRC38/mm10 reference genome and the Rattus norvegicus/Rnor_6.0 genome (spike-in) using the BWA. File conversion, sorting, removal of duplicates and mitochondrial reads were performed using SAMtools. For visualisation purposes bigwig files were generated using bamCoverage from deepTools. A scale factor was applied calculated as the number of uniquely mapped rat spike-in reads for each sample divided by the number of uniquely mapped reads in the sample with the lowest count. All replicates were pooled together into a single track using bigWigMerge from UCSC Tools. Mean profiles of the signal and heatmaps of these regions were plotted from these tracks using the ComputeMatrix, plotHeatmap and plotProfile functions from deepTools. Peak calling was performed using macs2 and SEACR (Meers et al., 2019) . SEACR was used in non-control mode with the stringent threshold setting (threshold = 0.01). Input BED files consisted of scaled fragment bedgraphs generated from CUT&RUN data. Scaling was performed using a factor based on the number of uniquely mapped rat spike-in reads for each sample, which was then adjusted relative to the sample with the fewest mapped reads. Motif analysis was conducted using HOMER. Differential accessibility analysis was also carried out using the DiffBind package, incorporating DESeq2 functionality. Differentially bound peaks were identified in pairwise comparisons to timepoint-0 using an adjusted p-value threshold of 0.05. Chromatin state enrichment analysis for differentially bound regions was performed using ChromHMM as for ATAC-seq.

MNase-seq

All MNase experiments, sequencing and data processing were carried out exactly as described (Bornelöv et al., 2018) except that nuclei were digested with 500 U/ml micrococcal nuclease (New England Biolabs) at 24°C for 15 minutes with shaking.

Accession numbers

ATAC-seq: E-MTAB-15037

RNAseq: E-MTAB-15102

nascent RNAseq: E-MTAB-15127

Cut&Run: E-MTAB-15606, E-MTAB-15607

Cut&Tag: E-MTAB-15625

Sall4 Depletion ATAC-seq: E-MTAB-15375

830  Sall4 Cut&Run: GSE203303 (Ru et al., 2022)

831  MNase: E-MTAB-6807 (Bornelöv et al., 2018), PRJNA1332303

833  **Acknowledgments**

834  We are grateful to Nick Owens, Vladimir Teif, Joel Mackay and past and present Hendrich

835 and Laue group members for helpful discussions; to David Klenerman and Ziwei Zhang for

836 help with imaging; to Marco Trizzino for support and to the CSCI facilities for expert technical

837 assistance.

839  **Funding**

840  This work was supported by PhD studentships from Wolfson College, Cambridge to A.K.,

841 from AstraZeneca to O.O. and from the Medical Research Council to D.S., and by grants from

842 the Medical Research Council to B.D.H. (MR/X018342/1 and MR/Y000595/1) and to E.D.L.

843 (MR/P019471/1 and MR/M010082/1), from the Wellcome Trust to E.D.L. (206291/Z/17/Z), from

844 the Isaac Newton Trust to B.D.H. (17.24(aa)), and core funding from the Wellcome/MRC

845 (203151/Z/16/Z) to the Cambridge Stem Cell Institute.

847  **Author Contributions**

848  AK, OO, IMB, DS and BH created and validated ES cell lines; AK, OO and NR generated

849 sequencing data; AK, OO and RR analysed sequencing data; DS and EDL conducted single

850 molecule imaging experiments and analyses; ML devised methodology; NR, RR, EDL and BH

851 supervised the project and EDL and BH obtained funding. The manuscript was written by BH

852 with input from all other authors.

854  **Competing Interests Statement**

855  The authors declare no competing interests.

857 **Rererences**

858 Abdulhay NJ, Hsieh LJ, McNally CP, Ostrowski MS, Moore CM, Ketavarapu M, Kasinathan S,
859 Nanda AS, Wu K, Chio US, Zhou Z, Goodarzi H, Narlikar GJ, Ramani V. 2023. Nucleosome
860 density shapes kilobase-scale regulation by a mammalian chromatin remodeler. *Nat Struct*
861 *Mol Biol* 1–11. doi:10.1038/s41594-023-01093-6

862 Ahringer J. 2000. NuRD and SIN3 histone deacetylase complexes in development. *Trends*
863  *Genet* **16**:351–356. doi:10.1016/s0168-9525(00)02066-7

Arends T, Dege C, Bortnick A, Danhorn T, Knapp JR, Jia H, Harmacek L, Fleenor CJ, Straign D, Walton K, Leach SM, Feeney AJ, Murre C, O'Connor BP, Hagman JR. 2019. CHD4 is essential for transcriptional repression and lineage progression in B lymphopoiesis. *Proc National Acad Sci* **116**:201821301. doi:10.1073/pnas.1821301116

Aughey GN, Forsberg E, Grimes K, Zhang S, Southall TD. 2023. NuRD-independent Mi-2 activity represses ectopic gene expression during neuronal maturation. *EMBO Rep* **24**:e55362. doi:10.15252/embr.202255362

Basu S, Shukron O, Hall D, Parutto P, Ponjavic A, Shah D, Boucher W, Lando D, Zhang W, Reynolds N, Sober LH, Jartseva A, Ragheb R, Ma X, Cramard J, Floyd R, Balmer J, Drury TA, Carr AR, Needham L-M, Aubert A, Communie G, Gor K, Steindel M, Morey L, Blanco E, Bartke T, Croce LD, Berger I, Schaffitzel C, Lee SF, Stevens TJ, Klenerman D, Hendrich BD, Holcman D, Laue ED. 2023. Live-cell three-dimensional single-molecule tracking reveals modulation of enhancer dynamics by NuRD. *Nat Struct Mol Biol* **30**:1628–1639. doi:10.1038/s41594-023-01095-4

Bates LE, Alves MRP, Silva JCR. 2021. Auxin-degron system identifies immediate mechanisms of OCT4. *Stem Cell Rep* **16**:1818–1831. doi:10.1016/j.stemcr.2021.05.016

Bornelöv S, Reynolds N, Xenophontos M, Gharbi S, Johnstone E, Floyd R, Ralser M, Signolet J, Loos R, Dietmann S, Bertone P, Hendrich B. 2018. The Nucleosome Remodeling and Deacetylation Complex Modulates Chromatin Structure at Sites of Active Transcription to Fine-Tune Gene Expression. *Mol Cell* **71**:56-72.e4. doi:10.1016/j.molcel.2018.06.003

Bracken AP, Brien GL, Verrijzer CP. 2019. Dangerous liaisons: interplay between SWI/SNF, NuRD, and Polycomb in chromatin regulation and cancer. *Gene Dev* **33**:936–959. doi:10.1101/gad.326066.119

Burgold T, Barber M, Kloet S, Cramard J, Gharbi S, Floyd R, Kinoshita M, Ralser M, Vermeulen M, Reynolds N, Dietmann S, Hendrich B. 2019. The Nucleosome Remodelling and Deacetylation complex suppresses transcriptional noise during lineage commitment. *EMBO J* **38**:e100788. doi:10.15252/embj.2018100788

Carr AR, Ponjavic A, Basu S, McColl J, Santos AM, Davis S, Laue ED, Klenerman D, Lee SF. 2017. Three-Dimensional Super-Resolution in Eukaryotic Cells Using the Double-Helix Point Spread Function. *Biophys J* **112**:1444–1454. doi:10.1016/j.bpj.2017.02.023

Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, Satpathy AT, Rubin AJ, Montine KS, Wu B, Kathiria A, Cho SW, Mumbach MR, Carter AC, Kasowski M, Orloff LA, Risca VI, Kundaje A, Khavari PA, Montine TJ, Greenleaf WJ, Chang HY. 2017. An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. *Nat Methods* **14**:959–962. doi:10.1038/nmeth.4396

Dieuleveult M de, Yen K, Hmitou I, Depaux A, Boussouar F, Dargham DB, Jounier S, Humbertclaude H, Ribierre F, Baulard C, Farrell NP, Park B, Keime C, Carrière L, Berlivet S, Gut M, Gut I, Werner M, Deleuze J-F, Olaso R, Aude J-C, Chantalat S, Pugh BF, Gérard M. 2016. Genome-wide nucleosome specificity and function of chromatin remodellers in ES cells. *Nature* **530**:113–116. doi:10.1038/nature16505

Dodonova SO, Zhu F, Dienemann C, Taipale J, Cramer P. 2020. Nucleosome-bound SOX2 and
SOX11 structures elucidate pioneer factor function. *Nature* **580**:669–672.
doi:10.1038/s41586-020-2195-y

Ernst J, Kellis M. 2017. Chromatin-state discovery and genome annotation with ChromHMM.
Nat Protoc **12**:2478–2492. doi:10.1038/nprot.2017.124

Etheridge TJ, Carr AM, Herbert AD. 2022. GDSC SMLM: Single-molecule localisation
microscopy software for ImageJ. *Wellcome Open Res* **7**:241.
doi:10.12688/wellcomeopenres.18327.1

Feng Q, Zhang Y. 2001. The MeCP1 complex represses transcription through preferential
binding, remodeling, and deacetylating methylated nucleosomes. *Gene Dev* **15**:827–832.
doi:10.1101/gad.876201

Gillotin S. 2018. Isolation of Chromatin-bound Proteins from Subcellular Fractions for
Biochemical Analysis. *BIO-Protoc* **8**:e3035. doi:10.21769/bioprotoc.3035

Gómez-del Arco P, Perdiguero E, Yunes-Leites PS, Acín-Pérez R, Zeini M, Garcia-Gomez A,
Sreenivasan K, Jiménez-Alcázar M, Segalés J, López-Maderuelo D, Ornés B, Jiménez-
Borreguero LJ, D'Amato G, Enshell-Seijffers D, Morgan B, Georgopoulos K, Islam ABMMK,
Braun T, de la Pompa JL, Kim J, Enriquez JA, Ballestar E, Muñoz-Cánoves P, Redondo JM.
2016. The Chromatin Remodeling Complex Chd4/NuRD Controls Striated Muscle Identity
and Metabolic Homeostasis. *Cell Metab* **23**:881–892. doi:10.1016/j.cmet.2016.04.008

Gourisankar S, Krokhotin A, Wenderski W, Crabtree GR. 2024. Context-specific functions of
chromatin remodellers in development and disease. *Nat Rev Genet* **25**:340–361.
doi:10.1038/s41576-023-00666-x

Grimm JB, English BP, Choi H, Muthusamy AK, Mehl BP, Dong P, Brown TA, Lippincott-
Schwartz J, Liu Z, Lionnet T, Lavis LD. 2016. Bright photoactivatable fluorophores for
single-molecule imaging. *Nat Methods* **13**:985–988. doi:10.1038/nmeth.4034

Gu Z, Eils R, Schlesner M. 2016. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**:2847–2849.
doi:10.1093/bioinformatics/btw313

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass
CK. 2010. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-
Regulatory Elements Required for Macrophage and B Cell Identities. *Mol Cell* **38**:576–589.
doi:10.1016/j.molcel.2010.05.004

Hendrich B, Guy J, Ramsahoye B, Wilson VA, Bird A. 2001. Closely related proteins MBD2 and
MBD3 play distinctive but interacting roles in mouse development. *Gene Dev* **15**:710–723.
doi:10.1101/gad.194101

Henikoff JG, Belsky JA, Krassovsky K, MacAlpine DM, Henikoff S. 2011. Epigenome
characterization at single base-pair resolution. *Proc National Acad Sci* **108**:18318–18323.
doi:10.1073/pnas.1110731108

Hota SK, Bruneau BG. 2016. ATP-dependent chromatin remodeling during mammalian
development. *Development* **143**:2882–2897. doi:10.1242/dev.128892

Janssens DH, Otto DJ, Meers MP, Setty M, Ahmad K, Henikoff S. 2022. CUT&Tag2for1: a
modified method for simultaneous profiling of the accessible and silenced regulome in
single cells. *Genome Biol* **23**:81. doi:10.1186/s13059-022-02642-w

Kaya-Okur HS, Wu SJ, Codomo CA, Pledger ES, Bryson TD, Henikoff JG, Ahmad K, Henikoff S.
2019. CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nat*
Commun **10**:1930. doi:10.1038/s41467-019-09982-5

Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D. 2002. The
Human Genome Browser at UCSC. *Genome Res* **12**:996–1006. doi:10.1101/gr.229102

Kim J, Lu C, Srinivasan S, Awe S, Brehm A, Fuller MT. 2017. Blocking promiscuous activation
at cryptic promoters directs cell type–specific gene expression. *Science* **356**:717–721.
doi:10.1126/science.aal3096

Klein DC, Troy K, Tripplehorn SA, Hainer SJ. 2023. The esBAF and ISWI nucleosome
remodeling complexes influence occupancy of overlapping dinucleosomes and fragile
nucleosomes in murine embryonic stem cells. *BMC Genom* **24**:201. doi:10.1186/s12864-
023-09287-4

Kloet SL, Baymaz HI, Makowski M, Groenewold V, Jansen PWTC, Berendsen M, Niazi H, Kops
GJ, Vermeulen M. 2015. Towards elucidating the stability, dynamics and architecture of
the nucleosome remodeling and deacetylase complex by using quantitative interaction
proteomics. *Febs J* **282**:1774–1785. doi:10.1111/febs.12972

Kong NR, Bassal MA, Tan HK, Kurland JV, Yong KJ, Young JJ, Yang Y, Li F, Lee JD, Liu Y, Wu C-S,
Stein A, Luo HR, Silberstein LE, Bulyk ML, Tenen DG, Chai L. 2021. Zinc Finger Protein
SALL4 Functions through an AT-Rich Motif to Regulate Gene Expression. *Cell Reports*
34:108574. doi:10.1016/j.celrep.2020.108574

Kubik S, Bruzzone MJ, Jacquet P, Falcone J-L, Rougemont J, Shore D. 2015. Nucleosome
Stability Distinguishes Two Different Promoter Types at All Protein-Coding Genes in Yeast.
Mol Cell **60**:422–434. doi:10.1016/j.molcel.2015.10.002

Kubota T, Nishimura K, Kanemaki MT, Donaldson AD. 2013. The Elg1 Replication Factor C-like
Complex Functions in PCNA Unloading during DNA Replication. *Mol Cell* **50**:273–280.
doi:10.1016/j.molcel.2013.02.012

Lai AY, Wade PA. 2011. Cancer biology and NuRD: a multifaceted chromatin remodelling
complex. *Nat Rev Cancer* **11**:588–596. doi:10.1038/nrc3091

Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, Chen X, Taipale J, Hughes TR,
Weirauch MT. 2018. The Human Transcription Factors. *Cell* **172**:650–665.
doi:10.1016/j.cell.2018.01.029

Lando D, Ma X, Cao Y, Jartseva A, Stevens TJ, Boucher W, Reynolds N, Montibus B, Hall D,
Lackner A, Ragheb R, Leeb M, Hendrich BD, Laue ED. 2024. Enhancer-promoter

interactions are reconfigured through the formation of long-range multiway hubs as mouse
ES cells exit pluripotency. *Mol Cell* **84**:1406-1421.e8. doi:10.1016/j.molcel.2024.02.015

Lauberth SM, Rauchman M. 2006. A Conserved 12-Amino Acid Motif in Sall1 Recruits the
Nucleosome Remodeling and Deacetylase Corepressor Complex. *J Biol Chem* **281**:23922–
23931. doi:10.1074/jbc.m513461200

Lenz J, Brehm A. 2023. Conserved mechanisms of NuRD function in hematopoetic gene
expression. *Enzym* 7–32. doi:10.1016/bs.enz.2023.07.006

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25**:1754–1760. doi:10.1093/bioinformatics/btp324

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Subgroup 1000 Genome Project Data Processing. 2009. The Sequence Alignment/Map
format and SAMtools. *Bioinformatics* **25**:2078–2079. doi:10.1093/bioinformatics/btp352

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**:923–930.
doi:10.1093/bioinformatics/btt656

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**:550. doi:10.1186/s13059-014-0550-8

Lu F, Lionnet T. 2021. Transcription Factor Dynamics. *Cold Spring Harb Perspect Biol*
13:a040949. doi:10.1101/cshperspect.a040949

Lu J, Jeong H, Kong N, Yang Y, Carroll J, Luo HR, Silberstein LE, YupoMa, Chai L. 2009. Stem
Cell Factor SALL4 Represses the Transcriptions of PTEN and SALL1 through an Epigenetic
Repressor Complex. *Plos One* **4**:e5577. doi:10.1371/journal.pone.0005577

Mahendrawada L, Warfield L, Donczew R, Hahn S. 2025. Low overlap of transcription factor
DNA binding and regulatory targets. *Nature* 1–9. doi:10.1038/s41586-025-08916-0

Martincorena I, Raine KM, Gerstung M, Dawson KJ, Haase K, Loo PV, Davies H, Stratton MR,
Campbell PJ. 2017. Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell*
171:1029-1041.e21. doi:10.1016/j.cell.2017.09.042

McDonel P, Costello I, Hendrich B. 2009. Keeping things quiet: Roles of NuRD and Sin3 co-
repressor complexes during mammalian development. *Int J Biochem Cell Biology* **41**:108–
116. doi:10.1016/j.biocel.2008.07.022

Meers MP, Bryson TD, Henikoff JG, Henikoff S. 2019. Improved CUT&RUN chromatin profiling
tools. *eLife* **8**:e46314. doi:10.7554/elife.46314

Miller A, Ralser M, Kloet SL, Loos R, Nishinakamura R, Bertone P, Vermeulen M, Hendrich B.
2016. Sall4 controls differentiation of pluripotent cells independently of the Nucleosome
Remodelling and Deacetylation (NuRD) complex. *Development* **143**:3074–3084.
doi:10.1242/dev.139113

Mindel V, Barkai N. 2025. On the overlap of transcription factor binding and regulatory targets:
functional and regulatory coherence of top-bound targets is masked by weakly bound
ones. *bioRxiv* 2025.10.12.681120. doi:10.1101/2025.10.12.681120

Montibus B, Ragheb R, Diamanti E, Dunn S-J, Reynolds N, Hendrich B. 2023. The Nucleosome
Remodelling and Deacetylation complex coordinates the transcriptional response to
lineage commitment in pluripotent cells. *Biol Open* **13**:bio060101. doi:10.1242/bio.060101

Morris SA, Baek S, Sung M-H, John S, Wiench M, Johnson TA, Schiltz RL, Hager GL. 2014.
Overlapping chromatin-remodeling systems collaborate genome wide at dynamic
chromatin transitions. *Nat Struct Mol Biol* **21**:73–81. doi:10.1038/nsmb.2718

Moshkin YM, Chalkley GE, Kan TW, Reddy BA, Ozgur Z, Ijcken WFJ van, Dekkers DHW,
Demmers JA, Travers AA, Verrijzer CP. 2012. Remodelers Organize Cellular Chromatin by
Counteracting Intrinsic Histone-DNA Sequence Preferences in a Class-Specific Manner.
Mol Cell Biol **32**:675–688. doi:10.1128/mcb.06365-11

Mulas C, Kalkan T, Meyenn F von, Leitch HG, Nichols J, Smith A. 2019. Defined conditions for
propagation and manipulation of mouse embryonic stem cells. *Development*
146:dev173146. doi:10.1242/dev.173146

Nabet B, Roberts JM, Buckley DL, Paulk J, Dastjerdi S, Yang A, Leggett AL, Erb MA, Lawlor MA,
Souza A, Scott TG, Vittori S, Perry JA, Qi J, Winter GE, Wong K-K, Gray NS, Bradner JE. 2018.
The dTAG system for immediate and target-specific protein degradation. *Nat Chem Biol*
14:431–441. doi:10.1038/s41589-018-0021-8

Narlikar GJ, Sundaramoorthy R, Owen-Hughes T. 2013. Mechanisms and Functions of ATP-
Dependent Chromatin-Remodeling Enzymes. *Cell* **154**:490–503.
doi:10.1016/j.cell.2013.07.011

Nishimura K, Fukagawa T, Takisawa H, Kakimoto T, Kanemaki M. 2009. An auxin-based
degron system for the rapid depletion of proteins in nonplant cells. *Nat Methods* **6**:917–
922. doi:10.1038/nmeth.1401

Nishinakamura R, Matsumoto Y, Nakao K, Nakamura K, Sato A, Copeland NG, Gilbert DJ,
Jenkins NA, Scully S, Lacey DL, Katsuki M, Asashima M, Yokota T. 2001. Murine homolog of
SALL1 is essential for ureteric bud invasion in kidney development. *Development*
128:3105–3115. doi:10.1242/dev.128.16.3105

Nocente MC, Karamitsos AM, Drouineau E, Soleil M, Albawardi W, Dulary C, Ribierre F,
Picaud H, Alibert O, Acker J, Kervella M, Aude J-C, Gilbert N, Ochsenbein F, Chantalat S,
Gérard M. 2024. cBAF generates subnucleosomes that expand OCT4 binding and function
beyond DNA motifs at enhancers. *Nat Struct Mol Biol* **31**:1756–1768. doi:10.1038/s41594-
024-01344-0

O'Shaughnessy-Kirwan A, Signolet J, Costello I, Gharbi S, Hendrich B. 2015. Constraint of
gene expression by the chromatin remodelling protein CHD4 facilitates lineage
specification. *Development* **142**:2586–2597. doi:10.1242/dev.125450

Ostapcuk V, Mohn F, Carl SH, Basters A, Hess D, Iesmantavicius V, Lampersberger L, Flemr
M, Pandey A, Thomä NH, Betschinger J, Bühler M. 2018. Activity-dependent
neuroprotective protein recruits HP1 and CHD4 to control lineage-specifying genes.
Nature **557**:739–743. doi:10.1038/s41586-018-0153-8

Pantier R, Chhatbar K, Quante T, Skourti-Stathaki K, Cholewa-Waclaw J, Alston G, Alexander-
Howden B, Lee HY, Cook AG, Spruijt CG, Vermeulen M, Selfridge J, Bird A. 2021. SALL4
controls cell fate in response to DNA base composition. *Mol Cell* **81**:845-858.e8.
doi:10.1016/j.molcel.2020.11.046

Pavani SRP, Greengard A, Piestun R. 2009. Three-dimensional localization with nanometer
accuracy using a detector-limited double-helix point spread function system. *Appl Phys
Lett* **95**:021103. doi:10.1063/1.3158923

Picelli S, Björklund ÅK, Reinius B, Sagasser S, Winberg G, Sandberg R. 2014. Tn5 transposase
and tagmentation procedures for massively scaled sequencing projects. *Genome Res*
24:2033–2040. doi:10.1101/gr.177881.114

Pierson TM, Otero MG, Grand K, Choi A, Graham JM, Young JI, Mackay JP. 2019. The NuRD
complex and macrocephaly associated neurodevelopmental disorders. *Am J Medical
Genetics Part C Seminars Medical Genetics* **181**:548–556. doi:10.1002/ajmg.c.31752

Pintacuda G, Wei G, Roustan C, Kirmizitas BA, Solcan N, Cerase A, Castello A, Mohammed S,
Moindrot B, Nesterova TB, Brockdorff N. 2017. hnRNPK Recruits PCGF3/5-PRC1 to the Xist
RNA B-Repeat to Establish Polycomb-Mediated Chromosomal Silencing. *Mol Cell* **68**:955-
969.e10. doi:10.1016/j.molcel.2017.11.013

Polo SE, Kaidi A, Baskcomb L, Galanty Y, Jackson SP. 2010. Regulation of DNA-damage
responses and cell-cycle progression by the chromatin remodelling factor CHD4. *Embo J*
29:3130–3139. doi:10.1038/emboj.2010.188

Pundhir S, Su J, Tapia M, Hansen AM, Haile JS, Hansen K, Porse BT. 2023. The impact of
SWI/SNF and NuRD inactivation on gene expression is tightly coupled with levels of RNA
polymerase II occupancy at promoters. *Genome Res* **33**:332–345.
doi:10.1101/gr.277089.122

Ragheb R, Gharbi S, Cramard J, Ogundele O, Kloet S, Burgold T, Vermeulen M, Reynolds N,
Hendrich B. 2020. Differential regulation of lineage commitment in human and mouse
primed pluripotent stem cells by the Nucleosome Remodelling and Deacetylation
Complex. *Stem Cell Res* **46**:101867. doi:10.1016/j.scr.2020.101867

Rainer J, Gatto L, Weichenberger CX. 2019. ensembldb: an R package to create and use
Ensembl-based annotation resources. *Bioinformatics* **35**:3151–3153.
doi:10.1093/bioinformatics/btz031

Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. 2014. deepTools: a flexible platform for
exploring deep-sequencing data. *Nucleic Acids Res* **42**:W187–W191.
doi:10.1093/nar/gku365

Reid XJ, Low JKK, Mackay JP. 2023. A NuRD for all seasons. *Trends Biochem Sci* **48**:11–25.
doi:10.1016/j.tibs.2022.06.002

Reid XJ, Zhong Y, Mackay JP. 2024. How does CHD4 slide nucleosomes? *Biochem Soc Trans*.
doi:10.1042/bst20230070

Reynolds N, Latos P, Hynes-Allen A, Loos R, Leaford D, O'Shaughnessy A, Mosaku O, Signolet
J, Brennecke P, Kalkan T, Costello I, Humphreys P, Mansfield W, Nakagawa K, Strouboulis
J, Behrens A, Bertone P, Hendrich B. 2012. NuRD Suppresses Pluripotency Gene
Expression to Promote Transcriptional Heterogeneity and Lineage Commitment. *Cell Stem
Cell* **10**:583–594. doi:10.1016/j.stem.2012.02.020

Ru W, Koga T, Wang X, Guo Q, Gearhart MD, Zhao S, Murphy M, Kawakami H, Corcoran D,
Zhang J, Zhu Z, Yao X, Kawakami Y, Xu C. 2022. Structural studies of SALL family protein
zinc finger cluster domains in complex with DNA reveal preferential binding to an AATA
tetranucleotide motif. *J Biol Chem* **298**:102607. doi:10.1016/j.jbc.2022.102607

Saotome M, Poduval DB, Grimm SA, Nagornyuk A, Gunarathna S, Shimbo T, Wade PA, Takaku
M. 2024. Genomic transcription factor binding site selection is edited by the chromatin
remodeling factor CHD4. *Nucleic Acids Res* **52**:3607–3622. doi:10.1093/nar/gkae025

Schep AN, Buenrostro JD, Denny SK, Schwartz K, Sherlock G, Greenleaf WJ. 2015. Structured
nucleosome fingerprints enable high-resolution mapping of chromatin architecture within
regulatory regions. *Genome Res* **25**:1757–1770. doi:10.1101/gr.192294.115

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden
C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P,
Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*
9:676–682. doi:10.1038/nmeth.2019

Serizay J, Ahringer J. 2021. Generating fragment density plots in R/Bioconductor with VplotR. *J
Open Source Softw* **6**:3009. doi:10.21105/joss.03009

Signolet J, Hendrich B. 2015. The function of chromatin modifiers in lineage commitment and
cell fate specification. *Febs J* **282**:1692–1702. doi:10.1111/febs.13132

Smeenk G, Wiegant WW, Vrolijk H, Solari AP, Pastink A, Attikum H van. 2010. The NuRD
chromatin–remodeling complex regulates signaling and repair of DNA damage. *J Cell
Biology* **190**:741–749. doi:10.1083/jcb.201001048

Sreenivasan K, Rodríguez-delaRosa A, Kim J, Mesquita D, Segalés J, Arco PG, Espejo I, Ianni A,
Croce LD, Relaix F, Redondo JM, Braun T, Serrano AL, Perdiguero E, Muñoz-Cánoves P.
2021. CHD4 ensures stem cell lineage fidelity during skeletal muscle regeneration. *Stem
Cell Rep* **16**:2089–2098. doi:10.1016/j.stemcr.2021.07.022

Srivastava D, Mahony S. 2020. Sequence and chromatin determinants of transcription factor
binding and the establishment of cell type-specific binding patterns. *Biochim Biophys Acta
(BBA) - Gene Regul Mech* **1863**:194443. doi:10.1016/j.bbagrm.2019.194443

Stevens TJ, Lando D, Basu S, Atkinson LP, Cao Y, Lee SF, Leeb M, Wohlfahrt KJ, Boucher W, O'Shaughnessy-Kirwan A, Cramard J, Faure AJ, Ralser M, Blanco E, Morey L, Sansó M, Palayret MGS, Lehner B, Croce LD, Wutz A, Hendrich B, Klenerman D, Laue ED. 2017. 3D structures of individual mammalian genomes studied by single-cell Hi-C. *Nature* **544**:59–64. doi:10.1038/nature21429

Tokunaga M, Imamoto N, Sakata-Sogawa K. 2008. Highly inclined thin illumination enables clear single-molecule imaging in cells. *Nat Methods* **5**:159–161. doi:10.1038/nmeth1171

Wade PA, Jones PL, Vermaak D, Wolffe AP. 1998. A multiple subunit Mi-2 histone deacetylase from Xenopus laevis cofractionates with an associated Snf2 superfamily ATPase. *Curr Biol* **8**:843–848. doi:10.1016/s0960-9822(98)70328-8

Watson JA, Pantier R, Jayachandran U, Chhatbar K, Alexander-Howden B, Kruusvee V, Prendecki M, Bird A, Cook AG. 2023. Structure of SALL4 zinc finger domain reveals link between AT-rich DNA binding and Okihiro syndrome. *Life Sci Alliance* **6**:e202201588. doi:10.26508/lsa.202201588

Wilczewski CM, Hepperla AJ, Shimbo T, Wasson L, Robbe ZL, Davis IJ, Wade PA, Conlon FL. 2018. CHD4 and the NuRD complex directly control cardiac sarcomere formation. *Proc National Acad Sci* **115**:201722219. doi:10.1073/pnas.1722219115

Williams CJ, Naito T, Arco PG, Seavitt JR, Cashman SM, Souza BD, Qi X, Keables P, Andrian UHV, Georgopoulos K. 2004. The Chromatin Remodeler Mi-2β Is Required for CD4 Expression and T Cell Development. *Immunity* **20**:719–733. doi:10.1016/j.immuni.2004.05.005

Wilson BG, Roberts CWM. 2011. SWI/SNF nucleosome remodellers and cancer. *Nat Rev Cancer* **11**:481–492. doi:10.1038/nrc3068

Wit E de, Nora EP. 2023. New insights into genome folding by loop extrusion from inducible degron technologies. *Nat Rev Genet* **24**:73–85. doi:10.1038/s41576-022-00530-4

Workman JL, Buchman AR. 1993. Multiple functions of nucleosomes and regulatory factors in transcription. *Trends Biochem Sci* **18**:90–95. doi:10.1016/0968-0004(93)90160-o

Wu H, Muñoz EN, Hsieh LJ, Chio US, Gourdet MA, Narlikar GJ, Cheng Y. 2023. Reorientation of INO80 on hexasomes reveals basis for mechanistic versatility. *Science* **381**:319–324. doi:10.1126/science.adf4197

Xi Y, Yao J, Chen R, Li W, He X. 2011. Nucleosome fragility reveals novel functional states of chromatin and poises genes for activation. *Genome Res* **21**:718–724. doi:10.1101/gr.117101.110

Xue Y, Wong J, Moreno GT, Young MK, Côté J, Wang W. 1998. NURD, a Novel Complex with Both ATP-Dependent Chromatin-Remodeling and Histone Deacetylase Activities. *Mol Cell* **2**:851–861. doi:10.1016/s1097-2765(00)80299-3

Yoshida T, Hu Y, Zhang Z, Emmanuel AO, Galani K, Muhire B, Snippert HJ, Williams CJ, Tolstorukov MY, Gounari F, Georgopoulos K. 2019. Chromatin restriction by the

nucleosome remodeler Mi-2β and functional interplay with lineage-specific transcription
regulators control B-cell differentiation. *Genes Dev* **33**:763–781.
doi:10.1101/gad.321901.118

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R Package for Comparing Biological
Themes Among Gene Clusters. *OMICS: A J Integr Biol* **16**:284–287.
doi:10.1089/omi.2011.0118

Zhang M, Jungblut A, Kunert F, Hauptmann L, Hoffmann T, Kolesnikova O, Metzner F, Moldt M,
Weis F, DiMaio F, Hopfner K-P, Eustermann S. 2023. Hexasome-INO80 complex reveals
structural basis of noncanonical nucleosome remodeling. *Science* **381**:313–319.
doi:10.1126/science.adf6287

Zhang W, Aubert A, Segura JMG de, Karuppasamy M, Basu S, Murthy AS, Diamante A, Drury
TA, Balmer J, Cramard J, Watson AA, Lando D, Lee SF, Palayret M, Kloet SL, Smits AH,
Deery MJ, Vermeulen M, Hendrich B, Klenerman D, Schaffitzel C, Berger I, Laue ED. 2016.
The Nucleosome Remodeling and Deacetylase Complex NuRD Is Built from Preformed
Catalytically Active Sub-modules. *J Mol Biol* **428**:2931–2942.
doi:10.1016/j.jmb.2016.04.025

Zhang Y, LeRoy G, Seelig H-P, Lane WS, Reinberg D. 1998. The Dermatomyositis-Specific
Autoantigen Mi2 Is a Component of a Complex Containing Histone Deacetylase and
Nucleosome Remodeling Activities. *Cell* **95**:279–289. doi:10.1016/s0092-8674(00)81758-4

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,
Brown M, Li W, Liu XS. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol*
9:R137. doi:10.1186/gb-2008-9-9-r137

Zhong Y, Paudel BP, Ryan DP, Low JKK, Franck C, Patel K, Bedward MJ, Torrado M, Payne RJ,
Oijen AM van, Mackay JP. 2020. CHD4 slides nucleosomes by decoupling entry- and exit-
side DNA translocation. *Nat Commun* **11**:1519. doi:10.1038/s41467-020-15183-2

Zhu F, Farnung L, Kaasinen E, Sahu B, Yin Y, Wei B, Dodonova SO, Nitta KR, Morgunova E,
Taipale M, Cramer P, Taipale J. 2018. The interaction landscape between transcription
factors and the nucleosome. *Nature* **562**:76–81. doi:10.1038/s41586-018-0549-5

+500 µM Auxin

Supplemental Figure 1. Cell cycle arrest and cell death following CHD4 depletion. Phase contrast images and example flow cytometry plots of CHD4-mAID ES cells in 2iL conditions after indicated times of Auxin addition. Cell cycle stage and relative percentage of cells are indicated on the flow cytometry plots.

1200

Supplemental Figure 2. **Schematic diagram of Vplots.** Adapted from (Henikoff et al., 2011) .
The Vplot is derived by plotting the midpoint of all recovered fragments (horizontal lines) onto
the graph, with midpoint position (white circle) on the x-axis and fragment length on the y-axis.
Red lines correspond to reads obtained from fragments with both ends (i.e. Tn5 integration
sites) in the NFR, while blue lines represent reads spanning a nucleosome. The vertical dotted
line indicates the position of the central hyperaccessible site. The inferred chromatin
structure of the locus is shown below.

1208