Rejuvenation-Responsive and Senolytic-Sensitive Muscle Stem Cells Unveiled by CD200 and CD63 in Geriatric Muscle

Ye Lynne Kim[1†], Young-Woo Jo[1†], Takwon Yoo[1†], Kyusang Yoo[1], Ji-Hoon Kim[2], Myungsun Park[3], In-Wook Song[1], Hyun Kim[1], Yea-Eun Kim[1], Sang-Hyeon Hann[1], Jong-Eun Park[3], Daehyun Baek[1], and Young-Yun Kong[1*].

[1]School of Biological Sciences, Seoul National University, 08826, Seoul, Korea.

[2]Molecular Recognition Research Center, Korea Institute of Science and Technology, Seoul, Republic of Korea.

[3]Graduate School of Medical Science and Engineering, KAIST, Daejeon, Republic of Korea.

[†] These authors contributed equally to this work.

[*] Corresponding author. Email: ykong@snu.ac.kr, Telephone: +82) 02-880-2638, FAX: +82) 02-872-1993

Author contributions:

Conceptualization, Y.L.K., Y.-W.J. and Y.-Y.K.; Data acquisition: Y.-W.J., Y.L.K., T.Y., J.-H.K. Y.-E.K., and S.-H.H.; Data analysis: Y.-W.J., Y.L.K., T.Y., and K.Y.; Technical support: T.Y., M.P., I.-W.S. J.-E.P., and D.B.; Writing: Y.L.K., Y.-W.J., T.Y., and Y.-Y.K.; Supervision: J.-E.P., D.B., Y.-Y.K.

Competing interests: Authors declare that they have no competing interests.

Abstract

Muscle stem cells (MuSCs) are parenchymal cells in skeletal muscle regeneration and maintenance. With aging, MuSCs experience a decline in their regenerative function and reduction in their number. However, recent evidence points to substantial heterogeneity within the aged MuSC population, raising questions about the underlying mechanisms of age-associated dysfunction. Here, we used Pax7^{CreERT2};Rosa^{YFP} mice (MuSC^{YFP}) to label Pax7-expressing MuSCs and chronologically traced MusCs until geriatric age. Genetic labeling and chronological tracing revealed that the number of YFP$^+$ MuSC remained comparable between young, middle and geriatric ages. At geriatric age, YFP$^+$ MuSCs exhibited reduced expression of traditional MuSC markers such as VCAM1 and PAX7. A previously unrecognized subpopulation emerged, characterized by loss of VCAM1 and low or absent PAX7. Despite their altered marker profile, these cells retained transcriptional signatures of quiescence and myogenic potential, but displayed significantly reduced proliferative and regenerative capacities. They displayed gene expression patterns indicative of senescence-like state and were selectively ablated by senolytic treatment. DHT restored regenerative function in aged mice and re-induced VCAM1 expression in YFP$^+$/Pax7$^{-/low}$/VCAM1$^-$ cells, indicating responsiveness to rejuvenation. Based on their emergence with aging, functional impairment and responsiveness to rejuvenation, we termed this population GERI-MuSCs (Geriatric Emerging Rejuvenation-responsive and Impaired MuSCs). CD63 and CD200 were identified as novel surface markers that together with VCAM1, reliably detect GERI-MuSCs as well as classical Pax7$^+$/VCAM1^{High} MuSCs, providing a tool for comprehensive isolation of MuSCs from aged wild-type mice. Together, our findings provide a refined framework for studying MuSC aging and offer new tools for isolating functionally distinct MuSC subsets from aged skeletal muscle.

Introduction

Skeletal muscle regeneration is orchestrated primarily by tissue resident stem cells, known as muscle stem cells (MuSCs) or satellite cells(Relaix et al., 2021). MuSCs reside beneath the basal lamina of myofibers in a quiescent state under homeostatic condition but become rapidly activated upon injury to proliferate, differentiate, and fuse into regenerating myofibers(Evano and Tajbakhsh, 2018). While the regenerative function of MuSCs is well maintained in youth, a decline in their number and regenerative competence is observed in aging, leading to impaired muscle repair and tissue maintenance(Blau et al., 2015).

The age-related decline in MuSC function arises from a combination of extrinsic and intrinsic factors. Extrinsic factors include the increased pro-inflammatory cytokines(Degens, 2007), extracellular matrix remodeling(Schuler et al., 2021) and altered mechanical properties of the niche microenvironment(Chakkalakal et al., 2012). MuSC intrinsic factors include disrupted metabolic homeostasis(Pala et al., 2018), loss of proteostasis(Kim et al., 2021b), altered epigenetic regulation(Hernando-Herraez et al., 2019), mitochondrial dysfunction(Rygiel et al., 2016) and aberrant activation of signaling pathways, including p38 MAPK(Bernet et al., 2014), JAK/STAT(Price et al., 2014) and mTORC1(Tang et al., 2019). As a result, aged MuSCs show reduced colony-forming capacity, failure to maintain quiescence and loss of self-renewal potential causing a significant decline in the MuSC pool. This reduction in the resident stem cell population directly undermines the muscle's capacity for effective repair(Blau et al., 2015, Day et al., 2010).

Efforts to reverse MuSC dysfunction have explored both systemic and cell-intrinsic interventions. Circulating factors identified in parabiosis experiments(Conboy et al., 2005), as well as

supplementation with molecules such as NAD$^+$(Zhang et al., 2016) and oxytocin(Elabd et al.,

2014), have shown potential in restoring regenerative capacity. Pharmacological modulation of

signaling pathways, such as inhibition of JAK/STAT(Price et al., 2014), p38 MAPK(Bernet et al.,

2014), mTORC1, TGF-β(F and al., 2021), or Wnt signaling(AS and al., 2007), has led to improved

regeneration in aged muscle. However, variation in the age of experimental animals complicates

the interpretation of these findings. Many rejuvenation studies define 'aged' mice as 12 to 24

months old. This range encompasses a period of heterogeneous functional decline, during which a

substantial fraction of MuSCs may still retain regenerative capacity. Therefore, since it is unclear

whether these models fully represent a truly geriatric state, questions remain as to whether these

interventions directly restore dysfunctional MuSCs, selectively expand a subset of aged MuSCs

that retain partial functionality, or delay or prevent the aging of MuSCs that have not yet undergone

substantial aging. To address this uncertainty, we chronologically traced the fate of Pax7-

expressing MuSCs in geriatric mice using a genetic labeling system. This allowed us to evaluate

the composition, functionality, and responsiveness of aged MuSC subsets.

In addition to interventions aimed at reversing MuSC dysfunction, an alternative strategy to restore

regenerative competence involves targeting senescent cells. Senescent cells accumulate in aged

muscle and are thought to impair regeneration through secretion of pro-inflammatory and matrix-

remodeling factors, collectively known as the senescence-associated secretory phenotype (SASP)

(MJ and al., 2023, Saito and Chikenji, 2021). Senolytic agents, such as dasatinib and quercetin,

which selectively eliminate senescent cells have been shown to improve tissue homeostasis and

extend muscle healthspan in preclinical models(Xu et al., 2018). However, although senescence is

broadly defined as a state of irreversible cell cycle arrest accompanied by transcriptional

reprogramming, how this process manifests in quiescent adult stem cells *in vivo* remains

incompletely understood. This uncertainty raises concerns regarding the use of senolytics, as such

treatment may inadvertently eliminate MuSC subsets that retain latent regenerative potential. A

better understanding of how senescence-like features emerge within the aged MuSC pool is

therefore critical for evaluating the risks and benefits of senolytic treatment in aging muscle.

In this study, our findings reveal an age-dependent accumulation of a previously unrecognized

population of MuSCs, which we termed GERI-MuSCs (Geriatric Emerging Regeneration-

capable and Impaired MuSCs). GERI-MuSCs are characterized by low expression of traditional

marker genes, impaired yet retained functionality, susceptibility to senolytic deletion, and

responsiveness to DHT rejuvenation. Moreover, CD63 and CD200 can be reliable markers that

can be used with VCAM1 to detect previously hard-to-detect MuSCs. This combinatorial

strategy provides a framework for future isolation and therapeutic targeting of GERI-MuSCs.

Together, our results refine the understanding of in vivo MuSC aging and highlight the

importance of studying stem cell subsets that are functionally impaired yet not irreversibly

senescent.

Results

Genetic labeling of Pax7$^+$ cells reveal marker-deficient MuSCs in aged muscle

Paired box protein 7 (*Pax7*) is a transcription factor widely recognized as a defining marker of

MuSCs (Sambasivan et al., 2011). To investigate how aging alters the identity of MuSCs, we used

Pax7^{CreERT2};Rosa^{YFP} mice (MuSC^{YFP}). In this model, tamoxifen administration at 3-5 months of

age irreversibly labels Pax7-expressing MuSCs. YFP$^+$ cells were isolated at young (3-5 months),

middle (10-14 months) and geriatric (>28 months) ages, using a conventional surface markers and

gating strategy: stem cell antigen-1 negative (SCA-1$^-$) and vascular cell adhesion protein 1 positive

(VCAM1$^+$) cells among lineage negative (CD31$^-$/CD45$^-$) and live cells (Liu et al., 2015) (Fig. 1A).

As previously reported, the number of VCAM1$^+$/SCA-1$^-$ MuSCs declined significantly with age,

from an average of 12.22% at young and 13.62% at middle age to 8.31% at geriatric age (Fig. 1B).

Interestingly, however, when MuSCs were defined by YFP expression alone, their number

remained comparable across ages (Fig. 1C).

To investigate the disparity in the numbers of surface marker labelled MuSCs and genetically

labelled MuSCs at geriatric age, we examined the expression of traditional MuSC surface markers,

VCAM1 and integrin alpha 7 (ITGA7) in YFP$^+$ population (Blanco-Bose et al., 2001, Sacco et al.,

2008). We observed a significant decline in VCAM1 expression, while ITGA7 expression

remained comparable (Fig. 1D). Notably, both VCAM1 and ITGA7 expression showed

significantly greater variation at geriatric age, in YFP$^+$ cells, likely due to inter-individual

variability during aging.

Since YFP$^+$ cells at geriatric age have varying marker gene expression, we investigated their

localization *in vivo* by conducting immunohistochemistry of *tibialis anterior* (TA) muscle sections

for PAX7. YFP$^+$ cells showed sublaminar localization and co-expressed PAX7 in middle aged

mice (Fig. 1E). In geriatric aged mice, however, the number of PAX7$^+$/YFP$^+$ cells decreased, while

YFP$^+$/PAX7$^{low/-}$ cells concomitantly increased (Fig. 1E).

To examine the relationship between PAX7 and varying MuSC marker expression, YFP$^+$ cells

were grouped by VCAM1 expression levels: VCAM1^{High}, VCAM1^{Low}, and VCAM1$^-$, and

immunostained for PAX7. The PAX7 fluorescence intensity was comparable between VCAM1^{High}

cells from middle-aged and geriatric mice, but was markedly reduced in VCAM1^{Low} and VCAM1$^-$

cells of geriatric mice (Fig. 1F). Together, these findings show an emerging population of YFP$^+$

cells that lose traditional markers like Pax7 and VCAM1 at geriatric age.

YFP$^+$/VCAM1$^-$ cells are quiescent MuSCs

To further explore the characteristics of YFP$^+$/VCAM1$^-$ cells in geriatric mice, we examined their

potential myogenic status. Given that loss of canonical markers might reflect lineage progression,

we assessed the expression of myogenin (MYOG), a marker of late-stage myogenic differentiation,

in YFP$^+$ cells from uninjured middle, geriatric ages and 5-days post injury. We found that MYOG

expression was largely absent in YFP$^+$ cells from uninjured aged muscle, even in geriatric

YFP$^+$/VCAM1^{Low} and YFP$^+$/VCAM1$^-$ cells (Fig. 2A-C). This contrasts with robust MYOG signal

detected at 5 days post injury, suggesting that YFP$^+$ cells are not myoblasts.

To further investigate cell identity, we performed single-cell transcriptome analysis of YFP$^+$ cells

from middle-aged and geriatric mice and integrated these data into public single-cell sequencing

data of uninjured and injured muscles.(De Micheli et al., 2020, Oprescu et al., 2020) (Fig. S1A-E).

Integration of public single-cell sequencing data yielded 5 clusters: quiescent MuSC, activated

MuSC, myoblast/myocyte, fast fiber myonuclei and slow fiber myonuclei (Fig. 2D). The

transcriptional signatures characterizing each cluster were consistent with known lineage identities,

supporting the validity of the clustering (Fig. 2E). Surprisingly, all YFP$^+$ cells from both middle

and geriatric age mapped to transcriptional clusters associated with quiescent MuSCs (Fig. 2F).

This suggests that even VCAM1^{Low} and VCAM1$^-$ cells from geriatric age did not cluster with

activated MuSCs or differentiating myoblasts or myocytes, but rather remained within the quiescent MuSC compartment. These findings support the possibility that these cells may retain core stem-like features, despite altered surface marker profiles.

Since quiescent MuSCs are generally smaller than activated progenitors due to reduced cytoplasmic volume, low metabolic activity, and compact chromatin organization(Liu et al., 2013), we also compared the cell size of VCAM1$^-$ cells with activated MuSCs from injured mice. Indeed, both VCAM1^{Low} and VCAM1$^-$ cells from geriatric mice were similar in diameter to YFP$^+$ cells from uninjured middle aged mice, but were significantly smaller than activated MuSCs from injured mice (Fig. S2A-B). These results are consistent with our interpretation that VCAM1^{Low} and VCAM1$^-$ cells from geriatric mice may remain in a quiescent state.

Given reports of smooth muscle mesenchymal cells (SMMCs) expressing overlapping surface markers with MuSCs, we considered whether the VCAM1$^-$ cells might reflect SMMC contamination(Giordani et al., 2019). mRNA sequence analysis revealed that this population exhibited low expression of SMMC markers, including Myl9, Acta2, and Pdgfrb (Fig S2C), suggesting they are transcriptionally distinct from SMMCs. Together, these results suggest that YFP$^+$/VCAM1$^-$ cells from geriatric age may retain quiescent features and myogenic identity, despite lacking typical MuSC markers.

Impaired functionality of YFP$^+$/VCAM1$^-$ MuSCs

Since YFP$^+$/VCAM1$^-$ cells retained molecular and morphological features consistent with quiescent MuSCs, we next investigated their functional potential. Colony-forming assays revealed that YFP$^+$/VCAM1^{High} cells from geriatric mice expanded at comparable rates to those from

middle-aged mice. In contrast, colony formation by YFP$^+$/VCAM1^{Low} and YFP$^+$/VCAM1$^-$ subsets

were progressively reduced, with the VCAM1$^-$ subsets forming smaller colonies (Fig. 3A-B).

These findings suggest that these cells retain proliferative ability, although their expansion

capacity was compromised.

To examine their regenerative potential *in vivo*, we transplanted equal numbers of YFP$^+$ cells from

each group into pre-injured TA of MuSC-deficient *Pax7^{CreERT2};Rosa^{DTA}* recipient mice (Fig. 3C).

In the absence of donor cells, regeneration was severely impaired, consistent with a lack of

endogenous MuSCs (Fig 3D). Transplantation of YFP$^+$/VCAM1^{High} cells from both middle-aged

and geriatric donors supported robust regeneration, resulting in centrally nucleated myofibers with

comparable morphology between age groups, consistent with heterotypic transplantation studies

(Fig. 3D-E) (Ikemoto-Uezumi et al., 2015).

Interestingly, both YFP$^+$/VCAM1^{Low} or YFP$^+$/VCAM1$^-$ cells were also capable of contributing to

myofiber regeneration, despite lacking traditional MuSC markers (Fig. 3E). However, regenerated

myofibers in these groups were significantly smaller than those derived from VCAM1^{High} cells

(Fig. 3F), suggesting that loss of VCAM1 expression may be associated with diminished

regenerative capacity.

Together, these results indicate that even the most marker-deficient MuSCs in geriatric muscle,

YFP$^+$/VCAM1$^-$ cells, retain a degree of regenerative competence. However, their reduced

proliferative output and attenuated *in vivo* contribution suggests functional heterogeneity within

the aged MuSC pool, wherein diminished marker expression may correlate with impaired

regenerative performance.

VCAM1⁻ MuSCs exhibit senescence-like features and sensitivity to senolytics

To investigate the molecular basis of impaired function in VCAM1⁻ MuSCs, we performed mRNA

sequence analysis on YFP⁺ MuSCs. Samples showed high quality and a strong correlation was

observed among biological replicates (Fig. S3A-C). Each group consistently reproduced similar

gene expression patterns in biological replicates in hierarchical clustering (Fig. S3D). In particular,

hierarchical clustering showed similar gene expression pattern between VCAM1^{High} MuSCs of

middle-aged and geriatric mice and between VCAM1^{Low} and VCAM1⁻ MuSCs (Fig. S3E),

consistent with our results of *in vivo* and *in vitro* functionality experiments (Fig. 3A-F).

Kyoto-encyclopedia of genes and genomes (KEGG) pathway enrichment analysis based on the

DEGs (Fig. S4E) (Kolberg et al., 2023) showed alterations in pathways related to cellular

senescence and PI3K-AKT pathway in VCAM1⁻ MuSCs of geriatric age as compared to

VCAM1^{High} MuSCs of middle-age (Fig. 4A). We further investigated the expression levels of

individual genes known to be altered in MuSCs of aged mice, such as genes related to extracellular

matrix (Schuler et al., 2021), Notch signaling (Liu et al., 2018, Conboy et al., 2003, Conboy et al.,

2005), and autophagy-lysosomal pathway (Garcia-Prat et al., 2016, Kim et al., 2021b), all of which

were more altered in VCAM1^{Low} and VCAM1⁻ MuSCs compared to VCAM1^{High} MuSCs of

geriatric mice (Fig. 4B). *Cdkn2a* (*p16^{INK4a}*) and *Cdkn2b* (*p15^{INK4b}*), which are cell cycle regulators

widely used as markers for senescent cells, showed highly altered expression patterns in VCAM1⁻

MuSCs compared to VCAM1^{Low} MuSCs (Fig. 4B) (Saito and Chikenji, 2021). However, the

expression levels of other well-known cellular senescence marker genes, such as *Rb1*, *Il6*, and

Trp53, were not altered much. *Pik3cb* and *Tsc1,* which are involved in the PI3K-Akt-mTOR

pathway, showed overall increased expression, while other genes in the PI3K-Akt pathway

remained relatively unchanged. Together, these results suggest a unique combination of *in vivo*

senescence-like molecular characteristics of VCAM1$^-$ MuSCs, distinct from molecular

characteristics of replicative cellular senescence.

Given their unique senescence associated transcriptome and partially elevated PI3K-Akt pathway,

we tested whether VCAM1$^-$ MuSCs were susceptible to senolytic elimination. Geriatric mice were

treated with senolytic agents dasatinib and quercetin (DQ) (Xu et al., 2018), and analyzed 2 weeks

later (Fig. 4C). Despite variation in baseline VCAM1^{High} and VCAM1$^{Low/-}$ population across aged

animals, a consistent inverse correlation was observed between VCAM1^{High} and VCAM1$^{Low/-}$

fractions (Fig. S3F). Following DQ treatment, VCAM1^{High} MuSCs remained comparable while

VCAM1$^{Low/-}$ MuSCs showed reduced abundance and lower variability (Fig. 4D). These results

indicate selective sensitivity of VCAM1$^{Low/-}$ MuSCs to senolytic agents.

Together, our findings reveal a distinct subset of quiescent MuSCs in geriatric muscle that exhibit

unique senescence-like molecular features and are amenable to senolytic clearance. We term these

cells GERI-MuSCs (Geriatric Emerging Regeneration-capable and Impaired-MuSCs).

Rejuvenation of GERI-MuSCs by DHT treatment

Studies have demonstrated that treatment with rejuvenating agents can restores myofiber

regeneration capacity in aged mice (Bernet et al., 2014, Cosgrove et al., 2014, Zhang et al., 2016,

Elabd et al., 2014, Price et al., 2014, Garcia-Prat et al., 2016). Previously, dihydrotestosterone

(DHT) supplementation has been shown to preserve MuSCs function and enhance regeneration in

antide treated mice by restoring autophagy activation (Kim et al., 2021b). Consistent with these

findings, implantation of DHT-filled silastic tube in 28-month-old geriatric mice improved muscle

regeneration, as evidenced by increased fiber cross sectional area (Fig. S4A-C).

To investigate whether DHT could affect GERI-MuSCs, we treated 28-month-old MuSC^{YFP} mice

with DHT and analyzed MuSC populations one month later (Fig. 4E). Remarkably, DHT treatment

resulted in a decrease in GERI-MuSCs with a corresponding expansion of VCAM1^{High} MuSCs

population. This phenotypic reversion suggests that DHT promotes restoration of canonical marker

expression in GERI-MuSCs.

To determine whether DHT acts through rejuvenation of GERI-MuSCs, we first depleted these

cells using DQ, then administered DHT. In mice pre-treated with DQ, DHT supplementation failed

to reconstitute the VCAM1^{High} MuSC pool (Fig. 4F). This finding suggests a model in which

GERI-MuSCs are not terminally senescent, but rather exist in a latent, hormone-responsive state.

Upon DHT exposure, these cells likely reacquire a Vcam1^{High} phenotype, underscoring their

potential as a therapeutic target in muscle aging.

Identification of GERI-MuSCs using CD63 and CD200

In non-reporter models, the detection and isolation of GERI-MuSCs pose a significant challenge,

as these cells lack expression of conventional MuSC markers such as VCAM1 and PAX7 at

geriatric age. To overcome this limitation, we conducted a transcriptomic screen to identify

alternative surface markers that remain stably expressed across the full spectrum of MuSC aging

states. Candidate genes were selected based on three criteria; high transcript abundance (RPKM),

consistent expression from VCAM1^{High} to VCAM1^{-} MuSCs, and predicted surface localization.

CD63 and CD200 sufficed all three criteria and were therefore chosen as candidate markers for

further validation (Fig. 5B).

We confirmed protein level expression of CD63 and CD200 expression in MuSCs by we co-

immunostaining TA cross-sections for CD63, CD200, YFP and PAX7. Both markers co-localized

with YFP and PAX7 beneath the basal lamina (Fig. 5C, D). Flow cytometry analysis further

demonstrated that 96.3% of $YFP^+/VCAM1^+$ cells from young $MuSC^{YFP}$ mice were

$CD63^{high}/CD200^{high}$ (Fig. 5E). Notably, among lineage-negative ($CD31^-/CD45^-$), $SCA-1^-$, and live

cells, 93.2% of $CD63^{high}/CD200^{high}$ cells were $YFP^+/VCAM1^+$ cells (Fig. 5F). Importantly,

96.5±2.8% of sorted $CD63^{high}/CD200^{high}$ cells were $PAX7^+$, validating these markers as a robust

strategy for MuSC enrichment.

To assess whether the CD63/CD200-based strategy could detect and isolate GERI-MuSCs in wild-

type mice, we applied the strategy to middle and geriatric aged mice. In middle aged mice, all

$CD63^{high}/CD200^{high}$ cells corresponded exclusively to $VCAM1^{High}$ MuSCs (Fig. 5H). In contrast,

geriatric mice exhibited a marked shift, with the majority of $CD63^{high}/CD200^{high}$ cells falling

within the $VCAM1^{Low/-}$ gate (Fig. 5H). These results indicate that CD63 and CD200 expression is

preserved in GERI-MuSCs, enabling their prospective identification and isolation in the absence

of genetic labeling. Therefore, these findings establish CD63 and CD200 as reliable surface

markers that can be used with VCAM1. Together, they can to detect the full spectrum of MuSCs,

including functionally latent populations that evade detection by conventional means. This strategy

offers a critical tool for future studies aiming to comprehensively characterize MuSC heterogeneity

and therapeutic responsiveness in the context of aging.

Discussion

Our study challenges the longstanding paradigm that muscle stem cell (MuSC) number declines irreversibly with age. While previous reports have described significant depletion of Pax7[+] MuSCs in aged muscle, these findings were largely based on immunohistochemical detection or flow cytometric profiling using conventional surface markers. Here, we demonstrate that genetically labeled MuSCs, tracked longitudinally *in vivo* using a Pax7-CreERT2;Rosa-YFP model, do not decrease in number even in geriatric mice. Instead, a substantial subset of aged MuSCs loses expression of canonical markers such as Pax7 and VCAM1, rendering them undetectable by traditional approaches.

We identify this overlooked population as GERI-MuSCs (Geriatric Emerging Regeneration-capable and Impaired MuSCs). Despite their diminished expression of hallmark MuSC markers, GERI-MuSCs exhibit features consistent with quiescence, rather than terminal differentiation or apoptosis. Transcriptomic profiling and functional assays reveal that these cells retain a partial capacity for proliferation and regeneration, albeit at reduced efficiency. These findings suggest that previous conclusions about MuSC depletion may have reflected loss of marker expression rather than true cellular loss.

GERI-MuSCs display a unique gene expression profile that includes upregulation of senescence-associated transcripts such as Cdkn2a but unaltered expression of other typical senescence-associated genes, such as Il6 and Mmp3, while showing preservation of certain stemness-related pathways. These cells do not fulfill the classical definition of cellular senescence, such as irreversible cell cycle arrest. Rather, GERI-MuSCs exhibit a hybrid state—retaining regenerative potential while expressing select markers traditionally associated with senescence. This finding

suggests that the criteria used to define senescence in proliferative cell types may not fully apply to quiescent stem cells like MuSCs, which spend most of their life in a non-proliferative state. Further studies are warranted to delineate *in vivo* senescence features specific to long-lived, tissue-resident stem cells.

Our results suggest that GERI-MuSCs could be functionally reactivated. Treatment with dihydrotestosterone (DHT) restored VCAM1 expression in GERI MuSCs and regenerative performance in aged mice, suggesting that their phenotype is reversible and hormonally responsive. This provides a compelling explanation for the efficacy of hormone-based interventions in aging muscle and highlights the potential for targeting latent regenerative capacity in aged stem cells. Notch signaling appears to be a key regulator of this reversible phenotype. We found diminished expression of Notch pathway components—including Notch1, Notch3, Hey1, and HeyL—in VCAM1Low and VCAM1– subsets, along with reduced transcription of Pax7 and VCAM1. Given that both Pax7 and VCAM1 are downstream targets of Notch signaling and that Notch activity can be modulated by androgen signaling, the restoration of MuSC function by DHT may act through this axis. Testing such mechanisms in future studies will be important for establishing therapeutic strategies.

The presence of GERI-MuSCs also raises caution about the use of senolytic agents. Although dasatinib and quercetin are used to clear senescent cells and have shown benefits in aged tissue, our data show that these agents also deplete GERI-MuSCs. Since GERI-MuSCs retain functionality and regenerative potential, their removal could compromise muscle maintenance and repair in elderly individuals. Given the critical role of MuSCs in preserving muscle mass,

supporting myonuclei turnover, and maintaining neuromuscular junctions, the long-term

application of senolytics in skeletal muscle warrants careful scrutiny.

Finally, we propose CD63 and CD200 as reliable surface markers that can be used with VCAM1

for identifying MuSCs across age groups. These markers remain expressed in both young and aged

MuSCs, including those with diminished canonical marker expression. Their use enables

comprehensive isolation of the full MuSC population and will be instrumental in improving the

resolution of future studies on stem cell heterogeneity and aging.

In conclusion, we identify a previously unrecognized population of MuSCs—GERI-MuSCs—that

persists with age despite loss of conventional markers. These cells remain quiescent, partially

functional, and responsive to rejuvenation cues. Their discovery challenges existing models of

MuSC depletion and senescence, and underscores the importance of revisiting stem cell aging with

improved detection strategies and nuanced interpretations of cellular function.

Materials and Methods

List of Antibodies for Flow Cytometry and FACS (Fluorescence-activated cell sorting

Marker	Fluorophore	Catalog No.	Company
CD31	APC	102510	BioLegend
CD31	Pacific Blue™	108120	BioLegend
CD45	APC	103112	BioLegend
CD45	Pacific Blue™	405206	BioLegend
VCAM1	Biotin	553331	BD Biosciences
Streptavidin	PE/Cy7	405206	BioLegend
Ly-6A/E (Sca-1)	Pacific Blue™	108120	BioLegend
Ly-6A/E (Sca-1)	FITC	553335	BD Biosciences
CD200	BV711	745548	BD Biosciences
CD63	APC	170631-82	eBioscience

Animals

All animal procedures conformed to the guidelines of Institutional Animal Care and Use Committee (IACUC) of Seoul National University, Korea. Male C57BL/6J mice were housed under a 12-h light/12-h dark cycle at 22°C with 40-60% humidity, and fed ad libitum. Pax7^{CreER} (017763), Rosa^{YFP} (006148), Rosa^{DTA} (009669) mice were obtained from the Jackson Laboratory (Bar, Harbor, ME, USA). Tamoxifen (25mg/mL in corn oil; Sigma-Aldrich, St. Louis, MO, USA) was administered orally for 5 consecutive days (240 mg/kg). For muscle injury, mice were anesthetized with avertin (250 μg/g, i.p.) and injected with 60 μL of 1.2% BaCl2 (Sigma-Aldrich)

into the Tibialis Anterior (TA) muscles. One-day post-injury, mice were anesthetized with avertin

and MuSCs were transplanted at indicated numbers.

Flow Cytometry and FACS (Fluorescence-activated cell sorting)

Limb muscles were mechanically dissected and enzymatically dissociated in Dulbecco's modified

Eagle's medium (DMEM) containing 10% horse serum (Hyclone, Logan, UT, USA), collagenase

type II (800U/mL; Worthington, Lakewood, NJ, USA), and dispase (1.1 U/mL; StemCell

Technologies, Vancouver, BC, Canada) at 37°C for 50 minutes. Digested suspensions were

triturated, washed with DMEM and filtered to obtain mononuclear cells. Cells were stained with

antibodies, analyzed and sorted using a FACS Aria III or FACS Aria Fusion (BD Bioscience). List

of antibodies used are provided as supplementary materials. Representative plots were processed

with FlowJo v10.9.

MuSC Culture

FACS-isolated MuSCs were cultured in DMEM/F12 + 20% FBS + 1% ZellShield (Minerva

Biolabs, Germany) + 2.5 ng/mL bFGF (F0291, Sigma Aldrich) on Matrigel (256231, BD

Bioscience)-coated plates at 37°C and 5% CO_2. Medium was refreshed every 3 days. MuSCs were

imaged daily using EVOS FL Auto 2 (Thermo Fisher, USA) and the number of cells in the same

region was counted.

Histochemistry and Immunohistochemistry

TA muscles were fixed overnight in 4% paraformaldehyde, embedded in the Tissue Tek OCT

compound (Sakura Fineteck, USA), and snap-frozen in liquid nitrogen. The muscles were

sectioned at 7μm and stained on the slides. H&E staining was performed according to basic

protocols. For immunostaining, sections were subjected to epitope retrieval in 50mM Tris-EDTA

(pH 9.0) at 105°C for 10 minutes, then blocked with 3% BSA + 0.1% Triton X-100. Sections were

incubated overnight with primary antibodies, mouse anti-PAX7 (DSHB, Pax7, 1:100 dilution),

chicken anti-GFP (abcam, ab13970, 1:300), rabbit anti-laminin (Sigma, L9393-2ML, 1:500), rat

anti-CD200 (BD Biosciences, 1:50), and rat anti-CD63 (eBioscience, 1:50). After washing,

sections were incubated with secondary antibodies, Alexa Fluor 647-conjugated anti-rat or anti

rabbit IgG, Alexa Fluor 594-conjugated anti mouse IgG, and Alexa Fluor 488-conjugated anti

chicken IgG. Nuclei were counterstained with Hoechst 33342 (Invitrogen, H3570). Slides were

mounted with VECTASHIELD (Vector Laboratories) and images were acquired using EVOS FL

Auto 2 (Thermo Fisher) and Leica TCS SP8 confocal microscope. Cross-sectional area (CSA) was

calculated by SMASH[23] and ImageJ programs, respectively.

Immunocytochemistry

Isolated MuSCs were cytospun onto slides (Cytospin 4, Thermo Scientific), fixed with 4%

paraformaldehyde for 10 minutes, washed with PBS and permeabilized with PBS containing 0.4%

Triton X-100 for 30 minutes. Slides were then incubated with primary antibodies, mouse anti-

PAX7 (DSHB) and mouse anti-MYOG (DSHB, F5D, 1:100 dilution) in blocking buffer.

Subsequent steps followed the immunohistochemistry protocol. Relative fluorescent intensity was

quantified using ImageJ.

Senolytics and DHT treatment

Dihydrotestosterone (DHT) implantation was performed as previously described[8]. Senolytic

agents Dasatinib (D-3307, LC laboratories, 5mg/kg) and Quercetin (Q4951, Sigma, 50 mg/kg) was

administered orally for 5 consecutive days. Experimental schemes of combined treatment are as

mentioned in the results section of this article.

Single-cell RNA sequence analysis

To obtain gene-specific UMI (Unique Molecular Identifier) counts in each cell, we processed raw

scRNA-seq data using the BD Rhapsody WTA analysis pipeline (v1.11.1)[24]. Scanpy (v1.9.3)[25]

was utilized for single-cell RNA sequence analysis. Cells with a gene count below 200 or a

mitochondrial gene percentage above 20% were considered low-quality cells and were excluded

from downstream analyses. We set the upper limit of each UMI count as (mean(total counts) + 2

standard deviation(total counts) value) to remove doublets. Normalization was applied to the count

matrix (scanpy.pp.normalize_total with target_sum=10^4 and scanpy.pp.log1p).

We characterized the enriched cells by merging our scRNA-seq data with public datasets

(GSE162172, GSE138826)[26,27]. From the public dataset, only the cells belonging to the

myogenic lineage (from quiescent MuSCs to Myonuclei (Type IIb or IIx)) were selected for

comparison. After merging the public dataset with our dataset

(scanpy.external.pp.harmony_integrate with max_iter_harmony=50), 2000 highly variable genes

were identified and scaled (scanpy.pp.highly_variable_genes with n_top_genes=2000,

scanpy.pp.scale with max_value=10).

We visualized the scRNA-seq data using uniform manifold approximation and projection (UMAP)

(scanpy.tl.umap) with a precomputed neighborhood graph (scanpy.pp.neighbors with n_pcs=50).

We clustered the UMAP-transformed data using the Louvain algorithm (scanpy.tl.louvain,

resolution=0.5). Each cluster was annotated using cell-specific markers (Fig. 1D, E). The

proportions of quiescent MuSCs, activated MuSCs, and Myoblasts/Progenitors were measured in

our samples (Mid and Geri) and the public dataset across various days-post-injury (DPI). Cases

with less than 50 cells per DPI were excluded during this analysis (Fig. S1A).

We utilized diffusion map, PAGA, and force-directed graph to analyze the trajectory of cells

belonging to the myogenic lineage. After dimension reduction using diffusion maps

(scanpy.tl.diffmap), we computed neighborhood graphs (scanpy.pp.neighbors) and abstracted the

graphs using PAGA (scanpy.tl.paga). We modeled the trajectory of the cells using the abstracted

graphs as initial positions by constructing a force-directed graph (Fig. S1B-E).

Representative Isoforms of Mouse mRNAs

To acquire a sole representative isoform for each gene, we utilized the filtering pipeline described

in a previous study(Kim et al., 2016). The mouse reference annotation (GRCm39 refFlat) was

downloaded from the UCSC website(Pruitt et al., 2007). mRNA sequences with accession

category 'NM' located on the 21 chromosomes (1-19, X, and Y) were selected for downstream

analyses. Entries with incorrect start or stop codons, internal stop codons, and ORF lengths not

divisible by 3, were removed. Nonsense-mediated mRNA decay candidates, defined as the

mRNAs with stop codons positioned less than 50 nucleotides before the final exon-exon junction,

were excluded (Nagy and Maquat, 1998). A total of 20,582 non-redundant representative isoforms

were selected as mouse reference mRNAs for subsequent analyses.

mRNA sequence analysis

We trimmed low-quality read ends using a custom algorithm. Each base position was assigned a

reward (+5) or a penalty (-1) based on its Phred score with a threshold of 30. To define high-quality

read ends, we calculated the cumulative sum from both ends, identified the positions with the

minimum value, and trimmed the reads at those positions. This process was repeated iteratively

until no further trimming was necessary. Reads shorter than 17 bases after trimming were excluded.

Subsequently, we removed adapter sequences using Cutadapt (v3.1) (Martin, 2011) with options

(-O 10 -m 18 --match-read-wildcards). We eliminated sequencing artifacts and PCR duplicates

using FASTX-Toolkit's fastx_artifact_filter with option (-Q 33) (hannonlab.cshl.edu/fastx_toolkit/,

v0.0.13). To eliminate reads derived from mouse non-coding RNA, we aligned the reads to non-

coding RNA sequences in RNAcentral(Consortium, 2021) using Bowtie2 (v2.1.0) (Langmead and

Salzberg, 2012) with options (-k 1 --norc --very-sensitive), and the reads aligned to non-coding

RNAs were excluded. Unpaired reads were also excluded.

Preprocessed reads were aligned to the mouse genome (GRCm39) utilizing the STAR alignment

software (v2.7.5b) (Dobin et al., 2013). The parameters used for the STAR alignment are described

in a previous study (Kim et al., 2021a). The reads were sorted and indexed using Samtools (v0.1.19)

(Danecek et al., 2021). The expression of each gene was quantified by calculating the fragment

per kilobase of transcript per million mapped reads (FPKM).

We applied the same analysis pipeline to both our experimental samples (Mid, Vh, Vl, V-) and

publicly available SMMC samples (GSE110878, SRR67542132-SRR67542134)(Giordani et al.,

2019) to calculate gene expression levels. Quantile-normalized expression levels were compared

between the two datasets (Fig. S2C).

The following analyses were performed exclusively on our experimental samples.

To assess the quality of our samples, the proportion of reads that were either removed, unmapped,

or successfully mapped was computed for each sample throughout the preprocessing and

alignment steps (Fig. S3B). The correlation between replicates was quantified using Spearman's

R (Fig. S3C).

We normalized all samples using the following procedure to ensure a fair inter-sample comparison.

We applied quantile normalization to equalize the FPKM distribution of all samples. To remove

individual mouse variations, we used the RUVs function (k=2) from the RUVSeq package (v1.34.0)

(Risso et al., 2014). Subsequently, genes with normalized expression higher than the 50^{th} percentile

were selected, and the expression values were z-normalized. We clustered the samples using the

clustermap function of the Seaborn package (v0.11.1) (Waskom, 2021) with correlation metric

(Fig. S3D).

We identified differentially expressed genes (DEGs) by comparing the expression between the

Mid, V^H, V^L, and V^- samples. We used DESeq2 (v1.40.2) (Love et al., 2014) to compute $\log_2$ fold

changes and false discovery rate (FDR) corrected Q values for individual genes. DEGs were

defined as genes with |$\log_2$ fold change| > 1.2 and FDR-corrected Q value < 0.05 (Fig. S3E).

To annotate the pathways linked to DEGs, we performed enrichment analysis for Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathways using the g:Profiler web server (Kolberg

et al., 2023, Kanehisa et al., 2016). Among the highly expressed 12,000 genes, DEGs were queried

using the g:GOst tool, and the g:SCS algorithm was employed for multiple testing correction.

Pathways with multiple testing corrected Q value < 0.05 were presented (Fig. 3A).

Statistical Analysis

All statistical analyses were performed using R (R-4.3.1) and GraphPad Prism. All variables were

confirmed to have a normal distribution by using the Shapiro-Wilk test and Kolmogorov-Smirnov

test. For comparison of significant differences in 2 groups of normally distributed data, Levene's

test for homogeneity of variance using the car package followed by a 2-tailed Student's t test or

Welch's t test was used. For comparison of multiple groups, one-way analysis of variance

(ANOVA) or two-way ANOVA followed by Bonferroni's multiple comparison test was

performed. All data are presented as mean ± s.d. n.s., not significant; $P > 0.05$; *$P < 0.05$; **$P < 0.01$;

***$P < 0.001$, ****$P < 0.0001$. *P value* of $<.05$ considered statistically significant at the 95%

confidence level. The number of biological replicates, p values, and detailed statistical method for

each data were indicated in the figure legends.

Acknowledgments

We thank C.Kang (Seoul National University) for providing senolytic agents, Dasatinib and

Quercetin.

Funding

This work was supported by the National Research Foundation of Korea (NRF) grant funded by

the Korea government (MIST) (RS-2020-NR049538)

Data and materials availability: The bulk RNA sequencing data and scRNA-seq data have been

deposited at Gene Expression Omnibus (GEO, GSE accession number: GSE254475 and

GSE254476, respectively) and are publicly available as of the date of publication. Source data

supporting the findings of this study are provided with this paper. Additional data that support the

findings of this study are available from the corresponding author upon reasonable request.

References

References

AS, B. & AL., E. 2007. Increased Wnt signaling during aging alters muscle stem cell fate and
increases fibrosis. *Science,* 317**,** 807–810.

BERNET, J. D., DOLES, J. D., HALL, J. K., KELLY TANAKA, K., CARTER, T. A. & OLWIN,
B. B. 2014. p38 MAPK signaling underlies a cell-autonomous loss of stem cell self-renewal in
skeletal muscle of aged mice. *Nat Med,* 20**,** 265-71.

BLANCO-BOSE, W. E., YAO, C. C., KRAMER, R. H. & BLAU, H. M. 2001. Purification of
mouse primary myoblasts based on alpha 7 integrin expression. *Exp Cell Res,* 265**,** 212-20.

BLAU, H. M., COSGROVE, B. D. & HO, A. T. 2015. The central role of muscle stem cells in
regenerative failure with aging. *Nat Med,* 21**,** 854-62.

CHAKKALAKAL, J. V., JONES, K. M., BASSON, M. A. & BRACK, A. S. 2012. The aged niche
disrupts muscle stem cell quiescence. *Nature,* 490**,** 355-60.

CONBOY, I. M., CONBOY, M. J., SMYTHE, G. M. & RANDO, T. A. 2003. Notch-mediated
restoration of regenerative potential to aged muscle. *Science,* 302**,** 1575-7.

CONBOY, I. M., CONBOY, M. J., WAGERS, A. J., GIRMA, E. R., WEISSMAN, I. L. &
RANDO, T. A. 2005. Rejuvenation of aged progenitor cells by exposure to a young systemic
environment. *Nature,* 433**,** 760-4.

CONSORTIUM, T. R. 2021. RNAcentral 2019: a hub of information for non-coding RNA
sequences. *Nucleic Acids Res,* 49**,** D212-D220.

COSGROVE, B. D., GILBERT, P. M., PORPIGLIA, E., MOURKIOTI, F., LEE, S. P., CORBEL,
S. Y., LLEWELLYN, M. E., DELP, S. L. & BLAU, H. M. 2014. Rejuvenation of the muscle stem
cell population restores strength to injured aged muscles. *Nat Med,* 20**,** 255-64.

DANECEK, P., BONFIELD, J. K., LIDDLE, J., MARSHALL, J., OHAN, V., POLLARD, M. O.,
WHITWHAM, A., KEANE, T., MCCARTHY, S. A., DAVIES, R. M. & LI, H. 2021. Twelve
years of SAMtools and BCFtools. *Gigascience,* 10.

DAY, K., SHEFER, G., SHEARER, A. & YABLONKA-REUVENI, Z. 2010. The depletion of
skeletal muscle satellite cells with age is concomitant with reduced capacity of single progenitors
to produce reserve progeny. *Dev Biol,* 340**,** 330-43.

DE MICHELI, A. J., LAURILLIARD, E. J., HEINKE, C. L., RAVICHANDRAN, H., FRACZEK,
P., SOUEID-BAUMGARTEN, S., DE VLAMINCK, I., ELEMENTO, O. & COSGROVE, B. D.
2020. Single-Cell Analysis of the Muscle Stem Cell Hierarchy Identifies Heterotypic
Communication Signals Involved in Skeletal Muscle Regeneration. *Cell Rep,* 30**,** 3583-3595 e5.

DEGENS, H. 2007. The role of systemic inflammation in age-related muscle wasting and
reparation. *J Physiol,* 583**,** 343--344.

DOBIN, A., DAVIS, C. A., SCHLESINGER, F., DRENKOW, J., ZALESKI, C., JHA, S., BATUT,
P., CHAISSON, M. & GINGERAS, T. R. 2013. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics, 29**,** 15-21.

ELABD, C., COUSIN, W., UPADHYAYULA, P., CHEN, R. Y., CHOOLJIAN, M. S., LI, J.,
KUNG, S., JIANG, K. P. & CONBOY, I. M. 2014. Oxytocin is an age-specific circulating
hormone that is necessary for muscle maintenance and regeneration. *Nat Commun,* 5**,** 4082.

EVANO, B. & TAJBAKHSH, S. 2018. Skeletal muscle stem cells in comfort and stress. *NPJ*
Regen Med, 3**,** 24.

F, G. & AL., E. 2021. TGFβ signaling curbs cell fusion and muscle regeneration. *Nature*
Communications, 12**,** 3131.

GARCIA-PRAT, L., MARTINEZ-VICENTE, M., PERDIGUERO, E., ORTET, L.,
RODRIGUEZ-UBREVA, J., REBOLLO, E., RUIZ-BONILLA, V., GUTARRA, S.,
BALLESTAR, E., SERRANO, A. L., SANDRI, M. & MUNOZ-CANOVES, P. 2016. Autophagy
maintains stemness by preventing senescence. *Nature,* 529**,** 37-42.

GIORDANI, L., HE, G. J., NEGRONI, E., SAKAI, H., LAW, J. Y. C., SIU, M. M., WAN, R.,
CORNEAU, A., TAJBAKHSH, S., CHEUNG, T. H. & LE GRAND, F. 2019. High-Dimensional
Single-Cell Cartography Reveals Novel Skeletal Muscle-Resident Cell Populations. *Mol Cell,* 74**,**
609-621 e6.

HERNANDO-HERRAEZ, I., EVANO, B., STUBBS, T., COMMERE, P. H., JAN BONDER, M.,
CLARK, S., ANDREWS, S., TAJBAKHSH, S. & REIK, W. 2019. Ageing affects DNA
methylation drift and transcriptional cell-to-cell variability in mouse muscle stem cells. *Nat*
Commun, 10**,** 4361.

IKEMOTO-UEZUMI, M., UEZUMI, A., TSUCHIDA, K., FUKADA, S., YAMAMOTO, H.,
YAMAMOTO, N., SHIOMI, K. & HASHIMOTO, N. 2015. Pro-Insulin-Like Growth Factor-II
Ameliorates Age-Related Inefficient Regenerative Response by Orchestrating Self-Reinforcement
Mechanism of Muscle Regeneration. *Stem Cells,* 33**,** 2456-68.

KANEHISA, M., SATO, Y., KAWASHIMA, M., FURUMICHI, M. & TANABE, M. 2016.
KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res,* 44**,** D457-62.

KIM, D., KIM, S., PARK, J., CHANG, H. R., CHANG, J., AHN, J., PARK, H., PARK, J., SON,
N., KANG, G., KIM, J., KIM, K., PARK, M. S., KIM, Y. K. & BAEK, D. 2021a. A high-resolution
temporal atlas of the SARS-CoV-2 translatome and transcriptome. *Nat Commun,* 12**,** 5120.

KIM, D., SUNG, Y. M., PARK, J., KIM, S., KIM, J., PARK, J., HA, H., BAE, J. Y., KIM, S. &
BAEK, D. 2016. General rules for functional microRNA targeting. *Nat Genet,* 48**,** 1517-1526.

KIM, J. H., PARK, I, SHIN, H. R., RHEE, J., SEO, J. Y., JO, Y. W., YOO, K., HANN, S. H.,
KANG, J. S., PARK, J., KIM, Y. L., MOON, J. Y., CHOI, M. H. & KONG, Y. Y. 2021b. The
hypothalamic-pituitary-gonadal axis controls muscle stem cell senescence through autophagosome
clearance. *J Cachexia Sarcopenia Muscle,* 12**,** 177-191.

KOLBERG, L., RAUDVERE, U., KUZMIN, I., ADLER, P., VILO, J. & PETERSON, H. 2023.
g:Profiler-interoperable web service for functional enrichment analysis and gene identifier
mapping (2023 update). *Nucleic Acids Res,* 51**,** W207-W212.

LANGMEAD, B. & SALZBERG, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat
Methods,* 9**,** 357-9.

LIU, L., CHARVILLE, G. W., CHEUNG, T. H., YOO, B., SANTOS, P. J., SCHROEDER, M. &
RANDO, T. A. 2018. Impaired Notch Signaling Leads to a Decrease in p53 Activity and Mitotic
Catastrophe in Aged Muscle Stem Cells. *Cell Stem Cell,* 23**,** 544-556 e4.

LIU, L., CHEUNG, T. H., CHARVILLE, G. W., HURGO, B. M., LEAVITT, T., SHIH, J.,
BRUNET, A. & RANDO, T. A. 2013. Chromatin modifications as determinants of muscle stem
cell quiescence and chronological aging. *Cell Rep,* 4**,** 189-204.

LIU, L., CHEUNG, T. H., CHARVILLE, G. W. & RANDO, T. A. 2015. Isolation of skeletal
muscle stem cells by fluorescence-activated cell sorting. *Nat Protoc,* 10**,** 1612-24.

LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.

MARTIN, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011, 17**,** 3.

MJ, S. & AL., E. 2023. Senescence atlas reveals an aged-like inflamed niche that blunts muscle
regeneration. *Nature,* 613**,** 169–178.

NAGY, E. & MAQUAT, L. E. 1998. A rule for termination-codon position within intron-
containing genes: when nonsense affects RNA abundance. *Trends Biochem Sci,* 23**,** 198-9.

OPRESCU, S. N., YUE, F., QIU, J., BRITO, L. F. & KUANG, S. 2020. Temporal Dynamics and
Heterogeneity of Cell Populations during Skeletal Muscle Regeneration. *iScience,* 23**,** 100993.

PALA, F., DI GIROLAMO, D., MELLA, S., YENNEK, S., CHATRE, L., RICCHETTI, M. &
TAJBAKHSH, S. 2018. Distinct metabolic states govern skeletal muscle stem cell fates during
prenatal and postnatal myogenesis. *J Cell Sci,* 131.

PRICE, F. D., VON MALTZAHN, J., BENTZINGER, C. F., DUMONT, N. A., YIN, H., CHANG,
N. C., WILSON, D. H., FRENETTE, J. & RUDNICKI, M. A. 2014. Inhibition of JAK-STAT
signaling stimulates adult satellite cell function. *Nat Med,* 20**,** 1174-81.

PRUITT, K. D., TATUSOVA, T. & MAGLOTT, D. R. 2007. NCBI reference sequences (RefSeq):
a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids*
Res, 35**,** D61-5.

RELAIX, F., BENCZE, M., BOROK, M. J., DER VARTANIAN, A., GATTAZZO, F.,
MADEMTZOGLOU, D., PEREZ-DIAZ, S., PROLA, A., REYES-FERNANDEZ, P. C., ROTINI,
A. & TAGLIETTI, T. 2021. Perspectives on skeletal muscle stem cells. *Nat Commun,* 12**,** 692.

RISSO, D., NGAI, J., SPEED, T. P. & DUDOIT, S. 2014. Normalization of RNA-seq data using
factor analysis of control genes or samples. *Nat Biotechnol,* 32**,** 896-902.

RYGIEL, K., PICARD, M. & TURNBULL, D. 2016. The ageing neuromuscular system and
sarcopenia: a mitochondrial perspective. *J Physiol,* 594**,** 4499--4512.

SACCO, A., DOYONNAS, R., KRAFT, P., VITOROVIC, S. & BLAU, H. M. 2008. Self-renewal
and expansion of single transplanted muscle stem cells. *Nature,* 456**,** 502-6.

SAITO, Y. & CHIKENJI, T. S. 2021. Diverse Roles of Cellular Senescence in Skeletal Muscle
Inflammation, Regeneration, and Therapeutics. *Front Pharmacol,* 12**,** 739510.

SAMBASIVAN, R., YAO, R., KISSENPFENNIG, A., VAN WITTENBERGHE, L., PALDI, A.,
GAYRAUD-MOREL, B., GUENOU, H., MALISSEN, B., TAJBAKHSH, S. & GALY, A. 2011.
Pax7-expressing satellite cells are indispensable for adult skeletal muscle regeneration.
Development, 138**,** 3647-56.

SCHULER, S. C., KIRKPATRICK, J. M., SCHMIDT, M., SANTINHA, D., KOCH, P., DI
SANZO, S., CIRRI, E., HEMBERG, M., ORI, A. & VON MALTZAHN, J. 2021. Extensive
remodeling of the extracellular matrix during aging contributes to age-dependent impairments of
muscle stem cell functionality. *Cell Rep,* 35**,** 109223.

TANG, H., INOKI, K., BROOKS, S. V., OKAZAWA, H., LEE, M., WANG, J., KIM, M.,
KENNEDY, C. L., MACPHERSON, P. C. D., JI, X., VAN ROEKEL, S., FRAGA, D. A., WANG,
K., ZHU, J., WANG, Y., SHARP, Z. D., MILLER, R. A., RANDO, T. A., GOLDMAN, D.,
GUAN, K. L. & SHRAGER, J. B. 2019. mTORC1 underlies age-related muscle fiber damage and
loss by inducing oxidative stress and catabolism. *Aging Cell,* 18**,** e12943.

WASKOM, M. L. 2021. seaborn: statistical data visualization. *JOSS,* 6**,** 3021.

XU, M., PIRTSKHALAVA, T., FARR, J. N., WEIGAND, B. M., PALMER, A. K., WEIVODA,
M. M., INMAN, C. L., OGRODNIK, M. B., HACHFELD, C. M., FRASER, D. G., ONKEN, J.
L., JOHNSON, K. O., VERZOSA, G. C., LANGHI, L. G. P., WEIGL, M., GIORGADZE, N.,
LEBRASSEUR, N. K., MILLER, J. D., JURK, D., SINGH, R. J., ALLISON, D. B., EJIMA, K.,
HUBBARD, G. B., IKENO, Y., CUBRO, H., GAROVIC, V. D., HOU, X., WEROHA, S. J.,
ROBBINS, P. D., NIEDERNHOFER, L. J., KHOSLA, S., TCHKONIA, T. & KIRKLAND, J. L.
2018. Senolytics improve physical function and increase lifespan in old age. *Nat Med,* 24**,** 1246-
1256.

ZHANG, H., RYU, D., WU, Y., GARIANI, K., WANG, X., LUAN, P., D'AMICO, D., ROPELLE,
E. R., LUTOLF, M. P., AEBERSOLD, R., SCHOONJANS, K., MENZIES, K. J. & AUWERX, J.
2016. NAD(+) repletion improves mitochondrial and stem cell function and enhances life span in
mice. *Science,* 352**,** 1436-43.

Figure. 1. The YFP$^+$ cells from *Pax7^{CreERT2};Rosa^{YFP}* mice do not decrease with age. (**A**)

Scheme of MuSC labeling in *Pax7^{CreERT2};Rosa^{YFP}* mice (MuSC^{YFP}) and analysis at middle (Mid)

and geriatric age (Geri) using fluorescence-activated cell sorting (FACS) strategy for isolation of

YFP$^+$ MuSCs using traditional marker, VCAM1. (**B, C**) Representative plot and quantification for

VCAM1$^+$ MuSCs (**B**) and YFP$^+$ cells (**C**) in lineage negative (Lin$^-$) cells of Young (n=6 and n=8

each), Mid (n = 7, n = 8, each) and Geri (n = 6, n = 11, each) MuSC^{YFP}, Welch's *t* test and *F*-test;

p = 0.0441 and 0.0460 for (**B**), respectively. (**D**) Representative FACS histogram and

quantification of VCAM1 and ITGA7 expression in YFP$^+$ cells of Mid and Geri MuSC^{YFP}.

Quantification of relative VCAM1 and ITGA7 mean fluorescence intensity (MFI) in YFP$^+$ cells

of Mid (n = 12, n = 6, each) and Geri (n = 14, n = 8, each) MuSC^{YFP}, Welch's *t* test and *F*-test; *p*

= 0.0009 and <0.0001. (**E**) Representative images and quantification of immunohistochemistry

staining for Hoechst (Blue), YFP (Green), PAX7 (Yellow), and Laminin (Red) in the *Tibialis*

Anterior (TA) of MuSC^{YFP}. Arrows indicate PAX7$^+$/YFP$^+$ cells and arrowheads indicate PAX7$^-$

/YFP$^+$ cells under basal lamina. Quantification of the number of Pax7$^+$/YFP$^+$ and PAX7$^-$/YFP$^+$

cells from IHC staining of TA in Mid (n = 7) and Geri (n = 9) MuSC^{YFP}, two-way ANOVA

followed by Bonferroni's multiple comparison test, PAX7$^+$/YFP$^+$; *p* = 0.0205, PAX7$^-$/YFP$^+$; *p* =

0.0051. (**F**) Division of YFP$^+$ cells according to their VCAM1 MFI as VCAM1^{High} (V^H),

VCAM1^{Low} (V^L), and VCAM1^{Neg} (V$^-$) followed by representative images and quantification of

immunocytochemistry (ICC) staining for Hoechst (Blue), PAX7 (Red) of YFP$^+$ cells.

Quantification of PAX7 fluorescence intensity from isolated YFP$^+$ cells of Mid V^H, Geri V^H, V$^-$

(n = 5, each), and V^L (n=6), one-way ANOVA followed by Bonferroni's multiple comparison test;

p = 0.0004 (vs Mid V^H), and 0.0179, 0.0123 (Geri V^H vs V^L, V$^-$, respectively).

A *Pax7^{CreERT2};Rosa^{YFP}*

YFP⁺ Cells

YFP

VCAM1 / FSC

Uninjured-Mid / Injured-5dpi / Uninjured-Geri

B Hoechst / MYOG

Injured-5dpi

Geri-V⁻

50μm

C

MYOG⁺ cells in YFP⁺ cells (%)

YFP⁺ 5dpi | Vᴴ Mid | Vᴴ | Vᴸ | V⁻ Geri

D

UMAP_2 / UMAP_1

Unbiased Clustering
- Quiescent MuSC
- Activated MuSC
- Myoblast/Myocyte
- Myonuclei (Slow fiber)
- Myonuclei (Fast fiber)

E

	MuSC Quiescence	Cd34
		Spry1
		Sdc4
		Pax7
MuSC Marker		Itga7
		Vcam1
		Myf5
Myoblast Myocyte		Myod1
		Myog
		Myh3
		Myh1
Myofiber		Myh4
		Ckm

Mean expression in group

F

Public only

UMAP_2

+Mid-YFP⁺

+Geri-YFP⁺

UMAP_1

Figure 2. YFP$^+$/VCAM1$^-$ cells are quiescent MuSCs. (**A**) Representative FACS plot for YFP$^+$

cells from uninjured Mid and Geri, and injured-5dpi young (20-weeks-old) mice. (**B**, **C**)

Representative images (**B**) of ICC staining for Hoechst (Blue) and MYOG (Red), and

quantification (**C**) of MYOG$^+$ cells in each YFP$^+$ cell, one-way ANOVA followed by Bonferroni's

multiple comparison test; p = 0.0139 between YFP+ cells of injured-5dpi and Mid V^H. (**D**)

Uniform manifold approximation and projection (UMAP) plot of YFP$^+$ cells integrated with public

scRNA-seq data of MuSC-lineage mononucleated cells in uninjured/injured muscle colored by

cell types. (**E**) Matrix plot for representative genes of specific cell types for determining the

identities of each cluster. (**F**) The localization of Mid and Geri YFP$^+$ cells in this study on UMAP

plot of Fig. 2D.

A

7 days in GM

Mid | Geri

V^H | V^H | V^L | V^-

100μm

B

Relative Cell Number

- ○ V^H ⊢Mid
- ● V^H
- ● V^L ⊢Geri
- ● V^-

2d 3d 5d 7d 10d

*** ** *

C

Pax7^{CreERT2}; Rosa^{DTA}

Tmx

2w

BaCl₂ injury

21dpi

PBS or
30,000 YFP⁺ cells T.P.

D

Inj

50μm

E

Hoechst / Laminin

Mid – V^H | Geri – V^H | Geri – V^L | Geri – V^-

50μm

F

Regenerated Fiber CSA (μm²)

2.5K
2.0K
1.5K
1.0K
0.5K
0.0K

**** **** ** n.s. * *

NA

| PBS | V^H | V^H | V^L | V^- |
| | Mid | Geri | | |

Figure 3. Compromised but retained functionality in VCAM1⁻ MuSCs. (**A**) Representative

bright field images of Mid V^H, Geri V^H, V^L, and V^- MuSCs at 7 days culture in MuSC growth

media. (**B**) Proliferation measured by relative cell number at 2, 3, 5, 7, and 10 days, two-way

ANOVA followed by Bonferroni's multiple comparison test; p = 0.0029 and 0.0002 (Mid V^H vs

V^L and V^-, respectively) and p= 0.0102 and 0.0006 (Geri V^H vs VL and V^-, respectively). (**C**)

Experimental schematics of transplantation of YFP^+ cells to TA muscle of $Pax7^{CreERT2};Rosa^{DTA}$

mice. (**D, E**) Representative images of IHC staining for Hoechst (Blue) and Laminin (Red) in PBS

treated TA muscle section following injury. (**D**) Left side of the dotted line show areas affected by

BaCl2 injury and right side shows uninjured areas of the muscle. (**E**) IHC staining shows

centralized myofibers in all images, with notably smaller myofibers in Geri V^-. (**F**) Quantification

of cross sectional area of regenerated fibers from TA muscle of $Pax7^{CreERT2};Rosa^{DTA}$ mice

transplanted with Mid V^H, Geri V^H, V^L, and V^- at 21 dpi, n = 3 for each group. Total 43 data points

are outside of the axis. Kruskal-Wallis test followed by Dunn's multiple comparison test; p <

0.0001 (Mid and Geri V^H vs V^-) and p = 0.0059 (Mid V^H vs Geri V^L), 0.0149 (Geri V^H vs V^L) and

0.0143 (Geri V^L vs V^-).

Figure 4. Senescence characteristics of VCAM1⁻/YFP⁺ cells and their rejuvenation by DHT

supplementation. (**A**) The statistical significance of Kyoto-Encyclopedia of Genes and Genomes

(KEGG) pathways, associated with differentially expressed genes (DEGs) between Mid V^H and

Geri V^H, V^L and V^-, visualized as a heatmap. (**B**) Heatmap of relative expressions of representative

genes known to be altered in aged MuSCs compared to Mid V^H. (**C**) Experimental schematics of

senolytics, dasatinib and quercetin (DQ), treatment in MuSC^{YFP} at geriatric age and representative

FACS histogram of VCAM1 expression Mid, Geri, and Geri treated with SL (Geri+DQ) mice. (**D**)

Quantification of the percentage of VCAM1^{High} and VCAM1$^{Low/-}$cells divided by VCAM1

expression levels in Mid (n = 8) and Geri mice with (n = 5) or without (n = 11) senolytics, 2 tailed

student's t test, $p = 0.0051$ for Mid vs Geri in VCAM1^{High} and F test: $p = 0.0026$ for Geriatric vs

Geriatric+DQ in VCAM1$^{Low/-}$. (**E**) Experimental schematics of dihydrotestosterone (DHT)

treatment and FACS plot of VCAM1 expression levels for YFP⁺ cells in MuSC^{YFP}. (**F**)

Experimental schematics of DQ and DHT treatment and quantification of the percentage of

VCAM1^{High} MuSCs in Live/Lin– cells in Mid and Geri B6 mice (n = 4 in each), 2 tailed student's

t test, $p = 0.0113$ for Young vs Geri+DQ+T.

A

RNA seq database

Mid - V^H
Geri - V^H
Geri - V^L
Geri - V^-

High RPKM (>100 in all samples)

V^H ≈ V^- or V^H < V^- in RPKM

Located in cell surface

Expressing cell type -Tabula Muris Senis (GSE132042)

CD63 & CD200

B

Cd63

Cd200

C

PAX7 / CD63 / Laminin / Hoechst

10μm

PAX7 / CD200 / Laminin / Hoechst

D

VCAM1 vs YFP

CD200 vs CD63   96.3%

E

CD200 vs CD63

VCAM1 vs YFP   93.2%

Hoechst / Pax7

96.5 ± 2.8%

50μm

F

Mid

CD200 vs CD63

VCAM1

Geri

CD200 vs CD63

VCAM1

Figure 5. Detection of GERI-MuSCs using CD63/CD200-based FACS strategy. (**A**) Vann Diagram schematics showing criteria used to select MuSC marker candidate CD63 and CD200, from RNA seq database. (**B**) Relative gene expressions of *Cd63* and *Cd200* shown in RPKMs from Mid V^H, Geri V^H, V^L, and V^- MuSCs. (**C**, **D**) Representative IHC staining images for Hoechst (Blue), Laminin (Green), PAX7 (Yellow) with CD63 (red, **C**) or CD200 (red, **D**) in the TA muscle of Mid MuSC^{YFP}. (**E**) Representative FACS plot from young mice showing 96.3% of VCAM1$^+$/YFP$^+$ cells are CD63^{high}/CD200^{high}. (**F**) Representative FACS plot from young mice showing 93.2% of CD63^{high}/CD200^{high} cells are VCAM1$^+$/YFP$^+$, and ICC staining of this population shows that 96.5±2.8% are PAX7$^+$. (**G**) Representative FACS plot of MuSCs of *B6* Mid and Geri mice gated by CD63 and CD200 and their VCAM1 expression level.

Fig. S1. Quiescent MuSC characteristics of YFP⁺ cells. (**A**) Stacked bar graph of myogenic cell cluster proportion determined in Fig. 2.at Mid, Geri, and each days-post-injury (DPI) of public data. (**B**) Force-directed graph (left) and partition-based graph abstraction (PAGA) connectivities (right) of YFP⁺ cells integrated with public scRNA-seq data described in Fig. 2, colored by cell types. (**C**) Force-directed graph showing identities of cells based on the days-post-injury. (**D**) Force-directed graph showing the locations of YFP⁺ cells of Mid and Geri in this study. (**E**) The expression levels of representative genes of specific cell types shown in the force-directed graph.

A

Mid-V^H   Geri-V^H

Geri-V^L   Geri-V⁻

10μm

5dpi-YFP⁺

10μm

B

C

SMMC
Young
VH
V-

Fig. S2. VCAM1⁻ cells resemble quiescent MuSCs in size and differ from SMMCs. (A)

Representative bright field image of freshly isolated YFP$^+$ cells of injured-5dpi young (20-weeks-

old), Mid V^H, Geri V^H, V^L, and V$^-$. (**B**) Quantification of cell diameter in YFP$^+$ cells of injured-

5dpi, Mid V^H, Geri V^H, V^L, and V$^-$, one-way ANOVA followed by Bonferroni's multiple

comparison test; p <0.0001 for injured-5dpi vs all YFP$^+$ cell groups of uninjured muscle. (**C**)

Normalized expression of MuSC marker genes, Pax7 and Vcam1, and smooth muscle

mesenchymal cells marker genes, Pdgfrb, Myl9 and Acta2.

Fig. S3. Validation and basic analysis of transcriptomic data of YFP$^+$ cells. (**A**) Total number of sequenced reads of each sample. (**B**) The percentage of trimmed reads of each sample for quality control. (**C**) Dot plots of gene expression showing linear correlations of between samples of the same group. The number in the plot represents Spearman Coefficient. (**D**) Hierarchical clustering of bulk RNA-seq samples with gene expression profiles, showing sample similarity (x-axis) and gene similarity (y-axis), illustrating inter-group similarity and similarity between V^H of Mid and Geri and between V^L and V$^-$ of Geri. (**E**) Volcano plots showing differentially expressed genes (DEGs) between groups at a threshold of LFC > 1.2 and adjusted p value < 0.05. The numbers in the plot represent the number of up-regulated (red) and down-regulated (blue) genes compared to the sample indicated in the right of the panel. (**F**) Correlation analysis between the percentages of VCAM1^{High} and VCAM1$^{Low/-}$ cells within YFP$^+$/Live/Lin$^-$ MuSCs isolated from geriatric mice. Each dot represents an individual mouse. A significant negative correlation was observed (Pearson's $r = -0.847$, $p = 0.001$), indicating that a higher proportion of VCAM1^{High} MuSCs is associated with a lower proportion of VCAM1$^{Low/-}$ MuSCs. Shaded area represents the 95% confidence interval of the linear regression.

A

B

Geri + Vehicle   Geri + DHT

C

Cross Sectional Area (µm²)

Fig. S4. Rehabilitation of myofiber regeneration capacity in geriatric mice after DHT

supplementation.

(**A**) Experimental schematics geriatric B6 mice with DHT or vehicle tube implantation before

BaCl2 injury. (**B**) Representative images of immunohistochemistry staining for laminin in geriatric

mice with vehicle or DHT tube implantation. (**C**) Quantification of the cross-sectional area of the

21 dpi TA muscle of geriatric mice after vehicle or DHT tube implantation.